Dissociating instructive from permissive roles of brain circuits with reversible neural activity manipulations

Daniel Quintana[1*], Hayley A. Bounds[1,2*], Jennifer Brown[1], May Wang[1], J. Simon Wiegert[3,4], Hillel Adesnik[1,2**]

[1]Department of Neuroscience, University of California, Berkeley

[2]The Helen Wills Neuroscience Institute

[3]University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[4]Department of Neurophysiology, MCTN, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany

*Authors contributed equally

**Corresponding author

SUMMARY

Recent work has demonstrated that both permanent lesions and acute inactivation experiments can lead to erroneous conclusions about the causal role of brain areas in specific behaviors, casting serious doubt on major avenues by which neuroscientists study the brain. To overcome this challenge, we developed a three-stage optogenetic approach which leverages the ability to precisely control the temporal period of regional inactivation with either brief or sustained illumination, enabling investigators to dissociate between putative 'permissive' and 'instructive' roles of brain areas in behavior. We applied this approach to the mouse primary visual cortex (V1) to probe whether V1 is permissive or instructive for the detection low contrast stimuli. Acute inactivation of V1 drastically suppressed performance, but during persistent inactivation, the animals' contrast detection recovered to pre-silencing levels. This recovery was itself reversible, as returning the animals to intermittent V1 inactivation reinstated the behavioral deficit. These results argue that V1 is the default circuit mice use to detect visual stimuli, but in its absence, other regions can compensate for it. This novel, temporally controllable optogenetic perturbation paradigm should be useful in other brain circuits to assess whether they are instructive or permissive in a brain function or behavior.

INTRODUCTION

Neuroscientists rely on targeted perturbations to causally map functions in the brain[1]. Yet since the brain is highly interconnected, manipulation of one area can impact behavior through indirect effects on other brain regions, complicating the interpretation of such results[2,3]. Permanent lesions of one area can lead to dysfunction or degeneration of other brain regions, making it difficult to determine whether the functional consequences of the lesion are due to loss of the targeted region or to those that are indirectly damaged. Moreover, the often observed recovery of behavioral performance after a permanent lesion can further complicate the interpretation of lesion experiments: either the lesioned area was never directly involved, or another brain area was able to compensate for its loss[4,5].

Recent studies have highlighted how the results of acute and irreversible inactivation can conflict[4–6]. For example, acute inactivation of forebrain motor or sensory areas in animals can lead to profound behavioral deficits, but animals with permanent lesions are either unaffected or can recover performance within just a few days[4–6]. In one striking example, the behavioral deficit observed during acute inactivation of one forebrain region of the songbird could instead be attributed to an indirect effect on activity in a downstream area[4]. Once the downstream area homeostatically adjusted its activity, the animal's behavior recovered. This led the authors to propose that a brain region can either be "instructive" – providing essential information that is used in the computation for a behavioral task, or "permissive" – acutely required, but not providing essential information or computation. A permissive region is only 'required' in that it might provide tonic excitatory or inhibitory tone to a downstream region to generally set that region's gain or sensitivity (Fig. 1a). In song production, the forebrain region was merely permissive in the behavior rather than instructive.

Similarly, another study found conflicting effects of acutely optogenetically silencing or permanently lesioning the barrel cortex on performance in a whisker-dependent task. While acute silencing of barrel cortex impaired behavior, animals could entirely recover from an irreversible lesion of the barrel cortex within ~2 days, a process that was experience dependent[5]. This study offered two possible conclusions: either barrel cortex is instructive but is redundant with another area that quickly compensates for its loss[7] (Fig. 1b); or barrel cortex is permissive and acute inactivation transiently disrupts the instructive area. Yet another study found that mice trained to discriminate pure tones recovered their performance following auditory cortex lesions in just a few hours, but did not recover performance during prolonged optogenetic silencing over the same timeframe, raising the possibility that lesions may uniquely engage compensatory mechanisms[6]. Together, these studies highlight that existing approaches for probing the roles of brain areas in behavior cannot easily resolve whether an area is instructive or permissive. Furthermore, when multiple brain areas can both be instructive in a task, they also cannot clearly determine which one might be the 'default' area the brain primarily relies on to solve it, and which is only secondary.

These studies demonstrate that acute manipulations alone can lead to erroneous or ambiguous conclusions about the instructive role of a brain area, such as the cortex, in a specific behavior.

Yet acute, rapid, and reversible manipulations, such as with optogenetics[8–10], have become
central to the investigation of the brain and behavior. Compared to irreversible lesions, acute
manipulations avoid potential indirect neurodegenerative effects[2,11], permit rigorously
interleaved control trials, and can address the timescales for when neural activity in specific
regions is necessary for behavioral output[12,13]. However, the current inability to properly
interpret the results of such acute manipulations is a major roadblock in neuroscience. Thus, we
sought to develop a new paradigm that can help resolve the critical ambiguities associated with
acute perturbations, providing crucial additional evidence to determine whether an area is
permissive or instructive in a task, and if instructive, whether it is the default area that normally
instructs the behavior.

RESULTS

A flexible optogenetic strategy for both acute and persistent silencing of neural circuits.

We reasoned that we could help address the ambiguities of existing perturbation approaches
by employing a flexible optogenetic inactivation strategy in which one could acutely or
chronically silence a brain area in the same animal. With such an approach, one could first
measure the acute effects of inactivation of a brain area on a behavior, then chronically silence
it to mimic an irreversible lesion, and finally return the brain area to acute inactivation (Fig. 1c).
In cases where acute inactivation impairs behavior, animals might recover behavioral function
during chronic optogenetic silencing, as they often do after irreversible lesions. In such cases,
one might be able resolve the apparent conflict by finally returning to acute, intermittent
silencing in the recovered animals. This last test could be critically informative: if recovered
animals no longer showed any effect of acute silencing, this would suggest that the brain region
only played a permissive role in the initial condition (Fig. 1c). Conversely, if a return to acute
silencing in recovered animals reinstates the behavioral deficit, the brain region is more likely to
directly contribute to the computational process in an instructive manner.

To develop and leverage the proposed optogenetic strategy, we focused on the primary visual
cortex (V1), which is among the best studied areas in the mammalian brain, but whose role in
different aspects of visual perception remains in doubt. Prior work has shown that inactivating
or lesioning V1 severely compromises visual function[3,12,14–16], including in contrast sensitivity
(particularly at higher spatial frequencies and/or small sizes)[17–20], but that visual performance
can recover spontaneously, with further training, or sometimes even with lesions to other brain
areas[21–23]. We first confirmed these findings in mice trained on a visual contrast detection
task[18,24]. We quantified performance using the detection threshold, or c50, estimated by fitting
a psychometric function to the behavioral data. We found that permanently lesioning V1
resulted in an acute reduction in task performance, followed by a substantial, albeit incomplete
recovery over the subsequent week (Fig. 2 a-b; Pre vs Day 1: $p<0.05$, Pre vs. Days 7-10: $p<0.05$,
Day 1 vs Days 7-10: $p<0.05$. n=7 mice). To assess the extent to which surgery itself may have
contributed to the observed deficits, we lesioned primary somatosensory cortex (S1) in a
separate cohort of mice. Lesioning S1 had a dramatically smaller effect on visual behavior (Fig.
S1 a-b), demonstrating that the initial deficit we observed following V1 lesion was specific and

not due to surgery *per se* (Day 1 post lesion detection threshold, S1 lesion versus V1 lesion:
p<0.05, n=10 mice). These data argue that following V1 lesion, mice exhibit substantial, but
incomplete recovery of contrast sensitivity, raising two important questions: first, why does
visual behavior recover at all? Behavior might recover because V1 is merely permissive and the
brain homeostatically scaled up the gain of the instructive area. Alternatively, V1 is instructive,
but behavior recovered because the animal learned to perform the task using alternative brain
structures that are not normally required. Second, why is recovery incomplete? Either V1 is
essential for a small component of behavioral performance, or because lesioning V1 leads to
degeneration of other visual areas that are. Indeed, inspection of the primary visual thalamus in
V1-lesioned mice showed a profound reduction in the number of thalamic neurons on the side
ipsilateral to the lesion (Fig. 2c-d p<0.01, n=7 mice). Thus, based on these lesion results alone,
we cannot conclude whether V1 is instructive or permissive in the contrast detection task, and
we cannot rule out that distal degeneration of other visual areas is the source of lasting the
behavioral deficit. These data highlight key problems with interpreting the results of
conventional lesioning studies.

To begin to address these challenges, we next sought to use a reversible perturbation to clarify
the role of V1 in basic visual functions in mice as other studies have done [18,19,25]. We silenced
excitatory neurons by optogenetically stimulating GABAergic neurons which caused a profound
reduction in contrast sensitivity on illumination trials (Fig. 2e; PV-Cre; Rosa-LSL-ChR2(Ai32):
p<0.0001, n=10 mice). Similar illumination in mice not expressing any opsin did not affect
performance (Fig. S1c, p = 0.86, paired t-test, n=4 mice), controlling for any direct effects of the
illumination on behavior. At face value, this acute inactivation experiment supports an
instructive role of V1 in contrast detection, but it seemingly conflicts with the substantial
recovery seen in mice with V1 entirely removed (Fig. 2b). Taken together, the results of
irreversible and acute, reversible lesions of V1 highlight the difficulty of drawing definitive
conclusions about its role in behavior even with a combination of approaches.

While optogenetic manipulations have conventionally been used to silence brain regions or cell
types only briefly, we reasoned that prolonged optogenetic silencing might add crucial
information by mimicking a permanent lesion, but ultimately still being reversible. For this
approach we employed direct optogenetic silencing of cortical excitatory neurons by expressing
the blue light silencer stGtaCR2 via the construct somBiPOLES[26]. To test the efficacy of silencing
neural activity with BiPOLES under both intermittent and persistent illumination conditions, we
recorded neural activity with multielectrode arrays across the layers of V1 while illuminating
the cortex briefly on every third trial, and then persistently for 40 continuous minutes
($2mW/mm^2$, 455 nm) (Fig. 3a-c). In BiPOLES-expressing mice, brief illumination resulted in
immediate and strong suppression of both spontaneous and visually evoked neural activity (Fig.
3c-e; 83±4% reduction at 100% contrast, p<0.05, Wilcoxon signed rank test). Next, we asked if
uninterrupted sustained illumination could effectively silence cortical activity for the entire
duration of the illumination, and whether neural activity would rapidly recover after cessation
of illumination to rule out potential photo-toxic effects. Indeed, during persistent illumination,
the mean firing rate averaged over all recorded units was suppressed by 76±4% (Fig. 3d, e; at
100% contrast, p<0.05, Wilcoxon sign-rank test). After the cessation of sustained illumination,

the mean activity was indistinguishable from baseline levels (Fig. 3e; p=0.53, Wilcoxon signed
rank test), arguing against potential issues with prolonged illumination[26] at this light level,
2mW/mm^2, or that prolonged flux of ionic currents might lead to cellular alterations[27]. These
experiments confirm the efficacy, stability, and reversibility of these acute and more persistent
optogenetic manipulations.
179
Acute and persistent effects of V1 inactivation in a visual detection task
Equipped with this flexible optogenetic approach, we probed how acute versus persistent
optogenetic inactivation of V1 would impact performance on the task. We trained somBiPOLES-
expressing mice on the same visual contrast detection task used above[18,24] (Fig. 3a) and
quantified performance using the c50, or detection threshold, as above. Consistent with results
above in which we silenced the cortex by stimulating PV neurons, intermittent silencing
excitatory neurons severely impaired performance (Fig. 4b, left; Fig. S2: p<0.001; paired t-test,
n=13 mice).
188
To better mimic a permanent lesion, we then illuminated V1 for the entire duration of each
behavioral session. Notably, this avoids irreversibly lesioning the tissue so that silencing can be
switched off in future sessions. Moreover, it avoids silencing V1 outside the task, such as in the
home cage, which might on its own cause local or downstream circuit reorganizations[30-33]. We
applied this persistent illumination scheme to the same mice as above to test if mice might
recover performance over time despite continued suppression of V1. Between the last day of
intermittent silencing and the first day of persistent silencing, we included a 'transition day' in
which the first half of the session employed intermittent silencing and the second half
employed persistent silencing to look for fast changes in the role of V1 in behavior. After this
day, the optogenetic light was on for the entirety of each session for six consecutive days
(session duration: 57±13 minutes; mean ± SD). On average, performance during the persistent
silencing was not significantly different from light on performance in the intermittent portion
(Fig. 4e; Transition Day 1 (T1); Intermittent light on (I_{on}) vs Persistent: p=0.12, I_{on} vs I_{off}: p<0.001,
I_{off} vs P: p<0.01; n=13 mice). However, on the first full day of persistent silencing ('P1'), the mice
substantially recovered performance on the task such that their detection threshold under
persistent illumination was not significantly different from their performance without
illumination on the last full day of intermittent silencing (Fig. 4f-h; Intermediate silencing day 6
(I6) vs persistent day 1 (P1); $I6_{on}$ vs P1: p<0.05, $I6_{on}$ vs $I6_{off}$: p<0.01, $I6_{off}$ vs P1: p=0.12; n=13
mice). These data demonstrate that mice can recover much of their contrast sensitivity even in
the continued suppression of V1 activity. Moreover, the very rapid recovery that we observed
on the first day of persistent silencing imply that mice can rapidly switch to using alternative
circuits to detect contrast. This, along with the recovery observed after mechanical lesion,
suggests that normal V1 activity is not absolutely required for task performance. Thus, so far,
the data leave it unclear whether V1 is instructive or permissive in the task.
213
Reversion to a reliance on V1 activity after cessation of persistent silencing.
The data above show that the mice can adapt to solve a contrast detection task when V1 is very
strongly suppressed or even entirely ablated by exploiting alternative brain circuits. Thus, V1 is

not required for task performance, but may still be instructive in the normal condition. We
reasoned that we could address this key question by reverting the persistently silenced and
behaviorally recovered mice to intermittent silencing. If intermittent silencing of V1 no longer
had a behavioral effect, this would imply that the cortex is probably not the default circuit
instructing performance on this visual task and is likely merely permissive. Conversely, if mice
instead return to failing on intermittent V1 silencing trials, this will imply that V1 is the default
circuit that actively instructs perceptual behavior.

To discriminate between these possibilities, we reverted the persistently silenced animals to
intermittent silencing the cortex only on 33% of trials, as before (intermittent repeat, IR). To
resolve any fast changes in behavioral performance, we started with a second transition day
composed of persistent silencing in the first half and intermittent silencing in the second half.
Then we began the intermittent repeat block of four sessions of intermittent silencing alone.
Surprisingly, we found that during the first day of intermittent repeat (IR1) block, on average
mice returned to showing a dramatic reduction in performance during intermittent silencing
trials (Fig. 4b,c,g IR1$_{on}$ vs P: p<0.05, IR$_{on}$ vs IR$_{off}$: p<0.05, I$_{off}$ vs P: p=0.15; n=11 mice). By the end
of the intermittent repeat sessions (IR4), the performance in intermittent silencing trials was
the same before as in at the end of the original intermittent silencing block (I6), before the
behavioral recovery during persistent illumination (Fig S3a; p=0.44, paired t-test, n=13 mice).
This reversion to requiring V1 activity is consistent with V1 playing an instructive rather than a
merely permissive role in the visual task.

Again, to address how fast this reversion to relying on V1 activity might occur, we examined the
transition session between the P and IR blocks. During the intermittent half of the session,
there was already a significant difference between light off and light on (Fig 4h; Transition Day 2
(T2); Pairwise comparisons: IR$_{on}$ vs P: p=0.18, IR$_{on}$ vs IR$_{off}$: p<0.05, IR$_{off}$ vs P: p=0.24; n=13 mice).
These findings emphasize how quickly mice go back to relying on V1 activity. More generally,
these results with a return to intermittent silencing demonstrate that the behavioral recovery
observed during persistent inactivation of V1 is itself reversible: once the animal regains access
to normal V1 activity it goes back to relying on it. This further implies that V1 is instructive,
rather than permissive.

Another advantage of our flexible optogenetic paradigm is that V1 has normal activity between
each behavioral session, which can help further disambiguate between instructive and
permissive roles. These 'off' periods, where V1 activity is unperturbed, might allow potential
homeostatic adjustments in other brain areas to revert between sessions. To test this, we
carefully analyzed performance across persistent silencing sessions. If homeostatic recovery of
other brain areas explains behavioral recovery during V1 silencing, we would expect that it
would likewise revert between persistent silencing sessions (when V1 activity is normal). In this
case, performance should be worse at the start of each persistent silencing session and recover
by its end. However, hit rates in the first fifth of a persistent session were comparable to those
at the end of the prior session (Fig. S3b,c, p=0.82, n=13 mice), suggesting that whatever
recovery occurs, it does not occur repeatedly within each session. This provides additional
evidence against a homeostatic explanation for the behavioral recovery during persistent

silencing, implying instead that the mice learn to use an auxiliary brain area to detect visual
stimuli in the absence of V1 activity.
263
DISCUSSION

Here, we present a simple and potentially broadly applicable three stage paradigm for using
optogenetics to help disambiguate whether a given brain circuit is instructive or permissive in a
behavior. In stage one, acute inactivation assesses whether the target circuit is involved at all
but does not discriminate between an instructive or permissive role. In stage two, sustained
inactivation assesses whether the behavior can recover, either due to relearning with
alternative circuits or due to homeostasis. Finally, when there is behavioral recovery, stage
three returns to acute inactivation to help determine whether the targeted circuit is instructive
or permissive.
273
Our data on mouse V1 supports a model in which V1 is instructive in perceptual behavior and
not merely permissive. It is the result from the return to intermittent inactivation that best
distinguishes between the multiple competing explanations for why acute V1 suppression
impairs behavior, but persistent silencing yields recovery. First, the performance recovery could
have been explained by homeostatic recovery of a downstream area (e.g., the superior
colliculus) whose normal activity level depends on sensory cortical input. If this were true, then
when the mice were reverted to intermittent silencing, one would have expected the mice to
be resistant to cortical inactivation since the downstream area had homeostatically become
independent of cortical drive. However, this was not the case as mice quickly reverted to
depending on V1 activity. It is possible that the homeostatic recovery could itself have
disappeared in the intervening time between the last persistent inactivation session and the
first intermittent inactivation session of the subsequent block. However, if this were true it
should have happened between each of the persistent inactivation sessions, and then mice
should have shown transient poor performance at the beginning of each persistent inactivation
session. However, this was not the case (Fig. S3c), arguing against any spontaneous recovery
between days of persistent illumination.
290
In a second scenario, the recovery of performance during persistent silencing could have been
explained by the animals permanently relearning the task using alternative circuits that do not
depend on primary sensory cortical activity[5]. In this scenario, task relevant information is
available to the animal in both the primary sensory cortex and alternative structures
independently, and though originally the animal depended on V1 activity to solve the task, over
several days of training in its absence, the animal would have relearned the task via these
alternate circuits. In this scheme, again one would have expected that the animals would be
resistant to reversion to intermittent inactivation of V1 since they had learned to solve the task
independent of the primary sensory cortex. However, this was not the case, as again, animals
very quickly returned to showing poor performance upon reversion to intermittent silencing
(Fig. 4b,c,g,h). This does not support the second explanation.
302

In a third scenario, V1 is part of the default circuit that instructs the animal's downstream decision circuits to solve the perceptual task. When V1 is inactivated intermittently, the animal fails on the task since it is relying on primary sensory cortical output for instructing performance. When V1 activity is unavailable for long enough, the downstream decision circuits eventually adapt to employing an alternate source of information to solve the task that may always have been present but was originally ineffective at driving these circuits. This learned reliance on alternate circuits is, unlike in scenario two, impermanent: when primary sensory cortical activity again becomes available, the downstream circuits revert to ignoring this alternative source of information even on trials in which the primary sensory cortex is acutely inactivated. Our results support this third scenario; they argue that V1 is the default source of instructive information for solving a simple contrast detection task.

Taken together, this new multi-stage optogenetic approach can generally help address the challenges posed by potential off-target effects of acute circuit perturbations. In principle, any study already using optogenetic inactivation can execute this paradigm without substantial additional technical complexity, although it is important to confirm the efficacy of sustained optogenetic suppression. The main advantage is that using this paradigm should lead to much more interpretable results. This should obviate the need to exhaustively search for potential alternative neural pathways with which an animal can solve a task (although such a search would still be valuable). Moreover, it may aid in understanding the mechanisms of behavioral recovery after brain lesions[28], helping to delineate the default versus accessory circuits that can contribute to specific behaviors.

ACKNOWLEDGEMENTS

The authors thank J. Beyer and K. Gopakumar for technical support, N. Bhatla for performing V1 suction lesions, and K. Liang for stereology. This work was funded by NIH R01EY023756 and RF1MH120680. H. Adesnik is a Chan-Zuckerberg Biohub investigator. J.S.W. was funded by the German Research Foundation (SPP 1926, Project Nr.: 315380903 WI4485/3-2 and acknowledges Kathrin Sauter and Ingke Braren for technical support.

AUTHOR CONTRIBUTIONS

D.Q., J.B., H.A.B. and H.A. conceived the experiments. D.Q. performed all behavioral experiments with assistance from M.W. H.A. performed all electrophysiological recordings. H.A.B. developed the task and performed quantitative analysis. J.B. performed an initial series of behavioral and optogenetics experiments to define illumination parameters. J.S.W. developed and contributed the AAV-somBiPOLES vector. H.A. wrote the manuscript with help from D.Q. and H.A.B.

DECLARATION OF INTERESTS

 The authors declare no competing interests.

346
Fig. 1. A three-stage optogenetic inactivation paradigm for dissociating instructive from
permissive roles of a targeted brain circuit.
A) A permissive model for the circuit. In this model, Area A is only needed to keep Area B in a
functional state so that Area B instructs correct performance. When Area A is removed or
permanently silenced (bottom), Area B can homeostatically recover (curved green arrow) from
its loss of drive from Area A and restore correct behavior.
B) An instructive model for the circuit. In this model, Area A directly instructs correct
performance, but when it is removed or permanently silenced, a redundant accessory circuit
through Area B restores performance (bottom).
C) An approach to distinguish among these models with chronic, but reversible inactivation:
three successive stages of intermittent, persistent, and repeat intermittent silencing of the
targeted circuit. Bottom: schematic plots of behavioral performance on a relevant task showing
performance in each stage for light off (gray) and light on (blue) trials. For stage 2, the dotted
lines are the schematized performance data from stage one replotted for comparison. In this
example set of results, in Stage 1 inactivation reduces performance, but in stage 2 performance
recovers. In stage 3, intermittent inactivation either returns to reducing performance (top) or
no longer has any effect (bottom).

Figure 2. Conflicting effects of lesion and acute inactivation of V1 on behavioral performance
in visual detection task
A) Coronal section from a representative mouse stained with anti-NeuN, showing the extent of
the V1 lesion. Dashed circles indicate the locations of the bilateral dLGNs. Scale bar: 1 mm.
B) Left, psychometric performance curves for mice with V1 lesions measured on the day before
lesioning (Day -1, black), the day after lesion (Day 1, yellow), and Day 10 post-lesion (orange).
Lines represent Weibull function fits to the behavioral data. Right, boxplot showing detection
thresholds averaged across six pre-lesion days, Day 1 post-lesion, and Days 7–10 post-lesion.
Statistical comparisons: repeated-measures ANOVA, $p < 0.05$; paired t-tests with multiple
comparison correction: Pre vs. Day 1: $p < 0.05$, Pre vs. Days 7–10: $p < 0.05$; Day 1 vs. Days 7–10:
$p < 0.05$; n = 7 mice.
C) Quantification of dLGN cell counts comparing ipsilateral (to the V1 lesion) and contralateral
(to the V1 lesion) hemispheres ($p < 0.01$, paired t-test; n = 7 mice).
D) Expanded view of the images of left (ipsilateral to the V1 lesion) and right (contralateral to
the V1 lesion) dLGN from the section in (D). Scale bar: 100 μm.
E) Left, psychometric performance curve for PV-Cre;Rosa-LSL-ChR2(Ai32) mice during
intermittent silencing. Lines are Weibull function fits. Blue, light on, black, light off. Right,
boxplot of detection thresholds during intermittent photostimulation; $p<0.0001$, paired t-test.
n=10 mice.
389
390
391

392
393
Figure 3. Acute and persistent optogenetic suppression of cortical activity.
A) Left: schematic of the recording configuration. Right: Example image of a post-mortem slice
of visual cortex expressing the inhibitory opsin throughout the cortical layers. Scale bar: 100
μm. Bottom: schematic of the illumination procedure with an intermittent illumination block,
followed by 40 minutes of continuous persistent illumination, followed by a return to
intermittent illumination.
B) Example raster plot for all spikes on an example electrode during the acute/persistent
optogenetic suppression protocol. Left: trials with no illumination during the intermittent
phases. Right: trials during the illumination phases (blue rectangle); intermittent (pre),
persistent, and intermittent (post), as indicated by the blue shaded regions.
C) Averaged normalized spiking activity across all recorded tetrodes across all experiments (3
mice, 6 sessions, 25 tetrodes) during each phase of the optogenetic protocol for the 100%
contrast stimulus. Intermittent illumination for ~40 trials, followed by continuous illumination
for ~130 trials, followed by a return to intermittent illumination. Black dots correspond to no
light trials, blue dots indicate illumination trials. Each phase of illumination is schematized by
the blue bars above the plot; broken lines: intermittent; solid bar: persistent light.

D) Average contrast response plots across all recorded units in each phase of the silencing
experiment. Left: control (black) and intermittent illumination trials (blue). Middle: persistent
illumination trials. Right: return to the intermittent illumination.
E) Scatter plots of recording unit firing rates between the indicated conditions; 3 mice, 6
sessions, 25 tetrodes; Wilcoxon sign rank test. All error bars are s.e.m.
415

Fig. 4. Reversible recovery from persistent inactivation of visual cortex.
A) Left: Schematic of visual detection task. Right: Optogenetic trial structure: Light is on
transiently (~2 seconds) for 30% of trials (intermittent and intermittent repeat, left) or
chronically (persistent, right) for the duration of the session. Gray bar represents stimulus
presentation time, black line represents response window, time referenced to the start of the
stimulus window.

B) Top, Schematic of silencing blocks, excluding transition days. In the intermittent blocks, silencing was applied on a subset of trials. In the persistent block, silencing was continuous throughout each session. The intermittent repeat block had the same structure as the intermittent block. Bottom, psychometric curves of performance for n = 10 mice tested at the most common contrast set. Left, intermittent block, light blue shows light on performance and black is light off performance. Center, persistent block, performance during silencing shown in dark blue, with dashed lines indicating intermittent performance replotted from the left panel for comparison. Right, intermittent repeat block, colors as in intermittent.

C) Detection threshold in each day of the testing protocol. Transition days (T1 & T2) had one half of the session intermittent, one half persistent. n=13 mice.

D) Detection threshold in light off (black) vs light on (blue) trials for intermittent day 1 (I1) ($p<0.001$, n=13 mice, paired t-test).

E) Detection threshold for T1. In the first half of the session intermittent silencing occurred on light on (light blue) trials interspersed within light off (black) trials. In the second half light was turned on for the entire period for persistent silencing (dark blue). $p<0.001$, repeated measures ANOVA (rmANOVA). Pairwise comparisons: I_{on} vs I_{off}: $p<0.001$, I_{on} vs P: $p=0.08$, I_{off} vs P: $p<0.01$; multiple-comparisons corrected paired t-test; n=13 mice.

F) Detection threshold during light off vs light on trials for intermittent day 6 (I6) vs persistent day 1 (P1). $P<0.001$, rmANOVA. Pairwise comparisons: $I6_{on}$ vs P1: $p=0.12$, $I6_{on}$ vs $I6_{off}$: $p<0.001$, I_{off} vs P: $p<0.01$, multiple-comparisons corrected paired t-test; n=13 mice.

G) Detection thresholds in persistent day 6 (P6) versus intermittent silencing day 1 (IR1). $p<0.01$ rmANOVA. $IR1_{on}$ vs P6: $p<0.05$, $IR1_{on}$ vs $IR1_{off}$: $p<0.05$, $IR1_{off}$ vs P6: $p=0.15$, multiple-comparisons corrected paired t-test; n=11 mice.

H) Detection thresholds for T2, which is the inverse of T1 (e). $p<0.01$ rmANOVA. IR_{on} vs P: $p=0.18$, IR_{on} vs IR_{off}: $p<0.05$, IR_{off} vs P: $p=0.24$; n=13 mice. B-C) Data are shown as group mean ± s.e.m. D-H) Boxplots show the median (center line), interquartile range (box limits: 25th to 75th percentile), and whiskers (extending to the most extreme data points within 1.5× the interquartile range). Outliers beyond the whiskers are shown as individual points.

454
Fig S1. Control experiments for the specificity of V1 lesion and optogenetic inactivation on
contrast detection
A) Psychometric curves of performance of mice with S1 lesions on the day immediately prior to
lesion (Day -1, black), day after lesion (Day 1, dark gray), and Day 10 (light gray) after lesion.
Lines, Weibull function fits of the psychometric curve.
B) Performance of mice for six days before lesion (circles), tested hours after lesion (Day 0), and
then tested for ten days after lesion (squares). Orange, V1 lesion, n=7 mice, gray, S1 lesion, n=3
mice.
C) Boxplot showing detection thresholds in opsin negative controls averaged 3 sessions per
mouse (p = 0.85, paired t-test, n=4 mice).

470
Fig. S2. Behavioral performance curves for each mouse. Psychometric curves of performance
for each mouse during each stage of the task, fit with a Weibull function. Data are group mean
± s.e.m.

Fig. S3. Intermittent versus intermittent repeat and within-session performance comparisons.

A) Comparison of light on detection thresholds for intermittent day 6 (I6) and intermittent repeat day 4 (IR4) p=0.44, paired t-test, n=13 mice.

B) Comparison of, for persistent silencing days, last bin of performance on prior days vs bin on next day (Fig. S3b,c, p=0.82, paired t- test; n=13 mice).

C) Binned performance over each block (top: intermittent, middle: persistent, bottom: intermittent repeat) and for each session (1 to 6; left to right) for no light (black) and light (color) trials. Group mean ± s.e.m.

MATERIALS AND METHODS

Animals

Mice used for experiments in this study were C57/BL6J (Jax stock# 000664) and PV-IRES-Cre (Jax stock# 008069) crossed with Rosa-LSL-ChR2 (PV-Cre;Rosa-LSL-ChR2(Ai32)) (Jax stock# 012569)[35,36]. Mice were housed in cohorts of five or fewer in a reverse light:dark cycle of 12:12 hours, with experiments occurring during the dark phase. Both female and male mice were used. For optogenetic silencing with somBiPOLES, a total of C57/BL6 13 mice were used (7 males) aged 10 – 20 weeks. For the V1 silencing in Figure 1 10 PV-Cre; Ai32 mice were used (5 males) aged 8 – 15 weeks. For lesion experiments, a total of 11 C57/BL6 mice were used (3 males) aged 12 – 30 weeks.

Viral injection

Neonatal C57/BL6 mice (P3-P5) were cryoanesthetized and their head was placed in a ceramic mold and secured with non-abrasive tape. Viral solutions (AAV9-CaMKII-somBiPOLES-mCerulean, Addgene) were loaded into beveled glass injection needles (WPI) and mounted in a NanoJect II (Drummond). The needle was inserted through the skin and scalp over the left V1 (2.3 mm lateral and 0.1 mm anterior to lambda) using a micromanipulator (Sutter MP-285) at three locations, each spaced approximately 0.3 mm apart. Virus was injected at four depths within V1 (400, 300, 200, and 100 μm), with 20 nl of viral solution delivered per site at a rate of 10 nl/second.

Headpost surgery

All experiments were performed in accordance with the guidelines and regulations of the Animal Care and Use Committee of the University of California, Berkeley. For head fixation during behavioral and physiological experiments, a small custom stainless-steel headplate was surgically implanted. Briefly, adult mice (P70 – P 155) were anesthetized with 2-3% isoflurane and mounted in a stereotaxic apparatus. Buprenorphine (0.05-0.1 mg/kg SC) was administered for analgesia and dexamethasone (2 mg/kg SC) to prevent cerebral edema. Body temperature was monitored and maintained at 37°C. The scalp was removed, the fascia retracted, and the skull lightly scored with a drill bit. Vetbond (3M) was applied to the skull surface, and the headplate was fixed to the skull with Metabond (C&B). A fine-point marker was used to note the approximate location of the left primary visual cortex (V1; 2.7mm lateral, 0mm posterior to lambda). For all behavioral experiments involving mice injected with somBipoles, a glass cranial window was implanted over the marked site to provide optical access to the cortex. In V1 silencing comparison experiments, the skull was instead thinned and covered with a single layer of Vetbond to enhance light penetration. Similarly, for electrophysiological recordings, the skull was thinned to transparency on the day of the experiment to ensure optimal electrode access and signal quality. Mice received buprenorphine and meloxicam for pain management and could recover for at least three days before being placed on water restriction.

Lesion surgery

Water restricted adult mice (P75-P180) previously headplated (see headpost surgery for procedure) and trained to perform the behavioral paradigm were anesthetized using isoflurane

2.5% and mounted in a stereotaxic apparatus. Buprenorphine (0.05-0.1 mg/kg SC) was
administered at anesthetic induction. The eyes of the animal are covered with ocular lubricant
to keep eyes moist. A dental burr was used to perform a 3 mm × 4 mm rectangular craniotomy
over the left visual cortex (V1; 2.7mm lateral, 0mm posterior to lambda). Exposed cortical tissue
was aspirated using a vacuum-attached blunt needle (16-30 awg). The lesion cavity is extended
ventrally until the alveus surface of the hippocampus is exposed. The cavity was covered and
sealed using a glass coverslip. Animal received meloxicam and buprenorphine post-surgery for
analgesia. Mice performed on the behavioral paradigm 2-4 hours post-surgery and
performance is reported as lesion day.
547
548
In vivo electrophysiology recordings
Mice were anesthetized with isoflurane (2.5%), and a small craniotomy (<250 microns) was
opened over V1 with a fine needle. Mice were transferred to the electrophysiology rig where
they were head fixed and allowed to wake and run freely on a circular treadmill. A 32-channel
multielectrode array (NeuroNexus, A1x32-5mm-25-177-a32) was guided into the brain using a
micromanipulator (Sutter Instruments) and a stereo microscope (Leica). Electrical activity was
amplified and digitized at 30kHz (spikeGadgets Inc.) and stored on a computer hard drive. The
bottom of the electrode was zeroed with the surface of the brain to determine z depth. The
electrode array was lowered ~650-750 microns into V1. A multimode fiber (400 µm diameter)
was positioned ~2-3mm from brain next to the recording electrode to deliver 455 nm LED light
during optogenetic illumination at a total of 2mW/mm^2 measured at the tip of the fiber.
560
Spike detection was performed across assigned tetrodes (four adjacent electrode contacts)
using SpikeGadgets threshold detection algorithm. Analysis was performed on all detected
spikes on each tetrode. Not all assigned tetrodes were used because the top tetrodes were
often located in L1 or outside of the brain. Statistically significant differences between
conditions were determined using the Wilcoxon sign rank test across all tetrodes. Since
mCerulean was not readily visible during adult surgeries, all experiments were initially
performed blind to successful opsin expression, and animals with no obvious physiological
optogenetic effects (strong activation by red light via the Chrimson in somBiPOLES), were
excluded, which correlated with weak, or no expression observed post-mortem.
570
Water restriction
Initial animal weight was recorded for 3 days before water restriction to establish baseline
weight. Mice were then placed water restriction. On training days, mice received most water
during the task. Mice were weighed after training and given additional water if their weight
dropped below 70% initial weight. Food was available *ad libitum*. The weight and health (quality
of the fur and nails, gait, posture) of the mice were monitored daily.
577
578 **Behavioral training on the contrast detection task:**
579
580 *Apparatus*
581

For the contrast detection task, the visual behavioral apparatus was controlled in real time
using an Arduino DUE (A000067) synchronized with Raspberry Pi 3+/4+
(BCM2837B0/SC0193(9)) and a custom software in Python, Arduino and Java[29]. The mouse was
head fixed bilaterally and placed on a 3D-printed holder tube. Mouse licking was measured
through the detection of changes in the capacitive load in a touch sensor (Adafruit,
AT42QT1070) caused by the contact of the mouse's tongue and a 0.05-inch diameter steel
lickport. Water was delivered through the lickport using a 2-way normally closed isolation valve
control (Neptune Research Inc.). Visual stimuli were displayed in a gamma corrected LCD
monitor (Podofo, 15.7 x 8.9 cm, 1024x600 pixels, 60 Hz refresh rate) positioned 12 cm from the
right eye with a 30-degree angle from the longitudinal axis.
592
593 *Behavioral training and task design*
594
595 Head fixed mice were trained to lick when they detected a visual stimulus of variable contrasts
596 displayed on the monitor. The training started after weight dropped below 80% of the initial
597 weight from water restriction. Prior to training, mice were habituated to head-fixation for two
598 daily sessions lasting 30 minutes each. The behavioral training protocol consisted of four stages.
599
In stage 1, mice were classically conditioned to lick in response to a visual stimulus that
consisted in a square patch with drifting sinusoidal grating (2Hz, 0.08 cycles/degree, 600 ms,
100% contrast, 34 visual degrees, black background luminescence). Each visual stimulus was
paired with a water reward delivered at the beginning of a 1000 ms response window. Mice
who achieved a performance of more than 90% of correct trials were transitioned to the next
phase.
606
In stage 2, mice were operantly conditioned to respond every time the same stimulus from
stage 1 was displayed (2Hz, 0.08 cycles/degree, 600 ms, 100% contrast, 34 visual degrees, black
background luminescence). Mice received a water reward when they licked within a 1000 ms
response window after the visual stimulus presentation. A no-lick period was enforced after
two seconds of the trial. Catch trials were introduced and consisted of 25% total trials. Mice
licking in a no-lick period, or a catch trial received a 5 to 9 seconds time out punishment. Mice
achieving performance with a hit rate >90% and false alarm rate <30% for two consecutive
sessions were switched to the next phase.
615
616 In stage 3, mice continued to train in the same behavioral paradigm with increased background
617 luminescence at different sessions until it reached mean luminescence. Then, the visual
618 stimulus size was decreased in different sessions until it reached 18 VD. The same performance
619 criteria as in stage 2 was used in every session before transitioning to any change.
620
In stage 4, mice started to detect stimuli of six different contrasts, pseudo-randomly
distributed. The intensities were customized for each mouse to provide a reliable detection
threshold. Mice were introduced to the optogenetic protocol once performance was stable for
more than 3 sessions. During photostimulation experiments, catch trials were 25% of all trials

and response window was reduced to 500 ms. On average, mice completed 375 trials per
session on non-transition days, and 420 trials for the entire session on transition days.
627
628   **Optogenetic stimulation during behavior:**
629
For V1 one-photon optogenetic experiments, an optical fiber (400 µm diameter, 0.39 nA;
M98L01, Thorlabs Inc.) coupled to a 455 nm LED (M4553, Thorlabs) and attached to a LED
driver (LEDD1B, Thorlabs), was placed above the mouse's window or skull thinned area using a
micromanipulator (MM3, Narishige). Light power was calibrated daily and measured from the
tip end of the fiber using a power meter (PM160T, Thorlabs). Photostimulation power was set
to 2.0 mW for somBiPOLES-injected mice, 5.0 mW for non-opsin-expressing controls, and
ranged from 2.0–8.0 mW for PV-Cre;Ai32 mice. During the intermittent blocks, optogenetic
stimulation were 33% of all trials, pseudo-randomly distributed in blocks. Light was delivered at
a random time prior to the stimulus onset, between 300 - 3000 ms, to avoid cueing the mouse
to the trial start. Light was sustained until the end of the response window. During the
persistent block, optogenetic stimulation was turned on prior to the beginning of the session
and remained on until the session ended. All som-BiPOLES mice underwent 14 consecutively
consisting of 6 sessions each I, P and IR blocks, with two additional transition days (T1 and T2)
between different blocks. However, only 4 IR sessions were included in the analysis, as some
mice were inadvertently exposed to incorrect contrast intensities during the final 2 sessions of
this stage. A custom designed removable light blocking system was attached to the head of the
mouse to prevent the mice from seeing the optogenetic stimulation light. A masking light and a
plastic eye patch over the left eye were also placed to prevent additional extraneous cues.
648
649   **Histology**
650
Mice were anesthetized with 5% isoflurane and ketamine in preparation for transcardial
perfusion with phosphate-buffered saline (PBS), followed by 4% paraformaldehyde (PFA).
Brains were removed, post-fixed overnight in PFA at 4°C, rinsed three times in PBS, then
cryoprotected in 30% sucrose for 1–2 days. Brains were monitored to ensure they initially
floated and subsequently sank, indicating complete cryoprotection. Afterward, a sliding
microtome (American Optical Company), frozen using dry ice and ethanol to ensure proper
adhesion and sectioning, was used to obtain 40 µm coronal sections of S1 and/or V1.
658
For NeuN staining, sections were incubated for 1 hour at 4°C on a rocker in blocking solution
containing 3% normal goat serum (NGS), 0.6% Triton X-100, 0.2% Tween-20, and 3% bovine
serum albumin (BSA) in PBS. Primary antibody incubation was performed overnight at 4°C using
Rabbit anti-NeuN (Abcam) diluted 1:10,000 in blocking buffer. The following day, sections were
washed in PBS with 0.25% Triton X-100 (PBS-T) and incubated for 1 hour at room temperature
with Alexa Fluor 405–conjugated Goat anti-Rabbit IgG H&L secondary antibody (1:1000;
Thermo Fisher Scientific). Sections were washed again in PBS-T, mounted onto glass microscope
slides, and allowed to dry before applying coverslips and Vectashield mounting medium (Vector
Laboratories) to reduce movement artifacts.
668

669 Imaging was conducted using an Olympus MVX10 Stereo Microscope with a 2× and 0.63x
670 objective lens, 1.8× zoom, 0.5 numerical aperture, and 500 ms exposure. Cell counting was
671 performed using a custom QuPath v0.2.3 pipeline automated for detection of NeuN-positive
672 cells within manually defined regions of interest across sequential brain slices, using intensity
673 thresholds and cell segmentation to quantify labeled nuclei.
674
675

Quantification and Statistical Analysis
Hit rate (% Hit) was defined as the number of trials go trials (contrast > 0) where the mouse
correctly licked within the response window divided by the total number of go trials. Days where
mice were run on the incorrect contrast levels due to experimenter error were excluded. For
detection threshold analysis, data for individual sessions were fit using a Weibull cumulative
distribution function that fit the false alarm (lower asymptote), lapse rate (upper asymptote),
slope and threshold parameter. Functions were fit separately for photostimulation and no-
photostimulation trials. Detection threshold was defined as the contrast at which the fit hit
ratewas halfway between the fit false alarm (lower asymptote) and fit lapse rate (upper
asymptote).
686
Statistical analyses were performed using MATLAB or python. The analyses performed were
repeated measures analysis of variance (rmANOVAs) with spherical correction when sphericity
was violated, paired t-tests with Holm-Bonferroni corrections where needed, independent t-
tests, and Wilcoxon rank sign tests. Unless otherwise noted, all tests were two-tailed and all plots
with error bars were reported as mean ± SEM, all boxplots show the median (center line),
interquartile range (box limits: 25th to 75th percentile), and whiskers (extending to the most
extreme data points within 1.5× the interquartile range). Outliers beyond the whiskers are shown
as individual points. Sample size was not predetermined using power analysis.
695

696 **Data availability**
697 All behavioral and electrophysiological data will be made available upon publication.

References

719
1. Lomber, S. G. The advantages and limitations of permanent or reversible deactivation

techniques in the assessment of neural function. *J Neurosci Methods* **86**, 109–117 (1999).

2. Carrera, E. & Tononi, G. Diaschisis: past, present, future. *Brain* **137**, 2408–2422 (2014).

3. Lashley, K. S. Mass action in cerebral function. *Science* **73**, 245–254 (1931).

4. Otchy, T. M. *et al.* Acute off-target effects of neural circuit manipulations. *Nature* **528**,

358–363 (2015).

5. Hong, Y. K., Lacefield, C. O., Rodgers, C. C. & Bruno, R. M. Sensation, movement and

learning in the absence of barrel cortex. *Nature* vol. 561 542–546 Preprint at

https://doi.org/10.1038/s41586-018-0527-y (2018).

6. O'sullivan, C., Weible, A. P. & Wehr, M. Auditory cortex contributes to discrimination of

pure tones. *eNeuro* **6**, (2019).

7. Cohen, J. D. & Castro-Alamancos, M. A. Early sensory pathways for detection of fearful

conditioned stimuli: tectal and thalamic relays. *J Neurosci* **27**, 7762–7776 (2007).

8. Fenno, L., Yizhar, O. & Deisseroth, K. The development and application of optogenetics.

Annu. Rev. Neurosci. **34**, 389–412 (2011).

9. Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G. & Deisseroth, K. Millisecond-timescale,

genetically targeted optical control of neural activity. *Nat. Neurosci.* **8**, 1263–1268 (2005).

10. Nagel, G. *et al.* Channelrhodopsin-2, a directly light-gated cation-selective membrane

channel. *Proc Natl Acad Sci U S A* **100**, 13940–13945 (2003).

11. Cowey, A., Alexander, I. & Stoerig, P. Transneuronal retrograde degeneration of retinal

ganglion cells and optic tract in hemianopic monkeys and humans. *Brain* **134**, 2149–2157

(2011).

12. Zatka-Haas, P., Steinmetz, N. A., Carandini, M. & Harris, K. D. Sensory coding and the causal

impact of mouse cortex in a visual decision. *Elife* **10**, (2021).

13. Resulaj, A., Ruediger, S., Olsen, S. R. & Scanziani, M. First spikes in visual cortex enable

perceptual discrimination. *Elife* **7**, (2018).

14. Petruno, S. K., Clark, R. E. & Reinagel, P. Evidence That Primary Visual Cortex Is Required

for Image, Orientation, and Motion Discrimination by Rats. *Plos One* **8**, (2013).

15. Burgess, C. P. *et al.* High-Yield Methods for Accurate Two-Alternative Visual Psychophysics

in Head-Fixed Mice. *Cell Rep.* **20**, 2513–2524 (2017).

16. Schneider, G. E. Two visual systems. *Science* **163**, 895–902 (1969).

17. Prusky, G. T. & Douglas, R. M. Characterization of mouse cortical spatial vision. *Vision Res*

44, 3411–3418 (2004).

18. Glickfeld, L. L., Histed, M. H. & Maunsell, J. H. R. Mouse Primary Visual Cortex Is Used to

Detect Both Orientation and Contrast Changes. *Journal of Neuroscience* **33**, 19416–19422

(2013).

19. Ruediger, S. & Scanziani, M. Learning speed and detection sensitivity controlled by distinct

cortico-fugal neurons in visual cortex. *Elife* **9**, (2020).

20. Del Rosario, J. *et al.* Lateral inhibition in V1 controls neural and perceptual contrast

sensitivity. *Nat. Neurosci.* **28**, 836–847 (2025).

21. Dean, P. Visual acuity in hooded rats: effects of superior collicular or posterior neocortical

lesions. *Brain Res* **156**, 17–31 (1978).

22. Sabel, B. A. & Kasten, E. Restoration of vision by training of residual functions. *Curr Opin*

Ophthalmol **11**, 430–436 (2000).

23. Sprague, J. M. Interaction of cortex and superior colliculus in mediation of visually guided

behavior in the cat. *Science* **153**, 1544–1547 (1966).

24. Histed, M. H., Carvalho, L. A. & Maunsell, J. H. R. Psychophysical measurement of contrast

sensitivity in the behaving mouse. *J. Neurophysiol.* **107**, 758–765 (2012).

25. Cone, J. J., Scantlen, M. D., Histed, M. H. & Maunsell, J. H. R. Different Inhibitory

Interneuron Cell Classes Make Distinct Contributions to Visual Contrast Perception. *Eneuro*

6, (2019).

26. Owen, S. F., Liu, M. H. & Kreitzer, A. C. Thermal constraints on in vivo optogenetic

manipulations. *Nat. Neurosci.* (2019) doi:10.1038/s41593-019-0422-3.

27. Mahn, M., Prigge, M., Ron, S., Levy, R. & Yizhar, O. Biophysical constraints of optogenetic

inhibition at presynaptic terminals. *Nat. Neurosci.* **19**, 554–556 (2016).

28. Krakauer, J. W., Carmichael, S. T., Corbett, D. & Wittenberg, G. F. Getting

neurorehabilitation right: what can be learned from animal models? *Neurorehabil Neural*

Repair **26**, 923–931 (2012).

35. Hippenmeyer, S. *et al.* A developmental switch in the response of DRG neurons to ETS

transcription factor signaling. *PLoS Biol.* **3**, e159 (2005).

36.   Madisen, L. *et al.* A toolbox of Cre-dependent optogenetic transgenic mice for light-

induced activation and silencing. *Nat. Neurosci.* **15**, 793–802 (2012).