Neural Correlates and Reinstatement of Recent and Remote Memory:

A Comparison Between Children and Young Adults

Iryna Schommartz[a,i], Philip F. Lembcke[b,c], Javier Ortiz-Tudela[a], Martin Bauer[b], M., Angela M.

Kaindl[c,d,e,f], Claudia Buss[b,h]*, and Yee Lee Shing[a,i]*

* Yee Lee Shing and Claudia Buss should be considered joint senior author.

[a]Department of Psychology, Goethe University Frankfurt, Frankfurt, Germany

[b]Charité – Universitätsmedizin Berlin, Department of Medical Psychology, Berlin, Germany

[c]Charité – Universitätsmedizin Berlin, Department of Pediatric Neurology, Berlin, Germany

[d]Charité – Universitätsmedizin Berlin, Center for Chronically Sick Children, Berlin, Germany

[e]Charité – Universitätsmedizin Berlin, Institute of Cell- and Neurobiology, Berlin, Germany

[f]Charité – Universitätsmedizin Berlin, Department of Pediatric Surgery, Berlin, Germany

[h]Development, Health and Disease Research Program, Department of Pediatrics, University of

California Irvine, USA

[i]Center for Individual Development and Adaptive Education of Children at Risk (IDeA),

Frankfurt, Germany

Corresponding authors at: Department of Psychology, Goethe University Frankfurt,

Frankfurt, Germany, Theodor-W.-Adorno-Platz, 6, 60323

E-mail addresses: schommartz@psych.uni-frankfurt.de (I. Schommartz),

shing@psych.uni-frankfurt.de (Y.L. Shing)

RESEARCH HIGHLIGHTS

- Children showed less robust memory consolidation across short and long delay compared to young adults.
- From short to long delay, children show differential neural upregulation for remote versus recent memory compared to young adults.
- Over time, both children and young adults showed reduced scene-specific reinstatement of neural patterns.
- Children relied more on gist-like neural representations in medial and ventrolateral prefrontal brain regions.

Abstract

Memory consolidation tends to be less robust in childhood than adulthood. However, little is known about the corresponding functional differences in the developing brain that may underlie age-related differences in retention of memories over time. This study examined system-level memory consolidation of object-scene associations after learning (immediate delay), one night of sleep (short delay), as well as two weeks (long delay) in 5-to-7-year-old children (n = 49) and in young adults (n = 39), as a reference group with mature consolidation systems. Particularly, we characterized how functional neural activation and reinstatement of neural patterns change over time, assessed by functional magnetic resonance imaging combined with representational similarity analysis (RSA). Our results showed that memory consolidation in children was less robust and strong (i.e., more forgetting) compared to young adults. Contrasting correctly retained remote versus recent memories across time delay, children showed less upregulation in posterior parahippocampal gyrus, lateral occipital cortex, and cerebellum than adults. In addition, both children and adults showed decrease in scene-specific neural reinstatement over time, indicating time-related decay of detailed differentiated memories. At the same time, we observed the emergence of generic gist-like neural representations in prefrontal brain regions uniquely in children, indicating qualitative difference in memory trace in children. Taken together, 5-to-7-year-old children, compared to young adults, show less robust memory consolidation, possibly due to difficulties in engaging in differentiated neural representations in neocortical mnemonic regions during retrieval of remote memories, coupled with relying more on gist-like generic neural representations.

Keywords: object-scene associations, memory consolidation, representational similarity analysis, neural reinstatement, neural representations, drift diffusion modelling

INTRODUCTION

Every day we form new memories that may become long-lasting through memory consolidation,

a complex process in flux between encoding and retrieval (Dudai, 2012; Josselyn et al., 2015;

Moscovitch & Gilboa, 2022; Semon, 1921).During systems-level consolidation, memory

representations and traces are reorganized across medial temporal lobe and neocortical brain

networks (Ranganath & Ritchey, 2012; Ritchey & Cooper, 2020). These networks include brain

regions that are involved both in initial encoding and in integration of new memories as time

passes (Axmacher & Rasch, 2017; Dudai, 2012; Moscovitch & Gilboa, 2022; Squire et al.,

2015). While decades of work have shed light on general neural mechanisms of memory

consolidation in adults (Moscovitch & Gilboa, 2022; Sekeres et al., 2017a; Winocur &

Moscovitch, 2011), much less is known about neural mechanisms that support memory

consolidation in children – a knowledge gap that we aimed to address with the current study.

Neural correlates of memory consolidation

Learning through repeated activation and reinstatement is one way to rapidly stabilize memory

traces and make them accessible upon retrieval (Dudai, 2004; Nader & Hardt, 2009; Teyler &

Rudy, 2007). For instance, in young adults, repeated exposure to word-image pairs during

encoding, compared to single exposure, was shown to accelerate memory consolidation. This is

achieved through enhanced replay of repeated events in the retrosplenial cortex (RSC) and the

medial prefrontal cortex (PFC), as well as via increased hippocampal (HC)-cortical replay that

promotes the associative word-object memories (Yu et al., 2022). In another study by Brodt et al.

(2016), it was found that during repeated spatial navigation in a virtual environment, activation

in the posterior parietal cortex (PPC), especially the precuneus, increased and remained elevated

after 24 hours, while HC activity and HC-PPC connectivity declined with repeated encoding

rounds (Brodt et al., 2016). In addition, neocortical plasticity measured by diffusion-weighted

magnetic resonance imaging in the PPC (Brodt et al., 2018) and the cerebellum (Stroukov et al.,

2022) supported rapid cortical storage of memory traces for object-location associations after

repeated exposure in young adults 1 hour and 12 hours post-learning. Taken together, these

findings indicate that repeated learning in young adults promotes fast creation of neural memory

representations, which can remain stable for at least 24 hours and predict behavioural mnemonic

performance.

Memory consolidation of well-learnt information does not end with the last learning
cycle, but undergoes further neural reorganizing and modification over time (Roüast &
Schönauer, 2023; Sekeres et al., 2017). For example, during cued recall of face-location
associations, young adults who were tested 24 hours after learning, compared to 15 minutes,
showed increased activation in the precuneus, inferior frontal gyrus (IFG), and fusiform gyrus,
whereas the hippocampus showed a decrease in activation (Takashima et al., 2009). Similarly,
increased activation in the anterior temporal cortex during the retrieval of studied figure pairs
eight weeks prior was observed, while increased activation in the HC was shown for pairs
learned immediately before retrieval (Yamashita et al., 2009). Furthermore, delayed retrieval of
naturalistic video clips after a delay of seven days in young adults was associated with increased
activations in the lateral and medial PFC and decreases in HC and parahippocampal (PHG)
activations over time (Sekeres et al., 2021). This is convergent with the notion that the role of the
prefrontal cortex increases during recollection as consolidation progresses over time (Milton et
al., 2011). Moreover, subsequently recollected memories showed higher post-rest HC- lateral
occipital cortex (LOC) connectivity specifically related to scene-related mnemonic content,
indicating the role of LOC in associative memory consolidation (Tambini et al., 2010). On the
other hand, HC activation has been reported to remain stable after seven days (Sekeres, Winocur,
Moscovitch, et al., 2018) three months (Harand et al., 2012) or even years (Söderlund et al.,
2012) for consistent episodic memories that retained contextual details.

To summarize, in alignment with the Multiple Trace Theory (Nadel et al., 2000; Nadel &
Moscovitch, 1997), studies have shown that memories of well-learned information increasingly
engage cortical regions over time. There regions include the prefrontal, parietal, occipital, and
anterior temporal brain areas, supporting the retrieval of general and schematic memories, as
well as complex associative information. In line with the Standard Consolidation Theory, some
studies have demonstrated a decrease in the recruitment of the HC over time (Squire & Alvarez,
1995). Conversely, and converging with the Contextual Binding Theory (Yonelinas et al., 2019)
and the Multiple Trace Theory, some studies have shown that hippocampal involvement lingers
over time, particularly for detailed and contextual memories. However, most research has
focused on only a selected delay window and solely on young adults.

Mnemonic transformation and reinstatement across consolidation

In addition to changes in neural activation during mnemonic retrieval over time, it is important to

characterize the transformations and reinstatement of neural representations – that is, distinctive

pattern of neural activity generated by a specific memory (Averbeck et al., 2006; Kriegeskorte,

2008; Kriegeskorte & Kievit, 2013) – as these multivariate patterns of neural activity may

change over time. For example, memory for perceptual details often declines over time, while

memory for gist may tends to remain more stable, suggesting differential temporal trajectories of

transformation (Sekeres et al., 2016). According to Fuzzy Trace Theory (Reyna & Brainerd,

1995, 1998) and Trace Transformation Theory (Moscovitch & Gilboa, 2022), detailed and gist-

like memories may be uniquely present or coexist, depending on the strength of formed

memories. For instance, Diamond et al. (2020) showed that the specific accurate nature may be

preserved for correctly recalled memories. In other instances, initially weak detailed memories

may be reorganized over time, with lingering specific memories and parallel creation of gist-like

generic memories. Further research supports suggest the idea that memory traces undergo

transformation and abstraction beyond simple perceptual reinstatement (Chen et al., 2017; St-

Laurent & Buchsbaum, 2019; Ye et al., 2020), pointing to the development of schematic, generic

representations. However, relatively little is known about how the neural representation of well-

learned memories change over the consolidation period – particularly how similar patterns of

neural activity are reactivated upon retrieved again (Clarke et al., 2022; Deng et al., 2021).

Using representational similarity analysis (RSA; Kriegeskorte, 2008), Tompary & Davachi

(2017) showed that a one-week delay led to differential memory reorganisation in HC and mPFC

for memories with and without overlapping features. Specifically, after a one-week mnemonic

representations became more similar for memories with overlapping features, indicating

consolidation-related gist-like neural reorganization. Moreover, the authors showed memory-

specific reinstatement of neural patterns for specific memories in the right HC, indicated by

significant encoding-retrieval similarity for remote but not recent memories. Comparing neural

reinstatement of visual clips during encoding, immediate, and delayed recall (after 1-week-

period), Oedekoven et al. (2017) showed reliable reinstatement in core retrieval networks,

including the precuneus, medial temporal gyrus, occipital gyrus, HC, and PHG among others. In

contrast to Tompary and Davachi (2018), this study found no time-related differences in

reinstatement effects. Therefore, the findings on memory reinstatement are mixed, and, to date,

no study have directly tracked the neural representations of memory traces for perceptual

together with more abstract, gist-like features (e.g., semantic categories).

Neural correlates of memory consolidation and mnemonic transformation and

reinstatement in middle childhood

Brain regions involved in memory consolidation show protracted developmental trajectories

from early to late childhood (Badre & Wagner, 2007b; Ghetti & Bunge, 2012c; Gogtay et al.,

2004; Keresztes et al., 2022; Lenroot & Giedd, 2006; Mills et al., 2016; Ofen et al., 2007; Shing

et al., 2008), which could lead to differences in neural activity and/or patterns and subsequently

mnemonic reinstatement between children and adults. For instance, univariate selectivity was

reduced in children, while fine-grained neural representational similarity along the ventral visual

stream was similar in 5-11 years old children and adults (Cohen et al., 2019; Golarai et al.,

2015). Fandakova et al. (2019) also showed that the neural representational distinctiveness of

information during encoding was similar in 8-to-15-year-old children and adults in the RSC,

LOC and PHG. The fidelity of neural representations was also associated with subsequent

memory in a similar way between children and adults. Overall, although these findings did not

address the question of neural reinstatement directly in children, they suggest that mnemonic

reinstatement may develop prior to univariate selectivity. However, it is yet to be investigated.

Moreover, it is unclear whether the age-related differences in neural activation and reinstatement

mentioned above are similar for memory consolidation. Specifically, to what extent does

consolidation-related transformation of neural representations occur, and how does it impact

neural reinstatement of mnemonic content in the developing brain?

In middle childhood, the trade-off between retaining vivid, detail-rich memories and their

transformation into vague, gist-like memories due to delay may be more pronounced. Brainerd et

al., (2002) demonstrated that, during development, specific memory and gist-memory for events

emerge together. However, as children mature, they exhibit more false memories based on gist in

the absence of exact memories for the events. On the other hand, Keresztes et al. (2018)

postulated that younger children tend to rely more on generalization when forming new

memories, while older children and adults use more specific detail-rich information, suggesting a

shift from generalization to specificity as children mature. Hence, there are some inconsistencies

in the theoretical postulations and findings regarding item-specific and gist-based memories that

may impact memory consolidation in middle childhood. Investigation on the neural

reinstatement patterns of item-specific and gist-like memories across time may add to the
understanding of these inconsistencies in children.

Aim of the current study

In this study, we examined the univariate neural activation and multivariate neural reinstatement
patterns of memories for object-location associations across a short delay (after one night of
sleep) and a long delay (after a 2-week period), relative to recently consolidated memories (after
30 minutes). Children (5-to-7-year-old) were compared to young adults serving as a reference
group with a mature memory consolidation system. We selected 5 to 7 years as the age range of
interest because previous studies showed a large improvement in associative memory around this
age (Sluzenski, Newcombe, & Kovacs, 2006). Practically, this is also the youngest age range in
which MRI scanning coupled with active task execution could be applied relatively successfully.

We hypothesized (i) according to the Multiple Trace Theory, an increasing involvement of
prefrontal, parietal, cerebellar, occipital and PHG brain regions over time in adults in comparison
to children, as these regions are still maturing in preschool and early school-aged children
(Ghetti & Bunge, 2012b; Keresztes et al., 2022; Lebel et al., 2012; Shing et al., 2008, 2010a); (ii)
according to the Contextual Binding Theory, the Multiple Trace Theory, and supported by the
evidence from Sekeres, Winocur, & Moscovitch (2018), a stable involvement of HC over time in
adults and children due to relative maturity of the HC in middle childhood and detailed
contextual nature of the repeatedly learned information, as our task emphasizes spatial-
contextual binding of objects within scenes (Keresztes et al., 2017; Nadel et al., 2000; Sekeres,
Winocur, Moscovitch, et al., 2018; Shing et al., 2008; Sluzenski et al., 2006; Yonelinas et al.,
2019); (iii) a decreasing neural reinstatement in all ROIs over time, with this decrease being
more pronounced in children compared to young adults (Cohen et al., 2019; Golarai et al., 2015);
(iv) qualitative differences in representational format between age groups. Specifically, we
expected more generic category-level gist-like memory representations in children, whereas
adults would retain more detailed item-specific reinstatement patterns over time due to
differences in the strength of formed memories and differences in underlying associative and
strategic components of memories (Reyna & Brainerd, 1995; Shing et al., 2008, 2010). This
assumption aligns with the Fuzzy Trace Theory (Brainerd & Reyna, 2002), which posits that
verbatim and gist representations are encoded in parallel and that verbatim memories can be
created without the extraction of gist. Our task design – involving repeated strategic learning –

may foster the emergence of consolidation-driven, categorical gist-like neural representation in
children. These are to be distinguished from mature semantic gist as defined by the FTT for
verbal material. Due to ongoing maturation of associative and strategic memory components and
their underlying neural substrates, children may be more inclined to extract generic category-
based gist information at the expense of detailed information.

RESULTS

Behavioural results

241   *Final Learning Performance*

Unique sets of object-location association pairs were learned on Day 0, Day 1, and Day 14.
During each initial encoding trial, participants were presented with an object within a congruent
scene (e.g., a fox in a spring pine tree forest), and were asked to memorize the exact location of
the object within the scene by creating a story and making "mental" pictures of the scene. The
choices for locations varied across scenes while they remained constant across time within
individuals. There were 18 unique key locations among which object could be distributed,
resulting in a heterogenous set of locations for objects. We employed an adaptive, repetitive
learning-to-criteria procedure to ensure initially strong memories (see Fig. 1A for the task
overview and Fig. 1B for experimental procedure overview).

251   Before the learning phase began, participants were instructed to create stories to help
them memorize the locations of objects within scenes. To familiarize themselves with this
strategy, they first practiced this strategy on two unique sets of five object-location associations.
Learning then commenced with the initial encoding block, followed by adaptively repeated
retrieval-encoding cycles to strengthen memory for the object locations. During these learning
cycles, participants were presented with the same scenes again, now with three rectangles
indicating possible locations for each previously learned object. The task followed a three-
alternative forced-choice task (3AFC) format, with the correct location randomly appearing on
the left, middle, or right. The rectangles were presented in close proximity within each scene,
requiring participants to recall location details with high precision. Participants were asked to
choose one rectangle that corresponded to the correct location of the object within the scene (Fig.
1A "Learning Cycles"). After each response – regardless of accuracy – the object was shown in
its correct location to reinforce learning. The learning cycles were repeated for a minimum of
two times and a maximum of four times, or until participants achieved at least 83% accuracy in

one cycle. This 83% threshold, established through pilot testing, served as a guideline for starting

the next learning cycle rather than as a strict learning criterion to exclude participants.

Participants who did not reach this threshold after four cycles were still included in the analysis

if their performance exceeded chance level (33%). All participants demonstrated at least average

cognitive abilities, as determined by a standardized intelligence test (see Table 1).

Figure 1

(A) Trial Structures in the Experimental Task. (i) *Initial Encoding*: Participants memorized object-location pairs

by creating a story or forming a "mental photo" of each scene, focusing on the exact location of the object within the

scene. (ii) *Learning Phase*: Participants selected one of three possible object locations and received feedback: a

happy face for correct responses, a sad face for incorrect ones, and a sleeping face for missed response. The correct

object-location pairing was then displayed again. (iii) *Retrieval Phase:* Conducted inside the MR scanner,

participants chose the object's location in the scene from three options without receiving feedback. **(B)**

Experimental Procedure. Testing took place across three days. On Day 0, participants learned 60 object-location

associations (*remote items*). On Day 1 (*short delay*), they learned 30 new object-location associations (*recent items*)

and retrieved 30 remote and 30 recent items. On Day 14 (*long delay*), participants learned another 30 new

associations and retrieved 30 remote and 30 recent items. Throughout all sessions, participants also completed socio-

demographic and psychometric questionnaires, which were distributed across sessions. *Note*: RT – reaction time; s –

second, fMRI – functional magnetic resonance imaging.

Concerning number of learning cycles, the linear mixed effects (LME) model revealed a

significant *Group* effect, $F_{(1,563)} = 7.09$, p .008, $w^2 = .01$, with children needing more learning

cycles to reach the learning criteria in comparison to adults, $b = -.43$, $t_{(563)} = -2.66$, $p = .008$,

within the defined minimum and maximum of learning cycles (Fig. 2A). Five child participants

did not reach the learning criteria after the fourth learning cycle and their final performance

ranged between 70% and 80%. The number of learning cycles did not differ between sessions as

revealed by non-significant *Session* effect and *Group x Session* interaction (all p > .40).

Final learning accuracy, operationalized as the percentage of correctly identified object

locations, was significantly higher in young adults than in children, $F_{(1,79)} = 94.31$, $p < .001$,

$w^2 = .53$, $t_{(185)} = 7.55$, $p < .001$ (Fig. 2B) , as revealed by the LME model. There was no

significant effect of *Session* (p = .79) and no *Session x Group* interaction (p = .96), indicating

that the learning accuracy was stable across sessions with different stimuli sets. Although the

learning procedure was adaptive, children showed consistently lower learning performance

compared to young adults.

Figure 2

(A) Overview of Learning Performance. Individual learning trajectories across up to four encoding-retrieval cycles for children and young adults on Day 0, Day 1 and Day 14. Each coloured dot represents a participant's accuracy (percentage of correct responses) at a given cycle. Transparent connecting lines illustrate within-person changes in accuracy across cycles. Across all sessions, children needed on average between two to four learning-retrieval cycles to reach the criterion of 83% correct responses, while young adults typically reached it within two cycles. (B) Final Learning Performance. Final learning accuracy is calculated as the percentage of correct responses during the last learning cycle for both children and young adults. For each group and session, distributions are visualized using half-eye plots (smoothed density estimates), overlaid with boxplots indicating the median and interquartile range. The shape and spread of density plot reflect individual data variability. Grey dashed line indicates the criteria of 83% correctly learned items.

Memory Retention Across Time

Changes in memory retention were assessed during the retrieval part of the memory task (Fig.

1A "Retrieval (fMRI)"). Participants were cued with the object and were instructed to recall as

vividly as possible the associated scene and the location of the object within the scene during the

fixation window, when no visual input was presented on the screen. The associated scene was

then presented with three choices and participants had to select a rectangle indicating the correct

location of the object in the scene (see Methods for more details).

First, we investigated whether retention rates for recently learned items (initially correctly

encoded on Day 1 and Day 14) differed between sessions in children and adults. The *Session x*

Group interaction was not significant, $F_{(1,75)} = 1.77$, p =.187, $w^2 = .001$, indicating that retention

rate differences across sessions did not vary significantly between groups (see Table S1A).

Based on that, we averaged recent retention rates across sessions within each group for

subsequent analysis.

Second, we examined changes in memory retention rates for items that were initially

correctly learned (i.e., strong initial memories), focusing on group differences in recent and

remote (short- and long-delay) memory retention relative to a 100% baseline (see Fig. 3 and

Table S1B for details). The linear mixed-effects model predicting retrieval accuracy for learned

object-location pairs explained a substantial portion of variance, $R^2 = .77$, 95% CI [.73 – .81].

We observed a significant main effects of *Item Type,* $F_{(3,250)} = 229.18$, p <.001, $w^2 = .73$. Post

hoc comparisons revealed no significant difference between recent memory retention and short-

delay remote memory retention, $b = 1.49$, $t_{(259)} = 1.26$, p = .754. However, recent memory

retention was significantly higher than long-delay remote retention, $b = 21.36$, $t_{(259)} = 17.59$,

p < .001, and short-delay remote retention was significantly higher than long-delay remote

memory retention, $b = 19.88$, $t_{(260)} = 16.16$, p < .00. Further, we observed a significant main

effect of *Group*, $F_{(1,85)} = 55.00$, p <.001, $w^2 = .38$. Post hoc comparisons revealed an overall

lower memory retention in children compared to young adults, $b = -11.1$, $t_{(91)} = -7.20$, p < .001.

Additionally, we observed a significant *Item Type* x *Group* interaction, $F_{(3,250)} = 17.35$, p < .001,

$w^2 = .16$. Model-based Sidak post hoc comparisons showed that the slope of memory retention

decline was significantly steeper in children compared to adults for recent items, $b = 15.26$, $t_{(254)}$

= 6.56, p < .001, for short delay remote items, $b = 11.41$, $t_{(255)} = 4.84$, p < .001, and for long

delay remote items, $b = 13.08$, $t_{(258)} = 5.38$, p < .001. In addition, memory retention rates

significantly increased (corrected for multiple comparisons with False Discovery Rate (FDR))
with age in the child group for recent items, b = .89, t = 2.62, p = .016, for short delay remote
items, b = .91, t = 2.67, p = .016, but not for long delay remote items, b = .15, t = .326, p = .747.
Of note, we conducted an additional analysis on a subsample that included only
participants who needed two learning cycles to reach the learning criterion (see Table S1.1, Fig.
S1, Table S1.2 for details). Twenty-one child participants were excluded, resulting in the final
subsample of n = 28 children. The results from this subsample fully replicated the findings from
the full sample, indicating that the amount of re-exposure to stimuli during encoding did not
affect consolidation-related changes in memory retrieval at the behavioral level.
Figure 3

353
Retention rates for initially correctly learned items. Memory accuracy is operationalized as the percentage of
correct responses in the retrieval task conducted during the MRI scanning sessions for items that were initially
correctly learned, indicating strong initial memories. Memory accuracy for recently consolidated items did not differ
between sessions in young adults and children and was collapsed across sessions. Overall, young adults show higher
and more stable memory accuracy than children, with memory declining over time for both groups, particularly for
long delay. All tests used Sidak correction for multiple comparisons. *p < .05; **p < .01; ***p < .001(significant
difference); non-significant differences were not specifically highlighted. The boxplot summarizes the distribution
of accuracy scores across sessions and delay conditions. In each boxplot, the central line indicates the median, the
box represents the interquartile range (25th - 75th percentile), and the whiskers extend to the range of values within
1.5 times the variability. The red dashed line at 34% indicates the threshold for chance-level performance.
364
Taken together, both age groups showed a decline in memory performance over time.
However, compared to young adults, children showed a steeper slope of memory decline for both
immediate recent and remote short- and long-delay memories. In sum, the results showed that

children had overall worse memory retention rates compared young adults, indicating less robust

memory consolidation in children.

fMRI Results

372   *Mean activation for remote > recent memory in ROIs.*

To investigate how neural activation for correctly recalled memories varied across different time

delays, we examined the contrast of **remote > recent** correct trials during object presentation at

retrieval (Fig. 4 "Retrieval fMRI).

We first tested whether the **remote > recent** contrast significantly differed from zero in

each age group and session (Day 1 and Day 14), as an indicator of differential engagement

during memory retrieval. FDR-adjusted results showed no significant results in the anterior and

posterior HC (Fig. 4A), anterior PHG (Fig. 4B), and RSC (Fig. 4G) across sessions and age

groups (all $p > .054$; see Table S6 for details). To rule out the possibility that these non-

significant differences reflect an overall absence of retrieval-related activation, we tested whether

mean activation for recent and remote items – each relative to the implicit baseline – was

significantly above zero. FDR-adjusted results revealed that activation in these ROIs was

significantly greater than zero (all $p < .031$), except in the recent Day 1 condition in children for

the posterior HC ($p > .141$) and the precuneus ($p > .056$, see Table S7 and Fig. S3 for details).

These findings indicate that the anterior and posterior HC, anterior PHG, and RSC are similarly

engaged during successful retrieval of both recent and remote memories, regardless of delay or

age group[1]. Other ROIs showed more differentiated patterns, which are discussed below[2].

To further explore the more differentiated patterns observed in other ROIs, we examined

changes in the **remote > recent** contrast across age groups and sessions (Day 1 and Day 14)

using LME models, controlling for sex, handedness, general intelligence, and mean reaction

---

[1] As a control analysis, we tested whether the anterior and posterior HC, anterior PHG, RSC were similarly engaged during retrieval of recent and remote items over time using the LME models. These models included mean activation relative to the implicit baseline, a Session x Delay x Group interaction, and Subject as a random intercept. The results were consistent with the earlier findings, showing no significant main effect of Delay (all $p > .106$), Group (all $p > .060$), or Session x Delay interaction (all $p > .340$), indicating comparable engagement of these ROIs across delays and age groups (see Table S8 for full statistical details).

[2] In contrast, the vlPFC, CE, posterior PHG and LOC, precuneus, and mPFC showed a significant main effect of *Delay* (all $p < .009$, see Table S8 for details), indicating time-related changes in the **remote > recent** contrast. These effects are examined in more detail below. Notably, these findings are consistent with results from the whole-brain analyses (Tables S3.1, 4.1, 5.1).

time. All main and interaction effects were FDR-adjusted, and all post hoc tests were Sidak-

corrected (see Table S9 for details).

For the **posterior PHG** (Fig. 4B), a significant *Session x Group* interaction, $F_{(1,83)} = 9.54$,

$p = .020$, $w^2 = .09$, indicated a more pronounced increase in **remote > recent** mean signal

difference over time in young adults compared to children, $b = .11$, $t_{(83)} = 3.09$, $p = .003$.

Similarly, also for **the cerebellum** (Fig. 4C) a significant *Session x Group* interaction,

$F_{(1,161)} = 7.68$, $p = .020$, $w^2 = .04$, indicated a stronger increase in **remote > recent** mean signal

difference over time in young adults compared to children, $b = .09$, $t_{(160)} = 2.77$, $p = .006$.

For **the mPFC** (Fig. 4D), a significant main effect of *Group*, $F_{(1,86)} = 7.61$, $p = .023$,

$w^2 = .07$, denoted that the overall **remote > recent** mean signal difference in children was higher

than in young adults, $b = -.10$, $t_{(86)} = -2.76$, $p = .007$.

For **the vlPFC** (Fig. 4E), a significant main effect of *Group*, $F_{(1,82)} = 31.35$, $p = <.001$,

$w^2 = .13$, indicated an overall lower **remote > recent** mean signal difference in children

compared to young adults, $b = -.125$, $t_{(108)} = -3.91$, $p < .001$. In addition, a significant main effect

of *Session*,

$F_{(1,99)} = 10.68$, $p = .005$, $w^2 = .09$, pointed out overall higher **remote > recent** mean signal

difference on Day 14 compared to Day 1, $b = .08$, $t_{(99)} = 3.27$, $p = .001$.

For **the precuneus** (Fig. 4F), a significant main effect of *Group*, $F_{(1,161)} = 5.09$, $p = .027$,

$w^2 = .02$, indicated an overall lower **remote > recent** mean signal difference in adults compared

to children, $b = -.05$, $t_{(160)} = -2.26$, $p = .037$. In addition, a significant main effect of *Session*,

$F_{(1,161)} = 6.50$, $p = .036$, $w^2 = .03$, denoted an overall lower **remote > recent** contrast for Day 14

compared to Day 1, $b = -.05$, $t_{(160)} = -2.55$, $p = .012$. Although the **remote > recent** contrasts

were mostly negative, the mean activation for recent and remote items — each relative to the

implicit baseline — was significantly greater than zero for all delays and group (all $p < .023$),

except for children's recent items on Day 1 ($p = .056$; see Table S7 for details).

For the **LOC** (Fig. 4H), a significant main effect of *Group*, $F_{(1,82)} = 9.12$, $p = .015$,

$w^2 = .09$, indicated a higher **remote > recent** mean signal difference in young adults compared

to children, $b = .07$, $t_{(82)} = 3.02$, $p = .003$. Additionally, a significant main effect of *Session*,

$F_{(1,97)} = 16.76$, $p = <.001$, $w^2 = .14$, showed an overall increase in **remote > recent** mean signal

difference on Day 14 compared Day 1, $b = .07$, $t_{(97)} = 4.10$, $p = <.001$. Furthermore, a significant

Session x Group interaction, $F_{(1,81)} = 6.42$, $p = .032$, $w^2 = .06$, demonstrated higher increase in

remote > recent mean signal difference over time in adults compared to children, b = .09,

t(81) = 2.53, p = .013.

Of note, we conducted an additional univariate analysis using a subsample that included

only participants who needed two learning cycles to reach the learning criteria (see Table S9.1

for details). The subsampled results fully replicated the findings from the full sample and

demonstrated that the amount of re-exposure to stimuli during encoding did not affect

consolidation-related changes in memory retrieval at the neural level.

In summary, our findings revealed distinct consolidation-related neural upregulation for

remote memory between children and adults. From Day 1 to Day 14, adults showed higher

increase in **remote > recent** signal difference for remembered items in the posterior PHG, LOC,

and cerebellum than children. Adults showed overall higher remote > recent difference in the

vlPFC than children, while children showed overall higher remote > recent difference in the

mPFC than adults. Furthermore, we observed a constant activation of anterior and posterior HC,

anterior PHG, and RSC in memory retrieval across age groups irrespective of memory type or

delay.

446   **Figure 4**

Mean Signal Differences Between Correct Remote and Recent Memories.

The figure presents mean signal difference for remote > recent contrast across sessions and groups during the object
presentation time window in **(A)** Anterior and Posterior Hippocampus; **(B)** Anterior and Posterior Parahippocampal
Gyrus; **(C)** Cerebellum; **(D)** Medial Prefrontal Cortex; **(E)** Ventrolateral Prefrontal Cortex; **(F)** Precuneus; **(G)**
Retrosplenial Cortex; **(H)** Lateral Occipital Cortex. *Note:* Bars indicate the group mean for each session (solid lines
for Day 1, dashed lines for Day 14), plotted separately for children and young adults. Error bars represent ±1
standard error of the mean. The colour indicated the age groups: purple for children and khaki yellow for young
adults. Across all panels, mean of individual subject data are shown with transparent points. The connecting faint
lines reflect within-subject differences across sessions. Orange asterisks denote significant difference of **remote >**
recent contrast from zero. An upward orange arrow indicates that this difference is greater than zero, while a
downward arrow indicates that this is less than zero. *$p < .05$; **$p < .01$; ***$p < .001$(significant difference); non-
significant differences were not specifically highlighted. Significant main and interaction effects are highlighted by
the corresponding asterisks. All main and interaction p-values were FDR-adjusted for multiple comparisons.
465
466

Neural-behavioural Correlation

We further investigated whether neural upregulation (i.e., remote > recent univariate signal

difference) is related to memory performance. Specifically, considering all ROIs simultaneously

and differential directionality of remote > recent signal differences, we investigated whether any

specific profile of ROI constellation of neural upregulation is related to variations in memory

performance. For this purpose, we employed the partial least square correlation analysis (PLSC;

Abdi, 2010; Abdi & Williams, 2013). With regard to the interconnectedness of the predefined

ROIs, the PLSC is a well-suited method to address multivariate associations between neural

measures and memory measures. Consequently, latent variables that represent differential

profiles of ROI's neural upregulations with robust relation with either short- or long-delay

variations in memory performance were extracted (for more detailed description of the PLSC

method, refer to Method section). In addition, we derived for each subject a value that denotes a

within-person robust expression of either short- or long-delay brain profile.

For each delay, the permutation test of significance resulted in a single latent variable that

reliably and optimally represents across age groups (i) the associations of short delay ROI neural

upregulations with variations in short-delay memory accuracy (Fig. 5A; r = .536, p = .0026); and

(ii) the associations of long delay ROI neural upregulations with variations in long-delay

memory accuracy (Fig. 5B; r = .542, p = .0024). With further bootstrapping, we identified Z-

scores estimates of robustness (larger/smaller than ± _1.96 (a < 0.05)) of the components within

the multivariate brain profiles across all participants. Thus, for short delay, we observed that

higher memory accuracy was robustly associated with greater neural upregulations in the anterior

PHG (Z-score = 2.161, r = .347) and vlPFC (Z-score = 3.457, r = .640), as well as with lesser

neural upregulation in precuneus (Z-score = -2.133, r = -.323) and cerebellum (Z-score = -2.166, r = -.371) across age groups. In contrast, for long delay, we observed that higher memory accuracy was robustly associated with greater neural upregulation in the vlPFC (Z-score = 3.702, r = .492), RSC (Z-score = 4.048, r = .524), and LOC (Z-score = 3.568, r = .455), and with lesser neural upregulation in mPFC (Z-score = -2.958, r = -.394) across age groups. The identified latent variables indicate that substantial amount of variance (short delay: r = .536 and long delay: r = .542) in either short- or long-delay memory performance was accounted by the identified differential functional profiles of brain regions.

Figure 5

Multivariate short- and long-delay brain profiles of neural upregulation (remote versus recent neural activation differences) are associated with variations in memory accuracy. A) *Short Delay Brain Profile*. Latent

variables weights or saliences for each ROI build up one latent variable that expresses a composite short-delay brain
profile across both age groups . B) *Long Delay Brain Profile*. Latent variables weights or saliences for each ROI
build up one latent variable that expresses a composite long-delay brain profile across both age groups. The bar plot
shows the bootstrap ration (BSR) values for the latent variable, reflecting the stability of the relationship between
brain activation and memory performance. Stability of salience elements is defined by Z-scores (depicted as red line:
a value larger/smaller than ± _1.96 is treated as reliably robust at (a < .05). C) Short Delay Brain Scores by Group.
D) Long Delay Brain Scores by Group. Each box represents the distribution of brain scores within a group, with
central lines indicating the median and boxes showing the interquartile range. Whiskers represent the full range of
non-outlier values. *Note:* PHGa – anterior parahippocampal gyrus; PHGp – posterior parahippocampal gyrus; HCa –
anterior hippocampus ; HCp – posterior hippocampus; PC– precuneus; vlPFC – ventrolateral prefrontal cortex;
mPFC – medial prefrontal cortex; RSC – retrosplenial cortex; LOC – lateral occipital cortex; CE – cerebellum; r –
Spearman's rank order correlation index.
513

Identified brain profiles across groups suggest shared patterns between neural mean signal

differences in differential sets of ROIs and memory accuracy are consistent across children and

adults. As this approach optimizes for consistent covariance pattern across the full sample, it

does not test for group-specific profiles. When conducting the same PLS models within each

group, no stable latent profile emerged (all p > .069). The reduced within-group sample may

have affected the bootstrap-based stability. To address this, we explored whether groups differ in

their expression of the common LV (i.e., brain scores). This analysis revealed that children

showed significantly lower brain scores than adults both in short delay, $t_{(83)}$ = -4.227, p = .0001

(Fig. 5C), and long delay, $t_{(74)}$ = -5.653, p < .001 (Fig. 5D), suggesting that while the brain-

behaviour profile was shared, its expression varied by group.

Taken together, differential short- and long-delay brain profiles of neural upregulation

were related to variations in memory accuracy. Despite age-related differences in the derived

brain scores, higher expression of within-participant brain score was associated with higher

memory retention rates in short and long delay similarly in children and young adults.

529   *Representational similarity results.*

In addition to distinct univariate neural upregulation for recent and remote memories, children

and adults may exhibit differences in neural representations of these memories. Over time, these

representations could also undergo consolidation-related transformations. To address this further,

we investigated both more differentiated detailed scene-specific reinstatement and more generic

category-based neural representations in children and adult.

3.2.2.1 Corrected scene-specific reinstatement.

To measure how scene-specific reinstatement at retrieval during fixation time window (after

short cue by object presentation; see Fig. 6A) changes over time as memories decay, we

computed a *scene-specific reinstatement index* for each neural representational similarity matrix.

We hypothesized that neural patterns evoked by reinstatement of a specific scene without any

visual input during fixation time window would be similar to neural patterns evoked by actual

presentation of the scene during the scene time window. Therefore, the scene time window was

used as a template against which the fixation period can be compared to. Participants were

explicitly instructed to recall and visualize the scene and location of the object during fixation

time window after being cued by the object. Since the locations were contextually bound to the

scene and each object had a unique location in each scene, the location of the object was always

embedded in the specific scene context.

To investigate how scene-specific reinstatement changes over time with memory

consolidation, all analyses were restricted to correctly remembered items. For each specific

scene, the correlation between neural patterns during fixation "*fixation period*" and neural

patterns when viewing the scene "*scene period*" was conducted (Fisher-transformed Pearson's r;

Fig. 6B). A *set-based reinstatement index* was calculated as an average distance between

"*fixation*" and "*scene*" period for a scene and every other scene within the stimuli set (Deng et

al., 2021; Ritchey et al., 2013; Wing et al., 2015). The set-based reinstatement index reflects the

baseline level of non-specific neural activation patterns during reinstatement. We then calculated

the *corrected scene-specific reinstatement index* as the difference between scene-specific and

set-based Fisher-transformed Pearson's r (Deng et al., 2021; Ritchey et al., 2013; Wing et al.,

2015). Given the temporal proximity of the fixation and scene time window, we refrain from

interpreting the absolute values of the observed scene-specific reinstatement index. However,

given that the retrieval procedure is the same over time and presumably similarly influenced by

the temporal autocorrelations, we focus primarily on the changes in reinstatement index for

correctly retrieved memories across immediate, short, and long delays. In other words, the focus

in the following analysis lies on the time-related change in the scene-specific reinstatement

index.

First, scene-specific reinstatement indices for recent items — tested on different days —

did not significantly differ, as indicated by non-significant main effects of *Session* (all p > .323)

and *Session x ROI* interactions (all p > .817) in either age group. This indicates that temporal

autocorrelation was consistent across scanning sessions. Based on that, we averaged the scene-

specific reinstatement indices for recent items across sessions. To investigate time-dependent

change in scene-specific reinstatement in children and young adults in the predefined ROIs, we

Figure 6

Scene-Specific Reinstatement

(A) Index Computation (Scene). A representational similarity index was computed by calculating the average
similarity between activation patterns in the fixation and scene time windows, separately for recent scenes, remote
scenes on Day 1, and remote scenes on Day 14. **(B) Scene-Specific Reinstatement**. A corrected scene-specific
reinstatement index was computed by assessing the average similarity within-trial similarity between the fixation
and scene time windows and subtracting the average between-trial (set-based) similarity across all other trials. This
controls for baseline similarity unrelated to specific scene content. S – scene time window; F – fixation time
window; r – Pearson's correlation index; Δz – difference between two Fisher transformed r values. * - Activation
patterns.

conducted LME models, with delay (recent, remote short and remote long delays), group

(children and young adults) for each ROI, controlling for ROI BOLD activation (Varga et al.,

2023) during corresponding sessions. All main and interaction effects were FDR-adjusted, and

all post hoc tests were Sidak-corrected for multiple comparisons.

Generally, in all predefined ROIs, we observed a significant main effect of *Session* (all p <

.001) and a significant effect of *Group* (all p <.004, Fig.7A-G), except for the LOC (p = .271,

Fig. 7J). The pattern of time-related decline was similar across age groups, as indicated by not

significant *Session x Group* interactions in all ROIs (all p > .159). There was no significant effect

of *BOLD activation* (all p > .136). The full statistical report on the LME-model is in

Supplementary Material in Table S10. A more detailed overview of the observed main effects

and their Sidak-corrected post-hoc tests are summarized in the Table 2.

Figure 7

Corrected scene-specific neural reinstatement.

Scene-specific neural reinstatement defined as the difference between Fisher-transformed scene-specific and set-
specific representational similarity. Scene-specific neural reinstatement index by group (children vs. adults) and
session (Day 0 – recent, Day 1 – remote short delay, Day 14 – remote long delay). Bars represent the mean
reinstatement index for each session within each group, with error bars indicating standard error of the mean.

Transparent dors show individual participant data points, jittered horizontally for visibility. The x-axis is grouped by
group and displays (**A**) Hippocampus Anterior; (**B**) Hippocampus Posterior; (**C**) Parahippocampal Gyrus Anterior;
(**D**) Parahippocampal Gyrus Posterior; (**E**) Cerebellum; (**F**) Medial Prefrontal Cortex; (**G**) Ventrolateral Prefrontal
Cortex; (**H**) Precuneus; (**I**) Retrosplenial Cortex; (**J**) Lateral Occipital Cortex. *$p < .05$; **$p < .01$;
***$p < .001$(significant difference). Error bars indicate standard error.
608

609 Table 2

610 *Statistical overview of LME-model based Sidak corrected post hoc comparisons for scene-specific*

611 *reinstatement analysis (based on LME-model described in Table S10).*

	Model-based post hoc comparisons*								
	YC > YA			Recent > Remote Day1			Remote Day 1 > Day 14		
ROI	*b*	$t_{(DF)}$	*p*	*b*	$t_{(DF)}$	*p*	*b*	$t_{(DF)}$	*p*
HCa	-.071	$-5.15_{(89)}$	**<.001**	.040	$4.35_{(162)}$	**<.001**	.095	$9.60_{(167)}$	**<.001**
HCp	-.068	$-5.14_{(91)}$	**<.001**	.040	$4.29_{(162)}$	**<.001**	.094	$9.45_{(168)}$	**<.001**
PHGa	-.069	$-4.75_{(90)}$	**<.001**	.039	$4.05_{(162)}$	**<.001**	.098	$9.62_{(167)}$	**<.001**
PHGp	-.055	$-3.91_{(90)}$	**<.001**	.040	$3.77_{(178)}$	**<.001**	.096	$9.07_{(172)}$	**<.001**
mPFC	-.049	$-2.94_{(92)}$	**.004**	.045	$4.16_{(162)}$	**<.001**	.093	$7.91_{(169)}$	**<.001**
vlPFC	-.058	$-3.84_{(93)}$	**<.001**	.053	$4.55_{(179)}$	**<.001**	.089	$7.79_{(169)}$	**<.001**
CE	-.044	$-3.05_{(89)}$	**.003**	.046	$3.97_{(166)}$	**<.001**	.086	$7.19_{(170)}$	**<.001**
RSC	-.041	$-2.99_{(90)}$	**.003**	.039	$3.72_{(162)}$	**<.001**	.094	$8.56_{(169)}$	**<.001**
PC	-.047	$-3.33_{(89)}$	**.001**	.044	$4.15_{(165)}$	**<.001**	.086	$7.89_{(168)}$	**<.001**
LOC	-.017	$-1.09_{(103)}$	.279	.045	$3.97_{(173)}$	**<.001**	.083	$7.07_{(174)}$	**<.001**

Notes. Degrees of freedom were adjusted based on Kenward-Roger methods. P-values were adjusted based on Sidak adjustment.
YA – young adults; CH – children; ROI – region of interest; HCa – anterior hippocampus ; HCp – posterior hippocampus;
PHGa – anterior parahippocampal gyrus; PHGp – posterior parahippocampal gyrus; mPFC – medial prefrontal
cortex; vlPFC – ventrolateral prefrontal cortex; CE – cerebellum; RSC – retrosplenial cortex; PC– precuneus; LOC
– lateral occipital cortex; b – Beta values; t – t-value; DF – degrees of freedom; p – p-value; CI – confidence interval; *p < .05;
** < .01, *** < .001 (significant difference).
618

To ensure that the observed scene-specific reinstatement effects were not driven by general

signal properties or artefacts unrelated to memory retrieval, we conducted several control

analyses.

First, we repeated the reinstatement analysis using the *"object period"* instead of the

"scene period". The rationale was that the object and the reinstated scene during fixation period

are expected to rely on distinct neural representations. In line with this, we did not expect a

delay-related decline in reinstatement. The derived object-specific similarity index, which is also

subject to temporal autocorrelation, showed no significant effect of Session or Delay in any ROI

(all p > .059; see Table S10.1, S10.2), supporting the specificity of the original reinstatement

effect.

Second, we tested whether the observed group and delay effects might reflect global or

non-specific BOLD signal fluctuations by analysing three control regions within the corpus

callosum (genu, body, and splenium), where no memory-related reinstatement is expected. The

LME models revealed no significant Group effects in any of the white matter ROIs (all p > .426),

indicating no difference between children and adults. Although we observed significant main

effects of Session (all p < .001), post hoc comparisons showed that these effects were driven by

differences between the recent (Day 0) and most remote (Day 14) sessions. Crucially, the key

contrasts of interest — recent vs. Day 1 remote and Day 1 remote vs. Day 14 remote — were not

significant (all p > .080; see Table S10.3, S10.4), in contrast to the robust decline observed in

key ROIs for scene-specific reinstatement.

Finally, we assessed whether the observed reinstatement effects were specific to successful

memory retrieval by examining item-based reinstatement for incorrectly remembered trials. This

analysis revealed no session-related decline in any ROI, further supporting the interpretation that

the reinstatement effects observed in correctly remembered trials are memory-related rather than

driven by unspecific signal changes (see Fig S5).

Taken together, scene-specific reinstatement declined significantly for overnight compared

to immediate memories and declined further after a 2-week delay across all ROIs. These results

indicate that the main decrease in scene-specific neural reinstatement for successfully

consolidated memories occurs already after a short overnight delay and continues with further

decline after a longer, fortnight delay.

Gist-like neural representation*s.*

Another way to evaluate the quality of neural representations during the post-cue fixation time

window is to examine potential shifts in the ongoing balance between differentiated detailed

("verbatim") and generalized generic ("gist") memory, as described by the Fuzzy Trace Theory

(Brainerd & Reyna, 2002). Although our associative memory paradigm was designed to foster

precise, detailed retrieval, it inherently also permits more generic, gist-like retrieval – for

example, some participants may recall "a field" without its unique details (yielding a generic

field representation), whereas others reinstate the full, specific features of the original scene.

Accordingly, to quantify gist-like representations of the scenes sharing the same category (e.g.,

field, forest, etc.) during the fixation time window following the object cueing (see Fig. 1A

Retrieval; Fig. 8), we computed a gist-like representation index.

First, a within-category similarity indices were computed by correlating the multivoxel

patterns during the fixation time window for all correctly remembered scene pairs from the same

category (i.e., field, water, housing, forest, infrastructure, indoor, farming), excluding self-

correlations. Category exemplars were evenly and randomly distributed across runs, preventing

clusters of temporally adjacent trials. By including only correctly recalled trials and building

representational similarity matrices from both within-run and cross-run scene pairs, we

substantially increased the number of independent pairwise comparisons in the representational

similarity analysis – and hence our sensitivity to detect effects. Next, a between-category

similarity indices were computed in the same way, but for scene pairs drawn from different

categories. Finally, a gist-like representation index was defined as the difference between

Fischer-transformed within- and between-category correlations (i.e., [within category$_{recent}$ r –

between category$_{recent}$ r] and [within category$_{remote}$ r – between category$_{remote}$ r] for each session,

Fig. 8). Thus, the gist-like representation index reflects the extent to which neural patterns during

the fixation window reactivate a generalized category representation (i.e., forest) – over and

above any nonspecific similarity to scenes from other categories.

The non-zero values in this index reflect gist-like representation, as the similarity

distance would be higher for pairs of trials within the same category, indicating more generic

representation (e.g., during representation of scenes belonging to a category "forest", participants

may tend to recall a generic image of some forest without any specific details). In other words,

the representation of a more generic, gist-like image of a forest across multiple trials should yield

more similar neural activation patterns. Not significant gist-like representation would indicate

that even within the same category, representation of specific scenes is sufficiently differential

and rich in details, rendering them dissimilar (e.g., participants may tend to recall detailed image

of forests: fall forest with yellow trees, dark pike-tree forest, light summer forest with young

birch trees, etc.).

Figure 8

A Index Computation (Gist)

Task: recollect scene associated with object

B Gist-Like Reistatement

Gist-like Reinstatement (Δz) = Fisher's z (Pearson's r $_{within-category}$) – Fisher's z (Pearson's r $_{between-categories}$)

Representational Similarity Analysis.

(A) Index Computation (Gist). A representational similarity index was computed by assessing the average similarity for fixation time window for within-category and between-category scenes separately for recent, remote (Day 1), and remote (Day 14) scenes based on both within-run and cross-run comparisons. The diagonal (similarity of fixation time window with itself) was excluded from the analysis. **(B) Gist-like Representation.** A gist-like representation index was computed by assessing the average similarity in fixation time window for the same-category pairs and subtracting from it the any-other-category pairs. S – scene time window; F – fixation time window; r – Pearson's correlation index. Δz – difference between two Fisher transformed r values.

First, we aggregated the gist-like representation indices for recent items on Day 1 and Day 14, as there were no significant differences between sessions in ROIs (all p > .231). Then we applied a one-sample permutation t-test to test for significance of all gist-like indices against zero in each ROI (for full overview see Table S12). FDR-corrected values revealed that young adults showed only remote Day 1 gist-like representation in LOC (p = .024), while significant remote gist-like representation was observed in children on Day 1 in precuneus (p = .044) and LOC (p = .024), and on Day 14 in the mPFC (p = .013) and vlPFC (p = .007). Following this, we further analysed group differences separately for each ROI that showed significant gist-like representation, controlling for the BOLD mean activation in each ROI during corresponding sessions.

Figure 9

Gist-like Representations

Bar plots show mean gist-like representation index (difference in Fisher's z-transformed (Δ z) similarity: *within-*
category — between-category) in each group (Children, Young Adults) and session (Day 0, Day 1, Day 14),
computed from combined within- and cross-run comparisons. Error bars indicate ±1 standard error of the mean. A
representation value above zero (denoted by red asterisks) reflect greater neural pattern similarity during fixation
time window between item from the same category than across categories. Bar positions are grouped by age group
(x-axis). Session-specific estimates (Day 0, 1, 14) are differentiated by line of bar border. (**A**) Medial Prefrontal
Cortex; (**B**) Ventrolateral Prefrontal Cortex; (**C**) Lateral Occipital Cortex (**D**) Precuneus; *$p < .05$; **$p < .01$;
***$p < .001$ (significant difference; FDR corrected for multiple comparisons); non-significant difference was not
specifically highlighted.

Second, we investigated the time-dependent change in gist-like representation in ROIs that

showed gist-like representation. For the mPFC (Fig. 9A), we observed a significant main effect

of *Group*, $F_{(1,244)} = 6.55$, $p = .011$, $\omega^2 = .02$, indicating significantly higher gist-like

representation in the mPFC in children compared to young adult, $b = .011$, $t_{(82)} = 2.52$, $p = .013$.

Additionally, a significant main effect of *Session*, $F_{(1,244)} = 3.89$, $p = .022$, $\omega^2 = .02$, indicated

higher remote Day 14 compared to remote Day 1 gist-like representation, $b = .014$, $t_{(180)} = 2.64$,

$p = .027$. For the vlPFC (Fig. 9B), we observed a significant effect of *Session*, $F_{(1,174)} = 4.45$,

$p = .013$, $\omega^2 = .04$, indicating higher remote Day 14 gist-like representation compared to recent

one, $b = .013$, $t_{(195)} = 2.91$, $p = .012$. A significant *Session x Group* interaction, $F_{(1,167)} = 3.04$,

$p = .05$, $\omega^2 = .02$, highlighting significantly higher remote Day 14 gist-like representation in

children compared to young adults, b = .017, $t_{(249)}$ = 2.52, p = .037. Neither LOC nor precuneus

showed any significant main or interaction effects (all p > .062; Fig. 9C and 9D). Taken together,

only the child group showed gist-like representation in the medial and ventrolateral prefrontal

brain regions that was significantly higher during retrieval of long delay remote memories,

indicating a reorganization of memory representations in children.

740   *Neural-behavioural Correlations*

741   Further, we also explored whether over time, recent, short- and long-delay scene-specific

742   reinstatement and gist-like representations are beneficial or detrimental for memory performance

743   by correlating the indices with memory retention rates. We derived, with a PLSC analysis, latent

744   brain pattern across implicated ROIs that share the most variance with delay-related variations in

745   memory accuracy.

746   *Neural-behavioural correlations (scene-specific reinstatement)*

747   For the scene-specific reinstatement all predefined ROIs in both age groups were included.

748   With further bootstrapping we identified Z-scores estimates of robustness (larger/smaller

749   than ± _1.96 (a < 0.05)) of the components within the multivariate brain profile.

750   First, for recent delay (30 minutes after learning), the permutation test of significance

resulted in a single latent variable that robustly represents the association of scene-specific

reinstatement brain profile and memory accuracy across both age groups (Fig. 10B, r = .293,

p = .007). Higher recent memory accuracy was robustly associated with greater scene-specific

reinstatement in the anterior PHG (Z-score = 3.010, r = .819), posterior PHG (Z-score = 2.575,

r = .367), anterior HC (Z-score = 2.629, r = .3713), posterior HC (Z-score = 3.009, r = .417), and

precuneus (Z-score = 2.206, r = .318) across age groups.

757   Second, for short delay, the permutation test of significance resulted in a single latent

variable that robustly represents the association of scene-specific reinstatement brain profile and

memory accuracy across both age groups (Fig. 10B, r = .339, p = .0017). Higher memory

accuracy was robustly associated with greater scene-specific reinstatement in the anterior PHG

(Z-score = 2.885, r = .371), posterior PHG (Z-score = 2.597, r = .342), anterior HC (Z-

score = 3.126, r = .399), posterior HC (Z-score = 2.844, r = .375), vlPFC (Z-score = 2.434,

r = .317), mPFC (Z-score = 2.753, r = .333), and LOC (Z-score = 2.176, r = .298) across age

groups.

Third, for long delay, the permutation test of significance resulted in a single latent

variable that robustly represents the association of scene-specific reinstatement brain profile and

memory accuracy across both age groups (Fig. 10C, $r = .455$, $p = <.001$). Higher memory

accuracy was robustly associated with greater scene-specific reinstatement in the anterior PHG

(Z-score = 6.213, $r = .414$), posterior PHG (Z-score = 4.810, $r = .334$), anterior HC (Z-

score = 5.353, $r = .389$), posterior HC (Z-score = 4.707, $r = .354$), precuneus (Z-score = 3.404,

$r = .281$), vlPFC (Z-score = 3.291, $r = .266$), RSC (Z-score = 3.72, $r = .293$), LOC (Z-

score = 3.288, $r = .282$), and cerebellum (Z-score = 3.842, r = .308) across age groups.

Identified brain profiles across groups suggest shared patterns between neural mean signal

differences in differential sets of ROIs and memory accuracy are consistent across children and

adults. As this approach optimizes for consistent covariance pattern across the full sample, it

does not test for group-specific profiles. When conducting the same PLS models within each

group, no stable latent profile emerged (all $p > .069$). The reduced within-group sample may

have affected the bootstrap-based stability. To address this, we explored whether groups differ in

their expression of the common LV (i.e., brain scores). This analysis revealed that children

showed significantly lower brain scores than adults both in immediate delay, $t_{(85)} = -3.971$,

$p = .0001$ (Fig. 10C), in short delay, $t_{(81)} = -2.973$, $p = .004$ (Fig. 10C), and long delay, $t_{(70)} = -$

2.659, $p = .01$ (Fig. 10D), suggesting that while the brain-behaviour profile was shared, its

expression varied by group.

These results underscore the importance of scene-specific reinstatement in positively

contributing to memory performance for detailed associative information both in children and

adult.

Figure 10

Multivariate short- and long-delay brain profiles of scene-specific reinstatement are associated with
variations in memory accuracy. A) *Recent Delay Brain Profile*. Latent variables weights or saliences for each ROI
build up one latent variable that expresses a composite immediate-delay scene-specific reinstatement brain profile.
B) *Short Delay Brain Profile*. Latent variables weights or saliences for each ROI build up one latent variable that
expresses a composite short-delay scene-specific reinstatement brain profile. C) *Long Delay Brain Profile*. Latent
variables weights or saliences for each ROI build up one latent variable that expresses a composite long-delay scene-
specific reinstatement brain profile. Stability of salience elements is defined by Z-scores (depicted as red line: a
value larger/smaller than $\pm$ _1.96 is treated as reliably robust at (a < .05). The bar plot shows the bootstrap ration
(BSR) values for the latent variable, reflecting the stability of the relationship between brain scene-specific neural
reinstatement and memory performance. D) Recent Delay Brain Scores. E) Short Delay Brain Scores. F) Long
Delay Brain Scores. Each box represents the distribution of brain scores within a group, with central lines indicating
the median and boxes showing the interquartile range. Whiskers represent the full range of non-outlier values. *Note:*
PHGa – anterior parahippocampal gyrus; PHGp – posterior parahippocampal gyrus; HCa – anterior hippocampus ;
HCp – posterior hippocampus; PC– precuneus; vlPFC – ventrolateral prefrontal cortex; mPFC – medial prefrontal

cortex; RSC – retrosplenial cortex; LOC – lateral occipital cortex; CE – cerebellum; r – Spearman's rank order
correlation index.
817

818   *Neural-behavioural correlations (gist-like representations)*

For gist-like representations, we included only those ROIs that showed significant gist-like

representations (i.e., for short delay: LOC, and precuneus across both age group; for long delay:

vlPFC and mPFC in child group).

First, for short delay, the permutation test of significance did not result in a single latent

variable that robustly represents the association gist-like representations brain profile and

memory accuracy in children (Fig. 11A, $r = .221$, $p = .065$). Second, for long delay, the

permutation test of significance resulted in a single latent variable that robustly represented the

association of gist-like representations brain profile and memory accuracy in the child group

(Fig. 11B, $r = .516$, $p = .0014$). The higher long delay Day 14 memory accuracy was robustly

associated with lower gist-like representations in the mPFC (Z-score $= -2.396$, $r = -.498$), and

vlPFC (Z-score $= -5.918$, $r = -.876$) in children.

The significant negative association between long delay gist-like representations in both

prefrontal brain regions and memory accuracy observed in children underscores that gist-like

representations was detrimental to memory performance for detailed associative information in

children in long delay. Taken together, more differentiated detail-rich neural reinstatement was

related to better memory retrieval in both children and young adults. On the other hand, more

gist-like neural representations, uniquely found in children, was related to worse memory

retrieval.

Figure 11

Multivariate short- and long-delay brain profiles of gist-like representations are associated with variations in
memory accuracy. A) *Short Delay Brain Profile*. Latent variables weights or saliences for each ROI build up one
latent variable that expresses a composite short-delay gist-like representations brain profile across age groups. B)
Long Delay Brain Profile. Latent variables weights or saliences for each ROI build up one latent variable that
expresses a composite long-delay gist-like representations brain profile in child group. The bar plot shows the
bootstrap ration (BSR) values for the latent variable, reflecting the stability of the relationship between brain gist-
like neural representations and memory performance. Stability of salience elements is defined by Z-scores (depicted
as red line: a value larger/smaller than ± 1.96 is treated as reliably robust at (a < .05). *Note:* vlPFC – ventrolateral
prefrontal cortex; mPFC – medial prefrontal cortex; LOC – lateral occipital cortex; r – Spearman's rank order
correlation index.

DISCUSSION

In the present study, we investigated system-level memory consolidation of object-location

associations across three delays: immediately after learning, after one night of sleep (short delay)

and after two weeks (long delay). We tracked changes in neural activation and multivariate

activation patterns over time, comparing 5-to-7-year-old children and young adults. Our main

findings are as follows: (i) Children showed greater decline in memory retention compared to

young adults at both short and long delays. (ii) Regarding neural upregulation – reflected as the

mean difference in activation between remote and recent retrieval – the two age groups showed

distinct changes over time. Young adults exhibited an increase in neural upregulation over time

in the posterior PHG, cerebellum and LOC, and an overall higher neural upregulation in the
vlPFC compared to children. In contrast, only children showed a decrease in neural upregulation
over time in the RSC and overall greater neural upregulation in the mPFC than adults. Distinct
neural upregulation profiles with specific sets of brain regions were related to immediate, short
and long delay memory accuracy. (iii) Using RSA, we found that differentiated scene-specific
reinstatement declined over time in both age group. Notably, more generic gist-like
representations were observed only in children, particularly in medial and ventrolateral prefrontal
regions. Importantly, higher scene-specific reinstatement was related to better memory retention
in both age groups, whereas greater gist-like representations were related to lower memory
retention at the long delay only in children.

Our study extends previous adult-based findings and, for the first time, demonstrates that the
memory retrieval after consolidation in children is accompanied by differential patterns of neural
activation of some of the core retrieval regions, attenuated neural reinstatement of detailed
memories, and the emergence of generic gist-like representations. These findings suggest that
adults leverage mature neural memory systems and extensive existing knowledge base to encode
and consolidate new complex information with detailed accuracy. In contrast, children – whose
neural system is still developing – may rely more on consolidating gist information as a
foundational scaffold for their still sketchy knowledge base, possibly at the cost of episodic
detailedness. At this developmental stage, focusing on precise detail may not be yet prioritized
(Keresztes et al., 2018). Each of these findings is discussed in detail in the following sections.
879

Less robust short and long delay memory retention in children compared to young
adults.

Our findings indicate that preschool children (5 to 7 year old) can encode and retain complex
associative and highly contextualized information successfully over extended periods following
adaptive learning. However, their overall learning and retrieval performance was lower
compared to young adults. Moreover, children exhibited more pronounced declines in memory
retention over both short and long delays for correctly learned information, suggesting less
robust memory consolidation compared to young adults.

Regarding learning, children needed more cycles to memorize object-scene associations
and showed lower learning performance after initial strategic encoding compared to young

adults. Although we did not expect children to reach adult-like learning rates given the complex
and associative nature of the task (Pressley et al., 1981), we aimed to maximize children's
learning capacities through adaptive learning procedures. To support this, attention allocation
and motivation during encoding and learning were closely monitored through the constant
presence of the experimenter and the use of feedback questionnaires. In addition, all participants
underwent training in elaborative encoding strategies to support later retrieval.

Overall, our findings on learning suggests that children were less adept that adults at
utilizing strategic control over encoding, such as creating and maintaining stories to aid their
retrieval. This is consistent with previous literature, showing continuous improvement in
children's ability to use elaborative strategies between ages 4 and 8 (Bjorkund et al., 2009;
Crowley & Siegler, 1999; Pressley, 1982). Additionally, children at this age may experience
difficulties in controlling (Ruggeri et al., 2019) and effectively using their learning strategies
over time (Brod, 2021; Shing et al., 2010). Observed lower learning rates may also be attributed
to less efficient binding processes in children compared to young adults (Shing et al., 2010;
Sluzenski et al., 2006). Although we included only stimuli from the primary school curriculum to
reduce age differences in knowledge availability, ongoing maturation of the memory brain
network in 5-to-7-year-old children may have attenuated their benefit from pre-existing
knowledge and memory aid through strategic elaboration (Ghetti & Bunge, 2012; Lenroot &
Giedd, 2006; Nishimura et al., 2015; Ofen, 2012; Shing et al., 2008). Despite these challenges, 5-
to-7-year-old children were capable of learning complex associative information to a
considerable extent, which aligns with their ability to gradually accumulate world knowledge
(Bauer, 2021; Brod & Shing, 2022; Wagner, 2010).

Concerning memory consolidation, our results are in line with previous studies that
reported worse memory retention for associative information in school age children compared to
adults (Østby et al., 2012; Schommartz et al., 2023, 2024). On the other hand, our results are not
in line with sleep-related beneficial effects on mnemonic performance of 7-to-12-year-old
children after one night delay (Peiffer et al., 2020; Wang et al., 2018) that were shown for novel
stimuli not related to any prior knowledge (in the sense of arbitrary stimuli). As we opted for
well-learned information that should allow for rapid creation of new schemas or integration of
new associations into already existing schemas, our findings indicate that the beneficial role of
sleep on memory consolidation in children compared to adults may not apply for repeatedly and

strategically learned information. Deliberate learning is potentially more advantageous for subsequent memory retention in young adults, as this information may be integrated into pre-existing knowledge structures faster (van Kesteren et al., 2013), with higher strategic control of memories upon retrieval and therefore greater accessibility of consolidated memories (Fandakova et al., 2017; Gaudreau et al., 2001). Taken together, our findings indicate that compared to young adults, 5-to-7-year-old children exhibit less robust memory consolidation for well-learned information, suggesting an overall reduced ability to retain detailed memories in children.

To better understand whether observed age-related differences in memory performance reflect changes or differences in memory strength per se, we conducted exploratory analysis using drift diffusion modelling (DDM; Lerche & Voss, 2019; Palada et al., 2016; Ratcliff et al., 2011, 2012; Ratcliff & McKoon, 2008; Zhou et al., 2021). DDM estimates the underlying cognitive mechanisms driving decision-making by jointly modelling accuracy and reaction time, offering a nuanced index of memory strength via the drift rate parameter. This approach allowed us to quantify trial-wise memory accessibility beyond raw performance measures (see Fig. S1 and section S2.1 in Supplementary Materials for detailed overview). The results revealed that children had significantly lower drift rates compared to young adults across all delays, indicating slower and noisier evidence accumulation – possibly due to weaker memory representations. As drift rate closely correlates with memory accuracy (Ratcliff et al., 2011), our findings on the memory strength align with those on memory accuracy during retrieval in both age groups. Crucially, drift rate decreased systematically from recent to remote conditions in both groups, but this decline was steeper in adults. This finding suggests that while adults started with stronger memory traces, these detailed, differentiated traces were also more susceptible to decay. In contrast, children's already lower drift rates remained relatively more stable over time. This pattern points to qualitative group differences in how memories are initially encoded and subsequently consolidated. The DDM analysis helped us to dissociate group differences in retrieval dynamics from surface-level accuracy. It revealed how response patterns varied as a function of memory strength across time, supporting the conclusion that developmental differences in memory consolidation cannot be fully explained by initial performance alone. Our neural findings suggest that differences in functional engagement of the retrieval network and the

characteristics of memory representations being created and retained may underlie the observed behavioural differences.

Differential upregulation of remote > recent neural activation over time in children in comparison to young adults.

Analyses of neural upregulation (i.e., remote > recent difference in neural activation) over time allowed us to control for the effects of rapid consolidation during repeated learning, while examining changes in short- or long-delay neural activation (Brodt et al., 2016b, 2018; Yu et al., 2022). First, we observed increased upregulation in the vlPFC over time in both age groups, with young adults showing greater vlPFC upregulation overall. Furthermore, we observed stable greater upregulation in the mPFC over time in children. On the one hand, this may indicate a stronger strategic control over retrieval processes in young adult, given the vlPFC's role in strategic remembering and retrieval of stored memories (Badre & D'Esposito, 2009; Kuhl et al., 2012). Such vlPFC upregulation was beneficial for memory retention. On the other hand, the observed higher mPFC upregulation in children may reflect less efficient suppression of the default mode network during effortful memory search (Chai et al., 2014; Fair et al., 2008). Over time, cognitive control during memory retrieval may increase as it requires greater effort to recollect elaborative stories to remember the associated spatial context. Strategic control over memories may be present but less pronounced in children due to the more protracted developmental trajectories of prefrontal cortex maturation (Ghetti & Bunge, 2012c; Gogtay et al., 2004; Ofen, 2012; Shing et al., 2010b).

In addition, our results indicate that a more pronounced schema-related retrieval that may be mediated by mPFC to a greater extent in children than in young adults. This extends previous findings on the involvement of mPFC in structured and schema-related retrieval of long-term memories (Takashima et al., 2006; Yamashita et al., 2009) to a child developmental cohort. Higher mPFC upregulation in long delay was negatively related to memory performance, suggesting that it is detrimental to the retention of detailed associative memories. In addition, it may suggest consolidation-related transformation of memory traces into less differentiated, more generic and gist-like memories (Gilboa & Marlatte, 2017; Gilboa & Moscovitch, 2021).

Second, in other constituents of the recollection network (Ranganath & Ritchey, 2012), we

observed increased upregulation from short to long delay in the posterior PHG and overall lower

upregulation in precuneus (i.e., remote > recent) in young adults. As young adults showed higher

memory retention rates for more detail-rich information, this superior memory may be mediated

by higher upregulation in the posterior PHG involved in contextual associations and scene

memory (Aminoff et al., 2013). In children, PHG undergoes prolonged maturation (Golarai et

al., 2007), and its increased functional maturation is related to long-term scene recollection

(Chai, 2010). In addition, higher mnemonic distinctiveness of more recent memories (i.e., higher

retention rates for detailed information) may also be mediated by RSC and precuneus activation

profiles, as these regions are involved in mnemonic vividness, spatial, and associative memory as

indicated by other findings from immediate delays (Brodt et al., 2016b; Hebscher et al., 2019;

Mitchell et al., 2018; Richter et al., 2016; Tambini & D'Esposito, 2020; Vann et al., 2009).

Moreover, lower precuneus upregulation after a short delay and higher RSC upregulation after a

long delay was related to better memory performance. Time-related decrease in activation of the

posterior brain regions observed in children aligns with previous findings of DeMaster and

Ghetti (2013), who reported that the engagement of parietal regions during recollection of correct

memories increased with age in 8-to-11-year-old children. Therefore, the continuing maturation

of parietal regions in 5-to-7-year-old children (Sowell et al., 2002) presumably contributes to the

age-related differences in consolidation-related upregulation observed in these regions.

Third, the observed increase in neural upregulation from short to long delay in the LOC

and the cerebellum in young adults is also in line with previous findings suggesting that the

cerebellum supports rapid cortical storage of memory traces after repeated exposure – even after

24 hours (Stroukov et al., 2022) – and shows upregulation of neural activation for long-term

episodic memory retrieval (Andreasen et al., 1999). Concerning the LOC, prior studies have

linked HC-LOC activation to scene-related associative memory consolidation (Tambini et al.,

2010) and to human object recognition (Grill-Spector et al., 2001). Moreover, the network

comprising the angular gyrus and LOC has been shown to enhance overnight retention of

schema-related memories in young adults (van der Linden et al., 2017). Consistent with these

findings, we also observed that greater LOC upregulation after a long delay was related to better

memory performance. The more pronounced upregulation from short to long delay in these

regions in adults suggests that the cerebellum and LOC support long-delay memory retention –
one that appears to be functionally immature in middle childhood.

Finally, our findings on age-group and delay-invariant activation in the anterior HC and
PHG, and posterior HC during the retrieval of detail-rich memories (i.e., the exact location of an
object within a scene) are in line with Nadel & Moscovitch (1997),who postulated that the
hippocampal formation and related structures remain involved in detail-rich memories upon their
retrieval, irrespective of memory age. For example, Du et al. (2019) reported stable hippocampal
involvement during retrieval of associative memory across delays of one day, one week and one
month in young adults. Tanrıverdi et al. (2022) also demonstrated that post-encoding
coactivation of hippocampal and cortical brain regions may lead to experience-dependent change
in memories, highlighting the importance of hippocampal involvement during consolidation.
Furthermore, the absence of age-related differences in HC and anterior PHG involvement are
also in line with developmental studies that have reported the relative maturity of the HC in
middle childhood (Keresztes et al., 2017; Lee et al., 2014; Shing et al., 2010b), which is
concomitant with an improvement in the ability to bind event features together into a coherent
representation around the age of six years (Sluzenski et al., 2006). Specifically, our finding on
hippocampal engagement being robust in children and adults extends the Multiple Trace Theory
and the Trace Transformation Theory to a child developmental cohort (Moscovitch & Gilboa,
2022; Nadel et al., 2000). Taken together, the similar engagement of medial-temporal cortex
over time in children and adults indicated that the retrieval of well-learned detail-rich memories
is mediated by these brain structures already in middle childhood.

To summarize, we provide novel evidence about changes in neural upregulation for
successfully consolidated memories over short and long delay, relative to immediately learned
memories. While children exhibited adult-like stable neural activation for recent and remote
memories in medial-temporal brain regions, young adults relied more on prefrontal, occipital,
cerebellar, and parietal brain regions over time, compared to more pronounced reliance on
medial prefrontal regions in children. Adults show more mature neocortical consolidation-related
engagement, resulting in stronger and more durable detailed memories over time while in
children immature neocortical engagement may lead to consequent reduction in memory
retention of detailed memories.

1042

Reduced scene-specific reinstatement over time in children and young adults.

We found that scene-specific reinstatement decreased over time both in children and young
adults, aligning with delay-related decrease in memory retention. Additionally, higher scene-
specific neural reinstatement was related to better memory performance in short and long delay
in both age groups.

Our findings contribute to the memory consolidation literature by demonstrating that
scene-specific neural reinstatement observed in neocortical, medial temporal and cerebellar brain
regions supports reinstatement of detailed specific contextual memories. This observation is
consistent with the Contextual Binding Theory (Yonelinas et al., 2019), which posits that
stronger reinstatement of contextual details can enhance memory retention. The similar decay of
these processes over time in both children and adults suggests that the basic mechanisms of
contextual binding are present early in development. Additionally, in line with the Trace
Transformation Theory (Moscovitch & Gilboa, 2022), our findings suggest that reinstatement
patterns continuously transform over time. This transformation, observed across all considered
memory-related regions, indicates a consistent and systematic consolidation-related reshaping of
the unique scene-specific memory representations over time (Chen et al., 2017).

Our findings on scene-specific reinstatement align with and add to the previous literature
that show reliable reinstatement of unique events. For example, our findings align with the
effects observed by Masís-Obando et al. (2022) for the immediate recall of story details in key
memory regions. Consistent with Oedekoven et al. (2017), our results show that memory
representations for unique events can be reliably detected through scene-specific reinstatement
even after extended delays. Furthermore, we build on Guo & Yang (2022) by demonstrating how
specific ROI-related profiles of neural reinstatement during retrieval correlate with long-term
memory retention. Unlike Oedekoven et al. (2017), who reported no time-related differences in
reinstatement effects and used the same video clips for immediate and delayed recall – which
could have inadvertently reinforced memory through reactivation – our study employed unique
stimulus sets for each retrieval sessions, preventing any reconsolidation of mnemonic
representations. This approach revealed a significant attenuation of reinstatement patterns after
an overnight delay, which further diminished after two weeks, highlighting the importance of
intentional reactivation for maintaining the specificity of neural reinstatement.

Our findings indicate similar patterns of scene-specific neural reinstatement between
children and young adults. Building on Fandakova et al. (2019), who found similar
distinctiveness of neural representations during encoding in 8-to-15-year-old children and adults,
our results suggest that this similarity extends to younger ages, showing comparable
distinctiveness of neural representations for unique memories from middle to late childhood and
early adolescence. Additionally, our research supports the presence of delay-related change in
scene-specific reinstatement in 5-to-7-year-old children, albeit at a lower level compared to
adults, aligning with and extending previous studies (Benear et al., 2022; Cohen et al., 2019;
Golarai et al., 2015), which demonstrated reliable mnemonic reinstatement for visual input (i.e.,
faces, movie clips) in 5-to-11-year-old children. Furthermore, we extend these findings, by
showing that successful of long-term memory retrieval is associated with more differentiated
neural reinstatement in both children and young adults, indicating similar mechanisms of detail-
rich memory consolidation present as early as 5-to-7 year.

Our results indicate that higher scene-specific neural reinstatement over time correlated
with better memory retention in both children and adults. This is in line with the neural fidelity
hypothesis (Xue, 2018), suggesting that more similar neural reinstatement reflect less noisy
representations of mnemonic information. Convergent evidence showed that higher fidelity of
neural representation across study episodes leads to successful memory (Xue et al., 2010, 2013).
Similarly, Masís-Obando et al. (2022) reported that more specific neural representations
predicted subsequent memory performance in young adults.

Of note, our study design, which resulted in temporal autocorrelation in the BOLD signal
between memory retrieval (i.e., fixation time window) and scene observation and response (i.e.,
scene time window), was consistent across all three delay windows. Since the retrieval procedure
remained unchanged over time and was similarly influenced by temporal autocorrelations in
addition to several control analyses, we attribute our RSA findings to differences in
reinstatement between recent and remote trials. Given that the scene time window for the 3AFC
task was constant, the brain signals should exhibit similar perception-based but variably
memory-based patterns across all delays.

Furthermore, all items, regardless of retrieval delay, underwent extensive learning and
showed successful consolidation, as evidenced by correct recall. This suggests that both the
fixation and scene time windows engaged memory-related neural processes. According to Brodt

et al., (2016, 2018), rapid consolidation-related neural reorganization can occur immediately
after learning, indicating that even during recent retrieval, scenes are processed in a memory-
oriented manner. Additionally, during the scene time window, participants engaged in retrieval
by selecting the correct object location within the scene. Thus, while the scene time window
involved perceptual processing, its impact is consistent across all items due to uniform exposure
to repeated learning, making them equally familiar to participants. Although our paradigm per se
cannot arbitrate between perception-based and memory-based nature of retrieval during scene
presentation, our exploratory univariate analysis during the scene presentations time window (see
Figure S6, Table S11 in Supplementary Materials) revealed higher neural engagement in the key
memory regions with passing time, supporting memory-related processing during the scene time
window.

Taken together, our findings provide novel evidence that although children exhibit more
attenuated scene-specific reinstatement compared to young adults, the consolidation-related
decrease in differentiated reinstatement follows similar patterns as in adults. This highlights that
despite less robust memory consolidation and lower memory strength, children's neural
transformations of distinct memories over time may share the same mechanisms as adults, with
scene-specific reinstatement proving beneficial for memory retention in both groups.

1121

Unique Gist-like Representations in Children.

The results showed that only children demonstrated the emergence of generic gist-like
representations in medial and ventrolateral prefrontal brain regions during successful long-delay
retrieval. Furthermore, greater long-delay gist-like representations were associated with poorer
long-delay memory accuracy in children. With these findings, we provide the first neural
empirical evidence to support the Fuzzy Trace Theory (Brainerd & Reyna, 2002; Reyna &
Brainerd, 1995), showing neural reorganization of memory representations in children.

The Fuzzy Trace Theory aims to characterize the shifts in ongoing balance between
precise, detailed "verbatim" memory and more generalize, simplified "gist" memory (Brainerd &
Reyna, 2002) from a developmental perspective. Our associative object-location task allowed the
investigation of these "dichotomy" as it was aimed to cultivate detailed, precise memories for
retrieval. Simultaneously, it enabled generalization by creating of more generic representations
due to the presence of related category-based information. Adults were able to build upon solid

pre-existing knowledge by embellishing them with details and integrating them into these
structures. Children, in contrast, with their sparser knowledge, may have focused more on
solidifying the structure with overlapping information. Aligning with the Fuzzy Trace Theory,
our results suggest that reliance on gist-like memory representations is less effective for long-
term retention of complex associative information compared to detailed verbatim memory, which
seems to be characteristic of adults.

We also observed short-delay gist-like representations in posterior brain regions: the
precuneus in children and the LOC in both age groups. These representations were not directly
related to memory accuracy. The LOC is involved in object and scene recognition (Golarai et al.,
2007; Grill-Spector et al., 2001) and has been shown to participate in schema-related
consolidation or in durable but less specific memories (van der Linden et al., 2017). Its
involvement in gist-like representations across age groups suggests occipital areas already
engage in some degree of categorical abstraction, especially when stimuli share common visual
or contextual features. In contrast, the additional precuneus involvement in children might reflect
broader and less differentiated cortical engagement. The precuneus is involved in mental
imagery, integration of visuospatial and self-referential information, and episodic simulation
(Hebscher et al., 2019; Plachti et al., 2023). In young children, where functional specialization is
still developing, memory representations may be less tightly constrained, leading to more diffuse
activation patterns across associative and imagery-related regions (Plachti et al., 2023). At short
delays, memories may still be relatively strong — as supported by our drift diffusion modelling
showing higher drift rates — thus allowing these emerging gist-like signals to coexist with
detailed memory traces, perhaps as a by-product of early consolidation.

The emergence of long-delay gist-like neural representations in both the mPFC and vlPFC
in children may reflect consolidation-related integration of memory representations into more
abstract, generic forms over time. This aligns with the mPFC's known role in integrating across
memories (Schlichting et al., 2015), the increase in semantically transformed representations for
related information over time in adults (Krenz et al., 2023), and the integration of new
information into schema (Gilboa & Marlatte, 2017; Preston & Eichenbaum, 2013). While gist-
like neural representations may support the generalization of information to bolster the sparse
knowledge structures in children, this occurs at the costs of memory precision (Reyna et al.,
2016). The involvement of the vlPFC in gist-like representations was also stable. The vlPFC has

been implicated in controlled semantic retrieval, selection among competing memory traces, and integration of overlapping information (Badre & Wagner, 2007; Simons & Spiers, 2003). In the context of developing memory systems, children's engagement of the vlPFC may reflect an effort to resolve interference among overlapping scene-object memories by drawing on more abstracted or semantically reduced representations. This aligns with the findings that the vlPFC may support the selection and organization of relevant features, especially under cognitive load or when representations are weak (Bunge et al., 2004; Sanefuji et al., 2011; Trelle et al., 2019). Thus, the coactivation of mPFC and vlPFC in children during long-delay memory retrieval suggests that gist-based retrieval strategies are not only present but possibly compensatory, reflecting an adaptive but less precise means of accessing complex memories.

Importantly, we found that gist-like neural representations in the mPFC and vlPFC at long delay were negatively associated with memory accuracy in children. This suggests that, while gist representations may serve a generalizing function, they are less effective for supporting retrieval of detailed object-location associations. In contrast to our findings, Masís-Obando et al. (2022) demonstrated that more schema-based representations in the mPFC were associated with better subsequent memory performance in adults. However, the study utilized stimuli with clearly differentiable schema and details components, whereas our design required the retention of both contextual details and object associations. It is important to note that in our study, gist-like representations were observed only for correctly remembered items, suggesting that children retained some core aspects of the memory trace, even if details were compromised. This aligns with the idea that gist-based representations preserve the overall meaning or category, but not the specific spatial or contextual bindings necessary for high-fidelity retrieval (Reyna et al., 2016). Thus, the negative correlation may not reflect an entirely detrimental effect, but rather a trade-off between generalization and detail preservation.

Overall, our results are in line with Brainerd et al. (2002), showing that in middle childhood, precise mnemonic representations (i.e., scene-specific reinstatement) and gist-like mnemonic representations can co-exist at the neural level. These findings also extend the adult literature, supporting the notion of qualitative transformations of memory traces, whereby detailed and more schematic, generic memories may simultaneously be present (Chen et al., 2017; St-Laurent & Buchsbaum, 2019; Ye et al., 2020). Building on the postulations from Keresztes et al. (2018) and Ngo et al. (2021), who showed that 5-to-7-year-old children tend to

rely more on generalization, our findings suggest that retaining memories with viewer specific
details may allow for faster integration of overlapping features into emerging knowledge
structures (Bauer, 2021; Gilboa & Marlatte, 2017). In contrast, adults may form strong, highly
detailed memories supported by effective strategic retrieval mechanisms, without the need to
form gist-like representations. Although category-level reinstatement has been documented in
adults (e.g., Kuhl & Chun, 2014; Tompary et al., 2020; Tompary & Davachi, 2017), the absence
of such effects in our adult group may reflect differences in study design, particularly our use of
non-repeated, cross-trial comparisons based on fixation events. It may also reflect different
consolidation strategies, with adults preserving more differentiated or item-specific
representations, while children tend to form more schematic or generalizable representations — a
pattern consistent with our interpretation and supported by prior work (Fandakova et al., 2019b;
Sekeres, Winocur, & Moscovitch, 2018) .

Taken together, our findings provide novel evidence children's memory consolidation is
characterized by a shift toward gist-based representations, supported by mPFC and vlPFC
engagement. While these representations may aid generalization and schema-building, they
appear less effective for detailed retrieval, especially over long delays. With this we provide the
first empirical evidence to support Fuzzy Trace Theory at the level of gist-like neural
representations in children. Future research may build on this approach to further explore
conditions under which schema-based representations enhance memory performance and how
these process differ across development.

1217

LIMITATIONS

Several limitations of the current study should be noted. First, our test for memory was based on
a 3-alternative forced choice procedure, which was intended to reduce the need for strategic
search (e.g., in free recall). As reorganization and stabilization in consolidation depend on the
psychological nature of mnemonic representations (Moscovitch & Gilboa, 2022), future studies
may employ more demanding recall-based memories to characterize memory consolidation more
comprehensively. Particularly, future studies may differentiate mnemonic accessibility vs.
precision (Murray et al., 2015; Richter et al., 2016), as they may show differential temporal
dynamics in the developing brain and involve differential neural mechanisms. Second, as we
included only stimuli congruent with prior knowledge, future studies may introduce knowledge-

incongruent information to investigate the beneficial effect of prior knowledge on memrory

consolidation more directly. Prior knowledge may impact learning and consolidation of

information over time differentially by development (McKenzie & Eichenbaum, 2011; van

Kesteren et al., 2013; Wang& Morris, 2010). Third, we concentrated on a limited age range in

middle childhood. To characterize how neural mechanisms of memory consolidation evolve over

time, future studies should include other developmental cohorts. Fourth, we acknowledge that

our study design leads to temporal autocorrelation in the BOLD signal when calculating RSA

between fixation and scene time windows. Although we argue that our results, given the identical

procedure over time, are more attributed to the delay-related changes in the neural reinstatement,

future studies should tailor the design of the retrieval procedure to warrant cross-run

comparisons. This could be achieved by introducing the same items repeatedly across different

runs. Fifth, our task may not have been demanding enough for young adults to fully challenge

their memory retention and encourage the formation of more gist-like representations. Future

studies could explore this further by using more challenging conditions to enhance the formation

of more generic memories in adults, avoid bias related to prior knowledge. Sixth, although we

focused on ROIs associated with the recollection network and implicated in retrieval of visual

information, we did not investigate the connectivity between these brain regions and how it

changes as memories age. Future studies should investigate consolidation-related neural

connectivity patterns and their temporal dynamics in the developing brain. Finally, children in

our sample were positively biased in socio-demographical score and IQ compared to young

adults, which may restrict the generalizability of our results.

1249

CONCLUSIONS

In this study, we present novel empirical evidence on the neural mechanisms underlying the less

robust memory retention of intentionally learned object-location associations in 5-to-7-year-old

children compared to young adults. Our findings reveal that, over time, children show attenuated

consolidation-related upregulation in neocortical and cerebellar brain regions during successful

retrieval. Furthermore, they appear to form different types of memory representations than young

adults: while both groups show delay-related change in detailed scene-specific reinstatement,

only children exhibit the emergence of more generic gist-like representations, particularly after a

longer delay. Our results suggest that, unlike the mature consolidation systems in young adults,

the developing brains of early school-age children supports only partially adult-like neural

changes over time. Children show less pronounced neural upregulation in core retrieval regions.

At the same time, they appear to rely more on gist-like representations, possibly as a

developmental mechanism to scaffold and accumulate schema-relevant knowledge despite

weaker detailed memory.

1264

1265

Data availability statement

The datasets generated and analysed during the current study are available from the

corresponding authors upon reasonable request.

Conflict of interest disclosure

We have no known conflict of interest to disclose.

Acknowledgements

We thank all the children and parents who participated in the study. This work conducted at the

Berlin Center for Advanced Neuroimaging (BCAN) and was supported by the Deutsche

Forschungsgemeinschaft (DFG; German Research Foundation, Project-ID 327654276, SFB

1315, "Mechanisms and Disturbances in Memory Consolidation: from Synapses to Systems").

The work of YLS was also supported by the European Union (ERC-2018-StG-PIVOTAL-

758898) and the Hessisches Ministerium für Wissenschaft und Kunst (HMWK; project 'The

Adaptive Mind'). We also thank Henriette Schultz and Nina Wald de Chamorro for their

assistance with study management and data collection.

Author contributions

Y.L.S, C.B., A.K secured funding. I.S and Y.L.S, C.B., A.K contributed to conception and

design of the study. I.S. and P.L. performed data collection and data curation. I.S., P.L., and J.O.-

T. performed the statistical analysis. I.S. wrote the first draft of the manuscript. All authors

contributed to manuscript revision, read, and approved the submitted version.

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

MATERIALS AND METHODS

Participants

Sixty-three typically developing children and 46 young adults were recruited to participate in the

study through advertisement in newspapers, on the university campus, word-of-mouth, and city

registry. All participants had normal vision with or without correction, no history of

psychological or neurological disorders or head trauma, average IQ > 85, and were term-born

(i.e., born after 37 weeks of pregnancy). Fourteen children were excluded due to : (i) incomplete

task execution and missing data (n = 2); (ii) poor quality of the data (n = 7); (iii) technical issues

during data acquisition (n = 5). Seven young adult participants were excluded due to incomplete

task execution and missing data (n = 5) or being identified as extreme outlier (n = 2) based on

interquartile range (IQR; above $Q3_{\text{upper quartile(75th percentile)}} + 3\text{xIQR}$ or below $Q1_{\text{lower quartile(25th}}$

$_{\text{percentile)}} - 3\text{xIQR}$ (Hawkins, 1980)) for memory behavioural measures. The excluded participants

were comparable in terms of age, sex, and socio-economic status to the final sample. The final

total sample consisted of 49 children (22 female, mean age: 6.34 years, age range: 5.3 – 7.1

years), and 39 young adults (19 female, mean age: 25.60 years, age range: 21.3 – 30.8 years; see

Table 1 for more details).

All participants or their legal guardians gave written informed consent prior to participation.

The study was approved by the ethics committee of the Goethe University Frankfurt am Main

(approval E 145/18). The participants received 100 Euro as compensation for taking part in the

study.

Table 1

Sample characteristics by age group

	Children	Young adults	Group effect
	(CH; N = 49)	(YA; N = 39)	(CH vs YA)

Demographic measures	*M*	*SD*	*M*	*SD*	*p-value*	ω^2
Age	6.34	.43	25.60	2.79	***	.96
Sex (M/F)	27/22	-	20/19	-	-	-
IQ Score	117.90	12.92	107.64	12.49	***	.13
Socioeconomical Status						
ISCED – Father	6.22	1.43	4.39	1.75	***	.29
ISCED - Mother	6.17	1.34	4.08	1.85	***	.24

1317 *Notes.* Income is based on a 1-7 Scale (1 = less than 15.000 €, 7 = more than 100.000 €); ISCED = International

1318 Standard Classification of Education 2011 (*International Standard Classification of Education,* 2011);

1319 IQ = Intelligence Quotient based on K-ABC (Kaufman & Kaufman, 2015) for children and WAIS-IV (Wechsler,

1320 2015) for young adults; M = mean; SD = standard deviation; ω^2 = omega squared; *p < .05; ** < .01, *** < .001

1321 (significant difference).

1322

Materials and Procedure

Object-location associations task

Stimuli for the object-location association task were chosen based on the social studies and

science curriculum for German primary school first and second graders (see similar procedure in

Brod & Shing, 2019). The themes were chosen based on ratings provided by four primary school

teachers on the familiarity of first graders with the topics. 60 different themes (e.g., classroom,

farm, etc.) were chosen, each belonging to one of seven categories (i.e., field, water, housing,

forest, infrastructure, indoor, farming). Four scene stimuli and four thematically congruent object

pictures were selected for each theme (see Fig. 1 for an example), resulting in 240 individual

scenes and 240 individual objects. The 240 object-scene pairs were assigned to versions A and

B, each containing 120 object-scene pairs. Each participant was randomly assigned either

version A or version B. There were six possible object locations across all scenes. Around each

location, there were three possible object placements. The distribution of locations across scenes

was controlled to ensure realistic placement of the objects within the scenes (for more detailed

information see Supplementary Methods section). The object-location association task consisted

of three phases (see Fig. 1):

(i) *Initial encoding phase* (Day 0, Day 1, Day 14). A total of 120 object-location pairs were used to

create the trials in this phase, with 60 pairs presented on Day 0, 30 pairs on Day 1, and 30

pairs on Day 14. The initially learned object-scene associations on Day 0 were split in

two halves based on their categories. Specifically, half of the pairs from the first set and half of the pairs from the second set of 30 object-scene associations were used to create the set 30 remote pair for Day 1 testing. A similar procedure was repeated for the remaining pairs to create a set of remote object-scene associations for Day 14 retrieval. We tried to equally distribute the categories of pairs between the testing sets. During each trial, participants viewed an object in isolation for 2 seconds, followed by the same object superimposed on a scene at a particular location for 10 seconds. After this, a blank screen with a fixation cross was presented for 1 second. Participants were instructed to memorize the object-location pairs and to remember the exact location of the object within the scene using elaborative encoding strategies, such as creating a story or making a "mental photo" of the scene. Such elaborative encoding strategies have been shown to improve memory performance in both children and adults (Craik & Tulving, 1975; Pressley, 1982; Pressley et al., 1981; Shing et al., 2008);

(ii) *Learning phase* (Day 0, Day 1, Day 14). Following the initial encoding phase, participants continued learning the correct location of the object within the scene through adaptively repeated retrieval-encoding cycles. The cycles continued until participants achieved at least 83% correct responses or until the maximum of four cycles had been completed. The number of cycles therefore ranged from a minimum of two to a maximum of four. The 83% threshold was established during piloting as a guideline to determine the appropriate number of learning-retrieval cycles, rather than as a strict learning criterion. It served to standardize task continuation, not to serve as a basis for excluding participants post hoc. Children who did not reach the 83% threshold after the fourth cycle were still included in the analysis if their performance exceeded chance level (33%). Excluding them would have biased the sample toward higher-performing children and reduced the ecological validity of our findings. Including them ensures a more representative view of children's performance under extended learning conditions.

(iii) During each trial, participants were first presented with an isolated object for 2 seconds, followed by a one-second blank screen with a fixation cross. They were then shown a scene containing three red-framed rectangles, indicating possible location choices. Participants had to select the correct location by choosing one of the rectangles within 12 seconds, and the chosen rectangle was highlighted for 0.5 seconds. After this, feedback in

1373	the form of a smiley face was given, with the happy face for a correct answer, a sad face
1374	for an incorrect answer, and a sleeping face for a missed answer. Following the feedback,
1375	correct object-location associations were displayed for two seconds if the choice was
1376	correct and for three seconds if the choice was incorrect or missed.
1377	(iv) *Retrieval phase* (Day 1 and Day 14). The retrieval phase was conducted inside the MRI scanner.
1378	Participants were presented with 30 recently learnt item and 30 remote items learned on
1379	Day 0. The remote item sets for Day 1 and Day 14 based on items from Day 0 did not
1380	differ in the learning accuracy in either age group (all p > .06 as based on the analysis of
1381	variance).
1382	Participants were instructed to recollect and visualize ("put in front of their mental
1383	eyes") as vividly as possible the location of the object within the scene. In this way we
1384	prompted the recall of the scene and the location of the object within this scene.
1385	Each trial began with a fixation cross jittered between 3 to 7 seconds (mean of 5
1386	seconds). Participants were then presented with an isolated object for 2 seconds, followed
1387	by the presentation of another fixation cross jittered between 2 to 8 seconds (mean of 5
1388	seconds). Following the fixation cross, participants were prompted with the associated
1389	scene and were required to recall the location of the object by selecting one of the three
1390	red rectangles on the scene within 7.5 seconds. If participants failed to respond within the
1391	deadline, the trial was terminated. No time-outs were recorded for young adults, while
1392	5,4 % of time-out trials were recorded for children and these trials were excluded for
1393	analysis. After a choice was made or the response deadline was reached, the scene
1394	remained on the screen for an additional 0.5 second. The jitters and the order of
1395	presentation of recent and remote items were determined using OptimizeXGUI (Spunt,
1396	2016)which followed an exponential distribution (Dale, 1999). Ten unique recently
1397	learned items (from the same testing day) and ten unique remotely learned items (from
1398	Day 0) were distributed withing each run (in total three runs) in the order as suggested by
1399	the software as the most optimal. There were three runs with unique sets of stimuli, each
1400	resulting in thirty unique recent and thirty unique remote stimuli overall.
1401	*Assessment of demographic and cognitive covariates*
1402	IQ scores were assessed using the German version of the "Kaufman Assessment Battery for
1403	Children – Second Edition" (K-ABC II; Kaufman & Kaufman, 2015) in children and the

1404   "Wechsler Adult Intelligence Scale – Fourth Edition" (WAIS -IV; Wechsler, 2015) in young

1405   adults. General socio-demographic questionnaires to assess socio-demographic characteristics of

1406   the participants were administered as well.

1407   *Experimental Procedure*

The testing was conducted over three days (see Fig. 1B). On Day 0, the experiment began with a

short training session aimed at familiarizing participants with the object-location associations

task and elaborative encoding strategy, using five object-location pairs. The experimental task

started with the initial encoding of unique sets of object-location associations. Participants had to

learn two unique sets comprised of 30 object-location associations each. After encoding each set,

participants engaged in a brief distraction task where they listened to and had to recall a string of

numbers. Next, they underwent a learning phase with retrieval-encoding cycles until they

reached a criterion of 83% (or a maximum of four cycles). This was done to minimize variances

attributed to encoding, allowing for more accurate comparison of subsequent memory

consolidation. Afterwards, the children visited a mock scanner to become familiar with the MRI

scanning environment. This procedure involved teaching the children the sounds of MRI

scanning and training them to stay still during scanning.

On Day 1, participants first learned a new set of 30 object-location associations, using the

same learning procedure as on Day 0. This was followed by retrieval in the MRI scanner, during

which they were required to recall 30 object-location associations learnt on Day 0 (short-delay,

remote) and another 30 learnt on Day 1 (recent). On Day 14, the same procedure was followed as

on Day 1, with a new set of 30 object-location associations. They were again required to recall

30 object-location associations learnt on Day 0 (long-delay, remote) and another 30 learnt on

Day 14 (recent). In total, participants completed 60 retrieval trials in the MR scanner on Day 1

and Day 14 each, which took approximately 15-20 minutes. Besides the primary task,

participants also completed other psychometric tests across all testing sessions. Additionally,

socio-demographic questionnaires were administered to young adults and legal guardians of

children.

1431

Data acquisition

Behavioural data acquisition

The task paradigm during all phases was presented using Psychtoolbox (Kleiner et al., 2007)

software in MATLAB 9.5, R2018b (MATLAB, 2018). During the encoding and learning phases,

stimuli were presented on a computer screen with the resolution of 1920x1080 pixels. During the

retrieval phase, an MR-compatible screen with identical resolution was used, and participants

used an MR-compatible button box with three buttons. To minimize head movements, foam

cushions were placed inside the head coil, and MR-compatible headsets and ear plugs were used

to reduce the scanner noise.

Magnetic resonance imaging data acquisition

MR images were acquired on a 3 Tesla SIEMENS PRISMA MRI scanner (Siemens Medical

Solutions, Erlangen, Germany) using a 64-channel head coil at Berlin Center for Advanced

Neuroimaging, Charité Universitätsmedizin Berlin. Each session started with the acquisition of a

localizer and head scout sequences for field-of-view-alignment (FoV) based on anatomical

landmarks. T1-weighted structural images were obtained with the magnetization prepared rapid

gradient echo (MP-RAGE) pulse sequence (TR = 2500 ms, echo time = 2.9 ms, flip angle = 8°,

FoV = 256 mm, voxel size = 1x1x1 mm^3, 176 slices). Functional images were acquired using

echo-planar imaging sequences (TR = 800 ms, echo time = 37 ms, flip angle = 52°,

FoV = 208 mm, 72 slices, voxel size = 2x2x2 mm^3, maximally 588 volumes). In addition,

gradient echo images (field maps) were acquired before each functional run for correction of

magnetic field inhomogeneities.

1453

Behavioural data analysis

Learning and Consolidation

The behavioural analyses were performed with R packages (R Core Team, 2022) in RStudio

2022.07.0 (RStudio, Inc.). Throughout the analyses, statistical significance level was set

at $< .05$.

All p-values were FDR-adjusted for multiple comparisons due to multiple ROIs. As a

measure of baseline memory performance, final learning accuracy was defined as the percentage

of correctly learned locations in relation to the total number of items at the end of the learning

phase of each day. To examine memory consolidation, we quantified memory retention across

delays, focusing on trials that were correctly learned on Day 0. From these trials, we calculated

the percentage of correct responses, separately for Day 1 and Day 14. We conducted a linear

mixed-effect model (LME model) for memory measures using the lmer function from the lme4

package in R (Bates et al., 2015) and lmerTest (Kuznetsova et al., 2017). All LME models were

calculated with maximum-likelihood estimation and Subject as the random intercept to account

for between-subject variability in retention accuracy.

First, to investigate baseline memory performance, we analysed whether final learning

accuracy in all three sessions differed between groups. For that, we included the within-subject

factor of *Session* (Day 0, Day 1, and Day 14) and the between-subject factor of *Group* (children

and young adults) in the LME model. Second, for memory retention rates, we included *Session*

(Day 1, Day 14), *Item Type* (recent, remote), and *Group* (children, young adults) as fixed factors

in the LME model. In addition, we added *Subjects* as random factor, as well as *IQ*, *Sex,* and

Handedness (Kang et al., 2017; Willems et al., 2014) as covariates. Degrees of freedom were

adjusted using the Satterthwaite's method (Kuznetsova et al., 2017) if the assumptions of

homogeneity of variances were violated. Significant effects were followed up with Sidak post-

hoc multiple comparisons. For further group differences in socio-demographic measures, we

performed one-way independent analysis of variance (ANOVA) or Games-Howell test (S. Lee &

Lee, 2018). The effect size estimation was performed using omega squared (w^2) as a less biased

estimate for reporting practical significance of observed effects (Okada, 2013). To determine the

amount of variance explained by the model, we used partR2 package (Stoffel et al., 2021).

1483

1484   *fMRI data pre-processing*

1485   Anatomical and functional MR data was pre-processed using fMRIPrep 22.0.0 (Esteban et al.,

1486   2019), based on Nipype 1.8.3 (Gorgolewski et al., 2011). Detailed description of the anatomical

1487   and functional data pre-processing can be found in Supplementary Methods section.

1488

fMRI data analysis

FMRI data analysis was conducted with FEAT in FSL (Version 6.0.1, FMRIB's Software

Library, Jenkinson et al., 2012; Smith et al., 2004; Woolrich et al., 2009). Prior to that, single

runs were excluded if there was (i) root-mean-square realignment estimates(Jenkinson et al.,

2002) exceeding 1mm; and (ii) framewise displacement (FD) > 1, and (iii) less than two correct

trials in the entire run. Based on these criteria, 14 single runs and two complete sessions in
children were excluded from further analysis.

General Linear Model for Mean Activation

For each participant's fMRI data, a first-level analysis was performed separately for each run
using a general linear model (GLM) with eight experimental regressors. The regressors
represented the onset and duration of the following events: (i) object recent$_{correct}$, (ii) object
remote$_{correct}$, (iii) scene recent$_{correct}$, (iv) scene remote$_{correct}$, (v) object recent$_{incorrect}$, (vi) object
remote$_{incorrect}$, (vii) scene recent$_{incorrect}$, (viii) scene remote$_{incorrect}$. The duration of object events
was two seconds, while the duration of scene events was dependent on the reaction time (RT).
The regressors were convolved with a hemodynamic response function, modelled with a double-
gamma function with first and second derivatives. Confounding regressors were also included in
the GLM and were calculated with fMRIPrep, namely six rigid body realignment parameters,
framewise displacement, and standardised DVARS (D, temporal derivatives over time courses;
VARS, variance over voxels). In addition, six anatomic component-based noise correction
(CompCor, a combination of cerebrospinal fluid and white matter) regressors, global signal[3] and
cosine drift terms were included, based on previous methodological studies (Ciric et al., 2017;
Esteban et al., 2020; Jones et al., 2021; Satterthwaite et al., 2013). The functional images were
spatially smoothed with SUSAN (Smallest Univalue Segment Assimilating Nucleus, Smith &
Brady, (1997)), applying a Gaussian kernel with a full-width at half-maximum of 6 mm. A high-
pass Gaussian filter with a cut-off period of 80 s was applied. Contrasts were defined for each
run per subject, and within-subject fixed-effects averaging across runs within each session was
conducted per subject. Group-level analysis was performed with FLAME1 (Woolrich et al.,
2004) within each session, based on the statistical maps obtained from the first-level analysis.
The main contrast of interest was *object remote > object recent*, as we were primarily interested
in the reinstatement of object-scene association before the scene was shown. Univariate analysis

---

[3] We re-run the entire first-level univariate analysis using the pipeline that excluded the global signal. The resulting activation maps (see Supplementary Figure S3.2, S4.2, S5.2, S9.2) differed notably from those obtained with the original pipeline. Specifically, group differences in cortical regions such as mPFC, cerebellum, and posterior PHG no longer reached significance, and the overall pattern of results appeared noisier. Additional analyses revealed that: (i) the global signal was not dependent on group or session in our sample; (ii) the global signal reduced inter-subject variability in children, likely reflecting improved signal quality; (iii) the global signal stabilized the signal and attenuated non-neuronal variability.

was performed with statistical tests voxel-wise and corrected for multiple comparisons with

cluster-based thresholding using a z threshold of z > 3.1 and a two-tailed probability of .001.

Several a priori regions of interest (ROI) were selected based on anatomical masks:

bilateral anterior/posterior hippocampus (HC), bilateral anterior/posterior parahippocampal gyrus

(PHG), and RSC. The masks for the medio-temporal lobe ROIs were taken from the Harvard-

Oxford Cortical and Subcortical Atlases (threshold at 30% probability; (Desikan et al., 2006)),

and the mask for RSC was taken from the Talairich Atlas (threshold at 30% probability;

Lancaster et al., 2000; Talairich & Tournoux, 1988) . For further ROIs in large cortical regions

(namely mPFC, precuneus, LOC, vlPFC, and cerebellum), anatomical masks derived from

Harvard-Oxford Cortical and Subcortical Atlases or Juelich Atlas (Amunts et al., 2020) were

combined with a functional task-related map, based on mean activation across recent and remote

objects across all participants and sessions, at voxel-wise threshold of z > 3.1 and a two-tailed

probability of .001. With these masks, the mean percent signal change (from the contrast of

object remote$_{correct}$ > object recent$_{correct}$) was extracted using FEAT in FSL for each session of

each participant, which were then submitted to statistical analysis in R. A linear mixed-effect

model was set up to model percent signal change. The linear mixed effect model was calculated

with maximum-likelihood estimation and *Subject* as random intercept to account for between-

subject variability. As fixed factors, we included *Session* (Day 1, Day 14) and *Group* (children,

young adults). We also added *IQ and sex and handedness* and mean reaction time as covariates

to the model.

1539   *Representational similarity analysis for neural reinstatement.*

For the multivariate analysis, single-event (i.e., for every event on each trial) β (beta) estimates [4]

were first computed by modelling BOLD time course with a series of Generalized Linear Models

(GLM) using the Least Square Separate method (LSS; Abdulrahman & Henson, 2016; Mumford

et al., 2012). Each trial contained three events (i.e., object, fixation, and scene), hence a total of

30 GLMs (i.e., ten for objects, ten for fixations, and ten for scenes) were computed for each run,

session, and participant. Each of the GLMs contained four experimental regressors: for instance,

one for the single fixation of interest and three more for the rest of the events (i.e., for all other

---

[4] Beta estimates were obtained from a Least Square Separate (LSS) regression model. Each event was modeled with their respective onset and duration and, as such, one beta value was estimated per event (with the lags between events differing from trial to trial). The jitter was included to enable an estimation of the patterns evoked by the events and all subsequent RSA analyses were conducted normally on these estimates without further controls.

fixations except the fixation of interest, for all objects, and for all scenes). The same set up was

followed for the object GLMs and the scene GLMs. The regressors were convolved with the

hemodynamic response function, which was modelled with a double-gamma function with first

and second derivatives. Additionally, the same confounding regressors as the ones for mean-

activation analysis were included.

Next, to assess whether mnemonic reinstatement during the fixation period, during which

participants were supposed to recollect the scenes associated with the objects, was more item-

specific or gist-like, we used the single-event beta estimates of each trial to compute two types of

Representational Similarity Matrices (RSMs; Kriegeskorte, 2008). Each RSM was computed

separately for each previously identified ROI. All subsequent analyses were performed with

homebrew scripts available at https://github.com/iryna1schommartz/memokid_fmri.

Scene-specific reinstatement: To measure the extent of scene reinstatement following object

presentation, we computed a *scene-specific reinstatement index* for each neural RSM, separately

for correctly remembered recent and correctly remembered remote scenes of each session (***see***

Figure 5A-B). For each specific scene, we computed the index as the average distance between

the "*fixation*" and "*scene period*" (Fisher-transformed Pearson's r; Fig. 5B), which was the

correlation between neural patterns during fixation and neural patterns when viewing the scene.

We averaged the index across all items, all runs within a session, and then within subjects,

resulting in a single value per predefined ROIs and sessions. In addition to scene-specific

reinstatement, we also calculated a *set-based reinstatement index* as a control analysis, which

was calculated as an average distance between "*fixation*" and "*scene period*" for a scene and

every other scene within the stimuli set (Deng et al., 2021; Ritchey et al., 2013; Wing et al.,

2015). The set-based reinstatement index reflects the baseline level of non-specific neural

activation patterns during reinstatement. We then calculated the *corrected scene-specific*

reinstatement index as the difference between set-based and scene-specific Fisher-transformed

Pearson's values (Deng et al., 2021; Ritchey et al., 2013; Wing et al., 2015). A higher value in

this index denotes more distinct scene reinstatement patterns. Only correctly retrieved items were

included for this analysis. We obtained the corrected scene-specific reinstatement indices for

recent items on Day 1 and Day 14 and tested them for session-related differences. If no

differences were observed, the set-corrected scene-specific reinstatement indices for recent

scenes on Day 1 and 14 were averaged to obtain a single value per ROI and participant. We then

conducted a final LME model, separately for each ROI, with *Subject* as the random factor and

Delay (recent, remote Day 1, remote Day 14) and *Group* (children, young adults) as fixed

factors. In addition, mean neural activation was added as a covariate into the model.

Gist-like representations: Seven overarching thematic categories were identified during stimuli

selection (i.e., field, water, housing, forest, infrastructure, indoor, farming). A within-category

similarity indices were computed based on fixation time window of correctly remembered items

belonging to the same category and excluding the similarity computation for the fixation time

windows of correctly remembered items with itself. A between-category similarity indices were

computed based on fixation time window of correctly remembered items belonging to different

categories. These indices were computed for each run and across runs, Z-standardized and then

averaged. A gist-like representations index was computed by subtracting between-categories

from within-categories Z-transformed distances ([within category$_{recent}$ – between category$_{recent}$]

and [within category$_{remote}$ – between category$_{remote}$] for each session, Fig. 7A-B) . The non-zero

values in this corrected index reflect gist-like representations*,* as the similarity distance would be

higher for pairs of trials with the same categories than for pairs with different categories. We

applied a one-sample permutation t-test to test for significance in each ROI. Similar to the

procedure described above, gist-like representations indices for recent items on Day 1 and Day

14 were averaged when no difference was found, obtaining a single value per ROI and

participant. We then conducted a final LME model, separately for each ROI, with *Subject* as the

random factor and *Delay* (recent, remote Day 1, remote Day 14) and *Group* (children, young

adults) as fixed factors and mean neural activation as a covariate, to analyse any delay-related

differences in gist-like representations index for successfully retrieved trials. Finally, we also

explored whether over time, long-delay item-specific and representations is beneficial or

detrimental for memory performance by correlating the index with memory retention rates. We

tested whether this correlation within each group differs based on ROI. If no differences were

observed, we averaged representations indices across ROIs that showed significant

representations in long delay.

Brain-behavioural relations

To examine the connections between brain function and behavior, we utilized brain metrics

generated via the application of a multivariate method known as Partial Least Square Correlation

(PLSC) (Abdi & Williams, 2013; McIntosh et al., 1996; Schommartz et al., 2023). This approach

focuses on multivariate links between specified neural measures in Regions of Interest (ROIs)

and fluctuations in memory performance over short and long delays across different age cohorts.

We argue that this multivariate strategy offers a more comprehensive understanding of the

relationships between brain metrics across various ROIs and memory performance, given their

mutual dependence and connectivity (refer to Genon et al. (2022) for similar discussions).

Initially, we established a cross-subject correlation matrix that included (i) a matrix (n x

10) comprising short and long delay brain indices (encompassing both neural upregulation,

scene-specific and gist-like indices) for all specified ROIs, and (ii) a vector (n-sized) that

represents a continuous assessment of either short-delay or long-delay memory performance

(RR): R = CORR (RR, ROIs). Prior to the correlation, all metrics were standardized. The

decomposition of this correlation matrix, R = USV', was performed using singular value

decomposition, yielding singular vectors U and V, or saliences. Here, the left singular vector

symbolizes the weights for short- or long-delay memory accuracy (U), while the right singular

vector represents ROI weights (V) indicating specific neural indices that optimally represent R,

with S being a matrix of singular values.

Subsequently, PLSC identifies a singular estimable latent variable (LV), uncovering pairs

of latent vectors with maximal covariance that best describe the association between memory

retention rates and ROI neural indices. Therefore, LV delineates distinct patterns of neural

indices across ROIs closely linked to either short- or long-delay retention rates. Moreover, we

computed a singular value for each participant, termed a within-person "profile," summarizing

the robust expression of the defined LV's pattern. This was achieved by multiplying the model-

derived ROI weight vector (V) with the within-person estimates of ROI neural metrics.

To verify the generalizability and significance of the saliences or LV, we performed 5000

permutation tests to derive a p-value. We also determined the stability of the within-LV weights

by bootstrapping with 5000 resamples, calculating a bootstrap ratio (BSRs) by dividing each

ROI's salience by its bootstrap standard error. BSRs, analogous to Z-scores, serve as normalized

robustness estimates; hence, values exceeding 1.96 (p < .05) indicate statistically stable

saliences. Utilizing PLSC for multivariate statistical analysis in one step eliminates the need for

multiple comparisons correction across all ROIs (McIntosh et al., 1996).

To avoid multicollinearity and redundancy, which might diminish the power to uncover

neural-behavioral links through conventional statistical approaches, we initially derived a single

metric per participant—a participant's expression of the latent brain pattern (i.e., brain score) for

neural indices that share the most variance with either short-delay or long-delay memory

accuracy variations. We further explored how these brain patterns correlate with memory

performance.

REFERENCES

Abdi, H. (2010). Partial least squares regression and projection on latent structure regression (PLS Regression). *WIREs Computational Statistics*, *2*(1), 97–106. https://doi.org/10.1002/wics.51

Abdi, H., & Williams, L. J. (2013). Partial Least Squares Methods: Partial Least Squares Correlation and Partial Least Square Regression. In B. Reisfeld & A. N. Mayeno (Eds.), *Computational Toxicology* (Vol. 930, pp. 549–579). Humana Press. http://link.springer.com/10.1007/978-1-62703-059-5_23

Abdulrahman, H., & Henson, R. N. (2016). Effect of trial-to-trial variability on optimal event-related fMRI design: Implications for Beta-series correlation and multi-voxel pattern analysis. *NeuroImage*, *125*, 756–766. https://doi.org/10.1016/J.NEUROIMAGE.2015.11.009

Aminoff, E. M., Kveraga, K., & Bar, M. (2013). The role of the parahippocampal cortex in cognition. *Trends in Cognitive Sciences*, *17*(8), 379–390. https://doi.org/10.1016/j.tics.2013.06.009

Amunts, K., Mohlberg, H., Bludau, S., & Zilles, K. (2020). Julich-Brain: A 3D probabilistic atlas of the human brain's cytoarchitecture. *Science*, *369*(6506), 988–992. https://doi.org/10.1126/SCIENCE.ABB4588/SUPPL_FILE/ABB4588_AMUNTS_SM.PDF

Andreasen, N. C., O'Leary, D. S., Paradiso, S., Cizadlo, T., Arndt, S., Watkins, G. L., Boles Ponto, L. L., & Hichwa, R. D. (1999). The cerebellum plays a role in conscious episodic memory retrieval. *Human Brain Mapping*, *8*(4), 226–234. https://doi.org/10.1002/(SICI)1097-0193(1999)8:4<226::AID-HBM6>3.0.CO;2-4

Averbeck, B. B., Latham, P. E., & Pouget, A. (2006). Neural correlations, population coding and computation. *Nature Reviews Neuroscience*, *7*(5), 358–366. https://doi.org/10.1038/nrn1888

Axmacher N, & Rasch B. (2017). *Cognitive Neuroscience of Memory Consolidation* (N. Axmacher & B. Rasch, Eds.). Springer International Publishing. https://doi.org/10.1007/978-3-319-45066-7

Badre, D., & D'Esposito, M. (2009). Is the rostro-caudal axis of the frontal lobe hierarchical? *Nature Reviews Neuroscience*, *10*(9), 659–669. https://doi.org/10.1038/nrn2667

Badre, D., & Wagner, A. D. (2007). Left ventrolateral prefrontal cortex and the cognitive control of memory. *Neuropsychologia*, *45*(13), 2883–2901. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2007.06.015

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/JSS.V067.I01

Bauer, P. J. (2021). We know more than we ever learned: Processes involved in accumulation of world knowledge. *Child Development Perspectives*, *15*(4), 220–227. https://doi.org/10.1111/cdep.12430

Benear, S. L., Horwath, E. A., Cowan, E., Camacho, M. C., Ngo, C. T., Newcombe, N. S., Olson, I. R., Perlman, S. B., & Murty, V. P. (2022). Children show adult-like hippocampal pattern similarity for familiar but not novel events. *Brain Research*, *1791*, 147991. https://doi.org/10.1016/j.brainres.2022.147991

Bjorkund, D. F., Dukes, C., & Brown, R. D. (2009). The development of memory strategies. In M. L. Courage & N. Cowan (Eds.), *The development of memory in infancy and childhood* (End ed., pp. 145–175). Psychology Press.

Brainerd, C. J., & Reyna, V. F. (2002). *Fuzzy-Trace Theory: Dual Processes in Memory, Reasoning, and Cognitive Neuroscience* (pp. 41–100). https://doi.org/10.1016/S0065-2407(02)80062-3

Brainerd, C. J., Reyna, V. F., & Forrest, T. J. (2002). Are Young Children Susceptible to the False-Memory Illusion? *Child Development*, *73*(5), 1363–1377. https://doi.org/10.1111/1467-8624.00477

Brod, G. (2021). Generative Learning: Which Strategies for What Age? *Educational Psychology Review*, *33*(4), 1295–1318. https://doi.org/10.1007/s10648-020-09571-9

Brod, G., & Shing, Y. L. (2019). A boon and a bane: Comparing the effects of prior knowledge on memory across the lifespan. *Developmental Psychology*, *55*(6), 1326–1337. https://doi.org/10.1037/dev0000712

Brod, G., & Shing, Y. L. (2022). Are there Age☐Related Differences in the Effects of Prior Knowledge on Learning? Insights Gained from the Memory Congruency Effect. *Mind, Brain, and Education*, *16*(2), 89–98. https://doi.org/10.1111/mbe.12320

Brodt, S., Gais, S., Beck, J., Erb, M., Scheffler, K., & Schönauer, M. (2018). Fast track to the neocortex: A memory engram in the posterior parietal cortex. *Science*, *362*(6418), 1045–1048. https://doi.org/10.1126/SCIENCE.AAU2528/SUPPL_FILE/AAU2528_S1.MP4

Brodt, S., Pöhlchen, D., Flanagin, V. L., Glasauer, S., Gais, S., & Schönauer, M. (2016a). Rapid and independent memory formation in the parietal cortex. *Proceedings of the National Academy of Sciences*, *113*(46), 13251–13256. https://doi.org/10.1073/pnas.1605719113

Brodt, S., Pöhlchen, D., Flanagin, V. L., Glasauer, S., Gais, S., & Schönauer, M. (2016b). Rapid and independent memory formation in the parietal cortex. *Proceedings of the National Academy of Sciences*, *113*(46), 13251–13256. https://doi.org/10.1073/pnas.1605719113

Bunge, S. A., Burrows, B., & Wagner, A. D. (2004). Prefrontal and hippocampal contributions to visual associative recognition: Interactions between cognitive control and episodic retrieval. *Brain and Cognition*, *56*(2), 141–152. https://doi.org/10.1016/j.bandc.2003.08.001

Chai, X. J. (2010). Scene complexity: Influence on perception, memory, and development in the medial temporal lobe. *Frontiers in Human Neuroscience*, *4*. https://doi.org/10.3389/fnhum.2010.00021

Chai, X. J., Ofen, N., Gabrieli, J. D. E., & Whitfield-Gabrieli, S. (2014). Development of deactivation of the default-mode network during episodic memory formation. *NeuroImage*, *84*, 932–938. https://doi.org/10.1016/j.neuroimage.2013.09.032

Chen, J., Leong, Y. C., Honey, C. J., Yong, C. H., Norman, K. A., & Hasson, U. (2017). Shared memories reveal shared structure in neural activity across individuals. *Nature Neuroscience*, *20*(1), 115–125. https://doi.org/10.1038/nn.4450

Ciric, R., Wolf, D. H., Power, J. D., Roalf, D. R., Baum, G. L., Ruparel, K., Shinohara, R. T., Elliott, M. A., Eickhoff, S. B., Davatzikos, C., Gur, R. C., Gur, R. E., Bassett, D. S., & Satterthwaite, T. D. (2017). Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. *NeuroImage*, *154*, 174–187. https://doi.org/10.1016/j.neuroimage.2017.03.020

Clarke, A., Crivelli-Decker, J., & Ranganath, C. (2022). *Behavioral/Cognitive Contextual Expectations Shape Cortical Reinstatement of Sensory Representations*. https://doi.org/10.1523/JNEUROSCI.2045-21.2022

Cohen, M. A., Dilks, D. D., Koldewyn, K., Weigelt, S., Feather, J., Kell, A. JE., Keil, B., Fischl, B., Zöllei, L., Wald, L., Saxe, R., & Kanwisher, N. (2019). Representational similarity precedes category selectivity in the developing ventral visual pathway. *NeuroImage*, *197*, 565–574. https://doi.org/10.1016/j.neuroimage.2019.05.010

Craik, F. I. M., & Tulving, E. (1975). Depth of processing and the retention of words in episodic memory. *Journal of Experimental Psychology: General*, *104*(3), 268–294. https://doi.org/10.1037/0096-3445.104.3.268

Crowley, K., & Siegler, R. S. (1999). Explanation and Generalization in Young Children's Strategy Learning. *Child Development*, *70*(2), 304–316. https://doi.org/10.1111/1467-8624.00023

Dale, A. M. (1999). Optimal experimental design for event-related fMRI. *Human Brain Mapping*, *8*(2–3), 109–114. http://www.ncbi.nlm.nih.gov/pubmed/10524601

DeMaster, D. M., & Ghetti, S. (2013). Developmental differences in hippocampal and cortical contributions to episodic retrieval. *Cortex*, *49*(6), 1482–1493. https://doi.org/10.1016/j.cortex.2012.08.004

Deng, L., Davis, S. W., Monge, Z. A., Wing, E. A., Geib, B. R., Raghunandan, A., & Cabeza, R. (2021). Age-related dedifferentiation and hyperdifferentiation of perceptual and mnemonic representations. *Neurobiology of Aging*, *106*, 55–67. https://doi.org/10.1016/j.neurobiolaging.2021.05.021

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980. https://doi.org/10.1016/j.neuroimage.2006.01.021

Diamond, N. B., Armson, M. J., & Levine, B. (2020). The Truth Is Out There: Accuracy in Recall of Verifiable Real-World Events. *Psychological Science*, *31*(12), 1544–1556. https://doi.org/10.1177/0956797620954812

Du, X., Zhan, L., Chen, G., Guo, D., Li, C., Moscovitch, M., & Yang, J. (2019). Differential activation of the medial temporal lobe during item and associative memory across time. *Neuropsychologia*, *135*, 107252. https://doi.org/10.1016/j.neuropsychologia.2019.107252

Dudai, Y. (2012). The Restless Engram: Consolidations Never End. *Annual Review of Neuroscience*, *35*(1), 227–247. https://doi.org/10.1146/annurev-neuro-062111-150500

Esteban, O., Ciric, R., Finc, K., Blair, R. W., Markiewicz, C. J., Moodie, C. A., Kent, J. D., Goncalves, M., DuPre, E., Gomez, D. E. P., Ye, Z., Salo, T., Valabregue, R., Amlien, I. K., Liem, F., Jacoby, N., Stojić, H., Cieslak, M., Urchs, S., … Gorgolewski, K. J. (2020). Analysis of task-based functional MRI data preprocessed with fMRIPrep. *Nature Protocols*, *15*(7), 2186–2202. https://doi.org/10.1038/s41596-020-0327-3

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature Methods*, *16*(1), 111–116. https://doi.org/10.1038/s41592-018-0235-4

Fair, D. A., Cohen, A. L., Dosenbach, N. U. F., Church, J. A., Miezin, F. M., Barch, D. M., Raichle, M. E., Petersen, S. E., & Schlaggar, B. L. (2008). The maturing architecture of the brain's default network. *Proceedings of the National Academy of Sciences*, *105*(10), 4028–4032. https://doi.org/10.1073/pnas.0800376105

Fandakova, Y., Leckey, S., Driver, C. C., Bunge, S. A., & Ghetti, S. (2019a). Neural specificity of scene representations is related to memory performance in childhood. *NeuroImage*, *199*, 105–113. https://doi.org/10.1016/j.neuroimage.2019.05.050

Fandakova, Y., Leckey, S., Driver, C. C., Bunge, S. A., & Ghetti, S. (2019b). Neural specificity of scene representations is related to memory performance in childhood. *NeuroImage*, *199*, 105–113. https://doi.org/10.1016/j.neuroimage.2019.05.050

Fandakova, Y., Selmeczy, D., Leckey, S., Grimm, K. J., Wendelken, C., Bunge, S. A., & Ghetti, S. (2017). Changes in ventromedial prefrontal and insular cortex support the development of metamemory from childhood into adolescence. *Proceedings of the National Academy of Sciences*, *114*(29), 7582–7587. https://doi.org/10.1073/pnas.1703079114

Gaudreau, H., Carrier, J., & Montplaisir, J. (2001). Age-related modifications of NREM sleep EEG: from childhood to middle age. *Journal of Sleep Research*, *10*(3), 165–172. https://doi.org/10.1046/j.1365-2869.2001.00252.x

Genon, S., Eickhoff, S. B., & Kharabian, S. (2022). Linking interindividual variability in brain structure to behaviour. *Nature Reviews Neuroscience*, *23*(5), 307–318. https://doi.org/10.1038/s41583-022-00584-7

Ghetti, S., & Bunge, S. A. (2012a). Neural changes underlying the development of episodic memory during middle childhood. *Developmental Cognitive Neuroscience*, *2*(4), 381–395. https://doi.org/10.1016/J.DCN.2012.05.002

Ghetti, S., & Bunge, S. A. (2012b). Neural changes underlying the development of episodic memory during middle childhood. *Developmental Cognitive Neuroscience*, *2*(4), 381–395. https://doi.org/10.1016/j.dcn.2012.05.002

Ghetti, S., & Bunge, S. A. (2012c). Neural changes underlying the development of episodic memory during middle childhood. *Developmental Cognitive Neuroscience*, *2*(4), 381–395. https://doi.org/10.1016/j.dcn.2012.05.002

Gilboa, A., & Marlatte, H. (2017). Neurobiology of Schemas and Schema-Mediated Memory. *Trends in Cognitive Sciences*, *21*(8), 618–631. https://doi.org/10.1016/j.tics.2017.04.013

Gilboa, A., & Moscovitch, M. (2021). No consolidation without representation: Correspondence between neural and psychological representations in recent and remote memory. *Neuron*, *109*(14), 2239–2255. https://doi.org/10.1016/j.neuron.2021.04.025

Gogtay, N., Giedd, J. N., Lusk, L., Hayashi, K. M., Greenstein, D., Vaituzis, A. C., Nugent, T. F., Herman, D. H., Clasen, L. S., Toga, A. W., Rapoport, J. L., & Thompson, P. M. (2004). Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(21), 8174–8179. https://doi.org/10.1073/PNAS.0402680101/SUPPL_FILE/02680MOVIE4.MPG

Golarai, G., Ghahremani, D. G., Whitfield-Gabrieli, S., Reiss, A., Eberhardt, J. L., Gabrieli, J. D. E., & Grill-Spector, K. (2007). Differential development of high-level visual cortex correlates with category-specific recognition memory. *Nature Neuroscience*, *10*(4), 512–522. https://doi.org/10.1038/nn1865

Golarai, G., Liberman, A., & Grill-Spector, K. (2015). Experience Shapes the Development of Neural Substrates of Face Processing in Human Ventral Temporal Cortex. *Cerebral Cortex*, bhv314. https://doi.org/10.1093/cercor/bhv314

Gorgolewski, K., Burns, C. D., Madison, C., Clark, D., Halchenko, Y. O., Waskom, M. L., & Ghosh, S. S. (2011). Nipype: A Flexible, Lightweight and Extensible Neuroimaging Data Processing Framework in Python. *Frontiers in Neuroinformatics*, *5*. https://doi.org/10.3389/fninf.2011.00013

Grill-Spector, K., Kourtzi, Z., & Kanwisher, N. (2001). The lateral occipital complex and its role in object recognition. *Vision Research*, *41*(10–11), 1409–1422. https://doi.org/10.1016/S0042-6989(01)00073-6

Guo, D., & Yang, J. (2022). Reactivation of schema representation in lateral occipital cortex supports successful memory encoding. *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhac475

Harand, C., Bertran, F., La Joie, R., Landeau, B., Mézenge, F., Desgranges, B., Peigneux, P., Eustache, F., & Rauchs, G. (2012). The Hippocampus Remains Activated over the Long Term for the Retrieval of Truly Episodic Memories. *PLoS ONE*, *7*(8), e43495. https://doi.org/10.1371/journal.pone.0043495

Hawkins, D. M. (1980). Identification of Outliers. *Identification of Outliers*. https://doi.org/10.1007/978-94-015-3994-4/COVER

Hebscher, M., Meltzer, J. A., & Gilboa, A. (2019). A causal role for the precuneus in network-wide theta and gamma oscillatory activity during complex memory retrieval. *ELife*, *8*, e43114. https://doi.org/10.7554/eLife.43114

Hill, P. F., King, D. R., & Rugg, M. D. (2021). Age Differences In Retrieval-Related Reinstatement Reflect Age-Related Dedifferentiation At Encoding. *Cerebral Cortex*, *31*(1), 106–122. https://doi.org/10.1093/cercor/bhaa210

International Standard Classification of Education. (2011). http://www.uis.unesco.org

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage*, *17*(2), 825–841. https://doi.org/10.1006/nimg.2002.1132

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL. *NeuroImage*, *62*(2), 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jones, J. S., the CALM Team, & Astle, D. E. (2021). A transdiagnostic data-driven study of children's behaviour and the functional connectome. *Developmental Cognitive Neuroscience*, *52*, 101027. https://doi.org/10.1016/j.dcn.2021.101027

Josselyn, S. A., Köhler, S., & Frankland, P. W. (2015). Finding the engram. *Nature Reviews Neuroscience*, *16*(9), 521–534. https://doi.org/10.1038/nrn4000

Kang, S. J., Kang, K. A., Jang, H., Lee, J. Y., Lee, K. Il, Kwoen, M. S., Kim, J. S., & Park, K. M. (2017). Brain morphology according to age, sex, and handedness. *Annals of Clinical Neurophysiology*, *19*(2), 93. https://doi.org/10.14253/acn.2017.19.2.93

Kaufman, A. S., & Kaufman, N. L. (2015). *Kaufman Assessment Battery for Children - Second Edition* (P. Melchers & M. Melchers, Eds.; 2004 Pearson, Inc.).

Keresztes, A., Bender, A. R., Bodammer, N. C., Lindenberger, U., Shing, Y. L., & Werkle-Bergner, M. (2017). Hippocampal maturity promotes memory distinctiveness in childhood and adolescence. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(34), 9212–9217. https://doi.org/10.1073/PNAS.1710654114/SUPPL_FILE/PNAS.201710654SI.PDF

Keresztes, A., Ngo, C. T., Lindenberger, U., Werkle-Bergner, M., & Newcombe, N. S. (2018). Hippocampal Maturation Drives Memory from Generalization to Specificity. *Trends in Cognitive Sciences*, *22*(8), 676–686. https://doi.org/10.1016/j.tics.2018.05.004

Keresztes, A., Raffington, L., Bender, A. R., Bögl, K., Heim, C., & Shing, Y. L. (2022). Longitudinal developmental trajectories do not follow cross-sectional age associations in hippocampal subfield and memory development. *Developmental Cognitive Neuroscience*, *54*, 101085. https://doi.org/10.1016/j.dcn.2022.101085

Kleiner, M., Brainard, D., Pelli, D., Ingling, A., Murray, R., & Broussard, C. (2007). What's new in psychtoolbox-3. *Perception*, *36*(14), 1–16.

Krenz, V., Alink, A., Sommer, T., Roozendaal, B., & Schwabe, L. (2023). Time-dependent memory transformation in hippocampus and neocortex is semantic in nature. *Nature Communications*, *14*(1), 6037. https://doi.org/10.1038/s41467-023-41648-1

Kriegeskorte, N. (2008). Representational similarity analysis – connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*. https://doi.org/10.3389/neuro.06.004.2008

Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: integrating cognition, computation, and the brain. *Trends in Cognitive Sciences*, *17*(8), 401–412. https://doi.org/10.1016/j.tics.2013.06.007

Kuhl, B. A., Bainbridge, W. A., & Chun, M. M. (2012). Neural Reactivation Reveals Mechanisms for Updating Memory. *Journal of Neuroscience*, *32*(10), 3453–3461. https://doi.org/10.1523/JNEUROSCI.5846-11.2012

Kuhl, B. A., & Chun, M. M. (2014). Successful Remembering Elicits Event-Specific Activity Patterns in Lateral Parietal Cortex. *Journal of Neuroscience*, *34*(23), 8051–8060. https://doi.org/10.1523/JNEUROSCI.4328-13.2014

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26. https://doi.org/10.18637/JSS.V082.I13

Lancaster, J. L., Woldorff, M. G., Parsons, L. M., Liotti, M., Freitas, C. S., Rainey, L., Kochunov, P. V., Nickerson, D., Mikiten, S. A., & Fox, P. T. (2000). Automated Talairach Atlas labels for functional brain mapping. *Human Brain Mapping*, *10*(3), 120–131. https://doi.org/10.1002/1097-0193(200007)10:3<120::AID-HBM30>3.0.CO;2-8

Lebel, C., Gee, M., Camicioli, R., Wieler, M., Martin, W., & Beaulieu, C. (2012). Diffusion tensor imaging of white matter tract evolution over the lifespan. *NeuroImage*, *60*(1), 340–352. https://doi.org/10.1016/J.NEUROIMAGE.2011.11.094

Lee, J. K., Ekstrom, A. D., & Ghetti, S. (2014). Volume of hippocampal subfields and episodic memory in childhood and adolescence. *NeuroImage*, *94*, 162–171. https://doi.org/10.1016/j.neuroimage.2014.03.019

Lee, S., & Lee, D. K. (2018). What is the proper way to apply the multiple comparison test? *Korean Journal of Anesthesiology*, *71*(5), 353. https://doi.org/10.4097/KJA.D.18.00242

Lenroot, R. K., & Giedd, J. N. (2006). Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neuroscience & Biobehavioral Reviews*, *30*(6), 718–729. https://doi.org/10.1016/J.NEUBIOREV.2006.06.001

Lerche, V., & Voss, A. (2019). Experimental validation of the diffusion model based on a slow response time paradigm. *Psychological Research*, *83*(6), 1194–1209. https://doi.org/10.1007/s00426-017-0945-8

Masís-Obando, R., Norman, K. A., & Baldassano, C. (2022). Schema representations in distinct brain networks support narrative memory during encoding and retrieval. *ELife*, *11*. https://doi.org/10.7554/eLife.70445

MATLAB. (2018). *9.7.0.1190202* (R2019b ed.). The MathWorks Inc.

McIntosh, A. R., Bookstein, F. L., Haxby, J. V, & Grady, C. L. (1996). Spatial Pattern Analysis of Functional Brain Images Using Partial Least Squares. *NeuroImage*, *3*(3), 143–157. https://doi.org/10.1006/nimg.1996.0016

McKenzie, S., & Eichenbaum, H. (2011). Consolidation and Reconsolidation: Two Lives of Memories? *Neuron*, *71*(2), 224–233. https://doi.org/10.1016/j.neuron.2011.06.037

Mills, K. L., Goddings, A. L., Herting, M. M., Meuwese, R., Blakemore, S. J., Crone, E. A., Dahl, R. E., Güroğlu, B., Raznahan, A., Sowell, E. R., & Tamnes, C. K. (2016). Structural

brain development between childhood and adulthood: Convergence across four longitudinal samples. *NeuroImage*, *141*, 273–281. https://doi.org/10.1016/J.NEUROIMAGE.2016.07.044

Milton, F., Muhlert, N., Butler, C. R., Smith, A., Benattayallah, A., & Zeman, A. Z. (2011). An fMRI study of long-term everyday memory using SenseCam. *Https://Doi.Org/10.1080/09658211.2011.552185*, *19*(7), 733–744. https://doi.org/10.1080/09658211.2011.552185

Mitchell, A. S., Czajkowski, R., Zhang, N., Jeffery, K., & Nelson, A. J. D. (2018). Retrosplenial cortex and its role in spatial cognition. *Brain and Neuroscience Advances*, *2*, 239821281875709. https://doi.org/10.1177/2398212818757098

Moscovitch, M., & Gilboa, A. (n.d.). *Systems consolidation, transformation and reorganization: Multiple Trace Theory, Trace Transformation Theory and their Competitors* (Vol. 1, Issue 2). Oxford University Press.

Moscovitch, M., & Gilboa, A. (2022). Systems consolidation, transformation and reorganization: Multiple Trace Theory, Trace Transformation Theory and their Competitors. *PsyArXiv Preprints*, 1–125.

Mumford, J. A., Turner, B. O., Ashby, F. G., & Poldrack, R. A. (2012). Deconvolving BOLD activation in event-related designs for multivoxel pattern classification analyses. *NeuroImage*, *59*(3), 2636–2643. https://doi.org/10.1016/J.NEUROIMAGE.2011.08.076

Murray, J. G., Howie, C. A., & Donaldson, D. I. (2015). The neural mechanism underlying recollection is sensitive to the quality of episodic memory: Event related potentials reveal a some-or-none threshold. *NeuroImage*, *120*, 298–308. https://doi.org/10.1016/j.neuroimage.2015.06.069

Nadel, L., & Moscovitch, M. (1997). Memory consolidation, retrograde amnesia and the hippocampal complex. *Current Opinion in Neurobiology*, *7*(2), 217–227. https://doi.org/10.1016/S0959-4388(97)80010-4

Nadel, L., Samsonovich, A., Ryan, L., & Moscovitch, M. (2000). Multiple trace theory of human memory: Computational, neuroimaging, and neuropsychological results. *Hippocampus*, *10*(4), 352–368. https://doi.org/10.1002/1098-1063(2000)10:4<352::AID-HIPO2>3.0.CO;2-D

Ngo, C. T., Benear, S. L., Popal, H., Olson, I. R., & Newcombe, N. S. (2021). Contingency of semantic generalization on episodic specificity varies across development. *Current Biology*, *31*(12), 2690-2697.e5. https://doi.org/10.1016/j.cub.2021.03.088

Nishimura, M., Scherf, K. S., Zachariou, V., Tarr, M. J., & Behrmann, M. (2015). Size Precedes View: Developmental Emergence of Invariant Object Representations in Lateral Occipital Complex. *Journal of Cognitive Neuroscience*, *27*(3), 474–491. https://doi.org/10.1162/jocn_a_00720

Oedekoven, C. S. H., Keidel, J. L., Berens, S. C., & Bird, C. M. (2017). Reinstatement of memory representations for lifelike events over the course of a week. *Scientific Reports*, *7*(1), 14305. https://doi.org/10.1038/s41598-017-13938-4

Ofen, N. (2012). The development of neural correlates for memory formation. *Neuroscience & Biobehavioral Reviews*, *36*(7), 1708–1717. https://doi.org/10.1016/j.neubiorev.2012.02.016

Ofen, N., Kao, Y. C., Sokol-Hessner, P., Kim, H., Whitfield-Gabrieli, S., & Gabrieli, J. D. E. (2007). Development of the declarative memory system in the human brain. *Nature Neuroscience 2007 10:9*, *10*(9), 1198–1205. https://doi.org/10.1038/nn1950

Okada, K. (2013). Is Omega Squared Less Biased? a Comparison of Three Major Effect Size Indices in One-Way Anova. *Behaviormetrika*, *40*(2), 129–147. https://doi.org/10.2333/bhmk.40.129

Østby, Y., Tamnes, C. K., Fjell, A. M., & Walhovd, K. B. (2012). Dissociating Memory Processes in the Developing Brain: The Role of Hippocampal Volume and Cortical Thickness in Recall after Minutes versus Days. *Cerebral Cortex*, *22*(2), 381–390. https://doi.org/10.1093/cercor/bhr116

Palada, H., Neal, A., Vuckovic, A., Martin, R., Samuels, K., & Heathcote, A. (2016). Evidence accumulation in a complex task: Making choices about concurrent multiattribute stimuli under time pressure. *Journal of Experimental Psychology: Applied*, *22*(1), 1–23. https://doi.org/10.1037/xap0000074

Peiffer, A., Brichet, M., De Tiège, X., Peigneux, P., & Urbain, C. (2020). The power of children's sleep - Improved declarative memory consolidation in children compared with adults. *Scientific Reports*, *10*(1), 9979. https://doi.org/10.1038/s41598-020-66880-3

Plachti, A., Latzman, R. D., Balajoo, S. M., Hoffstaedter, F., Madsen, K. S., Baare, W., Siebner, H. R., Eickhoff, S. B., & Genon, S. (2023). Hippocampal anterior- posterior shift in childhood and adolescence. *Progress in Neurobiology*, *225*, 102447. https://doi.org/10.1016/j.pneurobio.2023.102447

Pressley, M. (1982). Elaboration and Memory Development. *Child Development*, *53*(2), 296. https://doi.org/10.2307/1128972

Pressley, M., Samuel, J., Hershey, M. M., Bishop, S. L., & Dickinson, D. (1981). Use of a mnemonic technique to teach young children foreign language vocabulary. *Contemporary Educational Psychology*, *6*(2), 110–116. https://doi.org/10.1016/0361-476X(81)90039-4

Preston, A. R., & Eichenbaum, H. (2013). Interplay of Hippocampus and Prefrontal Cortex in Memory. *Current Biology*, *23*(17), R764–R773. https://doi.org/10.1016/j.cub.2013.05.041

R Core Team. (2022). *R: A language and environment for statistical computing* (4.1.2). R Foundation for Statistical Computing.

Ranganath, C., & Ritchey, M. (2012). Two cortical systems for memory-guided behaviour. *Nature Reviews Neuroscience*, *13*(10), 713–726. https://doi.org/10.1038/nrn3338

Ratcliff, R., Love, J., Thompson, C. A., & Opfer, J. E. (2012). Children Are Not Like Older Adults: A Diffusion Model Analysis of Developmental Changes in Speeded Responses. *Child Development*, *83*(1), 367–381. https://doi.org/10.1111/j.1467-8624.2011.01683.x

Ratcliff, R., & McKoon, G. (2008). The Diffusion Decision Model: Theory and Data for Two-Choice Decision Tasks. *Neural Computation*, *20*(4), 873–922. https://doi.org/10.1162/neco.2008.12-06-420

Ratcliff, R., Thapar, A., & McKoon, G. (2011). Effects of aging and IQ on item and associative memory. *Journal of Experimental Psychology: General*, *140*(3), 464–487. https://doi.org/10.1037/a0023810

Reyna, V. F., & Brainerd, C. J. (1995). Fuzzy-trace theory: An interim synthesis. *Learning and Individual Differences*, *7*(1), 1–75. https://doi.org/10.1016/1041-6080(95)90031-4

Reyna, V. F., & Brainerd, C. J. (1998). Fuzzy-Trace Theory and False Memory: New Frontiers. *Journal of Experimental Child Psychology*, *71*(2), 194–209. https://doi.org/10.1006/jecp.1998.2472

Reyna, V. F., Corbin, J. C., Weldon, R. B., & Brainerd, C. J. (2016). How fuzzy-trace theory predicts true and false memories for words, sentences, and narratives. *Journal of Applied Research in Memory and Cognition*, *5*(1), 1–9. https://doi.org/10.1016/j.jarmac.2015.12.003

Richter, F. R., Cooper, R. A., Bays, P. M., & Simons, J. S. (2016). Distinct neural mechanisms underlie the success, precision, and vividness of episodic memory. *ELife*, *5*. https://doi.org/10.7554/eLife.18260

Ritchey, M., & Cooper, R. A. (2020). Deconstructing the Posterior Medial Episodic Network. *Trends in Cognitive Sciences*, *24*(6), 451–465. https://doi.org/10.1016/j.tics.2020.03.006

Ritchey, M., Wing, E. A., LaBar, K. S., & Cabeza, R. (2013). Neural Similarity Between Encoding and Retrieval is Related to Memory Via Hippocampal Interactions. *Cerebral Cortex*, *23*(12), 2818–2828. https://doi.org/10.1093/cercor/bhs258

Roüast, N. M., & Schönauer, M. (2023). Continuously changing memories: a framework for proactive and non-linear consolidation. *Trends in Neurosciences*, *46*(1), 8–19. https://doi.org/10.1016/j.tins.2022.10.013

Ruggeri, A., Markant, D. B., Gureckis, T. M., Bretzke, M., & Xu, F. (2019). Memory enhancements from active control of learning emerge across development. *Cognition*, *186*, 82–94. https://doi.org/10.1016/j.cognition.2019.01.010

Sanefuji, M., Takada, Y., Kimura, N., Torisu, H., Kira, R., Ishizaki, Y., & Hara, T. (2011). Strategy in short-term memory for pictures in childhood: A near-infrared spectroscopy study. *NeuroImage*, *54*(3), 2394–2400. https://doi.org/10.1016/j.neuroimage.2010.09.090

Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., Eickhoff, S. B., Hakonarson, H., Gur, R. C., Gur, R. E., & Wolf, D. H. (2013). An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *NeuroImage*, *64*, 240–256. https://doi.org/10.1016/j.neuroimage.2012.08.052

Schlichting, M. L., Mumford, J. A., & Preston, A. R. (2015). Learning-related representational changes reveal dissociable integration and separation signatures in the hippocampus and prefrontal cortex. *Nature Communications*, *6*(1), 8151. https://doi.org/10.1038/ncomms9151

Schommartz, I., Kaindl, A. M., Buss, C., & Shing, Y. L. (2024). Short- and long-delay consolidation of memory accessibility and precision across childhood and young adulthood. *Developmental Psychology*. https://doi.org/10.1037/dev0001691

Schommartz, I., Lembcke, P. F., Pupillo, F., Schuetz, H., de Chamorro, N. W., Bauer, M., Kaindl, A. M., Buss, C., & Shing, Y. L. (2023). Distinct multivariate structural brain profiles are related to variations in short- and long-delay memory consolidation across children and young adults. *Developmental Cognitive Neuroscience*, *59*. https://doi.org/10.1016/J.DCN.2022.101192

Sekeres, M. J., Bonasia, K., St-Laurent, M., Pishdadian, S., Winocur, G., Grady, C., & Moscovitch, M. (2016). Recovering and preventing loss of detailed memory: differential rates of forgetting for detail types in episodic memory. *Learning & Memory*, *23*(2), 72–82. https://doi.org/10.1101/lm.039057.115

Sekeres, M. J., Moscovitch, M., & Winocur, G. (2017a). Mechanisms of Memory Consolidation and Transformation. In N. Axmacher & B. Rasch (Eds.), *Cognitive Neuroscience of Memory Consolidation* (pp. 17–44). Springer International Publishing. http://link.springer.com/10.1007/978-3-319-45066-7_2

Sekeres, M. J., Moscovitch, M., & Winocur, G. (2017b). Mechanisms of Memory Consolidation and Transformation. In N. Axmacher & B. Rasch (Eds.), *Cognitive Neuroscience of Memory Consolidation* (pp. 17–44). Springer International Publishing. http://link.springer.com/10.1007/978-3-319-45066-7_2

Sekeres, M. J., Moscovitch, M., Winocur, G., Pishdadian, S., Nichol, D., & Grady, C. L. (2021). Reminders activate the prefrontal⍰medial temporal cortex and attenuate forgetting of event memory. *Hippocampus*, *31*(1), 28–45. https://doi.org/10.1002/hipo.23260

Sekeres, M. J., Winocur, G., & Moscovitch, M. (2018). The hippocampus and related neocortical structures in memory transformation. *Neuroscience Letters*, *680*, 39–53. https://doi.org/10.1016/j.neulet.2018.05.006

Sekeres, M. J., Winocur, G., Moscovitch, M., Anderson, J. A. E., Pishdadian, S., Martin Wojtowicz, J., St⍰Laurent, M., McAndrews, M. P., & Grady, C. L. (2018). Changes in patterns of neural activity underlie a time⍰dependent transformation of memory in rats and humans. *Hippocampus*, *28*(10), 745–764. https://doi.org/10.1002/hipo.23009

Semon, R. W. (1921). *The mneme*. Allen & Unwin.

Shing, Y. L., Werkle-Bergner, M., Brehmer, Y., Müller, V., Li, S.-C., & Lindenberger, U. (2010a). Episodic memory across the lifespan: The contributions of associative and strategic components. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1080–1091. https://doi.org/10.1016/j.neubiorev.2009.11.002

Shing, Y. L., Werkle-Bergner, M., Brehmer, Y., Müller, V., Li, S.-C., & Lindenberger, U. (2010b). Episodic memory across the lifespan: The contributions of associative and strategic components. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1080–1091. https://doi.org/10.1016/j.neubiorev.2009.11.002

Shing, Y. L., Werkle-Bergner, M., Li, S.-C., & Lindenberger, U. (2008). Associative and strategic components of episodic memory: A life-span dissociation. *Journal of Experimental Psychology: General*, *137*(3), 495–513. https://doi.org/10.1037/0096-3445.137.3.495

Simons, J. S., & Spiers, H. J. (2003). Prefrontal and medial temporal lobe interactions in long-term memory. *Nature Reviews Neuroscience*, *4*(8), 637–648. https://doi.org/10.1038/nrn1178

Sluzenski, J., Newcombe, N. S., & Kovacs, S. L. (2006). Binding, relational memory, and recall of naturalistic events: A developmental perspective. *Journal of Experimental Psychology: Learning Memory and Cognition*, *32*(1), 89–100. https://doi.org/10.1037/0278-7393.32.1.89

Smith, S. M., & Brady, J. M. (1997). SUSAN—A New Approach to Low Level Image Processing. *International Journal of Computer Vision*, *23*(1), 45–78. https://doi.org/10.1023/A:1007963824710

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*, *23*, S208–S219. https://doi.org/10.1016/j.neuroimage.2004.07.051

Söderlund, H., Moscovitch, M., Kumar, N., Mandic, M., & Levine, B. (2012). As time goes by: Hippocampal connectivity changes with remoteness of autobiographical memory retrieval. *Hippocampus*, *22*(4), 670–679. https://doi.org/10.1002/hipo.20927

Sowell, E. R., Trauner, D. A., Gamst, A., & Jernigan, T. L. (2002). Development of cortical and subcortical brain structures in childhood and adolescence: a structural MRI study. *Developmental Medicine and Child Neurology*, *44*(01), 4. https://doi.org/10.1017/S0012162201001591

Spunt, B. (2016). *Easy-Optimize-X: Formal Release For Archiving On Zenodo*. Zenodo. https://zenodo.org/record/58616

Squire, L. R., & Alvarez, P. (1995). Retrograde amnesia and memory consolidation: a neurobiological perspective. *Current Opinion in Neurobiology*, *5*(2), 169–177. https://doi.org/10.1016/0959-4388(95)80023-9

Squire, L. R., Genzel, L., Wixted, J. T., & Morris, R. G. (2015). Memory Consolidation. *Cold Spring Harbor Perspectives in Biology*, *7*(8), a021766. https://doi.org/10.1101/cshperspect.a021766

St-Laurent, M., & Buchsbaum, B. R. (2019). How Multiple Retrievals Affect Neural Reactivation in Young and Older Adults. *The Journals of Gerontology: Series B*, *74*(7), 1086–1100. https://doi.org/10.1093/geronb/gbz075

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2021). partR2□: partitioning R^2 in generalized linear mixed models. *PeerJ*, *9*, e11414. https://doi.org/10.7717/peerj.11414

Stroukov, E., Kumral, D., & Schoenauer, M. (2022). Cerebellar involvement in memory formation. *Psychologie Und Gehirn*, 387–387.

Takashima, A., Nieuwenhuis, I. L. C., Jensen, O., Talamini, L. M., Rijpkema, M., & Fernandez, G. (2009). Shift from Hippocampal to Neocortical Centered Retrieval Network with Consolidation. *Journal of Neuroscience*, *29*(32), 10087–10093. https://doi.org/10.1523/JNEUROSCI.0799-09.2009

Takashima, A., Petersson, K. M., Rutters, F., Tendolkar, I., Jensen, O., Zwarts, M. J., McNaughton, B. L., & Fernández, G. (2006). Declarative memory consolidation in humans: A prospective functional magnetic resonance imaging study. *Proceedings of the National Academy of Sciences*, *103*(3), 756–761. https://doi.org/10.1073/pnas.0507774103

Talairich, J., & Tournoux, P. (1988). *Co-planar stereotaxic atlas of the human brain 3-dimentional proportional system: an approach to cerebral imaging 1988*. Georg Thieme Verlag.

Tambini, A., & D'Esposito, M. (2020). Causal Contribution of Awake Post-encoding Processes to Episodic Memory Consolidation. *Current Biology*, *30*(18), 3533-3543.e7. https://doi.org/10.1016/J.CUB.2020.06.063

Tambini, A., Ketz, N., & Davachi, L. (2010). Enhanced Brain Correlations during Rest Are Related to Memory for Recent Experiences. *Neuron*, *65*(2), 280–290. https://doi.org/10.1016/j.neuron.2010.01.001

Tanrıverdi, B., Cowan, E. T., Metoki, A., Jobson, K. R., Murty, V. P., Chein, J., & Olson, I. R. (2022). Awake Hippocampal-Cortical Co-reactivation Is Associated with Forgetting. *BioRxiv*, 2022.12.10.519896. https://doi.org/10.1101/2022.12.10.519896

Tompary, A., & Davachi, L. (2017). Consolidation Promotes the Emergence of Representational Overlap in the Hippocampus and Medial Prefrontal Cortex. *Neuron*, *96*(1), 228-241.e5. https://doi.org/10.1016/j.neuron.2017.09.005

Tompary, A., Zhou, W., & Davachi, L. (2020). Schematic memories develop quickly, but are not expressed unless necessary. *PsyArXiv*.

Trelle, A. N., Henson, R. N., & Simons, J. S. (2019). Neural evidence for age-related differences in representational quality and strategic retrieval processes. *Neurobiology of Aging*, *84*, 50–60. https://doi.org/10.1016/j.neurobiolaging.2019.07.012

van der Linden, M., Berkers, R. M. W. J., Morris, R. G. M., & Fernández, G. (2017). Angular Gyrus Involvement at Encoding and Retrieval Is Associated with Durable But Less Specific Memories. *The Journal of Neuroscience*, *37*(39), 9474–9485. https://doi.org/10.1523/JNEUROSCI.3603-16.2017

van Kesteren, M. T. R., Rijpkema, M., Ruiter, D. J., & Fernández, G. (2013). Consolidation Differentially Modulates Schema Effects on Memory for Items and Associations. *PLoS ONE*, *8*(2), e56155. https://doi.org/10.1371/journal.pone.0056155

Vann, S. D., Aggleton, J. P., & Maguire, E. A. (2009). What does the retrosplenial cortex do? *Nature Reviews Neuroscience*, *10*(11), 792–802. https://doi.org/10.1038/nrn2733

Varga, N. L., Rooms, H. E., Molitor, R. J., Martinez, L., Hipskind, E. M., Mack, M. L., Preston, A. R., & Schlichting, M. L. (2023). Differentiation of related events in hippocampus supports memory reinstatement in development. *BioRxiv*.

Wagner, L. (2010). The acquisition of semantics. *WIREs Cognitive Science*, *1*(4), 519–526. https://doi.org/10.1002/wcs.24

Wang, J.-Y., Weber, F. D., Zinke, K., Inostroza, M., & Born, J. (2018). More Effective Consolidation of Episodic Long-Term Memory in Children Than Adults-Unrelated to Sleep. *Child Development*, *89*(5), 1720–1734. https://doi.org/10.1111/cdev.12839

Wang, S.-H., & Morris, R. G. M. (2010). Hippocampal-Neocortical Interactions in Memory Formation, Consolidation, and Reconsolidation. *Annual Review of Psychology*, *61*(1), 49–79. https://doi.org/10.1146/annurev.psych.093008.100523

Wechsler, D. (2015). *Wechsler Adult Intelligence Scale - Fourth Edition* (F. Petermann, Ed.; 2004 Pearson).

Willems, R. M., der Haegen, L. Van, Fisher, S. E., & Francks, C. (2014). On the other hand: including left-handers in cognitive neuroscience and neurogenetics. *Nature Reviews Neuroscience*, *15*(3), 193–201. https://doi.org/10.1038/nrn3679

Wing, E. A., Ritchey, M., & Cabeza, R. (2015). Reinstatement of Individual Past Events Revealed by the Similarity of Distributed Activation Patterns during Encoding and Retrieval. *Journal of Cognitive Neuroscience*, *27*(4), 679–691. https://doi.org/10.1162/jocn_a_00740

Winocur, G., & Moscovitch, M. (2011). Memory Transformation and Systems Consolidation. *Journal of the International Neuropsychological Society*, *17*(05), 766–780. https://doi.org/10.1017/S1355617711000683

Woolrich, M. W., Behrens, T. E. J., Beckmann, C. F., Jenkinson, M., & Smith, S. M. (2004). Multilevel linear modelling for FMRI group analysis using Bayesian inference. *NeuroImage*, *21*(4), 1732–1747. https://doi.org/10.1016/j.neuroimage.2003.12.023

Woolrich, M. W., Jbabdi, S., Patenaude, B., Chappell, M., Makni, S., Behrens, T., Beckmann, C., Jenkinson, M., & Smith, S. M. (2009). Bayesian analysis of neuroimaging data in FSL. *NeuroImage*, *45*(1), S173–S186. https://doi.org/10.1016/j.neuroimage.2008.10.055

Xue, G. (2018). The Neural Representations Underlying Human Episodic Memory. *Trends in Cognitive Sciences*, *22*(6), 544–561. https://doi.org/10.1016/j.tics.2018.03.004

Xue, G., Dong, Q., Chen, C., Lu, Z., Mumford, J. A., & Poldrack, R. A. (2010). Greater Neural Pattern Similarity Across Repetitions Is Associated with Better Memory. *Science*, *330*(6000), 97–101. https://doi.org/10.1126/science.1193125

Xue, G., Dong, Q., Chen, C., Lu, Z.-L., Mumford, J. A., & Poldrack, R. A. (2013). Complementary Role of Frontoparietal Activity and Cortical Pattern Similarity in Successful Episodic Memory Encoding. *Cerebral Cortex*, *23*(7), 1562–1571. https://doi.org/10.1093/cercor/bhs143

Yamashita, K. -i., Hirose, S., Kunimatsu, A., Aoki, S., Chikazoe, J., Jimura, K., Masutani, Y., Abe, O., Ohtomo, K., Miyashita, Y., & Konishi, S. (2009). Formation of Long-Term Memory Representation in Human Temporal Cortex Related to Pictorial Paired Associates.

Journal of Neuroscience, *29*(33), 10335–10340. https://doi.org/10.1523/JNEUROSCI.1328-
09.2009
Ye, Z., Shi, L., Li, A., Chen, C., & Xue, G. (2020). Retrieval practice facilitates memory
updating by enhancing and differentiating medial prefrontal cortex representations. *ELife*, *9*,
1–51. https://doi.org/10.7554/ELIFE.57023
Yonelinas, A. P., Ranganath, C., Ekstrom, A. D., & Wiltgen, B. J. (2019). A contextual binding
theory of episodic memory: systems consolidation reconsidered. *Nature Reviews.*
Neuroscience, *20*(6), 364–375. https://doi.org/10.1038/S41583-019-0150-4
Yu, W., Zadbood, A., Chanales, A. J. H., & Davachi, L. (2022). Repetition accelerates neural
markers of memory consolidation. *BioRxiv*, 2022.12.14.520481.
https://doi.org/10.1101/2022.12.14.520481
Zhou, J., Osth, A. F., Lilburn, S. D., & Smith, P. L. (2021). A circular diffusion model of
continuous-outcome source memory retrieval: Contrasting continuous and threshold
accounts. *Psychonomic Bulletin & Review*, *28*(4), 1112–1130.
https://doi.org/10.3758/s13423-020-01862-0
2208

A Overview of Learning Perfromance

Day 0 Day 1 Day 14

Correct Responses [%]

Encoding-Retrieval Cycles

Children   Young Adults

B Final Learning Performance

Final Learning Accuracy [%]

Day 0 Day 1 Day 14

Learning Session

Children

Young Adults

Retrieval (fMRI)

5s [3-7s]   2s   5s [2-8s]   RT - 7.5s   0.5s

time

D **Medial Prefrontal Cortex** E **Ventrolateral Prefrontal Cortex** F **Precuneus** G **Retrosplenial Cortex** H **Lateral Occipital Cortex**

Mean Signal Difference [Remote > Recent]

D Group **

E Group ***
Session **

F Group **
Session *

H Group **
Session ***
Group x Session *

Children   Young Adults   ☐ Day 1   ⊡ Day 14

A Index Computation (Scene)

2s 5s [2-8s] RT – 7.5s

(F_i) (S_g)

Fixation Time Window Scene Time Window

Pearson's r

Recent (Day1/14) Remote (Day 1) Remote (Day 14)

B Scene-Specific Reinstatement

Scene-Specific Reinstatement =
Scene-Specific Reinstatement – Set-Based Reinstatement

Scene Time Window (S_g)

Fixation Time Window (F_i)

	F_1S_1	F_1S_2	F_1S_3	F_1S_n
F_2S_1	F_2S_2	F_2S_3	F_1S_n
F_3S_1	F_3S_2	F_3S_3	F_1S_n
F_nS_1	F_nS_2	F_nS_3	F_nS_n

Scene-Specific Pearson's r

Set-Based Pearson's r

Scene-Specific Reinstatement (Δ z) = Fisher's z (Pearson's r $_{scene\text{-}specific}$) – Fisher's z (Pearson's r $_{set\text{-}based}$)

A Hippocampus   **B** Parahippocampal Gyrus   **C** Cerebellum

Anterior   Posterior   Anterior   Posterior

Y-axis: Scene-specific Reinstatement Index

Panel A Anterior: Delay ***, Group ***
Panel A Posterior: Delay ***, Group ***
Panel B Anterior: Delay ***, Group ***
Panel B Posterior: Delay ***, Group ***
Panel C: Delay ***, Group **

A Index Computation (Gist)

| | 2s | 5s [2-8s] | RT – 7.5s |

Task: recollect scene associated with object

Forest 1 · Forest n · Field 1 · Field n

Pearson's r — Within-Category
Pearson's r — Within-Category
Pearson's r — Between-Category
Pearson's r — Between-Category

B Gist-Like Reistatement

Fixation Time Window (F_i)

		Forest 1	Field 1	Indoor 1	Forest 2	Indoor 2	Field 2	... n
Forest 1	✛		F_1F_2 between	F_1F_3 between	F_1F_4 within	F_1F_5 between	F_1F_6 between	F_1F_n
Field 1	✛	F_2F_1 between		F_2F_3 between	F_2F_4 between	F_2F_5 between	F_2F_6 within	F_2F_n
Indoor 1	✛	F_3F_1 between	F_3F_2 between		F_3F_4 between	F_3F_5 within	F_3F_6 between	F_3F_n
Forest 2	✛	F_4F_1 within	F_4F_2 between	F_4F_3 between		F_4F_5 between	F_4F_6 between	F_4F_n
Indoor 2	✛	F_5F_1 between	F_5F_2 between	F_5F_3 within	F_5F_4 between		F_5F_6 between	F_5F_n
Field 2	✛	F_6F_1 between	F_6F_2 within	F_6F_3 between	F_6F_4 between	F_6F_5 between		F_6F_n
... n	✛	F_nF_1	F_nF_2	F_nF_3	F_nF_4	F_nF_5	F_nF_6	

(Left vertical axis labels: Forest 1, Forest 2, Forest 3, Field 1, Field 2, Field 3 ... — **Fixation Time Window (F_i)**)

Gist-like Reinstatement (Δz) = Fisher's z (Pearson's r $_{within\text{-}category}$) – Fisher's z (Pearson's r $_{between\text{-}categories}$)

A Recent Delay Brain Profile

Scene-Specific Representations

B Short Delay Brain Profile

Scene-Specific Representations

C Long Delay Brain Profile

Scene-Specific Representations

D Recent Delay Brain Score

E Short Delay Brain Score

F Long Delay Brain Score

A Short Delay Brain Profile

Gist-Like Representations

B Long Delay Brain Profile

Gist-Like Representations