1 ***LTP2* hypomorphs show genotype-by-environment interaction in early seedling traits in**
2 ***Arabidopsis thaliana***

3 Cristina M Alexandre[1], Kerry L Bubb[1], Karla M Schultz[1], Janne Lempe [2], Josh T Cuperus[1,*] and
4 Christine Queitsch[1,3,*]
5
Affiliations:
1: Department of Genome Sciences, University of Washington, Seattle WA 98195, USA.
2: Julius Kühn Institute (JKI) - Federal Research Centre for Cultivated Plants, Institute for
Breeding Research on Fruit Crops, Dresden, Germany.
3: Brotman Baty Institute for Precision Medicine, Seattle, WA 98195, USA
*:Corresponding Authors,
Cuperusj@uw.edu
Queitsch@uw.edu

ABSTRACT

Isogenic individuals can display seemingly stochastic phenotypic differences, limiting the
accuracy of genotype-to-phenotype predictions. The extent of this phenotypic variation depends
in part on genetic background, raising questions about the genes involved in controlling
stochastic phenotypic variation. Focusing on early seedling traits in *Arabidopsis thaliana*, we
found that hypomorphs of the cuticle-related gene *LTP2* greatly increased variation in seedling
phenotypes, including hypocotyl length, gravitropism and cuticle permeability. Many *ltp2*
hypocotyls were significantly shorter than wild-type hypocotyls while others resembled the wild
type. Differences in epidermal properties and gene expression between *ltp2* seedlings with long
and short hypocotyls suggest a loss of cuticle integrity as the primary determinant of the
observed phenotypic variation. We identified environmental conditions that reveal or mask the
increased variation in *ltp2* hypomorphs, and found that increased expression of its closest
paralog *LTP1* is necessary for *ltp2* phenotypes. Our results illustrate how decreased expression
of a single gene can generate starkly increased phenotypic variation in isogenic individuals in
response to an environmental challenge.

Keywords hypocotyl; lipid transfer protein; *LTP2*; cuticle integrity; non-genetic variation;
phenotypic variation; stochasticity; phenotypic robustness, hypomorph; gene-by-environment
interaction

INTRODUCTION

Genetically identical individuals can develop different phenotypes. Understanding the
mechanistic underpinnings of this non-genetic phenotypic variation and the relative contributions
of environmental factors and stochasticity holds promise for more accurate genotype-phenotype
predictions. As early as 1920, Sewall Wright wrote that a sizable fraction of the environmental
contribution to phenotypic variation is likely missed experimentally, and suggested that
stochasticity may play an important role in shaping individual phenotypes (Wright, 1920).
Individuals sampled from an isogenic population can differ because of (1) stochastic differences

in gene expression (Elowitz *et al.*, 2002; Blake *et al.*, 2003; Raser & O'Shea, 2004; Volfson *et*
al., 2006; Lomvardas *et al.*, 2006; Gimelbrant *et al.*, 2007), protein levels (Feinerman *et al.*,
2008) or metabolic states (Smith *et al.*, 2007; Heerden *et al.*, 2014); (2) parental effects (Perez
et al., 2017); or (3) differences in microenvironments, relative position or other contextual
information (Eagle & Levine, 1967; Snijder *et al.*, 2009). None of these possible causes are
mutually exclusive. Even subtle differences can have large cumulative effects, because the
internal state of individuals affects how they respond to environment factors. A classic example
of environmentally-induced heterogeneity is the behavior of a collection of temperature-sensitive
cell-cycle mutants in yeast. When grown asynchronously, these mutants show heterogeneous,
non-heritable phenotypes due to individual cells experiencing the restrictive temperature
treatment at different stages in the cell cycle (Hartwell *et al.*, 1974). A multitude of studies in
animals and plants have shown that the extent of non-genetic phenotypic variation across
individuals and populations depends in part on genotype, with some genetic backgrounds of the
same species showing greater non-genetic variation (or less phenotypic robustness) than others
(Waddington, 1942; Whitlock & Fowler, 1999; Ros *et al.*, 2004; Hall *et al.*, 2007; Hill *et al.*, 2007;
Sangster *et al.*, 2008; Ansel *et al.*, 2008; Shen *et al.*, 2012; Ayroles *et al.*, 2015; Katsanos *et al.*,
2017).

The more that increased, non-genetic phenotypic variation occurs in a particular genetic
background, the lower will be our ability to predict phenotype from genotype, because the same
genetic variants will show different expressivity in different individuals (Queitsch *et al.*, 2002;
Eldar *et al.*, 2009; Raj *et al.*, 2010; Burga *et al.*, 2011; Casanueva *et al.*, 2012; Lachowiec *et al.*,
2016; Zabinsky *et al.*, 2019). Therefore, non-genetic phenotypic variation has wide-ranging
implications, from cancer drug resistance (Sharma *et al.*, 2010; Shaffer *et al.*, 2017; Márquez-
Jurado *et al.*, 2018; Emert *et al.*, 2021) to microbial bioproduction (Delvigne & Goffin, 2014; Xiao
et al., 2016). In agriculture, trait uniformity is particularly highly prized (Finch-Savage & Bassel,
2016; Tran *et al.*, 2017), and breeding programs rely on parental performance. A better
understanding of the genetic underpinnings of non-genetic phenotypic variation and their
interplay with environmental factors might inform targeted breeding of more robustly performing
varieties. In plants, several genes have been identified that affect non-genetic variation of traits
like growth (Joseph *et al.*, 2015; Illouz-Eliaz *et al.*, 2019), organ size or number (Hall *et al.*,
2007; Hong *et al.*, 2016), germination (Abley *et al.*, 2021), early seedling phenotypes (Queitsch
et al., 2002; Mason *et al.*, 2016; Lachowiec *et al.*, 2018; Lemus *et al.*, 2023) and defense
metabolites (Jimenez-Gomez *et al.*, 2011; Joseph *et al.*, 2015).

Here, we focus on hypocotyl elongation in the dark, an adaptive trait relevant for seedling
establishment. Hypocotyl elongation in the dark shows large non-genetic variation in *A. thaliana*
(~10% coefficient of variation in hypocotyl length(Maloof *et al.*, 2001; Queitsch *et al.*, 2002;
Borevitz *et al.*, 2002; Sangster *et al.*, 2008; Lachowiec *et al.*, 2018). We found that hypomorphs
of *LTP2* (*LIPID TRANSFER PROTEIN 2*/AT2G38530), a highly expressed gene in dark-grown
seedlings, show increased phenotypic variation under specific environmental conditions. Plant
Lipid Transfer Proteins (LTPs) are a family of small (~9 kDa) lipid-binding, cysteine-rich proteins
that are commonly found in the shoot epidermis (Kader, 1996; Yeats & Rose, 2008). While
structurally similar, LTPs have different expression patterns, suggesting functional specialization
(Arondel *et al.*, 2000; Chae *et al.*, 2010). LTPs have been associated with antimicrobial activity

and cuticle physiology, and are implicated in a wide variety of biological processes (Molina &
García-Olmedo, 1993; Buhot *et al.*, 2001; Maldonado *et al.*, 2002; Nieuwland *et al.*, 2005;
Cameron *et al.*, 2006; Debono *et al.*, 2009; Chae *et al.*, 2009; Potocka *et al.*, 2012; Finkina *et*
al., 2016; Gao *et al.*, 2016).

LTP2 is abundant in the epidermal cell wall of dark-grown hypocotyls (Irshad *et al.*, 2008; Jacq
et al., 2017), where it promotes cuticle integrity and desiccation tolerance (Jacq *et al.*, 2017).
We found that under certain environmental conditions, *LTP2* is necessary for full hypocotyl
elongation in the dark, and that a decrease in *LTP2* expression increases variation in hypocotyl
length, gravitropism and cuticle permeability in isogenic seedlings. Differences in epidermal
morphology and cuticle permeability between long and short *ltp2* hypocotyls and between
growth conditions that promote or mask trait variation suggest that loss of cuticle integrity in *ltp2*
hypocotyls is the main determinant of this background's increased non-genetic phenotypic
variation.

RESULTS

<u>Decreased *LTP2* expression increases non-genetic phenotypic variation in skotomorphogenesis</u>

Young seedlings grown in the dark show common skotomorphogenic phenotypes with
elongated hypocotyls, etiolated cotyledons and short roots (Gendreau *et al.*, 1997;
Vandenbussche *et al.*, 2005). *LTP2* is among the most highly expressed genes in dark-grown
shoots (99th percentile, Figure S1A, B), suggesting that its function is required during
skotomorphogenesis. To measure the impact of *LTP2* on the phenotypic variation of elongating
hypocotyls, we used two *ltp2* hypomorphs, *ltp2-1* and *ltp2-2,* harboring T-DNA insertions less
than 500 bp upstream of the *LTP2* transcriptional start site (Figure 1A). Both lines expressed
less than 25% of wild-type *LTP2* RNA levels in dark-grown shoots (Figure 1B), and showed
similar defects in skotomorphogenesis. When grown in the dark, *ltp2* hypocotyls were shorter
(Figure 1C, D) and more variable than wild type (Col-0; Figure 1C), in both length (Figure 1E)
and orientation (Figure 1F). Hypocotyl lengths were almost twice as variable in *ltp2-1* as in Col-0
wild type (merged coefficient of variation, CV, of 21% *vs* 13%), with *ltp2-2* being slightly less
variable (merged CV 18%; Figure 1E). The higher variation in the hypomorphs was not due to a
bimodal distribution but to a continuous, wider distribution of hypocotyl lengths (Figure 1D). An
even larger difference was measured for hypocotyl orientation (CV 34-37% *vs* 10%; Figure 1F),
a proxy for reduced negative gravitropism. However, there was no substantial correlation
between hypocotyl length and orientation of individual seedlings (Figure S1C). Consistent with a
non-genetic origin for this increased variation in hypocotyl length, the selfed offspring of *ltp2*
parents with either long or short hypocotyls showed similar mean hypocotyl lengths (Figure 1G).
Other notable *ltp2* phenotypes included longer roots than hypocotyls, resulting in a smaller
hypocotyl/root ratio per individual seedling than in Col-0 wild type (Figure 1H), and a tendency
for open and expanded cotyledons (Figure 1C arrowhead, 1I). Taken together, reduced levels of
LTP2 globally affect non-genetic variation in skotomorphogenesis.

<u>*ltp2* phenotypes show strong gene-by-environment interaction</u>

Next, we explored internal and external factors that might be associated with the hypocotyl
length of individual seedlings or modulate the extent of phenotypic variation in *ltp2* hypomorphs.
We started by examining the influence of seed batch, seed germination and seed size on
hypocotyl length. We found similar mean hypocotyl lengths (Figure S2A) and similarly high
coefficients of variation with different *ltp2* seed batches (Figure 1E). There was no germination
delay or increased heterogeneity in germination in the *ltp2* hypomorphs relative to Col-0 wild
type, as inferred from the correlation of hypocotyl and root length and its time-dependent drop in
seedling development (Figure S2B). Finally, hypocotyl length of individual seedlings was not
substantially explained by seed size in either the *ltp2* hypomorphs or Col-0 wild type (Figure
S2C).

We continued by examining the impact of the growth medium. In the dark, hypocotyls grow
longer if provided with sucrose. However, the presence of sucrose also modifies hypocotyl and
root growth kinetics (Kircher & Schopfer, 2012) (Figure S3A) and is associated with a decrease
in the hypocotyl/root ratio compared to growth conditions without sucrose (Kircher & Schopfer,
2012) (Figure S3B). We reasoned that the presence of sucrose might affect variation in
hypocotyl elongation. In our typical experimental setup, seedlings grow in vertical plates, with
hypocotyls in direct contact with the medium, and sucrose-driven hypocotyl elongation depends
on shoot uptake(Figure S3C). A comparison between *ltp2* seedlings grown on MS media,
MS+1% sucrose or MS+1% glucose revealed that *ltp2* phenotypes were strongly sucrose-
dependent. Either removing sucrose or replacing it with glucose was sufficient to rescue all *ltp2*
growth phenotypes, including differences in mean length and coefficient of variation (Figure
2A,B), hypocotyl negative gravitropism (Figure 2C) and hypocotyl length/root length ratio (Figure
2D). Therefore, *ltp2* hypocotyls can fully elongate under favorable conditions. Sucrose did not
cause an increase in the coefficient of variation of hypocotyl length in wild type Col-0, despite a
noticeable increase in mean length in this condition (Figure 2A, B).

We wondered whether the effects of sucrose on *ltp2* seedlings involved osmotic stress. To test
this possibility, we added mannitol to the growth medium. Mannitol causes strong osmotic stress
in *Arabidopsis* seedlings (Zwiewka *et al.*, 2015; Kalve *et al.*, 2020). Adding equimolar amounts
of mannitol (29mM) and sucrose (1%) together improved, rather than aggravated, the *ltp2*
phenotypic defects (Figure S4). We also considered whether *ltp2* seedlings were deficient in
sucrose uptake but concluded that this scenario is unlikely because of the following results: The
hypocotyls of *ltp2* seedlings remain sucrose-sensitive and show similar sucrose-driven
responses as those of wild-type (Figure S5). Further, doubling the amount of sucrose inhibited
hypocotyl elongation to the same extent in *ltp2* and wild-type seedlings (Figure S5D). Lastly,
ltp2 hypocotyls were shorter when grown on MS+1% sucrose compared to MS alone (Figure
2A), with a significant difference in mean values (*ltp2-1*: 3.03mm CI95% 2.62-345, *ltp2-2*:
3.96mm CI95% 3.54-4.38, *ltp2-1*: $t(258.75)=14.459$; $p <2.2e-16$, *ltp2-2*: $t(274.64)=18.5$; $p <2.2e-$
16), inconsistent with deficient sucrose uptake. We conclude that neither osmotic stress nor
sucrose uptake contribute substantially to the *ltp2* phenotypes.

<u>A sucrose-dependent increase in cuticle permeability is associated with short *ltp2* hypocotyls</u>

A comparison of *ltp2* hypocotyls grown with and without sucrose revealed epidermal features
that were associated with short hypocotyls. Compared to wild-type Col-0, the epidermal surface
of *ltp2* hypocotyls was not smooth, and instead was fuzzy or wrinkled (Figure 2E); this *ltp2*
phenotype was sucrose-specific and much stronger in short than in long *ltp2* hypocotyls (Figure
2E). Furthermore, growth with sucrose greatly increased permeability to the water-soluble dye
toluidine blue, an indicator of cuticle integrity (Tanaka *et al.*, 2004), in *ltp2* hypocotyls, but not in
Col-0 hypocotyls, compared to controls without sucrose. Most *ltp2* hypocotyls stained deeply,
and over more than 50% of their full length (Figure 2F, G), consistent with reduced cuticle
integrity (Jacq *et al.*, 2017). The extent of staining varied widely in *ltp2* hypocotyls, from 0 to
100% (Figure 2G). On average, heavily stained hypocotyls were smaller than those not stained
(Figure 2G), and *ltp2* seedlings with the wild-type-like hypocotyl length/root length ratio > 1 were
less stained overall (Figure 2G). In contrast, staining in Col-0 wild-type hypocotyls was generally
weak and spatially restricted (Figure 2F, G). These results suggest that altered cuticle integrity
contributes to the sucrose-dependent inhibition of hypocotyl elongation in *ltp2* seedlings.

<u>Hypocotyl length is associated with many transcriptional differences in *ltp2* seedlings</u>

To identify what molecular functions are associated with non-genetic variation in hypocotyl
length, we compared the transcriptional profiles of Col-0 and *ltp2-1* seedlings with short (S) and
long (L) hypocotyls (bottom and top 15th percentiles, respectively; Figure 3A). For each
genotype by length combination, we performed bulk RNA-seq on two replicate pools of 70
shoots each (Figure 3A; see Figure S6A-C for QC metrics). Principal component analysis
showed three clearly distinguishable clusters: one formed by Col-0 L and S samples, a second
formed only by *ltp2-1* L samples and a third containing the *ltp2-1* S samples (Figure 3B). Nearly
half of the global variance in gene expression (47%, PC1) was correlated with mean hypocotyl
length (Figure 3C). Over 20 times as many differentially expressed genes (DEGs, log2FC >=1
and p-adj < 0.01) were found between the short and long *ltp2* samples (584, Table S2)
compared to the Col-0 samples (24, Table S1). The differentially expressed genes associated
with hypocotyl length in *ltp2* samples were enriched in gene ontology (GO) terms related to cell
wall modification, response to stress and plant defense (Figure 3D, fold-enrichment >= 2, p-adj
< 0.05). *ltp2* samples with short hypocotyls showed downregulation of cell wall-related genes
that promote growth, like *PGX1* and *XTH20* (Miedes *et al.*, 2013; Xiao *et al.*, 2014), secondary
cell wall biosynthesis genes, including the three main laccases *LAC4*, *LAC11* and *LAC17,* and
several peroxidases and genes related to casparian strip deposition (Figure 3E). Although
casparian strip biology is not well studied outside of roots, the casparian strip is present in dark-
grown hypocotyls (Karahara, 2012; Geldner, 2013). None of the genes in the enriched term
"casparian strip" were differentially expressed between Col-0 wild-type and *ltp2* seedlings with
long hypocotyls (Figure 3E), suggesting that the down-regulation of these genes may contribute
to the phenotype of *ltp2* seedlings with short hypocotyls.

We also observed the upregulation of a variety of stress and defense-response genes related to
hypoxia, response to fungus, anthocyanin and jasmonate biosynthesis (Figure 3F). A few of
these genes (11), including several of the most highly upregulated genes in *ltp2* seedlings with
short hypocotyls (Figure S6E), were also upregulated in Col-0 seedlings with short hypocotyls
compared to Col-0 seedlings with long hypocotyls (Figure S6D). This result supports the idea

that the phenotypic impact of perceived stress on hypocotyl length for a given individual can be
genotype-independent; however, *ltp2* individuals experience this impact far more frequently and
far more severely.

The majority of the differentially expressed genes (474, 81%) between the long and short *ltp2*
samples were also differentially expressed between the short samples of *ltp2* and Col-0. This
result likely reflects that the difference in mean hypocotyl length between the short *ltp2* and the
short wild-type samples is about as large as the difference between the long and the short *ltp2*
samples (Figure S6F). However, we found more than twice as many differentially expressed
genes (1638) in the comparison of the short *ltp2* and short wild-type samples (Figure S6G).
These differentially expressed genes showed similar gene ontology enrichments as found for
the comparison of long and short *ltp2* samples, related to hypoxia, oxidative stress, plant
defense and cell wall metabolism, with additional growth-related terms such as response to
auxin (Figure S6H). Indeed, we found that many auxin-responsive genes, including some with
known roles in hypocotyl elongation and/or gravitropism like *SAUR19/23/24/32* (Park *et al.*,
2007; Spartz *et al.*, 2012), *ARGOS* (Rai *et al.*, 2015), *SHY2* (Reed *et al.*, 1998; Tian *et al.*, 2002)
and *HAT2* (Sawa *et al.*, 2002), were downregulated in *ltp2* seedlings with short hypocotyls
compared to wild-type seedlings with short hypocotyls, but not compared to *ltp2* seedlings with
long hypocotyls (Table S5). Most of the differentially expressed genes in the short sample
comparison (996/1164, 86%) did not overlap with the differentially expressed genes in the long
sample comparison. Because the *ltp2* seedlings with long hypocotyls and Col-0 wild-type
seedlings with long hypocotyls differ in genotype but little in mean hypocotyl length, we
conclude that the gene expression differences across samples are strongly associated with
hypocotyl length, in particular with the strongly reduced hypocotyl length of short *ltp2* seedlings.

ltp2 phenotypes depend on upregulation of its closest paralog *LTP1*
Focusing on cuticle-related genes (see Methods; from (Li-Beisson *et al.*, 2013)), we found that
21 out of the 160 expressed in dark-grown shoots were deregulated in *ltp2* relative to Col-0
seedlings (30% more than expected, p=0.019; Table S6). Most of these genes (19/21) were
upregulated in *ltp2*, including six other LTP paralogs (Figure 4A). To explore to what extent
close paralogs may compensate for reduced *LTP2* expression, we examined the role of its
nearest paralog, *LTP1* (Figure 4B, Figure S7A). Analysis of *LTP1* and *LTP2* expression in light-
and dark-grown wild-type seedlings showed a reciprocal expression pattern that suggests non-
redundant roles during hypocotyl elongation: whereas *LTP1* was about 100 times more
abundant than *LTP2* in the light, it was 1/10 times as abundant in the dark (Figure 4C). In dark-
grown seedlings, *LTP2* accounts for more than 86% of global Type I LTP expression compared
to only 5% for *LTP1* (Figure S7B). In stark contrast, *LTP1* tends to be the more abundantly
expressed paralog in many other conditions, organs, and developmental stages (41/56
samples; Figure S7C). Moreover, we found that *LTP1* expression was no longer repressed in
the dark in *ltp2* seedlings (Figure 4C, Figure S7B). To rule out that this trend was an artifact of
bulk expression analysis, we measured *LTP1* and *LTP2* expression in individual wild-type Col-0
and *ltp2-1* seedlings and measured the length of their hypocotyls. *LTP1* expression levels in
individual *ltp2-1* seedlings recapitulated our bulk observations, and their hypocotyl lengths were
negatively correlated with *LTP1* expression (Figure S8; Spearman rho = -0.401; p=.0108). We
further found that *LTP1* and *LTP2* expression levels were correlated across individual seedlings

in both wild-type and *ltp2-1* seedlings (Figure S8A; Col-0: 0.832; p< 2.2e-16; *ltp2-1*: 0.933;
p<2.2e-16), suggesting that the *LTP1/LTP2* expression ratio may matter for phenotype. Indeed,
we observed that *LTP1/LTP2* expression ratios were weakly but significantly correlated with
hypocotyl length across individual *ltp2-1* and wild-type seedlings (Figure S8E; Spearman's rho =
0.484, p = 0.0018 *ltp2-1*; Spearman's rho = 0.437, p= 0.0059 Col-0).

We next tested whether the altered expression ratio of the paralogs might contribute to the *ltp2*
phenotypes in transgenic lines. We used an artificial microRNA to simultaneously knock-down
LTP1 and *LTP2* gene expression while aiming for a similar *LTP1/LTP2* ratio as found in Col-0
wild-type (Figure 4D). Indeed, the mean value and variation of hypocotyl length of the *ltp1 ltp2*
double knock-down seedlings closely resembled those of Col-0 wild-type seedlings with both
distributions largely overlapping (Figure 4E, F). The *ltp1 ltp2* double knock-down seedlings were
also similar to Col-0 wild-type seedlings in hypocotyl/root ratio (Figure 4G) and hypocotyl cuticle
integrity (Figure 4H). This result is even more remarkable considering that the expression levels
of the individual *LTP1* and *LTP2* genes differed substantially in the *ltp1 ltp2* double knock-down
seedlings from those observed in wild-type Col-0 seedlings (Figure S9), consistent with their
expression ratio as a determinant of the hypomorph phenotype. We conclude that the
upregulation of *LTP1* compared to low expression levels of *LTP2* in dark-grown seedlings
contributes to the variable skotomorphogenesis phenotypes of the *ltp2* hypomorphs.

DISCUSSION

Here, we show that the gene *LTP2* plays an important role in shaping skotomorphogenesis
traits. Hypomorphs of *LTP2* show increased phenotypic variation in three hypocotyl traits:
negative gravitropism, length and cuticle permeability. For the latter two, the increased variation
was accompanied by altered mean values relative to wild type, with most *ltp2* dark-grown
seedlings having short hypocotyls with highly permeable cuticles. Variation in hypocotyl length
increases in dark-grown hypocotyls upon perturbation of the chaperone Hsp90 (Queitsch *et al.*,
2002; Sangster *et al.*, 2008), or its client protein BEH4 (Lachowiec *et al.*, 2018), as well as in
AGO1 hypomorphs (Lemus *et al.*, 2023). As for the *ltp2* hypomorphs described here, in these
three cases, the increased variation in hypocotyl length was also accompanied by decreased
length means. This concordance of changes in mean and variation might be expected if wild-
type hypocotyls reach lengths close to their maximum physiological limit under these
experimental conditions.

In *ltp2* hypocotyls, a strong defect in cuticle permeability was associated with reduced
elongation and increased phenotypic variation compared to wild type; however, these
phenotypes depended on exposure to sucrose. This remarkably strong gene-by-environment
effect is unlikely due to a higher requirement for LTP2 function, as hypocotyls elongate more
when provided with an exogenous carbon source. *LTP2* is required in dark-grown hypocotyls to
seal the cuticle and prevent water loss (Jacq *et al.*, 2017). Moreover, increased cuticle
permeability is linked to structural defects and poor adhesion between the cuticle and the rest of
the cell wall. Our finding that *ltp2* hypocotyls were much more susceptible to sucrose-induced
dye uptake than wild type is consistent with *LTP2*'s role in cuticle sealing and suggests that
sucrose acts as a cuticle stress. Sucrose may trigger gene expression changes that modify

cuticle composition or act directly to increase cuticle hydration, the mechanical strain associated
with water accumulating in the cuticle and in gaps between the cuticle and the cell wall. This
added strain could lead to further cuticle detachment in sensitized *ltp2* hypocotyls, thus
aggravating their documented cuticle integrity defect (Jacq *et al.*, 2017). The sucrose-induced
loss of cuticle integrity is likely the primary determinant of the shorter *ltp2* hypocotyls and their
increased variation in length.

Dark-grown long hypocotyls have thick cuticles (Gendreau *et al.*, 1997), and shorter hypocotyls
are associated with pharmacological or genetic disruptions in cuticular wax biosynthesis and
deposition (Narukawa *et al.*, 2016). The greater need for cuticle integrity during hypocotyl
elongation in the dark may reflect a seedling's push upwards through the soil while minimizing
abrasion and the need for structural reinforcement due to cell wall thinning in very long cells
(Derbyshire *et al.*, 2007). Loss of cuticle barrier function causes water loss and activates cuticle-
dependent defense priming (Chassot *et al.*, 2007; Bessire *et al.*, 2007; L'Haridon *et al.*, 2011;
Serrano *et al.*, 2014), which inhibits growth, and is consistent with the gene ontology
enrichments observed among upregulated genes in *ltp2* seedlings with short hypocotyls
compared to those with long ones. However, the precise mechanisms by which *LPT2* facilitates
the elongation of dark-grown hypocotyls remain unknown.

In contrast to phenotypic buffers, which act on gene regulation (Lachowiec *et al.*, 2016, 2018;
Lemus *et al.*, 2023) or protein folding (Queitsch *et al.*, 2002; Sangster *et al.*, 2008; Lachowiec *et*
al., 2016; Zabinsky *et al.*, 2019), *LTP2* appears to act structurally by sealing the cuticle. The
integrity of this seal determines the effectiveness of the cuticle as an insulating barrier, thus
providing a simple mechanism of phenotypic robustness against environmental insults.
Biophysical and regulatory complexity makes plant cuticles liable to harbor considerable non-
genetic variation in their composition, ultra-structure and properties, in particular when
considering the complexity of the environments plants face. Upon damage or genetic
perturbation, this intrinsic variation is amplified as shown by the disorganized cuticles of mutants
with cutin defects or increased cuticle permeability (Lolle *et al.*, 1992; Sieber *et al.*, 2000;
Wellesen *et al.*, 2001; Chen *et al.*, 2003; Schnurr *et al.*, 2004; Kurdyukov *et al.*, 2006; Takahashi
et al., 2010).

Loss of cuticle integrity may not fully explain the increased phenotypic variation of *ltp2*
seedlings. We cannot rule out that the increased phenotypic variation depends on another, yet
undiscovered *LTP2*- function. However, our results that show genetic interaction between the
close paralogs *LTP1* and *LTP2*, and that their expression ratio is a determinant of increased
cuticle permeability and phenotypic variation during skotomorphogenesis, strongly support our
interpretation. In crown gall tumors, the only other context outside of skotomorphogenesis
where *ltp2* phenotypes have been identified (Jacq *et al.*, 2017), *LTP1* is also highly upregulated
(Deeken *et al.*, 2016). This finding is consistent with our result that the upregulation of *LTP1*
compared to *LTP2* expression is predictive of the deleterious *ltp2* phenotypes.

While paralogs with redundant or partially redundant functions can confer genetic robustness
(Gu *et al.*, 2003; Kafri *et al.*, 2005; DeLuna *et al.*, 2008, 2010; Dean *et al.*, 2008; Diss *et al.*,
2014), this is not universally observed (Ihmels *et al.*, 2007; DeLuna *et al.*, 2010; Diss *et al.*,
2017; Dandage & Landry, 2019). In fact, incomplete functional compensation can be a source of
increased phenotypic variation (Burga *et al.*, 2011; Bauer *et al.*, 2015). For example, *BEH4*, the
earliest diverged member of the *BZR/BEH* family of transcription factors, governs phenotypic
robustness of hypocotyl length by integrating regulatory cross talk among the six gene family
members (Lachowiec *et al.*, 2018). Thus, even among these closely related, partially redundant
paralogs, increased trait variation arises when the activity of *BEH4* is lost. The loss of properly
integrated regulatory cross talk as a cause of increased phenotypic variation is consistent with
our findings that upregulation of *LTP1* in the dark is associated with the *ltp2* phenotypes.

Subtle changes in gene expression that percolate through gene regulatory networks and amplify
each other to affect expression of certain core genes are thought to underlie complex diseases
and complex traits in humans (the omnigenic model, (Boyle *et al.*, 2017; Liu *et al.*, 2019)). The
genetic variants found to be associated with complex diseases and traits in genome-wide
association studies (GWAS) typically reside in regulatory regions, likely resulting in
hypomorphs. The trait heritability explained by GWAS variants tends to be small, and these
variants have little power to predict the disease risk of individuals (Manolio *et al.*, 2009; Eichler
et al., 2010; Gibson, 2012; Khera *et al.*, 2018). The low power to predict phenotype from
genotype is consistent with high non-genetic trait variation (Queitsch *et al.*, 2012). We speculate
that this non-genetic variation arises because regulatory variants cause small expression
changes that are integrated differently among individuals. In turn, these differences in
integrating expression changes will sensitize certain individuals but not others to environmental
factors, resulting in different phenotypes. At least for *LTP2*, this interpretation holds: the
upregulation of *LTP1* is not sufficient for the observed phenotypes as all *ltp2* seedlings exhibit it.
Likewise, all *ltp2* seedlings experience exposure to sucrose; however, not all seedlings have
short hypocotyls and show loss of cuticle integrity. The loss of barrier (*i.e.* cuticle) function in this
plant example likely holds lessons for studies of human traits and diseases and points to
genotype-by-environment effects as a major contributor to non-genetic variation.

Our study highlights that even highly inbred, *de facto* homozygous genetic backgrounds
maintain a physiologically relevant reservoir of phenotypic variation, which can be exposed by
stress. While stress often increases phenotypic variation in isogenic and inbred populations
(Thattai & van Oudenaarden, 2004; Newman *et al.*, 2006; Braendle & Félix, 2008; Tokatlidis *et*
al., 2010; Uyttewaal *et al.*, 2012; Holland *et al.*, 2014; Mitosch *et al.*, 2017; Sandner *et al.*, 2021;
de Groot *et al.*, 2022), phenotypic robustness (*i.e.*, low non-genetic variation) is associated with
stress tolerance and vigor in crops and lifestock (Tollenaar & Lee, 2002; Blasco *et al.*, 2017;
Elgersma *et al.*, 2018). A common strategy to achieve phenotypic robustness coupled with high
performance in agricultural settings has been the use of F1 hybrids, which are often more
uniform in phenotype than their inbred parental lines (Lewis, 1953; Smith *et al.*, 1955; Phelan &
Austad, 1994). A better understanding of the mechanistic underpinnings of non-genetic
phenotypic variation might lead to the development of crops and livestock that combine

uniformity of phenotype with broad stress tolerance. This better understanding of non-genetic
phenotypic variation will also facilitate efforts to unravel the complexity of non-Mendelian human
traits and diseases.
378

MATERIALS AND METHODS

Plant Material *Arabidopsis thaliana*, all lines are in the Col-0 background. *ltp2-1* is
SALK_026257 (ABRC), previously described (Jacq *et al.*, 2017). *ltp2-1* plants homozygous for
the T-DNA insertion (Chr2:16,128,007 SALK project) were confirmed by PCR analysis with
primer pairs CA340 (Chr2:16,128,062) + CA103 (Chr2:16,128,340) and CA249
(Chr2:16,127,7702) + CA103 (primer sequences in Table S7). *ltp2-2* is line DT7-3 (Marjorie
Matzke lab) previously described (Kanno *et al.*, 2004); contains a transgene insertion located
upstream of *LTP2*, mapped in Kanno et al., 2004. Approximate location is Chr2:16,128,261,
determined by sequencing the PCR fragment CA340+CA103. *ltp2-2* plants were confirmed
homozygous for the transgene insertion by PCR genotyping with the same primers as *ltp2-1*.
The *ltp1ltp2* double knock-down line was generated by Agrobacterium-mediated (GV3101) co-
transformation of Col-0 plants with the helper plasmid pSOUP (CD3-1124, ABRC) and the
plasmid CSHL_0103F2 (ABRC), containing an artificial microRNA against both *LTP2* and *LTP1.*
T0 seeds were selected on 15mg/ml phosphinotricin (PPT/BASTA); PPT-resistant T1 plants
were then propagated on 25mg/ml PPT; only generations T3 and above, homozygous for the
transgene, were used in hypocotyl assays.

Hypocotyl Assays *(1) Hypocotyl and root length:* Unless otherwise stated, all seed batches were
single-seed descent. Seeds were sterilized, resuspended in 0.1% (w/v) Bacto agar (BD,
Diagnostics) and spotted in a well-spaced fashion onto square plates containing 1x MS media
(Murashige & Skoog Basal Salt Mixture) pH 5.8, with 0.5g/l MES (Sigma-Aldrich), 0.3%
Phytagel (Sigma-Aldrich) and 1% sucrose (w/v). Plates were double-sealed with Micropore
surgical tape (3M) and stratified in the dark for 4 days at 4C. After stratification, plates were
exposed to 3 hours of light, then placed in vertical racks, wrapped in aluminum foil and kept at
22C in a Conviron chamber (50% RH; 16h light/8h dark; ~ 100 µmol m2 s) for 7 days. For
certain experiments, the age and media composition varied as indicated elsewhere. Genotype
placement was randomized across racks, and each genotype was distributed over multiple
plates per experiment. At 7d after stratification, plates were opened and imaged on a fixed
stand. ImageJ was used to trace and measure hypocotyl and root length for every seedling. The
hypocotyl was scored from the collet (hypocotyl/root transition zone) till the shoot apical
meristem. Root length was used to infer late germination; outlier hypocotyls with < 5mm (at 7
days) were removed from the analysis. Despite our best efforts, hypocotyl elongation proved
highly sensitive to random environmental effects, and differences of a few millimeters in mean
were not uncommon between replicates, even when controlling for seed batch (see Figure S2A)

(2) Hypocotyl negative gravitropism: The angle between the shoot apical meristem and an
imaginary vertical line, drawn starting at the base of the hypocotyl, was used to quantify the
negative gravitropic response in dark-grown hypocotyls at 7d post-stratification. Data was

expressed as deviations off vertical (in degrees), with vertical representing perfect negative
gravitropism. Measurements were done on ImageJ, as described for hypocotyl length.

(3) Hypocotyl epidermal imaging: The epidermal surface of dark-grown hypocotyls was imaged
at 7d post-stratification. Whole seedlings were placed directly on a glass slide and the hypocotyl
imaged using a Zeiss Axioplan microscope at 50x magnification.

(4)Toluidine Blue Assay Protocol based on Tanaka et al., 2003 but with a lower concentration of
dye to avoid saturation in *ltp2*. Whole dark-grown seedlings were immersed in an aqueous
solution of 0.02% (w/v) toluidine blue (Sigma) for 2mins with gentle shaking, then washed 3x
with distilled water and left in water until imaging. ImageJ was used to determine hypocotyl
length and the fraction of the total length stained with toluidine blue.

Seed Size Matched datasets of seed size and hypocotyl length were obtained by imaging the
same plate twice: (1) after stratification, when seeds were fully imbibed, and (2) after 7days of
growth in the dark. The position of the seed and the hypocotyl are the same in both pictures.
Between 10-20 seeds were spotted per plate and several plates were employed per genotype.
For imaging seeds, pictures were taken under a stereomicroscope, at 5x magnification. Seed
area was measured in ImageJ using > 8-bit > Threshold (auto) > Analyze Particles, with
settings: area, 5000-infinity; circularity >= 0.7.

qRT-PCR Seedling samples were either single individuals or pools of about 30 individuals.
Unless otherwise stated, only the shoot was harvested (includes hypocotyl, cotyledons and
shoot apical meristem). For roots, rosette leaves and flowers, all samples were pools; roots
were excised from 7d old seedlings; rosette leaves and flowers from mature plants. RNA was
extracted from LN2 frozen material using TRIZOL (Invitrogen) and DNase I treated. cDNA was
synthesized from 250-500mg of total RNA with the RevertAid First Strand cDNA Synthesis kit
(Thermo Scientific). qRT-PCR was performed with LightCycler 480 SYBR Green I Master Mix
(Roche). Relative gene expression was calculated as 2^-(Ct target - Ct reference), with either
AP2M/AT5G46630, *UBC21*/AT5G25760 or *PP2A*/AT1G13320 as reference genes. Primers are
listed in Table S7.

RNA-seq data
(1) Setup and sequencing: For each replicate of Col-0 and *ltp2-1*, seven hypocotyl assays were
set up in parallel, each containing 64-70 seedlings. At day 7 after stratification, hypocotyl length
was scored for each of the seven assays and from each, only the shoots of the 10 most extreme
seedlings at either tail of the distribution (bottom and top 15[th] percentile) were collected as
"short" and "long" samples, respectively. Each sample contained a total of 70 shoots, and
corresponds to one replicate. The process was repeated to obtain a second biological replicate,
in a total of eight samples (Col-L1, Col-L2, Col-S1, Col-S2, *ltp2*-L1, *ltp2*-L2, *ltp2*-S1, *ltp2*-S2).
RNA was extracted from all samples in parallel using the SV Promega Total RNA System
followed by NaCl/EtOH precipitation. Generation of RNA-seq libraries, multiplexing and
sequencing was outsourced to GENEWIZ (GENEWIZ Inc., NJ). Each library was sequenced on
four lanes and two flowcells of an Illumina HiSeq2500 in a 1x50bp SE format.
454

(2) Pseudoalignment of reads and estimation of transcript abundance was done using Kallisto
(Bray *et al.*, 2016).. The kallisto index was built with Arabidopsis thaliana TAIR10 cDNA models
ftp://ftp.ensemblgenomes.org/pub/plants/release-
50/fasta/arabidopsis_thaliana/cdna/Arabidopsis_thaliana.TAIR10.cdna.all.fa.gz. Transcript
abundances were quantified with parameters –single -l 200 -s 20. Kallisto abundance files were
parsed into R (v. 3.6.1) using tximport(). The tx2gene file and the TxDb object used TAIR10
annotations ftp://ftp.arabidopsis.org/home/tair/
Genes/TAIR10_genome_release/TAIR10_gff3/TAIR10_GFF3_genes_transposons.gff.
The output was a matrix of estimated counts with 26923 rows (genes) and 8 columns (samples).
464
(3) Differentially Expressed Genes (DEGs) After non-specific filtering to remove all non-
expressed genes (zero counts across all samples; 2,917 genes), and all genes which did not
have at least 1 count in all samples (4,023), a filtered matrix of 19,983 genes x 8 samples was
converted to integers and used as input for DESeq2 (Love *et al.*, 2014), with parameters:
condition = genotype x hypocotyl length, with 2 biological replicates per condition. Only genes
with a log2FC >=1 & adjusted p-value <= 0.01 were called as DEGs. As high fold-changes are
more frequent in genes with low baseline expression, we specifically chose a less stringent fold-
change cut-off to avoid discarding genes with very high expression like *LTP2.* DEGs were
obtained for four different pairwise comparisons: (1) Col-L *vs* Col-S, (2) Col-L *vs ltp2*-L, (3) Col-
S *vs* ltp2-S and (4) *ltp2*-L *vs ltp2*-S (Tables S1-S4).
475
476 *(4) Principal Component Analysis (PCA)* prcomp() was applied to a transposed standardized
477 matrix of log10(counts) with 19,983 rows x 8 columns.
478
479 *(5) Gene Ontology (GO) enrichments* were obtained with gProfiler (Raudvere *et al.*, 2019)
480 https://biit.cs.ut.ee/gprofiler/gost, with the input to DESeq2 (a list of 19,983 genes) as
481 background. Significant GO terms were shortlisted if fold-enrichment >= 2 & Bonferroni
482 corrected p-value < 0.05. Query genesets were (1) the list of 584 DEGs between *ltp2*-L and
483 *ltp2*-S and (2) the list of 1164 DEGs unique to the comparison Col-S and *ltp2*-S.
484
(6) Cuticle genes From the dataset in Li-Beisson et al.,(2013) we selected only genes with
annotated roles in cutin or wax biosynthesis and/or deposition (groups Cuticle Synthesis &
Transport 1, Fatty Acid Elongation & Wax Biosynthesis and Fatty Acid Synthesis). The curated
list contained 224 genes (Table S6), of which 160 were present in our RNA-seq dataset (Table
S6), and 21 were differentially expressed between Col-0 and *ltp2* (union set between Col-L vs
ltp2-L and Col-S vs *ltp2*-S, 1717 DEGs); the hypergeometric test indicates this is a modest over-
representation (p= 0.019).

<u>Protein sequence tree of PR-14/LTP Type I proteins</u> Protein sequences for PR-14/Type I LTP
proteins (Arondel12 *et al.*, 2000) were retrieved from NCBI: NP_181388.1 LTP1, NP_181387.1
LTP2, NP_568905.1 LTP3, NP_568904.1 LTP4, NP_190728.1 LTP5, NP_187489.1 LTP6,
NP_973466.1 LTP7, NP_179428.1 LTP8, NP_179135.2 LTP9, NP_195807.1 LTP10,
NP_680758.3 LTP11, NP_190727.1 LTP12, NP_001078707.1 LTP13, NP_001078780.1
LTP14, NP_192593.3 LTP15. Multiple sequence alignment was done with ClustalW in the msa

package (Bodenhofer *et al.*, 2015) and the neighbor-joining tree with the ape package (Paradis
et al., 2004).

LTP1 and *LTP2* global expression pattern We used the Digital Expression Explorer 2 repository
(Ziemann *et al.*, 2019) https://dee2.io/ to retrieve uniformly processed RNA-seq data from
Arabidopsis thaliana. We curated a dataset of 56 samples, spanning several organs, contexts
and developmental stages (Table S8), and kept only genes with at least 3 counts in one sample
out of the 56 (26,372 genes x 56 samples).

DATA ACCESS

ACKNOWLEDGEMENTS

This work was supported by the National Science Foundation (NSF MCB-1242744,
RESEARCH-PGR grant 1748843, PlantSynBio 2240888 to C.Q.) and the National Institute of
Health (NIH NHGRI 5RM1HG010461, NIH NIGMS R35GM139532 to C.Q.)

Abley K, **Formosa-Jordan P**, **Tavares H**, **Chan EY**, **Afsharinafar M**, **Leyser O**, **Locke JC**.
2021. An ABA-GA bistable switch can account for natural variation in the variability of
Arabidopsis seed germination time. *eLife* **10**.

Ansel J, **Bottin H**, **Rodriguez-Beltran C**, **Damon C**, **Nagarajan M**, **Fehrmann S**, **François J**,
Yvert G. **2008**. Cell-to-cell stochastic variation in gene expression is a complex genetic trait.
PLoS genetics **4**: e1000049.

Arondel V V, **Vergnolle C**, **Cantrel C**, **Kader J**. **2000**. Lipid transfer proteins are encoded by a
small multigene family in Arabidopsis thaliana. *Plant science: an international journal of*
experimental plant biology **157**: 1–12.

Ayroles JF, **Buchanan SM**, **O'Leary C**, **Skutt-Kakaria K**, **Grenier JK**, **Clark AG**, **Hartl DL**, **de**
Bivort BL. **2015**. Behavioral idiosyncrasy reveals genetic control of phenotypic variability.
Proceedings of the National Academy of Sciences of the United States of America **112**: 6706–
6711.

Blake WJ, **KAErn M**, **Cantor CR**, **Collins JJ**. **2003**. Noise in eukaryotic gene expression.
Nature **422**: 633–637.

Bodenhofer U, **Bonatesta E**, **Horejš-Kainrath C**, **Hochreiter S**. **2015**. msa: an R package for
multiple sequence alignment. *Bioinformatics (Oxford, England)* **31**: 3997–3999.

Borevitz JO, **Maloof JN**, **Lutes J**, **Dabi T**, **Redfern JL**, **Trainer GT**, **Werner JD**, **Asami T**,
Berry CC, **Weigel D**, *et al.* **2002**. Quantitative trait loci controlling light and hormone response in
two accessions of Arabidopsis thaliana. *Genetics* **160**: 683–696.

Buhot N, **Douliez JP**, **Jacquemard A**, **Marion D**, **Tran V**, **Maume BF**, **Milat ML**, **Ponchet M**,
Mikès V, **Kader JC**, *et al.* **2001**. A lipid transfer protein binds to a receptor involved in the
control of plant defence responses. *FEBS letters* **509**: 27–30.

Burga A, **Casanueva MO**, **Lehner B**. **2011**. Predicting mutation outcome from early stochastic
variation in genetic interaction partners. *Nature* **480**: 250–253.

Cameron KD, **Moskal WA**, **Smart LB**. **2006**. A second member of the Nicotiana glauca lipid
transfer protein gene family, NgLTP2, encodes a divergent and differentially expressed protein.
Functional plant biology: FPB **33**: 141–152.

Casanueva MO, **Burga A**, **Lehner B**. **2012**. Fitness trade-offs and environmentally induced
mutation buffering in isogenic C. elegans. *Science* **335**: 82–85.

Chae K, **Gonong BJ**, **Kim S-C**, **Kieslich CA**, **Morikis D**, **Balasubramanian S**, **Lord EM**. **2010**.
A multifaceted study of stigma/style cysteine-rich adhesin (SCA)-like Arabidopsis lipid transfer
proteins (LTPs) suggests diversified roles for these LTPs in plant growth and reproduction.
Journal of experimental botany **61**: 4277–4290.

Chae K, **Kieslich CA**, **Morikis D**, **Kim S-C**, **Lord EM**. **2009**. A gain-of-function mutation of
Arabidopsis lipid transfer protein 5 disturbs pollen tube tip growth and fertilization. *The plant cell*
21: 3902–3914.

Debono A, **Yeats TH**, **Rose JKC**, **Bird D**, **Jetter R**, **Kunst L**, **Samuels L**. **2009**. Arabidopsis
LTPG is a glycosylphosphatidylinositol-anchored lipid transfer protein required for export of
lipids to the plant surface. *The plant cell* **21**: 1230–1238.

Delvigne F, **Goffin P**. **2014**. Microbial heterogeneity affects bioprocess robustness: dynamic
single-cell analysis contributes to understanding of microbial populations. *Biotechnology journal*
9: 61–72.

Eagle H, **Levine EM**. **1967**. Growth regulatory effects of cellular interaction. *Nature* **213**: 1102–
1106.

Eldar A, **Chary VK**, **Xenopoulos P**, **Fontes ME**, **Losón OC**, **Dworkin J**, **Piggot PJ**, **Elowitz**
MB. **2009**. Partial penetrance facilitates developmental evolution in bacteria. *Nature* **460**: 510–
514.

Elowitz MB, **Levine AJ**, **Siggia ED**, **Swain PS**. **2002**. Stochastic gene expression in a single
cell. *Science (New York, N.Y.)* **297**: 1183–1186.

Emert BL, **Cote CJ**, **Torre EA**, **Dardani IP**, **Jiang CL**, **Jain N**, **Shaffer SM**, **Raj A**. **2021**.
Variability within rare cell states enables multiple paths toward drug resistance. *Nature*
biotechnology **39**: 865–876.

Feinerman O, **Veiga J**, **Dorfman JR**, **Germain RN**, **Altan-Bonnet G**. **2008**. Variability and
robustness in T cell activation from regulated heterogeneity in protein levels. *Science (New*
York, N.Y.) **321**: 1081–1084.

Finch-Savage WE, **Bassel GW**. **2016**. Seed vigour and crop establishment: extending
performance beyond adaptation. *Journal of experimental botany* **67**: 567–591.

Finkina EI, **Melnikova DN**, **Bogdanov IV**, **Ovchinnikova TV**. **2016**. Lipid transfer proteins as
components of the plant innate immune system: Structure, functions, and applications. *Acta*
naturae **8**: 47–61.

Gao S, **Guo W**, **Feng W**, **Liu L**, **Song X**, **Chen J**, **Hou W**, **Zhu H**, **Tang S**, **Hu J**. **2016**. LTP3

contributes to disease susceptibility in Arabidopsis by enhancing abscisic acid (ABA)
biosynthesis. *Molecular plant pathology* **17**: 412–426.

Geldner N. **2013**. The endodermis. *Annual review of plant biology* **64**: 531–558.

Gendreau E, **Traas J**, **Desnos T**, **Grandjean O**, **Caboche M**, **Höfte H**. **1997**. Cellular basis of
hypocotyl growth in Arabidopsis thaliana. *Plant physiology* **114**: 295–305.

Gimelbrant A, **Hutchinson JN**, **Thompson BR**, **Chess A**. **2007**. Widespread monoallelic
expression on human autosomes. *Science (New York, N.Y.)* **318**: 1136–1140.

Hall MC, **Dworkin I**, **Ungerer MC**, **Purugganan M**. **2007**. Genetics of microenvironmental
canalization in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences of the*
United States of America **104**: 13717–13722.

Hartwell LH, **Culotti J**, **Pringle JR**, **Reid BJ**. **1974**. Genetic control of the cell division cycle in
yeast. *Science (New York, N.Y.)* **183**: 46–51.

Heerden JHV, **Wortel MT**, **Bruggeman FJ**, **Heijnen JJ**, **Bollen YJ**, **Planqué R**, **Hulshof J**,
O'Toole TG, **Wahl SA**, **Teusink B**. **2014**. Fatal attraction in glycolysis: how Saccharomyces
cerevisiae manages sudden transitions to high glucose. *Microbial cell (Graz, Austria)* **1**: 103–
106.

Hill WG, **Mulder HA**, **Zhang X-S**. **2007**. The quantitative genetics of phenotypic variation in
animals. *Acta agriculturae Scandinavica. Section A, Animal science* **57**: 175–182.

Hong L, **Dumond M**, **Tsugawa S**, **Sapala A**, **Routier-Kierzkowska A-L**, **Zhou Y**, **Chen C**,
Kiss A, **Zhu M**, **Hamant O**, *et al.* **2016**. Variable cell growth yields reproducible organ
development through spatiotemporal averaging. *Developmental cell* **38**: 15–32.

Illouz-Eliaz N, **Ramon U**, **Shohat H**, **Blum S**, **Livne S**, **Mendelson D**, **Weiss D**. **2019**. Multiple
gibberellin receptors contribute to phenotypic stability under changing environments. *The plant*
cell **31**: 1506–1519.

Irshad M, **Canut H**, **Borderies G**, **Pont-Lezica R**, **Jamet E**. **2008**. A new picture of cell wall
protein dynamics in elongating cells of Arabidopsis thaliana: confirmed actors and newcomers.
BMC plant biology **8**: 94.

Jacq A, **Pernot C**, **Martinez Y**, **Domergue F**, **Payré B**, **Jamet E**, **Burlat V**, **Pacquit VB**. **2017**.
The Arabidopsis Lipid Transfer Protein 2 (AtLTP2) Is Involved in Cuticle-Cell Wall Interface
Integrity and in Etiolated Hypocotyl Permeability. *Frontiers in plant science* **8**: 263.

Jimenez-Gomez JM, **Corwin JA**, **Joseph B**, **Maloof JN**, **Kliebenstein DJ**. **2011**. Genomic
analysis of QTLs and genes altering natural variation in stochastic noise. *PLoS genetics* **7**:
e1002295.

Joseph B, **Corwin JA**, **Kliebenstein DJ**. **2015**. Genetic variation in the nuclear and organellar
genomes modulates stochastic variation in the metabolome, growth, and defense. *PLoS*
genetics **11**: e1004779.

Kader J-C. **1996**. Lipid-transfer proteins in plants. *Annual review of plant physiology and plant*
molecular biology **47**: 627–654.

Kalve S, **Sizani BL**, **Markakis MN**, **Helsmoortel C**, **Vandeweyer G**, **Laukens K**, **Sommen M**,
Naulaerts S, **Vissenberg K**, **Prinsen E**, *et al.* **2020**. Osmotic stress inhibits leaf growth of
Arabidopsis thaliana by enhancing ARF-mediated auxin responses. *The new phytologist* **226**:
1766–1780.

Kanno T, **Mette MF**, **Kreil DP**, **Aufsatz W**, **Matzke M**, **Matzke AJM**. **2004**. Involvement of
putative SNF2 chromatin remodeling protein DRD1 in RNA-directed DNA methylation. *Current*
biology: CB **14**: 801–805.

Karahara I. **2012**. The pea stem: a unique experimental system to study the development of the
Casparian strip. *Plant signaling & behavior* **7**: 1182–1184.

Katsanos D, **Koneru SL**, **Mestek Boukhibar L**, **Gritti N**, **Ghose R**, **Appleford PJ**, **Doitsidou**
M, **Woollard A**, **van Zon JS**, **Poole RJ**, *et al.* **2017**. Stochastic loss and gain of symmetric
divisions in the C. elegans epidermis perturbs robustness of stem cell number. *PLoS biology* **15**:
e2002429.

Lachowiec J, **Mason GA**, **Schultz K**, **Queitsch C**. **2018**. Redundancy, Feedback, and
Robustness in the Arabidopsis thaliana BZR/BEH Gene Family. *Frontiers in genetics* **9**: 523.

Lachowiec J, **Queitsch C**, **Kliebenstein DJ**. **2016**. Molecular mechanisms governing
differential robustness of development and environmental responses in plants. *Annals of botany*
117: 795–809.

Lemus T, **Mason GA**, **Bubb KL**, **Alexandre CM**, **Queitsch C**, **Cuperus JT**. **2023**. AGO1 and
HSP90 buffer different genetic variants in Arabidopsis thaliana. *Genetics* **223**.

Li-Beisson Y, **Shorrosh B**, **Beisson F**, **Andersson MX**, **Arondel V**, **Bates PD**, **Baud S**, **Bird**
D, **DeBono A**, **Durrett TP**, *et al.* **2013**. Acyl-lipid metabolism. *The Arabidopsis book* **11**: e0161.

Lomvardas S, **Barnea G**, **Pisapia DJ**, **Mendelsohn M**, **Kirkland J**, **Axel R**. **2006**.
Interchromosomal interactions and olfactory receptor choice. *Cell* **126**: 403–413.

Maldonado AM, **Doerner P**, **Dixon RA**, **Lamb CJ**, **Cameron RK**. **2002**. A putative lipid transfer
protein involved in systemic resistance signalling in Arabidopsis. *Nature* **419**: 399–403.

Maloof JN, **Borevitz JO**, **Dabi T**, **Lutes J**, **Nehring RB**, **Redfern JL**, **Trainer GT**, **Wilson JM**,
Asami T, **Berry CC**, *et al.* **2001**. Natural variation in light sensitivity of Arabidopsis. *Nature*
genetics **29**: 441–446.

Márquez-Jurado S, **Díaz-Colunga J**, **das Neves RP**, **Martinez-Lorente A**, **Almazán F**,
Guantes R, **Iborra FJ**. **2018**. Mitochondrial levels determine variability in cell death by
modulating apoptotic gene expression. *Nature communications* **9**: 389.

Mason GA, **Lemus T**, **Queitsch C**. **2016**. The mechanistic underpinnings of an ago1-mediated,
environmentally dependent, and stochastic phenotype. *Plant physiology* **170**: 2420–2431.

Miedes E, **Suslov D**, **Vandenbussche F**, **Kenobi K**, **Ivakov A**, **Van Der Straeten D**, **Lorences**
EP, **Mellerowicz EJ**, **Verbelen J-P**, **Vissenberg K**. **2013**. Xyloglucan
endotransglucosylase/hydrolase (XTH) overexpression affects growth and cell wall mechanics
in etiolated Arabidopsis hypocotyls. *Journal of experimental botany* **64**: 2481–2497.

Molina A, **García-Olmedo F**. **1993**. Developmental and pathogen-induced expression of three

barley genes encoding lipid transfer proteins. *The Plant journal: for cell and molecular biology* **4**:
983–991.

Nieuwland J, **Feron R**, **Huisman BAH**, **Fasolino A**, **Hilbers CW**, **Derksen J**, **Mariani C**. **2005**.
Lipid transfer proteins enhance cell wall extension in tobacco. *The plant cell* **17**: 2009–2019.

Paradis E, **Claude J**, **Strimmer K**. **2004**. APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics (Oxford, England)* **20**: 289–290.

Park J-E, **Kim Y-S**, **Yoon H-K**, **Park C-M**. **2007**. Functional characterization of a small auxin-up
RNA gene in apical hook development in Arabidopsis. *Plant science: an international journal of*
experimental plant biology **172**: 150–157.

Perez MF, **Francesconi M**, **Hidalgo-Carcedo C**, **Lehner B**. **2017**. Maternal age generates
phenotypic variation in Caenorhabditis elegans. *Nature* **552**: 106–109.

Potocka I, **Baldwin TC**, **Kurczynska EU**. **2012**. Distribution of lipid transfer protein 1 (LTP1)
epitopes associated with morphogenic events during somatic embryogenesis of Arabidopsis
thaliana. *Plant cell reports* **31**: 2031–2045.

Queitsch C, **Sangster TA**, **Lindquist S**. **2002**. Hsp90 as a capacitor of phenotypic variation.
Nature **417**: 618–624.

Rai MI, **Wang X**, **Thibault DM**, **Kim HJ**, **Bombyk MM**, **Binder BM**, **Shakeel SN**, **Schaller GE**.
2015. The ARGOS gene family functions in a negative feedback loop to desensitize plants to
ethylene. *BMC plant biology* **15**: 157.

Raj A, **Rifkin SA**, **Andersen E**, **van Oudenaarden A**. **2010**. Variability in gene expression
underlies incomplete penetrance. *Nature* **463**: 913–918.

Raser JM, **O'Shea EK**. **2004**. Control of stochasticity in eukaryotic gene expression. *Science*
(New York, N.Y.) **304**: 1811–1814.

Raudvere U, **Kolberg L**, **Kuzmin I**, **Arak T**, **Adler P**, **Peterson H**, **Vilo J**. **2019**. g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists (2019 update).
Nucleic acids research **47**: W191–W198.

Reed JW, **Elumalai RP**, **Chory J**. **1998**. Suppressors of an Arabidopsis thaliana phyB mutation
identify genes that control light signaling and hypocotyl elongation. *Genetics* **148**: 1295–1310.

Ros M, **Sorensen D**, **Waagepetersen R**, **Dupont-Nivet M**, **SanCristobal M**, **Bonnet J-C**,
Mallard J. **2004**. Evidence for genetic control of adult weight plasticity in the snail Helix aspersa.
Genetics **168**: 2089–2097.

Sangster TA, **Salathia N**, **Lee HN**, **Watanabe E**, **Schellenberg K**, **Morneau K**, **Wang H**,
Undurraga S, **Queitsch C**, **Lindquist S**. **2008**. HSP90-buffered genetic variation is common in
Arabidopsis thaliana. *Proceedings of the National Academy of Sciences* **105**: 2969–2974.

Sawa S, **Ohgishi M**, **Goda H**, **Higuchi K**, **Shimada Y**, **Yoshida S**, **Koshiba T**. **2002**. The
HAT2 gene, a member of the HD-Zip gene family, isolated as an auxin inducible gene by DNA
microarray screening, affects auxin response in Arabidopsis. *The Plant journal: for cell and*
molecular biology **32**: 1011–1022.

Shaffer SM, **Dunagin MC**, **Torborg SR**, **Torre EA**, **Emert B**, **Krepler C**, **Beqiri M**, **Sproesser**
K, **Brafford PA**, **Xiao M**, *et al.* **2017**. Rare cell variability and drug-induced reprogramming as a
mode of cancer drug resistance. *Nature* **546**: 431–435.

Sharma SV, **Lee DY**, **Li B**, **Quinlan MP**, **Takahashi F**, **Maheswaran S**, **McDermott U**, **Azizian**
N, **Zou L**, **Fischbach MA**, *et al.* **2010**. A chromatin-mediated reversible drug-tolerant state in
cancer cell subpopulations. *Cell* **141**: 69–80.

Shen X, **Pettersson M**, **Rönnegård L**, **Carlborg Ö**. **2012**. Inheritance beyond plain heritability:
variance-controlling genes in Arabidopsis thaliana. *PLoS genetics* **8**: e1002839.

Smith MCA, **Sumner ER**, **Avery SV**. **2007**. Glutathione and Gts1p drive beneficial variability in
the cadmium resistances of individual yeast cells. *Molecular microbiology* **66**: 699–712.

Snijder B, **Sacher R**, **Rämö P**, **Damm E-M**, **Liberali P**, **Pelkmans L**. **2009**. Population context
determines cell-to-cell variability in endocytosis and virus infection. *Nature* **461**: 520–523.

Spartz AK, **Lee SH**, **Wenger JP**, **Gonzalez N**, **Itoh H**, **Inzé D**, **Peer WA**, **Murphy AS**,
Overvoorde PJ, **Gray WM**. **2012**. The SAUR19 subfamily of SMALL AUXIN UP RNA genes
promote cell expansion. *The Plant journal: for cell and molecular biology* **70**: 978–990.

Tanaka T, **Tanaka H**, **Machida C**, **Watanabe M**, **Machida Y**. **2004**. A new method for rapid
visualization of defects in leaf cuticle reveals five intrinsic patterns of surface defects in
Arabidopsis. *The Plant journal: for cell and molecular biology* **37**: 139–146.

Tian Q, **Uhlir NJ**, **Reed JW**. **2002**. Arabidopsis SHY2/IAA3 inhibits auxin-regulated gene
expression. *The plant cell* **14**: 301–319.

Tran DT, **Hertog MLATM**, **Tran TLH**, **Quyen NT**, **Van de Poel B**, **Mata CI**, **Nicolaï BM**. **2017**.
Population modeling approach to optimize crop harvest strategy. The case of field tomato.
Frontiers in plant science **8**: 608.

Vandenbussche F, **Verbelen J-P**, **Van Der Straeten D**. **2005**. Of light and length: regulation of
hypocotyl growth in Arabidopsis. *BioEssays: news and reviews in molecular, cellular and*
developmental biology **27**: 275–284.

Volfson D, **Marciniak J**, **Blake WJ**, **Ostroff N**, **Tsimring LS**, **Hasty J**. **2006**. Origins of
extrinsic variability in eukaryotic gene expression. *Nature* **439**: 861–864.

Waddington CH. **1942**. Canalization of development and the inheritance of acquired
characters. *Nature* **150**: 563–565.

Whitlock MC, **Fowler K**. **1999**. The changes in genetic and environmental variance with
inbreeding in Drosophila melanogaster. *Genetics* **152**: 345–353.

Wright S. **1920**. The relative importance of heredity and environment in determining the piebald
pattern of guinea-pigs. *Proceedings of the National Academy of Sciences of the United States*
of America **6**: 320–332.

Xiao Y, **Bowen CH**, **Liu D**, **Zhang F**. **2016**. Exploiting nongenetic cell-to-cell variation for
enhanced biosynthesis. *Nature chemical biology* **12**: 339–344.

Xiao C, **Somerville C**, **Anderson CT**. **2014**. POLYGALACTURONASE INVOLVED IN

EXPANSION1 functions in cell elongation and flower development in Arabidopsis. *The Plant cell*
26: 1018–1035.

Yeats TH, **Rose JKC**. **2008**. The biochemistry and biology of extracellular plant lipid-transfer
proteins (LTPs). *Protein science: a publication of the Protein Society* **17**: 191–198.

Zabinsky RA, **Mason GA**, **Queitsch C**, **Jarosz DF**. **2019**. It's not magic – Hsp90 and its effects
on genetic and epigenetic variation. *Seminars in cell & developmental biology* **88**: 21–35.

Zwiewka M, **Nodzyński T**, **Robert S**, **Vanneste S**, **Friml J**. **2015**. Osmotic stress modulates
the balance between exocytosis and clathrin-mediated endocytosis in Arabidopsis thaliana.
Molecular plant **8**: 1175–1187.

FIGURE LEGENDS

Figure 1. *LTP2* **hypomorphs increase phenotypic variation in skotomorphogenesis traits. A**.
Approximate location of the T-DNA insertion (LB) relative to the *LTP2* gene in *ltp2-1* and *ltp2-2*
lines. **B**. *LTP2* relative expression in the shoots of wild-type (Col-0), *ltp2-1* and *ltp2-2* seedlings
grown in the dark for 7 days (d) after stratification. In parenthesis are the % of Col-0 *LTP2*
transcript levels detected in the *ltp2* lines. **C**. Representative image of Col-0 (top), *ltp2-1*
(middle) and *ltp2-2* (bottom) dark-grown seedlings at 7 days after stratification; the roots were
trimmed. **D**. Density lines of the distributions of hypocotyl length scored at 7 days post-
stratification in Col-0 and *ltp2* seedlings grown in MS media with 1% sucrose; each line
represents the merged distribution of 5 biological replicates (n=78-210 each) per genotype. **E**.
Coefficient of variation (sd/mean) in hypocotyl length, a measure of variation, for (1) 5
biological replicates (open circles) and (2) merged for all 5 replicates (filled circle, same data as
in **D**). **F**. Hypocotyl negative gravitropism measured as the deviation from vertical (in degrees)
of the hypocotyl apex in 7d dark-grown seedlings grown on MS+1% sucrose. The density
distribution of measured angles for all genotypes is plotted. **G**. Comparison between the
hypocotyl length of individual *ltp2* parents (5 long and 5 short) and the mean hypocotyl length of
their offspring; all lengths measured at 7d post-stratification on seedlings grown on MS+1%
sucrose. **H**. Beanplots of hypocotyl (left side) and root (right side) length from the merged
dataset used in **D** and **E**. **I**. Percentage of dark-grown seedlings with an open cotyledon
phenotype, in wild-type (Col-0), *ltp2-1* and *ltp2-2* genotypes, at 7 days after stratification on
MS+ 1% sucrose. Two replicates are shown (n=79-208).

Figure 2. Sucrose-dependent loss of cuticle integrity in *ltp2* **is associated with short**
hypocotyls. **A**. Density lines of the distribution of hypocotyl length for seedlings grown on
plates containing MS medium, MS+1% glucose or MS+1% sucrose for 7 days post-stratification.
Shown are the merged distributions of two biological replicates per condition/per genotype. **B**.
Coefficient of variation in hypocotyl length for the distributions shown in **A**. **C**. Representative
shoots of 7d dark-grown seedlings grown on plates containing MS medium, MS+1% glucose or
MS+1% sucrose. Hypocotyl negative gravitropism is largely rescued in *ltp2* seedlings grown
without sucrose. **D**. Boxplots comparing the hypocotyl/root ratio for every seedling in **A** across

different growth media. **E**. Magnified images of the epidermal surface of Col-0 and *ltp2-1*
hypocotyls at 7d post-stratification when grown with (left) or without (right) 1% sucrose. **F**.
Comparing the effects of 1% sucrose on hypocotyl cuticle permeability: dark-grown seedlings of
Col-0 and *ltp2-1* were stained with toluidine blue after growing for 5 days on MS+1% sucrose
(left) or 7 days on MS alone (right). **G**. Relationship between hypocotyl length at 7 days after
stratification on MS+1% sucrose and toluidine blue staining coverage expressed in % of
hypocotyl length. Filled dots represent seedlings with the wild-type developmental pattern of
hypocotyl length > root length.

Figure 3. Hypocotyl length is associated with many transcriptional differences in *ltp2* but
not Col-0 seedlings, mostly related to stress, defense and growth. A. Experimental setup
for collecting RNA-seq samples. For Col-0 and *ltp2-1*, the first boxplot shows the distribution of
hypocotyl length from one out of seven hypocotyl assays done per replicate; the second set of
four boxplots (two replicates with short and two replicates with long hypocotyls) shows the
actual distribution of hypocotyl lengths (n=70) from the selected individuals used to generate
RNA-seq libraries. **B**. Biplot of Principal Component Analysis showing the first two PCs. The
percentage of the total variance explained by PC1 and PC2 is indicated on the top right corner.
C. Correlation between mean hypocotyl length for the eight RNA-seq samples and PC1
loadings; shown on the bottom left corner is the Spearman rho. **D**. Shortlisted GO enrichments
(fold-enrichment >=2 and adjusted p-value < 0.05) for the set of 584 differentially expressed
genes between *ltp2*-L and *ltp2*-S, split by down- and upregulated genes. Shown are the
log10(fold-enrichment) and the number of DEGs per enriched term. **E-F** Row-scaled heatmap
visualizations of a subset of DEGs downregulated (**E**) or upregulated (**F**) in *ltp2*-S relative to
ltp2-L. Pink dots indicate whether each gene was also a DEG on other pairwise-comparisons.

Figure 4. The *ltp2* phenotype depends on the expression ratio of *LTP2* and its close
paralog *LTP1* A. Row-scaled heatmap visualization of cuticle-related genes differentially
expressed between Col-0 and *ltp2* samples. **B**. Detail of a tree depicting amino acid sequence
similarity among PR14/Type I LTP proteins (full tree in Figure S5A). **C**. Relative expression of
LTP1 and *LTP2* in the shoots of light or dark-grown (gray box) Col-0 and *ltp2* seedlings grown
for 7 days after stratification on MS+ 1% sucrose. **D**. Comparison of the *LTP1/LTP2* expression
ratio between Col-0, *ltp2-1* and an *ltp1ltp2* double mutant. **E-G** Hypocotyl length distribution (**E**),
coefficient of variation of hypocotyl length (**F**) and (**G**) hypocotyl/root ratio for Col-0, *ltp2-1* and
an *ltp1 ltp2* double knockdown; all seedlings were grown in the dark for 7d after stratification on
MS+ 1% sucrose. **H**. Representative images of toluidine blue stained etiolated hypocotyls from
Col-0, *ltp2-1* and an *ltp1 ltp2* double mutant grown for 7 days after stratification on MS+ 1%
sucrose
805
Figure S1. A. *LTP2* is induced in the dark. Shown is the relative expression of *LTP2* in light vs
dark-grown shoots at 7d post-stratification (MS+1% sucrose). B. Boxplot of log10(counts+1) for
19,987 genes expressed in our RNA-seq dataset from dark-grown shoots at 7d post-
stratification; the black dot indicates *LTP2*. In Col-0 wild type, *LTP2* was the 7th most highly
expressed gene. **C**. Scatterplot of hypocotyl length vs orientation for each individual, measured
at 7d post-stratification (MS+1% sucrose); little correlation was found; indicated is Spearman ρ
and p-value.

Figure S2. A. Boxplots of hypocotyl length, at 7d post-stratification on MS+1% sucrose, for all 5
replicates from Figure 1, showing similar distributions. Replicates 1-3 are from different seed
batches, replicates 4-5 are the same seed batch done on different days. **B**. (top) Time-
dependent decrease in the Pearson correlation coefficient between hypocotyl and root length at
3, 4, 5 and 7d post-stratification (MS+1% sucrose). Values are similar for *ltp2* and Col-0
supporting similar timing and uniformity of germination in *ltp2*. **C**. Scatterplot of hypocotyl length
vs seed area (arbitrary units), showing little correlation; indicated is the Spearman ρ and p-
value.

Figure S3. A. Hypocotyl (top) and root length (bottom) as a function of time since stratification
in Col-0 seedlings grown on MS (open circles) or MS+1%sucrose (filled circles). B. Boxplots
comparing the distribution of individual hypocotyl/root ratios for Col-0 seedlings grown on MS
(open circles) and MS+1%sucrose (filled circles). C. Boxplots of hypocotyl length for seedlings
grown on MS+1% sucrose on either vertical plates (shoot in contact with the media) or
horizontal plates (only the roots contact the media); shows that direct sucrose uptake from the
shoot is involved in sucrose-induced hypocotyl elongation.

Figure S4. Mannitol improves hypocotyl elongation of sucrose-grown *ltp2* seedlings. **A.**
Density lines of the distributions of hypocotyl length for Col-0 (grey), *ltp2-1* (blue) and *ltp2-2*
(green) seedlings grown on either MS+1% sucrose or MS+1%sucrose+29mM mannitol for 7
days post-stratification.The x- and y-axis range is the same in all graphs. **B**. Dosage-
response curves of mean hypocotyl length (y-axis) at 7d post-stratification relative to
increasing amounts of mannitol (0-150mM). **C**. Coefficient of variation of hypocotyl length
for the distributions shown in **A. D.** Boxplots showing the hypocotyl/root ratio of every
seedling shown in **A.**

Figure S5. *ltp2* hypocotyls show wild-type-like sucrose-induced responses. **A.** Hypocotyl length
as a function of time since stratification in Col-0, *ltp2-1* and *ltp2-2* seedlings grown either on MS
(open circles) or on MS+ 1%sucrose (filled circles). **B.** Boxplots comparing the distribution of
individual hypocotyl/root ratios for Col-0, *ltp2-1* and *ltp2-2* seedlings grown on MS (positions 1, 3
and 5) and MS+1%sucrose (positions 2, 4 and 6). **C.** Lugol staining of Col-0 (WT), *ltp2-1* and
ltp2-2 etiolated seedlings grown for 7d after stratification on MS (left), MS+1%sucrose (center)
or MS+1%glucose. Seedlings were stained for 2min with undiluted lugol solution, rinsed, and
left in distilled H$_2$O for ~1hr until imaging. **D.** Boxplots show the distribution of hypocotyl lengths
for etiolated Col-0 (WT) and *ltp2-1* seedlings grown for 7d after stratification on either MS+1%
sucrose (left, no fill) or MS+2% sucrose (right, filled). Inset (top, right) shows the mean per
genotype and condition, and the ratio of the means (2% suc /1% suc) in each genotype.

Figure S6. A. Similar numbers of reads (>40 Mio) were obtained for each of the 8 RNA-seq
samples. **B**. **left:** Correlation matrix for all 8 x 8 sample comparisons; **right:** Pearson correlation
coefficients between replicates are all > 0.98 (see top right corner). **C**. Boxplots of log10(counts)
for all 8 samples showing no major difference in count distributions between Col-0 and *ltp2*. **D**.
(left) Venn-diagram showing the overlap in DEGs (black; 11) between *ltp2*-L vs *ltp2*-S and Col-L

vs Col-S; (right) Fold-change and p-value for the 10 DEGs common to long x short comparisons
in Col-0 and *ltp2*. **E**. Boxplot of log10(counts) for all 19,983 genes in our dataset; orange circles:
10 genes from (**D**). **F**. Distance in log10 of the number of DEGs plotted against distance in
mean hypocotyl length (7d post-stratification, MS+ 1%sucrose) for each of four sample
pairwise-comparisons. **G**. Venn-diagram showing the overlap in DEGs (gray; 474) between the
comparison *ltp2*-L vs *ltp2*-S and Col-S vs *ltp2*-S. H. GO term enrichments (fold-enrichment >= 2,
p-adj < 0.05) for the set of DEGs unique to Col-S vs *ltp2*-S (1164; see **G**).
864
Figure S7. A. Tree depicting the aminoacid similarity between PR14/Type I LTP proteins. **B**.
Plot shows all the Type I LTPs expressed in our RNA-seq dataset of dark-grown shoots (x-axis)
and the contribution of each (in %; y-axis) to the subfamily total in these samples. For this
analysis, the counts from all four Col-0 samples and all four *ltp2* samples were averaged.
C. Plot shows the gene expression rank of *LTP2* and *LTP1* relative to all 26,372 genes, across
a curated dataset of 56 Arabidopsis RNA-seq samples (see Methods); samples are ordered
from highest to lowest; open circle is *LTP1*; filled circle is *LTP2*. In 41 out of 56 samples *LTP1*
was higher than *LTP2*, indicating that *LTP1* is the dominant paralog by expression level.
D. The *LTP1/LTP2* expression ratio is fairly constant across individuals, regardless of their
hypocotyl length; shown is the relation between hypocotyl length (at 7d post-stratification on
MS+1% sucrose) and the log10(*LTP1/LTP2*) expression ratio for individual shoots. **E**. The
LTP1/LTP2 expression ratio changes during development; dark-grown seedlings have the
lowest ratio compared to all other tested samples (light-grown seedlings, roots, leaves and
flowers).
879
Figure S8. *LTP1* is upregulated across individual *ltp2-1* seedlings. **A.** Scatterplot comparing
LTP1 (y-axis) with *LTP2* expression (x-axis) in individual etiolated shoots grown for 7d post
stratification on MS+1%sucrose. **B.** Stacked bar chart depicting the percentage of *LTP1*
expression (white) or *LTP2* expression (black) in individual etiolated shoots sorted by increasing
hypocotyl length (left to right) and harvested 7d post stratification. **C.** Scatterplot of the
relationship between hypocotyl length (y-axis) and *LTP1* expression (x-axis) for a set of wild-
type (Col-0, grey) and *ltp2-1* (blue) individual seedlings collected at 7d post stratification (only
shoots were used for RNA extraction). Dashed line denotes the line of regression for each
genotype. **D.** Scatterplot of the relationship between hypocotyl length (y-axis) and *LTP2*
expression (x-axis) for a set of wild-type (Col-0, grey) and *ltp2-1* (blue) individual seedlings
collected at 7d post stratification (only shoots were used for RNA extraction). Dashed line
denotes the line of regression for each genotype. **E.** Scatterplot of the relationship between
hypocotyl length (y-axis) and the log10 of the LTP1/LTP2 expression ratio (x-axis) in wild-type
(Col-0, gray, left) and ltp2-1 (blue, right); data are from the same individuals shown in **C** and **D**.
894
895 **Figure S9.** *LTP1* and *LTP2* expression is low in the double *ltp1 ltp2* seedlings. Shown are the
896 separate expression levels of *LTP1* (open circles) and *LTP2* (filled circles) used to calculate the
897 *LTP1*/*LTP2* ratio shown in **Figure 4D**. Each dot represents a bulk sample of about 30 dark-
898 grown shoots collected 7 days post stratification.
899

900   **Table S1-S4**. Differential expressed genes between Col-L and Col-S (Table S1), *ltp2*-L and *ltp2*-
901   S (Table S2), Col-L and *ltp2*-L (Table S3), Col-S and *ltp2*-L (Table S4).
902
903   **Table S5**. Differentially expressed auxin-responsive genes not shared between Col-S vs *ltp2*-S
904   and *ltp2*-L vs *ltp2*-S.
905
906   **Table S6**. Cuticle-related genes from Li-Beisson et al., 2013 expressed in our RNA-seq dataset.
907
908   **Table S7**. List of primers used in genotyping and qRT-PCR.
909   **Table S8**. List of publicly available RNA-seq samples used to establish global expression
910   patterns for *LTP1* and *LTP2*.
911

A

ltp2-1   ltp2-2   **LTP2**/AT2G38530

T-DNA

-370bps   -116bps   TSS Chr2:16,128,378

B

log10 (*LTP2*)

○ Col-0, WT (100%)
○ *ltp2-1* (~24%)
○ *ltp2-2* (~1%)

n = 2-3 pools of ~ 30 shoots each

C

Col-0

ltp2-1

ltp2-2

D

density

n= 711-717
5 replicates
merged

— Col-0
— *ltp2-1*
— *ltp2-2*

hypocotyl length (mm)

E

coefficient of variation (CV) of hypocotyl length

○ Col-0
○ *ltp2-1*
○ *ltp2-2*
●●● merged

n= 5 replicates
each 78-210
hypocotyls

F

density

— Col-0
— *ltp2-1*
— *ltp2-2*
n = 147-177

+   -

degrees off vertical

G

parent hypocotyl length (mm)

○ *ltp2-1*
○ *ltp2-2*

offspring mean hypocotyl length (mm)

n offspring = 69-70 hypocotyls
n parents = 5 short, 5 long

H

length (mm)

hypocotyl   root

Col-0   *ltp2-1*   *ltp2-2*

n= 711-717,
5 replicates merged

I

percentage of open cotyledons

○ Col-0
○ *ltp2-1*
○ *ltp2-2*
n= 79-208
2 replicates

Figure 1

Figure 2

Figure 3

A

samples →

cuticle-related genes

GPAT2/AT1G02390
LACS3/AT1G64400
LTP2/AT2G38530
LTP1/AT2G38540
AT2G38995
LACS1/AT2G47240
LTP6/AT3G08770
CER26-like/AT3G23840
PES2/AT3G26840
FAR4/AT3G44540
AT3G53980
ABCG18/AT3G55110
CYP96A9/AT4G39480
LTP10/AT5G01870
KCS19/AT5G04530
WSD7/AT5G12420
PEL3/AT5G23940
CUS2/AT5G33370
LTP4.4/AT5G55450
LTP4/AT5G59310
LTP3/AT5G59320

down   up

-1 0 1
row Z-score

Col-L Col-S ltp2-L ltp2-S

B

LTP5
LTP7
LTP1
LTP2
LTP12
LTP3
LTP4

LTP2   LTP1
AT2G38530   AT2G38540

Chr2:16,128,377-16,131,174

C

LTP1   LTP2

log10(expression)

○ Col-0
○ ltp2-1
○ ltp2-2

n= 2 replicates
each dot ~ 30 shoots

light dark   light dark

D

○ Col-0
○ ltp2-1
○ ltp1 ltp2

log10(LTP1/LTP2)

n= 7-10 replicates
each dot ~ 30 shoots

E

density

── Col-0
── ltp2-1
── ltp1 ltp2

n= 73-100

hypocotyl length (mm)

F

○ Col-0
○ ltp2-1
○ ltp1 ltp2

coefficient of variation
of hypocotyl length

G

hypocotyl/ root

── Col-0
── ltp2-1
── ltp1 ltp2

H

Col-0   ltp2-1   ltp1 ltp2

Figure 4