**Conserving a threatened North American walnut: a chromosome-scale reference genome for butternut (*Juglans cinerea*)**

**Running head:** A reference genome for *Juglans cinerea*

**Authors:** Cristopher R. Guzman-Torres[*1,3], Emily Trybulec[*2,3], Hannah LeVasseur[*1,3], Harshita Akella[1,3], Maurice Amee[1,3], Emily Strickland[1], Nicole Pauloski[2,3], Martin Williams[4], Jeanne Romero-Severson[5], Sean Hoban[6], Keith Woeste[7], Carolyn C. Pike[8], Karl C. Fetter[**1,3], Cynthia N. Webster[**1,3], Michelle L. Neitzey[**2,3], Rachel J. O'Neill[**2,3], Jill L. Wegrzyn[**1,3]

[1]Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA 06269
[2]Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT, USA 06269
[3]Biodiversity and Conservation Genomics Center, Institute for Systems Genomics, University of Connecticut, Storrs, CT USA 06269
[4]Atlantic Forestry Center, Canadian Forest Service, Natural Resources Canada, Fredericton, NB, Canada
[5]Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA 46556
[6]The Center for Tree Science, The Morton Arboretum, Lisle, IL, USA 60532
[7]USDA Forest Service, Northern Research Station, West Lafayette, IN, USA 47906
[8]USDA Forest Service, Eastern Region State, Private and Tribal Forestry, West Lafayette, IN USA 47906

[*]First co-authors (undergraduate team)

[**]Corresponding co-authors (mentors):
Karl Fetter kcf@uconn.edu
Cynthia N. Webster cynthia.webster@uconn.edu
Michelle L. Neitzey michelle.neitzey@uconn.edu
Rachel J. O'Neill rachel.oneill@uconn.edu
Jill L. Wegrzyn jill.wegrzyn@uconn.edu

**Keywords:** reference genome, butternut, conservation genetics, long-read, nanopore, genome evolution, Juglandaceae, genome annotation, methylation, phylogeny

**Abstract**

With the advent of affordable and more accurate third generation sequencing
technologies and the associated bioinformatic tools, it is now possible to sequence, assemble,
and annotate more species of conservation concern than ever before. *Juglans cinerea*,
commonly known as butternut or white walnut, is a member of the walnut family, native to the
Eastern United States and Southeastern Canada. The species is currently listed as Endangered
on the IUCN Red List due to decline from an invasive fungus known as *Ophiognomonia*
*clavigignenti-juglandacearum* (Oc-j) that causes butternut canker. Oc-j creates visible sores on
the trunks of the tree which essentially starves and slowly kills the tree. Natural resistance to
this pathogen is rare. Conserving butternut is of utmost priority due to its critical ecosystem role
and cultural significance. As part of an integrated undergraduate and graduate student training
program in biodiversity and conservation genomics, the first reference genome for *Juglans*
*cinerea* is described here. This chromosome-scale 539 Mb assembly was generated from over
100X coverage of Oxford Nanopore long reads and scaffolded with the *Juglans mandshurica*
genome. Scaffolding with a closely related species oriented and ordered the sequences in a
manner more representative of the structure of the genome without altering the sequence.
Comparisons with sequenced Juglandaceae revealed high levels of synteny and further
supported *J. cinerea's* recent phylogenetic placement. Comparative assessment of gene family
evolution revealed a significant number of contracting families, including several associated with
biotic stress response.

## 1 **Introduction**

Temporal analysis of biodiversity loss indicates that we are in the midst of the Earth's
sixth mass extinction event (Ceballos et al. 2017). Habitat destruction and degradation,
overexploitation, pollution, invasive species, climate change, and other anthropogenic factors
are the greatest threats to Earth's flora and fauna. The International Union for the Conservation
of Nature (IUCN) Red List noted that nearly 30% of the species assessed are at risk of
extinction (IUCN 2022), including about 30% of trees.
Preserving adaptive potential through conservation genetics is widely recognized as
important to ensuring long-term survival, especially in the face of climate change (Kardos et al
2021). Conservation managers and geneticists can work together to monitor population genetic
diversity and develop restoration plans including assisted breeding. However, several studies
have acknowledged a "conservation genetics gap" stemming from, among other things, a lack of
training, barriers to collaboration, discipline-specific terminology, and access to resources
(Taylor et al. 2017; Sandstrom et al. 2019). The slow advancement of tool development and
implementation in conservation is further impacted by the lack of genomic resources for the vast
majority of threatened species, with less than 1% of all eukaryotic species sequenced to date
(Lewin et al. 2022). As such, the implementation of genetic tools in conservation and/or
restoration benefits from the development of a reference genome (DeWoody et al. 2022).
Butternut (*Juglans cinerea* L.) is a deciduous, outcrossing, and wind-pollinated canopy
tree native to eastern North America. Once valued for its wood and edible nuts, today, it is
threatened by butternut canker, a disease caused by the fungus *Ophiognomonia clavigignenti-*
*juglandacearum* (Nair et al. 1979), discovered in the late 1960s and now found throughout
butternut's range. The disease had already killed more than 80% of butternut trees in numerous
states by the 1990s (the last detailed censuses available) and is threatening the species with
extinction. Diseased trees typically die within 5 to 10 years, and little resistance has been
identified outside of hybrids with the introduced Japanese walnut (*Juglans ailantifolia Carrière*).
The search for butternut individuals or populations with less susceptibility to butternut canker is
confounded by environmental factors. For example, the severity of butternut canker as well as
Armillaria root rot is greater in low-elevation sites with heavy soils and in humid sites with heavy
canopies (LaBonte et al. 2015). Dramatic declines over the past few decades have resulted in
the listing of butternut as endangered in Canada (Nielsen et al. 2003) and Endangered on the
IUCN Red list in 2018. Several US states have also included butternut as protected or
endangered, including Minnesota and Wisconsin (Smith 2018; WDNR 2021).
Butternut shows high genetic diversity (moderate cline from south to north) (Hoban et al
2008; 2010). In more isolated populations, lower genetic diversity and inbreeding are observed.
Chloroplast markers have revealed population-level genetic differentiation across an east–west
gradient that may have originated from multiple Ice Age refugia (Laricchia et al 2015). Butternut
is sympatric with *Juglans nigra Linnaeus* through much of its range however it cannot hybridize
with this species, but it easily hybridizes with its close relative, the non-native *J. ailantifolia.*
Hybrids with *J. ailantifolia* are more tolerant of butternut canker, grow faster/larger, and are
more productive (Boraks and Broders 2014; Brennan et al. 2020). Hybrids between the species
occur at low frequencies in numerous forests in several states (Hoban et al. 2012). Given the
limited resistance observed among pure *J. cinerea* individuals, a plan proposed by practitioners

seeks to conserve native germplasm and introduce genes for resistance from *J. ailantifolia* (Pike
et al. 2020), similar to efforts in American chestnut (Fernandes et al. 2022). The development of
reliable markers and the materials for forward and/or reverse breeding approaches, and for
analyzing adaptive introgression of hybrids in the wild, will rely on a high quality reference
genome.

To address the conservation genetics gap, undergraduate and graduate students trained
in the newly formed Biodiversity and Conservation Genomics Center at the University of
Connecticut worked with practitioners to generate the first reference genome for the threatened
*J. cinerea*. This study represents the intersection of industry (Oxford Nanopore), federal
research agencies (USDA Forest Service, Canadian Forest Service (NRCAN)), forest tree
conservation (Morton Arboretum), and academia. The chromosome-scale reference was
annotated with existing transcriptomic (RNA-Seq) resources and examined in an evolutionary
context to recently sequenced members of the Juglandaceae from around the globe.

## Materials & Methods

*Plant Material, Extraction & Sequencing*

Leaves were collected from a single individual from a wild population of trees in
Sheffield, New Brunswick, Canada (45.88 N, -66.36 W) (Fig. 1A) that appeared healthy (Fig.
1B). A vouchered specimen was deposited at the Connell Memorial Herbarium, University of
New Brunswick (UNB #69382). Leaves were flash frozen in liquid nitrogen, and stored at -80°C
until extraction. Leaf tissue weighing 750mg was bulked from a single accession and ground
with a mortar and pestle in liquid nitrogen. High molecular weight (HMW) genomic DNA
extraction was then conducted following Vaillancourt and Buell CTAB-Genomic-tip protocol
(Vaillancourt et al. 2019). In brief, the aqueous layer containing the DNA was collected and
mixed with isopropanol to precipitate the DNA, which was then resuspended in Buffer G2 and
purified using a Qiagen Genomic-tip column. After being eluted in Buffer QF, the purified DNA
was further purified by precipitation with isopropanol, followed by a wash with ethanol and
elution in Buffer EB overnight at room temperature. DNA was stored at 4°C until shipped to
University of Connecticut for library preparation and sequencing. The DNA was sheared to
approximately 25 kb using a Covaris® G-tube (#520079, Covaris, LLC, Woburn, MA, USA). The
sequencing library was prepared using the ligation sequencing DNA v14 method (SQK-LSK114,
Oxford Nanopore Technologies, Oxford, UK). Three separate loads of the library were run on
one flow cell (FLO-PRO114M) over 90 hours, resulting in 5.2 million reads with MinKNOW® (v.
22.08.6).

*Assembly*

Kmer-freq and GCE were run to estimate genome size at k-mer size 17, taking the long
reads (FASTQ) as input (Wang et al. 2020; Liu et al. 2013). A haploid chromosome-scale
assembly was generated using Oxford Nanopore PromethION long reads at 100X coverage,
based on the estimated genome size of 570 Mb (Fig. 1C; Fig. S1). The raw reads were
basecalled by Guppy (v.6.3.8) in two modes: (1) high accuracy (Q>9) duplex (HAC) and (2)
super accuracy (Q>10) simplex (SUP). Both sets were independently assessed through the first
stages of QC and assembly. To remove potential contaminants from both read sets, including

bacteria and fungal associations, Centrifuge (v.1.0.4-beta) was used (Table S1) (Kim et al. 2016). Quality assessment was performed with NanoPlot before and after contaminant removal (De Coster et al. 2018). Two long-read assembly methods, Flye (v.2.9) and Canu (v.2.2), were evaluated (Kolmogorov et al. 2019; Koren et al. 2017). Following *de novo* assembly, Medaka (v.1.7.2) was used for error correcting (polishing) the Flye-generated assemblies (*medaka*). Polishing was not independently run on the Canu assembly outside of Canu's approach. Completeness and contiguity were assessed at each stage with BUSCO (v.5.0.0) against the Embryophyta single-copy ortholog database (OrthoDB v.10.0), QUAST (v.5.0.2), Minimap2 (v.2.15), and PycoQC (v.2.5.2) (Manni et al. 2021; Kriventseva et al. 2019; Gurevich et al. 2013; Li 2018; Leger and Leonard 2019). These metrics led to the selection of the SUP assemblies for further refinement. Following error-correction, Purge Haplotigs (v.1.1.2) was applied to the Canu and Flye assemblies to reduce allele-based duplications (Roach et al. 2018). Both assemblies were subsequently scaffolded with the chromosome-scale *Juglans mandshurica* reference (BioProject: PRJCA006358) using RagTag (v.2.1.0) in *scaffold mode* (Li et al. 2022b; Alonge and Leonard 2021). The assemblies were then filtered to include the 16 longest scaffolds that aligned to the 16 chromosomes. Using the same assessments listed above, the scaffolded Canu assembly was selected for downstream annotation and analysis.

*Annotation*

   Repeat identification on the selected Canu assembly was performed with RepeatModeler (v.2.0.4), which provided a set of consensus sequences that served as input to RepeatMasker (v.4.1.4) (Table S2) (Flynn et al. 2020). Regions identified as repeats were softmasked in preparation for genome annotation. Six public Illumina RNA-Seq libraries were accessed from NCBI (PRJNA413418) for *J. cinerea* leaf tissue (Table S3). Each library was adapter trimmed with FastP (v.0.23.2) (Chen et al. 2018). The adapter trimmed reads were aligned to the softmasked reference with HISAT2 (v.2.2.1) to provide evidence for gene prediction with an 85% mapping rate cutoff, leading us to exclude two libraries (Kim et al. 2019). The RNA alignments and the softmasked genome served as the input to both BRAKER (v2.1.6) and TSEBRA (v.1.0.1) to generate reference annotations (Bruna et al. 2021; Gabriel et al. 2021). BRAKER was run with transcriptomic alignments (RNA-only mode) and viridiplantae protein input from OrthoDB (v.10) (protein-only mode). Both sets of predictions were fed into TSEBRA. This resulted in one transcriptomic set of predictions from BRAKER and one combined run from TSEBRA. The resulting GFF files were summarized with AGAT (v.1.0.0) and functionally annotated with EnTAP (v.0.10.8) (Dainat 2022; Hart et al. 2020). This provided a reciprocal BLAST estimate against NCBI's RefSeq Plant Protein database (v208) as well as annotations derived from EggNOG and Gene Ontology (Ashburner et al. 2000; Huerta-Cepas et al. 2019). Both sets of predictions were filtered by the functional annotations; candidates without an EggNOG assignment were removed. The final annotation was selected from a combination of BUSCO, AGAT statistics, and the annotation rate reported by EnTAP.

*Methylation & Circos Visualization*

Because CpG and non-CpG methylation are central to plant gene regulation, 5mC
methylation was called in all contexts. Bonito-remora (v.0.6.2) with rerio modified base model
5mC_all_context_sup_r1041_e82 was used to estimate methylation (*bonito* n.d.; *rerio* n.d.). The
resulting calls were summarized with modbam2bed (v0.9.5) and filtered to >= 10X coverage per
site (*modbam2bed*). CpG content was quantified with fastaRegexFinder (v.0.2.0) (Beraldi n.d.).
For visualization with ShinyCircos, 5mC methylation was averaged across 100 kb windows, and
CpG, repeat, and gene coverage were summed across 100 kb, 10 kb, and 10 kb windows,
respectively (Yu et al. 2018).

*Synteny*

A macrosynteny analysis was conducted using MCscan (Python version) and jcvi
(v.1.3.3) with available *Juglans* species, including *J. cinerea* (PRJEB56451), *J. nigra* (Zhou et
al. 2023), *J. regia* (PRJNA291087), and *J. mandshurica* (PRJCA006358) (Tang et al. 2008)
(Table 1). If not pre-filtered, any unclassified scaffolds were eliminated from both the genome
and annotation files, resulting in a set of 16 chromosomes for each *Juglans* species. Protein-
coding sequences from chromosome-scale assemblies, along with their respective annotations
(GFF), were used as input for MCscan. The jcvi format module was used to first filter input
FASTA files by longest isoform and convert GFF files to BED format. Next, a pairwise synteny
search was conducted across *J. cinerea* and *J. mandshurica*, *J. mandshurica* and *J. regia,* and
*J. regia* and *J. nigra* respectively. The resulting anchors were then clustered into synteny blocks
with a minimum span of 30 and visualized by the jcvi karyotype graphical module. The final
chromosome order was determined by length decreasing from left to right. *J. regia* had eight
noticeable inversions when compared to *J. mandshurica*; however, upon further inspection
these chromosomes were found to be reverse complemented. To address this, the strand
orientation in the simple anchors file was manually adjusted.
26
*Gene Family Analysis*

Orthologous gene families were identified with OrthoFinder (v2.5.4) (Emms and Kelly
2019) (Table S4) using 10 species with reference genomes among *Juglans (J. cinerea, J. nigra,*
*J. regia, J. sigillata, J. microcarpa, J. hindsii, J. mandshurica, J. cathayensis)*, *Carya (Carya*
*illinoinensis)*, and *Pterocarya (Pterocarya stenoptera)*. The species tree, based on resulting
gene trees, was constructed in OrthoFinder and rooted using the species tree inference from all
genes (STAG) methodology (Emms and Kelly 2018). Support values for internal nodes reflect
the proportion of orthogroup trees where the bipartition is observed, and ranges from 0 to 1.
Orthogroups only representing one species and orthogroups with oversized representatives
from a single species (>100 genes) were removed. The remaining 17,471 orthogroups and
ultrametric tree were provided to CAFE v5 (Mendes et al. 2021) to identify gene families with
evidence of significant expansion or contraction at p-value < 0.01 using the base model. The
longest gene from each orthogroup was used to assign a putative function from EnTAP. Gene
families associated with retroelements were filtered from the final analysis.

**Results & Discussion**

*Assembly*

A total of 59.7 Gb of high accuracy-basecalled (HAC) duplex reads and 59.9 Gb of super-basecalled (SUP) simplex Oxford Nanopore reads were generated using a PromethION sequencer, which each provided 105X coverage at an estimated genome size of 570 Mb (Fig. 1C). Contamination filtering removed primarily bacteria, with the highest proportion of reads associated with *Acinetobacter baumannii* (Fig. S2). Filtering resulted in 58.0 Gb of HAC reads with an N50 length of 19.5 Kb, and 58.0 Gb of SUP reads with an N50 length of 22.4 Kb.

With the SUP reads, Flye initially produced a slightly less complete but more contiguous assembly than Canu; 98.9% vs. 99.2% BUSCO completeness and N50 of 2.55 Mb (1,320 scaffolds) vs. N50 of 2.39 Mb (2,286 contigs), respectively (Table S5; Fig. 1D). Polishing of Flye with Medaka yielded no increase in overall BUSCO completeness, but increased single-copy completeness by 0.1%. Purge Haplotigs increased the contiguity of both assemblies while largely maintaining completeness (N50 of 2.59 Mb in 667 scaffolds for Flye and N50 of 3.02 kb in 345 contigs for Canu). RagTag scaffolding with the *J. mandshurica* assembly drastically increased the contiguity of both assemblies, leaving Flye with 175 scaffolds and Canu with 58 scaffolds. The longest 16 scaffolds from each assembly were extracted and determined to be chromosome-level based on genome size, completeness, and read alignment. *Juglans mandshurica* was chosen for its close phylogenetic position, chromosome conservation, and ability to hybridize (Mu et al. 2017). The assemblies were fairly similar in terms of completeness, contiguity, and overall accuracy (Table S5). The scaffolded Canu assembly was selected for downstream analyses and distinguished only by fewer gaps when compared to the scaffolded Flye assembly (5.32 vs. 9.18 Ns per 100 kb). The final 16 pseudo-chromosome assembly reported is 539 Mb in length with an N50 of 34.86 Mb and an embryophyta BUSCO completeness of 99.0% (Table S5; Fig. 1D). To ensure the assembly was not negatively impacted by the scaffolding process, missassembly was evaluated by examining the percentage of unmapped and soft-clipped bases from read alignment with minimap2. Before scaffolding, 12% of bases were unmapped and 3.58% were soft-clipped. After scaffolding, the percentage of unmapped bases remained the same, and 5.77% of bases were soft-clipped (Fig. S3). This supports minimal impact on overall quality through the scaffolding process.

This exclusively nanopore-assembled genome benefitted from high quality base calls (SUP reads) and >100X coverage to generate a genome of 539 Mb (slightly less than the estimated 570 Mb) (Table S5). The initial genome size estimates were produced from uncorrected long-reads and may not be as reliable of an estimate but were consistent with C-value estimates from Genomes on a Tree (GoaT) (Challis et al. 2023). *Juglans cinerea* is the 10th member of the Juglandaceae to have its whole genome sequenced. Comparing our final *J. cinerea* assembly with that of *J. mandshurica* to which it was scaffolded (Li et al. 2022b), *J. cinerea* exhibits a smaller scaffold N50 size of 34.86 Mb, higher BUSCO completeness score of 99.0%, larger genome size of 539 Mb, and higher GC content of 36.64% (Table 1).

*Annotation*

Prior to structural annotation, repeat sequences in the assembly were identified, resulting in 2,353 RepeatScout/RECON families and 1,117 LTRPipeline families. A total of 298 of the LTR families were determined to be redundant and removed, resulting in 3,172 unique repeats. Repeatmasker used this library to softmask 302 Mb (56.02%) of the butternut genome (Fig. 1E). Among the softmasked regions, 28.72% represented retroelements, most of which (22.46%) were predicted to be long terminal repeats (LTRs). Gypsy elements accounted for 8.13% of the genome, whereas Copia elements made up 11.12%. As expected, a large proportion of the repeats (22.42%) were unclassified (Table S2). *J. cinerea* exhibited repeat content that is consistent with other sequenced *Juglans*, with a value that fell between *J. sigillata* (50.06%) and *J. mandshurica* (60.99%) (Ning et al. 2020; Li et al., 2022). Retroelements–mostly LTRs–compose the highest classified repeat content in all three genomes. As reported in the *J. regia* x *Juglans microcarpa* hybrid and *J. regia* genomes, the telomeric repeat CCCTAAA was identified in *J. cinerea* at the ends of each chromosome (Zhu et al. 2019; Marrano et al. 2020).

After trimming, the six RNA-Seq libraries had an average mapping rate of 91.2% against the assembled *J.cinerea* genome (Table S4). Gene prediction was attempted with both BRAKER and TSEBRA on the softmasked assembly (Table S6). BRAKER provided a high quality annotation with only transcriptomic data, resulting in 38,377 predicted genes, 41,357 transcripts, a mono:multi ratio of 0.286, a BUSCO completeness score of 97.2% and an annotation rate of 77.1%. TSEBRA, which combined transcriptomic and protein BRAKER runs, resulted in a slightly lower quality annotation with 35,437 genes, 38,928 transcripts, 0.54 mono:mult ratio, 96.8% BUSCO completeness score, and 86.2% annotation rate. After filtering with EggNOG, the quality of both runs increased, and the number of predicted genes and transcripts was reduced to 30,526 and 33,275 respectively in BRAKER, and 30,328 and 33,694 in TSEBRA. Both programs resulted in a 98.1% annotation rate after filtering, although the BRAKER annotation was closer to the expected mono:multi ratio of plants at 0.21, outperforming that of TSEBRA, 0.36 (Vuruputoor et al. 2022). As a result, BRAKER was chosen to represent the final structural annotation for *J. cinerea*. These annotation results are similar to other *Juglans* species, including the *J. mandshurica* genome released in 2022, which has 39,466 predicted genes, 47,355 transcripts, a mono:multi ratio of 0.347, 95.2% BUSCO completeness, and an annotation rate of 84.9% in chromosome-level scaffolds. Likewise, *J. nigra* and *J. regia*, which have 29,949 and 30,709 genes respectively, 29,949 and 45,959 transcripts, BUSCO completeness of 96.4% and 99.6%, mono:multi of 0.210 and 0.206, and an annotation rate of 96.3% and 99.9% (Table 1). The impressive BUSCO completeness score of the first version of *J. cinerea*, ranking second to version three of *J. regia*, is a testament to the effectiveness of our methodology.

All context methylation (based on 5mC) revealed a central peak pattern to each chromosome that did not correlate with CpG content (Fig. 1E). However, it does appear to correspond with small peaks of repeat content, possibly representing centromeres. Future analyses should analyze methylation by type (CpG, CHG, CHH) in association with repeats and across gene lengths.

*Synteny*

The success of our genome assembly is in part due to the quality of the *J. mandshurica* genome, which was used to scaffold our *de novo* assembly. However, utilizing the *scaffold mode* of RagTag and not the *correction mode* allowed some rearrangements to be retained in the *J. cinerea* assembly, as highlighted by the pairwise macrosynteny analysis (Fig. 2). Conserved synteny at chromosome-level was observed in a phylogenetic context, comparing *J. cinerea* to *J. mandshurica*, *J. mandshurica* to *J. nigra,* and *J. nigra* to *J. regia*. A total of 52,993 (NR:38,556) seed synteny blocks (anchors) were found in 294 clusters between *J. cinerea* and *J. mandshurica*, 46,251 (NR:37,316) anchors were identified in 297 clusters between *J.mandshurica* and *J. nigra*, and 47,683 (NR:37,354) anchors were found in 247 cluster across *J.nigra* and *J. regia*. Considering the total number of protein-coding genes in each pairwise comparison, respectively, 68.1%, 66.6%, and 78.6% of compared genomes were collinear, indicating strong support for ancestral collinearity (Li et al. 2022b).

*Gene Family Analysis*

The OrthoFinder analysis conducted with ten Juglandaceae assigned 384,404 genes out of 401,209 (95.8%) into 33,922 orthogroups (Fig. 3A; Table S4). A total of 13,173 orthogroups (38.8%) were observed in all species and 4,076 orthogroups (12%) were specific to a single species. There were 807 single-copy orthogroups and 16,805 unassigned genes. In *J. cinerea*, 32,743 genes (98.4%) were assigned to an orthogroup. Of the total number of orthgroups found across the 10 species, 22,863 (67.4%) included *J. cinerea*. A total of 132 orthogroups (0.39%) containing 328 genes were unique to *J. cinerea*, and only 532 genes (1.6%) could not be assigned to an orthogroup. *J. cinerea* shares more one-to-one orthologs outside of the *Trachycaryon-Cardiocaryon* clade than within the clade, possibly reflecting isolation and divergence of the ancestral *Cardiocaryon* population to the ancestral *J. cinerea* population (Fig.3A; Table S4).

Members of the Juglandaceae assessed here belong to one of four sections. The North American species *J. nigra*, *J. hindsii* and *J. microcarpa* are members of sect. *Rhysocaryon*. The sampled Eurasian species (sect. *Juglans)* include *J. sigillata and J. regia*, the latter being the predominant cultivated species for walnut production. *Juglans cathayensis and J. mandshurica* are members of sect. *Cardiocaryon*. *J. ailantifolia*, also in section *Cardiocaryon,* but not included here, is sometimes described as a sub-species of *J. mandshurica* (Mu et al. 2017). This is of note given the ability of *J. cinerea* to hybridize with *J. ailantifolia* but not with the sympatric *J. nigra* (Brennan et al. 2020).

Phylogenetic relationships among Juglandaceae remain enigmatic despite multiple sequence and morphology-based analyses. Using protein sequences sourced from whole genome assemblies, *Juglans* was found to be sister to *Carya* with *Pterocarya* as the outgroup (Fig. 3B). Recent phylogenomic analyses relying on sequence data from *de novo* aligned SNPs (Mu et al. 2020), RADseq SNPs mapped to the *J. regia* genome (Zhou et al. 2021), and a deep analysis of subgenomes (Ding et al. 2023) support *Juglans* as sister to *Pterocarya, with Carya* as the outgroup (Fig. 3C). The incongruent results may be an artifact of reduced sampling in this study, however further research is warranted.

Within *Juglans,* phylogenetic discordance is frequently observed between trees built from different datasets, including whole genome sequencing, and the above mentioned SNP sets, subgenome analyses, and plastid versus nuclear markers. The tree generated from this study places *Juglans* section *Rhysocaryon* sister to sect. *Juglans*, in turn, are sister to a clade containing sections *Cadiocaryon* and *Trachycaryon* (Fig. 3C). However, RADseq SNPs dervied from *de novo* alignments generate a tree with section *Juglans* sister to *Cardiocaryon* and section *Rhysocaryon* sister to *Trachycaryon* (Mu et al. 2020), a topology in complete discordance with ours. Topologies made from nuclear SNPs mapped to the *J. regia* genome place *Cardiocaryon* sister to *Trachycaryon*, as observed here; however, section *Juglans* subtends the *Cardiocaryon-Trachycarhon* clade with *Rhysocaryon* as the outgroup in the nuclear SNP-based topology (Zhou et al. 2021).

The placement of *J. cinerea* has varied from the earliest plastid and ITS trees (Stanford et al. 2000), to more recent trees built from whole chloroplast genomes and low-copy nuclear markers (Dong et al. 2017). Characteristic of these trees is the uncertainty of the placement of *J. cinerea* with either Asian or North American *Juglans*. Phylogenetic evidence from whole plastid genomes of *Juglans* indicates *J. cinerea* captured a North American chloroplast genome, while the analysis conducted here places *J. cinerea* sister to the Asian Butternut clade (Fig. 3B). Clearly given the discordance within *Juglans* and *Juglandaceae,* there is a pressing need for additional research using multiple phylogenetic approaches and simulation models.

The overall gene family analysis with CAFE identified a mean gene family birth-death rate ($\lambda$) for the whole tree of 0.88, with a tri-modal distribution of gene family expansion/contraction probabilities (Fig. S4). The species with the most rapidly evolving gene families (significant at p-value < 0.01) was *J. regia* with 523 gene families. *J. cinerea* was the species within *Juglans* with the lowest number of significantly evolving gene families, with a total of 71, and the majority of these families were contracting (N = 52, Fig. 3B). After removing orthogroups consisting of retroelements, a total of 54 orthogroups were identified from CAFE as rapidly expanding (9) and rapidly contracting (45) in *J. cinerea* (Fig. 4; Table S7). Among the small set expanding, COG categories broadly described them as cytoskeleton-related, transcription regulation, post-translational modification, cell cycle control, and ion transport. Among those contracting, functions related to transcription regulation, signal transduction, secondary metabolite biosynthesis, carbohydrate transport, disease resistance, and cytoskeleton-related were represented. Examining expanded gene families related to biotic-stress (pathogen/pest resistance), revealed two gene families, one is associated with Auxin response factors (ARFs) and the other with Ripening-related proteins. The pathogenesis-related contracting families were far more numerous and included: NB-LRRs, Wall-associated receptor kinases (WAK), Germin-like proteins, Cytochrome P450s, glycosyl hydrolases, lipolytic acyl hydrolase (LAH), mTERFs, Class III peroxidases (PRX3), UPF0481 proteins, and PAN-like domain proteins. Of these, the following are actually absent in *J. cinerea*: one NB-LRR gene family, one of the three contracting WAK gene families, one cytochrome P450, one of three contracting glycosyl hydrolases gene families, lipolytic acyl hydrolase (LAH), one of two PRX3s, and both families associated with UPF0481 proteins.

Wall-associated receptor kinases (WAKs) are increasingly associated with plant immunity, including neo- and sub-functionalization in response to changing selection pressure

(Stephens et al. 2022; Zhou et al. 2023). Among *Juglans*, they are believed to be absent in *J.*
*hindsii* and *J. sigillata* (Trouern-Trend et al. 2020), rapidly contracting in *J. mandshurica*, and
expanding in *J. regia* (Yan et al. 2021). Evaluation of WAK gene families in *J. mandshurica* and
*J. regia* found they are under purifying selection (Yan et al. 2021). Expression levels of WAK
gene family members were observed to be higher in anthracnose-resistant varieties of *J. regia*
(Li et al. 2022a). In *J. cinerea*, a strong signature of contraction across two WAK gene families
and absence in the third, was observed. Further investigation of WAKs and other pathogenesis-
related families in a population context may reveal more about their role in resistance to specific
fungal pathogens.

*Conclusions*
The present study reports *Juglans cinerea*'s first high-quality reference genome,
generated using Oxford Nanopore technology and a combination of assembly tools. The *J.*
*mandshurica* genome enabled pseudo-chromosome generation and chromosome-level
analyses. A preliminary comparative view of *J. cinerea* was presented in context to other
species in the Juglandaceae family to assess both synteny and gene family evolution. The
construction of the genome was connected to a year-long training program initiated with high
molecular weight extraction of the target tree. Students worked closely with graduate students
and faculty members to understand the challenges associated with genome assembly and
annotation. In doing so, the genome benefitted from robust methods testing. Finally, the
program intersected with practitioners who provided input on the reference accession and a
framework that will guide the application of this high quality reference in the future. Future
studies expanded to population-level analyses of *J. cinerea*, and *J. ailantifolia* hybrid resistance
to Oc-j, will be needed to develop conservation strategies informed by genetics.

**Table 1. Comparison of Juglans chromosome-level genome assemblies and annotations.**
*Assembly and annotations not yet publicly available.

| | *J. cinerea* | *J. mandshurica* | *J. nigra* | *J. regia* |
|---|---|---|---|---|
| **Bioproject** | PRJEB56451 | PRJNA730664 | * | PRJNA291087 |
| **Biosample** | SAMEA111438462 | SAMN19238346 | * | SAMN03704235 |
| **GenBank** | * | GCA_022457165.1 | * | GCA_001411555.2 |
| **Sequence method** | Oxford Nanopore PromethION | PacBio HiFi, Hi-C, Illumina HiSeq | Illumina HiSeq; Pacbio SMRTbell; Hi-C, Illumina NovaSeq6000 | Illumina HiSeq2500; Oxford Nanopore MinION; Hi-C, Illumina HiSeq4000 |
| ***Assembly*** | | | | |
| **Genome size (Mb)** | 539.43 | 528.15 | 530.24 | 545.78 |
| **Number of chromosomes** | 16 | 16 | 16 | 16 |
| **N50 (Mb)** | 34.86 | 35.38 | 35.14 | 37.11 |
| **GC content** | 36.64% | 36.49% | 36.36% | 36.08% |
| **Complete BUSCO** | 99.0% | 98.8% | 99.0% | 99.0% |
| ***Annotation*** | | | | |
| **Number of genes** | 30,526 | 39,466 | 29,949 | 30,709 |
| **Number of transcripts** | 33,275 | 47,355 | 29,949 | 45,959 |
| **Average gene length (bp)** | 4,187 | 3,748 | 5,175 | 5,309 |
| **Average exon length (bp)** | 209 | 287 | 297 | 291 |
| **Complete BUSCO** | 97.2% | 95.2% | 96.4% | 99.6% |
| **Mono:multi ratio** | 0.207 | 0.347 | 0.210 | 0.206 |
| **Annotation rate** | 98.1% | 84.9% | 96.3% | 99.9% |

**Figure 1.** *Range map and sample location of the Juglans cinerea reference tree (UNB #69382).*
*Range information from the BIEN database **(A)**. Photographs of the sampled reference tree*
*from the Sheffield, NB population exhibiting a healthy crown and trunk absent of cankers **(B)**. k-*

*mer plot (length=17bp) of k-mer count (x-axis) versus frequency (y-axis). Genome size from*
*long-read data initially estimated at 570 Mb **(C)**. Scatter plot comparing assembly length (Mb)*
*and N50 (kb) obtained from QUAST for Flye and Canu throughout the assembly process.*
*Assemblies represented by color (Canu=yellow; Flye=blue) and shape (drafts=circles; selected*
*final assembly pre-chromosome filtering=star) with abbreviated assembly software*
*(MD=Medaka; PH=Purge Haplotigs). BUSCO single (S) and duplicated (D) completeness*
*scores in parentheses **(D)**. Circos plot representing 0) assembly chromosomes 1) average 5mC*
*methylation in all sequence contexts (CG, CHG, and CHH) over 100 kb windows 2) CpG site*
*count over 100 kb windows 3) gene coverage over 10 kb windows 4) repeat coverage over 10*
*kb windows **(E)**.*

***Figure 2.** Pairwise macro-synteny analysis in phylogenetic order comparing chromosomes*
*(boxes) and syntenic clusters (grey lines) in J. cinerea (purple) and J. mandshurica (green), J.*
*mandshurica and J. nigra (blue) and J. nigra and J. regia (orange). Published chromosome*
*identifiers and orientation from annotation are maintained, but chromosomes are re-ordered by*
*size (largest to smallest).*

***Figure 3.** Results of OrthoFinder and the phylogenomic analysis of 33,922 orthrogroup gene*
*trees. Upset plot of the numbers of genes shared by different intersections of 10 species **(A)**.*
*Consensus phylogeny from 33,922 orthogroup gene trees. STAG support values at internal*
*nodes and the number significantly expanding or contracting gene families (p-value < 0.01) on*
*branches **(B)**. Alternate hypotheses of relationships between Carya, Juglans, and Pterocarya*
*using data from RADseq SNPs with de novo alignment (Mu et al. 2020), RADseq SNPs*
*mapped to the J. regia genome (Zhou et al. 2021), subgenomes (Ding et al. 2023) versus whole*
*genome sequencing and orthogroup gene tree analysis (this study) **(C)**.*

***Figure 4.** Annotations of J. cinerea significantly contracting (pink) and expanding (green) gene*
*families, excluding retroelement families.*

**Data Availability Statement**
The ONT raw reads are available on the European Nucleotide Archive (PRJEB56451). Code,
genome assembly, and genome annotation files are available on GitLab:
https://doi.org/10.5281/zenodo.5634263.

**Funder Information**
Funding was provided by the University of Connecticut, College of Liberal Arts and Sciences,
through the Earth and its Future initiative that enabled the initiation of the Biodiversity and
Conservation Genomics Training Program. Funding was also provided through National
Science Foundation Awards DBI-1943371. Support for nanopore sequencing of the IUCN-listed
*Juglans cinerea* was provided by the OrgOne project by Oxford Nanopore Technologies. This
research was also funded in part by the USDA Forest Service.The use of trade names is for the
information and convenience of the reader and does not imply official endorsement or approval

by the U. S. Department of Agriculture or the Forest Service of any product to the exclusion of
others that may be suitable.

**Acknowledgements**
We thank the Institute for Systems Genomics Center for Genome Innovation at the University of
Connecticut for molecular resources and support. We would also like to thank the
Computational Biology Core for access to HPC resources. Additional thanks to Patrick G.S.
Grady for genome assembly consultation.

**REFERENCES:**

Alonge, M., & Leonard, A. (2021). *malonge/RagTag: RagTag v2.1.0*.
https://doi.org/10.5281/zenodo.5634263

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P.,
Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L.,
Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M., &
Sherlock, G. (2000). Gene ontology: tool for the unification of biology. The Gene
Ontology Consortium. *Nature genetics*, *25*(1), 25–29. https://doi.org/10.1038/75556

Beraldi, D. (n.d.). *fastaRegexFinder at master · dariober/bioinformatics-cafe*. Github. Retrieved
May 4, 2023, from https://github.com/dariober/bioinformatics-cafe

*bonito: A PyTorch Basecaller for Oxford Nanopore Reads*. (n.d.). Github. Retrieved May 4,
2023, from https://github.com/nanoporetech/bonito

Boraks, A., & Broders, K. D. (2014). Butternut (Juglans cinerea) health, hybridization, and
recruitment in the northeastern United States. *Canadian Journal of Forest Research.*
*Journal Canadien de La Recherche Forestiere*, *44*(10), 1244–1252.

Brennan, A. N., McKenna, J. R., Hoban, S. M., & Jacobs, D. F. (2020). Hybrid Breeding for
Restoration of Threatened Forest Trees: Evidence for Incorporating Disease Tolerance
in Juglans cinerea. *Frontiers in Plant Science*, *11*, 580693.

Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M., & Borodovsky, M. (2021). BRAKER2:
automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS
supported by a protein database. *NAR Genomics and Bioinformatics*, *3*(1), lqaa108.

*Butternut (Juglans cinerea)*. (2021). Wisconsin Department of Natural Resources (WDNR).
Retrieved May 19, 2023, from

https://dnr.wi.gov/topic/EndangeredResources/Plants.asp?mode=detail&SpecCode=PD
J UG02030

Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing sixth
mass extinction signaled by vertebrate population losses and declines. *Proceedings of*
*the National Academy of Sciences of the United States of America*, *114*(30), E6089–
E6096.

Challis, R., Kumar, S., Sotero-Caio, C., Brown, M., & Blaxter, M. (2023). Genomes on a Tree
(GoaT): A versatile, scalable search engine for genomic and sequencing project
metadata across the eukaryotic tree of life. *Wellcome Open Research*, *8*, 24.

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* , *34*(17), i884–i890.

Dainat, J., Hereñú, D., Davis, E., Crouch, K., LucileSol, Agostinho, N., pascal-git, & tayyrov.
(2022). *NBISweden/AGAT: AGAT-v1.0.0*. https://doi.org/10.5281/zenodo.7255559
De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., & Van Broeckhoven, C. (2018). NanoPack:
visualizing and processing long-read sequencing data. *Bioinformatics* , *34*(15), 2666–
2669.

DeWoody, J. A., Jeon, J. Y., Bickham, J. W., Heenkenda, E. J., Janjua, S., Lamka, G. F.,
Mularo,
A. J., Black, A., Brüniche-Olsen, A., & Willoughby, J. R. (2022). The Threatened Species
Imperative: Conservation assessments would benefit from population genomic insights
[Review of *The Threatened Species Imperative: Conservation assessments would*
*benefit from population genomic insights*]. *Proceedings of the National Academy of*
*Sciences of the United States of America*, *119*(35), e2210685119.

Ding, Y.-M., Pang, X.-X., Cao, Y., Zhang, W.-P., Renner, S. S., Zhang, D.-Y., & Bai, W.-N.
(2023). Genome structure-based Juglandaceae phylogenies contradict alignment-based
phylogenies and substitution rates vary with DNA repair genes. *Nature Communications*,
*14*(1), 617.

Dong, W., Xu, C., Li, W., Xie, X., Lu, Y., Liu, Y., Jin, X., & Suo, Z. (2017). Phylogenetic
Resolution in Juglans Based on Complete Chloroplast Genomes and Nuclear DNA
Sequences. *Frontiers in Plant Science*, *8*, 1148.

Emms, D. M., & Kelly, S. (2018). STAG: Species Tree Inference from All Genes. In *bioRxiv* (p.
267914). https://doi.org/10.1101/267914

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology*, *20*(1), 238.

Fernandes, P., Colavolpe, M. B., Serrazina, S., & Costa, R. L. (2022). European and American chestnuts: An overview of the main threats and control efforts. *Frontiers in Plant Science*, *13*, 951844.

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F. (2020). RepeatModeler2 for automated genomic discovery of transposable element families. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(17), 9451–9457.

Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M., & Stanke, M. (2021). TSEBRA: transcript selector for BRAKER. *BMC Bioinformatics*, *22*(1), 1–12.

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment tool for genome assemblies. *Bioinformatics* , *29*(8), 1072–1075.

Hart, A. J., Ginzburg, S., Xu, M. S., Fisher, C. R., Rahmatpour, N., Mitton, J. B., Paul, R., & Wegrzyn, J. L. (2020). EnTAP: Bringing faster and smarter functional annotation to non-model eukaryotic transcriptomes. *Molecular Ecology Resources*, *20*(2), 591–604.

Hoban, S., Anderson, R., McCleary, T., Schlarbaum, S., & Romero-Severson, J. (2008). Thirteen nuclear microsatellite loci for butternut (Juglans cinerea L.). *Molecular Ecology Resources*, *8*(3), 643–646.

Hoban, S. M., Borkowski, D. S., Brosi, S. L., McCleary, T. S., Thompson, L. M., McLachlan, J. S., Pereira, M. A., Schlarbaum, S. E., & Romero-Severson, J. (2010). Range-wide distribution of genetic diversity in the North American tree Juglans cinerea: a product of range shifts, not ecological marginality or recent population decline. *Molecular Ecology*, *19*(22), 4876–4891.

Hoban, S. M., McCleary, T. S., Schlarbaum, S. E., Anagnostakis, S. L., & Romero-Severson, J. (2012). Human-impacted landscapes facilitate hybridization between a native and an introduced tree. *Evolutionary Applications*, *5*(7), 720–731.

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., & Bork, P. (2019). eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic acids research*, *47*(D1), D309–D314. https://doi.org/10.1093/nar/gky1085

*IUCN 2021 : International Union for Conservation of Nature annual report.* (2022).
https://portals.iucn.org/library/node/49944

Kardos, M., Armstrong, E. E., Fitzpatrick, S. W., Hauser, S., Hedrick, P. W., Miller, J. M.,
Tallmon, D. A., & Funk, W. C. (2021). The crucial role of genome-wide genetic variation
in conservation. *Proceedings of the National Academy of Sciences of the United States*
*of America*, *118*(48). https://doi.org/10.1073/pnas.2104642118

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*,
*37*(8), 907–915.

Kim, D., Song, L., Breitwieser, F. P., & Salzberg, S. L. (2016). Centrifuge: rapid and sensitive
classification of metagenomic sequences. *Genome Research*, *26*(12), 1721–1729.

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads
using repeat graphs. *Nature Biotechnology*, *37*(5), 540–546.

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017).
Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and
repeat separation. *Genome Research*, *27*(5), 722–736.

Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., & Zdobnov,
E. M. (2019). OrthoDB v10: sampling the diversity of animal, plant, fungal, protist,
bacterial and viral genomes for evolutionary and functional annotations of orthologs.
*Nucleic Acids Research*, *47*(D1), D807–D811.

LaBonte, N. R., Ostry, M. E., Ross-Davis, A., & Woeste, K. E. (2015). Estimating heritability of
disease resistance and factors that contribute to long-term survival in butternut (Juglans
cinerea L.). *Tree Genetics & Genomes*, *11*(3), 63.

Laricchia, K. M., McCleary, T. S., Hoban, S. M., Borkowski, D., & Romero-Severson, J. (2015).
Chloroplast haplotypes suggest preglacial differentiation and separate postglacial
migration paths for the threatened North American forest tree Juglans cinerea L. *Tree*
*Genetics & Genomes*, *11*(2). https://doi.org/10.1007/s11295-015-0852-3

Leger, A., & Leonardi, T. (2019). pycoQC, interactive quality control for Oxford Nanopore
Sequencing. *Journal of Open Source Software*, *4*(34), 1236.

Lewin, H. A., Richards, S., Lieberman Aiden, E., Allende, M. L., Archibald, J. M., Bálint, M.,
Barker, K. B., Baumgartner, B., Belov, K., Bertorelle, G., Blaxter, M. L., Cai, J.,

Caperello, N. D., Carlson, K., Castilla-Rubio, J. C., Chaw, S. M., Chen, L., Childers, A. K., Coddington, J. A., Conde, D. A., … Zhang, G. (2022). The Earth BioGenome Project 2020: Starting the clock. *Proceedings of the National Academy of Sciences of the United States of America*, *119*(4), e2115635118. https://doi.org/10.1073/pnas.2115635118

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* , *34*(18), 3094–3100.

Li, M., Ma, J., Liu, H., Ou, M., Ye, H., & Zhao, P. (2022). Identification and Characterization of Wall-Associated Kinase (WAK) and WAK-like (WAKL) Gene Family in Juglans regia and Its Wild Related Species Juglans mandshurica. *Genes*, *13*(1). https://doi.org/10.3390/genes13010134

Li, X., Cai, K., Zhang, Q., Pei, X., Chen, S., Jiang, L., Han, Z., Zhao, M., Li, Y., Zhang, X., Li, Y., Zhang, S., Chen, S., Qu, G., Tigabu, M., Chiang, V. L., Sederoff, R., & Zhao, X. (2022). The Manchurian Walnut Genome: Insights into Juglone and Lipid Biosynthesis. *GigaScience*, *11*. https://doi.org/10.1093/gigascience/giac057

Liu, B., Shi, Y., Yuan, J., Hu, X., Zhang, H., Li, N., Li, Z., Chen, Y., Mu, D., & Fan, W. (2013). Estimation of genomic characteristics by analyzing k-mer frequency in de novo genome projects. In *arXiv [q-bio.GN]*. arXiv. http://arxiv.org/abs/1308.2012

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*, *38*(10), 4647–4654.

Marrano, A., Britton, M., Zaini, P. A., Zimin, A. V., Workman, R. E., Puiu, D., Bianco, L., Pierro, E. A. D., Allen, B. J., Chakraborty, S., Troggio, M., Leslie, C. A., Timp, W., Dandekar, A., Salzberg, S. L., & Neale, D. B. (2020). High-quality chromosome-scale assembly of the walnut (Juglans regia L.) reference genome. *GigaScience*, *9*(5). https://doi.org/10.1093/gigascience/giaa050

*medaka: Sequence correction provided by ONT Research*. (n.d.). Github. Retrieved May 4, 2023, from https://github.com/nanoporetech/medaka

Mendes, F. K., Vanderpool, D., Fulton, B., & Hahn, M. W. (2021). CAFE 5 models variation in evolutionary rates among gene families. *Bioinformatics* , *36*(22-23), 5516–5518.

*modbam2bed*. (n.d.). Github. Retrieved May 4, 2023, from https://github.com/epi2me-labs/modbam2bed

Mu, X.-Y., Sun, M., Yang, P.-F., & Lin, Q.-W. (2017). Unveiling the Identity of Wenwan Walnuts
and Phylogenetic Relationships of Asian Juglans Species Using Restriction Site-
Associated DNA-Sequencing. *Frontiers in Plant Science*, *8*, 1708.

Mu, X.-Y., Tong, L., Sun, M., Zhu, Y.-X., Wen, J., Lin, Q.-W., & Liu, B. (2020). Phylogeny and
divergence time estimation of the walnut family (Juglandaceae) based on nuclear RAD-
Seq and chloroplast genome data. *Molecular Phylogenetics and Evolution*, *147*, 106802.

Nair, V. M. G., Kostichka, C. J., & Kuntz, J. E. (1979). Sirococcus
Clavigignenti-Juglandacearum: An Undescribed Species Causing Canker on Butternut.
*Mycologia*, *71*(3), 641–646.

Nielsen, C., M. Cherry, B. Boysen, A. Hopkin, J. McLaughlin, T. Beardmore. (2003). *COSEWIC*
*status report on the butternut Juglans cinerea in Canada in COSEWIC assessment and*
*status report on the butternut Juglans cinerea in Canada* (pp. 1–32). Committee on the
Status of Endangered Wildlife in Canada.

Ning, D.-L., Wu, T., Xiao, L.-J., Ma, T., Fang, W.-L., Dong, R.-Q., & Cao, F.-L. (2020).
Chromosomal-level assembly of Juglans sigillata genome using Nanopore, BioNano,
and Hi-C analysis. *GigaScience*, *9*(2). https://doi.org/10.1093/gigascience/giaa006

Pike, C. C., Williams, M., Brennan, A., Woeste, K., Jacobs, J., Hoban, S., Moore, M., &
Romero-Severson, J. (2020). Save Our Species: A Blueprint for Restoring Butternut
(Juglans cinerea) across Eastern North America. *Journal of Forestry*, *119*(2), 196–206.

R. Taylor, H., Dussex, N., & van Heezik, Y. (2017). Bridging the conservation genetics gap by
identifying barriers to implementation for conservation practitioners. *Global Ecology and*
*Conservation*, *10*, 231–242.

*rerio: Research release basecalling models and configurations*. (n.d.). Github. Retrieved May 4,
2023, from https://github.com/nanoporetech/rerio

Roach, M. J., Schmidt, S. A., & Borneman, A. R. (2018). Purge Haplotigs: allelic contig
reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics*, *19*(1), 460.

Sandström, A., Lundmark, C., Andersson, K., Johannesson, K., & Laikre, L. (2019).
Understanding and bridging the conservation-genetics gap in marine conservation.
*Conservation Biology: The Journal of the Society for Conservation Biology*, *33*(3), 725–
728.

Smith, W. (2018). *Juglans cinerea : Butternut | Rare Species Guide*. Minnesota Department of
   Natural Resources.
   https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=PD
   JUG02030

Stanford, A. M., Harden, R., & Parks, C. R. (2000). Phylogeny and biogeography of Juglans
   (Juglandaceae) based on matK and ITS sequence data. *American Journal of Botany*,
   *87*(6), 872–882.

Stephens, C., Hammond-Kosack, K. E., & Kanyuka, K. (2022). WAKsing plant immunity, waning
   diseases. *Journal of Experimental Botany*, *73*(1), 22–37.

Tang, H., Bowers, J. E., Wang, X., Ming, R., Alam, M., & Paterson, A. H. (2008). Synteny and
   collinearity in plant genomes. *Science*, *320*(5875), 486–488.

Trouern-Trend, A. J., Falk, T., Zaman, S., Caballero, M., Neale, D. B., Langley, C. H.,
   Dandekar, A. M., Stevens, K. A., & Wegrzyn, J. L. (2020). Comparative genomics of six
   Juglans species reveals disease-associated gene family contractions. *The Plant Journal:
   For Cell and Molecular Biology*, *102*(2), 410–423.

Vaillancourt, B., & Robin Buell, C. (2019). High molecular weight DNA isolation method from
   diverse plant species for use with Oxford Nanopore sequencing. In *bioRxiv* (p. 783159).
   https://doi.org/10.1101/783159

Vuruputoor, V. S., Monyak, D., Fetter, K. C., Webster, C., Bhattarai, A., Shrestha, B., Zaman,
   S., Bennett, J., McEvoy, S. L., Caballero, M., & Wegrzyn, J. L. (2022). Welcome to the
   big leaves: best practices for improving genome annotation in non-model plant
   genomes. In *bioRxiv* (p. 2022.10.03.510643). https://doi.org/10.1101/2022.10.03.510643

Wang, H., Liu, B., Zhang, Y., Jiang, F., Ren, Y., Yin, L., Liu, H., Wang, S., & Fan, W. (2020).
   Estimation of genome size using k-mer frequencies from corrected long reads. In *arXiv
   [q-bio.GN]*. arXiv. http://arxiv.org/abs/2003.11817

Yan, F., Xi, R.-M., She, R.-X., Chen, P.-P., Yan, Y.-J., Yang, G., Dang, M., Yue, M., Pei, D.,
   Woeste, K., & Zhao, P. (2021). Improved de novo chromosome-level genome assembly
   of the vulnerable walnut tree Juglans mandshurica reveals gene family evolution and
   possible genome basis of resistance to lesion nematode. *Molecular Ecology Resources*,
   *21*(6), 2063–2076.

Yu, Y., Ouyang, Y., & Yao, W. (2018). shinyCircos: an R/Shiny application for interactive
   creation of Circos plot. *Bioinformatics* , *34*(7), 1229–1231.

Zhou, H., Hu, Y., Ebrahimi, A., Liu, P., Woeste, K., Zhao, P., & Zhang, S. (2021). Whole
genome based insights into the phylogeny and evolution of the Juglandaceae. *BMC*
*Ecology and Evolution*, *21*(1), 191.

Zhou, H., Yan, F., Hao, F., Ye, H., Yue, M., Woeste, K., Zhao, P., & Zhang, S. (2023). Pan-
genome and transcriptome analyses provide insights into genomic variation and
differential gene expression profiles related to disease resistance and fatty acid
biosynthesis in eastern black walnut (Juglans nigra). *Horticulture Research*, *10*(3),
uhad015.

Zhu, T., Wang, L., You, F. M., Rodriguez, J. C., Deal, K. R., Chen, L., Li, J., Chakraborty, S.,
Balan, B., Jiang, C.-Z., Brown, P. J., Leslie, C. A., Aradhya, M. K., Dandekar, A. M.,
McGuire, P. E., Kluepfel, D., Dvorak, J., & Luo, M.-C. (2019). Sequencing a Juglans
regia × J. microcarpa hybrid yields high-quality genome assemblies of parental species.
*Horticulture Research*, *6*, 55.

| COG Description | EggNOG Description | Contracting | Expanding |
|---|---|---|---|
| Amino acid transport and metabolism | Lipoxygenase | 1 | |
| Carbohydrate transport and metabolism | glycosyl hydrolase | 3 | |
| | UDP-glycosyltransferase family | 1 | |
| Cell cycle control and mitosis | Meprin and TRAF domain | | 1 |
| Cytoskeleton | Tubulin | 1 | |
| | ADF (actin binding) | | 1 |
| Energy production and conversion | (S)-2-hydroxy-acid oxidase | 1 | |
| Function unknown | Cysteine-rich receptor-like protein kinase | 5 | |
| | RBD (RNA-binding domain) | 4 | |
| | Plant protein of unknown function | 3 | |
| | UPF0481 protein | 2 | |
| | FAR1-RELATED SEQUENCE | 2 | |
| | zinc-binding in reverse transcriptase | 1 | |
| | Myb/SANT-like DNA-binding domain | 1 | |
| | Germin-like protein subfamily T member | 1 | |
| | Uncharacterized | | 2 |
| | Ripening-related protein | | 1 |
| Inorganic ion transport and metabolism | Calcium-selective channel | | 1 |
| Lipid transport and metabolism | lipolytic acyl hydrolase (LAH) | 1 | |
| Posttranslational modification, protein turnover, chaperones | Peptidase C1 family | 1 | |
| | heat shock protein 70 | 1 | 1 |
| Replication, recombination and repair | zinc finger | 1 | |
| Secondary metabolites biosynthesis, transport and catabolism | Peroxidase class III family | 2 | |
| | Dehydrogenase | 1 | |
| | cytochrome P450 | 1 | |
| Signal transduction mechanisms | Wall-associated receptor kinase | 3 | |
| | disease resistance | 2 | |
| | Protein kinase | 1 | |
| | PAN-like domain | 1 | |
| | NB-LRR | 1 | |
| | EF hand | | 1 |
| Transcription | mTERF | 1 | |
| | mitochondrial transcription termination factor protein | 1 | |
| | ARF (Auxin response factors) | | 1 |