Title:

- Regional Centromere Configuration in the Fungal Pathogens of *Pneumocystis* Genus

Authors

Ousmane H. Cissé [1]∗, Shelly Curran [1], H. Diego Folco [2], Yueqin Liu [1], Lisa Bishop [1], Honghui Wang [1], Elizabeth R. Fischer [3], A Sally Davis [4], Christian Combs [5], Sabrina Thapar [1], John P. Dekker [6], Shiv Grewal [2], Melanie Cushion [7], Liang Ma [1], and Joseph A. Kovacs [1]

Affiliations

[1] Critical Care Medicine Department, Clinical Center, National Institutes of Health, Bethesda, Maryland, USA.
[2] Laboratory of Biochemistry and Molecular Biology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, USA.
[3] Microscopy Unit, Research Technologies Branch, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, Montana, USA.
[4] Diagnostic Medicine/Pathobiology, Kansas State University College of Veterinary Medicine, Manhattan, USA.
[5] National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD
[6] Bacterial Pathogenesis and Antimicrobial Resistance Unit, National Institute of Allergy, and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA
[7] Department of Internal Medicine, College of Medicine, University of Cincinnati, Cincinnati, Ohio, USA.

Correspondence to: ousmane.cisse@nih.gov and/or jkovacs@mail.nih.gov

One sentence summary: Insights into the formation of genic centromeres in fungal pathogens.

Abstract
Centromeres are constricted chromosomal regions that are essential for cell division. In eukaryotes, centromeres display a remarkable architectural and genetic diversity. The basis of centromere accelerated evolution remains elusive. Here we focused on *Pneumocystis* species, a group of Mammalian-specific fungal pathogens that form a sister taxon with that of the *Schizosaccharomyces pombe*, an important genetic model for centromere biology research. Methods allowing reliable continuous culture of *Pneumocystis* species do not currently exist, precluding genetic manipulation. CENP-A, a variant of histone H3, is the epigenetic marker that defines centromeres in most eukaryotes. Using heterologous complementation, we show that the *Pneumocystis* CENP-A ortholog is functionally equivalent to CENP-A^{Cnp1} of *Schizosaccharomyces pombe*. Using organisms from a short-term *in vitro* culture or infected animal models and ChIP-Seq, we identified CENP-A bound regions in two *Pneumocystis* species that diverged ~35 million years ago. Each species has a unique short regional centromere (< 10kb) flanked by heterochromatin in 16-17 monocentric chromosomes. They span active genes and lack conserved DNA sequence motifs and repeats. These features suggest an epigenetic specification of centromere function. Analysis of centromeric DNA across multiple *Pneumocystis* species suggest a vertical transmission at least 100 million years ago. Common ancestry of

48 *Pneumocystis* and *S. pombe* centromeres is untraceable at the DNA level but the overall
49 architectural similarity could be the result of functional constraint for successful chromosomal
50 segregation.
51
Significance Statement
Pneumocystis species offer a suitable genetic system to study centromere evolution in pathogens
because of their phylogenetic proximity with the nonpathogenic yeast *Schizosaccharomyces*
pombe, a popular model for cell biology. We used this system to explore how centromeres have
evolved after divergence of the two clades ~460 million years ago. To address this question, we
established a protocol combining short-term culture and ChIP-Seq to characterize centromeres in
multiple *Pneumocystis* species. We show that *Pneumocystis* have short epigenetic centromeres
that function differently from those in *S. pombe*.
60
MAIN TEXT
62
Introduction
Centromeres are genomic locations where spindles attach during chromosomal segregation. They
are essential for cell division. Errors in the chromosomal segregation can result in aneuploidy and
cell division defects with disastrous consequences such as cancer. In fungi, centromeres are
fragile sites that are often involved in karyotype variability (*1, 2*) and drug resistance (*3*). In most
eukaryotes, centromeres are defined by the presence of CENP-A nucleosomes, a centromere
specific histone H3 variant that is essential for the localization of known kinetochore components
(*4*). There is a remarkable diversity in centromere structures ranging from sequence-dependent
(point) centromeres to epigenetically-regulated (regional) centromeres.
Point centromeres are exemplified by *Saccharomyces cerevisiae* with genetically defined
125 bp long centromeres containing conserved DNA elements (I, II and III) that serve as anchors
for the recruitment of the centromeric DNA binding factor 3 (CBF3) complex (*5, 6*). Regional
centromeres are much longer, lack universally conserved DNA patterns, and are often made up of
repetitive DNA (*7*). Regional centromeres are not strictly defined by DNA but are rather
epigenetically regulated; however, which factors contribute to this epigenetic specification of
centromeres remains elusive. In fungi, regional centromeres are classified as short (<20 kb, e.g.
Candida albicans (*8*)), intermediate (40 – 110 kb; e.g. *Schizosaccharomyces pombe* (*9*) and long
(150 – 300 kb, e.g., *Cryptococcus* (*10*)).
Pneumocystis is a genus of pathogenic fungi that exclusively infect mammals and remain
unculturable. They belong to the Taphrinomycotina subphylum and form a monophyletic clade
with fission yeasts *Schizosaccharomyces*, with a separation time estimated at ~460 million years
ago (*11, 12*). In addition to these two classes there are other classes, all of which are plant or soil
adapted organisms. Compared to *S. pombe*, which is a tractable organism that is widely used as
model for chromosome and cell biology, *Pneumocystis* species have streamlined genomes due to
substantial gene losses during their transition to animal parasitism. Recently, the genomes of
multiple *Pneumocystis* species have been sequenced (*12-15*). However, centromeres were not
defined in these because they could not be predicted bioinformatically without reference for
synteny based detection. Alternative methods such as chromosome conformation capture assay
(4C), that have been used to predict centromere loci in fungal genomes (*16*) are not applicable to
Pneumocystis because they require large quantities of pure DNA from synchronized cell cultures.
No long-term culture system exists and during purification of *Pneumocystis* organisms from
infected lungs it is virtually impossible to completely eliminate host cell DNA. *Pneumocystis*
have retained CENP-A (*12, 15*), which presumably binds to centromeres, and most of the
kinetochore proteins. Some data suggest that *Pneumocystis* centromeres differ structurally from
those of *S. pombe*. For example, the heterochromatin protein Swi6/HP1 (*17*), which is required

for <u>activation of</u> replication origins at the centromeres flanking regions (pericentromeres) has not
been found in *Pneumocystis* genomes (*18*). The RNA interference (RNAi) pathway, which is
essential to centromere function, has been lost in *Pneumocystis* (*19*), further supporting
mechanistic difference with *S. pombe*.
In this work, we undertook to characterize *Pneumocystis* centromeres and compare them
to *Schizosaccharomyces pombe* centromeres. <u>We found that *Pneumocystis* have small regional</u>
<u>centromeres that are defined by CENP-A and flanked by heterochromatin, resembling those of *S. pombe*.</u>
<u>However, *Pneumocystis* species lack orthologs for genes required for centromere function and</u>
<u>maintenance in *S. pombe*, suggesting that its centromeres may function differently.</u>
107
Results
Survey of genes related to centromere function in Taphrinomycotina fungi
To understand the kinetochore evolution in Taphrinomycotina fungi, we screened the predicted
proteomes of 14 species spanning six orders, *Pneumocystis* ($n = 7$), *Schizosaccharomyces* ($n = 3$),
Taphrina ($n = 1$), *Saitoella* ($n = 1$), *Protomyces* ($n = 1$) and *Neolecta* ($n = 1$), as well as other
fungi to have a broad overview of kinetochore evolution.
Using a combination of protein domain hidden Markov models and BLAST scans
followed by phylogenetic assignment (see methods), we searched for homologs of CENP-A[Cnp1],
CENP-C[Cnp3], CCAN (Constitutive Centromere Associated Network) and KMN (KNL-1/Mis12
complex/Ndc80 complex) proteins. We extended our searches to pathways required for
centromere function and chromosomal segregation (heterochromatin, RNAi, and DNA
methylation) (**Supplementary figure 1**).
CENP-A, CENP-C, and outer kinetochore proteins are conserved across
Taphrinomycotina. The CENP-A histones from *Pneumocystis* species, henceforth referring to as
*pn*CENP-A, display a high overall conservation (89% average protein identity). Two genes of the
inner kinetochore (CCAN) that are not found in *Pneumocystis* are innovations in *S. pombe* (*fta2*
and *fta3* genes). The *fta2* and *fta3* gene products associate with the central core (cnt) and
innermost repeats (imr) region of the centromere in *S. pombe* (*20*). No ortholog for the *mis*17
gene can be found in *Pneumocystis*. In *S. pombe*, *mis*17 encodes a member of the Mis6-Mal2-
Sim4 multiprotein required for CENP-A recruitment (*21*). The *S. pombe* centromere-associated
protein B genes (*cbp1*, *cbh1* and *cbh2*), which are derived from the domestication of *pogo*-like
transposases (*22*) have no identified homologs in other fungi.
The chromosomal passenger complex (CPC) is a heterotetrametric complex composed of
the Aurora B kinase Ark1 and three regulatory components Pic1 (inner centromere protein), Bir1p
(Survivin) and Nbl1 (Borealin) which mediates chromosome segregation and cytokinesis (*23*).
This pathway is conserved in *Pneumocystis* and *S. pombe*.
CENP-A recruiting complex, which includes Mis18 and Mis16 (*24*) as well as the CENP-
A histone chaperone Scm3 required for CENP-A loading at the centromere (*25*) are conserved in
Pneumocystis, as are the monopolin complex genes that coordinate the kinetochore microtubules
attachment during mitosis (*26*).
The network for heterochromatin formation in *Pneumocystis* seems intact with the
presence of the Clr4 methyltransferase complex (Clr4/SUV39H). *Pneumocystis* species have a
single chromo domain containing protein that bear similarity with *S. pombe* Swi6 and Chp2 which
are HP1 inparalogs (**Supplementary figure 2**).
DNA methyltransferases (DMTs) are divided into five classes (DNMT1, DNMT2,
DNMT3/DRM, Masc1/RID and DNMT5)(*27*). Canonical DNA methylase (DMTs) are not
detected in any of the seven sequenced *Pneumocystis* species (**Supplementary table 1**), which is
consistent with previous studies (*28, 29*).
Overall, the *Pneumocystis* kinetochore (CCAN) protein catalog is relatively conserved and
similar to those other Taphrinomycotina fungi.
148

*pn*CENP-A[Cnp1] localizes to the nucleus

Centromeres are identified by tracking CENP-A binding regions. Precise nuclear localization of CENP-A is required for accurate chromosomal segregation. To determine if *pn*CENP-A localizes in the nucleus, we used a anti – *Pneumocystis* CENP-A antibody, in conjunction with an anti-*Pneumocystis* antibody (*30*) and DAPI. Using indirect confocal immunofluorescence microscopy, we found that *pn*CENP-A displays a nuclear peripheral localization in replicating organisms from infected lung tissue and cultures (**Figures 1A and 1B**).

Fig. 1. Nuclear peripheral localization of CENP-A in *Pneumocystis* cells. (A) confocal immunofluorescence microscopy shows that *pn*CENP-A (red) is present a nuclear periphery (blue) inside *Pneumocystis carinii* organisms (green). A cluster of organisms from an-infected rat lung tissue section is presented and labeled with an anti CENP-A antibody (red), an anti-*Pneumocystis carinii* antibody 7C4 (green) and 4′,6-diamidino-2-phenylindole (DAPI, blue). The image is a maximum projection of a 5 µm thick z-stack. Bottom panels 1-8 are magnified fields. Most anti *pn*CENP-A labels overlap or are close in proximity with DAPI (represented by a pink color produced by the overlap between red and blue colors). Some organisms are not labelled by 7C4 antibody (e.g., box 5), which suggest that the epitope is not expressed. This field does not contain any host cell (rat). Scale bar =□2 µm. (**B)** in plane and orthogonal views of immunofluorescence labelling of *P. carinii* organisms co-cultured with mammalian cells for seven days. Organisms were triple labeled using the same an anti CENP-A antibody (red), a different anti-*Pneumocystis* antibody (RAE7) targeting *Pneumocystis* major surface glycoproteins (green) and DAPI (blue). Panels 1-3 are magnified fields. CENP-A is located inside *Pneumocystis* cells at the periphery of the nucleus. Few c are only labeled with the DAPI and do not express detectable levels of CENP-A and major surface glycoprotein. A rat cell nucleus is labelled with the letter 'H'. Scale bar =□1 µm.

*pn*CENP-A is functional in *S. pombe,* and supports viability and centromere loading

To confirm that the putative *cenp-a* gene from *P. murina* (*PmCENP-A*) encodes a functional CENP-A protein, we expressed *PmCENP-A* in *S. pombe*. *P. carinii CENP-A* has only one amino acid difference with *P. murina* (**Figure 2A**). *PmCENP-A* was codon-optimized, expressed under the endogenous *S. pombe cnp1*[+], the CENP-A ortholog, gene promoter and subcloned into a standard *LEU2*-based multicopy plasmid. In addition, the gene product was tagged with GFP at their N-terminus since it was shown that C-terminal tagging (*i.e.*, *cnp1*[+]*-GFP*) displays growth retardation (*31*). We first assessed the ability of *PmCENP-A* to rescue the lethality of *cnp1Δ* cells. We performed plasmid shuffling assays, in which *cnp1Δ* cells were initially kept viable by supplying a copy of *cnp1*[+] on a plasmid carrying the *ura4*[+] selection marker. A rescue plasmid carrying *PmCENP-A* and the *LEU2* selection marker was then transformed into the cells, followed by growth on medium containing 5-Fluoroorotic acid (FOA), which selects for loss of the *ura4*[+]-marked *cnp1*[+] plasmid (**Figure 2B**). As seen with control cells expressing the *S. pombe* GFP-CENP-A gene, we found that cells expressing *PmCENP-A* also fully support viability of *cnp1Δ* cells as they grew upon FOA selection (**Figure 2C**). We also observed no differences in growth between *cnp1Δ* cells expressing either *S. pombe* or *P. murina* CENP-A (**Supplementary figure 3**). We next asked if *PmCENP-A* can also rescue lethality of *cnp1-76* cells, a thermosensitive mutant, whose mutation T74M lies at the CATD (CENP-A targeting domain) region (*32*). As previously reported (*33*), we found that the *S. pombe* CENP-A tagged with GFP restored growth

of *cnp1-76* cells at restrictive temperatures (i.e., 34°C). Importantly, *P. murina* counterpart was also able to rescue the thermosensitive phenotype (**Figure 2D**) proving the functionality of this gene. In interphase fission yeast displays the Rabl configuration, in which centromeres cluster to the spindle pole body while telomeres associate to each other near nuclear periphery (*34, 35*). As expected, expressing GFP-CENP-A from *S. pombe* resulted in cells displaying single fluorescence foci that colocalized with a *tetO* array inserted at *cen2* (NB: a TetR-tdTomato fusion is expressed to bind to the *tetO* array and this locus is referred to as *cen2-tetO-tdTomato*) (**Figure 2E**). Consistent with the mentioned rescue of *cnp1-ts* and *cnp1Δ* cells and further supporting their role as functional CENP-A protein, *P. murina* CENP-A protein also colocalize with *cen2-tetO-tdTomato* and remarkably exhibited single foci, albeit the *P. murina* CENP-A protein also colocalized with *cen2-tetO-tdTomato* and, remarkably, exhibited single foci, albeit the intensities of centromere foci were dimmer than that of the *S. pombe* counterpart (**Figure 2F**). Finally, to confirm the specific localization of *Pm* CENP-A to centromeres, we performed ChIP-qPCR (**Figure 2G**). Noteworthy, we observed that *Pm* CENP-A localizes to central core regions (i.e., *cc1&3, cc2:ura4$^+$*) similarly to the fission yeast counterpart, albeit the enrichment was lower consistent with the results from live-cell imaging. Additionally, *Pm* CENP-A localization seems to be specific as no detectable enrichment was found at pericentromeric outer repeats coated with heterochromatin (i.e., *dg*) and euchromatin locations (i.e., *fbp1*). Together, these results confirm that *P. murina* CENP-A gene encodes a bona fide CENP-A protein.

> **Fig. 2. *Pneumocystis* CENP-A supports viability and centromere loading in a heterologous system. (A)** Protein sequence alignment of full-length CENP-A^{cnp-1} orthologs in *Schizosaccharomyces pombe*, *Pneumocystis murina* and *P. carinii*. Each *Pneumocystis* species only infects a single mammalian host: *P. murina* (infecting mice) and *P. carinii* (rats). The ~25 amino acid insert in *Pneumocystis* CENP-A N-terminal regions is unique to this genus. (**B**) Schematic of the plasmid-shuffling assay used in C. The inviability of *cnp1Δ* cells is masked by the presence of *cnp1$^+$* on a *ura4$^+$* marked plasmid. After introducing the *LEU2* marked rescue plasmid, cells that have lost the *ura4$^+$* marked plasmid are selected on medium containing FOA, leaving only those with the rescue plasmid. NB: *S. cerevisiae* *LEU2* gene complements the *S. pombe leu1-32* mutant. (**C**) Rescue experiments using the plasmid-shuffling assay described in B to test the ability of multicopy plasmids expressing GFP-tagged CENP-A transgenes to rescue *cnp1Δ* cells. Growth was assayed at indicated media using 10-fold serial dilutions. (**D**) Rescue experiment of fission yeast *cnp1-76* thermosensitive cells using same plasmids as in C. Growth was assayed at indicated temperatures using 10-fold serial dilutions. The transgene encoding *S. pombe* CENP-A serves as a non-temperature-sensitive control. (**E**) Representative images of *cen2-tetO-TdTomato* strains expressing the indicated GFP-tagged CENP-A transgenes in multicopy plasmids. GFP and TdTomato fluorescence as well as brightfield images are merged. Magnified views of boxed regions are shown at the bottom. (**F**) Integrated fluorescence intensity of GFP foci was measured and plotted for the strains used in panel E. The median (bar) and interquartile range (error bars) are shown. n>409. AU, arbitrary units. (**G**) Anti-GFP ChIP-qPCR analysis of indicated loci for strains expressing GFP-tagged CENP-A transgenes as single copy integrations. %IP represents the percentage of input that was immunoprecipitated. Error bars denote the SD (n = 3). For each strain, comparison of %IP between loci was performed by one-way ANOVA and Holm-Sidak test for multiple comparisons (P<0.001). Mean values marked with letters (a or b) indicate results that are significantly different from each other. Abbreviations: *cc1&3*, *S. pombe* centromere central core 1 & 3;

249   *cc2:ura4⁺*, *ura4⁺* insertion at central core 2; *dg*, a class of heterochromatic outer
250   repeats within pericentromeric regions (control); *fbp1*, euchromatic locus (control).
251

pnCENP-A binds to single genomic foci in *Pneumocystis* replicating cells

In *S. pombe*, CENP-A$^{\text{Cnp1}}$ is deposited at the centromeres during the G2 phase of the cell cycle
(*31*). Although *Pneumocystis* cell replication is not fully understood, these organisms take 5 to 8
days to replicate *in vivo* (*36*). To determine pnCENP-A bound genome regions, we established a
short-term co-culture system for *P. murina* and *P. carinii* (**Figure 3A**). We assessed organism
growth using quantitative PCR targeting the single copy gene dihydrofolate reductase (*dhfr*); a
typical growth curve shows a decline from day 0 to 7 followed by a gradual increase (**Figures 3B
and 3C**). Population growth is further supported by the presence of mitotic or fusing cells 7 days
post culture, though some cells do not appear to be dividing (**Figure 3D**).

**Fig. 3. *Pneumocystis* CENP-A supports viability and centromere loading in a
heterologous system. *Pneumocystis* CENP-A binds to single genomic foci in
replicating cells. (A)** Study workflow and sample preparation. *P. murina* and *P.
carinii* were obtained from CD40 ligand knock-out female mice and
immunosuppressed Sprague-Dawley male rats, respectively, and cultured on a co-
culture of human lung adenocarcinoma cells (A549) and immortalized murine lung
epithelial type 1 cells (Let-1) for 14 days. Species phylogeny and estimated
speciation timing are presented. Animal icons were obtained
from http://phylopic.org under creative commons
licenses https://creativecommons.org/licenses/by/3.0/: mouse (Anthony Caravaggi;
license CC BY-NC-SA 3.0); and rat (by Rebecca Groom; license CC BY-NC-SA
3.0). (**B**) *P. murina* population growth in duplicate wells were measured by
quantitative PCR targeting a single copy dihydrofolate reductase (*dhfr*) gene over
14 days. Error bars represent the standard deviation (*n*=2). (**C**) *P. carinii* growth
measurement by qPCR targeting *dhfr* gene. Error bars represent the standard
deviation (*n*=2). (**D**) Electron micrograph showing a possibly dividing *P. murina*
trophic form seven days post culture (black arrow). The field also include a non-
dividing trophozoite (white arrow). 2 μm scale bar.

We mapped CENP-A bound genomic regions in all *P. carinii* and *P. murina* chromosomes using
ChIP-from cultured organisms collected days 0 (normalized initial inoculum), 7 and 14.
    In *P. murina*, each of the 17 chromosome-level scaffolds display a single region
containing peaks of CENP-A at all three time points (**Figures 4A and 4B, Supplementary figure
4**). We confirmed CENP-A enrichment by ChIP-qPCR (**Supplementary figure 5A**). CENP-A
enrichment has a bimodal distribution, suggesting two distinct populations of CENP-A in
chromosome arms. The two CENP-A peaks are separated by a 1-kb noncoding region that lacks a
conserved DNA motif. Although the significance of two peaks of CENP-A in *Pneumocystis* is
unclear, in *S. pombe* CENP-A cores are interspersed with H3K4me (*37*). CENP-A enrichment
levels fluctuate in according to the growth curve, that is the highest level is observed at day 0
followed by a decline at day 7 and a recovery at day 14. In *S. pombe*, CENP-A$^{\text{Cnp1}}$ enrichment at
the centromeres correlates with a depletion of histone H3 (*38*). At day 7, the H3 occupancy
(expressed as ratio between H3 and H4) is significantly reduced at centromeres relative to
flanking regions (30 kb) as estimated by ChIP-Seq (Wilcoxon test; $P < .0001$; **Supplementary
figure 6**), which suggest that CENP-A substitutes the canonical H3. However, H3 depletion at the
centromeres does not persist after 14 days of culture (**Supplementary figure 7A**).

P. carinii also displays monocentric chromosomes enriched with two CENP-A peaks at the three time points by ChIP-Seq (**Figures 4C and 4D, Supplementary figure 8**) and ChIP-qPCR (**Supplementary figure 5B**). H3 depletion is observed at day 14 (**Supplementary figure 7B**). In contrast to *P. murina*, we found a significant reduction of CENP-A enrichment at day 14 despite a population growth (**Figure 3C**). Also not observed in *P. murina*, at day 14 low levels of CENP-A are present outside the primary CENP-A binding region in several chromosomes (**Supplementary figure 8**), which correspond to a non-specific displacement of CENP-A. In budding yeasts and chicken cells, low levels of CENP-A molecules have been shown outside the core domain (CENP-A cloud), which contribute to centromere plasticity (*39*).

In both species CENP-A bound regions span 4.8 to 8.0 kb, with an average length of 6.7 kb, and have an average of 3.9% lower GC content than the rest of the genome (**Table 1**). Using dot plot analysis of a concatemer of different centromeres, each CENP-A binding region appears unique within the genome (**Figures 4E and 4F**) and lacks a shared conserved DNA motif (see methods). Because most centromeres in fungi are associated with repetitive DNA (e.g., DNA transposons and retrotransposons), we searched for known signatures of repeats and found no significant overlap of repeats with centromeres (**Supplementary figures 6 and 9**).

Fig. 4. *Pneumocystis* **displays seventeen centromeres. (A)** Scaled ideogram of 17 chromosomal level scaffolds (grey) of *P. murina* genome showing CENP-A binding regions (constricted areas). (**B**) CENP-A binding regions delineate putative centromeres (CENs) in *P. murina* genome. Color coded peaks represent enrichment of immunoprecipitated DNA (IP DNA) relative to controls (Input DNA) in *P. murina* organisms (input subtracted). Only ChIP-Seq data from *Pneumocystis* cocultured with A549 and Let-1 cells for seven days are presented. Data for the full experiment covering days 0, 7 and 14 are presented in Supplementary material. The 20-kb windows of the enrichment peak are presented. Each scaffold displays two peaks labelled M (Major) and m (minor) according to the enrichment level. (**C**) Scaled ideogram of 17 chromosomal level scaffolds (grey) of *P. carinii* genome showing CENP-A binding regions (constricted areas). (**D**) CENP-A binding regions delineate putative centromeres (CENs) in *P. carinii* genome. Color coded peaks represent enrichment of immunoprecipitated DNA (IP DNA) relative to controls (Input DNA) in *P. carinii* organisms cocultured with A549 and Let-1 cells for seven days (input subtracted). Data for the full experiment covering days 0, 7 and 14 are presented in Supplementary figure 6 and Supplementary dataset. (**E**)A DNA dot plot of CENP-A binding regions in *P. murina* showing regional self-similarity. The main diagonal represents the sequence alignment with itself. Lines off the main diagonal which are repetitive patterns within the sequences are not observed. The plot shows that each CEN is unique within the genome. (**F**) DNA dot plot of CENP-A binding regions in *P. carinii* showing that each CEN is unique and repeat-free.

Pneumocystis **centromeres contain active genes.**
By cross referencing annotated gene locations with those of *pn*CENP-A bound regions, we found that *P. murina* and *P. carinii* centromeres encode 74 and 58 genes respectively; all of them are conserved in the genomes of both species except a prefoldin gene that is lost in *P. carinii* (**Supplementary Table 2**). Of the 74 *P. murina* centromeric genes, 73 were expressed based on RNA-seq and four were further detected by protein mass spectrometry mapping (LC-MS); of the

58 *P. carinii* genes, 56 are expressed and 53 were detected by LC-MS. Analysis of the predicted function of these genes revealed housekeeping functions without major differences compared to randomly sampled genes.

Centromeres are flanked by heterochromatin.

In *S. pombe*, H3K9me2/3 marking of chromatin is associated with the repression of centromeres, subtelomeres, ribosomal rDNA and the mating locus (*37*). There are two types of chromatins: euchromatin, the lightly packed form of the chromatin enriched with H3K4me2 (di-methylation of the 4th lysine residue of the histone H3), which is associated with active transcription, and heterochromatin enriched with H3K9me2 and H3K9me3 (di and tri-methylation of the 9th lysine residue of the histone H3 protein). To test if this feature is shared in *Pneumocystis*, we performed ChIP-Seq with antibodies targeting histones H3 modifications (H3K9me2/3 and H3K4me2). In *Pneumocystis*, we detected broadly distributed peaks of H3K9me2 and H3K9me3 bordering the putative centromeres delineated by CENP-A narrow peaks (**Figures 5A and 5B**, **Supplementary figures 6 and 9**). This configuration of chromatin markers is specific to centromeric regions. H3K4me2 is correlated with H3K9me2 in the centromeres (Pearson rho = .72) but not at the whole genome level (r = .05). However, research has shown that the histone modifications H3K4me2 and H3K9me2 decorate euchromatin and heterochromatin regions, respectively (*40*). Therefore, the correlation observed between H3K4me2 and H3K9me2 may be due to differences in growth kinetics for the *Pneumocystis* population of cells used in the ChIP analysis. Furthermore, similar to *S. pombe* (*37*), a small amount of H3K4me2 is present at the centromeres in *Pneumocystis* (**Figures 5A and 5B**, **Supplementary figures 6 and 9**). These results suggest that *Pneumocystis* centromeres are flanked by high levels of heterochromatin.

DNA methylation is frequently associated with centromeres in fungi, where they silence repeats (*41, 42*). Similar to *S. pombe*, DNA methylation has been predicted to be absent in *Pneumocystis* based on the absence of DNA methyltransferases (DMTs) (*29, 43*). To determine whether DNA methylation plays a role in *Pneumocystis* centromeres, we performed bisulfite sequencing (5 methylcytosine) in *P. carinii*. The overall level of 5mC DNA methylation as measured by the average weighted methylation percentage is 0.6% for *P. carinii* at the CG dinucleotides (**Supplementary figure 10**). These levels are in range with reported levels for other fungi e.g., *Verticillium* (0.4%) (*44*). The presence of 5mC methylated DNA bases despite the absence of recognizable DNA methylases in *Pneumocystis* requires further investigation. To assess the potential role of DNA methylation in centromere function in *Pneumocystis*, we analyzed the DNA methylation patterns over different genomic features (genes, intergenic spacers, and centromeres). However, there is no significant difference among centromeric or pericentromeric regions (defined as 30 kb flanking the centromeres) (Mann-Whitney U-test *P*-value > 0.3), and the randomly selected genomic regions (genomic background). These results suggest that 5-mC DNA methylation is not required for centromere function in *Pneumocystis*. The absence of repeats in the centromeres and the presence of actively transcribed genes suggests that DNA methylation does not induce an epigenetic silencing in *Pneumocystis* centromeres.

Fig. 5. Centromeres are flanked by heterochromatin and contain active genes. (**A**) Genomic view of chromosome 2 of *P. murina* genome subsequently showing annotated genes (directed grey boxes), DNA repeats (not present), percent GC content (blue), ChIP-Seq read coverage distribution (BPM normalized over bins of 50 bp; input subtracted) of CENP-A, histones H3 and H4 ratio, heterochromatin-associated modifications (H3K9me2 and H3K9me3), euchromatin (H3K4me2) and gene expression (RNA-seq) in relation with centromeres. (**B**) Genomic view of

chromosome 2 of *P. carinii* genome with the same features presented as *P. murina*.
A duplicated copy of *copia*-retrotransposon is presented (red arrows), which is
present in syntenic regions in other *Pneumocystis* genomes. The two presented
chromosomes are syntenic.

Centromere conservation during speciation

Pneumocystis murina and P. carinii species have diverged ~ 35 Mya (*12*). Yet, orthologous
regions act as centromeres in both *species,* which suggest a vertical transmission (**Figure 6A**).
Syntenic regions can be found across five additional *Pneumocystis* species which speciation spans
at least 100 million years (**Supplementary figure 11**).

To investigate footprints of selection acting on centromeres, we computed conservation
scores (PhastCons) from whole genome alignments. Conservation scores range from 0 to 1 and
represent probabilities of negative selection. *Pneumocystis* genomes encode a large multicopy
major surface glycoprotein (MSG) gene family. MSGs are highly polymorphic and evolve rapidly
due to their role in antigenic variation during mammalian host infection (reviewed in (*45*)). Using
MSGs as control for evolutionary speed, we found that centromeres and flanking regions are
substantially more conserved than MSGs and similar to genomic background (**Figures 6B and
6C**). The levels of centromeric conservation are variable across the chromosomes, with the
flanking regions being more conserved than the cores (defined as a 1-2 kb region in the middle of
each centromere) (**Supplementary figure 12**). Given that macrosynteny at the centromeric
regions is conserved across multiple *Pneumocystis* species, this suggests that centromere locations
are maintained by positive selection. This would be consistent with the hypothesis that
centromere positioning tends to be conserved in obligate sexual fungi due to their role in meiosis
(*46*).

Centromeres contribute to karyotypic diversity in some fungi (*2*). However, we found
little support for this hypothesis here because most centromeres do not overlap with chromosomal
breaks (only 3 of 17 in the *P. carinii* versus *P. murina* pairwise comparison). This is reminiscent
of other fungi such as *Verticillium* species where centromeres do not account for most of the
karyotype variation (*47*).

**Fig. 6. Centromere locations, not sequences, are evolutionarily constrained by
positive selection. (A)** Genome organization and synteny in *Pneumocystis*. Circos
plots depicting pairwise *P. carinii* and *P. murina* genome synteny. Rodent
infecting *Pneumocystis* (*P. carinii* and *P. murina*) have 17 chromosomes. Colored
connectors indicate regions of synteny between species. Centromeres that overlap
with recent chromosomal breakpoints are indicated (*). The square highlights *P.
carinii* centromere 4 displayed in the panel b. (**B**) Genome view of centromere 4 in
the *P. carinii* genome. Genes are represented by directed boxes (grey for protein
coding genes and cyan for polymorphic major surface glycoprotein genes),
*pn*CENP-A binding region (centromere) and sequence conservation scores which
were calculated from whole genome alignments of *P. carinii. P. murina* and *P.
wakefieldiae* (PhasCons). The phastCons scores represent probabilities of negative
selection and range between 0 (no conservation) and 1 (total conservation). (**C**)
Boxplot of conservation scores per genomic context summarized for the
centromere 4 in *P. carinii* genome, 30-kb regions flanking the centromeres
(Cenflk), major surface glycoproteins encoding regions (Msg) and random
genomic background (Bckg). Msgs are fast evolving proteins potentially involved
in antigenic variation. Background data were obtained from randomly selected

145   intervals ($n = 1 \times 10^6$) from genomic regions excluding above mentioned regions
146   (CEN, Cenflk and Msg). Statistical differences for the indicated comparisons were
147   obtained using one-sided nonparametric Mann-Whitney test; *P*-values < .0001,
148   ****; *P* < .001, ***; *P* < .01, **; *P* < .05, *. Data for all 17 centromeres are
149   presented in Supplementary material.

Discussion

151
In the current study we have identified centromeres of two *Pneumocystis* species by showing that
the centromeric histone CENP-A binds to gene rich genomic regions that are flanked with
heterochromatin (**Figure 7**).

Fig. 7. Model of regional centromere structures in *Pneumocystis* and other
representative fungi. On the left is presented the phylogeny of selected fungi
inferred from maximum likelihood phylogenetic analysis of shared core protein
orthologs. The overall centromere structures for *Pneumocystis carinii* (details
supporting our model are provided in the text), *Schizosaccharomyces pombe* (*48*),
Candida albicans (*8*), *Zymoseptoria tritici* (*46*) and *Cryptococcus neoformans* (*49*)
are used as representative to showcase the diversity of regional centromeres in
fungi. Animal pathogens are highlighted in pale olive and the plant fungal
pathogen *Zymoseptoria tritici* is in pale green. For sake of brevity, only
Pneumocystis carinii is presented. In the middle are presented DNA structures of
centromeres. *P. carinii* has 17 centromeres (one for each of its 17 chromosomes)
that share the same overall architecture. Centromeres are delineated by a localized
enrichment of the centromeric histone CENP-A, which overlaps with a reduction
of the canonical histone H3 (inverted grey triangle). Centromeres are flanked by
heterochromatin H3K9me (stands here stands for both H3K9me2 and H3K9me3).
All 17 *Pneumocystis carinii* centromeres span active genes (dark grey boxes).
Each centromere sequence is different and lacks shared DNA sequence motif. *S.*
pombe has three centromeres that share the same overall structure in which a
central core (*cnt*) domain is surrounded by innermost repeats (*imr*) and outer
repeats (*otr*). The *imr* repeats incorporate clusters of transfer RNAs (tRNAs) that
play a role in restricting CENP-A spread. *S. pombe* centromeres are flanked by
heterochromatin (H3K9me). Genes are found 0.75-1.5 kb beyond the limits of the
centromeres. *C. albicans* has 8 unique and different centromeres that are gene free
and lack shared sequence motifs. *Z. tritici* has 21 centromeres ranging from 6 to 14
kb in size, that partially overlap with genes. *C. neoformans* has 14 centromeres that
are gene free and enriched with Tcn transposons.

This configuration suggests the presence of small epigenetically regulated centromeres (regional).
Within each genome, each chromosome has a unique non repetitive centromere. Syntenic regions
for each centromere are present in all seven currently sequenced *Pneumocystis* species. Cell
replication and chromosomal segregation pathways are unexplored in these species. Central to
these pathways are the centromeres.
As centromeres cannot be predicted bioinformatically because no suitable reference for
synteny analyses existed before this study, the locations and characteristics of *Pneumocystis*
centromeres were unknown. Moreover, there are several roadblocks for characterizing
centromeres in *Pneumocystis*, with the most significant one being the lack of continuous culture
and transfection tools. Here we utilized two *Pneumocystis* species in a short-term culture system
to determine where *pn*CENP-A, the epigenetic marker for centromeres, binds in the genome.

As a first step, we demonstrated that *Pn*CENP-A functions similarly to CENP-A^{Cnp1} in *S. pombe*, though less efficiently, validating its use in characterizing *Pneumocystis* centromeres. We then developed a *Pn*CENP-A based ChIP-Seq assay that showed a reproducible efficiency in defining centromeric regions of *Pneumocystis* species.

Unlike the phylogenetically related *S. pombe*, in *Pneumocystis* these regions overlap with active genes, but like *S. pombe*, these centromeres are flanked by heterochromatin. Our work also provides the first experimental evidence that DNA methylation occurs in these species, although it may not be involved in centromere function.

Despite their close phylogenetic relationship, *P. murina* and *P. carinii* have slightly different growth kinetics in our short-term culture system, which could explain the variations in the *Pn*CENP-A and H3 DNA binding profiles. The nonspecific binding of *Pn*CENP-A observed in *P. carinii* at day 14 is likely the result of a random dissociation of CENP-A molecules from the centromeres.

In *P. murina*, 5 out of the 17 chromosomes showed secondary CENP-A enrichment at non centromeric sites (Figure S4). These secondary CENP-A peaks are likely ChIP-Seq artefacts because they were only detected at day 7 post culture, and mostly overlap with highly expressed loci (Figure S8), which are susceptible to misleading signals in ChIP-Seq experiments (*50*).

The presence of genes within centromeres is rare and has only been described in rice (*51*) and the plant fungal pathogen *Zymoseptoria tritici* (*46*). Centromeres bearing genes are interpreted as young centromeres (neocentromeres), in which the genes are progressively inactivated. This hypothesis lacks support here because there is no sign of pseudogenization in these genes: nearly all are expressed and translated, they are conserved in all species, and many are involved in housekeeping cellular pathways that are presumably critical for organism survival.

Findings in *Pneumocystis* cannot be generalized to other Taphrinomycotina species. The determination of ancestral traits between *Pneumocystis* and fission yeast centromeres will require characterizing the centromeres in additional Taphrinomycotina species. The loss of RNAi is often associated with shortening of centromeres (*49*); consistent with this, *Pneumocystis* centromeres are much smaller than *Schizosaccharomyces* centromeres.

Our study will benefit from a microscopy validation of CENP-A loading kinetics when a reliable long term culture system becomes available. Live imaging would help to determine at which cell cycle stage CENP-A is loaded. A direct genetic confirmation of *Pneumocystis* centromeres will also be required when a long-term culture system and genetic manipulation tools will become available.

In summary, we have identified short regional centromeres in genetically intractable micro-organisms. Our results provide insights into the formation of centromeres in host-adapted fungal pathogens. Our ultimate goal is to use *Pneumocystis* centromeres to stabilize plasmids for future genetic manipulation. This is the first step along this path, which should lead to better understanding of the biology of *Pneumocystis* and facilitate the discovery of novel interventions to effectively control and prevent the disease caused by this pathogen.

Materials and Methods

Ethics and organism source
Studies involving mouse and rat samples were approved by the NIH Institutional Review Board (IRB) (IRB protocols 99-I-0084 and CCM 19-05, respectively), and rat *Pneumocystis* studies by the Animal Care and Use Committees of the Cincinnati VA Medical Center (protocol 20-11-08-01).

543   *P. carinii* organisms were collected from heavily infected lungs of corticosteroid-treated
544 Sprague-Dawley male rats; *P. murina* from heavily infected lungs of CD40 ligand knock out
545 mouse. The list of samples is presented in the **Supplementary table 3**.
546

Pneumocystis culture and growth quantification
P. murina and *P. carinii* were partially purified by Ficoll-Hypaque density gradient centrifugation
(*52*) and frozen at -80^{o}C in cell recovery media (Gibco). For short term *Pneumocystis* cultures,
A549 (ATCC) and LET1 (a gift from Dr. Paul Thomas, St Jude Children's Research Hospital)
cells were cultured in culture medium F12 media with 2.5% or 5% heat inactivated fetal bovine
serum (GIBCO) and Penicillin-Streptomycin (GIBCO), plated to approximately 60 – 80%
confluency and incubated for 24h at 37°C. The next day, frozen *Pneumocystis* vials were thawed,
washed in 50ml 1x PBS, and centrifuged at 2,000g for 20 minutes. *Pneumocystis* cell pellets were
resuspended in culture medium and added to the plated cells. Media were partially changed every
3 or 4 days. At set time points (days 7 and 14), wells were scraped for collection. Cell/organism
suspensions were centrifuged at 10,000g for 3 min, the supernatant was removed, and the pellets
were frozen at -80°C. Genomic DNA was extracted using QiAmp DNA Extraction Kit (Qiagen).
Quantitative PCR targeting the single copy *dhfr* gene was performed as described previously (*53*).
560

P. murina CENP-A^{Cnp1} complementation and ChIP-qPCR in *S. pombe*
Standard procedures were used for fission yeast growth, genetics, and manipulations (*54*). The
sequence encoding *P. murina* CENP-A gene was codon optimized for *S. pombe* codon bias,
synthesized commercially (GenScript, NJ) and subcloned into pS2 vector (*55*) generating
pPmCENP-A. *Nde*I-*Sph*I fragments from both pS2 and pPmCENP-A were subcloned into the
same sites of pREP81 vector to obtain *LEU2*-based multicopy plasmids that were used in
experiments presented in Fig. 2B-F. Alternatively, plasmids were linearized with *Nsi*I and
introduced by transformation to the *lys1$^+$* locus of *cen2-tetO-tdTomato* strain (*56*). These strains
bearing single-copy integrations were grown overnight at 30°C in rich medium YEA and used for
ChIP with 2 µl of anti-GFP (ab290, Abcam). PCR oligonucleotides detecting central core
(*cc1&3; ura4*), heterochromatic *dg* and euchromatin control *fbp1* were previously reported (*57*).
572

Orthologs identification
To find orthologs for kinetochore proteins, we constructed a database of 21 proteomes from fungi
15 Taphrinomycotina including *Pneumocystis* (*n* = 7) and 6 other representative fungi
(**Supplementary Table 1**). We queried the database with *S. pombe* proteins as seed using
BLASTp (*58*) with a minimum an e-value of 1.0 x 10^{-5} as threshold.
If many homologs were detected in *Pneumocystis* species, we separated orthologs from
out-paralogs by searching in orthologous groups inferred from the clustering of the whole
proteome dataset using OrthoFinder v2.5.2 (*59*). A sequence was considered as ortholog if it was
assigned to the same orthologous group as *S. pombe* protein query. We verified that the
assignment was consistent with EggNOG database v6.0 (*60*), which contains all *S. pombe*
proteins and a limited number of *Pneumocystis* proteins. If a *Pneumocystis* sequence was detected
by BLASTp but not assigned as an ortholog, we first identified conserved domains in all
sequences using InterProScan version 5.64-96.0 (*61*). Then we constructed multiple sequence
alignment containing all sequences including *S. pombe* query using MAFFT version v7.467 (*62*)
either with the option E-INS-I or L-INS-i depending on the protein domain architecture.
Alignments were manually curated using MacVector version 18.6.1 (https://macvector.com/) to
remove spurious aligned regions and used to infer an approximately maximum likelihood gene
tree using FastTree v2.1.10 (*63*). When necessary, a maximum likelihood phylogeny was
performed using IQ-TREE version 1.5.5 (*64*). For histones, we used protein sequences identified

by (*65*) to guide the classification. In case of inconclusive phylogenetic classification, we BLASTed *Pneumocystis* protein sequences against NCBI nr using a minimum e-value of 1.0 x 10^{-5} as threshold. Retrieved proteins were added to the existing alignment, and the phylogenetic inference was repeated.

If no homologs were found in *Pneumocystis* using *S. pombe* query, we used a different seed from another *Taphrinomycotina* if available (e.g., *Saitoella*). Alternatively, proteins from, *Saccharomyces cerevisiae*, *Neurospora crassa*, and *Cryptococcus neoformans* were used as seeds. If no homologs were detected, we used PSI-BLASTp (*58*). In parallel, if *S. pombe* query contains a conserved protein domain, we screened our fungal database with the corresponding Pfam domain using hmmsearch (HMMER version 3.3.2; http://hmmer.org)(*66*) with a cut off e-value of 0.1.

Because many genes could not be found in *Pneumocystis* proteomes, we scanned the genomes using TBLASTn (BLAST+ 2.13.0)(*67*) with an e-value of 0.1 as threshold. If putative genomic regions were found, we performed using a protein-to-genome alignment with Exonerate (*68*). Protein to genome alignments were translated in protein sequences, aligned, and used phylogenetic inference as described above.

To identify DNA methylases within the Taphrinomycotina, we identified annotated proteins with a DNA methylase domain (Pfam domain PF000145(*69*)) using hmmsearch (HMMER version 3.3.2; http://hmmer.org)(*66*) with a cut off e-value of 0.01. Although no hits were found in *Pneumocystis* proteomes, DMTs were detected in other fungi including *Neolecta irregularis* and *Saitoella complicata*, which is consistent with a previous study (*28*). *Pneumocystis* proteomes were queried using sequences from *N. irregularis* and *S. complicata* using BLASTp (*58*) with an e-value of 1 x 10^{-5} as cut off. This strategy identified 52 *Pneumocystis* proteins with local similarities, which were combined with all DMTs identified, aligned using Muscle v3.8.31 (*70*) and used for phylogenetic inference using FastTree. Then, we BLASTed *Pneumocystis* proteins against NCBI nr database (last accessed Thu Oct 12 12:07:56 EDT 2023), retrieved the top hit sequences and added them to the multiple alignment. All 52 *Pneumocystis* sequences belong related but distinct protein families (e. g. DNA helicases and SNF2 chromatin remodeling ATPases).

Electron microscopy
P. murina and A549-LET1 cells described above were grown on Thermanox™ coverslips (Ted Pella, Redding, CA, USA), harvested at day 7, fixed with 2% paraformaldehyde/2.5% glutaraldehyde in 0.1 M Sorenson's phosphate buffer and then post-fixed with 1.0% osmium tetroxide/0.8% potassium ferricyanide in 0.1 M sodium cacodylate buffer for 1 hour, washed with buffer then stained with 1% tannic acid in dH$_2$O for 1 hr. After additional buffer washes, the samples were further osmicated with 2% osmium tetroxide in 0.1M sodium cacodylate, then washed with dH$_2$O. Specimens were then stained overnight at 4C with 1% aqueous uranyl acetate. The cells were then washed with dH$_2$O and dehydrated with a graded ethanol series, prior to embedding in Spurr's resin. Thin sections were cut with a Leica UC7 ultramicrotome (Buffalo Grove, IL) prior to viewing at 120 kV on a FEI BT Tecnai transmission electron microscope (ThermoFisher/FEI, Hillsboro, OR). Digital images were acquired with a Gatan Rio camera (Gatan, Pleasanton, CA).

Antibodies
To obtain antibody against *pn*CENP-A, we created an alignment containing CENP-A orthologs from *Rattus norvegicus* (rat), *Mus musculus* (mouse), *Schizosaccharomyces pombe* and multiple *Pneumocystis* species (**Supplementary figure 13A**). We deliberately avoided the C-terminus, which contains the highly conserved histone fold to minimize the risk of cross reactivity with the

host histones in partially purified *Pneumocystis* protein preparations. Immunogenic peptides were selected from the N-terminal regional. Peptides with sequence similarities with rat and/or mouse proteins (NCBI accession no. GCF_015227675.2 and GCF_000001635.27, respectively) were excluded using BLASTp (*58*) with word size of 6 and a minimum e-value of 1 as cut off as well as with pattern matches using Perl regular expressions. Additional searches were performed online at https://www.uniprot.org/peptide-search. We select one immunogenic peptide conserved in both *P. carinii* and *P. murina*. Two rabbits were immunized with 20 µg of the affinity purified peptide. The polyclonal antibody was purified by affinity column (GenScript). No cross reactivity against the host cell lysates was seen by western blots (**Supplementary figure 13B**). The following commercial antibodies targeting histones were used at 1:100 dilution for ChIP-Seq: H3 (ab1791, Abcam), H3K4me (39142, Active Motif), H3K9me2 (ab1220, Abcam), H3K9me3 (ab8898, Abcam) H4 (ab1015, Abcam). We verified that peptides used for immunization for commercial antibodies were conserved in *Pneumocystis* proteins.

Western blots
Pneumocystis proteins (antigens) were prepared using glass beads. Partially purified organism pellets were resuspended in 1X PBS buffer. 0.65 ml of 0.5 mm glass beads (Biospec Products, Inc) was added to the suspensions and vortexed for 5 minutes at 4°C. Beads were allowed to settle for less 30 seconds and the supernatant transferred to a new tube for sonication. The samples were centrifuged for 20, 000xg for 10 minutes and the supernatants were transferred to a new tube. The pellet was resuspended in 20 µl PBS.

For western blots, cell lysates and formalin-fixed chromatin preparations were run on Tris-Glycine SDS 4-20% WedgeWell gel (Thermo Fisher Scientific) and transferred on nitrocellulose membranes by incubation at room temperature for 30 minutes, followed by 1 hour incubation in blocking buffer (1X PBS + 5% milk + 0.05% Tween 20) with shaking. Nitrocellulose blots were incubated with primary antibodies (1:100) diluted in blocking buffer for an hour with shaking, followed by a washing and then 1 hour incubation with secondary anti rabbit or anti mouse IgG HRP goat polyclonal antibody (1:2000) (PRF&L) and developed with Pierce 1-Step^{TM} Ultra TMB-Blotting solution (Pierce).

Immunofluorescence microscopy
For culture, cells were plated in chamber well slide (Millicell EZ slide, Millipore). When cells were ready, the medium was removed and washed with cold PBS 1X, fixed with 4% formaldehyde for 10 minutes at room temperature, and washed three times. Cells were permeabilized with 0.2% Triton X-100 for 30 minutes and followed by a washing with 1X PBST (PBS with 0.05% Tween20).
Pneumocystis carinii infected lung tissues fixed in HistoChoice (VWR Life Science) were embedded in paraffin. Five µm sections were labelled with rabbit anti *Pneumocystis* CENP-A antibody (1:200), mouse monoclonal anti *Pneumocystis carinii* antibodies (7C4 and RAE7) (*71, 72*). The 7C4 antibody reacts with one or two uncharacterized antigens of 40 000 daltons and RAE7 reacts with the major surface glycoproteins of *P. carinii* (RAE7 antibody is a gift from Drs. Michael Linke and Peter Walzer, University of Cincinnati College of Medicine) and stained with Vector TrueView autofluorescence quenching kit with DAPI (Vector Laboratories). Alexa Fluor 555 (1:200)– labeled goat anti rabbit IgG (A270039, Invitrogen) and Alexa Fluor 488 (1:200) – labeled goat anti-mouse (A11029, Thermo Fisher Scientific) were used as secondary antibodies. Images were acquired using either a Leica SP8 confocal microscope with x 63x (1.4 NA) HC PL APO objective or a Zeiss 880 confocal microscope with a 63x (1.4 NA) PL APO objective. Images from the Zeiss were taken using 405, 488, and 561 nm excitation with emission bandwidths of 415-480nm, 500-550nm, and 565-700nm. Images from the Leica were taken using 405, 488, and 555 nm excitation with emission bandwidths of 409-491nm, 504-550nm, and 560-

23

721nm. All images were taken with a format of 1024 x 1024 pixels with pixel size ranging from 69-73nm and a line average of 2. Interslice distance for z-stack imaging was set to 160nm and 250nm with pinhole settings of 0.7 airy units and 1 airy unit of images acquired on the Leica and Zeiss microscopes respectively. Image deconvolution was performed assuming an idealized point spread function using Hyugens software program (Scientific Volume Imaging, Netherlands). The software program Imaris (Oxford Instruments, Abingdon UK) was used for image visualization.

S. pombe cells were grown overnight at 30°C in minimal medium PMG lacking leucine until logarithmic phase and then mounted in PMG 2% agarose as described (*73*). Images were acquired on a Delta Vision Elite microscope (Applied Precision) with a 100X 1.35 NA oil lens (Olympus). Twenty 0.35-μm z sections were acquired generating maximum intensity projections. Further image processing including background-subtracted centromere foci intensity measurements, was performed using ImageJ (National Institutes of Health).

ChIP-Seq
Five independent culture experiments with three time points (day 0, 7 and 14) were performed for each species. Purified cells (~5 x 10^6 cells) and tissue preparations were fixed by 37% formaldehyde treatment. Chromatin preparation, immunoprecipitation, and Illumina sequencing libraires were prepared using the Low Cell ChIP-Seq kit (Active Motif, Carlsbad, CA) according to the manufacturer instructions. Pre-immune sera served as negative controls (Input). ChIP-Seq libraries were sequenced commercially (Psomagen Rockville, Maryland, USA) using Illumina NovaSeq 6000 (150-bp paired end reads).

RNA-Seq
Total RNA was extracted using RNeasy Mini kit (Qiagen). RNA quality and integrity were estimated using Bioanalyzer RNA 6000 Pico Assay (Agilent) and sequenced commercially using Illumina HiSeq 4000 (150-base paired end libraries, Novogene Inc, USA). Reads were mapped using STAR v.2.7.9a (*74*), and normalized to bins per million mapped reads (BPM). Transcripts counts were quantified using kallisto 0.44.0 (*75*).

ChIP-Seq peak calling and sequence analysis
Raw reads were quality-checked with multi-QC (*76*). We removed host DNA contamination from raw sequence reads by mapping against the rat and mouse genomes using Bowtie2 (*77*) with default parameters. Unmapped reads (0.3-5% of total reads) were aligned to the *Pneumocystis* genomes (*P. murina* (GenBank accession number GCA_000349005.2) and *P. carinii* (GCA_001477545.1)), using the Burrows-Wheeler Aligner BWA-MEM (*78*). Alignments were filtered and sorted using SAMTools (*79*) and duplicates marked using PICARD MarkDuplicates http://broadinstitute.github.io/picard). Quality controls were performed using deepTools plotFingerprint (*80*) and phantompeakqualtools (*81*), which estimate the normalized strand coefficient (NSC) and the relative strand correlation (RSC)**.** DeepTools fingerprint plots showed that most reads are mapped within a small fraction of the genome (~3% of the total 7-8 Mb), which indicates that the antibodies demonstrated a restricted binding (**Supplementary figure 13C**). ChIP-Seq coverage data were calculated as 50-bp bin and normalized per genomic content (1x normalization). Chromosome ideograms were generated using chromoMap v0.3 (*82*). Enriched regions following immunoprecipitation (IP) were inspected using IGV genome browser.
Because CENP-A is expected to produce sharp peaks CENP-A ChIP-Seq peak calling was performed using MACS3 ChIP enrichment peaks were detected using Model-Based Analysis for ChIP-Sequencing (*83*) (MACS3 v3.0.0a5) callpeak function with estimated genome sizes and sorted by fold-enrichment (cut off 1.2) and FDR values (broad cutoff 0.1; FDR < 0.05) (raw data are provided in **Supplementary dataset 4**). Enriched peaks were further inspected using deepTools bamCompare versus shuffle non centromeric genomic regions. Peaks were further

examined in conjunction with other data (e.g. conservation score, RNA-seq, methylation data, GC content, repeats) with the Integrative Genomics Viewer (*84*), ensuring that peaks are present in all IP samples and absent in controls. Because H3K9me2/3 and H3K4me2 are histone modifications with broad distribution, we used SICER (Spatial-clustering method for Identification of ChIP-Enriched Regions) (*85*) with default parameters for peak calling (raw data are provided in **Supplementary dataset 4**).

De novo DNA motif searches were performed using MEME v5.1.0 (*86*) and Homer v4.9 (*87*) with a p-value cut off of 10^{-10}. GC content across windows (bin) was computed using BEDTools (*88*). Normalized coverage and annotations across chromosomes were visualized using pyGenomeTracks (*89*).

Repeated elements (DNA transposons, retrotransposons and low complexity repeats) were identified using RepeatMasker (*90*) and RepBase (*91*).
Gene located in centromeres were identified by Genes located were extracted from NCBI annotated GFF3 and converted in BED format using custom scripts. Overlaps between gene regions and centromere regions in BED format were identified uysing BEDtools intersect.

Genome synteny and conservation

DNA alignments were generated using Satsuma2 (*92*) and visualized with Circos (*93*). To estimate the sequence conservation scores, we performed one-to-one pairwise whole genome alignments using LAST (*94*) with the MAM4 seeding scheme (*95*). We consider the trio *P. carinii*/*P. murina*/*P. wakefieldiae*. We generated a neutral model using phyloFit (*96*) from PHAST which fit the tree model to multiple sequence alignments by maximum likelihood using the REV substitution model. Sequence conservation scores were estimated using phastCons (*97*) (--target-coverage 0.25 --expected-length 20 --estimate-trees). Wig files and Bedgraph files were converted using 'wigToBigWig' and 'bigWigToBedGraph' tools (*98*).

Bisulfite sequencing

Pneumocystis DNA (5 μg) was extracted by a *Pneumocystis* DNA enrichment protocol (*15*). Bisulfite conversion, library preparation and sequencing (Illumina NovaSeq 150 bp paired end reads) were performed commercially (Novogene). Bisulfite conversion of non-methylated DNA was performed using EZ DNA Methylation kit. Data analysis was performed using BSMAP and methratio script (*99*). Only cytosine positions with more than 5-fold coverage were considered. We used weighted methylation percentage (*44*), which is calculated as the number of reads supporting methylation over the number of cytosines sequenced to quantify methylation levels. Methylation data were partitioned over different genomic compartments using BEDTools (*100*). Statistical comparisons were performed using R (https://www.R-project.org) to compute non-parametric Mann-Whitney U test and Bonferroni correction to adjust *P*-values for multiple comparisons.

ChIP-qPCR

CENP-A enrichment and H3 depletion relative to H4 within the centromeres were evaluated using primers specific to centromeric and non-centromeric loci (control region taken 200 kilobases away from centromere 1 of each species) (see **Supplementary Table 5** for primers). Primers were designed using primer3 (*101*) from regions visualized using IGV genome browser. Primers with potential matches against mammal genomes were excluded using NCBI BLASTn. Dilutions of 1:10 were used based on preliminary test runs. Each reaction contains 5 μl of iTaq universal Sybr green supermix (Biorad), 1.25 μl of primers (500 nM)) and 2.5 μl of DNA. All assays were run on CFX96 thermocycler (Biorad). Three technical replicates were taken for each assay, and the standard errors of the mean were calculated. The PCR program was as follow: initial denaturation for 2 min at 95°C, followed by 30 cycles of 30 seconds at 95°C, 60°C for 30s, and

72° for 30s. Locus ΔCt values were normalized using the formula: $(^{\Delta\Delta Ct}$ ChIP – $^{\Delta Ct}$ Input; https://www.sigmaaldrich.com/US/en/technical-documents/protocol/genomics/qpcr/chip-qpcr-data-analysis) and the fold enrichment of the ChIP DNA over the input was computed as log2 $(2^{\Delta\Delta Ct})$. The plots and statistical analyses were performed with GraphPad Prism 8.

ChIP-LC-MS
Formalin-fixed purified chromatin preparations were separately co-immunoprecipitated with anti CENP-A, H4 and pre immune serum (control) using Pierce™ Direct Magnetic IP/Co-IP Kit (ThermoFisher). Each IP reaction was performed separately on aliquots from the same chromatin preparation. Elution was performed using 1% formic acid. Extracts were frozen in dry ice, lyophilized for 1 hour and digested using trypsin, diluted, and injected to a Thermo Orbitrap Fusion LC-MS/MS to identify unique peptides. LC-MS/MS data were searched against the proteomes using Proteome Discoverer 2.4 (Thermo Fisher Scientific, Waltham, MA). Label-free quantification was analyzed based on the peak intensity of the precursor ion.

Acknowledgments
We thank Rene Costello for animal studies, Drs. Keytam Awad, Salina Gairhe and Shuibang Wang for discussion about ChIP-Seq. This work has been funded in whole or in part with federal funds from the Intramural Research Program of the US National Institutes of Health (NIH) Clinical Center, the National Institute of Allergy, and Infectious Diseases (NIAID) and the National Cancer Institute. The views expressed in this work are those of the authors and do not necessarily represent the official positions of the NIH or U.S. Government. This study used the Office of Cyber Infrastructure and Computational Biology (OCICB) High Performance Computing (HPC) cluster at the National Institute of Allergy and Infectious Diseases (NIAID), Bethesda, MD. This study also utilized the high-performance computational capabilities of the Biowulf Linux cluster at the NIH, Bethesda, MD (http://biowulf.nih.gov).

Author contributions
Conceptualization: OHC, JAK
Methodology: OHC, HDF, JAK, SC
Investigation: OHC, SC, HDF, YL, LB, HW, ERF. ASD, ST, JPD, MC, LM
Visualization: OHC, JAK, SC, LB, CB, HDF
Supervision: OHC, JAK, SG
Writing—original draft: OHC
Writing—review & editing: OHC, JAK, HDF, LM, SC, YL, LB, HW, ERF, ASD, CB, ST, JPD, SG, MC

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: ChIP-Seq datasets and associated peak calling outputs, RNA-seq and bisulfite sequencing data generated in this study are deposited at NCBI GEO (accession numbers GSE232453, GSE226536, GSE232357, GSE232355, GSE230598). Codes generated for this project are available at GitHub: https://github.com/ocisse/pneumo-cens

Figures and Tables

Table 1. Coordinates, length, and GC content (in %) of *Pneumocystis* centromeres identified by direct ChIP.

Species	NCBI accession no.	Chromosome Length (kb)	Chromosome GC (%) *	Centromere ID	Centromere length (kb)	Centromere Start	Centromere End	Centromere GC (%)	Difference GC (%) **	Synteny
P. carinii B80	NW_017264713.1	635.4	22.9	PcCEN1	6.5	233,731	240,226	21.4	-1.5	PmCEN5
	NW_017264714.1	590.5	23.1	PcCEN2	6.1	121,692	127,752	24.4	1.3	PmCEN2
	NW_017264715.1	543.2	26.2	PcCEN3	7.2	84,071	91,273	24.2	-2	PmCEN4
	NW_017264716.1	564.5	21.4	PcCEN4	7.3	506,597	513,868	19	-2.4	PmCEN6
	NW_017264717.1	566.0	24.5	PcCEN5	4.3	78,025	82,304	22.5	-2	PmCEN1
	NW_017264718.1	476.3	23.5	PcCEN6	5.9	430,075	436,005	21	-2.5	PmCEN13
	NW_017264719.1	465.1	25.7	PcCEN7	6.4	176,750	183,172	23.3	-2.4	PmCEN12
	NW_017264720.1	416.9	23.7	PcCEN8	5.6	50,911	56,559	21.3	-2.4	PmCEN8
	NW_017264721.1	419.0	25.9	PcCEN9	7.1	86,466	93,542	24.9	-1	PmCEN15
	NW_017264722.1	437.4	21.8	PcCEN10	6.3	277,837	284,141	19.8	-2	PmCEN3
	NW_017264723.1	377.5	25.8	PcCEN11	6.1	241,326	247,471	22.7	-3.1	PmCEN9
	NW_017264724.1	370.5	24.3	PcCEN12	5.3	148,081	153,334	22.7	-1.6	PmCEN11
	NW_017264725.1	359.5	23.8	PcCEN13	7.1	240,705	247,764	23.4	-0.4	PmCEN10
	NW_017264726.1	281.9	25.1	PcCEN14	5.3	206,934	212,243	22.5	-2.6	PmCEN17
	NW_017264727.1	446.5	23.7	PcCEN15	4.6	110,888	115,452	22.5	-1.2	PmCEN7
	NW_017264728.1	274.9	23.6	PcCEN16	6.5	45,069	51,578	20.9	-2.7	PmCEN16
	NW_017264729.1	268.2	24.5	PcCEN17	7.8	61,164	68,971	20.1	-4.4	PmCEN14
P. murina	NW_006920852	535.3	23.9	PmCEN1	5.7	458,129	463,864	22.6	-1.34	
	NW_006920853	584.1	23.2	PmCEN2	6.3	463,956	470,213	25.3	2.13	
	NW_006920854	584.1	23.7	PmCEN3	8.0	126,184	134,213	20.8	-2.91	
	NW_006920855	545.3	26.1	PmCEN4	6.7	450,901	457,584	24.5	-1.65	
	NW_006920856	532.6	23.4	PmCEN5	7.4	197,805	205,216	23.0	-0.36	
	NW_006920857	536.0	21.3	PmCEN6	7.5	18,391	25,875	20.5	-0.82	
	NW_006920858	446.5	23.8	PmCEN7	4.8	331,196	335,979	22.7	-1.1	
	NW_006920859	432.0	23.4	PmCEN8	6.6	359,990	366,546	22.2	-1.24	
	NW_006920860	378.7	25.4	PmCEN9	8.1	236,499	244,613	24.0	-1.37	
	NW_006920861	376.0	23.2	PmCEN10	5.1	244,652	249,709	23.6	0.38	
	NW_006920862	385.6	23.5	PmCEN11	6.3	221,796	228,057	24.6	1.06	
	NW_006920863	491.3	24.9	PmCEN12	6.3	310,161	316,424	22.7	-2.18	
	NW_006920864	333.4	23.7	PmCEN13	8.3	39,073	47,361	23.5	-0.16	
	NW_006920865	318.1	22.9	PmCEN14	6.8	38,876	45,692	21.6	-1.35	

NW_006920866	412.2	26.5	PmCEN15	7.8	311,726	319,558	26.8	0.26
NW_006920867	290.2	23.5	PmCEN16	5.5	49,567	55,084	20.0	-3.46
NW_006920868	292.7	25.6	PmCEN17	7.5	208,752	216,295	24.5	-1.09

Supplementary Materials

Figs. S1 to S10

Figs. S11 to S13 and Tables S1 to S4 are available at DOI:

https://doi.org/10.5281/zenodo.10574230

References

1. J. L. Gordon, K. P. Byrne, K. H. Wolfe, Mechanisms of Chromosome Number Evolution in Yeast. *Plos Genetics* **7**, (2011).
2. S. R. Sankaranarayanan *et al.*, Loss of centromere function drives karyotype evolution in closely related Malassezia species. *Elife* **9**, (2020).
3. A. Selmecki, A. Forche, J. Berman, Aneuploidy and isochromosome formation in drug-resistant Candida albicans. *Science* **313**, 367-370 (2006).
4. K. L. McKinley, I. M. Cheeseman, The molecular basis for centromere identity and function. *Nat Rev Mol Cell Biol* **17**, 16-29 (2016).
5. L. Clarke, J. Carbon, Isolation of a yeast centromere and construction of functional small circular chromosomes. *Nature* **287**, 504-509 (1980).
6. J. L. Gordon, K. P. Byrne, K. H. Wolfe, Mechanisms of chromosome number evolution in yeast. *Plos Genet* **7**, e1002190 (2011).
7. L. E. Kursel, H. S. Malik, Centromeres. *Curr Biol* **26**, R487-R490 (2016).
8. K. Sanyal, M. Baum, J. Carbon, Centromeric DNA sequences in the pathogenic yeast Candida albicans are all different and unique. *Proc Natl Acad Sci U S A* **101**, 11374-11379 (2004).
9. H. D. Folco, A. L. Pidoux, T. Urano, R. C. Allshire, Heterochromatin and RNAi are required to establish CENP-A chromatin at centromeres. *Science* **319**, 94-97 (2008).
10. V. Yadav *et al.*, RNAi is a critical determinant of centromere evolution in closely related fungi. *P Natl Acad Sci USA* **115**, 3108-3113 (2018).
11. X. X. Shen *et al.*, Genome-scale phylogeny and contrasting modes of genome evolution in the fungal phylum Ascomycota. *Sci Adv* **6**, (2020).
12. O. H. Cisse *et al.*, Genomic insights into the host specific adaptation of the Pneumocystis genus. *Commun Biol* **4**, (2021).
13. O. H. Cisse, M. Pagni, P. M. Hauser, De novo assembly of the Pneumocystis jirovecii genome from a single bronchoalveolar lavage fluid specimen from a patient. *mBio* **4**, e00428-00412 (2012).
14. B. E. Slaven *et al.*, Draft assembly and annotation of the Pneumocystis carinii genome. *J Eukaryot Microbiol* **53 Suppl 1**, S89-91 (2006).
15. L. Ma *et al.*, Genome analysis of three Pneumocystis species reveals adaptation mechanisms to life exclusively in mammalian hosts. *Nat Commun* **7**, 10740 (2016).
16. N. Varoquaux *et al.*, Accurate identification of centromere locations in yeast genomes using Hi-C. *Nucleic Acids Res* **43**, 5331-5339 (2015).
17. J. M. Almeida, O. H. Cisse, A. Fonseca, M. Pagni, P. M. Hauser, Comparative genomics suggests primary homothallism of Pneumocystis species. *MBio* **6**, (2015).
18. M. T. Hayashi, T. S. Takahashi, T. Nakagawa, J. Nakayama, H. Masukata, The heterochromatin protein Swi6/HP1 activates replication origins at the pericentromeric region and silent mating-type locus. *Nat Cell Biol* **11**, 357-U284 (2009).

388   19.   O. H. Cisse, M. Pagni, P. M. Hauser, Comparative Genomics Suggests That the Human
389   Pathogenic Fungus Pneumocystis jirovecii Acquired Obligate Biotrophy through Gene
390   Loss. *Genome Biology and Evolution* **6**, 1938-1948 (2014).
391   20.   X. Liu, I. McLeod, S. Anderson, J. R. Yates, 3rd, X. He, Molecular analysis of
392   kinetochore architecture in fission yeast. *EMBO J* **24**, 2919-2930 (2005).
393   21.   Y. Shiroiwa *et al.*, Mis17 Is a Regulatory Module of the Mis6-Mal2-Sim4 Centromere
394   Complex That Is Required for the Recruitment of CenH3/CENP-A in Fission Yeast. *Plos*
395   *One* **6**, (2011).
396   22.   C. Casola, D. Hucks, C. Feschotte, Convergent domestication of pogo-like transposases
397   into centromere-binding proteins in fission yeast and mammals. *Mol Biol Evol* **25**, 29-41
398   (2008).
399   23.   J. D. Leverson, H. K. Huang, S. L. Forsburg, T. Hunter, The Schizosaccharomyces pombe
900   aurora-related kinase Ark1 interacts with the inner centromere protein Pic1 and mediates
901   chromosome segregation and cytokinesis. *Mol Biol Cell* **13**, 1132-1143 (2002).
902   24.   T. Hayashi *et al.*, Mis16 and Mis18 are required for CENP-A loading and histone
903   deacetylation at centromeres. *Cell* **118**, 715-729 (2004).
904   25.   J. S. Williams, T. Hayashi, M. Yanagida, P. Russell, Fission yeast Scm3 mediates stable
905   assembly of Cnp1/CENP-A into centromeric chromatin. *Mol Cell* **33**, 287-298 (2009).
906   26.   K. D. Corbett *et al.*, The Monopolin Complex Crosslinks Kinetochore Components to
907   Regulate Chromosome-Microtubule Attachments. *Cell* **142**, 556-567 (2010).
908   27.   L. Ponger, W. H. Li, Evolutionary diversification of DNA methyltransferases in
909   eukaryotic genomes. *Mol Biol Evol* **22**, 1119-1128 (2005).
910   28.   A. J. Bewick *et al.*, Diversity of cytosine methylation across the fungal tree of life. *Nat*
911   *Ecol Evol* **3**, 479-490 (2019).
912   29.   M. Freitag, Histone Methylation by SET Domain Proteins in Fungi. *Annu Rev Microbiol*
913   **71**, 413-439 (2017).
914   30.   J. A. Kovacs *et al.*, Monoclonal-Antibodies to Pneumocystis-Carinii - Identification of
915   Specific Antigens and Characterization of Antigenic Differences between Rat and Human
916   Isolates. *Journal of Infectious Diseases* **159**, 60-70 (1989).
917   31.   D. Lando *et al.*, Quantitative single-molecule microscopy reveals that CENP-A(Cnp1)
918   deposition occurs during G2 in fission yeast. *Open Biol* **2**, (2012).
919   32.   A. G. Castillo *et al.*, Plasticity of fission yeast CENP-A chromatin driven by relative
920   levels of histone H3 and H4. *PLoS Genet* **3**, e121 (2007).
921   33.   H. D. Folco *et al.*, The CENP-A N-tail confers epigenetic stability to centromeres via the
922   CENP-T branch of the CCAN in fission yeast. *Curr Biol* **25**, 348-356 (2015).
923   34.   H. Funabiki, I. Hagan, S. Uzawa, M. Yanagida, Cell cycle-dependent specific positioning
924   and clustering of centromeres and telomeres in fission yeast. *J Cell Biol* **121**, 961-976
925   (1993).
926   35.   T. Mizuguchi *et al.*, Cohesin-dependent globules and heterochromatin shape 3D genome
927   architecture in S. pombe. *Nature* **516**, 432-435 (2014).
928   36.   V. H. Vestereng *et al.*, Quantitative real-time polymerase chain-reaction assay allows
929   characterization of Pneumocystis infection in immunocompetent mice. *Journal of*
930   *Infectious Diseases* **189**, 1540-1544 (2004).
931   37.   H. P. Cam *et al.*, Comprehensive analysis of heterochromatin- and RNAi-mediated
932   epigenetic control of the fission yeast genome. *Nat Genet* **37**, 809-819 (2005).
933   38.   J. Thakur, P. B. Talbert, S. Henikoff, Inner Kinetochore Protein Interactions with Regional
934   Centromeres of Fission Yeast. *Genetics* **201**, 543-561 (2015).
935   39.   K. C. Scott, K. S. Bloom, Lessons learned from counting molecules: how to lure CENP-A
936   into the kinetochore. *Open Biol* **4**, (2014).

40.   J. C. Yasuhara, B. T. Wakimoto, Molecular landscape of modified histones in Drosophila
heterochromatic genes and euchromatin-heterochromatin transition zones. *PLoS Genet* **4**,
e16 (2008).
41.   A. Miura *et al.*, Mobilization of transposons by a mutation abolishing full DNA
methylation in Arabidopsis. *Nature* **411**, 212-214 (2001).
42.   K. M. Smith, J. M. Galazka, P. A. Phatale, L. R. Connolly, M. Freitag, Centromeres of
filamentous fungi. *Chromosome Res* **20**, 635-656 (2012).
43.   A. J. Bewick *et al.*, Diversity of cytosine methylation across the fungal tree of life. *Nat*
Ecol Evol **3**, 479-+ (2019).
44.   D. E. Cook, H. M. Kramer, D. E. Torres, M. F. Seidl, B. Thomma, A unique chromatin
profile defines adaptive genomic regions in a fungal plant pathogen. *Elife* **9**, (2020).
45.   L. Ma, O. H. Cissé, J. A. Kovacs, A Molecular Window into the Biology and
Epidemiology of
spp. *Clin Microbiol Rev* **31**, (2018).
46.   K. Schotanus *et al.*, Histone modifications rather than the novel regional centromeres of
Zymoseptoria tritici distinguish core and accessory chromosomes. *Epigenetics Chromatin*
8, 41 (2015).
47.   M. F. Seidl *et al.*, Repetitive Elements Contribute to the Diversity and Evolution of
Centromeres in the Fungal Genus Verticillium. *mBio* **11**, (2020).
48.   A. L. Pidoux, R. C. Allshire, Kinetochore and heterochromatin domains of the fission
yeast centromere. *Chromosome Res* **12**, 521-534 (2004).
49.   V. Yadav *et al.*, RNAi is a critical determinant of centromere evolution in closely related
fungi. *Proc Natl Acad Sci U S A* **115**, 3108-3113 (2018).
50.   L. Teytelman, D. M. Thurtle, J. Rine, A. van Oudenaarden, Highly expressed loci are
vulnerable to misleading ChIP localization of multiple unrelated proteins. *Proc Natl Acad*
Sci U S A **110**, 18602-18607 (2013).
51.   K. Nagaki *et al.*, Sequencing of a rice centromere uncovers active genes. *Nat Genet* **36**,
138-145 (2004).
52.   J. A. Kovacs *et al.*, Identification of antigens and antibodies specific for Pneumocystis
carinii. *J Immunol* **140**, 2023-2031 (1988).
53.   Y. Liu *et al.*, MUC1 mediates Pneumocystis murina binding to airway epithelial cells. *Cell*
Microbiol **22**, e13182 (2020).
54.   S. A. Sabatinos, S. L. Forsburg, Molecular genetics of Schizosaccharomyces pombe.
Methods Enzymol **470**, 759-795 (2010).
55.   Y. Takayama *et al.*, Biphasic incorporation of centromeric histone CENP-A in fission
yeast. *Mol Biol Cell* **19**, 682-690 (2008).
56.   T. Sakuno, Y. Watanabe, [Kinetochore geometry defined by cohesin within the
centromere]. *Tanpakushitsu Kakusan Koso* **54**, 1842-1849 (2009).
57.   H. D. Folco, A. McCue, V. Balachandran, S. I. S. Grewal, Cohesin Impedes
Heterochromatin Assembly in Fission Yeast Cells Lacking Pds5. *Genetics* **213**, 127-141
(2019).
58.   S. F. Altschul *et al.*, Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Res* **25**, 3389-3402 (1997).
59.   D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**, 238 (2019).
60.   A. Hernandez-Plaza *et al.*, eggNOG 6.0: enabling comparative genomics across 12 535
organisms. *Nucleic Acids Res* **51**, D389-D394 (2023).
61.   P. Jones *et al.*, InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**, 1236-1240 (2014).

62.   K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066 (2002).

63.   M. N. Price, P. S. Dehal, A. P. Arkin, FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* **26**, 1641-1650 (2009).

64.   L. T. Nguyen, H. A. Schmidt, A. von Haeseler, B. Q. Minh, IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**, 268-274 (2015).

65.   J. Postberg, S. Forcob, W. J. Chang, H. J. Lipps, The evolutionary history of histone H3 suggests a deep eukaryotic root of chromatin modifying mechanisms. *BMC Evol Biol* **10**, 259 (2010).

66.   S. R. Eddy, Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**, e1002195 (2011).

67.   E. M. Gertz, Y. K. Yu, R. Agarwala, A. A. Schaffer, S. F. Altschul, Composition-based statistics and translated nucleotide searches: improving the TBLASTN module of BLAST. *BMC Biol* **4**, 41 (2006).

68.   G. S. Slater, E. Birney, Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics* **6**, 31 (2005).

69.   S. El-Gebali *et al.*, The Pfam protein families database in 2019. *Nucleic Acids Res* **47**, D427-D432 (2019).

70.   R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792-1797 (2004).

71.   J. A. Kovacs *et al.*, Monoclonal antibodies to Pneumocystis carinii: identification of specific antigens and characterization of antigenic differences between rat and human isolates. *J Infect Dis* **159**, 60-70 (1989).

72.   M. J. Linke, S. M. Sunkin, R. P. Andrews, J. R. Stringer, P. D. Walzer, Expression, structure, and location of epitopes of the major surface glycoprotein of Pneumocystis carinii f. sp. carinii. *Clin Diagn Lab Immunol* **5**, 50-57 (1998).

73.   P. T. Tran, A. Paoletti, F. Chang, Imaging green fluorescent protein fusions in living fission yeast cells. *Methods* **33**, 220-225 (2004).

74.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

75.   N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525-527 (2016).

76.   P. Ewels, M. Magnusson, S. Lundin, M. Kaller, MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048 (2016).

77.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012).

78.   H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*, 1303.3997 (2013).

79.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

80.   F. Ramirez *et al.*, deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

81.   S. G. Landt *et al.*, ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome Res* **22**, 1813-1831 (2012).

82.   C. M. R. L. Lakshay Anand, chromoMap: An R package for Interactive Visualization and Annotation of Chromosomes. *bioRxiv*, (2020).

83.   Y. Zhang *et al.*, Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

84.   J. T. Robinson *et al.*, Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26 (2011).

23

85. C. Zang *et al.*, A clustering approach for identification of enriched domains from histone modification ChIP-Seq data. *Bioinformatics* **25**, 1952-1958 (2009).

86. T. L. Bailey *et al.*, MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**, W202-208 (2009).

87. S. Heinz *et al.*, Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol Cell* **38**, 576-589 (2010).

88. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010).

89. L. Lopez-Delisle *et al.*, pyGenomeTracks: reproducible plots for multivariate genomic datasets. *Bioinformatics* **37**, 422-423 (2021).

90. H. R. Smit AFA., Green P. , RepeatMasker Open-4.0., (2013-2015).

91. W. Bao, K. K. Kojima, O. Kohany, Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob DNA* **6**, 11 (2015).

92. M. G. Grabherr *et al.*, Genome-wide synteny through highly sensitive sequence alignment: Satsuma. *Bioinformatics* **26**, 1145-1151 (2010).

93. M. Krzywinski *et al.*, Circos: an information aesthetic for comparative genomics. *Genome Res* **19**, 1639-1645 (2009).

94. S. M. Kielbasa, R. Wan, K. Sato, P. Horton, M. C. Frith, Adaptive seeds tame genomic sequence comparison. *Genome Res* **21**, 487-493 (2011).

95. M. C. Frith, L. Noe, Improved search heuristics find 20,000 new alignments between human and mouse genomes. *Nucleic Acids Res* **42**, e59 (2014).

96. A. Siepel, D. Haussler, Phylogenetic estimation of context-dependent substitution rates by maximum likelihood. *Mol Biol Evol* **21**, 468-488 (2004).

97. A. Siepel *et al.*, Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res* **15**, 1034-1050 (2005).

98. W. J. Kent, A. S. Zweig, G. Barber, A. S. Hinrichs, D. Karolchik, BigWig and BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**, 2204-2207 (2010).

99. Y. Xi, W. Li, BSMAP: whole genome bisulfite sequence MAPping program. *BMC Bioinformatics* **10**, 232 (2009).

100. A. R. Quinlan, BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Curr Protoc Bioinformatics* **47**, 11 12 11-34 (2014).

101. T. Koressaar *et al.*, Primer3_masker: integrating masking of template sequence with primer design software. *Bioinformatics* **34**, 1937-1938 (2018).

A Lungs (day 0)

1 DAPI | *Pneumocystis* | CENP-A

1 2 3 4 5 6 7 8

B Culture (day 7)

H

A

~35 Mya

~460 Mya

P. murina

P. carinii

S. pombe

Co culture on A549 and
Let-1 cells for 14 days

B

P. murina

DNA copies (log DHFR)

10^6

10^5

10^4

0   7   14

Days Post inoculation

C

P. carinii

DNA copies (log DHFR)

10^6

10^5

10^4

0   7   14

Days Post inoculation

D

P. murina (day 7)

A *P. murina*

NW_006920853

Genes

Repetitive DNA

Gypsy-40_MLP-I

0.8 / 0 — Percent GC content

342.5 / 0 — CENP-A (ChIP-seq)

20 / -20 — H3/H4 ratio (ChIP-seq)

100 / 0 — H3K9me2 (ChIP-seq)

100 / 0 — H3K9me3 (ChIP-seq)

100 / 0 — H3K4me2 (ChIP-seq)

29.1 / 0 — Gene expression (RNA-seq)

6.2 kb

B *P. carinii*

NW_017264714.1

Genes

Repetitive DNA

Copia-58_BG-I
Copia-58_BG-I

0.7 / -0.0 — Percent GC content

1115.6 / 0 — CENP-A (ChIP-seq)

10 / -10 — H3/H4 ratio (ChIP-seq)

40 / 0 — H3K9me2 (ChIP-seq)

40 / 0 — H3K9me3 (ChIP-seq)

40 / 0 — H3K4me2 (ChIP-seq)

400 / 0 — Gene expression (RNA-seq)

6.1 kb

A

~35 Mya

P. carinii
━━ Centromeres

P. murina
━━ Centromeres

B

CEN4

420   440   460   480   500   520   540   560 Kb

Genes

352.6

CENP-A

PhastCons

P. carinii
Chromosome 4

C

****

****

****

PhastCons score

CEN   CENflk   Msg   Bckg