The Host Adapted Fungal Pathogens of *Pneumocystis* Genus Utilize Genic

Regional Centromeres

Authors:

Ousmane H. Cissé [1], Shelly Curran [1], H. Diego Folco [2], Yueqin Liu [1], Lisa Bishop [1], Honghui Wang [1],

Elizabeth R. Fischer [3], A Sally Davis [4], Spenser Babb-Biernacki [5], Vinson P. Doyle [6], Jonathan K.

Richards [6], Sergio A. Hassan [7], John P. Dekker [8,9], Pavel P. Khil [8,9], Jason M. Brenchley [10], Shiv Grewal

[2], Melanie Cushion [11], Liang Ma [1] and Joseph A. Kovacs [1]

Affiliations:
[1] Critical Care Medicine Department, Clinical Center, National Institutes of Health, Bethesda,
Maryland, USA.
[2] Laboratory of Biochemistry and Molecular Biology, National Cancer Institute, National Institutes of
Health, Bethesda, Maryland, USA.
[3] Microscopy Unit, Research Technologies Branch, Rocky Mountain Laboratories, National Institute
of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, Montana, USA.
[4] Diagnostic Medicine/Pathobiology, Kansas State University College of Veterinary Medicine,
Manhattan, USA.
[5] Museum of Natural Science and Department of Biological Sciences, Louisiana State University,
Baton Rouge, Louisiana, USA.
[6] Department of Plant Pathology and Crop Physiology, Lousiana State University AgCenter, Baton
Rouge, Louisiana, USA.
[7] Bioinformatics and Computational Biosciences Branch, National Institute of Allergy and Infectious
Diseases, National Institutes of Health, Bethesda, Maryland, USA.
[8] Bacterial Pathogenesis and Antimicrobial Resistance Unit, National Institute of Allergy, and
Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA.
[9] Department of Laboratory Medicine, Clinical Center, National Institutes of Health, Bethesda,
Maryland, USA.
[10] Laboratory of Viral Diseases, National Institute of Allergy, and Infectious Diseases, National
Institutes of Health, Bethesda, Maryland, USA.
[11] Department of Internal Medicine, College of Medicine, University of Cincinnati, Cincinnati, Ohio,
USA.

32
33   Correspondence to: ousmane.cisse@nih.gov and/or jkovacs@mail.nih.gov
34
35

Abstract

Centromeres are genomic regions that coordinate accurate chromosomal segregation during mitosis

and meiosis. Yet, despite their essential function, centromeres evolve rapidly across eukaryotes.

Centromeres are often the sites of chromosomal breaks which contribute to genome shuffling and

promote speciation by inhibiting gene flow. How centromeres form in strongly host-adapted fungal

pathogens has yet to be investigated. Here, we characterized the centromere structures in closely

related species of mammalian-specific pathogens of the fungal phylum of Ascomycota. Methods

allowing reliable continuous culture of *Pneumocystis* species do not currently exist, precluding genetic

manipulation. CENP-A, a variant of histone H3, is the epigenetic marker that defines centromeres in

most eukaryotes. Using heterologous complementation, we show that the *Pneumocystis* CENP-A

ortholog is functionally equivalent to CENP-A^{Cnp1} of *Schizosaccharomyces pombe*. Using organisms

from a short-term *in vitro* culture or infected animal models and ChIP-seq, we identified centromeres

in three *Pneumocystis* species that diverged ~100 million years ago. Each species has a unique short

regional centromere (< 10kb) flanked by heterochromatin in 16-17 monocentric chromosomes. They

span active genes and lack conserved DNA sequence motifs and repeats. CENP-C, a scaffold protein

that links the inner centromere to the kinetochore appears dispensable in one species, suggesting a

kinetochore rewiring. Despite the loss of DNA methyltransferases, 5-methylcytosine DNA

methylation occurs in these species, though not related to centromere function. These features

suggest an epigenetic specification of centromere function.

Short summary

Pneumocystis species offer a suitable genetic system to study centromere evolution in pathogens

during host adaptation because of their unique specificity for mammals, and their phylogenetic

proximity with the nonpathogenic yeast *Schizosaccharomyces pombe*, a popular model for cell biology.

We used this system to explore how centromeres have evolved after divergence of the two clades ~460

million years ago. To address this question, we established a protocol combining short-term culture

and ChIP-seq to characterize centromeres in multiple *Pneumocystis* species. We show that

Pneumocystis have short epigenetic centromeres that function differently from those in *S. pombe* and

exhibit similarities to centromeres in more distantly related host adapted fungal pathogens.

Introduction

Centromeres are genomic locations where spindles attach during chromosomal segregation. They are

essential for cell division. Errors in the chromosomal segregation can result in aneuploidy and cell

division defects with disastrous consequences such as cancer. In fungi, centromeres are fragile sites

that are often involved in karyotype variability (Sankaranarayanan, Ianiri et al. 2020, Seidl, Kramer et

al. 2020) and drug resistance (Selmecki, Forche et al. 2006). In most eukaryotes, centromeres are

defined by the presence of CENP-A nucleosomes, a centromere specific histone H3 variant that is

essential for the localization of known kinetochore components (McKinley and Cheeseman 2016).

There is a remarkable diversity in centromere structures ranging from sequence-dependent (point)

centromeres to epigenetically-regulated (regional) centromeres.

Point centromeres are exemplified by *Saccharomyces cerevisiae* with genetically defined 125 bp

long centromeres containing conserved DNA elements (I, II and III) that serve as anchors for the

recruitment of the centromeric DNA binding factor 3 (CBF3) complex (Clarke and Carbon 1980,

Gordon, Byrne et al. 2011). Regional centromeres are much longer, lack universally conserved DNA

patterns, and are often made up of repetitive DNA (Kursel and Malik 2016). Regional centromeres

are epigenetically regulated; however, which factors contribute to this epigenetic specification of

centromeres remains elusive.

Pneumocystis is a genus of pathogenic fungi that exclusively infect mammals and remain

unculturable. They belong to the Taphrinomycotina subphylum and form a monophyletic clade with

fission yeasts *Schizosaccharomyces*, with a separation time estimated at ~460 million years ago (Shen,

Steenwyk et al. 2020, Cisse, Ma et al. 2021). In addition to these two classes there are other classes, all

of which are plant or soil adapted organisms. Compared to *S. pombe*, *Pneumocystis* species have

streamlined genomes due to substantial gene losses during their transition to animal parasitism.

Recently, the genomes of multiple *Pneumocystis* species have been sequenced (Slaven, Meller et al.

2006, Cisse, Pagni et al. 2012, Ma, Chen et al. 2016, Cisse, Ma et al. 2021). However, centromeres

were not defined in these because they cannot be predicted bioinformatically. *Pneumocystis* have

retained CENP-A, which presumably binds to centromeres, and most of the kinetochore proteins.

Some data suggest that *Pneumocystis* centromeres differ structurally from those of *S. pombe*. For

example, missing in *Pneumocystis* genomes is the heterochromatin protein Swi6/HP1 (Almeida, Cisse

et al. 2015), which is required for replication origins at the centromeres flanking regions

(pericentromeres) (Hayashi, Takahashi et al. 2009). The RNAi pathway is lost in *Pneumocystis* (Cisse,

Pagni et al. 2014), further supporting mechanistic difference with *S. pombe*. Similar to *S. pombe*, DNA

methylation has been predicted to be absent in *Pneumocystis* based on the absence of DNA

methyltransferases (DMTs) (Freitag 2017).

In this work, we undertook to characterize *Pneumocystis* centromeres and compare them to

Schizosaccharomyces pombe centromeres. We found that *Pneumocystis* species have small regional

centromeres that are structurally different from those of *S. pombe*.

Results

Survey of genes related to centromere function in Taphrinomycotina fungi.

To understand the kinetochore evolution in Taphrinomycotina fungi, we screened the predicted

proteomes of 14 species spanning six orders, *Pneumocystis* ($n = 7$), *Schizosaccharomyces* ($n = 3$), *Taphrina*

($n = 1$), *Saitoella* ($n = 1$), *Protomyces* ($n = 1$) and *Neolecta* ($n = 1$), as well as other fungi to have a broad

overview of kinetochore evolution.

We searched for homologs of CENP-A [Cnp1], CENP-C[Cnp3], CCAN (Constitutive Centromere

Associated Network) and KMN (KNL-1/Mis12 complex/Ndc80 complex) proteins. We extended our

searches to pathways required for centromere function and chromosomal segregation

(heterochromatin, RNAi, and DNA methylation).

CENP-A, CENP-C, and outer kinetochore proteins are conserved across Taphrinomycotina

(**Supplementary figure 1**). Two genes of the inner kinetochore (CCAN) that are absent in

Pneumocystis are clearly innovations in *S. pombe* (*fta2* and *fta3* genes). The *fta2*, *fta3* and *fta4* gene

products associate with the central core (cnt) and innermost repeats (imr) region of the centromere

in *S. pombe* (Liu, McLeod et al. 2005). *Pneumocystis* also lack the *mis17* gene exclusively found in *S.*

pombe, which encodes a member of the Mis6-Mal2-Sim4 multiprotein required for CENP-A

recruitment in *S. pombe* (Shiroiwa, Hayashi et al. 2011). The *S. pombe* centromere-associated protein

B genes (*cbp1*, *cbh1* and *cbh2*), which are derived from the domestication of *pogo*-like transposases

(Casola, Hucks et al. 2008) have no identified homologs in other fungi.

The chromosomal passenger complex (CPC) is a heterotetrametric complex composed of the

Aurora B kinase Ark1 and three regulatory components Pic1 (inner centromere protein), Bir1p

(Survivin) and Nbl1 (Borealin) which mediates chromosome segregation and cytokinesis (Leverson,

Huang et al. 2002). This pathway is conserved in *Pneumocystis* and *S. pombe* (**Supplementary figure**

1).

CENP-A recruiting complex, which includes Mis18 and Mis16 (Hayashi, Fujita et al. 2004) as

well as the CENP-A histone chaperone Scm3 required for CENP-A loading at the centromere

(Williams, Hayashi et al. 2009) are conserved in *Pneumocystis*, as are the monopolin complex genes

that coordinate the kinetochore microtubules attached during mitosis (Corbett, Yip et al. 2010).

The network for heterochromatin formation in *Pneumocystis* seems intact with the presence

of the Clr4 methyltransferase complex (Clr4/SUV39H). The only notable loss is Swi6, which with

Chp2 are HP1 orthologs, while Chp2 is conserved. DNA methylases (DMTs) are not present in any

of the Taphrinomycotina fungi sequenced so far, which suggests that DNA methylation does not

occur in this subphylum. Overall, the *Pneumocystis* kinetochore protein catalog is similar to those

other Taphrinomycotina fungi except *S. pombe* which seems to be an outlier with a few gene

additions. We conclude that *Pneumocystis* species have retained an ancestral version of the

kinetochore.

Pneumocystis **CENP-A**[cnp1] **localizes to the nucleus.**

Centromeres are identified by tracking CENP-A binding regions. There are currently seven

Pneumocystis species with fully sequenced genomes (Ma, Chen et al. 2016, Cisse, Ma et al. 2021). All

of them encode an ortholog of *S. pombe* CENP-A[cnp1,] (**Figure 1a**). The CENP-A histones from these

species, henceforth referring to as *pn*CENP-A, display a high overall conservation (89% average

protein identity) with most divergence in the N-terminal tail.

Precise nuclear localization of CENP-A is required for accurate chromosomal segregation. To

determine if *pn*CENP-A localizes in the nucleus, we generated specific antibodies using synthetic

immunogenetic peptides representing sequences located in the N-terminus of the protein, to

minimize cross reactivity with host histones (**Supplementary figure 2; Supplementary methods**).

We used anti – *Pneumocystis* CENP-A antibodies in conjunction with DAPI to demonstrate that

*pn*CENP-A colocalizes with DAPI which support a nuclear localization within the *Pneumocystis*

nucleus (**Figure 1b**).

*pn***CENP-A variants are functional in *S. pombe*, and support viability and centromere loading.**

To confirm that the putative *cenp-a* genes from *Pneumocystis* encode functional CENP-A

proteins, we expressed the *cenp-a* genes from *P. jirovecii* and *P. murina* in the related fission yeast *S.*

pombe, a tractable organism of the subdivision Taphrinomycotina widely used as model for

chromosome and cell biology. The transgenes were codon-optimized, expressed under the

endogenous *S. pombe cnp1*, the CENP-A ortholog, gene promoter and inserted in single copy at the

lys1 locus. In addition, the gene products were tagged with GFP at their N-termini since it has been

shown that C-terminal tagging (*i.e.*, *cnp1-GFP*) displays growth retardation (Lando, Endesfelder et al.

2012). As *Pneumocystis* CENP-A proteins harbor an intervening sequence at the N-terminal tail that

may potentially interfere with function in *S. pombe*, we also expressed truncated versions missing ~25

aa from the N-terminal tail (**Figure 1c**). We first assessed the ability of *Pneumocystis* CENP-A to

rescue lethality of *cnp1-76* cells, a thermosensitive mutant, whose mutation T74M lies at the CATD

(CENP-A targeting domain) region (Castillo, Mellone et al. 2007). As previously reported (Folco,

Campbell et al. 2015), we found that the control transgene (SpCENP-A) restored growth of *cnp1-76*

cells at restrictive temperatures (i.e., 32°C and 35°C). Importantly, *Pneumocystis* counterparts were

also able to rescue the thermosensitive phenotype, albeit partially (**Figure 1c**), nevertheless proving

the functionality of these genes. Results were similar between constructs harboring either intact or

truncated N-termini (**Figure 1c**). Therefore, we continued our analyses with the full-length proteins.

It is well established that in interphase, *S. pombe* displays the Rabl configuration, in which

centromeres cluster to the spindle pole body while telomeres associate with each other near nuclear

periphery (Funabiki, Hagan et al. 1993, Mizuguchi, Fudenberg et al. 2014). As expected, expressing *S.*

pombe CENP-A tagged with GFP resulted in cells displaying single fluorescence foci that colocalized

with a *tetO* array inserted at *cen2* (NB: a TetR-tdTomato fusion is expressed to bind to the *tetO* array

and this locus is referred to as *cen2-tetO-tdTomato*) (**Figure 1d**). Consistent with the mentioned rescue

of *cnp1-ts* cells and further supporting their role as functional CENP-A proteins, *Pneumocystis* CENP-

A proteins also colocalize with *cen2-tetO-tdTomato* and remarkably, they exhibited single foci,

although the intensities of centromere foci were dimmer than that of the *S. pombe* counterpart

(**Figure 1e**). Finally, to confirm the specific localization of *Pneumocystis* CENP-A proteins to *S. pombe*

centromeres, we performed ChIP-qPCR (**Figure 1f**). Noteworthily, we observed that *Pneumocystis*

CENP-A localized to central core regions (i.e., *cc1&3*, *cnt2:ura4+*) similarly to the *S. pombe*

counterpart, albeit the enrichment was lower, consistent with the results from live-cell imaging.

Additionally, *Pneumocystis* CENP-A localization seems to be specific as no detectable enrichment was

found at pericentromeric outer repeats coated with heterochromatin (i.e., *dg*) and euchromatin

locations (i.e., *fbp1*). Together, these results confirm that *Pneumocystis* CENP-A genes encode bona

fide CENP-A proteins.

pnCENP-A binds to single genomic foci in *Pneumocystis* replicating cells.

In *S. pombe*, CENP-A^{cnp1} is deposited at the centromeres during the G2 phase of the cell cycle (Lando,

Endesfelder et al. 2012). Although *Pneumocystis* cell replication is not fully understood, these

organisms take 5 to 8 days to replicate *in vivo* (Vestereng, Bishop et al. 2004). To determine *pn*CENP-

A bound genome regions, we established a short-term co-culture system for *P. murina* and *P. carinii*

(**Figure 2a**). We used quantitative PCR targeting the single copy gene dihydrofolate reductase (*dhfr*)

to assess organism growth; a typical growth curve shows a decline from day 0 to 7 followed by a

gradual increase (**Figures 2b and 2c**). Population growth is further supported by the presence of

mitotic or fusing cells 7 days post culture, though some cells do not appear to be dividing (**Figure**

2d). We assessed *Pn*CENP-A location on the *P. carinii* genome using ChIP-seq targeting chromatin

prepared from infected lung tissue lysates and cultured organisms harvested at three time points.

CENP-A binds to discrete areas, which is consistent with centromeres (**Figure 2e** and

Supplementary figure 3).

We mapped centromeres in all *P. carinii* and *P. murina* chromosomes by ChIP-seq using anti-

*pn*CENP-A and anti-*pn*CENP-C antibodies (**Supplementary figures 2a** and **b**) with cell cultures

collected from three timepoints: days 0 (normalized initial inoculum), 7 and 14; pre-immune sera

served as negative controls (Input). Five independent experiments were performed for each species.

We removed host DNA contamination from raw sequence reads by mapping against the rat and

mouse genomes; filtered reads (0.3-5% of total reads) were aligned to the *Pneumocystis* genomes. We

the genome-wide ratios of immunoprecipitated DNA sequences and matching controls (input

sequences) to identify significantly enriched genomic loci using a false discovery rate [FDR] < 0.05

and a minimum fold enrichment of 1.2. Enriched regions following immunoprecipitation (IP) were

inspected using IGV genome browser. DeepTools fingerprint plots show that most reads are mapped

within a small fraction of the genome (~3% of the total 7-8 Mb), which indicates that the antibodies

demonstrated a restricted binding (**Supplementary figure 2c**). Using ChIP-qPCR, we confirmed

*pn*CENP-A enrichment over regions flagged by ChIP-seq compared to a control region taken 200

kilobases away from centromere 1 of each species) for 16 of 17 *P. murina* chromosomes and all 17 *P.*

carinii chromosomes (**Supplementary figure 4**).

In *P. murina*, each of the 17 chromosome-level scaffolds display a single region with

overlapping peaks of CENP-A and CENP-C (**Figures 3a and 3b**). The *P. carinii* genome also displays

monocentric chromosomes enriched with CENP-A but they lack any CENP-C enrichment (**Figures**

3c and 3d).

Enriched regions in both species span 4.8 to 8.0 kb, with an average length of 6.7 kb, and

have an average of 3.9% lower GC content than the rest of the genome (**Figures 4a and 4b;**

Supplementary Table 1). *P. murina* and *P. carinii* centromeres encode 74 and 58 genes respectively;

all of them are conserved in the genomes of both species except a prefoldin gene that is lost in *P.*

carinii (**Supplementary Table 2**). Of the 74 *P. murina* centromeric genes, 73 were expressed based on

RNA-seq and four were further detected by protein mass spectrometry mapping (LC-MS); of the 58

P. carinii genes, 56 are expressed and 53 were detected by LC-MS. Analysis of the predicted function

of these genes revealed housekeeping functions without major differences compared to randomly

sampled genes.

To map *P. macacae* centromeres, we performed ChIP-seq on infected lung tissues from two

Simian immunodeficiency virus (SIV)-infected macaques with untreated *Pneumocystis* pneumonia (cf.

Supplementary Methods). We prepared chromatin directly from lung tissues without culture.

Because anti-CENP-A and anti-CENP-C antibodies specific to *P. macacae* (**Supplementary figures 2a**

and 2b) yielded nonspecific peaks in ChIP-seq experiments, we utilized two antibodies targeting *P.*

macacae Mis12 (**Supplementary figure 2d**), a member of the outer kinetochore that has been used as

replacement of CENP-A for centromeres identification in the fungal pathogen *Mucor circinelloides*

(Navarro-Mendoza, Perez-Arques et al. 2019). Mis12 ChIP-seq identified -4-6 peaks per chromosome

(**Figure 4c; Supplementary figure 5**). Mis12 peaks tend to occur in GC rich regions. It is unclear if

the multiple peaks correspond to ChIP-seq artifacts or if Mis12 truly binds to multiple locations per

chromosome. However, there is only one peak flanked by heterochromatin (H3K9me2) per

chromosome, which also flanked *P. carinii* and *P. murina* centromeres (**Figures 4a and 4b,**

Supplementary figures 6 and 7). DNA sequence similarity searches showed that this peak is

homologous to *P. carinii* and *P. murina* centromeres. We consider this peak to be a putative

centromere.

We were unable to map centromeres in *P. jirovecii* despite utilizing anti-CENP-A and anti-

CENP-C antibodies specific for this species (inconclusive ChIP-seq results), possibly because the only

samples available for Chip-seq studies were from patients previously treated with anti-*Pneumocystis*

drugs.

Centromeric nucleosomes assemble differently among species.

CENP-C binds to specific motifs in CENP-A in the C-terminal cupin domain involved in

dimerization (Kato, Jiang et al. 2013). To confirm the absence of CENP-C ChIP-seq peaks at *P. carinii*

centromeres, we analyzed chromatin by co-immunoprecipitation (Co-IP) with anti-CENP-A, anti-

CENP-C and anti-H4 antibodies followed by LC-MS analysis. While CENP-A was detected by

peptide mapping with high confidence (peptide: "<u>WQSTAILCLQEATEAFLVHLFEDTNLCAIHAK</u>",

Qvality q-value = 0.00553986), none of the CENP-C peptides were identified in either of the two

runs. Of note, we detected CENP-C in total proteins from both species by LC-MS and Western

blotting (**Supplementary figures 2e and f**). These results are consistent with ChIP-seq results and

suggest that *P. carinii* CENP-C does not localize at the *P. carinii* centromeres. *pn*CENP-C may be

dispensable in *P. carinii*, similar to *S. pombe* (Sridhar and Fukagawa 2022).

To gain further molecular insight into the CENP-A nucleosome differences, we performed

molecular dynamics (MD) simulations of centromeric complex models of *P. carinii*, *P, murina*, and *S.*

pombe (cf. **Supplementary Methods**). The analysis suggests that the observed differences can be

explained by the different histones H2A:H2B dimers recruited by each species at the centromeres

(**Figure 5a**). Comparative analysis shows that variation in the terminal regions of these proteins can

induce structural and dynamic changes in the region surrounding the central crevice formed by the

spatial confluence of four histones (cf. **Figure 5b and Supplementary animation D2**). This region

of the octameric complex is solvent-exposed and poised for binding to inner-kinetochore proteins

such as CENP-C. Slight variations in its average size or dynamics can affect protein selection and

binding process, hence the early-stage assembly mechanism, even when its amino acid composition is

identical, as is the case of *P. carinii* and *P. murina*. Compared to these species, *S. pombe* presents

several mutations lining the crevice (cf. **Figure 5a**), which adds physicochemical features to consider

when comparing the two organisms. For the *Pneumocystis* sequences modeled here, the structural and

dynamic differences between the two species stem from the longer C-terminal domain of H2A in *P.*

murina, which adopts a helical conformation with several nonpolar moieties. This helix is quite

mobile and tend to interact hydrophobically with the disordered N-terminus of CENP-A

(**Supplementary Table 3**). However, when free from these interactions, the helix can transiently

gravitate towards the core center, affecting the exposure of the crevice to the solvent, hence to other

proteins (cf. **Figure 5b**). Transient helices in intrinsically disordered segments can have functional

implications (Chen, Yang et al. 2021, Zhao, Nguyen et al. 2022), and hydrophobic helixes, in

particular, can play a role in molecular recognition and drive protein association and aggregation

(Zhao, Nguyen et al. 2022).

Pneumocystis **CENP-A replaces the canonical histone H3 at the centromeres.**

In *S. pombe*, CENP-A^{cnp1} enrichment at the centromeres correlates with a depletion of histone H3

(Thakur, Talbert et al. 2015). To determine if this feature is present in *Pneumocystis*, we quantified the

relative abundance of canonical histones H3 and H4 using ChIP-seq. We examined the 16-17

centromeres per species, their flanking regions (30 kb) and control regions selected 25 kb away from

centromeres. H3 occupancy (expressed as ratio between H3 and H4) was found to be significantly

reduced at centromeres relative to controls in *P. carinii* and *P. murina* but not *P. macacae* (Wilcoxon

test; $P < .0001$; **Supplementary figure 8**). In parallel to ChIP-seq, we found by ChIP-qPCR a

reduction of histone H3 in 15 of the 17 *P. carinii* centromeres, and in 2 of the 17 *P. murina*

centromeres compared to the controls (**Supplementary figure 9**). The discrepancy between *P.*

murina ChIP-seq and ChIP-qPCR is potentially due to variability in culture experiments.

Centromeres sequences are unique and lack repetitive sequences.

Centromeres are located in non-repetitive sequences within the *Pneumocystis* genomes

(**Figures 3e and 3f**). Motif searches also did not reveal any conserved motif either in the centromeric

regions as a whole or when focused on *pn*CENP-A peaks. To determine if repeats (DNA transposons

and retrotransposons) are associated to centromeres, we searched for known signatures of repeats. In

P. murina, we detected long terminal repeats (LTRs) (*copia* and *gypsy*) in the vicinity of 5 of the 17

centromeres but they are not directly flanking the CENs (**Supplementary figure 7**). In *P. carinii*, of

the 17 centromeres, the CEN2 contains a duplicated copy of *copia*-retrotransposons and CEN16 has a

gypsy repeat in its upstream region (**Figure 4b; Supplementary figure 6**). The repeated DNA

elements are likely nonfunctional for further transposition because they lack essential domains

required for transposition activity (LTRs, reverse transcriptase, integrase) and they are integrated into

genes. No repeat is present in the vicinity of *P. macacae* centromeres (**Supplementary figure 5**).

Centromeres are flanked by heterochromatin and lack DNA methylation marks.

In *S. pombe*, H3K9me2 marking of chromatin is associated with the repression of centromeres,

subtelomeres, ribosomal rDNA and the mating locus (Cam, Sugiyama et al. 2005). There are two

types of chromatins: euchromatin, the lightly packed form of the chromatin enriched with H3K4me2

(di-methylation of the 4th lysine residue of the histone H3), which is associated with active

transcription, and heterochromatin enriched with H3K9me2 and H3K9me3 (di and tri-methylation

of the 9th lysine residue of the histone H3 protein). Heterochromatin regulates mating and

sporulation gene expression in *S. pombe*, in addition to controlling mating-type switching (Allshire

and Ekwall 2015). To test if this feature is shared in *Pneumocystis*, we performed ChIP-seq with

antibodies targeting histones H3 modifications (H3K9me2/3 and H3K4me2). In *Pneumocystis*, peaks

of H3K9me2 and H3K9me3 are only present at the pericentromeres (Supplementary figures 5,6 and

7). In all three species analyzed (*P. carinii*, *P. murina* and *P. macacae)*, H3K9me2 enrichment is

significantly more pronounced in pericentromeric regions compared to centromeres (**Figure 4;**

Wilcoxon test, *P* < .00001; **Supplementary figure 8**). H3K9me3 enrichment at pericentromeric

regions compared to centromeres is only significant in *P. carinii*. We observed a reduced

transcriptional activity by RNA-seq analysis only in *P. carinii* pericentromeres (**Supplementary**

figure 8). H3K4me2 is correlated with H3K9me2 in the centromeres (Pearson rho = .72) but not at

the whole genome level (r = 0.05). However, research has shown that the histone modifications

H3K4me2 and H3K9me2 decorate heterochromatin and euchromatin regions, respectively (Yasuhara

and Wakimoto 2008). Therefore, the correlation observed between H3K4me2 and H3K9me2 may be

due to the *Pneumocystis* population of cells used in the ChIP analysis. Furthermore, similar to *S.*

pombe (Cam, Sugiyama et al. 2005)*,* a small amount of H3K4me2 is present at the CENP-A

containing of centromeres in *Pneumocystis* (**Supplementary figure 8**). These results suggest that

Pneumocystis centromeres are flanked by heterochromatin.

DNA methylation is frequently associated with centromeres in fungi, where they silence

repeats (Miura, Yonebayashi et al. 2001, Smith, Galazka et al. 2012). To determine whether DNA

methylation plays a role in *Pneumocystis* centromeres, we performed bisulfite sequencing (5

methylcytosine) in *P. carinii* and *P. macacae*. The overall level of 5mC DNA methylation as measured

by the average weighted methylation percentage is 0.6% for *P. carinii* and 2% for *P. macacae* at the CG

dinucleotides (**Supplementary figure 10**). These levels are in range with reported levels for other

fungi e.g., *Verticillium* (0.4%) (Cook, Kramer et al. 2020) and *Neurospora* (2.5%) (Hosseini, Meunier et

al. 2020). To assess the potential role of DNA methylation implication in centromere function, we

analyzed the DNA methylation patterns over different genomic features (genes, intergenic spacers,

and centromeres). DNA methylation levels are higher in genes compared to centromeres (Mann-

Whitney U-test *P* = .005; **Supplementary figure 10**). However, there is no significant difference

among centromeric or pericentromeric regions (defined as 30 kb flanking the centromeres: P-value >

0.3), and the randomly selected genomic regions (genomic background). These results suggest that *5-*

mC DNA methylation is not required for centromere function in *Pneumocystis*, consistent with the

absence of repeats in the centromeres.

Centromere conservation during speciation

Pneumocystis species diverged about 100 Mya (**Figure 2a**). Yet, homologous regions act as

centromeres in *P. carinii*, *P. murina* and *P. macacae* (**Figure 6a, Supplementary table 4**). Centromeres

in these three species have similar DNA sequences and in *P. jirovecii* as well as *P. wakefieldiae*, *P.*

oryctolagi and *P. canis* (**Supplementary figure 11**), which indicates that sequences are vertically

transmitted.

To investigate footprints of selection acting on centromeres, we computed conservation

scores (PhastCons) from whole genome alignments. PhastCons conservation scores range from 0 to 1

and represent probabilities of negative selection. Centromeric regions are more conserved than

genomic background (Wilcoxon test *P* < 2.2e-16**; Figures 6b and 6c**). This suggests that centromeres

are maintained by negative selection. This is consistent with the hypothesis that centromere

positioning tends to be conserved in obligate sexual fungi due to their role in meiosis (Schotanus,

Soyer et al. 2015).

Centromeres may contribute to karyotypic diversity in fungi. However, we found little

support for this hypothesis here because most centromeres do not overlap with chromosomal breaks

(only 3 of 17 in the *P. carinii* versus *P. murina* pairwise comparison) (**Figure 6a**).

Discussion

In the current study we have identified putative centromeres of multiple *Pneumocystis* species and

demonstrated that they are short regional centromeres with pericentromeric heterochromatin and

span active genes (**Figure 7**). Cell replication and chromosomal segregation pathways are unexplored

in these species. Central to these pathways are the centromeres.

As centromeres cannot be predicted bioinformatically, the locations and characteristics of

Pneumocystis centromeres were unknown. Moreover, there are several roadblocks for characterizing

centromeres in *Pneumocystis*, with the most significant one being the lack of continuous culture and

transfection tools. Here we utilized two species of organisms in a short-term culture system to

determine where *pn*CENP-A, the epigenetic marker for centromeres, binds in the genome. We

demonstrated that *Pn*CENP-A functions similarly to CENP-A^{Cnp1} in *S. pombe*, though less efficiently,

validating its use in characterizing *Pneumocystis* centromeres. Unlike the phylogenetically related *S.*

pombe, in *Pneumocystis* these regions overlap with active genes, but like *S. pombe*, these centromeres

are flanked by heterochromatin. Our work provides the first experimental evidence that DNA

methylation occurs in these species, although it may not be involved in centromere function. Taken

together, our results suggest that *Pneumocystis* has small epigenetically regulated centromeres.

Individual centromeres are conserved among different species, though not across chromosomes

within species.

Differences between *P. carinii* and other species analyzed here (*P. murina* and *P. macacae*)

need to be clarified. *P. carinii* and *P. murina* centromeres are homologous, yet there seems to be

heterochromatin silencing only in *P. carinii*. Also, CENP-C is not linked to CENP-A or centromeres

in *P. carinii*, which may instead be using another histone linker (e.g., CENP-T) to connect the

centromere to the kinetochore (Sridhar and Fukagawa 2022).

The presence of genes within centromeres is rare and has only been described in rice (Nagaki,

Cheng et al. 2004) and the plant fungal pathogen *Zymoseptoria tritici* (Schotanus, Soyer et al. 2015).

Centromeres bearing genes are interpreted as young centromeres (neocentromeres), in which the

genes are progressively inactivated. This hypothesis lacks support here because there is no sign of

pseudogenization in these genes: nearly all are expressed and translated, they are conserved in all

species, and many are involved in housekeeping cellular pathways that are presumably critical for

organism survival.

Findings in *Pneumocystis* cannot be generalized to other Taphrinomycotina species. The

determination of ancestral traits between *Pneumocystis* and fission yeast centromeres will require

characterizing the centromeres in additional Taphrinomycotina species. The loss of RNAi is often

associated with shortening of centromeres (Yadav, Sun et al. 2018); consistent with this, *Pneumocystis*

centromeres are much smaller than *Schizosaccharomyces* centromeres.

Our study will benefit from a microscopy validation of CENP-A loading kinetics when a

reliable long term culture system becomes available. Live imaging would help to determine at which

cell cycle stage CENP-A is loaded.

In summary, we have identified atypical short regional centromeres in genetically intractable

micro-organisms. Our results provide insights into the formation of centromeres in host-adapted

fungal pathogens. Our ultimate goal is to use *Pneumocystis* centromeres to stabilize plasmids for

future genetic manipulation. This is the first step along this path, which should lead to better

understanding of the biology of *Pneumocystis* and facilitate the discovery of novel interventions to

effectively control and prevent the disease caused by this pathogen.

MATERIAL AND METHODS

Ethics and organism source

Studies involving human and mouse samples were approved by the NIH Institutional Review Board

(IRB) (IRB protocols 99-I-0084 and CCM 19-05, respectively), rat *Pneumocystis* studies by the Animal

Care and Use Committees of the Cincinnati VA Medical Center (protocol 20-11-08-01) and

macaque *Pneumocystis* studies by the National Institute of Allergy and Infectious Diseases (NIAID)

Division of Intramural Research (protocol LVD 26).

P. carinii organisms were collected from heavily infected lungs of corticosteroid-treated

Sprague-Dawley male rats; *P. murina* from heavily infected lungs of CD40 ligand knock out mouse;

P. macacae from heavily infected lungs of two Simian Immunodeficiency Virus (SIV) infected

macaques (P2C and P3C); and *P. jirovecii* organisms were isolated from a single bronchoalveolar

lavage from one patient and 3 heavily infected autopsy lung tissues from three different patients. The

list of samples is presented in the key resource file.

Pneumocystis culture and growth quantification

P. murina and *P. carinii* were partially purified by Ficoll-Hypaque density gradient centrifugation

(Kovacs, Halpern et al. 1988) and frozen at -80°C in cell recovery media (Gibco). For short term

Pneumocystis cultures, A549 (ATCC) and LET1 (a gift from Dr. Paul Thomas, St Jude Children's

Research Hospital) cells were cultured in culture medium F12 media with 2.5% or 5% heat

inactivated fetal bovine serum (GIBCO) and Penicillin-Streptomycin (GIBCO), plated to

approximately 60 – 80% confluency and incubated for 24h at 37°C. The next day, frozen *Pneumocystis*

vials were thawed, washed in 50ml 1x PBS, and centrifuged at 2,000g for 20 minutes. *Pneumocystis*

cell pellets were resuspended in culture medium and added to the plated cells. Media were partially

changed every 3 or 4 days. At set time points, wells were scraped for collection. Cell/organism

suspensions were centrifuged at 10,000g for 3 min, the supernatant was removed, and the pellets

were frozen at -80°C. Genomic DNA was extracted using QiAmp DNA Extraction Kit (Qiagen).

Quantitative PCR targeting the single copy *dhfr* gene was performed as described previously (Liu,

Davis et al. 2020).

PnCENP-A complementation and ChIP-qPCR in *S. pombe*

Standard procedures were used for fission yeast growth, genetics, and manipulations (Sabatinos and

Forsburg 2010). *P. jirovecii* and *P. murina* CENP-A full-length and N-terminal truncated DNA

sequences were codon optimized for *S. pombe* codon bias (**Supplementary dataset 1**), synthesized

commercially (GenScript, NJ) and subcloned into pS2 vector (Takayama, Sato et al. 2008).

Recombinant plasmids were isolated from *E. coli* strain *DH5α* colonies (Invitrogen) using QIAprep

Spin Miniprep Kit (Qiagen), linearized with *Nsi*I and introduced by transformation at the *lys1+* locus

of *cnp1-76* (Castillo, Mellone et al. 2007) and *cen2-tetO-tdTomato* (Sakuno and Watanabe 2009). For

ChIP-qPCR experiments, *S. pombe* cells were grown overnight at 30°C in rich medium YEA and 2 μl

of anti-GFP (ab290, Abcam) was used for each immunoprecipitation. PCR oligonucleotides detecting

central core (*cc1&3; ura4*), heterochromatic *dg* and euchromatin control *fbp1* were previously

reported (Folco, McCue et al. 2019).

Ortholog identification

Fungal proteomes were retrieved from NCBI. Ortholog identification was performed using

OrthoFinder (Emms and Kelly 2019). Remote homology detection was performed using PSI-BLASTp

(Altschul, Madden et al. 1997) and HMMER v.3.x.x (http://www.hmmer.org) (E-value < 10^{-5}) using *S.*

pombe and *S. cerevisiae* proteins as seeds. Ortholog results were further validated by BLAST searches

on NCBI website, and searches in eggNOG 6.0 (Hernandez-Plaza, Szklarczyk et al. 2023). Gene

reannotations were performed using protein-to-genome alignment with Exonerate (Slater and Birney

2005) followed by manual adjustments if necessary (cf. **Supplementary Methods**). *P. macacae* strain

P3C was sequenced using Oxford nanopore sequencing (cf. **Supplementary Methods**). Protein

multiple sequence alignments were generated using Clustal Omega (Sievers, Wilm et al. 2011).

Maximum likelihood phylogeny was performed using RAxML-ng (Kozlov, Darriba et al. 2019).

Predicted histones phylogenetic classification was performed with reference protein sequences

identified by (Postberg, Forcob et al. 2010). Immunogenetic peptides were selected for *Pneumocystis*

CENP-A, CENP-C and mis12 orthologs, used for antibody production and quality control

(**Supplementary Methods**).

Light, electron and immunofluorescence microscopy

P. murina and A549-LET1 cells described above were grown on Thermanox™ coverslips (Ted Pella,

Redding, CA, USA), harvested at day 7, fixed with 2% paraformaldehyde/2.5% glutaraldehyde in 0.1

M Sorenson's phosphate buffer and then post-fixed with 1.0% osmium tetroxide/0.8% potassium

ferricyanide in 0.1 M sodium cacodylate buffer for 1 hour, washed with buffer then stained with

1% tannic acid in dH_2O for 1 hr. After additional buffer washes, the samples were further

osmicated with 2% osmium tetroxide in 0.1M sodium cacodylate, then washed with dH_2O.

Specimens were then stained overnight at 4C with 1% aqueous uranyl acetate. The cells were

then washed with dH_2O and dehydrated with a graded ethanol series, prior to embedding in

Spurr's resin. Thin sections were cut with a Leica UC7 ultramicrotome (Buffalo Grove, IL) prior

to viewing at 120 kV on a FEI BT Tecnai transmission electron microscope (ThermoFisher/FEI,

Hillsboro, OR). Digital images were acquired with a Gatan Rio camera (Gatan, Pleasanton,

CA).

Fluorescence labelling of *Pneumocystis*-infected tissues was performed by Histoserv, Inc

(Germantown, Maryland) and visualized using Leica DMi8 microscope (DMI Medical Inc).

S. pombe cells were grown overnight at 32°C in minimal medium PMG until logarithmic

phase and then mounted in PMG 2% agarose as described (Tran, Paoletti et al. 2004). Images were

acquired on a Delta Vision Elite microscope (Applied Precision) with a 100X 1.35 NA oil lens

(Olympus). Twenty 0.35-μm z sections were acquired generating maximum intensity projections.

Further image processing including background-subtracted centromere foci intensity measurements,

was performed using ImageJ (National Institutes of Health).

ChIP-seq

Purified cells (~5 x 10^6 cells) and tissue preparations were fixed by 37% formaldehyde treatment.

Chromatin preparation, immunoprecipitation and Illumina sequencing libraires were prepared

using the Low Cell ChiP-Seq kit (Active Motif, Carlsbad, CA) according to the manufacturer

instructions. ChIP-seq libraries were sequenced commercially (Psomagen Rockville, Maryland, USA)

using Illumina NovaSeq 6000 (150-bp paired end reads).

RNA-Seq

Total RNA was extracted using RNeasy Mini kit (Qiagen). RNA quality and integrity were

estimated using Bioanalyzer RNA 6000 Pico Assay (Agilent) and sequenced commercially using

Illumina HiSeq 4000 (150-base paired end libraries, Novogene Inc, USA). Reads were mapped using

STAR v.2.7.9a (Dobin, Davis et al. 2013), and normalized to bins per million mapped reads (BPM).

Transcripts counts were quantified using kallisto 0.44.0 (Bray, Pimentel et al. 2016).

ChIP-seq peak calling and sequence analysis

Raw reads were quality-checked with multi-QC (Ewels, Magnusson et al. 2016). Host reads (human,

macaque, rat, mice) were removed by mapping using Bowtie2 (Langmead and Salzberg 2012) with

default parameters. Unmapped reads were retrieved and mapped to *Pneumocystis* reference genomes

(*P. murina* (GenBank accession number GCA_000349005.2), *P. carinii* (GCA_001477545.1), *P.*

macacae (GCA_018127085.1) and *P. jirovecii* (GCA_001477535.1)) using the Burrows-Wheeler

Aligner BWA-MEM (Li 2013). Alignments were filtered and sorted using SAMTools (Li, Handsaker

et al. 2009) and duplicates marked using PICARD MarkDuplicates

http://broadinstitute.github.io/picard). Quality controls were performed using deepTools

plotFingerprint (Ramirez, Ryan et al. 2016) and phantompeakqualtools (Landt, Marinov et al. 2012),

which estimate the normalized strand coefficient (NSC) and the relative strand correlation (RSC)**.**

Chip-Seq coverage data were calculated as 50-bp bin and normalized per genomic content (1x

normalization). Chromosome ideograms were generated using chromoMap v0.3 (Lakshay Anand

2020).

ChIP enrichment peaks were detected using <u>M</u>odel-Based <u>A</u>nalysis for <u>Ch</u>IP-<u>S</u>equencing

(Zhang, Liu et al. 2008)(MACS3 v3.0.0a5) callpeak function with default parameters and estimated

genome sizes and sorted by fold-enrichment and FDR values (broad cutoff 0.1; FDR < 0.05).

Enriched peaks were further inspected using deepTools bamCompare versus shuffle non centromeric

genomic regions. Peaks were further examined in conjunction with other data (e.g. conservation

score, RNA-seq, methylation data, GC content, repeats) with the <u>I</u>ntegrative <u>G</u>enomics <u>V</u>iewer

(Robinson, Thorvaldsdottir et al. 2011), ensuring that peaks are present in all IP samples and absent

in controls. *De novo* DNA motif searches were performed using MEME v5.1.0 (Bailey, Boden et al.

2009) and Homer v4.9 (Heinz, Benner et al. 2010) with a p-value cut off of 10^{-10}. GC content across

windows (bin) was computed using BEDTools (Quinlan and Hall 2010). Normalized coverage and

annotations across chromosomes were visualized using pyGenomeTracks (Lopez-Delisle, Rabbani et

al. 2021).

Repeated elements (DNA transposons, retrotransposons and low complexity repeats) were

identified using RepeatMasker (Smit AFA. 2013-2015) and RepBase (Bao, Kojima et al. 2015).

Genome synteny and conservation

DNA alignments were generated using Satsuma2 (Grabherr, Russell et al. 2010) and visualized with

Circos (Krzywinski, Schein et al. 2009). To estimate the sequence conservation scores, we performed

one-to-one pairwise whole genome alignments using LAST (Kielbasa, Wan et al. 2011) with the

MAM4 seeding scheme (Frith and Noe 2014). We consider two trios (1) *P. jirovecii/P. macacae/P.*

oryctolagi and (2) *P. carinii/P. murina/P. wakefieldiae.* We generated a neutral model using phyloFit

(Siepel and Haussler 2004) from PHAST which fit the tree model to multiple sequence alignments by

maximum likelihood using the REV substitution model. Sequence conservation scores were

estimated using phastCons (Siepel, Bejerano et al. 2005) (--target-coverage 0.25 --expected-length 20 --

estimate-trees). Wig files and Bedgraph files were converted using 'wigToBigWig' and

'bigWigToBedGraph' tools (Kent, Zweig et al. 2010).

Bisulfite sequencing

Pneumocystis DNA (5 µg) was extracted by a *Pneumocystis* DNA enrichment protocol (Ma, Chen et al.

2016). Bisulfite conversion, library preparation and sequencing (Illumina NovaSeq 150 bp paired end

reads) were performed commercially (Novogene). Bisulfite conversion of non-methylated DNA was

performed using EZ DNA Methylation kit. Data analysis was performed using BSMAP and

methratio script (Xi and Li 2009). Only cytosine positions with more than 5-fold coverage were

considered. We used weighted methylation percentage (Cook, Kramer et al. 2020), which is

calculated as the number of reads supporting methylation over the number of cytosines sequenced to

quantify methylation levels. Methylation data were partitioned over different genomic compartments

using BEDTools (Quinlan 2014). Statistical comparisons were performed using R (https://www.R-

project.org) to compute non-parametric Mann-Whitney U test and Bonferroni correction to adjust *P*-

values for multiple comparisons.

ChIP-qPCR

CENP-A enrichment and H3 depletion relative to H4 within the centromeres were evaluated using

primers specific to centromeric and non-centromeric loci (see **Supplementary Table 5** for primers).

Primers were designed using primer3 (Koressaar, Lepamets et al. 2018) from regions visualized using

IGV genome browser. Primers with potential matches against mammal genomes were excluded

using NCBI BLASTn. Dilutions of 1:10 were used based on preliminary test runs. Each reaction

contains 5 μl of iTaq universal Sybr green supermix (Biorad), 1.25 μl of primers (500 nM)) and 2.5 μl

of DNA. All assays were run on CFX96 thermocycler (Biorad). Three technical replicates were taken

for each assay, and the standard errors of the mean were calculated. The PCR program was as follow:

initial denaturation for 2 min at 95°C, followed by 30 cycles of 30 seconds at 95°C, 60°C for 30s, and

72° for 30s. Locus ΔCt values were normalized using the formula: ($^{\Delta\Delta Ct}$ ChIP – $^{\Delta Ct}$ Input;

https://www.sigmaaldrich.com/US/en/technical-documents/protocol/genomics/qpcr/chip-qpcr-data-

analysis) and the fold enrichment of the ChIP DNA over the input was computed as log2 ($2^{\Delta\Delta Ct}$). The

plots and statistical analyses were performed with GraphPad Prism 8.

ChIP-LC-MS

Formalin-fixed purified chromatin preparations were separately co-immunoprecipitated with anti

CENP-A, CENP-C, H4 and pre immune serum (control) using Pierce™ Direct Magnetic IP/Co-IP

Kit (ThermoFisher). Each IP reaction was performed separately on aliquots from the same chromatin

preparation. Elution was performed using 1% formic acid. Extracts were frozen in dry ice, lyophilized

for 1 hour and digested using trypsin, diluted, and injected to a Thermo Orbitrap Fusion LC-MS/MS

to identify unique peptides. LC-MS/MS data were searched against the proteomes using Proteome

Discoverer 2.4 (Thermo Fisher Scientific, Waltham, MA). Label-free quantification was analyzed

based on the peak intensity of the precursor ion.

Data availability

ChIP-seq datasets and associated MACS2 outputs generated in this study (CENP-A, CENP-C, Mis12,

H3, H4, H3K9me2/3, H3K4me2) are deposited at NCBI GEO (accession numbers GSE226536,

GSE230598).

Code availability

Codes generated for this project are available at GitHub: https://github.com/ocisse/pneumo-cens

Multi panel figures were assembled in Inkscape using original plots.

Acknowledgements

We thank Rene Costello for animal studies, Drs. Keytam Awad, Salina Gairhe and Shuibang Wang

for discussion about ChIP-seq. This work has been funded in whole or in part with federal funds

from the Intramural Research Program of the US National Institutes of Health (NIH) Clinical Center

and the National Institute of Allergy and Infectious Diseases (NIAID). The views expressed in this

work are those of the authors and do not necessarily represent the official positions of the NIH or

U.S. Government. This study used the Office of Cyber Infrastructure and Computational Biology

(OCICB) High Performance Computing (HPC) cluster at the National Institute of Allergy and

Infectious Diseases (NIAID), Bethesda, MD. This study also utilized the high-performance

computational capabilities of the Biowulf Linux cluster at the NIH, Bethesda, MD

(http://biowulf.nih.gov).

Figure legends

Figure 1. *Pneumocystis* CENP-A variants localize to the nucleus, and support viability and centromere loading in a heterologous system.

(**a**) Protein sequence alignment of full-length CENP-A^{cnp-1} homologs in *Schizosaccharomyces pombe* and seven *Pneumocystis* species. Each *Pneumocystis* species only infects a single mammalian host: *P. jirovecii* (infecting humans), *P. macacae* (macaques), *P. oryctolagi* (rabbits), *P. canis* (dogs), *P. murina* (mice), *P. carinii* (rats), and *P. wakefieldiae* (rats). The ~25 amino acid insert in *Pneumocystis* CENP-A N-terminal regions is unique to this genus.

(**b**) Immunofluorescence microscopy showing that anti-*Pneumocystis murina* CENP-A (*pn*CENP-A) reacts with *P. murina* organisms. *P. murina*-infected CD40 ligand knockout lung tissue sections were dual labeled using an anti CENP-A antibody(magenta) and 4′,6-diamidino-2-phenylindole (DAPI, light blue). The image is a maximum projection of a 5 µm thick z-stack. The images were minimally adjusted to enhance the contrast. The green box is magnified in the inset showing *pn*CENP-A inside *Pneumocystis* organisms (small dots). *pn*CENP-A signals overlap or are close in proximity with DAPI (represented by a white color produced by the overlap between magenta and light blue colors).

(**c**) Left, rescue experiment of fission yeast *cnp1-76* thermosensitive cells using the GFP-tagged CENP-A transgenes. Right, protein sequence alignment of *Pneumocystis* CENP-A N-terminal regions (*P. jirovecii* and *P. murina*) with conserved amino acids highlighted (grey areas). Plasmids containing full length and N-terminal region truncated (**) versions of *Pneumocystis* CENP-A were constructed and integrated as single copy. Growth was assayed at indicated temperatures using 10-fold serial dilutions plated on rich YEA medium. The transgene encoding *S. pombe* CENP-A serves as a non-temperature-sensitive control.

(**d**) Representative images of *cen2-tetO-TdTomato* strains expressing the indicated GFP-tagged CENP-A transgenes. On the bottom, GFP and tdTomato fluorescence as well as brightfield images are merged.

(**e**) Integrated fluorescence intensity of GFP foci was measured and plotted for the strains used in panel e. Mean (bars) and SD (error bars) are shown. n>176. AU, arbitrary units.

(**f**) Anti-GFP ChIP-qPCR analysis of indicated loci for strains used in panel e. %IP represents the percentage of input that was immunoprecipitated. Error bars denote the SD (n = 3). Abbreviations: cc1 &3, *S. pombe* centromere central core 1 & 3; cnt2:ura4$^+$, ura4$^+$ insertion at central core 2; *dg*, a class of heterochromatic outer repeats within pericentromeric regions (control); *fbp1*, euchromatic locus (control)

Figure 2. *Pneumocystis* CENP-A binds to single genomic foci in replicating cells

a) Study workflow and sample preparation. Four *Pneumocystis* species infecting mice, rats, macaques, and humans respectively were analyzed. *P. murina* and *P. carinii* were obtained from CD40 ligand knock-out female mice and immunosuppressed Sprague-Dawley male rats, respectively, and cultured on a co-culture of human lung adenocarcinoma cells (A549) and immortalized murine lung epithelial type 1 cells (Let-1) for 14 days. *P. macacae* organisms were obtained from a Simian Immunodeficiency Virus infected macaque (P2C), and *P. jirovecii* organisms were obtained from a single bronchoalveolar lavage from a non-infected HIV patient and three autopsy samples from patients with HIV infection. Species phylogeny and estimated speciation timing are presented. Animal icons were obtained from http://phylopic.org under creative commons licenses https://creativecommons.org/licenses/by/3.0/: mouse (Anthony Caravaggi; license CC BY-NC-SA 3.0); and rat (by Rebecca Groom; license CC BY-NC-SA 3.0).

b) *P. murina* population growth in duplicate wells were measured by quantitative PCR targeting a single copy dihydrofolate reductase (*dhfr*) gene over 14 days. Error bars represent the standard deviation (*n*=2).

c) *P. carinii* growth measurement by qPCR targeting *dhfr* gene. Error bars represent the standard deviation (*n*=2).

d) Electron micrograph showing a possibly dividing *P. murina* trophic form seven days post culture (black arrow). The field also include a non-dividing trophozoite (white arrow). 2 μm scale bar.

e) *pn*CENP-A ChIP-seq study in *P. carinii* organisms from infected lungs and cultures at days 0, 7 and 14. CENP-A remains strongly enriched at a single genomic location up to 7 days. Only one representative chromosome is presented. The remaining 16 chromosomes have similar ChIP-seq profiles and presented in supplementary figure 4.

Figure 3. *Pneumocystis* displays seventeen centromeres.

a) Scaled ideogram of 17 chromosomal level scaffolds (grey) of *P. murina* genome showing
CENP-A binding regions (constricted areas) and telomeric repeat motif (red).
b) CENP-A and CENP-C binding regions delineate putative centromeres (CENs) in *P. murina*
genome. Color coded peaks represent enrichment of immunoprecipitated DNA (IP DNA)
relative to controls (Input DNA). The 1-kb flanking sequences from the center of the
enrichment peak are presented. Each scaffold displays two peaks labelled M (Major) and m
(minor) according to the enrichment level.
c) Scaled ideogram of 17 chromosomal level scaffolds (grey) of *P. carinii* genome showing
CENP-A binding regions (constricted areas) and telomeric repeat motif (red).
d) CENP-A binding regions delineate putative centromeres (CENs) in *P. carinii* genome. Note
that no peak was observed with anti-CENP-C antibody.
e) A DNA dot plot of CENP-A binding regions in *P. murina* showing regional self-similarity.
The main diagonal represents the sequence alignment with itself. Lines off the main diagonal
which are repetitive patterns within the sequences are not observed. The plot shows that each
CEN is unique and repeat-free.
f) DNA dot plot of CENP-A binding regions in *P. carinii* showing that each CEN is unique and
repeat-free.

Figure 4. Centromeres are flanked by heterochromatin and contain active genes.

All three scaffolds are homologous.

    a) Genomic view of chromosome 2 of *P. murina* genome subsequently showing annotated genes (directed grey boxes), DNA repeats (not present), percent GC content (blue), ChIP-seq read coverage distribution (BPM normalized over bins of 50 bp; input subtracted) of CENP-A, CENP-C, histones H3 and H4 ratio, heterochromatin-associated modifications (H3K9me2 and H3K9me3), euchromatin (H3K4me2) and gene expression (RNA-seq) in relation with centromeres. Genome views of remaining 16 chromosomes are presented in Supplementary dataset figure 7.

    b) Genomic view of chromosome 2 of *P. carinii* genome with the same features presented as *P. murina*. A duplicated copy of *copia*-retrotransposon is presented (red arrows), which is present in syntenic regions in other *Pneumocystis* genomes. Genome views of all remaining 16 chromosomes are presented in supplementary figure 6.

    c) Genomic view of chromosome 14 of *P. macacae* genome showing the distribution of gene density, repeats (DNA transposons), homology region with GC content, mis12 ChIP-seq read coverage, chromatin-associated modifications (H3K9me2 and H3K9me3 and H3K4me2) and RNA-seq in relation with centromeres. The white box indicates a DNA sequence homologous to *P. carinii* centromere 2. A single copy of the DNA transposon EnSPM (CACTA family) and the class 2DNA transposon Helitron are present upstream the centromere (forward red arrows). These elements are not present in syntenic regions in other *Pneumocystis* genomes. Genome views of all remaining 15 chromosomes are presented in Supplementary figure 5.

Figure 5. Modeling of Molecular dynamics of CENP-A nucleosomes in *P. carinii*, *P. murina* and *S. pombe.*

a) Upper panel: Sequence alignment of the centromeric histones used in modeling and simulations. Differences between *P. carinii* and *P. murina* colored green and red; differences between *S. pombe* and *pneumocystis* colored blue. Lower panel: surface representation of the centromeric octamers of *P. carinii* (left), *P. murina* (middle), and *S. pombe* (right), showing the spatial location of the mutations (snapshots from the simulations); DNA (not simulated) shown only as a reference based on the crystal structure of the human centromere nucleosome, PDB: 3AN2). The white circles show the central crevices, the behaviors of which are expected to dictate the assembly of the inner kinetochore in each species.

b) Aggregation of snapshots along the dynamic trajectory illustrating the extent of species-specific conformational variations of the central crevices (see Animation in Supplementary dataset 2); histones color code is presented at the bottom). The C-terminal tip of CENP-A is located at the center of the core (white ribbon; residues 146-151 in *Pneumocystis* and 120 in *S. pombe*; see alignment in panel a). The observed differences between *P. carinii* and *P. murina* are due to structural and dynamic changes induced by the different (H2A:H2B) dimers (cf. alignment in panel a); the differences between these species and *S. pombe* are likely the result of more extensive sequence variations in the four histones, including several solvent-exposed mutations at the core center (cf. panel a). Early-stage assembly of the inner kinetochore is thus expected to be strongly determined by features of this crevice, including its size, dynamics, and other physicochemical features.

747 **Figure 6. Centromere locations, not sequences, are evolutionary constrained by negative**
748 **selection.**

    a) Genome organization and synteny in *Pneumocystis*. Circos plots depicting pairwise *P. carinii*,
750   *P. murina* and *P. macacae* genome synteny. Rodent infecting Pneumocystis (*P. carinii* and *P.*
751   *murina*) and macaques infecting *Pneumocystis* (*P. macacae*) have 17 and 16 chromosomal level
752   scaffolds, respectively. Colored connectors indicate regions of synteny between species.
753   Centromeres that overlap with recent chromosomal breakpoints are indicated (*). The square
754   highlights *P. carinii* centromere 4 displayed in the panel b.

    b) Genome view of centromere 4 in the *P. carinii* genome. Genes are represented by directed
756   boxes (grey for protein coding genes and cyan for polymorphic major surface glycoprotein
757   genes), *pn*CENP-A binding region (centromere) and sequence conservation scores which
758   were calculated from whole genome alignments of *P. carinii*. *P. murina* and *P. wakefieldiae*
759   (PhasCons). The phastCons scores represent probabilities of negative selection and range
760   between 0 (no conservation) and 1 (total conservation).

    c) Boxplot of conservation scores per genomic context summarized over all *P. carinii*
762   centromeres (CEN), 30-kb regions flanking the centromeres (Cenflk), major surface
763   glycoproteins (Msg) and random genomic background (Bckg). Statistical differences for the
764   indicated comparisons were obtained using one-sided nonparametric Mann-Whitney test; *P*-
765   values < .0001, ****; *P* < .001, ***; *P* < .01, **; *P* < .05, *; ns, not significant.

768
Figure 7. Model of regional centromere structures in *Pneumocystis* and other representative
fungi.
On the left is presented the phylogeny of selected fungi inferred from maximum likelihood
phylogenetic analysis of shared core protein orthologs. The overall centromere structures for
Pneumocystis carinii (details supporting our model are provided in the text), *Schizosaccharomyces pombe*
(Pidoux and Allshire 2004), *Candida albicans* (Sanyal, Baum et al. 2004), *Zymoseptoria tritici*
(Schotanus, Soyer et al. 2015) and *Cryptococcus neoformans* (Janbon, Ormerod et al. 2014) are used as
representative to showcase the diversity of regional centromeres in fungi. Animal pathogens are
highlighted in pale olive and the plant fungal pathogen *Zymoseptoria tritici* is in pale green. For sake
of brevity, only *Pneumocystis carinii* is presented for brevity, although there are differences with other
Pneumocystis species. In the middle are presented DNA structures of centromeres. *P. carinii* has 17
centromeres (one for each of its 17 chromosomes) that share the same overall architecture.
Centromeres are delineated by a localized enrichment of the centromeric histone CENP-A, which
overlap with a reduction of the canonical histone H3 (inverted grey triangle). Centromeres are
flanked by heterochromatin H3K9me (stands here stands for both H3K9me2 and H3K9me3). All 17
Pneumocystis carinii centromeres span active genes (dark grey boxes). Each centromere sequence is
different and lacks shared DNA sequence motif. *S. pombe* has three centromeres that share the same
overall structure in which a central core (*cnt*) domain is surrounded by innermost repeats (*imr*) and
outer repeats (*otr*). The *imr* repeats incorporate clusters of transfer RNAs (tRNAs) that play a role in
restricting CENP-A spread. *S. pombe* centromeres are flanked by heterochromatin (H3K9me). Genes
are found 0.75-1.5 kb beyond the limits of the centromeres. *C. albicans* has 8 unique and different
centromeres that are gene free and lack shared sequence motifs. *Z. tritici* has 21 centromeres ranging
from 6 to 14 kb in size, that partially overlap with genes. *C. neoformans* has 14 centromeres that are
gene free and enriched with Tcn transposons.
793
794

829   **Literature Cited**
830
Allshire, R. C. and K. Ekwall (2015). "Epigenetic Regulation of Chromatin States in
Schizosaccharomyces pombe." <u>Cold Spring Harb Perspect Biol</u> **7**(7): a018770.
Almeida, J. M., O. H. Cisse, A. Fonseca, M. Pagni and P. M. Hauser (2015). "Comparative genomics
suggests primary homothallism of Pneumocystis species." <u>MBio</u> **6**(1).
Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller and D. J. Lipman (1997).
"Gapped BLAST and PSI-BLAST: a new generation of protein database search programs." <u>Nucleic</u>
<u>Acids Res</u> **25**(17): 3389-3402.
Bailey, T. L., M. Boden, F. A. Buske, M. Frith, C. E. Grant, L. Clementi, J. Ren, W. W. Li and W. S.
Noble (2009). "MEME SUITE: tools for motif discovery and searching." <u>Nucleic Acids Res</u> **37**(Web
Server issue): W202-208.
Bao, W., K. K. Kojima and O. Kohany (2015). "Repbase Update, a database of repetitive elements in
eukaryotic genomes." <u>Mob DNA</u> **6**: 11.
Bray, N. L., H. Pimentel, P. Melsted and L. Pachter (2016). "Near-optimal probabilistic RNA-seq
quantification." <u>Nat Biotechnol</u> **34**(5): 525-527.
Cam, H. P., T. Sugiyama, E. S. Chen, X. Chen, P. C. FitzGerald and S. I. Grewal (2005).
"Comprehensive analysis of heterochromatin- and RNAi-mediated epigenetic control of the fission
yeast genome." <u>Nat Genet</u> **37**(8): 809-819.
Casola, C., D. Hucks and C. Feschotte (2008). "Convergent domestication of pogo-like transposases
into centromere-binding proteins in fission yeast and mammals." <u>Mol Biol Evol</u> **25**(1): 29-41.
Castillo, A. G., B. G. Mellone, J. F. Partridge, W. Richardson, G. L. Hamilton, R. C. Allshire and A.
L. Pidoux (2007). "Plasticity of fission yeast CENP-A chromatin driven by relative levels of histone
H3 and H4." <u>PLoS Genet</u> **3**(7): e121.
Chen, J. W., L. Yang, C. Santos, S. A. Hassan, P. L. Collins, U. J. Buchholz and C. Le Nouen (2021).
"Reversion mutations in phosphoprotein P of a codon-pair-deoptimized human respiratory syncytial
virus confer increased transcription, immunogenicity, and genetic stability without loss of
attenuation." <u>PLoS Pathog</u> **17**(12): e1010191.
Cisse, O. H., L. Ma, J. P. Dekker, P. P. Khil, J. H. Youn, J. M. Brenchley, R. Blair, B. Pahar, M. Chabe,
K. K. A. Van Rompay, R. Keesler, A. Sukura, V. Hirsch, G. Kutty, Y. Q. Liu, L. Peng, J. Chen, J. Song,
C. Weissenbacher-Lang, J. Xu, N. S. Upham, J. E. Stajich, C. A. Cuomo, M. T. Cushion and J. A.
Kovacs (2021). "Genomic insights into the host specific adaptation of the Pneumocystis genus."
<u>Communications Biology</u> **4**(1).
Cisse, O. H., M. Pagni and P. M. Hauser (2012). "De novo assembly of the Pneumocystis jirovecii
genome from a single bronchoalveolar lavage fluid specimen from a patient." <u>MBio</u> **4**(1): e00428-
00412.
Cisse, O. H., M. Pagni and P. M. Hauser (2014). "Comparative Genomics Suggests That the Human
Pathogenic Fungus Pneumocystis jirovecii Acquired Obligate Biotrophy through Gene Loss."
<u>Genome Biology and Evolution</u> **6**(8): 1938-1948.

Clarke, L. and J. Carbon (1980). "Isolation of a yeast centromere and construction of functional small

circular chromosomes." <u>Nature</u> **287**(5782): 504-509.

Cook, D. E., H. M. Kramer, D. E. Torres, M. F. Seidl and B. Thomma (2020). "A unique chromatin

profile defines adaptive genomic regions in a fungal plant pathogen." <u>Elife</u> **9**.

Corbett, K. D., C. K. Yip, L. S. Ee, T. Walz, A. Amon and S. C. Harrison (2010). "The Monopolin

Complex Crosslinks Kinetochore Components to Regulate Chromosome-Microtubule Attachments."

<u>Cell</u> **142**(4): 556-567.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson and T. R.

Gingeras (2013). "STAR: ultrafast universal RNA-seq aligner." <u>Bioinformatics</u> **29**(1): 15-21.

Emms, D. M. and S. Kelly (2019). "OrthoFinder: phylogenetic orthology inference for comparative

genomics." <u>Genome Biol</u> **20**(1): 238.

Ewels, P., M. Magnusson, S. Lundin and M. Kaller (2016). "MultiQC: summarize analysis results for

multiple tools and samples in a single report." <u>Bioinformatics</u> **32**(19): 3047-3048.

Folco, H. D., C. S. Campbell, K. M. May, C. A. Espinoza, K. Oegema, K. G. Hardwick, S. I. S. Grewal

and A. Desai (2015). "The CENP-A N-tail confers epigenetic stability to centromeres via the CENP-T

branch of the CCAN in fission yeast." <u>Curr Biol</u> **25**(3): 348-356.

Folco, H. D., A. McCue, V. Balachandran and S. I. S. Grewal (2019). "Cohesin Impedes

Heterochromatin Assembly in Fission Yeast Cells Lacking Pds5." <u>Genetics</u> **213**(1): 127-141.

Freitag, M. (2017). "Histone Methylation by SET Domain Proteins in Fungi." <u>Annu Rev Microbiol</u>

71: 413-439.

Frith, M. C. and L. Noe (2014). "Improved search heuristics find 20,000 new alignments between

human and mouse genomes." <u>Nucleic Acids Res</u> **42**(7): e59.

Funabiki, H., I. Hagan, S. Uzawa and M. Yanagida (1993). "Cell cycle-dependent specific positioning

and clustering of centromeres and telomeres in fission yeast." <u>J Cell Biol</u> **121**(5): 961-976.

Gordon, J. L., K. P. Byrne and K. H. Wolfe (2011). "Mechanisms of chromosome number evolution

in yeast." <u>PLoS Genet</u> **7**(7): e1002190.

Grabherr, M. G., P. Russell, M. Meyer, E. Mauceli, J. Alfoldi, F. Di Palma and K. Lindblad-Toh

(2010). "Genome-wide synteny through highly sensitive sequence alignment: Satsuma."

<u>Bioinformatics</u> **26**(9): 1145-1151.

Hayashi, M. T., T. S. Takahashi, T. Nakagawa, J. Nakayama and H. Masukata (2009). "The

heterochromatin protein Swi6/HP1 activates replication origins at the pericentromeric region and

silent mating-type locus." <u>Nature Cell Biology</u> **11**(3): 357-U284.

Hayashi, T., Y. Fujita, O. Iwasaki, Y. Adachi, K. Takahashi and M. Yanagida (2004). "Mis16 and

Mis18 are required for CENP-A loading and histone deacetylation at centromeres." <u>Cell</u> **118**(6): 715-

729.

Heinz, S., C. Benner, N. Spann, E. Bertolino, Y. C. Lin, P. Laslo, J. X. Cheng, C. Murre, H. Singh and

C. K. Glass (2010). "Simple Combinations of Lineage-Determining Transcription Factors Prime cis-

Regulatory Elements Required for Macrophage and B Cell Identities." <u>Molecular Cell</u> **38**(4): 576-589.

Hernandez-Plaza, A., D. Szklarczyk, J. Botas, C. P. Cantalapiedra, J. Giner-Lamia, D. R. Mende, R.

Kirsch, T. Rattei, I. Letunic, L. J. Jensen, P. Bork, C. von Mering and J. Huerta-Cepas (2023).

"eggNOG 6.0: enabling comparative genomics across 12 535 organisms." Nucleic Acids Res **51**(D1): D389-D394.

Hosseini, S., C. Meunier, D. Nguyen, J. Reimegard and H. Johannesson (2020). "Comparative analysis of genome-wide DNA methylation in Neurospora." Epigenetics **15**(9): 972-987.

Janbon, G., K. L. Ormerod, D. Paulet, E. J. Byrnes, 3rd, V. Yadav, G. Chatterjee, N. Mullapudi, C. C. Hon, R. B. Billmyre, F. Brunel, Y. S. Bahn, W. Chen, Y. Chen, E. W. Chow, J. Y. Coppee, A. Floyd-Averette, C. Gaillardin, K. J. Gerik, J. Goldberg, S. Gonzalez-Hilarion, S. Gujja, J. L. Hamlin, Y. P. Hsueh, G. Ianiri, S. Jones, C. D. Kodira, L. Kozubowski, W. Lam, M. Marra, L. D. Mesner, P. A. Mieczkowski, F. Moyrand, K. Nielsen, C. Proux, T. Rossignol, J. E. Schein, S. Sun, C. Wollschlaeger, I. A. Wood, Q. Zeng, C. Neuveglise, C. S. Newlon, J. R. Perfect, J. K. Lodge, A. Idnurm, J. E. Stajich, J. W. Kronstad, K. Sanyal, J. Heitman, J. A. Fraser, C. A. Cuomo and F. S. Dietrich (2014). "Analysis of the genome and transcriptome of Cryptococcus neoformans var. grubii reveals complex RNA expression and microevolution leading to virulence attenuation." PLoS Genet **10**(4): e1004261.

Kato, H., J. Jiang, B. R. Zhou, M. Rozendaal, H. Feng, R. Ghirlando, T. S. Xiao, A. F. Straight and Y. Bai (2013). "A conserved mechanism for centromeric nucleosome recognition by centromere protein CENP-C." Science **340**(6136): 1110-1113.

Kent, W. J., A. S. Zweig, G. Barber, A. S. Hinrichs and D. Karolchik (2010). "BigWig and BigBed: enabling browsing of large distributed datasets." Bioinformatics **26**(17): 2204-2207.

Kielbasa, S. M., R. Wan, K. Sato, P. Horton and M. C. Frith (2011). "Adaptive seeds tame genomic sequence comparison." Genome Res **21**(3): 487-493.

Koressaar, T., M. Lepamets, L. Kaplinski, K. Raime, R. Andreson and M. Remm (2018). "Primer3_masker: integrating masking of template sequence with primer design software." Bioinformatics **34**(11): 1937-1938.

Kovacs, J. A., J. L. Halpern, J. C. Swan, J. Moss, J. E. Parrillo and H. Masur (1988). "Identification of antigens and antibodies specific for Pneumocystis carinii." J Immunol **140**(6): 2023-2031.

Kozlov, A. M., D. Darriba, T. Flouri, B. Morel and A. Stamatakis (2019). "RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference." Bioinformatics **35**(21): 4453-4455.

Krzywinski, M., J. Schein, I. Birol, J. Connors, R. Gascoyne, D. Horsman, S. J. Jones and M. A. Marra (2009). "Circos: an information aesthetic for comparative genomics." Genome Res **19**(9): 1639-1645.

Kursel, L. E. and H. S. Malik (2016). "Centromeres." Curr Biol **26**(12): R487-R490.

Lakshay Anand, C. M. R. L. (2020). "chromoMap: An R package for Interactive Visualization and Annotation of Chromosomes." bioRxiv.

Lando, D., U. Endesfelder, H. Berger, L. Subramanian, P. D. Dunne, J. McColl, D. Klenerman, A. M. Carr, M. Sauer, R. C. Allshire, M. Heilemann and E. D. Laue (2012). "Quantitative single-molecule microscopy reveals that CENP-A(Cnp1) deposition occurs during G2 in fission yeast." Open Biology **2**.

Landt, S. G., G. K. Marinov, A. Kundaje, P. Kheradpour, F. Pauli, S. Batzoglou, B. E. Bernstein, P. Bickel, J. B. Brown, P. Cayting, Y. Chen, G. DeSalvo, C. Epstein, K. I. Fisher-Aylor, G. Euskirchen, M. Gerstein, J. Gertz, A. J. Hartemink, M. M. Hoffman, V. R. Iyer, Y. L. Jung, S. Karmakar, M. Kellis,

P. V. Kharchenko, Q. Li, T. Liu, X. S. Liu, L. Ma, A. Milosavljevic, R. M. Myers, P. J. Park, M. J.
Pazin, M. D. Perry, D. Raha, T. E. Reddy, J. Rozowsky, N. Shoresh, A. Sidow, M. Slattery, J. A.
Stamatoyannopoulos, M. Y. Tolstorukov, K. P. White, S. Xi, P. J. Farnham, J. D. Lieb, B. J. Wold and
M. Snyder (2012). "ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia."
<u>Genome Res</u> **22**(9): 1813-1831.
Langmead, B. and S. L. Salzberg (2012). "Fast gapped-read alignment with Bowtie 2." <u>Nat Methods</u>
9(4): 357-359.
Leverson, J. D., H. K. Huang, S. L. Forsburg and T. Hunter (2002). "The Schizosaccharomyces pombe
aurora-related kinase Ark1 interacts with the inner centromere protein Pic1 and mediates
chromosome segregation and cytokinesis." <u>Mol Biol Cell</u> **13**(4): 1132-1143.
Li, H. (2013). "Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM."
<u>arXiv</u>: 1303.3997.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin
and S. Genome Project Data Processing (2009). "The Sequence Alignment/Map format and
SAMtools." <u>Bioinformatics</u> **25**(16): 2078-2079.
Liu, X., I. McLeod, S. Anderson, J. R. Yates, 3rd and X. He (2005). "Molecular analysis of kinetochore
architecture in fission yeast." <u>EMBO J</u> **24**(16): 2919-2930.
Liu, Y., A. S. Davis, L. Ma, L. Bishop, O. H. Cisse, G. Kutty and J. A. Kovacs (2020). "MUC1 mediates
Pneumocystis murina binding to airway epithelial cells." <u>Cell Microbiol</u> **22**(6): e13182.
Lopez-Delisle, L., L. Rabbani, J. Wolff, V. Bhardwaj, R. Backofen, B. Gruning, F. Ramirez and T.
Manke (2021). "pyGenomeTracks: reproducible plots for multivariate genomic datasets."
<u>Bioinformatics</u> **37**(3): 422-423.
Ma, L., Z. Chen, W. Huang da, G. Kutty, M. Ishihara, H. Wang, A. Abouelleil, L. Bishop, E. Davey,
R. Deng, X. Deng, L. Fan, G. Fantoni, M. Fitzgerald, E. Gogineni, J. M. Goldberg, G. Handley, X.
Hu, C. Huber, X. Jiao, K. Jones, J. Z. Levin, Y. Liu, P. Macdonald, A. Melnikov, C. Raley, M. Sassi, B.
T. Sherman, X. Song, S. Sykes, B. Tran, L. Walsh, Y. Xia, J. Yang, S. Young, Q. Zeng, X. Zheng, R.
Stephens, C. Nusbaum, B. W. Birren, P. Azadi, R. A. Lempicki, C. A. Cuomo and J. A. Kovacs (2016).
"Genome analysis of three Pneumocystis species reveals adaptation mechanisms to life exclusively in
mammalian hosts." <u>Nat Commun</u> **7**: 10740.
McKinley, K. L. and I. M. Cheeseman (2016). "The molecular basis for centromere identity and
function." <u>Nat Rev Mol Cell Biol</u> **17**(1): 16-29.
Miura, A., S. Yonebayashi, K. Watanabe, T. Toyama, H. Shimada and T. Kakutani (2001).
"Mobilization of transposons by a mutation abolishing full DNA methylation in Arabidopsis."
<u>Nature</u> **411**(6834): 212-214.
Mizuguchi, T., G. Fudenberg, S. Mehta, J. M. Belton, N. Taneja, H. D. Folco, P. FitzGerald, J.
Dekker, L. Mirny, J. Barrowman and S. I. S. Grewal (2014). "Cohesin-dependent globules and
heterochromatin shape 3D genome architecture in S. pombe." <u>Nature</u> **516**(7531): 432-435.
Nagaki, K., Z. Cheng, S. Ouyang, P. B. Talbert, M. Kim, K. M. Jones, S. Henikoff, C. R. Buell and J.
Jiang (2004). "Sequencing of a rice centromere uncovers active genes." <u>Nat Genet</u> **36**(2): 138-145.

Navarro-Mendoza, M. I., C. Perez-Arques, S. Panchal, F. E. Nicolas, S. J. Mondo, P. Ganguly, J.
Pangilinan, I. V. Grigoriev, J. Heitman, K. Sanyal and V. Garre (2019). "Early Diverging Fungus
Mucor circinelloides Lacks Centromeric Histone CENP-A and Displays a Mosaic of Point and
Regional Centromeres." Curr Biol **29**(22): 3791-3802 e3796.
Pidoux, A. L. and R. C. Allshire (2004). "Kinetochore and heterochromatin domains of the fission
yeast centromere." Chromosome Res **12**(6): 521-534.
Postberg, J., S. Forcob, W. J. Chang and H. J. Lipps (2010). "The evolutionary history of histone H3
suggests a deep eukaryotic root of chromatin modifying mechanisms." BMC Evol Biol **10**: 259.
Quinlan, A. R. (2014). "BEDTools: The Swiss-Army Tool for Genome Feature Analysis." Curr Protoc
Bioinformatics **47**: 11 12 11-34.
Quinlan, A. R. and I. M. Hall (2010). "BEDTools: a flexible suite of utilities for comparing genomic
features." Bioinformatics **26**(6): 841-842.
Ramirez, F., D. P. Ryan, B. Gruning, V. Bhardwaj, F. Kilpert, A. S. Richter, S. Heyne, F. Dundar and
T. Manke (2016). "deepTools2: a next generation web server for deep-sequencing data analysis."
Nucleic Acids Res **44**(W1): W160-165.
Robinson, J. T., H. Thorvaldsdottir, W. Winckler, M. Guttman, E. S. Lander, G. Getz and J. P.
Mesirov (2011). "Integrative genomics viewer." Nat Biotechnol **29**(1): 24-26.
Sabatinos, S. A. and S. L. Forsburg (2010). "Molecular genetics of Schizosaccharomyces pombe."
Methods Enzymol **470**: 759-795.
Sakuno, T. and Y. Watanabe (2009). "[Kinetochore geometry defined by cohesin within the
centromere]." Tanpakushitsu Kakusan Koso **54**(14): 1842-1849.
Sankaranarayanan, S. R., G. Ianiri, M. A. Coelho, M. H. Reza, B. C. Thimmappa, P. Ganguly, R. N.
Vadnala, S. Sun, R. Siddharthan, C. Tellgren-Roth, T. L. J. Dawson, J. Heitman and K. Sanyal (2020).
"Loss of centromere function drives karyotype evolution in closely related Malassezia species." Elife
9.
Sanyal, K., M. Baum and J. Carbon (2004). "Centromeric DNA sequences in the pathogenic yeast
Candida albicans are all different and unique." Proc Natl Acad Sci U S A **101**(31): 11374-11379.
Schotanus, K., J. L. Soyer, L. R. Connolly, J. Grandaubert, P. Happel, K. M. Smith, M. Freitag and E.
H. Stukenbrock (2015). "Histone modifications rather than the novel regional centromeres of
Zymoseptoria tritici distinguish core and accessory chromosomes." Epigenetics Chromatin **8**: 41.
Seidl, M. F., H. M. Kramer, D. E. Cook, G. L. Fiorin, G. C. M. van den Berg, L. Faino and B.
Thomma (2020). "Repetitive Elements Contribute to the Diversity and Evolution of Centromeres in
the Fungal Genus Verticillium." mBio **11**(5).
Selmecki, A., A. Forche and J. Berman (2006). "Aneuploidy and isochromosome formation in drug-
resistant Candida albicans." Science **313**(5785): 367-370.
Shen, X. X., J. L. Steenwyk, A. L. LaBella, D. A. Opulente, X. Zhou, J. Kominek, Y. Li, M.
Groenewald, C. T. Hittinger and A. Rokas (2020). "Genome-scale phylogeny and contrasting modes
of genome evolution in the fungal phylum Ascomycota." Sci Adv **6**(45).

Shiroiwa, Y., T. Hayashi, Y. Fujita, A. Villar-Briones, N. Ikai, K. Takeda, M. Ebe and M. Yanagida (2011). "Mis17 Is a Regulatory Module of the Mis6-Mal2-Sim4 Centromere Complex That Is Required for the Recruitment of CenH3/CENP-A in Fission Yeast." Plos One **6**(3).

Siepel, A., G. Bejerano, J. S. Pedersen, A. S. Hinrichs, M. Hou, K. Rosenbloom, H. Clawson, J. Spieth, L. W. Hillier, S. Richards, G. M. Weinstock, R. K. Wilson, R. A. Gibbs, W. J. Kent, W. Miller and D. Haussler (2005). "Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes." Genome Res **15**(8): 1034-1050.

Siepel, A. and D. Haussler (2004). "Phylogenetic estimation of context-dependent substitution rates by maximum likelihood." Mol Biol Evol **21**(3): 468-488.

Sievers, F., A. Wilm, D. Dineen, T. J. Gibson, K. Karplus, W. Li, R. Lopez, H. McWilliam, M. Remmert, J. Soding, J. D. Thompson and D. G. Higgins (2011). "Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega." Mol Syst Biol **7**: 539.

Slater, G. S. and E. Birney (2005). "Automated generation of heuristics for biological sequence comparison." BMC Bioinformatics **6**: 31.

Slaven, B. E., J. Meller, A. Porollo, T. Sesterhenn, A. G. Smulian and M. T. Cushion (2006). "Draft assembly and annotation of the Pneumocystis carinii genome." J Eukaryot Microbiol **53 Suppl 1**: S89-91.

Smit AFA., H. R., Green P. (2013-2015). "RepeatMasker Open-4.0.".

Smith, K. M., J. M. Galazka, P. A. Phatale, L. R. Connolly and M. Freitag (2012). "Centromeres of filamentous fungi." Chromosome Res **20**(5): 635-656.

Sridhar, S. and T. Fukagawa (2022). "Kinetochore Architecture Employs Diverse Linker Strategies Across Evolution." Front Cell Dev Biol **10**: 862637.

Takayama, Y., H. Sato, S. Saitoh, Y. Ogiyama, F. Masuda and K. Takahashi (2008). "Biphasic incorporation of centromeric histone CENP-A in fission yeast." Mol Biol Cell **19**(2): 682-690.

Thakur, J., P. B. Talbert and S. Henikoff (2015). "Inner Kinetochore Protein Interactions with Regional Centromeres of Fission Yeast." Genetics **201**(2): 543-561.

Tran, P. T., A. Paoletti and F. Chang (2004). "Imaging green fluorescent protein fusions in living fission yeast cells." Methods **33**(3): 220-225.

Vestereng, V. H., L. R. Bishop, B. Hernandez, G. Kutty, H. H. Larsen and J. A. Kovacs (2004). "Quantitative real-time polymerase chain-reaction assay allows characterization of Pneumocystis infection in immunocompetent mice." Journal of Infectious Diseases **189**(8): 1540-1544.

Williams, J. S., T. Hayashi, M. Yanagida and P. Russell (2009). "Fission yeast Scm3 mediates stable assembly of Cnp1/CENP-A into centromeric chromatin." Mol Cell **33**(3): 287-298.

Xi, Y. and W. Li (2009). "BSMAP: whole genome bisulfite sequence MAPping program." BMC Bioinformatics **10**: 232.

Yadav, V., S. Sun, R. B. Billmyre, B. C. Thimmappa, T. Shea, R. Lintner, G. Bakkeren, C. A. Cuomo, J. Heitman and K. Sanyal (2018). "RNAi is a critical determinant of centromere evolution in closely related fungi." Proc Natl Acad Sci U S A **115**(12): 3108-3113.

Yasuhara, J. C. and B. T. Wakimoto (2008). "Molecular landscape of modified histones in Drosophila heterochromatic genes and euchromatin-heterochromatin transition zones." PLoS Genet **4**(1): e16.

Zhang, Y., T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C. Nusbaum, R. M.
Myers, M. Brown, W. Li and X. S. Liu (2008). "Model-based analysis of ChIP-Seq (MACS)." <u>Genome</u>
<u>Biol</u> **9**(9): R137.
Zhao, H., A. Nguyen, D. Wu, Y. Li, S. A. Hassan, J. Chen, H. Shroff, G. Piszczek and P. Schuck
(2022). "Plasticity in structure and assembly of SARS-CoV-2 nucleocapsid protein." <u>PNAS Nexus</u>
1(2): pgac049.
Zhao, H., A. Nguyen, D. Wu, Y. Li, S. A. Hassan, J. Chen, H. Shroff, G. Piszczek and P. Schuck
(2022). "Plasticity in structure and assembly of SARS-CoV-2 nucleocapsid protein." <u>bioRxiv</u>.
1073

Figure 1. *Pneumocystis* CENP-A variants localize to the nucleus, and support viability and centromere loading in a heterologous system.

(**a**) Protein sequence alignment of full-length CENP-A^{cnp-1} homologs in *Schizosaccharomyces pombe* and seven *Pneumocystis* species. Each *Pneumocystis* species only infects a single mammalian host: *P. jirovecii* (infecting humans), *P. macacae* (macaques), *P. oryctolagi* (rabbits), *P. canis* (dogs), *P. murina* (mice), *P. carinii* (rats), and *P. wakefieldiae* (rats). The ~25 amino acid insert in *Pneumocystis* CENP-A N-terminal regions is unique to this genus.

(**b**) Immunofluorescence microscopy showing that anti-*Pneumocystis murina* CENP-A (*pn*CENP-A) reacts with *P. murina* organisms. *P. murina*-infected CD40 ligand knockout lung tissue sections were dual labeled using an anti CENP-A antibody(magenta) and 4′,6-diamidino-2-phenylindole (DAPI, light blue). The image is a maximum projection of a 5 μm thick z-stack. The images were minimally adjusted to enhance the contrast. The green box is magnified in the inset showing *pn*CENP-A inside *Pneumocystis* organisms (small dots). *pn*CENP-A signals overlap or are close in proximity with DAPI (represented by a white color produced by the overlap between magenta and light blue colors).

(**c**) Left, rescue experiment of fission yeast *cnp1-76* thermosensitive cells using the GFP-tagged CENP-A transgenes. Right, protein sequence alignment of *Pneumocystis* CENP-A N-terminal regions (*P. jirovecii* and *P. murina*) with conserved amino acids highlighted (grey areas). Plasmids containing full length and N-terminal region truncated (**) versions of *Pneumocystis* CENP-A were constructed and integrated as single copy. Growth was assayed at indicated temperatures using 10-fold serial dilutions plated on rich YEA medium. The transgene encoding *S. pombe* CENP-A serves as a non-temperature-sensitive control.

(**d**) Representative images of *cen2-tetO-TdTomato* strains expressing the indicated GFP-tagged CENP-A transgenes. On the bottom, GFP and tdTomato fluorescence as well as brightfield images are merged.

(**e**) Integrated fluorescence intensity of GFP foci was measured and plotted for the strains used in panel e. Mean (bars) and SD (error bars) are shown. n>176. AU, arbitrary units.

(**f**) Anti-GFP ChIP-qPCR analysis of indicated loci for strains used in panel e. %IP represents the percentage of input that was immunoprecipitated. Error bars denote the SD (n = 3). Abbreviations: cc1 &3, *S. pombe* centromere central core 1 & 3; cnt2:ura4⁺, ura4⁺ insertion at central core 2; *dg*, a class of heterochromatic outer repeats within pericentromeric regions (control); *fbp1*, euchromatic locus (control).

a

S. pombe	
P. jirovecii	
P. macacae	
P. oryctolagi	
P. canis	
P. murina	
P. carinii	
P. wakefieldiae	

MARAKNTARKSAGR SEG RTARKTLAGA GN . EPGDPI PRGK . RRYRPGTVALKE IR . YQKSTDLL IRKLPFARVVRD IATDYVAS FSTDFGLRWQS TA ILCLQEATEAFLVHLFEDTNLCA IHAKRVT IMQKD IQLARR IRGGWGGLS

b

*pn*CENP-A   20 μm   DAPI   20 μm   Merged   5 μm

c

26°C   32°C   35°C   **CENP-A N-terminus**

cnp1-76 (ts)

(–)	
P. jirovecii	MARAKNTARKSAGRPSDA.PRGRKR.PAGSIERMEAGDPVPRAKGR
*P. jirovecii***	MARAKNTA....................EAGDPVPRAKGR
P. murina	MARAKNTARKSAGRLSEGSTRAARKTLAGASGNIEPGDPIPRGKAR
*P. murina***	MARAKNTA....................EPGDPIPRGKAR
S. pombe	MAKKSLMA....................EPGDPIPRPRKK

d

S. pombe   *P. jirovecii*   *P. murina*

GFP-
CENP-A

GFP-
CENP-A
+
cen2-tetO-
tdTomato
+
Brightfield   10 μm

Expression in
S. pombe cnp1⁺
cen2-tetO-tdTomato

e

Fluorescence (A.U.)

Sp   Pj   Pm
GFP–CENP-A

f

ChIP-qPCR

IP %

cc1&3   *cnt2:ura4⁺*   *dg*   *fbp1*

■ *S. pombe*
■ *P. jirovecii*
■ *P. murina*

Figure 2. *Pneumocystis* CENP-A binds to single genomic foci in replicating cells

a) Study workflow and sample preparation. Four *Pneumocystis* species infecting mice, rats, macaques, and humans respectively were analyzed. *P. murina* and *P. carinii* were obtained from CD40 ligand knock-out female mice and immunosuppressed Sprague-Dawley male rats, respectively, and cultured on a co-culture of human lung adenocarcinoma cells (A549) and immortalized murine lung epithelial type 1 cells (Let-1) for 14 days. *P. macacae* organisms were obtained from a Simian Immunodeficiency Virus infected macaque (P2C), and *P. jirovecii* organisms were obtained from a single bronchoalveolar lavage from a non-infected HIV patient and three autopsy samples from patients with HIV infection. Species phylogeny and estimated speciation timing are presented. Animal icons were obtained from http://phylopic.org under creative commons licenses https://creativecommons.org/licenses/by/3.0/: mouse (Anthony Caravaggi; license CC BY-NC-SA 3.0); and rat (by Rebecca Groom; license CC BY-NC-SA 3.0).

b) *P. murina* population growth in duplicate wells were measured by quantitative PCR targeting a single copy dihydrofolate reductase (*dhfr*) gene over 14 days. Error bars represent the standard deviation (*n*=2).

c) *P. carinii* growth measurement by qPCR targeting *dhfr* gene. Error bars represent the standard deviation (*n*=2).

d) Electron micrograph showing a possibly dividing *P. murina* trophic form seven days post culture (black arrow). The field also include a non-dividing trophozoite (white arrow). 2 μm scale bar.

e) *pn*CENP-A ChIP-seq study in *P. carinii* organisms from infected lungs and cultures at days 0, 7 and 14. CENP-A remains strongly enriched at a single genomic location up to 7 days. Only one representative chromosome is presented. The remaining 16 chromosomes have similar ChIP-seq profiles and presented in supplementary figure 4.

a

Co culture on A549 and Let-1 cells for 14 days

P. murina

P. carinii

~35 Mya

~100 Mya

P. macacae

no culture

~65 Mya

P. jirovecii

no culture

~460 Mya

S. pombe

b

P. murina

10⁶

DNA copies (log DHFR)

10⁵

10⁴

0   7   14

Days Post inoculation

c

P. carinii

10⁶

DNA copies (log DHFR)

10⁵

10⁴

0   3   7   14

Days Post inoculation

d

P. murina (day 7)

e

PcCEN12

50   75   100   125   150   175   200   225   250 Kb

CENP-A (lungs)

CENP-A (day 0)

CENP-A (day 7)

CENP-A (day 14)

Figure 3. *Pneumocystis* **displays seventeen centromeres.**

a) Scaled ideogram of 17 chromosomal level scaffolds (grey) of *P. murina* genome showing CENP-A binding regions (constricted areas) and telomeric repeat motif (red).

b) CENP-A and CENP-C binding regions delineate putative centromeres (CENs) in *P. murina* genome. Color coded peaks represent enrichment of immunoprecipitated DNA (IP DNA) relative to controls (Input DNA). The 1-kb flanking sequences from the center of the enrichment peak are presented. Each scaffold displays two peaks labelled M (Major) and m (minor) according to the enrichment level.

c) Scaled ideogram of 17 chromosomal level scaffolds (grey) of *P. carinii* genome showing CENP-A binding regions (constricted areas) and telomeric repeat motif (red).

d) CENP-A binding regions delineate putative centromeres (CENs) in *P. carinii* genome. Note that no peak was observed with anti-CENP-C antibody.

e) A DNA dot plot of CENP-A binding regions in *P. murina* showing regional self-similarity. The main diagonal represents the sequence alignment with itself. Lines off the main diagonal which are repetitive patterns within the sequences are not observed. The plot shows that each CEN is unique and repeat-free.

f) DNA dot plot of CENP-A binding regions in *P. carinii* showing that each CEN is unique and repeat-free.

Figure 4. Centromeres are flanked by heterochromatin and contain active genes.

All three scaffolds are homologous.

a) Genomic view of chromosome 2 of *P. murina* genome subsequently showing annotated genes (directed grey boxes), DNA repeats (not present), percent GC content (blue), ChIP-seq read coverage distribution (BPM normalized over bins of 50 bp; input subtracted) of CENP-A, CENP-C, histones H3 and H4 ratio, heterochromatin-associated modifications (H3K9me2 and H3K9me3), euchromatin (H3K4me2) and gene expression (RNA-seq) in relation with centromeres. Genome views of remaining 16 chromosomes are presented in Supplementary dataset figure 7.

b) Genomic view of chromosome 2 of *P. carinii* genome with the same features presented as *P. murina*. A duplicated copy of *copia*-retrotransposon is presented (red arrows), which is present in syntenic regions in other *Pneumocystis* genomes. Genome views of all remaining 16 chromosomes are presented in supplementary figure 6.

c) Genomic view of chromosome 14 of *P. macacae* genome showing the distribution of gene density, repeats (DNA transposons), homology region with GC content, mis12 ChIP-seq read coverage, chromatin-associated modifications (H3K9me2 and H3K9me3 and H3K4me2) and RNA-seq in relation with centromeres. The white box indicates a DNA sequence homologous to *P. carinii* centromere 2. A single copy of the DNA transposon EnSPM (CACTA family) and the class 2DNA transposon Helitron are present upstream the centromere (forward red arrows). These elements are not present in syntenic regions in other *Pneumocystis* genomes. Genome views of all remaining 15 chromosomes are presented in Supplementary figure 5.

Figure 5. Modeling of Molecular dynamics of CENP-A nucleosomes in *P. carinii*, *P. murina* and *S. pombe.*

a) Upper panel: Sequence alignment of the centromeric histones used in modeling and simulations. Differences between *P. carinii* and *P. murina* colored green and red; differences between *S. pombe* and *pneumocystis* colored blue. Lower panel: surface representation of the centromeric octamers of *P. carinii* (left), *P. murina* (middle), and *S. pombe* (right), showing the spatial location of the mutations (snapshots from the simulations); DNA (not simulated) shown only as a reference based on the crystal structure of the human centromere nucleosome, PDB: 3AN2). The white circles show the central crevices, the behaviors of which are expected to dictate the assembly of the inner kinetochore in each species.

b) Aggregation of snapshots along the dynamic trajectory illustrating the extent of species-specific conformational variations of the central crevices (see Animation in Supplementary dataset 2); histones color code is presented at the bottom). The C-terminal tip of CENP-A is located at the center of the core (white ribbon; residues 146-151 in *Pneumocystis* and 120 in *S. pombe*; see alignment in panel a). The observed differences between *P. carinii* and *P. murina* are due to structural and dynamic changes induced by the different (H2A:H2B) dimers (cf. alignment in panel a); the differences between these species and *S. pombe* are likely the result of more extensive sequence variations in the four histones, including several solvent-exposed mutations at the core center (cf. panel a). Early-stage assembly of the inner kinetochore is thus expected to be strongly determined by features of this crevice, including its size, dynamics, and other physicochemical features.

b

P. carinii   *P. murina*   *S. pombe*

CENP-A H4 H2A H2B

Figure 6. Centromere locations, not sequences, are evolutionary constrained by negative selection.

    a) Genome organization and synteny in *Pneumocystis*. Circos plots depicting pairwise *P. carinii*, *P. murina* and *P. macacae* genome synteny. Rodent infecting Pneumocystis (*P. carinii* and *P. murina*) and macaques infecting *Pneumocystis* (*P. macacae*) have 17 and 16 chromosomal level scaffolds, respectively. Colored connectors indicate regions of synteny between species. Centromeres that overlap with recent chromosomal breakpoints are indicated (*). The square highlights *P. carinii* centromere 4 displayed in the panel b.

    b) Genome view of centromere 4 in the *P. carinii* genome. Genes are represented by directed boxes (grey for protein coding genes and cyan for polymorphic major surface glycoprotein genes), *pn*CENP-A binding region (centromere) and sequence conservation scores which were calculated from whole genome alignments of *P. carinii*. *P. murina* and *P. wakefieldiae* (PhasCons). The phastCons scores represent probabilities of negative selection and range between 0 (no conservation) and 1 (total conservation).

    c) Boxplot of conservation scores per genomic context summarized over all *P. carinii* centromeres (CEN), 30-kb regions flanking the centromeres (Cenflk), major surface glycoproteins (Msg) and random genomic background (Bckg). Statistical differences for the indicated comparisons were obtained using one-sided nonparametric Mann-Whitney test; *P*-values < .0001, ****; $P < .001$, ***; $P < .01$, **; $P < .05$, *; ns, not significant.

Figure 7. Model of regional centromere structures in *Pneumocystis* and other representative fungi.

On the left is presented the phylogeny of selected fungi inferred from maximum likelihood phylogenetic analysis of shared core protein orthologs. The overall centromere structures for *Pneumocystis carinii* (details supporting our model are provided in the text), *Schizosaccharomyces pombe* (Pidoux and Allshire 2004), *Candida albicans* (Sanyal, Baum et al. 2004), *Zymoseptoria tritici* (Schotanus, Soyer et al. 2015) and *Cryptococcus neoformans* (Janbon, Ormerod et al. 2014) are used as representative to showcase the diversity of regional centromeres in fungi. Animal pathogens are highlighted in pale olive and the plant fungal pathogen *Zymoseptoria tritici* is in pale green. For sake of brevity, only *Pneumocystis carinii* is presented for brevity, although there are differences with other *Pneumocystis* species. In the middle are presented DNA structures of centromeres. *P. carinii* has 17 centromeres (one for each of its 17 chromosomes) that share the same overall architecture. Centromeres are delineated by a localized enrichment of the centromeric histone CENP-A, which overlap with a reduction of the canonical histone H3 (inverted grey triangle). Centromeres are flanked by heterochromatin H3K9me (stands here stands for both H3K9me2 and H3K9me3). All 17 *Pneumocystis carinii* centromeres span active genes (dark grey boxes). Each centromere sequence is different and lacks shared DNA sequence motif. *S. pombe* has three centromeres that share the same overall structure in which a central core (*cnt*) domain is surrounded by innermost repeats (*imr*) and outer repeats (*otr*). The *imr* repeats incorporate clusters of transfer RNAs (tRNAs) that play a role in restricting CENP-A spread. *S. pombe* centromeres are flanked by heterochromatin (H3K9me). Genes are found 0.75-1.5 kb beyond the limits of the centromeres. *C. albicans* has 8 unique and different centromeres that are gene free and lack shared sequence motifs. *Z. tritici* has 21 centromeres ranging from 6 to 14 kb in size, that partially overlap with genes. *C. neoformans* has 14 centromeres that are gene free and enriched with Tcn transposons.