Choline supplementation in early life improves and low levels of choline can impair outcomes in a mouse model of Alzheimer's disease.

Elissavet Chartampila[1,2], Karim S. Elayouby[1,3], Paige Leary[1,4], John J. LaFrancois[1,], David Alcantara-Gonzalez[1], Swati Jain[1], Kasey Gerencer[1,5], Justin J. Botterill[1,6], Stephen D. Ginsberg[1,4,7,9], and Helen E. Scharfman[1,4,7,8,9]

[1]Center for Dementia Research
The Nathan Kline Institute for Psychiatric Research
Orangeburg, NY 10962
lafranco@nki.rfmh.org
david.alcantarag@nki.rfmh.org
swati.jain@nki.rfmh.org

[2] Current address:
Department of Cell Biology and Physiology
University of North Carolina
Chapel Hill, NC 27510
echartampila@unc.edu

[3]Current address:
Department of Neurology
Mount Sinai School of Medicine
New York, NY 10029
karim.elayouby@mssm.edu

[4]Department of Neuroscience and Physiology
New York University Grossman School of Medicine
New York, NY 100016
paige.leary@nyulangone.org

[5]Current address:
Department of Psychology
University of Maine
Orono, ME 04469
kathleen.gerencer@maine.edu

[6]Current address:
Department of Anatomy, Physiology, & Pharmacology
College of Medicine
Saskatoon, SK S7N 5E5
j.botterill@usask.ca

[7]Department of Psychiatry
New York University Grossman School of Medicine
New York, NY 10016
stephen.ginsberg@nki.rfmh.org

47 **8**Departments of Child and Adolescent Psychiatry
48 New York University Grossman School of Medicine
49 New York, NY 10016
50
51 **9**NYU Neuroscience Institute
52 New York University Grossman School of Medicine
53 New York, NY 10016
54
Address for correspondence:
Hele E. Scharfman
Center for Dementia Research
The Nathan Kline Institute for Psychiatric Research
Orangeburg, NY 10962
helen.scharfman@nki.rfmh.org

Key words: dementia, maternal choline supplementation, hippocampus, dentate
gyrus, hilus, interictal spike, novel object location, novel object recognition
64

65 **Word count:**
66 Abstract: 246
67 Introduction: 737
68 Discussion: 1730
69

70 **Number of figures:** 7
71 Number of supplemental figures: 4
72

73 **Acknowledgements:**
74 Supported by NIH R01 AG055328, MH109305 and AG077103 and the New York
75 State Office of Mental Health. We thank Dr. Christos Lisgaras for comments on
76 the manuscript and Dr. Korey Kam for contributions at an early stage of this
77 project.

78
79

ABSTRACT

Maternal choline supplementation (MCS) improves cognition in Alzheimer's disease (AD). However, the effects of MCS on neuronal hyperexcitability in AD are unknown. Therefore, we investigated the effects of MCS in a well-established mouse model of AD, the Tg2576 mouse. Like many AD mouse models and patients, Tg2576 mice exhibit hyperexcitability, typically generalized EEG spikes (interictal spikes; IIS). Hyperexcitability is also reflected by elevated expression of the transcription factor ΔFosB in the principal cells of the dentate gyrus (DG), granule cells (GCs). We also studied the hilus of the DG because hilar neurons regulate GC excitability. We found that hilar neurons reduce expression of the neuronal marker NeuN in Tg2576 mice, which other studies have shown is a sign of oxidative stress or pathology.

Tg2576 breeding pairs received a diet with a relatively low, intermediate or high concentration of choline. After weaning, all mice received the intermediate diet. In offspring of mice fed the high choline diet, IIS frequency declined, GC ΔFosB expression was reduced, and NeuN expression restored. Spatial memory improved. In contrast, offspring of mice fed the relatively low choline diet had several adverse effects, such as increased mortality, suggesting poor health. The results provide new evidence that a diet high in choline in early life can improve outcomes in a mouse model of AD, and relatively low choline can have negative consequences. This is the first study showing that dietary choline can regulate hyperexcitability, hilar neurons, and spatial memory in an animal model of AD.

I. INTRODUCTION

Diet has been suggested to influence several aspects of brain health. One dietary intervention that has been studied extensively in rodent and human populations is supplementation of the maternal diet with the nutrient choline, (maternal choline supplementation; MCS). MCS improves many aspects of brain health in humans (Zeisel & da Costa, 2009; Jiang et al., 2014). In normal rats, MCS also is beneficial, with numerous studies showing improved behavior in the adult offspring (for review, see (Meck & Williams, 2003)).

In Alzheimer's disease (AD), changes to the diet have been recommended (Bourre, 2006; Power et al., 2019; Mao et al., 2021; Lobo et al., 2022), including MCS (Strupp et al., 2016; Velazquez et al., 2020). The recommendation to use MCS is based primarily on data from the Ts65Dn mouse model of Down syndrome (DS) and AD, where MCS led to improved memory and attention in Ts65Dn offspring (Strupp et al., 2016; Powers et al., 2017; Powers et al., 2021). In addition, degeneration of basal forebrain cholinergic neurons (BFCNs) in Ts65Dn mice, a hallmark of DS and AD, was reduced (Kelley et al., 2016; Powers et al., 2017).

Based on the studies of Ts65Dn mice, we asked if MCS would improve another impairment that has been found in AD patients and mouse models of AD, hyperexcitability. We chose to study Tg2576 mice because hyperexcitability is robust, and BFCNs are likely to play a role (Kam et al., 2016). In our past work, the primary type of hyperexcitability was interictal spikes (IIS), which are named because they occur in between seizures (ictal events) in epilepsy. They are studied by EEG. We also studied hyperexcitability using ΔFosB expression, a transcription factor that is increased when neurons have been highly active over the prior 10-14 days (McClung et al., 2004). We studied ΔFosB in granule cells (GCs) of the dentate gyrus (DG), because numerous ΔFosB-expressing GCs occur when there are IIS (You et al., 2017; You et al., 2018).

We also asked if behavior would improve if MCS reduced IIS frequency in Tg2576 mice. The basis for this question is in studies of epilepsy, where IIS disrupt cognition (Rausch et al., 1978; Aarts et al., 1984; Holmes & Lenck-Santini, 2006; Kleen et al., 2010) (Kleen et al., 2013; Gelinas et al., 2016). Also, IIS in the Tg2576 mouse start early in life (by 1 month of age), before memory impairment, which occurs after 3 months of age (Kam et al., 2016). Therefore, IIS might disrupt activity sufficiently to promote behavioral impairments. We previously found that the novel object location (NOL) task is impaired at 3-4 months of age in Tg2576 mice (Duffy et al., 2015), so we evaluated NOL. We further evaluated a related task, novel object recognition (NOR; (Vogel-Ciernia & Wood, 2014). We also studied expression of NeuN, a neuronal marker that is reduced in numerous disease conditions (Buckingham et al., 2008; Kadriu et al., 2009; Matsuda et al., 2009; Duffy et al., 2011; Duffy et al., 2015). Here we report loss of NeuN-ir in the DG hilus of Tg2576 mice, which is important because impaired hilar neurons could cause GC hyperexcitability (Sperk et al., 2007; Scharfman & Myers, 2012). Therefore, we asked if MCS would restore NeuN expression.

We fed dams one of three diets, which were relatively low, intermediate or
high in choline. The high choline diet provided levels of choline similar to other
studies of MCS in rodents (Meck et al., 1988; Loy et al., 1991; Holler et al., 1996;
Meck & Williams, 1999; Sandstrom et al., 2002; Mellott et al., 2004; Glenn et al.,
2012; Kelley et al., 2019). After weaning the intermediate diet was used.
Offspring were implanted with electrodes for EEG at 1 month of age, and 24 hr-
long recordings began after 1 week for recovery, followed by 24 hr-long recording
each month. At 3 and 6 months, NOL and NOR were tested, and then we
perfusion-fixed mice, sectioned the brains, and evaluated ΔFosB and NeuN
protein expression (Figure 1A).
The results showed a remarkable effect of the high choline diet. IIS and
spatial memory were improved, as was ΔFosB and NeuN expression.
Interestingly, the low choline diet had mixed effects, reducing IIS frequency, but
increasing mortality and having other adverse outcomes, suggesting IIS were
probably reduced due to poor health. In summary, we make a strong argument
for choline supplementation in early life to improve outcomes in an AD model,
and potentially AD.
164
165　**II. METHODS**
166
A. Animals
All experimental procedures followed the guidelines set by the National
Institute of Health (NIH) and were approved by the Institutional Animal Care and
Use Committee (IACUC) at the Nathan Kline Institute for Psychiatric Research.
Tg2576 mice express a mutant form of human APP (isoform 695) with a mutation
found in a Swedish cohort (Lys670Arg, Met671Leu). The expression is driven by
the hamster prion protein promoter . Mice were bred in-house from heterozygous
Tg2576 males and non-transgenic female mice (C57BL6/SJL F1 hybrid). WT
mice were obtained from Jackson Laboratories (Stock# 100012). At weaning,
genotypes were determined using a protocol to detect APP695.
Mice were housed in standard mouse cages with corn cob bedding and a 12-
hr light-dark cycle. Food and water were provided *ad libitum*.
Breeding pairs were randomly assigned to receive one of three diets with
different concentrations of choline chloride: 1.1 g/kg (AIN-76A, Dyets), 2.0 g/kg
(Purina 5008, W.F. Fisher and Son Inc.) and 5.0 g/kg (AIN-76A, Dyets;
Supplemental Table 1). These diets ("low choline" diet, "intermediate choline" diet
and "high choline" diet, respectively) were used until weaning (25-30 days of
age). After weaning, all mice were fed the intermediate diet. Mice were housed
with others of the same sex (1-4 per cage).
186
B. Behavior
1. *General information*
Mice were housed in the room where they would be tested behaviorally
starting at least 24 hr prior to all experiments. Both NOL and NOR included three
acclimations (2-3 min each) followed by a training and a testing session (5 min
each; Figure 1B-D). The interval between training and testing was 60 minutes

and therefore tested short-term memory (Vogel-Ciernia & Wood, 2014). NOL and NOR tests were separated by 7 days to ensure effects of one task did not affect the next. The order was randomized. Prior studies have failed to find an effect on one task on the next when a 7 day-long interval is used (Botterill et al., 2021).

Video recordings of all training and testing sessions were captured using a USB camera (Logitech HD Pro C920, Logitech). Acclimations were conducted between 10:00 a.m. and 12:00 p.m. for all animals, while training and testing was conducted between 1:00 p.m. that were selected and 4:00 p.m. All equipment was cleaned using 70% ethanol before each use. In pilot studies, we chose objects that would be explored equally by animals. As reported in the Results, animals showed no preference for either of the objects (i.e., LEGO object and pineapple, Supplemental Figure 1).

2. *NOL*

The task was conducted in a standard rat cage (26 cm wide x 40 cm long x 20 cm high) with different pictures on 3 sides of the cage to provide a consistent context and therefore foster acclimation. Pictures included several shapes and colors. The dimensions of the pictures were ~5 cm x 5 cm x 5 cm. During the acclimations, animals were allowed to freely explore the cage. During the training session, mice were placed in the cage where they were acclimated, with two identical objects, 1 in a corner of the cage (e.g., left top) and the second in the adjacent corner (e.g., left bottom, Supplemental Figure 1). They were then removed and placed in their home cage for 1 hr. During the test session, one of the objects was moved to the opposite end of the cage (e.g., right top; Supplemental Figure 1). The identical objects were identical pineapple-like objects made of hard material that was painted and approximately 3 x 3 x 5 cm (Supplemental Figure 1; Botterill et al., 2021).

3. *NOR*

The task was conducted in a standard rat cage as described above. During the acclimations, animals were allowed to freely explore the cage. During the training session, mice were placed in the cage used for acclimation with two identical objects centered between the one of the cage walls (Supplemental Figure 1). The mouse was then removed and placed in their home cage for 1 hr. During the testing session, one of the objects was replaced with a different one. The identical objects were 2 pineapple-like objects and the new object was made of 2 green LEGO pieces approximately 5 x 7 x 7 cm (Supplemental Figure 1; Botterill et al., 2021).

4. *Quantification*

The experimenter who conducted the analysis was blind to the genotype of the offspring and which object was novel or familiar. Videos of the training and testing sessions were analyzed offline. Exploration was quantified based on the guidelines of Vogel-Ciernia & Wood (2014) but in addition to time spent exploring the number of approaches to an object was also quantified. Time spent exploring each object was defined as the duration of time the nose was pointed at the

object and the nose was located within 2 cm of the object. In addition, exploration
time included the time animals spent on top of the object if they were looking
down and sniffing it. An approach was defined by a movement toward the object
and ending at least 2 cm from the edge of the object.
Animals that remembered the objects explored during the training phase
were expected to demonstrate an increased preference for novelty in the test
phase (Ennaceur & Delacour, 1988). In other words, exploration of the novel
object during testing was expected to be higher than 50% of the total time of
object exploration.
248
C. Anatomy
1. *Perfusion-fixation and sectioning*
Mice were initially anesthetized by isoflurane inhalation (NDC# 07-893-1389,
Patterson Veterinary), followed by urethane (2.5 g/kg; i.p.; Cat# U2500, Sigma-
Aldrich). Under deep anesthesia, the abdominal and heart cavities were opened
and the animal was perfused transcardially using 10 mL of room temperature
(RT) saline (0.9% sodi um chloride in double distilled (dd) H_2O) using a peristaltic
pump (Minipuls1; Gilson), followed by 30 mL of 4^{o}C 4% paraformaldehyde (PFA,
Cat# 19210, Electron Microscopy Sciences) in 0.1 M phosphate buffer (PB; pH
7.4). The brains were immediately extracted, hemisected, and post-fixed for at
least 24 hr in 4% PFA (at 4^{o}C).
Following post-fixation, the left hemisphere was sectioned in the coronal
plane and the right hemisphere was sectioned in the horizontal plane (50 µm-
thick sections) using a vibratome (VT 1000P, Leica Biosystems). Sections were
collected serially and stored at 4^{o}C in 24-well tissue culture plates containing
cryoprotectant solution (30% glycerol, 30% ethylene glycol, 40% 0.1 M
phosphate buffer, pH 6.7). Throughout the hippocampus, every 6th section was
processed with an antibody to NeuN or ΔFosB.
267
2. *Immunohistochemistry- NeuN and ΔFosB*
Free floating sections were washed in 0.1 M Tris Buffer (TB; 3 x 5 min),
followed by a 3 min wash in 1% H_2O_2. They were then washed in 0.1 M TB (3 x 5
min) and incubated for 60 min in 5% horse serum for NeuN (Cat# S-2000,
Vector) or 5% goat serum for ΔFosB (Cat# S-1000, Vector), diluted in a solution
of 0.25% (volume/volume or v/v) Triton-X 100, and 1% (weight/volume or w/v)
bovine serum albumin in 0.1 M TB. They were then incubated overnight at 4^{o}C in
primary antibody to NeuN (mouse monoclonal, 1:5000; Cat# MAB377, Chemicon
International) or anti-ΔFosB (rabbit monoclonal, 1:1000; Cat# D3S8R, Cell
Signaling), diluted in a solution of 0.25% (v/v) Triton-X 100, and 1% (w/v) bovine
serum albumin in 0.1M TB. Both NeuN and ΔFosB have been well-characterized
as antigens (Mullen et al., 1992; H. K. Wolf et al., 1996; Chen et al., 1997; Sarnat
et al., 1998) and the antibodies we used have been commonly employed in the
past (Chen et al., 1997; Duffy et al., 2013; Corbett et al., 2017).
On the following day, sections were washed in 0.1 M TB (3 x 5 min) and
then incubated for 60 min in biotinylated horse anti-mouse secondary antibody
for NeuN (1:500, Cat# BA-2000, Vector) or biotinylated goat anti-rabbit

secondary antibody for ΔFosB (1:500, Cat# BA-1000), diluted in a solution of
0.25% (v/v) Triton-X 100, and 1% (w/v) bovine serum albumin in 0.1M TB. The
sections were then washed in 0.1 M TB (3 x 5 min) and incubated in avidin-biotin
complex for 2 hr (1:1000; Cat# PK-6100, Vector). They were washed in 0.1 M TB
(3 x 5 min) and then reacted in a solution containing 0.5 mg/mL 3, 3'-
diaminobenzidine (DAB; Cat# D5905, Sigma-Aldrich), 40 µg/mL ammonium
chloride (Cat# A4514, Sigma-Aldrich), 25 mg/mL D(+)-glucose (Cat# G5767,
Sigma-Aldrich), and 3 g/mL glucose oxidase (Cat# G2133, Sigma-Aldrich) in 0.1
M TB. The purpose of this method was to slow the reaction time so that the
reaction could be temporally controlled, meaning that the reaction could be
stopped when the immunoreactivity was robust but background was still low. The
sections were then washed in 0.1 M TB (3 x 5 min), mounted on gelatin-coated
slides (1% bovine gelatin, Cat# G9391, Sigma-Aldrich) and dried at RT overnight.
The following day they were dehydrated in increasing concentrations of ethanol
(90%, 10 min; 95% 10 min; 100%, 10 min; 100% again, 10 min), washed in
Xylene (10 min; Cat# 534056, Sigma-Aldrich) and cover-slipped with Permount
(Cat# 17986-01, Electron Microscopy Sciences).
302
2. *Analysis*
Photomicrographs were acquired using ImagePro Plus V7.0 (Media
Cybernetics) and a digital camera (Model RET 2000R-F-CLR-12, Q-Imaging).
NeuN and ΔFosB staining were quantified from micrographs using ImageJ V1.44
(National Institutes of Health). All images were first converted to grayscale and in
each section, the hilus was traced, defined by zone 4 of Amaral (1978). A
threshold was then calculated to identify the NeuN-stained cell bodies but not
background. Then NeuN-stained cell bodies in the hilus were quantified
manually. Note that the threshold was defined in Image J using the distribution of
intensities in the micrograph. A slider was pushed from the low level of staining to
the location where staining intensity makes a sharp rise. Below that was
considered background and above that was defined as above threshold.
To count ΔFosB-stained cells, images were converted to grayscale and in
each section, the dentate gyrus granule cell layer was outlined and defined as a
region of interest (ROI). A threshold was then set to identify the dark cells but not
the background, analogous to the was threshold was defined in the preceding
paragraph. Two thresholds were used. The first threshold was most inclusive of
immunoreactive cells even if staining was not too dark. Presumably this group
reflected cells with less neuronal activity. Then counts were made a second time
with a higher threshold. This group probably corresponded to cells with the
highest activity. ImageJ was used to calculate the area (in pixels) within the ROI
that was above threshold.
325
326 **D. Video-electroencephalographic (video-EEG) recordings**
327 **1.** *Stereotaxic surgery*
328   EEG electrodes were implanted in Tg2576 animals at 4 weeks of age. WT
329 littermates were not implanted because previous work from our laboratory has
330 shown that WT mice have no IIS (Kam et al., 2016). The animals were

anesthetized by isoflurane inhalation (3% isoflurane. 2% oxygen for induction). The Isoflurane for maintenance during surgery was $\leq$ 1.75%, The flow rate was 1.2 L/ min. Mice were placed in a stereotaxic apparatus (David Kopf Instruments). Prior to the implantation of the electrodes, animals were injected with the analgesic Buprenex (0.2 mg/kg, s.c.; buprenorphine hydroxide, NDC# 12496-0757-5, Reckitt Benckiser), which was diluted in saline (0.03 mg/mL in sterile 0.9% sodium chloride solution, Vedco Inc.). The skull was exposed with a midline incision and six holes were drilled for the placement of subdural screw electrodes (Cat# 8209, 0.10" stainless steel screws, Pinnacle Technology).

The coordinates for the electrode placement were: right occipital cortex (AP -3.5 mm, ML 2.0 mm); left frontal cortex (AP -0.5 mm, ML -1.5 mm); left hippocampus (AP -2.5 mm, ML -2.0 mm); right hippocampus (AP -2.5 mm, ML 2.0 mm; (Kam et al., 2016). Two additional screws were placed, one over the right olfactory bulb as ground (AP 2.3 mm, ML 1.8 mm) and another near the midline over the cerebellum as a reference (relative to Lambda: AP -1.5 mm, ML -0.5 mm). Screws were attached to an eight-pin connector (Cat# ED85100-ND, Digi-Key Corporation), which was placed over the skull and secured with dental cement (Cat# 4734FIB, Lang Dental Mfg. Co., Inc.).

After surgery, animals were placed on a heating blanket overnight and injected with lactated Ringer's solution (50 mL/kg at 31^{o}C; NDC# 099355000476, Aspen Veterinary Resources Ltd). They were then transferred to the room where the video-EEG is recorded. Implanted animals were allowed to recover for one week following the surgery.

2. *Video-EEG recording*

Video-EEG was recorded at 1.25 months (5 weeks) of age. The animals were then recorded at 2, 3, 4, 5 and 6 months of age. Each recording session lasted 24 hr so that a long period of sleep could be acquired. IIS occur primarily in sleep so only sleep was analyzed.

Mice were placed into 21 cm x 19 cm transparent cages (similar to standard laboratory cages) with food and water provided *ad libitum* and corncob bedding. A pre-amplifier was inserted into the eight-pin connector on the skull, which was connected to a multichannel commutator (Cat# 8408, Pinnacle Technology). This arrangement allowed for free range of movement throughout the recording. EEG signals were acquired at 2000 Hz and bandpass filtered at 0.5 – 200 Hz using Sirenia Acquisition V2.0.4 (Pinnacle Technology). Simultaneous video recordings were captured using an infrared camera (Cat# AP-DCS100W, Apex CCTV).

3. *Analysis*

EEG recordings were analyzed offline with Neuroscore V3.2.1 (Data Science International). IIS were defined as large amplitude, 10-75 msec deflections occurring synchronously in all four leads (Kam et al., 2016). They were quantified by first setting an amplitude threshold (calculated separately for each recording) and a duration criterion (10-75 msec). To determine the amplitude threshold, we calculated the root mean square (RMS) amplitude for noise in each recording during a 60 sec segment of basal activity that did not include any artifacts. The

threshold for IIS was set at 9 standard deviations above the RMS amplitude. This
threshold was selected because it identified IIS extremely well and excluded
movement or other artifacts. The IIS in one of the hippocampal channels was
used because of data suggesting IIS begin in the hippocampus (Kam et al.,
2016). Following the automatic analysis of the IIS, the software's detection
accuracy was verified manually for each recording to ensure that all spikes
appeared in all four channels and that no artifacts were included in the final IIS
count.
385
E. Statistical comparisons
Data are expressed as mean ± standard error of the mean (SEM). P-value
was set to <0.05 prior to all experiments. Parametric data comparing two groups
used unpaired two-tailed t-tests. For >2 groups, one-way ANOVA was used
followed by Tukey-Kramer post-hoc tests that corrected for multiple comparisons.
For data with two main factors, two-way ANOVA was followed by Tukey-Kramer
post-hoc tests. Interactions are not reported unless they were significant.
For the analysis of IIS frequency, a repeated measures ANOVA was not
possible because animal numbers were not the same at all ages. This was due
to mortality in low choline-treated mice or other unidentified reasons. Therefore, a
mixed effects analysis was conducted.
For non-parametric data, Mann-Whitney U tests were used to compare two
groups and Kruskal-Wallis for >2 groups. The post-hoc tests were Dunn's test for
multiple comparisons. Tests were conducted using Prism (Graphpad).
400

III. RESULTS

A. Behavior

Tg2576 mice develop impairments in synaptic plasticity and behavior very early in life, prior to Aβ deposition (Duffy et al., 2015) (Ohno et al., 2004; Jacobsen et al., 2006). In this study, we examined short-term spatial memory and recognition memory using the NOL and the NOR tasks respectively. These tasks, unlike others previously used to evaluate cognitive deficits in >9 months-old Tg2576 mouse model (i.e., Morris Water Maze, Radial Arm Water Maze, Y-Maze; (Yassine et al., 2013; A. Wolf et al., 2016)), provided an opportunity to capture cognitive deficits in animals as young as three months-old (Duffy et al., 2015).

Figure 1B-C shows a schematic of the specific experimental procedures followed for NOL and NOR.

1. *NOL*

The results of the NOL task are presented in Figure 2. Exploration of the novel object is expressed as a percentage of the total exploration of both objects. During the training session, animals did not show preference for one object over the other and this result was not dependent on genotype. Thus, a two-way ANOVA revealed no main effects of genotype on novel object exploration during training at three ($F(1,97) = 0.40$, $p = 0.53$) or six months of age ($F(1,70) = 1.94$, $p = 0.17$). There also was no effect of maternal diet on training at three ($F(2,97) = 0.06$, $p = 0.94$) or six months of age ($F(2,70) = 2.94$, $p = 0.06$).

Next, we asked it Tg2576 mice were deficient in the test phase of NOL. We used a two-way ANOVA with genotype and task phase (training versus testing) as main factors. In animals that received the intermediate diet, a two-way ANOVA revealed a significant main effect of the phase of the task (training or testing; $F(1,24) = 7.48$, $p = 0.01$). Consistent with previous studies (Duffy et al., 2015), post-hoc analyses revealed a significant increase in the exploration of the novel object during testing in WT ($p = 0.02$) but not Tg2576 ($p = 0.50$; Figure 2A). Thus, Tg2576 mice were impaired in NOL at 3 and 6 months of age when fed the intermediate diet.

In animals that received the high choline diet, a two-way ANOVA revealed a main effect of the task phase ($F(1,36) = 48.23$, $p < 0.001$). Post-hoc analyses revealed a significant increase in the exploration of the novel object during testing in both WT ($p < 0.001$) and Tg2576 ($p < 0.001$). Therefore, high choline supplementation improved memory in Tg2576 mice.

In animals treated with the low choline diet, there was no effect of phase of the task ($F(1,37) = 1.39$, $p = 0.25$). These data suggesting that low levels of choline during early life impaired spatial memory in both WT and Tg2576 mice (Figure 2A). Thus, low choline had a significant adverse effect on NOL. Adverse effects are supported by survival plots showing that there was more mortality at earlier ages in low choline-treated mice (Supplemental Figure 1).

Figure 2B shows the results in 6 months-old WT and Tg2576 mice. In animals that received the intermediate diet, a two-way ANOVA showed no main

effect of genotype (F(1,18) = 0.01, p = 0.92) or task phase (F(1,18) = 3.40, p =
0.08), revealing age-related memory deficits even in WT mice at six months of
age (Figure 2B). In contrast, the high choline group showed a main effect of task
phase (F(1,29) = 28.32, p < 0.001). Post-hoc analyses showed a significant
increase in the exploration of the novel object during testing in both WT (p <
0.001) and Tg2576 (p = 0.02). In animals that received the low choline diet, there
was no effect of task phase (F(1,25) = 3.00, p = 0.10; Figure 2B). Thus, low
choline had adverse effects on NOL in both WT and Tg2576 mice.
Taken together, these results demonstrate the lasting beneficial effects of the
high choline diet and adverse effects of low choline on a spatial memory task in
Tg2576 as well as WT littermates.
458
2. *NOR*
The results of the NOR task are presented in Figure 3. During the training
session animals did not show preference for one object over the other, exploring
both objects equally. A two-way ANOVA revealed no main effects of genotype at
three months (F(1,71) = 2.59, p = 0.54) and no effect of diet (F(2,71) = 0.22, p =
0.81). The same result was obtained at six months of age (genotype: (F(1,66) =
0.11, p = 0.75); diet: (F(2,66) = 0.98, p = 0.38). Thus, novel object exploration
during training approached 50% in all treatment groups, independent of genotype
and maternal diet, and at both ages.
Figure 3A shows the results of the testing phase of the NOR task in 3
months-old WT and Tg2576 mice. In animals that received the intermediate
choline diet, a two-way ANOVA revealed a significant main effect of task phase
(F(1,19) = 40.53, p < 0.01) and no effect of genotype (F(1,19) = 0.78, p = 0.39).
Post-hoc analyses revealed a significant increase in the exploration of the novel
object during testing compared to training for both WT (p < 0.01) and Tg2576 (p
< 0.01; Figure 3A). These results suggest that Tg2576 did not have a deficit in
NOR at 3 months of age.
Mice fed the high choline diet also showed no deficit in NOR at 3 months of
age. Thus, a two-way ANOVA revealed a main effect of task phase (F(1,26) =
17.51, p < 0.01) but not genotype (F(1,26) = 3.53, p = 0.07). Post-hoc analyses
revealed a significant increase in the exploration of the novel object during
testing in both WT (p < 0.01) and Tg2576 (p = 0.03; Figure 3A).
In animals treated with low choline, there was no impairment in NOR at 3
months of age either (task phase (F(1,26) = 22.61, p < 0.01; genotype (F(1,26) =
0.02, p = 0.89). There was a significant increase in the exploration of the novel
object in testing both in WT (p = 0.03) and Tg2576 (p < 0.01).
Figure 3B shows that there were no impairments at 6 months of age. In
animals that received the intermediate diet, a two-way ANOVA revealed a
significant main effect of task phase (F(1,19) = 13.65, p < 0.01) but no main
effect of genotype (F(1,19) = 0.28, p = 0.60). Animals explored the novel object
significantly more during testing in WT (p = 0.03) and Tg2576 mice (p = 0.04;
Figure 3B). In animals that received the other diets, results were the same. For
the high choline group, there was an effect of task phase (F(1,27) = 16.26, p <
0.01) but not genotype (F(1,27) = 0.24, p = 0.63) and a significant increase in the

exploration of the novel object during testing in both WT (p = 0.04) and Tg2576
(p < 0.01; Figure 3B). Similarly, in animals treated with the low choline diet, there
was a main effect of task phase (F(1,26) = 32.66, p < 0.01), no effect of genotype
(F(1, 6) = 0.05, p = 0.82), and more exploration of the novel object during the test
phase (WT, p < 0.01; Tg2576, p < 0.01).
Thus, at both 3 and 6 months of age, WT and Tg2576 mice performed well in
the NOR task. These results suggest that young Tg2576 mice are less sensitive
to NOR than NOL. The greater sensitivity to NOL is consistent with past
demonstrations that the dentate gyrus contributes to NOL (Sahay et al., 2011;
Kesner et al., 2015; Spyrka & Hess, 2018; Vandrey et al., 2020; Gulmez Karaca
et al., 2021; GoodSmith et al., 2022), and early signs of adverse effects in the
dentate gyrus in mouse models of AD (Jacobsen et al., 2006; Palop et al., 2011).
Our implementation of NOL may have increased the dentate gyrus-dependence
of the task by making the object locations relatively close together, because the
dentate gyrus is critical to distinguishing small differences in locations. In studies
by Pofahl and colleagues, it was shown that distances between objects like those
we used made NOL dentate gyrus-dependent (Pofahl et al., 2021).
510
511 **3.** *Exploration time*
512   Total object exploration (TOE) was measured to address any effects of
513 genotype or diet on exploration, independent of the object (Figure 4).
514
a. 3 months of age
For NOL at 3 months of age, there were effects of genotype (F(1,67) = 6.891,
p = 0.01) but not maternal diet (F(2,87) = 0.67, p = 0.63; Figure 4A). Tukey-
Kramer post-hoc tests showed that Tg2576 mice fed the low choline diet showed
more exploration than WT mice fed the same diet but the effect was on the
border of significance (p = 0.049). Three month-old mice tested with NOR
showed no significant effect of genotype (F(1,76) = 1.35, p = 0.25) or diet
(F(2,76) = 0.30, p = 0.61; Figure 4C).
523
b. 6 months of age
1) Exploration
NOL and NOR showed significant effects of diet on exploration at 6 months
of age (Figure 4B,D). For NOL, there was no effect of genotype (F(1,66) = 0.33; p
= 0.57) but there was an effect of maternal diet (F(2,66) = 8.82; p = 0.0004). For
NOR, the same was true (genotype: F(1,72) = 0.96, p = 0.33; diet: F(2,72) =
8.50, p = 0.0005). We then pooled genotypes and asked to determine which diets
differed. For NOL, mice treated with the low choline diet had less exploration
than mice fed the high choline diet (one-way ANOVA, F(2,77) = 6.90; p = 0.002;
Tukey post-hoc test, p = 0.005) and the mice that were fed the intermediate diet
also had less exploration than mice fed the high choline diet (Tukey post-hoc
test, p = 0.008; Figure 4B). Results for NOR were the same: mice treated with
the low choline diet had less exploration than mice fed the high choline diet (one-
way ANOVA, F(2,74) =4.81; p = 0.02; Tukey post-hoc test, p = 0.02) and the

mice that were fed the intermediate diet also had less exploration than mice fed
the high choline diet (Tukey post-hoc test, p = 0.03; Figure 4D).
540

2) Approaches
To gain insight into the differences in exploration, we measured total object
approaches (TOA; Figure 4A-D). TOA was defined as the number of approaches
to the familiar object + number of approaches to the novel object. There were no
differences by genotype (two-way ANOVA, $F_{(1,73)}$ = 2.930, p = 0.09) or maternal
diet ($F_{(2,73)}$ = 2.36, p = 0.10; Figure 4A-D).
Taken together, the results indicate that, in 6 months-old mice, animals that
received the high choline diet spent more time with objects at each approach.
This may underlie increased object location memory in high choline-treated mice
by increasing time for information processing during an approach. It may also
help in the NOR task with object recognition memory.
Another possibility is that as the Tg2576 mice age they show compensatory
changes that enhance memory. This has been suggested for procedural learning
(Middei et al., 2004) but not for memory of objects.
Interestingly, it has been suggested that the high choline diet reduces anxiety
(Glenn et al., 2012; Langley et al., 2015; McCall et al., 2015). Reduced anxiety
may reduce fear of exploring objects, and as a result, animals may spend more
time with objects.
For 3 months-old mice there were no effects of maternal diet on exploration,
suggesting that increased time with objects for a given approach is a change that
develops with age. To our knowledge, this age-related change has not been
previously reported. It could be a result of impairments in Tg2576 mice with age,
or a strategy to cope with impairments that are developing as AD pathology
increases.
565

B. Anatomy
1. NeuN
To further analyze the effects of dietary choline early in life, an antibody
against a neuronal nuclear antigen, NeuN. Previous studies have found that
reduced expression of NeuN often occurs in neurons after insults or injury (e.g.,
ischemia, toxicity and even aging (Lind et al., 2005; Portiansky et al., 2006;
Buckingham et al., 2008; Kadriu et al., 2009; Matsuda et al., 2009; Won et al.,
2009; Duffy et al., 2011). When NeuN is reduced after a reduction in
ARMS/Kiddins, a protein critical to neurotrophin signaling, neurons in the
entorhinal cortex demonstrate additional signs of toxicity such as pkynosis (Duffy
et al., 2011; Duffy et al., 2015). Since NeuN is reduced when Aβ levels are
elevated (Wu et al., 2016) and NeuN is reduced in AD patients (Camporez et al.,
2021), we used NeuN expression to ask if the high choline diet could rescue the
loss of NeuN-ir.
As shown in Figure 5, relatively weak NeuN-ir was observed in Tg2576
animals in the hilus, an area known to be highly vulnerable to insult and injury
(Scharfman, 1999). The most NeuN loss appeared to be in mice that received
the low choline diet and the intermediate diet compared to the high choline diet

(Figure 5B). Therefore, we conducted a two-way ANOVA with diet and
septotemporal levels as main factors. To examine the septotemporal axis,
quantification from 2-3 dorsal coronal sections were averaged to provide a value
for dorsal DG and 2-3 ventral horizontal sections were averaged for a ventral
value. There was no effect of diet ($F_{(2,30)}$ = 2.11, p = 0.14) nor septotemporal
level ($F_{(1,30)}$ = 0.02, p = 0.88) but there was a significant interaction of diet and
septotemporal level ($F_{(2,30)}$ = 3.88, p = 0.04). A one-way ANOVA using dorsal
values showed a significant effect of diet ($F_{(2,21)}$= 7.58, p = 0.003) but this was
not the case for ventral values ($F_{(2,23)}$ = 0.13; p = 0.88; Figure 5D). In the dorsal
DG, there was less NeuN-ir in mice fed the low choline diet vs the high choline
diet (Tukey-Kramer post hoc test, p = 0.03) and less NeuN in mice fed the
intermediate vs high choline diet (p = 0.003; Figure 5D). The results suggest that
choline enrichment protected hilar neurons from NeuN loss in Tg2576 mice.
597
2. ΔFosB
To complement the video-EEG recordings we used a marker of elevated
neuronal activity, ΔFosB (Figure 6). ΔFosB is a truncated variant of the
transcription factor FosB, which is increased by neuronal activity; ΔFosB has a
half-life of approximately 8 days (Ulery-Reynolds et al., 2009), so when ΔFosB is
elevated, it reflects increased neuronal activity over the last 10-14 days. Previous
studies have shown that when the J20 mouse model of AD is examined with an
antibody to ΔFosB, the GC layer shows extremely high levels of ΔFosB
expression, similar to a mouse with chronic spontaneous seizures (epilepsy;
REF). This is not surprising since the J20 mice have recurrent seizures (REF).
Therefore, we asked if Tg2576 mice would have robust ΔFosB in the GC layer,
and choline supplementation would reduce it.
There was strong expression of ΔFosB in Tg2576 GCs especially in mice fed
the low choline diet (Figure 6). Two-way ANOVA showed a main effect of
maternal diet ($F_{(2,18)}$ = 3.76, p = 0.04) but there was not a main effect of
septotemporal level ($F_{(1,18)}$ = 1.3, p = 0.26).
To clarify the effect of maternal diet, a one-way ANOVA was conducted. For
dorsal sections, there was an effect of diet ($F_{(2,9)}$ = 10.74, p = 0.004) with post-
hoc tests showing significantly less GC ΔFosB expression in high choline-treated
mice compared to low choline (p = 0.003). Low-choline-treated mice were not
significantly different from mice that were fed the intermediate diet (p = 0.08). For
the ventral DG, there was no significant effect of diet ($F_{2,9}$) = 0.79, p = 0.52;
Figure 6).
ΔFosB quantification was repeated with a lower threshold to define a ΔFosB-
ir GCs (see Methods) and results were the same (Figure 6). One-way ANOVA
showed a significant effect of diet ($F_{(2,41)}$ = 14.66, p < 0.0001), and low choline
mice had greater ΔFosB-ir than high choline mice (p =0.003). Mice fed the
intermediate diet had higher expression than mice fed the high choline diet (p =
0.0001). These data suggest that high choline in the diet early in life can reduce
hyperexcitability of GCs in offspring later in life. In addition, low choline has an
opposite effect, again suggesting this maternal diet has adverse effects.
629

C. IIS

Previous research has shown that Tg2576 mice exhibit IIS (Figure 7A) starting at a very young age (4 weeks, (Kam et al., 2016); 6 weeks, (Bezzina et al., 2015)). Therefore, we performed video-EEG using cortical (left frontal, right occipital) and hippocampal (left hippocampal, right hippocampal) electrodes in freely behaving WT and Tg2576 mice (Figure 7B-C). Animals were recorded for 24 hr each session so that there was sufficient time to capture major behavioral states (exploration, awake rest, and sleep). Consistent with previous studies (Bezzina et al., 2015; Kam et al., 2016) we observed IIS in Tg2576 mice but not WT littermates.

As shown in Figure 7B-C, animals that received the intermediate diet had a significantly higher number of IIS in the 24 hr recording periods compared to animals that received the high choline and the low choline diets. A mixed model showed that there was a significant effect or age ($F_{(2,37)} = 3.38$; $p = 0.04$) and maternal diet ($F_{(2,36)} = 8.12$; $p = 0.01$). At the 1.25-month recording, Tukey post-hoc analyses showed that IIS frequency in animals treated with the intermediate diet was higher compared to animals treated with the low ($p = 0.03$) and high choline diets ($p = 0.04$). This also was true for 2 months and 3 months (low choline, high choline, $p < 0.05$). At 4 months, the low choline groups had significantly reduced IIS frequency compared to the intermediate diet ($p = 0.009$) but not the high choline group ($p = 0.98$). At 5-6 months, IIS frequencies were not significantly different in the mice fed the maternal diets (all $p > 0.05$).

As shown previously (Kam et al., 2016), and in Figure 7, at 4-6 months of age there was high variability in IIS frequency between animals. It is possible that IIS frequency becomes more variable because we found that seizures occur in the Tg2576 mice after 2 months of age (Kam et al., 2016), and there was a sharp increase in IIS during the day before and after a seizure (Supplemental Figure 20.

D. Seizures and premature mortality in mice that had been fed the low choline diet

We found that mice that had been fed the low choline diet had greater ΔFosB-ir in GCs and the hilus showed very low NeuN-ir. Therefore, we asked whether low choline-treated mice had more seizures than mice fed the other diets. We recorded 8 mice by video-EEG for 5 days each (at 6 months of age) to examine seizures and found 2 mice from the low choline group had seizures (11 seizures over 2 of the 5 days in one mouse, 1 seizure in the other mouse) whereas none of the mice in the intermediate or high choline group had seizures (n=0/4, 2 intermediate and 2 high choline).

These values are probably an underestimate for the low choline group because many mice in this group appeared to die in a severe seizure prior to 6 months of age. Therefore, the survivors at 6 months probably were the subset with few seizures.

The reason that low choline-treated mice appeared to die in a seizure was that they were found in a specific posture in their cage which occurs when a severe seizure leads to death (Supplemental Figure 2). They were typically found

676 in a prone posture with extended, rigid limbs (Supplemental Figure 2).
677 Regardless of how the mice died, there was greater mortality in the low choline
678 group compared to mice that had been fed the high choline diet (Supplemental
679 Figure 2).
680
E. Sex differences
As shown in Figures 2-3, there appeared to be no sex differences in NOL or
NOR. To confirm, we studied NOL further, since that was the task that showed
effects of diet. A three-way ANOVA with sex, genotype and task phase as factors
showed no significant effect of sex at 3 months of age (low choline diet, $F_{(1,35)}$ =
0.0001; p = 0.99). Results were the same for the 3 months-old mice fed the
intermediate diet ($F_{(1,22)}$ = 0.062; p = 0.81) and 3 months-old mice that were fed
the high choline diet ($F_{(1,34)}$ = 0.43; p = 0.52). At 6 months of age there also
were no effects of sex (low choline diet, $F_{(1,23)}$ = 0.34; p = 0.57; intermediate
diet, $F_{(1,16)}$ = 2.49; p = 0.13; high choline diet, $F_{(1,27)}$ = 0.29; p = 0.87).
For IIS, we did not find sex differences (Supplemental Figure 4). A two-way
ANOVA with sex and diet as factors showed no effect of sex ($F_{(1,20)}$ =2.48; p =
0.13) or diet ($F_{(1,20)}$ = 3.56; p = 0.07). (Note that for this comparison we used
the low choline and high choline groups but not the intermediate group because
sex information was unavailable for these mice.) The results for months 2-3 also
showed no effect of sex (2 months: $F_{(1,17)}$ = 0.64; p = 0.43; 3 months: $F_{(1,15)}$ =
0.19; p = 0.67).
698
699 **IV. DISCUSSION**
700
Summary
This study showed that early life choline supplementation had several
beneficial effects in Tg2576 mice. The high choline diet led toz reduced IIS
frequency, improved behavior in the NOL task, and improved expression of
ΔFosB.
We showed for the first time that Tg2576 mice have pathology in the hilus at
6 months of age in that hilar neurons are deficient in immunostaining with an anti-
NeuN antibody. Tg2576 offspring on the high choline diet showed a restoration of
hilar NeuN staining. Because hilar neurons play a role in NOL (Bui et al., 2018), it
is possible that rescue of hilar neurons by the high choline diet led to improved
spatial memory. Hilar neurons have been suggested to contribute to excitability
of GCs; Scharfman, 1999, 2016), so phenotypic rescue of hilar neurons may also
have contributed to the ability of the high choline diet to reduce hyperexcitability.
We also found surprising effects of treating mice with relatively low choline in
the maternal diet. On the one hand, IIS frequency was reduced. However, these
mice showed premature mortality. Furthermore, in WT mice where no
impairments in NOL normally occur, WT mice fed the low choline diet were
impaired. In Tg2576 mice, the group that had the low maternal choline diet had
the highest ΔFosB expression and the lowest NeuN expression of any
experimental group. Reasons for these surprising effects are discussed below.
721

Benefits of high choline

The results of this study are consistent with previous reports that prenatal or postnatal choline supplementation improves object memory. One study of iron deficiency showed that choline supplementation improved NOR (Kennedy et al., 2014). Wistar rats treated with a diet high in choline had improved NOR relative to rats that had been fed a diet low in choline (Moreno et al., 2013). In rats that were aged to 24 months, animals that had been fed a high choline diet showed improved NOR, but only in females (Glenn et al., 2008).

Although few studies have examined the effects of MCS on hyperexcitability, our results showing the benefits of high choline are consistent with reports that a methyl-enriched diet (including choline but also betaine, folic acid, vitamin B12, L-methionine and zinc) reduced spike wave discharges in the offspring (Sarkisova et al., 2023) and audiogenic seizure severity (Poletaeva et al., 2014). Perinatal ethanol treatment that increased excitability was mitigated by choline chloride injection from postnatal day 10-30 (Grafe et al., 2022).

To our knowledge our study is the first to show that MCS can exert effects of IIS and ΔFosB. However, prior studies have shown that MCS in rats improved memory after severe seizures. Using a convulsant to induce several hours of severe seizures (status epilepticus, SE), there is usually extensive hippocampal neuronal loss and when tested in the subsequent days and weeks, memory impairment. MCS reduced the impairment in the Morris water maze (Yang et al., 2000; Holmes et al., 2002; Wong-Goodrich et al., 2011).

Our results are consistent with previous studies of mouse models of AD. In the APP/PS1 mouse model, it was shown that life-long choline supplementation, to the level we used (5.0 g/kg) improved memory in aged mice compared to the 1.1 g/kg diet (Velazquez et al., 2019). Postnatal choline supplementation also improved memory in APP/PS1 mice (Wang et al., 2019). Other improvements were also shown in the APP/PS1 mice, such as improved expression of choline acetyl-transferase (Mellott et al., 2017; Velazquez et al., 2019).

Adverse effects of relatively low choline

The effects of the relatively low choline diet were surprising because several adverse effects were found. This is surprising because 1.1 g/kg choline has not been considered low by others who study rats. The past studies suggest that there could be a species difference. Also, some of the past studies used the diet between day 11 and 17 of gestation, which is shorter than what we used, and therefore could have led to fewer adverse effects.

One adverse effect was weak NeuN expression in hilar neurons of mice fed the low choline diet. Also, when WT mice were tested, NOL was impaired in the low choline group but not in the mice exposed to diets with higher levels of choline.

One of the surprising findings was a high mortality and appeared to die from seizures. The surviving mice had very high ΔFosB expression in the GCs, supporting the idea they had more seizures. Indeed, the survivors did have more seizures than mice fed the other diets. These data suggest that the offspring of

mothers fed the low choline diet were unhealthy, possibly leading to or because
of seizures.
On the other hand, IIS were reduced in frequency in the same mice. There
are several possible explanations for this apparent paradox. The most
parsimonious explanation is the mice were unhealthy. If principal cells or
synapses degenerate, IIS would be reduced. Another explanation is seizures
disrupted the behavioral states when IIS occur, such as REM sleep. If REM
would be consistently disrupted, one would expect that IIS would be reduced.
Also, seizures in the Tg2576 mice were followed by a depression of the EEG that
would reduce IIS (postictal depression; Supplemental Figure 3).
A more radical explanation is that the intermediate diet promoted IIS rather
than low choline reducing IIS. The idea is attractive because it would potentially
explain why low and high choline groups had low IIS frequency compared to the
intermediate group. Thus, choline may not affect IIS very much but something
else in the diet that was very different in the intermediate group led to a
facilitation of IIS. This idea is also consistent with the data showing that the low
IIS in months 1-3 rose in months 4-6 as time when they were fed the
intermediate diet increased and there was a longer time since exposure to the
low/high choline diets.
786
NeuN
As mentioned above, NeuN is a neuronal nuclear antigen that can be
phosphorylated, and when that occurs the antibody to NeuN no longer binds to
NeuN. NeuN is phosphorylated in response to oxidative damage, brain injury,
and oxidative stress (Lind et al., 2005). Therefore, it was of interest when we saw
that hilar neurons of the dentate gyrus showed reduced NeuN in Tg2576 mice.
Hilar neurons are mainly glutamatergic mossy cells and somatostatin
(SOM)/neuropeptide Y (NPY)-expressing GABAergic neurons (HIPP cells;
(Houser, 2007; Scharfman & Myers, 2012; Scharfman, 2016), and both neuronal
types are implicated in spatial memory functions of the GCs, as well as their
excitability (Myers & Scharfman, 2009, 2011; Scharfman & Myers, 2012; Jinde et
al., 2013; Scharfman, 2016; Raza et al., 2017; Bui et al., 2018; GoodSmith et al.,
2019; Li et al., 2021). If damaged by the high intracellular APP/Aβ levels in young
Tg2576 mice, the hilar neurons would be expected to show NeuN loss.   This
idea is consistent with prior studies of the HIPP cells in AD, stained using NPY.
The prior studies showed deficits in SOM and NPY-stained cells (Chan-Palay,
1987), indicating a vulnerability.
The rescue of dorsal hilar but not ventral hilar NeuN with the high choline diet
is unclear. However, there was a trend for the high choline mice to have greater
NeuN in ventral tissue sections. Therefore, the dorsal neurons appear to have
shown a more robust change. The greater sensitivity of dorsal neurons may be
related to differences in dorsal and ventral dentate gyrus gene expression
patterns, since some of the genes that are differentially expressed could affect
vulnerability to insult or injury (Cembrowski et al., 2016; Zhang et al., 2018). It is
possible that dorsal hilar neurons are inherently more resilient to insults,
especially in the context of AD. Resilience of vulnerable neuronal populations,

especially in the hippocampus, is a high interest topic across the AD spectrum,
and in relevant AD models, including the Tg2576 mouse.
815
ΔFosB
Given that seizures are associated with elevated GC expression of ΔFosB
(Chen et al., 1997; McClung et al., 2004; Corbett et al., 2017; You et al., 2017),
the differences in ΔFosB expression levels observed here could be the result of
increased seizures in animals that received the low choline diet. Therefore we
suggest that the high ΔFosB in Tg2576 mice fed the low choline diet reflects
more seizures than in the other two diets.
ΔFosB is a transcription factor that is linked to cognition. In the J20 mouse
model of AD, elevated ΔFosB in GCs led to reduced cognition, and when ΔFosB
was selectively reduced the cognition improved. Therefore, the reduction in
ΔFosB by the high choline diet was important to show hyperexcitability was
reduced and also important because it showed how high choline may benefit
cognition.
829
Additional considerations
Although the low and high choline diets only had differences in choline, the
intermediate diet had different amounts of choline as well as other constituents
(Supplementary Table 1). The intermediate diet was useful however, because
numerous studies in AD mouse models employ this standard diet.
There was high variability in the IIS data in the mice fed the intermediate diet
after 3 months of age, and this may have led to an underestimation of effects.
This variability was observed before and it may be due to the emergence of
aspects of AD pathophysiology that emerge after 3 months of age (Kam et al.,
2016).
Because IIS occur primarily in sleep, it is possible that the effects of choline
on IIS were indirectly due to their effects on sleep. For example, if the high
choline diet led to reduced sleep, it would reduce IIS. However, there is no
evidence that MCS influences sleep in humans or laboratory animals. The one
study that looked at the sleep/wake cycle in rats did not find effects (Cheng et al.,
2008).
Although we failed to detect sex differences, there may actually be
differences if females had been separated by stage of the estrous cycle at death.
This possibility is raised by prior data showing that rats and mice on diestrus
have low excitability in hippocampal slices. At other cycle stages hyperexcitability
occurs.
Limitations of the Tg2576 model should be considered because the mice do
not recapitulate all characteristics of AD, such as tau pathology. Also, the
Swedish mutation does not represent the majority of AD, which is sporadic.
However, MCS has now demonstrated structural/functional benefits in several
AD-relevant models, namely Tg2576 (this report), APP-PS1 (Mellott et al., 2017),
and Ts65Dn (Strupp, 2016; Powers 2017, 2021; Alldred 2021, 2023).
857

Figure 1. Schematic of the behavioral procedures for the NOL and NOR tasks.

A. Experimental timeline. Dams were fed one of three diets for mating, gestation, and offspring consumed the diet until weaning at 25-30 days of age. At 1 month-old, offspring were implanted with electrodes for EEG and recorded for a 24 hr period at 1.2, 2, 3, 4, 5, and 6 months of age. At 3 and 6 months of age behavior was tested. At 6 months of age, after behavioral testing, mice were perfused, brains were sectioned, and sections were processed with antibodies against NeuN and ΔFosB.

B. Prior to NOL or NOR, animals were acclimated to the testing arena. There were three acclimation sessions separated by 24hr during which animals were allowed to freely explore for 5 min.

C. In the NOL task, animals were placed in a cage with two identical objects and allowed to freely explore for 5 min (Training). After one hour they were brought back to the cage, where one object was displaced, and allowed to freely explore for 5 min (Testing).

D. In the NOR task. animals were placed in a cage with two identical objects and allowed to freely explore for 5 min (Training). After one hour they were brought back to the cage, where one object was replaced, and allowed to freely explore for 5 min (Testing).

Figure 2. Choline enrichment reduced spatial memory deficits.
A. Three months-old mice.
1. WT and Tg2576 mice exposed to the low choline diet showed spatial memory deficits.
2. Tg2576 exposed to the intermediate diet showed spatial memory deficits but not WT.
3. Spatial memory was improved in animals exposed to the high choline diet.
B. Six months-old mice.
1. WT and Tg2576 mice exposed to the low choline diet showed spatial memory deficits.
2. WT and Tg2576 mice exposed to the intermediate diet showed spatial memory deficits.
3. WT and Tg2576 mice exposed to the high choline diet had improved spatial memory.

A. Novel Object Location: 3 months-old

1. Low choline   2. Intermediate choline   3. High choline

B. Novel Object Location: 6 months-old

1. Low choline   2. Intermediate choline   3. High choline

Figure 3. WT and Tg2576 mice showed recognition memory regardless of diet.
A. Three months-old mice.
1-3. WT and Tg2576 mice treated performed the NOR task.
B. Six months-old mice.
1-3. WT and Tg2576 mice performed the NOR task.

Figure 4. Tg2576 mice treated with the choline supplemented diet spent
more time with the objects at each approach.
A-B. NOL.
There was no effect of diet on time exploring or approaches to the novel object in
the NOL task at 3 months of age (A) but there were significant differences at 6
months of age (B). Mice exposed to the high choline diet spent more time
exploring than mice that had been fed the low choline or the intermediate diet.
C-D. NOR.
There was no effect of diet on time exploring or approaches to the novel object in
the NOR task at 3 months of age (C) but there were significant differences at 6
months of age (D). Mice that had been fed the high choline diet spent more time
exploring than mice that had been fed the low choline or the intermediate diet.

917
918

Figure 5. Choline supplementation improves NeuN immunoreactivity (ir) in hilar cells in Tg2576 animals

A. A sagittal view of the brain illustrating the coronal sectioning plane.

B. Representative images of NeuN-ir staining in the dorsal dentate gyrus of Tg2576 animals from each treatment group.

C. Quantification methods. Representative images demonstrating the thresholding criteria established to quantify NeuN-ir.

1. A NeuN-stained section shows an inset (arrow) expanded at the top left to show 3 hilar cells. NeuN-ir cells are marked by blue arrows. NeuN-ir that was too close to background staining to be considered positive is marked by a green arrow.

2. After converting the image to grayscale, a threshold was set so that all cells stained well with NeuN antibody were red. The inset shows that the two cells that were darkest met the threshold while the weakly stained cell did not.

D. Six months-old Tg2576 mice treated with the choline supplemented diet had significantly more NeuN-ir cells in the hilus in the dorsal dentate gyrus (1) compared to other treatment groups but not ventral dentate gyrus (2).

Figure 6. Choline supplementation reduced ΔFosB expression in dorsal GCs.

A. A sagittal view of the brain illustrating the coronal sectioning plane.

B. Representative images of ΔFosB staining in the anterior dentate gyrus of Tg2576 animals from each treatment group.

C. Quantification methods. Representative images demonstrating the thresholding criteria established to quantify ΔFosB.

1. A ΔFosB -stained section shows strongly-stained cells (white arrows).

2. A strict thresholding criteria was used to make only the darkest stained cells red.

D. Six months-old Tg2576 mice treated with the choline supplemented diet had significantly more ΔFosB -ir pixels above threshold in the dorsal dentate gyrus (1) compared to other treatment groups but not ventral dentate gyrus (2).

E. Methods are shown using a threshold that was less strict.

1. Some of the stained cells that were included are not as dark as those used for the strict threshold (white arrows).

2. All cells above the less conservative threshold are shown in red.

F. With the less conservative threshold, the results were the same as D. Mice that were fed the high choline diet had less ΔFosB-ir pixels than the mice that were fed the other diets. However, this was true for dorsal (1) but not ventral (2) sections.

Figure 7. The high choline diet reduced IIS frequency in Tg2576 animals.

A. 1. Representative example of an IIS (red arrow). IIS were defined as occurring in all four channels (generalized) to distinguish them from focal spikes that can often be artifact.

2-3. The IIS shown in A1 is expanded.

B. 1. Scatter plot of IIS frequency at each age of recording.

2. Means and sem are plotted. The high choline diet group had fewer IIS than the intermediate diet group at ages 1-3 months ($, p<0.05) and the low choline group had less IIS than the intermediate diet at ages 1-4 months (@, p<0.05). The high and low choline diet groups were significantly different at ages 3 and 4 months (*, p<0.05).

976 **Supplemental Table 1. Comparison of the three diets used in this study.**

	Low choline	Intermediate choline	High choline
Choline	1.1g/kg	2.0g/kg	5.0g/kg
Carbohydrate	690	570	690
Protein	177	236	177
Fat	120	170	120
Thiamin	0.020	0.015	0.020
Riboflavin	0.600	0.005	0.600
Pyridoxine	0.700	0.006	0.700
Niacin	3.000	0.780	3.000
Folic Acid	0.200	0.003	0.200
Biotin	0.020	0.002	0.020

Supplemental Figure 1. Details of behavioral tasks.
A standard mouse cage is diagrammed with the locations of objects inside it. For
NOL, there were two pineapple-like objects and one was moved across the cage
for the test session. For the NOR test session, one pineapple-like object was
removed and an object made of LEGO was placed in the location where the
original pineapple-like object had been.

986
987

Supplemental Figure 2. Mortality was high in ice treated with the low
choline diet.
A. Survival curves are shown for mice fed the low choline diet and mice fed the
high choline diet.
B. A photo of a mouse after sudden unexplained death. The mouse was found in
a posture consistent with death during a convulsive seizure.
994
995

Supplemental Figure 3. IIS frequency before and after seizures.
A. Representative EEG trace of a seizure in a 5 months-old Tg2576 mouse.
B. IIS frequency was quantified from continuous video-EEG for 4 days. During
the first 24 hr, IIS frequency was low. The next day a seizure occurred (arrow),
and IIS frequency increased approximately 5 fold, and for the next 24 hr period.
On day 4, IIS frequency returned to lower levels.

A. Seizure

1002
1003
1004

1005 **Supplemental Figure 4. IIS frequency was similar for each sex.**
1006 **A.** At 1.2 months-old, female and male Tg2576 mice were not significantly
1007 different in IIS frequency for the low and high choline diets.
1008 **B-C.** There were no significant sex differences at 2 (B) and 3 (C) months of age.
1009

REFERENCES

Aarts JH, Binnie CD, Smit AM, & Wilkins AJ (1984) Selective cognitive impairment during focal and generalized epileptiform EEG activity. Brain. 107 (Pt 1): 293-308.

Bezzina C, Verret L, Juan C, Remaud J, Halley H, Rampon C, & Dahan L (2015) Early onset of hypersynchronous network activity and expression of a marker of chronic seizures in the Tg2576 mouse model of Alzheimer's disease. PLoS One. 10: e0119910.

Botterill JJ, Vinod KY, Gerencer KJ, Teixeira CM, LaFrancois JJ, & Scharfman HE (2021) Bidirectional regulation of cognitive and anxiety-like behaviors by dentate gyrus mossy cells in male and female mice. J Neurosci. 41: 2475-2495.

Bourre JM (2006) Effects of nutrients (in food) on the structure and function of the nervous system: Update on dietary requirements for brain. Part 2 : Macronutrients. J Nutr Health Aging. 10: 386-399.

Buckingham BP, Inman DM, Lambert W, Oglesby E, Calkins DJ, Steele MR, Vetter ML, Marsh-Armstrong N, & Horner PJ (2008) Progressive ganglion cell degeneration precedes neuronal loss in a mouse model of glaucoma. J Neurosci. 28: 2735-2744.

Bui AD, Nguyen TM, Limouse C, Kim HK, Szabo GG, Felong S, Maroso M, & Soltesz I (2018) Dentate gyrus mossy cells control spontaneous convulsive seizures and spatial memory. Science. 359: 787-790.

Camporez D, Belcavello L, Almeida JFF, Silva-Sena GG, Pimassoni LHS, Morelato RL, do Carmo Pimentel Batitucci M, & de Paula F (2021) Positive association of a sirt1 variant and parameters of oxidative stress on Alzheimer's disease. Neurol Sci. 42: 1843-1851.

Cembrowski MS, Wang L, Sugino K, Shields BC, & Spruston N (2016) Hipposeq: A comprehensive RNA-seq database of gene expression in hippocampal principal neurons. Elife. 5: e14997.

Chan-Palay V (1987) Somatostatin immunoreactive neurons in the human hippocampus and cortex shown by immunogold/silver intensification on vibratome sections: Coexistence with neuropeptide y neurons, and effects in Alzheimer-type dementia. J Comp Neurol. 260: 201-223.

Chen J, Kelz MB, Hope BT, Nakabeppu Y, & Nestler EJ (1997) Chronic Fos-related antigens: Stable variants of ΔFosb induced in brain by chronic treatments. J Neurosci. 17: 4933-4941.

Cheng RK, Williams CL, & Meck WH (2008) Oscillatory bands, neuronal synchrony and hippocampal function: Implications of the effects of prenatal choline supplementation for sleep-dependent memory consolidation. Brain Res. 1237: 176-194.

Corbett BF, You JC, Zhang X, Pyfer MS, Tosi U, Iascone DM, Petrof I, Hazra A, Fu CH, Stephens GS, Ashok AA, Aschmies S, Zhao L, Nestler EJ, & Chin J (2017) ΔFosB regulates gene expression and cognitive dysfunction in a mouse model of Alzheimer's disease. Cell Rep. 20: 344-355.

Duffy AM, Morales-Corraliza J, Bermudez-Hernandez KM, Schaner MJ, Magagna-Poveda A, Mathews PM, & Scharfman HE (2015) Entorhinal cortical

defects in Tg2576 mice are present as early as 2-4 months of age. Neurobiol
Aging. 36: 134-148.
Duffy AM, Schaner MJ, Chin J, & Scharfman HE (2013) Expression of c-fos in
hilar mossy cells of the dentate gyrus in vivo. Hippocampus. 23: 649-655.
Duffy AM, Schaner MJ, Wu SH, Staniszewski A, Kumar A, Arevalo JC, Arancio
O, Chao MV, & Scharfman HE (2011) A selective role for ARMS/kidins220
scaffold protein in spatial memory and trophic support of entorhinal and frontal
cortical neurons. Exp Neurol. 229: 409-420.
Ennaceur A, & Delacour J (1988) A new one-trial test for neurobiological studies
of memory in rats. 1: Behavioral data. Behav Brain Res. 31: 47-59.
Gelinas JN, Khodagholy D, Thesen T, Devinsky O, & Buzsaki G (2016) Interictal
epileptiform discharges induce hippocampal-cortical coupling in temporal lobe
epilepsy. Nat Med. 22: 641-648.
Glenn MJ, Kirby ED, Gibson EM, Wong-Goodrich SJ, Mellott TJ, Blusztajn JK, &
Williams CL (2008) Age-related declines in exploratory behavior and markers
of hippocampal plasticity are attenuated by prenatal choline supplementation
in rats. Brain Res. 1237: 110-123.
Glenn MJ, Adams RS, & McClurg L (2012) Supplemental dietary choline during
development exerts antidepressant-like effects in adult female rats. Brain Res.
1443: 52-63.
GoodSmith D, Lee H, Neunuebel JP, Song H, & Knierim JJ (2019) Dentate gyrus
mossy cells share a role in pattern separation with dentate granule cells and
proximal CA3 pyramidal cells. J Neurosci. 39: 9570-9584.
GoodSmith D, Kim SH, Puliyadi V, Ming GL, Song H, Knierim JJ, & Christian KM
(2022) Flexible encoding of objects and space in single cells of the dentate
gyrus. Curr Biol. 32: 1088-1101 e1085.
Grafe EL, Wade MMM, Hodson CE, Thomas JD, & Christie BR (2022) Postnatal
choline supplementation rescues deficits in synaptic plasticity following
prenatal ethanol exposure. Nutrients. 14.
Gulmez Karaca K, Brito DVC, Kupke J, Zeuch B, & Oliveira AMM (2021) Engram
reactivation during memory retrieval predicts long-term memory performance
in aged mice. Neurobiol Aging. 101: 256-261.
Hauser WA (1992) Seizure disorders: The changes with age. Epilepsia. 33 Suppl
4: S6-14.
Holler T, Cermak JM, & Blusztajn JK (1996) Dietary choline supplementation in
pregnant rats increases hippocampal phospholipase d activity of the offspring.
FASEB J. 10: 1653-1659.
Holmes GL, & Lenck-Santini PP (2006) Role of interictal epileptiform
abnormalities in cognitive impairment. Epilepsy Behav. 8: 504-515.
Holmes GL, Yang Y, Liu Z, Cermak JM, Sarkisian MR, Stafstrom CE, Neill JC, &
Blusztajn JK (2002) Seizure-induced memory impairment is reduced by
choline supplementation before or after status epilepticus. Epilepsy Res. 48: 3-
13.
Houser CR (2007) Interneurons of the dentate gyrus: An overview of cell types,
terminal fields and neurochemical identity. Prog Brain Res. 163: 217-232.

Jacobsen JS, Wu CC, Redwine JM, Comery TA, Arias R, Bowlby M, Martone R, Morrison JH, Pangalos MN, Reinhart PH, & Bloom FE (2006) Early-onset behavioral and synaptic deficits in a mouse model of Alzheimer's disease. Proc Natl Acad Sci U S A. 103: 5161-5166.

Jiang X, West AA, & Caudill MA (2014) Maternal choline supplementation: A nutritional approach for improving offspring health? Trends Endocrinol Metab. 25: 263-273.

Jinde S, Zsiros V, & Nakazawa K (2013) Hilar mossy cell circuitry controlling dentate granule cell excitability. Front Neural Circuits. 7: 14.

Kadriu B, Guidotti A, Costa E, & Auta J (2009) Imidazenil, a non-sedating anticonvulsant benzodiazepine, is more potent than diazepam in protecting against dfp-induced seizures and neuronal damage. Toxicology. 256: 164-174.

Kam K, Duffy AM, Moretto J, LaFrancois JJ, & Scharfman HE (2016) Interictal spikes during sleep are an early defect in the Tg2576 mouse model of beta-amyloid neuropathology. Sci Rep. 6: 20119.

Kelley CM, Ash JA, Powers BE, Velazquez R, Alldred MJ, Ikonomovic MD, Ginsberg SD, Strupp BJ, & Mufson EJ (2016) Effects of maternal choline supplementation on the septohippocampal cholinergic system in the Ts65Dn mouse model of Down syndrome. Curr Alzheimer Res. 13: 84-96.

Kelley CM, Ginsberg SD, Alldred MJ, Strupp BJ, & Mufson EJ (2019) Maternal choline supplementation alters basal forebrain cholinergic neuron gene expression in the Ts65Dn mouse model of Down syndrome. Dev Neurobiol. 79: 664-683.

Kennedy BC, Dimova JG, Siddappa AJ, Tran PV, Gewirtz JC, & Georgieff MK (2014) Prenatal choline supplementation ameliorates the long-term neurobehavioral effects of fetal-neonatal iron deficiency in rats. J Nutr. 144: 1858-1865.

Kesner RP, Taylor JO, Hoge J, & Andy F (2015) Role of the dentate gyrus in mediating object-spatial configuration recognition. Neurobiol Learn Mem. 118: 42-48.

Kleen JK, Scott RC, Holmes GL, Roberts DW, Rundle MM, Testorf M, Lenck-Santini PP, & Jobst BC (2013) Hippocampal interictal epileptiform activity disrupts cognition in humans. Neurology. 81: 18-24.

Kleen JK, Scott RC, Holmes GL, & Lenck-Santini PP (2010) Hippocampal interictal spikes disrupt cognition in rats. Ann Neurol. 67: 250-257.

Langley EA, Krykbaeva M, Blusztajn JK, & Mellott TJ (2015) High maternal choline consumption during pregnancy and nursing alleviates deficits in social interaction and improves anxiety-like behaviors in the BTBR t+itpr3tf/j mouse model of autism. Behav Brain Res. 278: 210-220.

Li X, Chen W, Yu Q, Zhang Q, Zhang T, Huang X, Li H, He A, Yu H, Jing W, Du H, Ke X, Zhang B, Tian Q, Liu R, & Lu Y (2021) A circuit of mossy cells controls the efficacy of memory retrieval by Gria2i inhibition of Gria2. Cell Rep. 34: 108741.

Lind D, Franken S, Kappler J, Jankowski J, & Schilling K (2005) Characterization of the neuronal marker neun as a multiply phosphorylated antigen with discrete subcellular localization. J Neurosci Res. 79: 295-302.

Lobo F, Haase J, & Brandhorst S (2022) The effects of dietary interventions on brain aging and neurological diseases. Nutrients. 14.

Loy R, Heyer D, Williams CL, & Meck WH (1991) Choline-induced spatial memory facilitation correlates with altered distribution and morphology of septal neurons. Adv Exp Med Biol. 295: 373-382.

Mao XY, Yin XX, Guan QW, Xia QX, Yang N, Zhou HH, Liu ZQ, & Jin WL (2021) Dietary nutrition for neurological disease therapy: Current status and future directions. Pharmacol Ther. 226: 107861.

Matsuda S, Umeda M, Uchida H, Kato H, & Araki T (2009) Alterations of oxidative stress markers and apoptosis markers in the striatum after transient focal cerebral ischemia in rats. J Neural Transm (Vienna). 116: 395-404.

McCall N, Mahadevia D, Corriveau JA, & Glenn MJ (2015) Adult emotionality and neural plasticity as a function of adolescent nutrient supplementation in male rats. Pharmacol Biochem Behav. 132: 125-135.

McClung CA, Ulery PG, Perrotti LI, Zachariou V, Berton O, & Nestler EJ (2004) ΔFosB a molecular switch for long-term adaptation in the brain. Mol Brain Res. 132: 146-154.

Meck WH, & Williams CL (1999) Choline supplementation during prenatal development reduces proactive interference in spatial memory. Dev Brain Res. 118: 51-59.

Meck WH, & Williams CL (2003) Metabolic imprinting of choline by its availability during gestation: Implications for memory and attentional processing across the lifespan. Neurosci Biobehav Rev. 27: 385-399.

Meck WH, Smith RA, & Williams CL (1988) Pre- and postnatal choline supplementation produces long-term facilitation of spatial memory. Dev Psychobiol. 21: 339-353.

Mellott TJ, Huleatt OM, Shade BN, Pender SM, Liu YB, Slack BE, & Blusztajn JK (2017) Perinatal choline supplementation reduces amyloidosis and increases choline acetyltransferase expression in the hippocampus of the appsweps1de9 Alzheimer's disease model mice. PLoS One. 12: e0170450.

Mellott TJ, Williams CL, Meck WH, & Blusztajn JK (2004) Prenatal choline supplementation advances hippocampal development and enhances mapk and creb activation. FASEB J. 18: 545-547.

Middei S, Geracitano R, Caprioli A, Mercuri N, & Ammassari-Teule M (2004) Preserved fronto-striatal plasticity and enhanced procedural learning in a transgenic mouse model of Alzheimer's disease overexpressing mutant happswe. Learn Mem. 11: 447-452.

Moreno HC, de Brugada I, Carias D, & Gallo M (2013) Long-lasting effects of prenatal dietary choline availability on object recognition memory ability in adult rats. Nutr Neurosci. 16: 269-274.

Mullen RJ, Buck CR, & Smith AM (1992) Neun, a neuronal specific nuclear protein in vertebrates. Development. 116: 201-211.

Myers CE, & Scharfman HE (2011) Pattern separation in the dentate gyrus: A role for the CA3 backprojection. Hippocampus. 21: 1190-1215.

Myers CE, & Scharfman HE (2009) A role for hilar cells in pattern separation in the dentate gyrus: A computational approach. Hippocampus. 19: 321-337.

Ohno M, Sametsky EA, Younkin LH, Oakley H, Younkin SG, Citron M, Vassar R,
& Disterhoft JF (2004) BACE1 deficiency rescues memory deficits and
cholinergic dysfunction in a mouse model of Alzheimer's disease. Neuron. 41:
27-33.
Olafsson E, Ludvigsson P, Gudmundsson G, Hesdorffer D, Kjartansson O, &
Hauser WA (2005) Incidence of unprovoked seizures and epilepsy in iceland
and assessment of the epilepsy syndrome classification: A prospective study.
Lancet Neurol. 4: 627-634.
Palop JJ, Mucke L, & Roberson ED (2011) Quantifying biomarkers of cognitive
dysfunction and neuronal network hyperexcitability in mouse models of
Alzheimer's disease: Depletion of calcium-dependent proteins and inhibitory
hippocampal remodeling. Methods Mol Biol. 670: 245-262.
Pofahl M, Nikbakht N, Haubrich AN, Nguyen T, Masala N, Distler F, Braganza O,
Macke JH, Ewell LA, Golcuk K, & Beck H (2021) Synchronous activity patterns
in the dentate gyrus during immobility. Elife. 10.
Poletaeva, II, Surina NM, Ashapkin VV, Fedotova IB, Merzalov IB, Perepelkina
OV, & Pavlova GV (2014) Maternal methyl-enriched diet in rat reduced the
audiogenic seizure proneness in progeny. Pharmacol Biochem Behav. 127:
21-26.
Portiansky EL, Barbeito CG, Gimeno EJ, Zuccolilli GO, & Goya RG (2006) Loss
of neun immunoreactivity in rat spinal cord neurons during aging. Exp Neurol.
202: 519-521.
Power R, Prado-Cabrero A, Mulcahy R, Howard A, & Nolan JM (2019) The role
of nutrition for the aging population: Implications for cognition and Alzheimer's
disease. Annu Rev Food Sci Technol. 10: 619-639.
Powers BE, Velazquez R, Strawderman MS, Ginsberg SD, Mufson EJ, & Strupp
BJ (2021) Maternal choline supplementation as a potential therapy for Down
syndrome: Assessment of effects throughout the lifespan. Front Aging
Neurosci. 13: 723046.
Powers BE, Kelley CM, Velazquez R, Ash JA, Strawderman MS, Alldred MJ,
Ginsberg SD, Mufson EJ, & Strupp BJ (2017) Maternal choline
supplementation in a mouse model of Down syndrome: Effects on attention
and nucleus basalis/substantia innominata neuron morphology in adult
offspring. Neuroscience. 340: 501-514.
Rausch R, Lieb JP, & Crandall PH (1978) Neuropsychologic correlates of depth
spike activity in epileptic patients. Arch Neurol. 35: 699-705.
Raza SA, Albrecht A, Caliskan G, Muller B, Demiray YE, Ludewig S, Meis S,
Faber N, Hartig R, Schraven B, Lessmann V, Schwegler H, & Stork O (2017)
HIPP neurons in the dentate gyrus mediate the cholinergic modulation of
background context memory salience. Nat Commun. 8: 189.
Sahay A, Scobie KN, Hill AS, O'Carroll CM, Kheirbek MA, Burghardt NS, Fenton
AA, Dranovsky A, & Hen R (2011) Increasing adult hippocampal neurogenesis
is sufficient to improve pattern separation. Nature. 472: 466-470.
Sandstrom NJ, Loy R, & Williams CL (2002) Prenatal choline supplementation
increases NGF levels in the hippocampus and frontal cortex of young and
adult rats. Brain Res. 947: 9-16.

Sarkisova KY, Fedosova EA, Shatskova AB, Rudenok MM, Stanishevskaya VA, & Slominsky PA (2023) Maternal methyl-enriched diet increases dnmt1, hcn1, and th gene expression and suppresses absence seizures and comorbid depression in offspring of WAG/Rij rats. Diagnostics (Basel). 13.

Sarnat HB, Nochlin D, & Born DE (1998) Neuronal nuclear antigen (neun): A marker of neuronal maturation in early human fetal nervous system. Brain Dev. 20: 88-94.

Scharfman HE (2016) The enigmatic mossy cell of the dentate gyrus. Nat Rev Neurosci. 17: 562-575.

Scharfman HE, & Myers CE (2012) Hilar mossy cells of the dentate gyrus: A historical perspective. Front Neural Circuits. 6: 106.

Scharfman HE (1999) The role of nonprincipal cells in dentate gyrus excitability and its relevance to animal models of epilepsy and temporal lobe epilepsy. Adv Neurol. 79: 805-820.

Sperk G, Hamilton T, & Colmers WF (2007) Neuropeptide Y in the dentate gyrus. Prog Brain Res. 163: 285-297.

Spyrka J, & Hess G (2018) Repeated neck restraint stress bidirectionally modulates excitatory transmission in the dentate gyrus and performance in a hippocampus-dependent memory task. Neuroscience. 379: 32-44.

Strupp BJ, Powers BE, Velazquez R, Ash JA, Kelley CM, Alldred MJ, Strawderman M, Caudill MA, Mufson EJ, & Ginsberg SD (2016) Maternal choline supplementation: A potential prenatal treatment for Down syndrome and Alzheimer's disease. Curr Alzheimer Res. 13: 97-106.

Vandrey B, Garden DLF, Ambrozova V, McClure C, Nolan MF, & Ainge JA (2020) Fan cells in layer II of the lateral entorhinal cortex are critical for episodic-like memory. Curr Biol. 30: 169-175 e165.

Velazquez R, Ferreira E, Knowles S, Fux C, Rodin A, Winslow W, & Oddo S (2019) Lifelong choline supplementation ameliorates Alzheimer's disease pathology and associated cognitive deficits by attenuating microglia activation. Aging Cell. 18: e13037.

Velazquez R, Ferreira E, Winslow W, Dave N, Piras IS, Naymik M, Huentelman MJ, Tran A, Caccamo A, & Oddo S (2020) Maternal choline supplementation ameliorates Alzheimer's disease pathology by reducing brain homocysteine levels across multiple generations. Mol Psychiatry. 25: 2620-2629.

Vogel-Ciernia A, & Wood MA (2014) Examining object location and object recognition memory in mice. Curr Protoc Neurosci. 69: 8 31 31-17.

Wang Y, Guan X, Chen X, Cai Y, Ma Y, Ma J, Zhang Q, Dai L, Fan X, & Bai Y (2019) Choline supplementation ameliorates behavioral deficits and Alzheimer's disease-like pathology in transgenic APP/PS1 mice. Mol Nutr Food Res. 63: e1801407.

Wolf A, Bauer B, Abner EL, Ashkenazy-Frolinger T, & Hartz AM (2016) A comprehensive behavioral test battery to assess learning and memory in 129s6/Tg2576 mice. PLoS One. 11: e0147733.

Wolf HK, Buslei R, Schmidt-Kastner R, Schmidt-Kastner PK, Pietsch T, Wiestler OD, & Blumcke I (1996) Neun: A useful neuronal marker for diagnostic histopathology. J Histochem Cytochem. 44: 1167-1171.

Won SY, Choi SH, & Jin BK (2009) Prothrombin kringle-2-induced oxidative
stress contributes to the death of cortical neurons in vivo and in vitro: Role of
microglial nadph oxidase. J Neuroimmunol. 214: 83-92.
Wong-Goodrich SJ, Glenn MJ, Mellott TJ, Liu YB, Blusztajn JK, & Williams CL
(2011) Water maze experience and prenatal choline supplementation
differentially promote long-term hippocampal recovery from seizures in
adulthood. Hippocampus. 21: 584-608.
Wu XG, Wang SS, Miao H, Cheng JJ, Zhang SF, & Shang YZ (2016) Scutellaria
barbata flavonoids alleviate memory deficits and neuronal injuries induced by
composited abeta in rats. Behav Brain Funct. 12: 33.
Yang Y, Liu Z, Cermak JM, Tandon P, Sarkisian MR, Stafstrom CE, Neill JC,
Blusztajn JK, & Holmes GL (2000) Protective effects of prenatal choline
supplementation on seizure-induced memory impairment. J Neurosci. 20:
RC109.
Yassine N, Lazaris A, Dorner-Ciossek C, Despres O, Meyer L, Maitre M,
Mensah-Nyagan AG, Cassel JC, & Mathis C (2013) Detecting spatial memory
deficits beyond blindness in Tg2576 Alzheimer mice. Neurobiol Aging. 34:
716-730.
You JC, Muralidharan K, Park JW, Petrof I, Pyfer MS, Corbett BF, LaFrancois JJ,
Zheng Y, Zhang X, Mohila CA, Yoshor D, Rissman RA, Nestler EJ, Scharfman
HE, & Chin J (2017) Epigenetic suppression of hippocampal calbindin-D28k by
ΔFosB drives seizure-related cognitive deficits. Nat Med. 23: 1377-1383.
You JC, Stephens GS, Fu CH, Zhang X, Liu Y, & Chin J (2018) Genome-wide
profiling reveals functional diversification of ΔFosB gene targets in the
hippocampus of an Alzheimer's disease mouse model. PLoS One. 13:
e0192508.
Zeisel SH, & da Costa KA (2009) Choline: An essential nutrient for public health.
Nutr Rev. 67: 615-623.
Zhang TY, Keown CL, Wen X, Li J, Vousden DA, Anacker C, Bhattacharyya U,
Ryan R, Diorio J, O'Toole N, Lerch JP, Mukamel EA, & Meaney MJ (2018)
Environmental enrichment increases transcriptional and epigenetic
differentiation between mouse dorsal and ventral dentate gyrus. Nat Commun.
9: 298.

1318