Population genomics of harbour seal *Phoca vitulina* from northern British

Columbia through California and comparison to the Atlantic subspecies

Ben J. G. Sutherland[1,2], Claire Rycroft[2], Ashtin Duguid[2], Terry D. Beacham[2], Strahan Tucker[2]

[1] Sutherland Bioinformatics, Lantzville, British Columbia, Canada, V0R 2H0.

[2] Fisheries and Oceans Canada, Pacific Biological Station, 3190 Hammond Bay Road, Nanaimo, BC V9T

6N7, Canada.

Author for correspondence: BJGS

Current Address:

Sutherland Bioinformatics, Lantzville BC, Canada, V0R 2H0

Email: Sutherland.Bioinformatics@protonmail.com

Author for correspondence: ST

Fisheries and Oceans Canada, Pacific Biological Station, 3190 Hammond Bay Road, Nanaimo, BC V9T

6N7, Canada

Email: Strahan.Tucker@dfo-mpo.gc.ca

Abstract

The harbour seal *Phoca vitulina* is a ubiquitous pinniped species found throughout coastal waters of the Northern Hemisphere. Harbour seal impacts on ecosystem dynamics may be significant due to their high abundance and food web position. Two subspecies exist in North America, *P. v. richardii* in the Pacific Ocean, and *P. v. vitulina* in the Atlantic. Strong natal philopatry of harbour seals can result in fine-scale genetic structure and isolation-by-distance. Management of harbour seals is expected to benefit from improved resolution of seal population structure and dynamics. Here we use genotyping-by-sequencing to genotype 146 harbour seals from the eastern Pacific Ocean (i.e., British Columbia (BC), Oregon, and California) and the western Atlantic Ocean (i.e., Québec, Newfoundland, and Labrador). Using 12,742 identified variants, we confirm the recently identified elevated genetic diversity in the eastern Pacific relative to the western Atlantic and greatest differentiation between the subspecies. Further, we demonstrate that this is independent of reference genome bias or other potential technical artefacts. Coast-specific analyses with 8,933 and 3,828 variants in Pacific and Atlantic subspecies, respectively, identify divergence between BC and Oregon-California, and between Québec and Newfoundland-Labrador. Unexpected PCA outlier clusters were observed in two populations due to cryptic relatedness of individuals; subsequently, closely related samples were removed. Admixture analysis indicates an isolation-by-distance signature where Oregon seals contained some of the BC signature, whereas California did not. Additional sampling is needed in the central and north coast of BC to determine whether a discrete separation of populations exists within the region.

Keywords: breeding colony; genetic diversity; genotyping; harbour seal; marine mammal; *Phoca vitulina*; population genomics; RADseq; relatedness

Introduction

The harbour seal (*Phoca vitulina* Linnaeus 1758) is the most widely distributed pinniped species globally

(Andersen & Olsen, 2010; Blanchet et al., 2021). The reported number of coastal marine harbour seal

subspecies varies from three to five, but typically the recognized subspecies include *P. v. stejnegeri* (Allen

1902) in the western Pacific Ocean, *P. v. richardii* (Gray 1864) in the eastern Pacific Ocean, *P. v. vitulina*

(Linnaeus 1758; formerly *P. v. concolor*; De Kay 1842) in the western Atlantic, and *P. v. vitulina* in the

eastern Atlantic (Andersen & Olsen, 2010). Deep divergence exists between the Pacific and Atlantic

subspecies (Liu et al., 2022), where the separation is estimated to have occurred 1.7-2.2 million years ago

(Andersen & Olsen, 2010). Secondly, strong differentiation exists between western and eastern subspecies

within each ocean (Liu et al., 2022; Stanley et al., 1996). Historical events and glaciations have had large

influences on the present day population genetics of harbour seals (Stanley et al., 1996). A step-wise

reduction in genetic diversity occurs, whereby the highest diversity is in the eastern Pacific Ocean, followed

by the western Atlantic and western Pacific, and finally the lowest in the eastern Atlantic Ocean, possibly

indicating colonization (Liu et al., 2022) and/ or recolonization events (Stanley et al., 1996). These

historical population dynamics, along with contemporary population structure, are important to consider

for species management and conservation (Olsen et al., 2014).

Harbour seals have been challenged by habitat loss, interspecific competition (Cordes et al., 2017),

predation, epizootics, and human hunting or culling activity; different stressors have impacted different

regions, but generally most regions have seen historical population declines, many of which have rebounded

in recent years (Brown et al., 2005; DFO, 2022; Jeffries et al., 2003; Muto et al., 2021; Olesiuk, 2010). In

the eastern Pacific Ocean, large-scale predator control programs and harvests in the late 1800s to mid 1900s

reduced the subspecies down to approximately 10,000 seals (Olesiuk, 2010). Harbour seals were then

protected in the early 1970s, after which their numbers dramatically increased to over 100,000 animals and

then stabilized at the expected carrying capacity (Brown et al., 2005). By comparison, the population in

eastern Canada (excluding Newfoundland) is comprised of 8,000-12,000 seals (reviewed by Blanchet et

al., 2021). Throughout the range, declines and bottlenecks may have left impacts on the genetic diversity

of the species (Burg et al., 1999; Westlake & O'Corry-Crowe, 2002). Severe reductions in population size

in the eastern Atlantic Ocean from overhunting, environmental pollution, and epizootics, may have resulted

in bottlenecks and resultant low genetic diversity (Andersen et al., 2011; Goodman, 1998; Swart et al.,

1996), leading to calls for immediate monitoring in some areas (Andersen et al., 2011).

Standing genetic variation is important for rapid adaptation (Barrett & Schluter, 2008). In harbour

seals, this may preserve the evolutionary potential for handling disease outbreaks or environmental

perturbations (Andersen & Olsen, 2010), especially at the edges of the range where changes in climate may

have a more drastic impact (Andersen et al., 2011; Blanchet et al., 2021). Genetic diversity has been directly

connected to fitness in harbour seals, where lower heterozygosity was associated with increased lungworm
burden in juvenile Wadden Sea harbour seals (Rijks et al., 2008). Further, birth weight and survival-to-
weaning correlate with genetic diversity in harbour seal pups on Sable Island, Nova Scotia (Coltman et al.,
1998). Quantifying genetic diversity of populations may help estimate adaptive potential (Barrett &
Schluter, 2008), and preserving this diversity is an important conservation goal that is expected to guide
management decisions.

Isolation-by-distance (IBD) is a main driver of population structure in harbour seals (Goodman,
1998; Huber et al., 2012; Liu et al., 2022; Stanley et al., 1996; Westlake & O'Corry-Crowe, 2002), with
increased genetic separation occurring with discontinuous distribution of colonies (Goodman, 1998).
Increased differentiation is also observed in smaller populations due to elevated impacts of genetic drift
(Westlake & O'Corry-Crowe, 2002). Unexpectedly high differentiation has been observed in relatively
close proximities (e.g., western and eastern Baltic Sea; Stanley et al., 1996). Harbour seals can travel great
distances, but they also exhibit high philopatry to breeding colony (Goodman, 1998; Härkönen & Harding,
2001). From eastern North America to Europe, 12 regional populations have been identified, although the
true number is likely greater (Andersen & Olsen, 2010), which has been further explored recently (Liu et
al., 2022). Closer inspection of regions typically identifies more fine-scale structure; the United Kingdom
is now delineated into northwest, northeast, and southeast populations (Olsen et al., 2017), and Ireland is
delineated into northwest/north, eastern, and southwest populations (Steinmetz et al., 2023). Similarly,
eastern North America was originally considered a single population, fitting with continuous clinal
variation in pupping season (Temte et al., 1991), but has now been further characterized (see Liu et al.,
2022). Western North America has previously been considered to be comprised of three coastal harbour
seal populations, delineated as southeast Alaska through northern BC; southern BC; and Washington
through California (Burg et al., 1999). A genetically distinct population was identified in Puget Sound,
Washington State (Lamont et al., 1996), and inland waters of Washington State may in fact contain three
discrete populations (Huber et al., 2012). Additional population structure occurs throughout Alaska,
following isolation-by-distance and stepping-stone models of genetic distance (Westlake & O'Corry-
Crowe, 2002). In general, most areas still require characterization of localized regions to understand
population dynamics at this finer scale (Blanchet et al., 2021).

Clear identification of genetic units is essential for understanding population status and
implementing management schemes (see Andersen & Olsen, 2010), which requires reliable and integrated
data, possibly with demographic information and population viability analyses (Olsen et al., 2014; Olsen et
al., 2017). In BC and Washington, perceived impacts of harbour seals on salmon have renewed discussions
on regional population control measures (Trzcinski, 2020). Moreover, increased interest in Food, Social or
Ceremonial (FSC) harvests in BC has led managers to consider regional harvest allocations, but there is

incomplete information on population structure or status. In BC, harbour seals are managed as one stock
(DFO, 2022). Population structure and management unit designation are particularly important to
understand at a fine-scale if the population is subject to harvest (Andersen & Olsen, 2010). A coastwide
assessment of population genetic structure and diversity could support the definition of units relevant for
management and conservation efforts (Waples & Gaggiotti, 2006), although other considerations, such as
local demography, can also be integrated into such analyses (Olsen et al., 2014). Population health is
important to consider in a metapopulation context, where some healthy populations may act as sources of
seals into other areas, and other populations in decline may act as sinks (Carroll et al., 2020).

Genetic studies on harbour seals have typically used allozymes (Swart et al., 1996), mitochondrial
DNA (e.g., Stanley et al., 1996; Westlake & O'Corry-Crowe, 2002), and microsatellites (e.g., Goodman,
1998; Huber et al., 2012). Recently, single nucleotide polymorphisms (SNPs) by restriction site-associated
DNA sequencing (RAD-seq; Baird et al., 2008; Peterson et al., 2012) were analyzed at a global scale (Liu
et al., 2022). SNPs have some known benefits over microsatellites (Hauser et al., 2011), such as ease and
automation of genotyping, interoperability between labs, and fewer individuals required to estimate allele
frequencies (Beacham et al., 2011). SNPs also enable screening of large numbers of markers to study both
neutral and adaptive variation (Andrews et al., 2016). Although high throughput methods such as RAD-
sequencing require some bioinformatic expertise (Shafer et al., 2017), once variants are identified they can
be integrated into amplicon panels, which can be easier to analyze to provide consistent and effective results
for conservation genomics applications (Meek & Larson, 2019). Broad-scale population genetic
assessments could benefit from a consistent and high throughput genotyping method, as well as
systematically sampled natal sites (Huber et al., 2012).

In the present study we genotype harbour seals from broad geographic regions in the eastern Pacific
Ocean (i.e., northern British Columbia (BC), Strait of Georgia (SOG), Oregon, and California), and in the
western Atlantic Ocean (i.e., Newfoundland, Labrador, and eastern Quebec). We analyze population
differentiation and diversity differences, shared ancestry fractions, as well as within-population relatedness,
and discuss the results in relation to the recently published global analysis (Liu et al., 2022), testing both
datasets for any impacts of reference bias on diversity differences. The high quality multilocus genotypes
generated here are expected to be useful for the continued characterization of the population genetics and
dynamics of this species.

Methods

Sample collection, DNA extraction, library preparation and sequencing

Harbour seal samples were obtained as skin or blood samples from collections in northern British Columbia (NBC) and the Strait of Georgia (SOG) in southern BC, Oregon (ORE), California (CAL), Québec (QUE), Newfoundland (NFL), and Labrador (LAB) (Figure 1; Table 1; Figure S1; Figure S2). Serum samples were also obtained from harbour seals in Alaska, but extractions failed to yield sufficient DNA and therefore were not included. Samples were often opportunistically collected and came from a variety of sources that included mortalities, strandings, rehabilitating captive seals, and research surveys. Agencies or organizations that collected the samples are indicated in Additional File S1, and all collections were made according to permits obtained by the individual organization as required (see details in Table S1).

DNA was extracted using magnetic bead extraction on a BioSprint automated DNA extraction system, following manufacturer's instructions (QIAGEN). Extracted purified DNA was then quantified by Qubit 2.0 with the dsDNA BR assay kit (Thermo Fisher) and normalized to 20 ng/µl in 10 µl volume. The normalized DNA was submitted to the Institute of Integrative Biology and Systems (IBIS) Genomic Analyses Platform of Université Laval, where libraries were prepared using double-digest restriction-site associated DNA sequencing (ddRAD-seq). The enzymes *Nsi*I and *Msp*I were used based on *in silico* cut site assessment using SimRAD (Lepais & Weir, 2014). Libraries were sequenced on an Ion Torrent sequencer (Thermo Fisher) in batches of 20 samples per sequencing chip. Raw single-end sequence data of variable read lengths with a mode of 177 bp were obtained for downstream analyses.

Bioinformatics

Raw sequence data were inspected for quality using fastqc (v.0.11.4; Andrews, 2010) and visualized using multiqc (v.1.14; Ewels et al., 2016). In general, the pipeline stacks_workflow (commit 868b410; E. Normandeau, GitHub) was followed, as explained below in brief. Adapters and any reads shorter than 50 bp were removed using cutadapt (v.1.9.1; Martin, 2011). Sequence data were then de-multiplexed using *process_radtags* of Stacks2 (v.2.62; Rochette et al., 2019) in parallel (Tange, 2022) trimming to a uniform length of 80 bp and de-multiplexing based on cut sites from the appropriate restriction enzymes. Samples were re-inspected for quality, then aligned to the harbour seal reference genome (i.e., GCA_004348235.1_GSC_HSeal_1.0; sourced from a female Pacific subspecies) using bwa (v.0.7.12-r1039; Li, 2013) and converted to bam format using samtools (v.1.17; Danecek et al., 2021). Numbers of reads and alignments per sample were quantified using custom scripts (see Data Availability). Two samples were found to be extreme outliers in a downstream principal components analysis (PCA) and were removed (i.e., NFL_102; NFL_121), as were two samples with fewer than 1 M reads (i.e., NBC_110, ORE_104).

Several approaches to genotyping were applied. First, the reference genome-based approach used

gstacks with software defaults. The *populations* module was then used to filter, retaining any locus that was

present in at least 70% of individuals, in all seven regional groupings, and at a minor allele frequency

(MAF) ≥ 0.01. Second, to evaluate possible effects of the reference genome on genotyping of the different

subspecies, *de novo* genotyping approaches were conducted within Stacks2 for each coast, using a balanced

design of 18 samples from each of SOG, ORE (eastern Pacific), and EQB, and NFL (western Atlantic). The

datasets were further balanced by normalizing all samples to 1.25 M reads each using seqtk's sample

function (v1.3-r106; Heng Li). The analytic pipeline for the project with all code is available on GitHub

(see Data Availability, *ms_harbour_seal*).

Population genetics

The percentage of polymorphic genomic sites, and per-population average observed heterozygosity (H_{OBS})

and nucleotide diversity (π) were calculated considering all genotyped loci in Stacks2. A variant call format

(VCF) file was generated using a single SNP exported per locus, and the mean genotype depth per locus

per sample and per population was calculated from the VCF file using the *extract.gt* function of vcfR (Knaus

& Grünwald, 2017). For downstream analysis, a genepop file was also generated including a single SNP

per locus using the Stacks2 populations module, and this was read into the R environment (R Core Team,

2023) using adegenet (v.2.1.8; Jombart & Ahmed, 2011). The data were also exported from Stacks2

including all variants in radpainter format for use in fineRADstructure (Malinsky et al., 2018). The

fineRADstructure analysis used software defaults and tutorials provided by the developers (see Data

Availability, *ms_harbour_seal*). The single-SNP per RAD-tag dataset was converted to genlight format

using dartR (v.2.0.4; Gruber et al., 2018) and coast-specific datasets were generated (i.e., Atlantic or

Pacific), each of which included a coast-specific MAF filter (i.e., MAF > 0.01).

Unsupervised PCAs were generated for the global and coast-specific datasets. Due to unexpected

clustering of samples within regional collections, inter-individual relatedness was estimated between

samples using related (v.1.0; Pew et al., 2015) and the Ritland statistic (Ritland, 1996) to characterize

potential cryptic relatedness. This analysis was conducted in the coast-wide data, as well as within PCA

groupings (e.g., all BC samples) to determine the impact of population structure on the estimated

relatedness. Within-population outlier pairs were identified based on the coast-wide data and defined as

pairs with Ritland relatedness values above the third quartile + 1.5x the interquartile range for that

population. For each outlier pair, one of the individuals was removed from the dataset until no more outlier

pairs remained above the cutoff per population, as described above. After purging the putative local

relatives, each dataset was filtered again for MAF (i.e., MAF > 0.01). Loci were then evaluated for Hardy-

Weinberg proportions within each population using the function *hw.test* of pegas, removing any variant that

was not conforming to HWP (p ≤ 0.01) in any one of the seven regional groupings. H_{OBS} and F_{ST} were then

calculated per locus for each dataset using the *summary* function of adegenet and the function *Fst* within

pegas (v.1.1; Paradis, 2010), respectively, and any locus with excess observed heterozygosity ($H_{OBS} > 0.5$)

was removed. These datasets were then used as inputs to coast-specific PCAs, genetic differentiation

assessment between populations with 95% confidence intervals of F_{ST} (Weir & Cockerham, 1984) in

hierfstat (v.0.5-11; Goudet & Jombart, 2022), and a neighbour-joining tree using the edwards.dist distance

metric (Cavalli-Sforza & Edwards, 1967) was generated using 10,000 bootstraps with the aboot function

of poppr (v.2.9.3; Kamvar et al., 2014). The retained loci and individuals in the final datasets were used to

subset the VCF file and vcftools (Danecek et al., 2011) was used to prepare bed, bim, and fam plink files

to be used in ADMIXTURE (Alexander et al., 2009). ADMIXTURE was run using *K* values from 1-6 for

the full dataset and for the east coast dataset. Optimal *K* was evaluated using a combination of lowest cross-

validation error, optimal *K* metrics MedMed K, MedMean K, and MaxMed K (Puechmaille, 2016), and

clumpak visualizations (Kopelman et al., 2015) as implemented in StructureSelector (Li & Liu, 2018), as

well as custom R scripts (see Data Availability). Private alleles were also identified between specific

groupings of samples using poppr. Analyses were conducted using functions within the GitHub repository

simple_pop_stats with analytic R scripts available within the repository *ms_harbour_seal* (see Data

Availability).

Re-analysis of Liu et al. 2022

A recently published RAD-sequencing dataset using the *Pst*I enzyme that used genotype likelihood calls to

investigate global trends in harbour seal genomic diversity (Liu et al., 2022) was downloaded using the

RunSelector from the Short Read Archive (SRA) of NCBI and the SRA Toolkit (NCBI, 2023). Specifically,

14 samples from each of the following collections were downloaded: Endicott Arm (END) and Kodiak,

Alaska (KOD); Île du Bic (BIC) in the St. Lawrence River, Quebec; Newfoundland (NFL); Orkney,

Scotland (ORK); and Wadden Sea, Netherlands (WNL). Samples were analyzed using the reference-based

genotyping methods described above, as well as by region-specific, individual *de novo* analyses. All reads

were truncated to 70 bp and any read shorter than 70 bp was removed from the analysis, due to Stacks2

developers' recommendation to keep all reads the same length for *de novo* analyses. In brief, the all-region

approach retained loci present in all six populations in at least 70% of individuals per population, with

global MAF ≥ 0.01. The three separate *de novo* analyses were also filtered requiring both populations in

each dataset to be genotyped in at least 70% of the individuals in each population, and all retained variants

to have global MAF ≥ 0.01. Population-level average H_{OBS} was calculated by the Stacks2 populations

module.

Results

Genotyping

The 146 samples retained in the analysis had an average (± s.d.) of 3.8 ± 2.0 M reads each and were comprised in total of 561.6 M reads (see Table 1 for totals per population). No outlier samples were identified at this stage (but see Methods for earlier outlier removal). Alignment rates against the reference genome, which was generated from a Pacific subspecies female, ranged from 88.5-95.1% for Pacific samples and 85.9-87.3% for Atlantic samples. Genotyping guided by reference genome alignments used 96% of the 508.6 M primary alignments, discarding 2.8% due to low mapping quality and 1.2% due to excessive soft clipping. This resulted in the formation of 1.11 M loci (i.e., RAD-tags) comprised of 488.1 M reads with an average effective coverage of 27.5 ± 27.9x prior to filtering. Filtering for missing data and minor allele frequency (MAF; see Methods) resulted in 53,684 loci retained with a mean length of 86.6 bp. For the genotyped loci, per individual missing data was on average 2.9 ± 3.0%. These final loci were comprised of 4.6 M genomic sites, and 12,742 variants were observed, or 10,847 variants if only retaining a single SNP per locus. The mean genotype depth for all samples and loci was 35.1x, with a wide variance between samples (Figure S3), with each population ranging from a mean genotype depth of 13.3x (SOG) to 83.3x (ORE; Figure S4). For sub-species-specific analyses, data were then separated into west and east coast datasets, and once separated, each coast was filtered again for MAF, retaining 8,933 variants in the Pacific dataset and 3,828 variants in the Atlantic.

Subspecies genetic differentiation and diversity differences

The two subspecies were found to be highly differentiated (Figure 1C), separating across PC1 (33.3% of total dataset variation; Figure 2A), similar to that recently reported (Liu et al., 2022). As an example, CAL and NFL 95% confidence interval (c.i.) F_{ST} was 0.40-0.42 (Table S2).

Percentages of polymorphic genomic sites also differed greatly between the subspecies, and when considering populations with at least 15 individuals ranged from 0.180-0.194% in Pacific populations and 0.083-0.087% in Atlantic populations (see Table 2 for population-specific metrics). Observed heterozygosity (H_{OBS}) was 1.9-fold higher in the Pacific than in the Atlantic subspecies for these collections (mean: 0.00045 and 0.00024, respectively), consistent with recently reported results (Liu et al., 2022).

To further evaluate the large differences in polymorphism between the coasts, we re-analyzed the data per coast using a consistent sample size and read depth per sample, and without the reference genome (i.e., *de novo*; see Methods). For each of the two coasts, 45 M reads were used in total, and after filters, 17,490 loci comprising 1.4 M genomic sites and 3,373 variants were observed for the Pacific subspecies (mean percent polymorphic sites = 0.183%), and 40,909 loci comprising 3.3 M genomic sites and 3,483 variants for the Atlantic subspecies (mean percent polymorphic sites = 0.085%). This demonstrates that the

cause of the elevated diversity was not due to reference genome bias. Notably, in the balanced and

normalized dataset, ORE showed higher observed heterozygosity than SOG (H_{OBS} = 0.00048 and 0.00039,

respectively; Table S3).

A subset of the data from Liu et al. (2022) was also re-analyzed to confirm the trends viewed here.

Using the reference-guided genotyping on the subset of the samples described (see Methods), Pacific H_{OBS}

ranged from 0.00031 to 0.00034, whereas the western Atlantic was estimated at 0.00019 (i.e., 1.7-fold

higher H_{OBS} in the Pacific), and 0.00009-0.00010 in the eastern Atlantic (Table S3). The *de novo* approach

applied to a subset of the Liu et al. (2022) data resulted in a Pacific H_{OBS} of 0.00074 and Atlantic H_{OBS} of

0.00035-0.00037 (i.e., 1.8-fold higher H_{OBS} in Pacific). Collectively, this indicates that the differences in

diversity reported previously and that are observed here are not due to differences in sample size, read

depth, nor genetic divergence of individuals from the reference genome (i.e., reference bias).

Cryptic relatedness within populations and genetic differentiation between populations

Coast-specific datasets were each inspected by PCA, and this showed an outlier cluster with nine

individuals in the SOG, all of which were sampled in Burrard Inlet, and an outlier grouping comprised of

harbour seals from the southernmost sampled region in CAL (Figure S5). Each population was then

inspected for relatedness in a coastwide analysis (Figure S6), and it was determined that both PCA outlier

clusters were comprised of individuals with elevated relatedness (outlier pairs Ritland metric $\geq$ 0.0975 and

0.1027 in CAL and SOG, respectively). Notably, when analyzing relatedness within a single regional

grouping (e.g., BC only) rather than using all samples on the coast, the absolute value of relatedness metric

decreased, but the same trend of outliers remained (see Figure S7).

Private alleles were identified for the Burrard Inlet cluster, which identified a private allele

observed nine times in seven of the individuals (i.e., 586731_19.04), two private alleles observed six times

(i.e., 729912_41.01, 807825_8.01), and three private alleles observed five times (i.e., 71641_26.04,

503353_64.04, 839341_28.01). By comparison, analyzing a haphazard selection of nine SOG individuals

that were not from the Burrard Inlet cluster found no private alleles observed more than three times.

A population tree derived from the co-ancestry matrix based on shared microhaplotypes between

individuals indicated two shallow branches for the Burrard Inlet cluster (Figure S8), but there was no

difference between or within these two branches in the inter-individual relatedness statistic (Additional File

S2), and therefore individuals within the Burrard Inlet cluster were considered approximately equally

related, with no evidence for further family structure. Elevated relatedness outlier pairs were then identified

and one individual from each outlier pair was removed to remove putative close relatives (see Methods;

Table 2; Table S4), resulting in the retention of 73 and 33 individuals in the Pacific and Atlantic datasets,

respectively (total n = 106). MAF filters were then reimposed (Pacific retained: 8,721; Atlantic retained:

3,658), and loci not conforming to HWP or with excess H_{OBS} were removed (Table 2), resulting in 7,699

and 3,134 variants retained in the Pacific and Atlantic datasets, respectively. Of these remaining variants,

56.3% and 46.5% were between 0.01-0.10 MAF in the Pacific and Atlantic harbour seal datasets,

respectively (Figure 3).

The Pacific dataset shows clear differentiation between the southern and northern samples,

explaining 7% of the variation across PC1 and separating SOG/NBC from ORE/CAL (Figure 2B; Figure

S9 for Scree plots), with significant genetic differentiation (e.g., SOG-CAL 95% c.i. F_{ST} = 0.065-0.071;

Table 3A). Within BC, overlap of confidence interval ellipses is observed for NBC and SOG, as well as for

ORE and CAL in the USA. These collections show similar levels of differentiation, with NBC-SOG F_{ST} =

0.014-0.021 and ORE-CAL F_{ST} = 0.006-0.009 (Table 3A). Due to the low sample size of the NBC

collection, the SOG-NBC F_{ST} value should be taken with caution. Notably, the closest SOG samples in the

PCA to the NBC samples were some of the most northern samplings within the SOG collection, including

Nelson Island on the Sunshine Coast (SOG_118), and Anvil Island in Howe Sound (SOG_123). Sample

NBC_111, sampled at northern Haida Gwaii was one of the most distant samples geographically, and the

most distant sample along PC1 from the SOG samples (Figure 2B; Additional File S3). Putative relatives

(removed in final analysis), NBC_112 and NBC_113, collected from Prince Rupert and Massett (Haida

Gwaii), respectively, were also grouped together with NBC_111 distantly from the SOG, but this is also

expected due to the estimated relatedness to NBC_111. However, while some geographic trend is therefore

apparent, given the low number of characterized samples from northern BC, there is no clear delineation

yet observed in the data between the SOG and the more northern samples (Figure 2B).

The Atlantic subspecies analysis has a generally strong separation between the EQB and the

LAB/NFL collections along PC1 (5.3% of the variation; Figure 2C; Figure S9). After removing putative

siblings from the EQB collections, the three samples that were highest on PC2 were sampled at or near the

Métis harbour seal colony (EQB), and all those lower on PC2 were from the Le Bic colony (EQB), which

appears to be explaining the separation of these samples across both PC2 and PC1. The LAB and NFL

collections were completely overlapping with each other and have an F_{ST} value overlapping zero,

suggesting that these are not separate populations. The separation of EQB and NFL is however reflected in

the significant F_{ST} between EQB and NFL (95% c.i. F_{ST} = 0.016-0.023; Table 3B).

Admixture analysis using data from both coasts after removing putative relatives was conducted at

K = 1-6 (all major modes shown in Figure S10). The lowest cross-validation (cv) error occurred at K = 3

(median cv error = 0.3019; Figure S11), and although K = 4 had a slightly higher value (cv error = 0.3208),

the minor modes of K = 4 (4/10 runs) clearly separated EQB from NFL/LAB (Figure 4; Figure S12). The

major modes of K = 4 (6/10 runs) separated the SOG into mixed ancestry individuals. When analyzing the

east coast dataset by itself, although the lowest CV error was at K = 1 (Figure S13), StructureSelector found

355 MedMed K, MedMean K, and MaxMean K all supporting the optimal $K = 2$ in the east coast data (Figure

356 S14), which separated EQB from LAB/NFL as per the minor modes of $K = 4$ in the both coast dataset

357 (Figure S15). Notably, the minor modes of $K = 4$ in the both coast data indicated that the SOG samples

358 were nearly completely comprised of the SOG ancestry, whereas the NBC samples had a small fraction of

359 ancestry shared with the other coastal samples from ORE and CAL. The lack of ORE/CAL ancestry in the

360 SOG was common in both the major and minor modes of $K = 4$. The ORE samples showed a small fraction

361 of the BC (SOG) signature, whereas the CAL samples were comprised completely of the CAL ancestry,

362 suggesting that ORE is more similar to BC than CAL is to BC. Both the PCA and the admixture analysis

363 indicate that ORE is most similar to CAL, even though its distance from BC and CAL are approximately

364 similar (Figure 1A).

366 *Future work and utility of identified markers*

367 Markers were identified here that may be useful for further characterization of population structure (i.e.,

368 loci with elevated F_{ST} between southern and northern BC or between BC and Oregon) or for individual

369 identification (i.e., elevated per locus H_{OBS} in both the BC and US collections; Additional File S4).

370 Contrasting per locus H_{OBS} between the BC and US datasets indicates high correlation in H_{OBS} between the

371 datasets (Figure S16). These markers could be implemented to design genotyping panels as needed, and

372 those with consistently elevated H_{OBS} in multiple populations may have utility on a coastwide scale, at least

373 in the regions assessed here.

Discussion

376 *Subspecies divergence and differential genetic diversity*

377 Species conservation and management of harbour seal will likely depend on effective characterization of

378 population structure and contemporary standing genetic variation (including functional variation), that is

379 considered in the context of large-scale historical demographic events (as per Beichman et al., 2023), as

380 well as contemporary and local demography (Olsen et al., 2014). In this study, we confirm the existence of

381 elevated genetic diversity in the eastern Pacific subspecies of harbour seal *P. v. richardii* over the western

382 Atlantic subspecies *P. v. vitulina* as was recently reported by Liu and co-workers (2022). We observe the

383 elevated diversity of the Pacific subspecies in new regions including British Columbia (BC) and Oregon,

384 whereas the previous analysis included Alaska and California for the eastern Pacific subspecies (Liu et al.,

385 2022). Further, we evaluate and find that this trend is not caused by reference genome bias in genotyping,

386 where individuals with increased divergence from the reference genome may exhibit biased population

387 genetic metrics (Bohling, 2020). Reference bias is typically considered in the context of divergent species,

388 but as more local genomes are produced, these issues may become more relevant within species (Beichman

et al., 2023; Thorburn et al., 2023). Further, given the high divergence between the subspecies here, it was

important to test whether *P. v. vitulina* metrics were impacted by the alignment to the *P. v. richardii*

genome. The diversity difference of around 1.8x higher (evaluated by H_{OBS}) in the Pacific over the Atlantic

North American subspecies remains when genotyping without a reference genome (i.e., *de novo*) with a

normalized number of samples and reads.

The ultimate cause of the diversity difference was proposed by Liu and co-workers (2022) to be

due to stepwise founder effects during colonization from the Pacific Ocean to the western Atlantic Ocean,

and then from the western Atlantic Ocean to the eastern Atlantic Ocean, with a decrease in genetic diversity

at each step. Proposed west to east colonization within each basin is supported by previous work (Stanley

et al., 1996), although some evidence also exists for an east to west radiation in the Pacific (Westlake &

O'Corry-Crowe, 2002). Stanley and co-workers (1996) proposed that the European population may have

been eliminated during the last Ice Age (~18,000 y.a.), and that it was recolonized following the recession

of the ice sheets. These estimates and additional demographic history of these populations will be important

to explore with next-generation sequencing data (Beichman et al., 2023), albeit with caution given the

susceptibility of demographic inference to artefacts from pipeline parameters (Shafer et al., 2017). The use

of these multiple, independent RAD-seq datasets may aid in this analysis. The relatively deep sequencing

and therefore highly reliable genotypes expected in the present analysis, and the numerous RAD-tags

generated through restriction enzyme selection (Lepais & Weir, 2014), may enable the analysis of the allele

frequency spectra of the two subspecies to characterize fluctuations in effective population size (Ne) and

therefore approximate the timing of the reduced genetic diversity event (e.g., Beichman et al., 2023; Liu &

Fu, 2020). Uncertainty levels around the demographic inference approaches may not be robust enough for

conservation and management decisions, but results of these approaches may increase general

understanding of putative and broad biogeographic hypotheses (Shafer et al., 2017).

The striking diversity differences between the subspecies may provide a model to study the effect

of environmental changes and adaptive potential in the two subspecies, and their responses to environmental

challenges. Furthermore, the large divergence between the Pacific and the Atlantic subspecies raises the

question of whether the subspecies are in fact separate species (Liu et al., 2022). F_{ST} values observed by

Liu et al. (2022) ranged from 0.38 to 0.69, and the authors noted that this is higher than most other marine

megafaunal subspecies to date. Our results agree with these levels, and although we did not have samples

from the eastern Atlantic Ocean, here the 95% lower and upper F_{ST} values for all contrasts between the

west and east coast of North America (excluding the low sample size NBC) ranged from $F_{ST} = 0.36$ to 0.43.

Population structure in the eastern Pacific

Harbour seals from the eastern Pacific Ocean along North America have been considered to be comprised

of three coastal populations: (1) southeast Alaska through northern BC; (2) southern BC; and (3)

Washington through California (Burg et al., 1999). In the present study, the separation between BC and

Oregon/California is clear. Additionally, here, based on PCA, admixture analysis, and F_{ST} some separation

is viewed among samples from southern BC and northern BC, and among samples from Oregon and

California, but there are no clear break points or strong separations observed between the putative

populations aside from the separation between BC and Oregon/California. Continuous distributions of

breeding animals along coastlines has previously been described to have little local substructuring, although

a continued increase in differentiation occurs with distance (Goodman, 1998). In the case of clinal isolation-

by-distance, sampling at extreme ends of a geographical region may result in apparent clusters, whereas if

the sampling was continuous throughout the range, a cline would be observed (Patterson et al., 2006). In

this regard, continuous sampling along coastlines would help refine whether breakpoints exist or if a

continuous distribution explains the coastal population structure. In addition to the continuous distribution

of breeding animals occurring along the coast, the detection of substructure may also be impeded due to

high population abundance (Burg et al., 1999). The location of a putative dividing line between southern

and northern BC has yet to be identified (Burg et al., 1999). In the present study we also do not identify this

line, which may be due to having too few northern BC samples genotyped here, and the location may be

only identifiable with a larger survey of coastal breeding colonies. Furthermore, the samples used in the

present study were mainly opportunistically collected, and this type of sampling has been proposed to blur

the lines between populations as opposed to careful sampling of unweaned pups to ensure exact natal

locations (Huber et al., 2012). In any case, markers used here did show some spread of samples in the PCA

and some differences in admixture fractions, between northern and southern BC, and between Oregon and

California. A subset of markers identified here, applied to a higher number of individuals in future sample

collections using amplicon sequencing approaches for example, may help resolve some of these questions.

A focus on systematic sampling of natal sites within the Central Coast region may be key in identifying a

potential dividing line within BC (Burg et al., 1999), if it exists.

The Strait of Georgia (SOG) has the highest density of harbour seals on the BC coast with nearly

40,000 animals (i.e., 37% of BC population), and is used for population surveys since the implementation

of protections (DFO, 2022; Majewski & Ellis, 2022). Notably, of the 12 harbour seals sampled in the

Burrard Inlet area, nine were found to have high estimated genetic relatedness. These harbour seals were

all pups, and were all sampled in the same summer. As female harbour seals only produce a single pup per

year (Schaeff et al., 1999), it is possible that the samples could be half-sibs, sharing a sire. However, harbour

seals are not expected to mate in harems, but rather in-water in a lek-type mating system (Boness et al.,

2006), which makes the observation of so many putatively related pups surprising. Colonies having

different levels of overall relatedness have been previously reported (Olsen et al., 2014). Regardless, the

unexpected relatedness (i.e., cryptic relatedness) had significant effects on the initial PCA (Yao & Ochoa,

2023), highlighting the importance of removing putative relatives to avoid erroneous identification of

population clusters (Patterson et al., 2006), given their overrepresentation of certain alleles (Elhaik, 2022).

This feature of population characterization, which is expected in other methods and not only PCA (Patterson

et al., 2006), should be considered in future sampling designs. It is interesting to note that the coastal BC

samples in northern BC showed a fraction of admixture signature that was shared with ORE/CAL, whereas

the SOG samples largely did not contain any of this signal (except one sample with a small fraction). This

further suggests isolation of the SOG samples, along with the increased relatedness signatures observed.

The presence or extent of any genetic separation existing between this colony (or colonies) and others is

unknown, and would need follow-up sampling to understand further. The BC signature was viewed in a

minor fraction of the ORE samples, but not the CAL samples, providing evidence of isolation-by-distance

in harbour seals that are furthest south from BC. In harbour seals, isolation-by-distance may not explain all

genetic separation, and significant differentiation can occur within only a few hundred kilometers,

potentially being driven by other demographic processes (Goodman, 1998; Stanley et al., 1996).

Inland populations in Washington State that are genetically distinct from the coastal population

were observed to have the most diverse mitochondrial haplotypes (Lamont et al., 1996). These populations

have different pupping times and cranial morphometrics, although are geographically close, and may be

more closely related to harbour seals from northern populations, including the SOG (reviewed by Lamont

et al., 1996). Late birthing of pups has been observed in Puget Sound and Vancouver Island that is

discontinuous with clinal variation observed further south, again suggesting some unique characteristics in

this area (Temte et al., 1991). Furthermore, Burg and co-workers (1999) identified a unique mitochondrial

haplotype signature in harbour seals sampled from southern Vancouver Island, but did not see this group

as a separate population in their microsatellite analysis. Similarly, in the present work, we noted high

frequency private alleles in the Burrard cluster of putatively related individuals prior to their removal.

Unique signatures were suggested to potentially originate from a separate colonization event for these

populations (Burg et al., 1999; Lamont et al., 1996). Whether the Burrard cluster is simply an artefact of

chance sampling of individuals with high relatedness from a fairly genetically homogenous colony in the

inlet, skewing allele frequencies and enriching rare alleles (Elhaik, 2022), or in fact is evidence of impeded

gene flow, requires further study with systematically obtained samples over time. Additional samples from

the inland populations of Washington State (Lamont et al., 1996) to compare with the Burrard cluster would

also be important to include in future analyses. A southern California cluster was also observed, and

similarly this group was found to have high estimated relatedness, and therefore was required to have close

relatives removed to not appear as artefactual population structure in the PCA (Patterson et al., 2006). This

provides additional support for the hypothesis that the family effect is driving the intra-collection clusters.
However, it is not clear why related individuals were specifically sampled in this area through the
opportunistic sampling methods applied. A caution regarding the removal of the putative close relatives is
that we are unable to determine whether this is an isolated colony (without significant gene flow to
neighbouring colonies); the highly related nature of the animals may simply be a characteristic of an isolated
population (Waples & Anderson, 2017), and the lack of the signature of ORE or CAL in the SOG samples
in the admixture analysis further indicates the isolation of this population, even in contrast with other
samples further north in BC.

California harbour seals are managed separately from Oregon and coastal Washington State, but
this is due to jurisdictional or political separation rather than due to genetic information (Brown et al.,
2005). In the present results, the level of genetic differentiation between ORE and CAL (i.e., $F_{ST} = 0.008$-
0.011) was lower than that between NBC and SOG (i.e., $F_{ST} = 0.016$-0.021), or EQB and NFL (i.e., $F_{ST} =$
0.019-0.024), but see above regarding caution for NBC allele frequencies. Given the expected gene flow
between Oregon and neighbouring states, this may explain how the reduction to approximately 500 harbour
seals in Oregon before protections were implemented in the early 1970s (reviewed by Brown et al., 2005)
did not result in a suppression of genetic diversity in the contemporary population. Similarly, in sea otter
Enhydra lutris, simulations suggest that accumulated genetic load through bottlenecks and low Ne
decreases more rapidly with restoration of gene flow (Beichman et al., 2023). As described above, the
demographic history and Ne fluctuations of Pacific and Atlantic subspecies, including within each basin,
would be a valuable approach to understand how recorded historic events compare with the genetic
signatures observed.

Marker selection for an amplicon panel
Marker panels, such as a low-density 500 marker panel, are useful tools for conservation genetics (Meek &
Larson, 2019), including for characterizing population structure, conducting genetic assignments and
parentage, as well as for providing some estimates of genetic diversity. With the present data, markers
identified here could be used to generate such a panel. To select highly differentiating markers that will
separate populations along the range of the Pacific harbour seal, markers could be selected based on high
F_{ST} both between Northern BC and SOG, and between SOG and Oregon/California. However, a limitation
of this approach is that regions not sampled in the present study may be missed in such a design. Additional
goals of individual identification could be considered as well, and other markers with elevated
heterozygosity would be useful in this regard to include in a panel (e.g., markers with elevated H_{OBS} in both
the US and Canadian populations). Removal of the highly related outlier cluster is likely to be a good
approach for identifying markers, as was done here, but it is also likely valuable to include several of the

high frequency private alleles identified in the Burrard cluster in the panel for broader investigation. These

genetic efforts are expected to be useful for future planning of management units for the Pacific harbour

seal.

Conclusions

The present study characterizes 12,742 variants in 146 harbour seals sampled from seven different regions

in the east and west coast of North America. Top differentiating or high heterozygosity variants can be used

for future studies, for example to design an amplicon panel, to further characterize population structure, for

parentage analysis, or to enable repeat detections of individuals. This study confirms the recent observation

of elevated diversity in Pacific harbour seals over the western Atlantic subspecies, and verifies that this

trend is not due to analytic artefacts such as reference bias, sequencing depth, or sample size. Although

northern BC samples share some ancestry with coastal seals from Oregon and California, those sampled in

the Strait of Georgia largely did not. The Oregon seals shared some of the BC ancestry signature, whereas

further south, the California seals did not; this suggests isolation-by-distance, known to be a factor in

population structure of harbour seals. Cryptic relatedness was observed in seals sampled in Burrard Inlet

and southern California, and was subsequently removed from the dataset. Cryptic relatedness may have

broader implications for future sampling efforts from colonies to characterize discrete populations.

Additional samples, systematically characterized over time will be helpful to determine the extent of gene

flow between putative colonies. Samples of expected parents could also help to resolve the nature of the

relationships among samples in colonies through parentage analyses. Future studies using the multiple

independent datasets now available could explore the demographic history of both subspecies to determine

timing of Ne fluctuations on either coast to further investigate the causes of the striking differences in

genomic diversity between the subspecies.

Acknowledgements

This work was supported by Fisheries and Oceans Canada Pinniped Research Program and the Species At

Risk Program. The authors are grateful to sample providers, including Garry Stenson of DFO Atlantic

(Newfoundland and Labrador), Mike Hammill and Xavier Bordeleau of DFO Québec (eastern Québec),

the Marine Mammal Rescue Centre, Vancouver and DFO Pacific (Strait of Georgia, northern BC), James

Rice of the Oregon Marine Mammal Stranding Network (Oregon), Barbie Halaska of the Marine Mammal

Center of Sausalito (California), and Lauri Jemison and Lori Polasek from Alaska Department of Fish and

Game (ADFG). Thanks to the Molecular Genetics Lab of DFO Pacific for technical support and

discussions. Thanks to Jamieson Gorrell for discussions on admixture analyses. Thanks to Thierry Gosselin,

Morten Tange Olsen, four anonymous reviewers, and the subject Editor for their constructive comments on
earlier versions of the manuscript.
559

Competing Interests

561 Ben Sutherland is affiliated with Sutherland Bioinformatics. The author has no competing financial interests
562 to declare. The other authors declare that no competing interests exist.
563

Data Availability

All code required to analyze the project: https://github.com/bensutherland/ms_harbour_seal
Functions used for analysis: https://github.com/bensutherland/simple_pop_stats
Demultiplexed sequence data is available on NCBI through the SRA within BioProject PRJNA948428,
BioSamples SAMN33902654- SAMN33902801.
Associated files on FigShare including compressed GitHub analysis pipeline, sample information file
needed for *stacks_workflow*, genotypes for all populations from the reference-guided analysis (all samples
or normalized) in genepop and VCF formats (both are single SNP per locus) as well as RADpainter input
(microhaplotypes), and the balanced, normalized, *de novo* genotyping genepops (single SNP per locus)
and VCF files (multiple SNP per locus): https://doi.org/10.6084/m9.figshare.22290472
574

Benefit-sharing statement

576 This research addresses the population genetics of harbour seals in the eastern Pacific and western Atlantic.
577 We have ensured that all raw data, output genotype data, and metadata is provided, as well as well-
578 documented scripts and instructions for data analyses. All individuals and/or organizations who provided
579 samples for the project are included in the Acknowledgements section.
580

Supplemental Information

In addition to the Additional Files shown below, the Supplemental Results section contains supplemental
figures and tables.
Additional File S1. All sample metadata including location, sex, and age information.
Additional File S2. Pairwise relatedness statistics displaying histogram of all pairwise values and the top
interindividual relatedness pairs for both Pacific and Atlantic datasets.
Additional File S3. Coast-specific analyses PCA scores, sample IDs, location, and GPS coordinates.
Additional File S4. Per locus marker statistics for (a) Pacific populations and (b) Atlantic populations,
both with all putative relatives removed, as well as between (c) SOG and NBC and (d) SOG and ORE.

Figures and Tables

592
Figure 1. Sampling locations for the eastern Pacific (A), and western Atlantic (B) oceans, where each
individual harbour seal sampled is displayed as a coloured point when individual-level sampling
coordinates were available, or as a highlighted region when sample location details were not available (i.e.,
Labrador and Newfoundland). Genetic similarity among the populations is shown in the dendrogram (C)
along with bootstrap support at the nodes. Acronyms: NBC = northern British Columbia; SOG = Strait of
Georgia; CAL = California; ORE = Oregon; EQB = eastern Quebec; NFL = Newfoundland; and LAB =
Labrador.

Figure 2. Principal components analysis following removal of putative close relatives for (A) all samples from both coasts (n = 106); (B) the Pacific subspecies (*P. v. richardii*) collections with 7,699 filtered variants; and (C) the Atlantic subspecies (*P. v. vitulina*) with 3,134 filtered variants. Ellipses represent 95% confidence level for the multivariate t-distribution.

Figure 3. Minor allele frequency distribution for (A) Pacific or (B) Atlantic datasets after removing low MAF markers within each coast-specific dataset (i.e., retain MAF ≥ 0.01). Markers with MAF between 0.01 to 0.10 comprise 56.3% and 46.5% of the Pacific and Atlantic datasets, respectively.

Figure 4. Admixture analysis with $K = 4$ including individuals from both coasts (n = 106 individuals) after removal of individuals with elevated estimated genetic relatedness. This plot represents a single run, and plots with other values of K and multiple runs are shown in the supplemental materials. Pacific samples (i.e., NBC, SOG, ORE, CAL) are arranged by latitude, with the leftmost samples being those that were sampled from the most northern locations. Atlantic samples are grouped only by region, as most did not have exact sampling coordinates available.

Table 1. Per regional collection sample size (n), total reads (M; million) and per sample average (± s.d.) per collection, total alignments (M), per sample average (± s.d.), and average alignment rate (%; ± s.d.) per collection. Values were calculated after extreme outliers (n = 2) and low coverage outliers (n = 2) were removed.

Collection	n	Reads per collection (M)	Per sample average (M)	Alignments per collection (M)	Per sample average (M)	Alignment rate (%)
NBC	7	30.2	4.3 ± 3.0	27.5	3.9 ± 2.8	90.0 ± 2.2
SOG	41	138.6	3.4 ± 1.4	122.7	3.0 ± 1.2	88.5 ± 1.2
ORE	19	104.5	5.5 ± 2.4	99.5	5.2 ± 2.3	95.1 ± 0.8
CAL	30	108.7	3.6 ± 1.9	102.8	3.4 ± 1.8	94.1 ± 1.8
EQB	20	73.8	3.7 ± 2.0	63.9	3.2 ± 1.8	86.2 ± 2.0
LAB	7	29.0	4.1 ± 2.3	25.6	3.7 ± 2.1	87.3 ± 3.3
NFL	22	76.8	3.5 ± 2.0	66.5	3.0 ± 1.9	85.9 ± 2.7

Table 2. Summary statistics from genotyping 146 individuals at 12,742 variant sites in 4.6 Mbp indicates elevated percentage of polymorphic loci and observed heterozygosity (H_{OBS}) in the Pacific subspecies relative to the Atlantic subspecies. This dataset is not restricted to a single SNP per RAD locus. The number of individuals with elevated genetic relatedness that were removed from each collection is shown along with the number of loci removed in the final single-SNP per locus dataset that did not conform to Hardy-Weinberg proportions. The total variants remaining in Pacific and Atlantic after all filters were 7,699 and 3,134, respectively.

Pop	Avg. n per locus	Polymorph. variants	Percent polymorph. (%)	H_{OBS}	Nucleotide diversity (π)	Related cutoff (Ritland)	Related indiv. rem.	Loci non-HWP
NBC	6.8	6400	0.138	0.00044	0.00044	NA*	2	0
SOG	39.8	9018	0.194	0.00043	0.00044	0.1027	12	360
ORE	18.9	8358	0.180	0.00046	0.00046	0.0921	2	289
CAL	29.7	8528	0.184	0.00045	0.00046	0.0975	8	281
EQB	19.5	3875	0.083	0.00024	0.00024	0.1061	6	123
LAB	6.9	3257	0.070	0.00025	0.00024	0.0433	2	0
NFL	21.4	4044	0.087	0.00024	0.00024	0.0461	8	131

*no outlier pairs, applied same cutoff as SOG

634 **Table 3.** Genetic differentiation (Weir-Cockerham F_{ST}) between populations in the (A) Pacific subspecies
635 dataset comprised of 7,699 variants; and in the (B) Atlantic subspecies dataset comprised of 3,134 variants.
636 Values in the upper and lower sections are upper and lower 95% confidence limits of the estimated F_{ST},
637 respectively. Any negative values should be considered $F_{ST} = 0$, or no differentiation.

638 **(A)**

	NBC	SOG	ORE	CAL
NBC	-	0.021	0.052	0.064
SOG	0.014	-	0.057	0.071
ORE	0.043	0.052	-	0.009
CAL	0.055	0.065	0.006	-

640 **(B)**

	EQB	NFL	LAB
EQB	-	0.023	0.032
NFL	0.016	-	0.002
LAB	0.020	-0.007	-

REFERENCES

Alexander, DH, Novembre, J, & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research, 19*(9), 1655-1664. doi:10.1101/gr.094052.109

Andersen, L, & Olsen, MT. (2010). Distribution and population structure of North Atlantic harbour seals (*Phoca vitulina*). *NAMMCO Scientific Publications, 8*(0), 15-35. doi:10.7557/3.2669

Andersen, LW, Lydersen, C, Frie, AK, Rosing-Asvid, A, Hauksson, E, & Kovacs, KM. (2011). A population on the edge: genetic diversity and population structure of the world's northernmost harbour seals (*Phoca vitulina*). *Biological Journal of the Linnean Society, 102*(2), 420-439. doi:10.1111/j.1095-8312.2010.01577.x

Andrews, KR, Good, JM, Miller, MR, Luikart, G, & Hohenlohe, PA. (2016). Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews Genetics, 17*(2), 81-92. doi:10.1038/nrg.2015.28

Andrews, S. (2010). FastQC: A quality control tool for high throughput sequence data.

Baird, NA, Etter, PD, Atwood, TS, Currey, MC, Shiver, AL, Lewis, ZA, et al. (2008). Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLOS ONE, 3*(10), e3376. doi:10.1371/journal.pone.0003376

Barrett, RDH, & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends in Ecology & Evolution, 23*(1), 38-44. doi:10.1016/j.tree.2007.09.008

Beacham, TD, McIntosh, B, & Wallace, CG. (2011). A comparison of polymorphism of genetic markers and population sample sizes required for mixed-stock analysis of sockeye salmon (*Oncorhynchus nerka*) in British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences, 68*(3), 550-562. doi:10.1139/f10-167

Beichman, AC, Kalhori, P, Kyriazis, CC, DeVries, AA, Nigenda-Morales, S, Heckel, G, et al. (2023). Genomic analyses reveal range-wide devastation of sea otter populations. *Molecular Ecology, 32*(2), 281-298. doi:https://doi.org/10.1111/mec.16334

Blanchet, M-A, Vincent, C, Womble, JN, Steingass, SM, & Desportes, G. (2021). Harbour seals: population structure, status, and threats in a rapidly changing environment. *Oceans, 2*(1), 41-63.

Bohling, J. (2020). Evaluating the effect of reference genome divergence on the analysis of empirical RADseq datasets. *Ecology and Evolution, 10*(14), 7585-7601. doi:https://doi.org/10.1002/ece3.6483

Boness, DJ, Bowen, WD, Buhleier, BM, & Marshall, GJ. (2006). Mating tactics and mating system of an aquatic-mating pinniped: the harbor seal, *Phoca vitulina*. *Behavioral Ecology and Sociobiology, 61*(1), 119-130. doi:10.1007/s00265-006-0242-9

Brown, RF, Wright, BE, Riemer, SD, & Laake, J. (2005). Trends in abundance and current status of harbor seals in Oregon: 1977-2003. *Marine Mammal Science, 21*(4), 657-670. doi:https://doi.org/10.1111/j.1748-7692.2005.tb01258.x

Burg, TM, Trites, AW, & Smith, MJ. (1999). Mitochondrial and microsatellite DNA analyses of harbour seal population structure in the northeast Pacific Ocean. *Canadian Journal of Zoology, 77*(6), 930-943. doi:10.1139/z99-057

Carroll, EL, Hall, A, Olsen, MT, Onoufriou, AB, Gaggiotti, OE, & Russell, DJ. (2020). Perturbation drives changing metapopulation dynamics in a top marine predator. *Proceedings of the Royal Society B: Biological Sciences, 287*(1928), 20200318. doi:doi:10.1098/rspb.2020.0318

Cavalli-Sforza, LL, & Edwards, AW. (1967). Phylogenetic analysis. Models and estimation procedures. *American Journal of Human Genetics, 19*(3 Pt 1), 233-257.

Coltman, DW, Bowen, WD, & Wright, JM. (1998). Birth weight and neonatal survival of harbour seal pups are positively correlated with genetic variation measured by microsatellites. *Proceedings of the Royal Society of London. Series B: Biological Sciences, 265*(1398), 803-809. doi:doi:10.1098/rspb.1998.0363

Cordes, LS, O'Corry-Crowe, G, & Small, RJ. (2017). Surreptitious sympatry: Exploring the ecological
and genetic separation of two sibling species. *Ecology and Evolution, 7*(6), 1725-1736.
doi:https://doi.org/10.1002/ece3.2774
Danecek, P, Auton, A, Abecasis, G, Albers, CA, Banks, E, DePristo, MA, et al. (2011). The variant call
format and VCFtools. *Bioinformatics, 27*(15), 2156-2158. doi:10.1093/bioinformatics/btr330
Danecek, P, Bonfield, JK, Liddle, J, Marshall, J, Ohan, V, Pollard, MO, et al. (2021). Twelve years of
SAMtools and BCFtools. *GigaScience, 10*(2). doi:10.1093/gigascience/giab008
DFO. (2022). *Stock assessment of Pacific harbour seals (Phoca vitulina richardsi) in Canada in 2019.* .
DFO Can. Sci. Advis. Sec. Sci. Advis. Rep.
Elhaik, E. (2022). Principal component analyses (PCA)-based findings in population genetic studies are
highly biased and must be reevaluated. *Scientific Reports, 12*(1), 14683. doi:10.1038/s41598-022-
14395-4
Ewels, P, Magnusson, M, Lundin, S, & Käller, M. (2016). MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics, 32*(19), 3047-3048.
doi:10.1093/bioinformatics/btw354
Goodman, SJ. (1998). Patterns of extensive genetic differentiation and variation among European harbor
seals (*Phoca vitulina vitulina*) revealed using microsatellite DNA polymorphisms. *Molecular
Biology and Evolution, 15*(2), 104-118. doi:10.1093/oxfordjournals.molbev.a025907
Goudet, J, & Jombart, T. (2022). hierfstat: estimation and tests of hierarchical F-statistics.
Gruber, B, Unmack, PJ, Berry, OF, & Georges, A. (2018). dartr: An r package to facilitate analysis of
SNP data generated from reduced representation genome sequencing. *Molecular Ecology
Resources, 18*(3), 691-699. doi:https://doi.org/10.1111/1755-0998.12745
Härkönen, T, & Harding, KC. (2001). Spatial structure of harbour seal populations and the implications
thereof. *Canadian Journal of Zoology, 79*(12), 2115-2127. doi:10.1139/z01-172
Hauser, L, Baird, M, Hilborn, R, Seeb, LW, & Seeb, JE. (2011). An empirical comparison of SNPs and
microsatellites for parentage and kinship assignment in a wild sockeye salmon (*Oncorhynchus
nerka*) population. *Molecular Ecology Resources, 11*(s1), 150-161.
doi:https://doi.org/10.1111/j.1755-0998.2010.02961.x
Huber, HR, Dickerson, BR, Jeffries, SJ, & Lambourn, DM. (2012). Genetic analysis of Washington State
harbor seals (*Phoca vitulina richardii*) using microsatellites. *Canadian Journal of Zoology,
90*(12), 1361-1369. doi:10.1139/cjz-2012-0047
Jeffries, S, Huber, H, Calambokidis, J, & Laake, J. (2003). Trends and status of harbor seals in
Washington State: 1978-1999. *The Journal of Wildlife Management, 67*(1), 207-218.
doi:10.2307/3803076
Jombart, T, & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide SNP data.
Bioinformatics, 27(21), 3070-3071. doi:10.1093/bioinformatics/btr521
Kamvar, ZN, Tabima, JF, & Grünwald, NJ. (2014). poppr: an R package for genetic analysis of
populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ, 2*, e281.
doi:10.7717/peerj.281
Knaus, BJ, & Grünwald, NJ. (2017). vcfr: a package to manipulate and visualize variant call format data
in R. *Molecular Ecology Resources, 17*(1), 44-53. doi:https://doi.org/10.1111/1755-0998.12549
Kopelman, NM, Mayzel, J, Jakobsson, M, Rosenberg, NA, & Mayrose, I. (2015). Clumpak: a program
for identifying clustering modes and packaging population structure inferences across K.
Molecular Ecology Resources, 15(5), 1179-1191. doi:https://doi.org/10.1111/1755-0998.12387
Lamont, MM, Vida, JT, Harvey, JT, Jeffries, S, Brown, R, Huber, HH, et al. (1996). Genetic substructure
of the Pacific harbor seal (*Phoca vitulina richardsi*) off Washington, Oregon, and California.
Marine Mammal Science, 12(3), 402-413. doi:https://doi.org/10.1111/j.1748-
7692.1996.tb00592.x
Lepais, O, & Weir, JT. (2014). SimRAD: an R package for simulation-based prediction of the number of
loci expected in RADseq and similar genotyping by sequencing approaches. *Molecular Ecology
Resources, 14*(6), 1314-1321. doi:https://doi.org/10.1111/1755-0998.12273

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM,
arXiv:1303.3997. Retrieved from https://ui.adsabs.harvard.edu/abs/2013arXiv1303.3997L
Li, Y-L, & Liu, J-X. (2018). StructureSelector: A web-based software to select and visualize the optimal
number of clusters using multiple methods. *Molecular Ecology Resources, 18*(1), 176-177.
doi:https://doi.org/10.1111/1755-0998.12719
Liu, X, & Fu, Y-X. (2020). Stairway Plot 2: demographic history inference with folded SNP frequency
spectra. *Genome Biology, 21*(1), 280. doi:10.1186/s13059-020-02196-9
Liu, X, Rønhøj Schjøtt, S, Granquist, SM, Rosing-Asvid, A, Dietz, R, Teilmann, J, et al. (2022). Origin
and expansion of the world's most widespread pinniped: Range-wide population genomics of the
harbour seal (*Phoca vitulina*). *Molecular Ecology, 31*(6), 1682-1699.
doi:https://doi.org/10.1111/mec.16365
Majewski, SP, & Ellis, GM. (2022). *Abundance and distribution of harbour seals (Phoca vitulina) in the*
Strait of Georgia, British Columbia; synthesis of the 2014 aerial survey and long-term trends. .
Canadian Science Advisory Secretariat (CSAS).
Malinsky, M, Trucchi, E, Lawson, DJ, & Falush, D. (2018). RADpainter and fineRADstructure:
Population Inference from RADseq Data. *Molecular Biology and Evolution, 35*(5), 1284-1290.
doi:10.1093/molbev/msy023
Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal, 17(1), 3. doi:10.14806/ej.17.1.200
Meek, MH, & Larson, WA. (2019). The future is now: Amplicon sequencing and sequence capture usher
in the conservation genomics era. *Molecular Ecology Resources, 19*(4), 795-803.
doi:https://doi.org/10.1111/1755-0998.12998
Muto, MM, Helker, VT, Delean, BJ, Young, NC, Freed, JC, Angliss, RP, et al. (2021). *Alaska marine*
mammal stock assessments, 2020. . NOAA Tech. Memo. NMFS-AFSC.
NCBI. (2023). SRA Toolkit: National Center for Biotechnology Information (NCBI). Retrieved from
https://github.com/ncbi/sra-tools
Olesiuk, PF. (2010). *An assessment of population trends and abundance of harbour seals (Phoca vitulina)*
in British Columbia. Nanaimo, B.C.: Canadian Science Advisory Secretariat Retrieved from
http://www.dfo-mpo.gc.ca/csas/.
Olsen, MT, Andersen, LW, Dietz, R, Teilmann, J, Härkönen, T, & Siegismund, HR. (2014). Integrating
genetic data and population viability analyses for the identification of harbour seal (*Phoca*
vitulina) populations and management units. *Molecular Ecology, 23*(4), 815-831.
doi:https://doi.org/10.1111/mec.12644
Olsen, MT, Islas, V, Graves, JA, Onoufriou, A, Vincent, C, Brasseur, S, et al. (2017). Genetic population
structure of harbour seals in the United Kingdom and neighbouring waters. *Aquatic*
Conservation: Marine and Freshwater Ecosystems, 27(4), 839-845.
doi:https://doi.org/10.1002/aqc.2760
Paradis, E. (2010). pegas: an R package for population genetics with an integrated–modular approach.
Bioinformatics, 26(3), 419-420. doi:10.1093/bioinformatics/btp696
Patterson, N, Price, AL, & Reich, D. (2006). Population structure and eigenanalysis. *PLoS Genetics,*
2(12), e190. doi:10.1371/journal.pgen.0020190
Peterson, BK, Weber, JN, Kay, EH, Fisher, HS, & Hoekstra, HE. (2012). Double digest RADseq: an
inexpensive method for *de novo* SNP discovery and genotyping in model and non-model species.
PLOS ONE, 7(5), e37135. doi:10.1371/journal.pone.0037135
Pew, J, Muir, PH, Wang, J, & Frasier, TR. (2015). related: an R package for analysing pairwise
relatedness from codominant molecular markers. *Mol Ecol Resour, 15*(3), 557-561.
doi:10.1111/1755-0998.12323
Puechmaille, SJ. (2016). The program structure does not reliably recover the correct population structure
when sampling is uneven: subsampling and new estimators alleviate the problem. *Molecular*
Ecology Resources, 16(3), 608-627. doi:https://doi.org/10.1111/1755-0998.12512
R Core Team. (2023). R: A Language and Environment for Statistical Computing. In. Vienna, Austria.

Rijks, JM, Hoffman, JI, Kuiken, T, Osterhaus, ADME, & Amos, W. (2008). Heterozygosity and
lungworm burden in harbour seals (*Phoca vitulina*). *Heredity, 100*(6), 587-593.
doi:10.1038/hdy.2008.18
Ritland, K. (1996). Estimators for pairwise relatedness and individual inbreeding coefficients. *Genetics*
Research, 67(2), 175-185. doi:10.1017/S0016672300033620
Rochette, NC, Rivera-Colón, AG, & Catchen, JM. (2019). Stacks 2: Analytical methods for paired-end
sequencing improve RADseq-based population genomics. *Molecular Ecology, 28*(21), 4737-
4754. doi:https://doi.org/10.1111/mec.15253
Schaeff, CM, Boness, DJ, & Bowen, WD. (1999). Female distribution, genetic relatedness, and fostering
behaviour in harbour seals, *Phoca vitulina*. *Animal Behaviour, 57*(2), 427-434.
doi:https://doi.org/10.1006/anbe.1998.1001
Shafer, ABA, Peart, CR, Tusso, S, Maayan, I, Brelsford, A, Wheat, CW, et al. (2017). Bioinformatic
processing of RAD-seq data dramatically impacts downstream population genetic inference.
Methods in Ecology and Evolution, 8(8), 907-917. doi:https://doi.org/10.1111/2041-210X.12700
Stanley, HF, Casey, S, Carnahan, JM, Goodman, S, Harwood, J, & Wayne, RK. (1996). Worldwide
patterns of mitochondrial DNA differentiation in the harbor seal (*Phoca vitulina*). *Molecular*
Biology and Evolution, 13(2), 368-382. doi:10.1093/oxfordjournals.molbev.a025596
Steinmetz, K, Murphy, S, Cadhla, OÓ, Carroll, EL, Onoufriou, AB, Russell, DJF, et al. (2023).
Population structure and genetic connectivity reveals distinctiveness of Irish harbour seals (*Phoca*
vitulina) and implications for conservation management. *Aquatic Conservation: Marine and*
Freshwater Ecosystems, 33(2), 160-178. doi:https://doi.org/10.1002/aqc.3909
Swart, JAA, Reijnders, P, J. H. , & Van Delden, W. (1996). Absence of genetic variation in harbor seals
(*Phoca vitulina*) in the dutch Wadden Sea and the British Wash. *Conservation Biology, 10*(1),
289-293.
Tange, O. (2022). GNU Parallel 20220822: Zenodo. Retrieved from
https://doi.org/10.5281/zenodo.7015730
Temte, JL, Bigg, MA, & Wiig, Ø. (1991). Clines revisited: The timing of pupping in the harbour seal
(*Phoca vitulina*). *Journal of Zoology, 224*(4), 617-632. doi:https://doi.org/10.1111/j.1469-
7998.1991.tb03790.x
Thorburn, D-MJ, Sagonas, K, Binzer-Panchal, M, Chain, FJJ, Feulner, PGD, Bornberg-Bauer, E, et al.
(2023). Origin matters: Using a local reference genome improves measures in population
genomics. *Molecular Ecology Resources, n/a*(n/a). doi:https://doi.org/10.1111/1755-0998.13838
Trzcinski, MK. (2020). *Synthesizing scientific knowledge about population dynamics and diet preferences*
of harbour seals, Stellar sea lions and California sea lions, and their impacts on salmon in the
Salish Sea Workshop 2: November 20-21, 2019, Bellingham, WA. .
Waples, RS, & Anderson, EC. (2017). Purging putative siblings from population genetic data sets: a
cautionary view. *Molecular Ecology, 26*(5), 1211-1224. doi:https://doi.org/10.1111/mec.14022
Waples, RS, & Gaggiotti, O. (2006). What is a population? An empirical evaluation of some genetic
methods for identifying the number of gene pools and their degree of connectivity. *Molecular*
Ecology, 15(6), 1419-1439. doi:https://doi.org/10.1111/j.1365-294X.2006.02890.x
Weir, BS, & Cockerham, CC. (1984). Estimating F-statistics for the analysis of population structure.
Evolution, 38(6), 1358-1370. doi:10.2307/2408641
Westlake, RL, & O'Corry-Crowe, GM. (2002). Macrogeographic structure and patterns of genetic
diversity in harbor seals (*Phoca vitulina*) from Alaska to Japan. *Journal of Mammalogy, 83*(4),
1111-1126. doi:10.1644/1545-1542(2002)083<1111:Msapog>2.0.Co;2
Yao, Y, & Ochoa, A. (2023). Limitations of principal components in quantitative genetic association
models for human studies. *eLife, 12*, e79238. doi:10.7554/eLife.79238
840