Regulation of coronavirus nsp15 cleavage specificity by RNA structure

Indranil Salukhe[1], Ryan Choi[2], Wesley Van Voorhis[2], Lynn Barrett[2], Jennifer L. Hyde[1]*

[1]Department of Microbiology, University of Washington School of Medicine, Seattle, WA, United

States of America

[2]Division of Allergy and Infectious Diseases, Department of Medicine, Center for Emerging and

Reemerging Infectious Diseases (CERID), University of Washington School of Medicine,

Seattle, WA, United States of America

*Corresponding author: Jennifer L. Hyde, PhD, Department of Microbiology, University of

Washington School of Medicine, Seattle WA 98109 USA Tel: 206-543-2820, Email:

jlhyde4@uw.edu

Figures 3

Abstract

SARS-CoV-2, the etiologic agent of the COVID-19 pandemic, has had an enduring

impact on global public health. However, SARS-CoV-2 is only one of multiple pathogenic human

coronaviruses (CoVs) to have emerged since the turn of the century. CoVs encode for several

nonstructural proteins (NSPS) that are essential for viral replication and pathogenesis. Among

them is nsp15, a uridine-specific viral endonuclease that is important in evading the host

immune response and promoting viral replication. Despite the established function of nsp15 as

a uridine-specific endonuclease, little is known about other determinants of its cleavage

specificity. In this study we investigate the role of RNA secondary structure in SARS-CoV-2

nsp15 endonuclease activity.

Using a series of *in vitro* endonuclease assays, we observed that thermodynamically

stable RNA structures were protected from nsp15 cleavage relative to RNAs lacking stable

structure. We leveraged the s2m RNA from the SARS 3'UTR as a model for our structural

studies as it adopts a well-defined structure with several uridines, two of which are unpaired and

thus high probably targets for nsp15 cleavage. We found that SARS-CoV-2 nsp15 specifically

cleaves s2m at the unpaired uridine within the GNRNA pentaloop of the RNA. Further

investigation revealed that the position of uridine within the pentaloop also impacted nsp15

cleavage efficiency, suggesting that positioning within the pentaloop is necessary for optimal

presentation of the scissile uridine and alignment within the nsp15 catalytic pocket. Our findings

indicate that RNA secondary structure is an important determinant of nsp15 cleavage and

provides insight into the molecular mechanisms of recognition of RNA by nsp15.

Introduction

Introduction

Coronaviruses (CoVs) are a diverse group of enveloped, positive sense single-stranded

RNA (+ssRNA) viruses within the *Nidovirales* order which infect a wide range of host species

and are associated with mild to severe disease in livestock and humans (1-3). The

Coronaviridae family is composed of four genera: Alpha-, Beta-, Gamma-, and

Deltacoronaviruses (4). Since the turn of the century, several pathogenic CoVs have emerged in

the human population including SARS, MERS, and SARS-CoV-2 with the latter being the

causative agent of the ongoing COVID-19 pandemic (1, 2).

Replication of the large (~30kb) CoV genome is facilitated by multiple nonstructural

proteins (nsp1-16) which are produced from two open reading frames encoded by the first two

thirds of the viral genome. These proteins encode several key functions necessary for synthesis

and post-translational processing of the viral genome, including RNA-dependent RNA

polymerase (nsp12), helicase (nsp13), RNA M7 and 2'-O-methyltransferase (nsp14,

nsp16/nsp10), protease (nsp3, nsp5), and exonuclease (nsp14) activities. Structural proteins

which assemble to form the virion, as well as accessory proteins, are encoded in a set of nested

subgenomic RNAs which are coterminal with the 3' end of the genome.

The viral replication/transcription complex (RTC) is housed within membrane structures

derived from the host endoplasmic reticulum (ER) which are induced by expression of nsp3, 4,

and 6 (5-12). Like other +ssRNA viruses, these double-membrane vesicles (DMVs) provide a

microenvironment that supports efficient viral replication and sequester double-stranded RNA

(dsRNA) replication intermediates from detection by host pathogen recognition receptors

(PRRs) (13, 14). Previous studies have demonstrated a key role for the interferon (IFN)

response in restriction of CoVs. Accordingly, CoVs have evolved several mechanisms to evade

antiviral immunity (e.g., 2'-O-methylation) and encode several proteins which antagonize the

IFN response (nsp1, ORF6) (15). Nsp15 is a uridine-specific endonuclease (endoU) encoded by

CoVs and other Nidoviruses which has been implicated in evasion of host-mediated RNA

sensing and IFN activation (16, 17). While nsp15 has been shown to be non-essential for viral

replication of several CoVs, absence of nsp15 endoU activity (Δnsp15) leads to greater

accumulation of dsRNA replication intermediates which trigger IFN activation in immune

competent cells such as macrophages (16). During CoV infection, dsRNA sensing occurs

primarily via MDA-5, though other RNA sensors (PKR) have also been implicated (16, 18).

While originally described as a uridine-specific endonuclease, nsp15 has been shown to

preferentially cleave pyrimidine-adenine dinucleotides ($U^{\downarrow}A$ or $C^{\downarrow}A$) (19). More recently, viral

targets of nsp15 have been identified in both the polyU tract of tract of 5' negative-sense RNA,

as well as at several sites throughout the CoV genome (17, 19). Although multiple nsp15
cleavage sites have been mapped throughout the viral genome, no other sequence
determinants have yet been identified. Despite the abundance of pyrimidine-adenine
dinucleotides across the SARS-CoV-2 genome, only a fraction of these sites are cleaved by
nsp15, suggesting that other RNA determinants likely affect cleavage specificity of viral RNA.
Earlier *in vitro* studies have shown that SARS nsp15 cleaves a single unpaired uridine in the
stem-loop II motif (s2m) RNA structural element found in the 3'-UTR of the viral genome
suggesting that the structural context in which the scissile uridine is presented may play a role in
determining nsp15 cleavage specificity (20). In this study, we use SARS s2m as a model RNA
to explore the secondary structure requirements for SARS-CoV-2 nsp15 cleavage. Our findings
reveal that thermodynamically stable RNAs are protected from SARS-CoV-2 nsp15 cleavage
and cleaved less efficiently. We show for the first time that SARS-CoV-2 nsp15 preferentially
cleaves unpaired bases. and show that the positioning and sequence context of the scissile
uridine play a role in determining SARS-CoV-2 nsp15 cleavage specificity. This is consistent
with previous studies of SARS nsp15, suggesting that SARS-CoV-2, and CoVs more broadly,
may use RNA structure to regulate nsp15 cleavage specificity. These data shed further light on
requirements for viral endoU target recognition and demonstrate an underappreciated role for
viral RNA structure in evasion of innate immunity.
93

Results

RNA secondary structure modulates nsp15 cleavage efficiency

Nsp15 cleavage sites have previously been mapped in the MHV genome following viral
infection (17, 19). Despite the abundance of pyrimidine-adenine dinucleotides throughout the
genome, MHV nsp15 was found to cleave only a fraction of these sites, suggesting that other
determinants play a role in CoV nsp15 cleavage specificity. We analyzed sequences upstream
and downstream of MHV nsp15 cleavage sites to identify shared motifs that might contribute to
nsp15 recognition and specificity. We reasoned that putative nsp15 recognition motifs present in
the genome of MHV may also be conserved across other betacoronaviruses, including SARS-
CoV-2. Using the MEME-suite XSTREME motif analysis and discovery tool (21) we analyzed
200 nt sequences corresponding to 100 nt upstream and downstream of the MHV nsp15
cleavage sites identified by Ancar et al. Although some motifs were found to be enriched in
sequences surrounding the MHV nsp15 cleavage sites, they were not uniformly conserved
across all cleavage sites. Moreover, many of these motifs were found at other sites in the viral

genome which were not previously identified as nsp15 targets and were not found to be
conserved in SARS-CoV-2 or other human CoVs.

As RNA structure has previously been implicated in nsp15 cleavage specificity (20), we
next compared the underlying predicted secondary structure of the same 200 nt fragments
using LocRNA software (22, 23). We did not identify any conserved structural elements at
nsp15 cleavage sites in MHV nor did we identify conserved structural elements upstream or
downstream of these sites. Despite the absence of any conserved sequence or structural
motifs, we hypothesized that the presence of thermodynamically stable RNA elements would
prevent cleavage of RNA by nsp15.

To test this hypothesis, we used RNAfold (24) to identify regions across the SARS-CoV-
2 genome predicted to have varying degrees of thermodynamic stability (Fig. 1A; RNA 1, RNA
2, RNA 3) (25, 26). We identified representative regions of high, moderate, and low
thermodynamic stability as indicated by their relative ΔG values (Fig. 1B), with more stable
predicted structures having lower ΔG values (27). 100 nt RNAs corresponding to these regions
were synthesized *in vitro*, and cleavage efficiency of these RNAs was assessed using a SARS-
CoV-2 nsp15 endonuclease assay (28) (Fig. 1C and D). Folded RNAs were incubated with
recombinant SARS-CoV-2 nsp15 for varying times (1, 15, 60, and 240 minutes) and cleavage
products analyzed by polyacrylamide gel electrophoresis (Fig. 1C). Cleavage of full-length RNA
was additionally quantified by densitometry by comparing the abundance of full-length RNA at
each time point to a denatured nsp15 control (no cleavage, (-) far-right lane) (Fig. 1D). As we
hypothesized, we observed that RNAs with relatively moderate or low predicted thermodynamic
stability were cleaved more rapidly than RNA with relatively high thermodynamically stability
(Fig. 1C, compare abundance of full-length transcript at 1, 15 and 60 min [red arrow]; Fig. 1D
compare percentage uncleaved RNA). Notably, RNA with relatively low thermodynamic stability
showed rapid cleavage as early as 1 min following the addition of nsp15. While each RNA
yielded different cleavage products owing to the differences in RNA sequence, we observed the
appearance of distinct cleavage products that either decreased (green arrows) or increased
(blue arrows) in abundance over time. These likely represent cleavage intermediates or end
products respectively, suggesting that other structural features or specific sequences also likely
contribute to the higher efficiency of nsp15 cleavage of some RNA sequences over others.
Collectively, these data suggest that RNA secondary structure impacts nsp15-dependent RNA
cleavage.

SARS-CoV-2 nsp15 cleaves unpaired bases in structured RNAs

In vitro biochemical studies by Bhardwaj et al. previously showed that SARS nsp15
specifically cleaves 3' of unpaired uridines in s2m, a conserved structural element found in the
3'UTR of the SARS genome and some other related CoVs (20, 29, 30). In addition to our results
described above (Fig. 1), this study suggests that the context in which the cleaved uridine is
presented may be important for determining the cleavage specificity of nsp15. In the case of
s2m, the two unpaired uridines (U25 and U30) are located in two unstructured loops (Fig. 2A,
highlighted in blue). Structural resolution of s2m has shown that U25, which is located within the
GNRNA sequence of the pentaloop of s2m, is highly disordered relative to other nucleotides,
and that U30 is also moderately disordered (29). Based on previous findings and the availability
of structural information, we posited that s2m would be a tractable model RNA for studying the
determinants of cleavage specificity of SARS-CoV-2 nsp15.

We first sought to determine whether SARS-CoV-2 nsp15 exhibits cleavage specificity
for unpaired uridines in s2m similar to that previously observed for SARS nsp15 (20). To identify
whether nsp15 preferentially cleaves at U25, U30, or both, we generated wild type (wt) SARS
s2m RNA, as well as single and double s2m mutants containing U-to-A substitutions at U25 and
U30 (Fig. 2A) and compared cleavage efficiency of these RNAs via endonuclease assay (Fig.
2A and B). In contrast to wt s2m, cleavage of the double mutant ($s2m_{mut3}$) was noticeably
impaired, consistent with previous studies showing nsp15 cleavage preference for unpaired
uridines. However, in contrast to previous findings with SARS nsp15, we observed that mutation
of U30 (U30A; $s2m_{mut2}$) did not impact RNA cleavage, whereas mutation of U25 (U25A; $s2m_{mut1}$)
did. Although quantitative differences in full-length RNA cleavage were not found to be
statistically significant (Fig 2B), accumulation of cleavage products in the U25A ($s2m_{mut1}$) and
double mutant ($s2m_{mut3}$) was clearly impaired relative to wt and U30A ($s2m_{mut2}$) RNAs.
Interestingly, we also observed accumulation of distinct cleavage products in $s2m_{mut1}$ and
$s2m_{mut3}$. This may indicate that step wise cleavage of RNA and RNA intermediates is altered in
the absence of the scissile U25. However, while mutation of U25 was not predicted to impact
the structure of s2m using RNAfold analysis (Fig. 2C), we cannot rule out the possibility that
mutation of U25 leads to changes in s2m RNA structure which impact nsp15-mediated
cleavage.

Due to the presence of multiple uridines in s2m, accurate identification of cleavage
products and intermediates was challenging. In order to eliminate extraneous cleavage products
to clarify interpretation of endonuclease data and further dissect RNA structural determinants of
cleavage efficiency, we designed truncated s2m RNAs (Δs2m) which lack the base of the stem
and all uridines except U25 and U30 (Fig. 2D). Similar to the experiments described above, we

also engineered Δs2m mutants with U-to-A substitutions at U13 (Δs2m$_{mut1}$) and U18 (Δs2m$_{mut2}$)

(equivalent to U25 and U30 in wt s2m), or both U13 and U18 (Δs2m$_{mut3}$). As with the full-length

s2m RNAs, the free energies and ensemble diversities of the Δs2m mutant RNAs were similar

to each other (Fig. 2C). We analyzed cleavage of wt and mutant Δs2m RNAs over time by

endonuclease assay as described above (Fig 2D and E). In keeping with experiments with full-

length s2m RNA, we observed rapid accumulation of cleavage products in wt Δs2m and

Δs2m$_{mut2}$ RNAs. In contrast, mutation of both U13 and U18 (U25 and U30 equivalent), or U13

(U25 equivalent) alone, resulted in delayed and reduced nsp15 cleavage of RNA. Interestingly,

we still observed some modest cleavage of Δs2m$_{mut1}$ suggesting that while U13 is the

predominant cleavage site in the Δs2m RNA, nsp15 is also able to recognize and cleave at

other sites. While mutation of both U13 and U18 led to a further reduction in nsp15-mediated

cleavage, some cleavage products were still detected in the double mutant, lending further

support to the hypothesis that nsp15 can also cleave non-uridine bases, albeit at lower

efficiency. Indeed, while nsp15 displays a strong preference of cleavage 3' of uridines, previous

work has shown that cleavage at other bases (especially cytosine) also occurs (31).

We hypothesized that RNA secondary structure contributes to nsp15 cleavage specificity

by presenting the scissile uridine in a structural context favorable for nsp15 cleavage. However,

an alternative explanation could be that RNA structure serves to sterically hinder or promote

nsp15-RNA binding. To address whether differences we observed in cleavage of wt and mutant

Δs2m RNAs was due to differences in protein-RNA binding alone, we compared nsp15-RNA

binding affinities using differential radial capillary of action ligand assay (DRaCALA) (32, 33).

Here, 5'-end radiolabeled Δs2m RNAs were incubated with increasing concentrations of

recombinant mutant nsp15 (nsp15 (K290A)), complexes applied to nitrocellulose membrane,

and the fraction of bound vs unbound RNA quantified (Fig. 2F). Nsp15 (K290A) was used to

eliminate endonuclease activity and contains a mutation of one of the conserved catalytic

residues necessary for endoU activity (20, 34-36). We compared binding of wt and mutant

Δs2m RNA, in addition to full-length wt s2m. The latter was chosen as a control, as it is

presently unknown what the minimum length requirement is for RNA recognition and binding by

nsp15. We observed binding of nsp15 (K290A) to wt Δs2m (K$_D$=1.5μM). When we compared

binding of K290A to each mutant Δs2m RNA, we observed only slight decreases in binding

affinity (approximately 1.3-fold; Δs2m$_{mut1}$ K$_D$=2.0μM, Δs2m$_{mut2}$ K$_D$=1.9μM, Δs2m$_{mut3}$ K$_D$=8.5μM),

which were not found to be statistically significant. Notably, we observed a modest increase in

binding affinity of K290A for the full-length s2m RNA (approximately 4-fold; s2m K$_D$=0.3μM)

suggesting that other RNA-protein contacts in longer RNAs may modestly contribute to binding

affinity. As nsp15 exhibited similar binding affinities for all Δs2m RNAs we concluded that

differences in cleavage efficiency of Δs2m RNAs is not due to differences in protein-RNA

binding, but likely driven by positioning of the scissile uridine in these differently structured

RNAs.

Flexible uridine nucleotides in structured pentaloops are susceptible to SARS-CoV-2

nsp15 cleavage

Recent structural studies have suggested that SARS-CoV-2 nsp15 uses a base flipping

mechanism to position the scissile uridine in the catalytic pocket, thus enabling cleavage of

dsRNA (37). The crystal structure of s2m shows that U25 in the GNRNA pentaloop adopts a

similar flipped out orientation (29) suggesting that this mechanism could also account for the

specificity of nsp15 cleavage for U25 in s2m. RNA pentaloops adopt a characteristic structure

which consists of a sheared G-A base pair that closes the pentaloop and induces base stacking

of first N, R, and A nucleotides. Notably, this conformation induces extrusion (or 'flipping') of the

second N (GNR**N**A) in the pentaloop (38, 39). Thus, we postulated that the position of the

scissile uridine in the pentaloop (GAG**U**A) might be necessary to confer the correct orientation

required for nsp15 cleavage. To test whether the relative position of U25 is necessary for

cleavage specificity, we constructed two additional Δs2m mutants in which the position of

scissile uridine has been altered (Fig. 3A, top), and compared cleavage efficiency in the endoU

assay (Fig 3A, bottom). In Δs2m$_{mut4}$ the adjacent guanidine and uridine in the pentaloop were

swapped (GA**UG**A), such that the scissile uridine now sits at the apex of the pentaloop. In

Δs2m$_{mut5}$ the scissile uridine was positioned downstream at the 3' end of the pentaloop and

adjacent to closing GC base pair of the helix (GAGA**U**). The second uridine in the bulge of

Δs2m (U18) was also mutated in both RNAs in order to assess the impact of U13 (U25

equivalent in full-length s2m) cleavage only. We predicted that in Δs2m$_{mut4}$ positioning of the

scissile uridine at the apex would have minimal effect on cleavage but that placement of uridine

adjacent to the helix would place structural constraints on this base such that it would be unable

to adopt the necessary conformation (flipped out) required for positioning within the catalytic

pocket of nsp15. When we compared cleavage of each mutant, we observed that Δs2m$_{mut4}$

(GA**U**GA) was cleaved to completion earlier than both Δs2m$_{mut2}$ (control) and Δs2m$_{mut5}$

(GAGA**U**) (Figure 3B). In contrast, we observed almost no cleavage of Δs2m$_{mut5}$ supporting the

hypothesis that the scissile uridine might be more structurally constrained at this position and

unable to engage with the catalytic pocket of nsp15. From these data, we concluded that uridine

position within the pentaloop influences nsp15 cleavage activity, likely through structural

constraints.

Discussion

Nsp15 is a viral endonuclease conserved across the *Nidovirales* and plays a crucial role

in CoV evasion of host innate immunity. The crystal structure of several CoV nsp15 proteins has

been solved to date, including SARS-CoV-2 (34, 36, 40-44). Structural and biochemical

analyses have revealed key aspects of nsp15 function, including the requirement of Mn^{2+} for

endonuclease activity at neutral pH as well as formation of higher order oligomers (hexamers)

which is also necessary for endoU function (20, 34, 35, 40, 44, 45). Nsp15 primarily functions to

evade innate immune responses by cleaving dsRNA replicative intermediates and preventing

activation of MDA-5, and possibly other RNA sensors in the host cell (16-19, 46-50). Some

studies also suggest that CoV nsp15 may play a role in viral transcriptional regulation during

viral replication (19) and that endoU-independent activities additionally contribute to modulation

of host responses (49, 51-54).

At present, the RNA targets of nsp15 endoU from across the *Coronaviridae* are poorly

defined. This can be attributed in part to the fact that in vitro studies have used different RNA

species (dsRNA vs ssRNA) and different RNA sequences and lengths, making cross-

comparison of these data challenging. Nsp15 has been shown to cleave 3' of pyrimidines, in

particular uridine, and exhibit a preference for weak bases (A or U) 3' of the scissile nucleotide.

However, to date no specific nsp15 recognition motifs have been described. Indeed, our

analysis of previously identified MHV nsp15 cleavage sites (19) failed to identify novel sequence

motifs in and around the cleavage site (data not shown). In the absence of a sequence motif, we

postulate that RNA secondary structure may function to regulate both RNA binding and

cleavage efficiency of nsp15.

To this end, recent work has shown that SARS-CoV-2 nsp15 manipulates the scissile uridine for

cleavage of dsRNA (37). However, the extent to which RNA secondary structure is involved in

modulating SARS-CoV-2 nsp15 cleavage activity remains to be determined. Thus, in this study

we sought to determine the role of RNA structure in the regulation of SARS-CoV-2 nsp15-

mediated cleavage and define specific structural features of RNAs that contribute to this. We

demonstrated that increased thermodynamic stability of RNA is associated with decreased

nsp15 cleavage efficiency (Fig. 1) and postulated that stable secondary structure may contribute

to nsp15 cleavage by two modes: 1) steric hinderance; and 2) inducing optimal positioning of

the scissile uridine in the nsp15-RNA complex. To test this second hypothesis, we used s2m as a model RNA to explore the RNA determinants of nsp15 cleavage specificity.

Using a genetic approach, we first demonstrated that nsp15 exhibits specificity for cleavage of the uridine nucleotide located in the GNRNA pentaloop of s2m (Fig. 2, Fig. 3). This contrasts with previous data from Bhardwaj et al., in which the authors observed preferential cleavage of U30, located in the bulge sequence downstream of the pentaloop (20). It should be noted that these previous studies were performed with SARS nsp15, which shares 88% identity with SARS-CoV-2 nsp15. One explanation for the discrepancy in our data could be structural differences between these proteins which lead to altered cleavage specificity.

Notably, the specificity for cleavage of the pentaloop uridine (GAG$\underline{U}$A) was not due to differences in nsp15-RNA binding (Fig. 2F), thus we postulated that the sequence context of the scissile uridine might be important for this specificity. The characteristic structure adopted by GNRNA pentaloops induces extrusion, or base flipping, of the scissile uridine away from the body of the RNA (38, 39), and this base flipping mechanism has been suggested to be important for correct positioning of uridine in the catalytic pocket of nsp15 (55). Pentaloop motifs have also been implicated in base editing mediated by $\underline{A}$denosine $\underline{D}$eaminases that $\underline{A}$ct on $\underline{R}$NA 2 (ADAR2) (56), though in this instance the GCUMA pentaloop is located downstream of the edited A, and is implicated in recruitment of ADAR2 via interactions between the dsRNA binding motif of ADAR2 and the target RNA. We speculated that if structural conformation of the scissile uridine was important for base recognition and cleavage that altering the position (and conformation) of the uridine base would lead to altered cleavage efficiency. Indeed, repositioning of the scissile uridine to the apex of the pentaloop (Δs2m$_{mut4}$) slightly enhanced cleavage efficiency RNA, whereas repositioning adjacent to the closing G-C base pair (Δs2m$_{mut5}$) significantly reduced RNA cleavage (Fig. 3). The marked decrease in Δs2m$_{mut5}$ cleavage is likely due to altered positioning of the scissile uridine which we predict is no longer extruded from the pentaloop, as well as decreased flexibility of this base which is now structurally constrained by the neighboring G-C base pair. Another possibility is that cleavage efficiency is impacted by altering the sequence context of the scissile uridine. Nsp15 has been shown to preferentially cleave uridines flanked by a weak base (A/U) which is found in wt Δs2m and Δs2m$_{mut2}$ (U$^{\downarrow}$A) but is changed to a strong base in Δs2m$_{mut5}$ (U$^{\downarrow}$C). However, we observed a modest increase in cleavage efficiency of Δs2m$_{mut4}$ which also has a strong base at the 3' position (U$^{\downarrow}$G) and lends support to the hypothesis that the structural context of the scissile uridine, not sequence context, may be more relevant in dictating cleavage specificity and efficiency.

Collectively, our data suggests that RNA secondary structure plays two distinct roles in

determining nsp15 cleavage specificity. First, thermodynamically stable structures likely

sterically hinder nsp15-RNA interactions, thus preventing engagement and cleavage of some

RNA sequences. Second, susceptible bases are presented in a structural context which

facilitates optimal positioning of the scissile nucleotide in the catalytic pocket of nsp15 (e.g.,

extrusion of the scissile U25 in the GNRNA pentaloop). However, several questions remain

unanswered, most notably, what are the biologically relevant targets (host and viral) of nsp15

cleavage during CoV infection and are viral targets conserved across different CoVs? Two

studies to date have identified cleavage sites within the positive-sense genomic RNA and 3'

polyU tails of the negative-sense RNA of MHV (17, 19). Despite this, no recognition motif or

other regulatory element associated with nsp15 cleavage has been identified. While the in vitro

assays performed here and in other studies provide a tractable system for dissecting the

molecular mechanism of nsp15-RNA interactions and cleavage determinants, nsp15 forms part

of a larger replication-transcription complex (RTC) which contains other viral proteins and host

factors. How nsp15 protein-protein interactions and RNA interactions with other replicase

proteins (e.g., nsp12, nsp14, nsp16) impacts nsp15 cleavage of RNA still remains unsolved.

Indeed, recent models of the RTC suggest the hexameric structure of nsp15 provides the

central structure and arrangement of a hexameric RTC (57). When nsp15 is bound in the RTC

structure, there may be differences in its cleavage of viral RNAs due to its 3-D interaction with

the nascent RNA as opposed to when nsp15 is free from the RTC as was the case in this study.

The importance of examining nsp15-RNA interactions in their relevant biological context is

further highlighted by recent studies showing that the requirement for Mn^{2+} and the structural

conformation of nsp15 is highly dependent on the pH of the environment (58). Given that the

redox state of the viral RTC of +ssRNA viruses is altered during replication, the conditions under

which nsp15 endoU activity is examined is highly relevant. To address this our future studies will

explore the impact of pH and redox state on nsp15 structure and function in vitro.

Overall, our findings suggest a role for RNA secondary structure in regulation of nsp15

cleavage. Identification of bona fide nsp15 cleavage sites in SARS-CoV-2 RNA during

replication and examination of how perturbation of these sites impacts the outcome of

replication and pathogenesis will provide key insight into the broader roles for nsp15 endoU

activity during infection (e.g., viral transcriptional/translational regulation, host translation).

Furthermore, defining key structural features of RNAs which are resistant or susceptible to

nsp15 cleavage may provide avenues for development of antiviral therapeutics.

Acknowledgements

The authors declare that they have no conflict of interest.

Materials and Methods

Generation of SARS-CoV-2 RNA segments: 100nt segments of SARS-CoV-2 genome

were amplified using the primers listed in Table 1. Amplicons were purified using Mag-Bind

Total Pure NGS beads (Omega Bio-Tek). 1.5 x bead volume relative to amplification reactions

was used and purified as per manufacturer's instructions. Purified amplicons were subcloned

into a vector containing the HDV ribozyme (59). These constructs were linearized and purified

prior to transcription.

Approximately 5 pmol of purified DNA was used as a template for T7 transcription

reactions. RNAs were transcribed using the HiScribe T7 High Yield RNA Synthesis Kit (New

England Biolabs). Reactions were set up as per manufacturer's instructions with the addition of

4 units of Murine RNase Inhibitor (NEB) and 5% DMSO. Transcriptions were incubated at 42^{o}C

overnight after which, the reactions were treated with DNaseI (NEB) as per manufacturer's

instructions. DNase-treated transcriptions were then purified using Mag-Bind Total Pure NGS

beads as described above.

Transcriptions were gel purified by running samples on urea gels. 15% TBE-Urea gels

were made using SequaGel UreaGel 29:1 Denaturing Gel System (National Diagnostics) Urea

was added to the samples to a final concentration of 4M before running on the gels at 200V for

90 minutes. Gels were briefly stained with 0.02% Methylene Blue in TBE until ladder was

visible. Gels were imaged on Gel Doc XR+(Bio-Rad), and bands corresponding to 100nt

fragments were cut out and shredded by centrifugation. RNAs were eluted overnight on rotator

in 3x volume of elution buffer (10mM Tris-HCl, 1mM EDTA, 300mM NaCl, 0.1% SDS) relative to

gel weight. Gel slurries were applied to cellulose acetate columns and frozen at -80^{o}C for 10

minutes. Samples were centrifuged at 10,000x g for 3 minutes. Eluate was adjusted up to 600μL

with nuclease-free water and purified by acid phenol-chloroform extraction.

Table 1

RNA	Forward Primer	Reverse Primer
RNA 1	CATGAGCTTA AGTAATACGA CTCACTATAG GCTACTAACA ATGCCAT	CATGAGGCTA GCAAGATTGT GTCCGCTTAA AA

| RNA 2 | CATGAGCTTA AGTAATACGA CTCACTATAG ACCTGAGCAT AGTCTTG | CATGAGGCTA GCGAGAACAC ACAGCCTC |
| RNA 3 | CATGAGCTTA AGTAATACGA CTCACTATAG TGAAGTGCTG TCTGAC | CATGAGGCTA GCGTTACACG ATAACCAGTA AAG |

Endonuclease Assay: Indicated amounts of RNAs were diluted in folding buffer (50mM

Tris-HCl, 50mM KCl, 5mM $MgCl_2$, 5mM DTT). RNAs were incubated at $95^{\circ}C$ for 10 minutes

then cooled to $25^{\circ}C$ at a rate of $0.1^{\circ}C$/sec. Purified nsp15 was diluted in assay buffer (50mM

Tris-HCl, 50mM KCl, 50mM $MnCl_2$, 5mM $MgCl_2$, 5mM DTT) to the appropriate concentration for

indicated nsp15:RNA ratios. Recombinant nsp15 was expressed and purified as previously

described (60). To denature nsp15, the protein was incubated at $95^{\circ}C$ for 5 minutes. Folded

RNA was aliquoted into individual tubes (1 tube/timepoint), and diluted nsp15 was mixed in

equal amounts to each RNA simultaneously. Reactions were arrested at indicated timepoints

with a final concentration of 20mM EDTA before flash freezing in liquid nitrogen. Denatured

nsp15 reactions were incubated for 240 minutes.

TBE-Urea gels were made using SequaGel UreaGel 29:1 Denaturing Gel Syst

em (National Diagnostics). 15% or 22.5% gels were poured as indicated in each experiment.

RNA samples were thawed and mixed with an equal volume of 2x RNA Loading Dye (NEB) and

loaded onto the gels. Gels were run at 200V for 1 hour in 1x TBE. Gels were stained with SYBR

Green (ThermoFisher) diluted 1:5000 in TBE for 30 minutes while shaking and imaged on Gel

Doc XR+(Bio-Rad). Full-length bands for at each time point were calculated by densitometry

using Image Lab software (Bio-Rad). Density of bands were normalized to denatured nsp15

control for each reaction.

DRaCALA: DRaCALAs were performed as described previously (32). Briefly, RNAs

were dephosphorylated and 5' radiolabeled with 30µCi ATP. Radiolabeled RNAs were set up in

reactions with the following components: DRaCALA buffer (50mM HEPES, 100mM KCl), 5mM

$MgCl_2$, 50ng/µL yeast tRNA, 1mM DTT, 5% glycerol. RNAs were heated to $95^{\circ}C$ for 2 minutes

and cooled at RT for 10 minutes. Serial dilutions of nsp15 K290A were added to the reactions

which were incubated for 15 minutes at RT and spotted onto a 0.45µm nitrocellulose

membrane. Blots were exposed to a phosphor screen overnight and imaged on Sapphire

Molecular Imager (Azure Biosystems). Binding was measured by densitometry as previously

described (33) using Fiji.

FIGURE LEGENDS

Figure 1. Secondary structure protects RNA from cleavage by SARS-CoV-2 nsp15. A. Predicted minimum free energy structures (RNAfold) of 100 nt RNAs from the SARS-CoV-2 genome of varying thermodynamic stability (high, medium, and low). Unpaired uridines are highlighted in blue, and paired uridines are outlined in blue. All RNA structures were predicted in RNAfold and designed in RNA2Drawer (27, 61). **B.** The locations within the SARS-CoV-2 genome are indicated for each structure, and thermodynamic stability is indicated as -ΔG. Ensemble diversity is listed as predicted by RNAfold. **C.** Endonuclease assays of RNAs from (A). Full-length (uncut) RNAs are indicated by the red arrows. Diminishing cleavage products (cleavage intermediates) are indicated by green arrows. Accumulating cleavage products (cleavage end products) indicated by blue arrows. Representative images from one of three total experiments are shown. **D.** Quantitation of data from (C). The percentage of full-length RNA remaining was measured by densitometry. Most thermodynamically stable RNA (RNA 1) was cleaved least rapidly relative to other RNAs. Percentage of uncut RNA was calculated by normalizing to a denatured nsp15 control (indicated by (-) in the far-right lane). Area under the curve (AUC) was calculated for each RNA and one-way ANOVA with multiple comparisons was performed on the AUC. Data represents three independent experiments.

Figure 2. SARS-CoV-2 nsp15 cleaves structured RNAs at specific sites. A. Endonuclease assays of full-length s2m mutant RNAs. Full-length bands indicated at the top of the gel with cleavage products below. Representative images from one experiment. **B.** Percent of full-length RNA remaining as measured by densitometry. Percentage of uncut RNA was calculated by normalizing to a denatured nsp15 control (indicated by (-) in the far-right lane). RNAs with uridines present in the pentaloop (s2m and s2m$_{mut2}$) cleaved more efficiently than s2m$_{mut}$1 and s2m$_{mut3}$. AUC was calculated for each RNA and one-way ANOVA with multiple comparisons was performed on the AUC. AUC of s2m mutant RNAs are compared to wt s2m. Data represents three independent experiments. **C.** Thermodynamic stability and ensemble diversity of s2m and mutant RNAs indicated as predicted by RNAfold. **D.** Endonuclease assays of modified s2m RNAs. Full-length bands indicated at the top of the gel with cleavage products below. Representative images from one experiment. **E.** Percent of full-length RNA remaining as measured by densitometry. Percentage of uncut RNA was calculated by normalizing to a denatured nsp15 control (indicated by (-) in the far-right lane). RNAs with uridines present in the pentaloop (Δs2m and Δs2m$_{mut2}$) cleaved more efficiently than Δs2m$_{mut}$1 and Δs2m$_{mut3}$. AUC was calculated for each RNA and one-way ANOVA with multiple comparisons was performed on the

AUC. AUC of Δs2m mutant RNAs are compared to wt Δs2m. Data represents three independent

experiments. **F.** Nsp15 binding curves of Δs2m RNAs as compared to wt s2m. Δs2m RNA

binding curves overlap with each other while wt s2m curve is shifted left indicating improved

binding with nsp15. Dissociation constant (K_D) was calculated for each RNA and one-way

ANOVA with multiple comparisons was performed on the K_D. AUC of wt full-length s2m and

Δs2m mutant RNAs are compared to wt Δs2m. Data represents three independent experiments.

Figure 3. Uracil position in loop structures impacts SARS-CoV-2 nsp15 cleavage

efficiency. A. Endonuclease assays of s2m pentaloop mutant RNAs. Full-length bands

indicated at the top of the gel with cleavage products below. Representative images from one

experiment. **B.** Thermodynamic stability and ensemble diversity of s2m and mutant RNAs

indicated as predicted by RNAfold. **C.** Percent of full-length RNA remaining as measured by

densitometry. Percentage of uncut RNA was calculated by normalizing to a denatured nsp15

control (indicated by (-) in the far-right lane). RNAs with uridines positioned closer to the apex of

the pentaloop cleaved more rapidly than Δs2m$_{mut5}$ in which the uridine is at the end of the

pentaloop. AUC was calculated for each RNA and one-way ANOVA with multiple comparisons

was performed on the AUC. Data represents three independent experiments.

References

1.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak
associated with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270-3.
2.   Helmy YA, Fawzy M, Elaswad A, Sobieh A, Kenney SP, Shehata AA. The COVID-19
Pandemic: A Comprehensive Review of Taxonomy, Genetics, Epidemiology, Diagnosis,
Treatment, and Control. J Clin Med. 2020;9(4).
3.   Weiss SR, Navas-Martin S. Coronavirus pathogenesis and the emerging pathogen severe
acute respiratory syndrome coronavirus. Microbiol Mol Biol Rev. 2005;69(4):635-64.
4.   Luk HKH, Li X, Fung J, Lau SKP, Woo PCY. Molecular epidemiology, evolution and
phylogeny of SARS coronavirus. Infect Genet Evol. 2019;71:21-30.
5.   Ricciardi S, Guarino AM, Giaquinto L, Polishchuk EV, Santoro M, Di Tullio G, et al. The
role of NSP6 in the biogenesis of the SARS-CoV-2 replication organelle. Nature.
2022;606(7915):761-8.
6.   Oudshoorn D, Rijs K, Limpens R, Groen K, Koster AJ, Snijder EJ, et al. Expression and
Cleavage of Middle East Respiratory Syndrome Coronavirus nsp3-4 Polyprotein Induce the
Formation of Double-Membrane Vesicles That Mimic Those Associated with Coronaviral RNA
Replication. mBio. 2017;8(6).
7.   Lundin A, Dijkman R, Bergstrom T, Kann N, Adamiak B, Hannoun C, et al. Targeting
membrane-bound viral RNA synthesis reveals potent inhibition of diverse coronaviruses
including the middle East respiratory syndrome virus. PLoS Pathog. 2014;10(5):e1004166.

8.   Angelini MM, Akhlaghpour M, Neuman BW, Buchmeier MJ. Severe acute respiratory
syndrome coronavirus nonstructural proteins 3, 4, and 6 induce double-membrane vesicles.
mBio. 2013;4(4).

9.   Gosert R, Kanjanahaluethai A, Egger D, Bienz K, Baker SC. RNA replication of mouse
hepatitis virus takes place at double-membrane vesicles. J Virol. 2002;76(8):3697-708.

10.   van der Meer Y, Snijder EJ, Dobbe JC, Schleich S, Denison MR, Spaan WJ, et al.
Localization of mouse hepatitis virus nonstructural proteins and RNA synthesis indicates a role
for late endosomes in viral replication. J Virol. 1999;73(9):7641-57.

11.   Shi ST, Schiller JJ, Kanjanahaluethai A, Baker SC, Oh JW, Lai MM. Colocalization and
membrane association of murine hepatitis virus gene 1 products and De novo-synthesized viral
RNA in infected cells. J Virol. 1999;73(7):5957-69.

12.   Prentice E, Jerome WG, Yoshimori T, Mizushima N, Denison MR. Coronavirus replication
complex formation utilizes components of cellular autophagy. J Biol Chem.
2004;279(11):10136-41.

13.   Overby AK, Popov VL, Niedrig M, Weber F. Tick-borne encephalitis virus delays
interferon induction and hides its double-stranded RNA in intracellular membrane vesicles. J
Virol. 2010;84(17):8470-83.

14.   Knoops K, Kikkert M, Worm SH, Zevenhoven-Dobbe JC, van der Meer Y, Koster AJ, et al.
SARS-coronavirus replication is supported by a reticulovesicular network of modified
endoplasmic reticulum. PLoS biology. 2008;6(9):e226.

15.   Kamitani W, Narayanan K, Huang C, Lokugamage K, Ikegami T, Ito N, et al. Severe acute
respiratory syndrome coronavirus nsp1 protein suppresses host gene expression by promoting
host mRNA degradation. Proc Natl Acad Sci U S A. 2006;103(34):12885-90.

16.   Deng X, Hackbart M, Mettelman RC, O'Brien A, Mielech AM, Yi G, et al. Coronavirus
nonstructural protein 15 mediates evasion of dsRNA sensors and limits apoptosis in
macrophages. Proc Natl Acad Sci U S A. 2017;114(21):E4251-E60.

17.   Hackbart M, Deng X, Baker SC. Coronavirus endoribonuclease targets viral polyuridine
sequences to evade activating host sensors. Proc Natl Acad Sci U S A. 2020;117(14):8094-103.

18.   Kindler E, Gil-Cruz C, Spanier J, Li Y, Wilhelm J, Rabouw HH, et al. Early endonuclease-
mediated evasion of RNA sensing ensures efficient coronavirus replication. PLoS Pathog.
2017;13(2):e1006195.

19.   Ancar R, Li Y, Kindler E, Cooper DA, Ransom M, Thiel V, et al. Physiologic RNA targets
and refined sequence specificity of coronavirus EndoU. RNA. 2020;26(12):1976-99.

20.   Bhardwaj K, Sun J, Holzenburg A, Guarino LA, Kao CC. RNA recognition and cleavage by
the SARS coronavirus endoribonuclease. J Mol Biol. 2006;361(2):243-56.

21.   Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. Nucleic Acids Res.
2015;43(W1):W39-49.

22.   Will S, Joshi T, Hofacker IL, Stadler PF, Backofen R. LocARNA-P: Accurate boundary
prediction and improved detection of structural RNAs. Rna. 2012;18(5):900-14.

23.   Raden M, Ali SM, Alkhnbashi OS, Busch A, Costa F, Davis JA, et al. Freiburg RNA tools: a
central online resource for RNA-focused research and teaching. Nucleic Acids Research.
2018;46(W1):W25-W9.

24.   Hofacker IL. Vienna RNA secondary structure server. Nucleic Acids Res.
2003;31(13):3429-31.

25.   Tavares RCA, Mahadeshwar G, Wan H, Huston NC, Pyle AM. The global and local
distribution of RNA structure throughout the SARS-CoV-2 genome. J Virol. 2021;95(5).
26.   Andrews RJ, O'Leary CA, Tompkins VS, Peterson JM, Haniff HS, Williams C, et al. A map
of the SARS-CoV-2 RNA structurome. NAR Genom Bioinform. 2021;3(2):lqab043.
27.   Gruber AR, Lorenz R, Bernhart SH, Neubock R, Hofacker IL. The Vienna RNA websuite.
Nucleic Acids Res. 2008;36(Web Server issue):W70-4.
28.   Choi R, Zhou M, Shek R, Wilson JW, Tillery L, Craig JK, et al. High-throughput screening of
the ReFRAME, Pandemic Box, and COVID Box drug repurposing libraries against SARS-CoV-2
nsp15 endoribonuclease to identify small-molecule inhibitors of viral activity. PLoS One.
2021;16(4):e0250019.
29.   Robertson MP, Igel H, Baertsch R, Haussler D, Ares M, Jr., Scott WG. The structure of a
rigorously conserved RNA element within the SARS virus genome. PLoS biology. 2005;3(1):e5.
30.   Tengs T, Kristoffersen AB, Bachvaroff TR, Jonassen CM. A mobile genetic element with
unknown function found in distantly related viruses. Virol J. 2013;10:132.
31.   Frazier MN, Dillard LB, Krahn JM, Perera L, Williams JG, Wilson IM, et al.
Characterization of SARS2 Nsp15 nuclease activity reveals it's mad about U. Nucleic Acids Res.
2021;49(17):10136-49.
32.   Patel DK, Gebbie MP, Lee VT. Assessing RNA interactions with proteins by DRaCALA.
Methods in enzymology. 2014;549:489-512.
33.   Orr MW, Lee VT. Differential Radial Capillary Action of Ligand Assay (DRaCALA) for High-
Throughput Detection of Protein-Metabolite Interactions in Bacteria. Methods Mol Biol.
2017;1535:25-41.
34.   Bhardwaj K, Palaninathan S, Alcantara JM, Yi LL, Guarino L, Sacchettini JC, et al.
Structural and functional analyses of the severe acute respiratory syndrome coronavirus
endoribonuclease Nsp15. J Biol Chem. 2008;283(6):3655-64.
35.   Ricagno S, Egloff MP, Ulferts R, Coutard B, Nurizzo D, Campanacci V, et al. Crystal
structure and mechanistic determinants of SARS coronavirus nonstructural protein 15 define an
endoribonuclease family. Proc Natl Acad Sci U S A. 2006;103(32):11892-7.
36.   Xu X, Zhai Y, Sun F, Lou Z, Su D, Xu Y, et al. New antiviral target revealed by the
hexameric structure of mouse hepatitis virus nonstructural protein nsp15. J Virol.
2006;80(16):7909-17.
37.   Frazier MN, Wilson IM, Krahn JM, Butay KJ, Dillard LB, Borgnia MJ, et al. Flipped over U:
structural basis for dsRNA cleavage by the SARS-CoV-2 endoribonuclease. Nucleic Acids Res.
2022.
38.   Scharpf M, Sticht H, Schweimer K, Boehm M, Hoffmann S, Rosch P. Antitermination in
bacteriophage lambda. The structure of the N36 peptide-boxB RNA complex. European journal
of biochemistry / FEBS. 2000;267(8):2397-408.
39.   Legault P, Li J, Mogridge J, Kay LE, Greenblatt J. NMR structure of the bacteriophage
lambda N peptide/boxB RNA complex: recognition of a GNRA fold by an arginine-rich motif.
Cell. 1998;93(2):289-99.
40.   Joseph JS, Saikatendu KS, Subramanian V, Neuman BW, Buchmeier MJ, Stevens RC, et al.
Crystal structure of a monomeric form of severe acute respiratory syndrome coronavirus
endonuclease nsp15 suggests a role for hexamerization as an allosteric switch. J Virol.
2007;81(12):6700-8.

41.   Zhang L, Li L, Yan L, Ming Z, Jia Z, Lou Z, et al. Structural and Biochemical
Characterization of Endoribonuclease Nsp15 Encoded by Middle East Respiratory Syndrome
Coronavirus. J Virol. 2018;92(22).
42.   Zheng A, Shi Y, Shen Z, Wang G, Shi J, Xiong Q, et al. Insight into the evolution of
nidovirus endoribonuclease based on the finding that nsp15 from porcine Deltacoronavirus
functions as a dimer. J Biol Chem. 2018;293(31):12054-67.
43.   Kim Y, Jedrzejczak R, Maltseva NI, Wilamowski M, Endres M, Godzik A, et al. Crystal
structure of Nsp15 endoribonuclease NendoU from SARS-CoV-2. Protein Sci. 2020;29(7):1596-
605.
44.   Pillon MC, Frazier MN, Dillard LB, Williams JG, Kocaman S, Krahn JM, et al. Cryo-EM
structures of the SARS-CoV-2 endoribonuclease Nsp15 reveal insight into nuclease specificity
and dynamics. Nature communications. 2021;12(1):636.
45.   Guarino LA, Bhardwaj K, Dong W, Sun J, Holzenburg A, Kao C. Mutational analysis of the
SARS virus Nsp15 endoribonuclease: identification of residues affecting hexamer formation. J
Mol Biol. 2005;353(5):1106-17.
46.   Zhao J, Sun L, Zhao Y, Feng D, Cheng J, Zhang G. Coronavirus Endoribonuclease Ensures
Efficient Viral Replication and Prevents Protein Kinase R Activation. J Virol. 2020.
47.   Volk A, Hackbart M, Deng X, Cruz-Pulido Y, O'Brien A, Baker SC. Coronavirus
Endoribonuclease and Deubiquitinating Interferon Antagonists Differentially Modulate the Host
Response during Replication in Macrophages. J Virol. 2020;94(11).
48.   Deng X, Buckley AC, Pillatzki A, Lager KM, Faaberg KS, Baker SC. Inactivating Three
Interferon Antagonists Attenuates Pathogenesis of an Enteric Coronavirus. J Virol. 2020;94(17).
49.   Liu X, Fang P, Fang L, Hong Y, Zhu X, Wang D, et al. Porcine deltacoronavirus nsp15
antagonizes interferon-beta production independently of its endoribonuclease activity.
Molecular immunology. 2019;114:100-7.
50.   Shokri S, Mahmoudvand S, Taherkhani R, Farshadpour F. Modulation of the immune
response by Middle East respiratory syndrome coronavirus. J Cell Physiol. 2019;234(3):2143-51.
51.   Lei Y, Moore CB, Liesman RM, O'Connor BP, Bergstralh DT, Chen ZJ, et al. MAVS-
mediated apoptosis and its inhibition by viral proteins. PLoS One. 2009;4(5):e5466.
52.   Bhardwaj K, Liu P, Leibowitz JL, Kao CC. The coronavirus endoribonuclease Nsp15
interacts with retinoblastoma tumor suppressor protein. J Virol. 2012;86(8):4294-304.
53.   Athmer J, Fehr AR, Grunewald ME, Qu W, Wheeler DL, Graepel KW, et al. Selective
Packaging in Murine Coronavirus Promotes Virulence by Limiting Type I Interferon Responses.
mBio. 2018;9(3).
54.   Wu Y, Zhang H, Shi Z, Chen J, Li M, Shi H, et al. Porcine Epidemic Diarrhea Virus nsp15
Antagonizes Interferon Signaling by RNA Degradation of TBK1 and IRF3. Viruses. 2020;12(6).
55.   Frazier MN, Wilson IM, Krahn JM, Butay KJ, Dillard LB, Borgnia MJ, et al. Flipped over U:
structural basis for dsRNA cleavage by the SARS-CoV-2 endoribonuclease. Nucleic Acids Res.
2022;50(14):8290-301.
56.   Stefl R, Allain FH. A novel RNA pentaloop fold involved in targeting ADAR2. RNA.
2005;11(5):592-7.
57.   Perry JK, Appleby TC, Bilello JP, Feng JY, Schmitz U, Campbell EA. An atomistic model of
the coronavirus replication-transcription complex as a hexamer assembled around nsp15. J Biol
Chem. 2021;297(4):101218.

58.   Godoy AS, Nakamura AM, Douangamath A, Song Y, Noske GD, Gawriljuk VO, et al.
Allosteric regulation and crystallographic fragment screening of SARS-CoV-2 NSP15
endoribonuclease. Nucleic Acids Res. 2023.
59.   Avis JM, Conn GL, Walker SC. Cis-acting ribozymes for the production of RNA in vitro
transcripts with defined 5' and 3' ends. Methods Mol Biol. 2012;941:83-98.
60.   Deng X, Baker SC. An "Old" protein with a new story: Coronavirus endoribonuclease is
important for evading host antiviral defenses. Virology. 2018;517:157-63.
61.   Johnson PZ, Kasprzak WK, Shapiro BA, Simon AE. RNA2Drawer: geometrically strict
drawing of nucleic acid structures with graphical structure editing and highlighting of
complementary subsequences. RNA Biol. 2019;16(12):1667-71.
618

1A

RNA 1   RNA 2   RNA 3

1B

RNA	SARS-CoV-2 gene	Thermodynamic Stability	ΔG of Minimum Free Energy (kcal/mol)	Ensemble Diversity
1	*nsp3*	High	-27.8	4.81
2	*nsp2*	Moderate	-23.4	25.80
3	*nsp13*	Low	-14.2	39.93

1C

RNA 1   RNA 2   RNA 3

100nt

Time (minutes)   1 15 60 240   1 15 60 240   1 15 60 240

1D

3A

3B

RNA	ΔG of Minimum Free Energy (kcal/mol)	Ensemble Diversity
Δs2m$_{mut2}$	-7.9	1.14
Δs2m$_{mut4}$	-7.9	1.13
Δs2m$_{mut5}$	-6.9	2.32

3C