Title: Transposable elements impact the population divergence of rice blast fungus

Magnaporthe oryzae

Lianyu Lin[a, #], Ting Sun[a, #], Jiayuan Guo[a,b], Lili Lin[a], Meilian Chen[b], Zhe Wang[a], Jiandong Bao[c],

Justice Norvienyeku[d], Dongmei Zhang[a], Yijuan Han[b], Guodong Lu[a], Christopher Rensing[e],

Huakun Zheng[a*], Zhenhui Zhong[a*], Zonghua Wang[a,b *]

[a] State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, College of Life

Science, Fujian Agriculture and Forestry University, Fuzhou 350002, China.

[b] Fuzhou Institute of Oceanography, Minjiang University, Fuzhou 350108, China.

[c] State Key Laboratory for Managing Biotic and Chemical Treats to the Quality and Safety of

Agro-Products, Institute of Plant Protection and Microbiology, Zhejiang Academy of Agricultural

Sciences, Hangzhou 310021, China.

[d] Key Laboratory of Green Prevention and Control of Tropical Plant Diseases and Pests, Ministry

of Education, College of Plant Protection, Hainan University, Haikou 570228, China.

[e] Institute of Environmental Microbiology, College of Resource and Environment, Fujian

Agriculture and Forestry University, Fuzhou 350002, China.

# These authors contributed to this work equally.

* To whom correspondence should be addressed. Email: wangzh@fafu.edu.cn (Z.W.),

zhenhuizhong@gmail.com (Z.Z.), huakunzheng@163.com (H.Z.)

ABSTRACT

Dynamic transposition of transposable elements (TEs) in fungal pathogens have significant impact on genome stability, gene expression, and virulence to the host. In *Magnaporthe oryzae*, genome plasticity resulting from TE insertion is a major driving force leading to the rapid evolution and diversification of this fungus. Despite their importance in *M. oryzae* population evolution and divergence, our understanding of TEs in this context remains limited. Here we conducted a genome-wide analysis of TE transposition dynamics in the 11 most abundant TE families in *M. oryzae* populations. Our results show that these TEs have specifically expanded in recently isolated *M. oryzae* rice populations, with the presence/absence polymorphism of TE insertions highly concordant with population divergence on Geng/*Japonica* and Xian/*Indica* rice cultivars. Notably, the genes targeted by clade-specific TEs showed clade-specific expression patterns and are involved in the pathogenic process, suggesting a transcriptional regulation of TEs on targeted genes. Our study provides a comprehensive analysis of TEs in *M. oryzae* populations and demonstrates a crucial role of recent TE bursts in adaptive evolution and diversification of the *M. oryzae* rice-infecting lineage.

IMPORTANCE

M. oryzae is the causal agent of the destructive blast disease, which caused massive loss of yield annually worldwide. The fungus diverged into distinct clades during adaptation toward two rice subspecies, Xian/indica and Geng/japonica. Although the role of TEs in the adaptive evolution was well established, mechanisms underlying how TEs promote the population divergence of *M. oryzae* remains largely unknown. In this study, we reported that TEs shape the population divergence of *M. oryzae* by differentially regulating gene expression between Xian/*Indica*-infecting and Geng/*Japonica*-infecting populations. Our results revealed a TE insertion mediated gene expression adaption that led to the divergence of *M. oryzae* population infecting different rice subspecies.

INTRODUCTION

Rice blast disease, caused by the ascomycete filamentous fungus *Magnaporthe oryzae* (syn: *Pyricularia oryzae*), poses a significant threat to rice production worldwide, resulting in annual yield losses of 10-30% (1, 2). The deployment of resistant rice varieties is the most cost-effective and environmentally friendly strategy for controlling rice blast disease. However, the effectiveness of such resistance can rapidly be diminished due to rapid mutation accumulation in avirulence genes (3, 4). Therefore, it is crucial to uncover the mechanisms by which *M. oryzae* rapidly evolves and evades the rice immune system.

Based on the two-speed genome model, filamentous fungi genomes had been shown to display a bipartite architecture comprising of a gene-rich compartment, which evolves slowly containing core genes encoding essential functions and metabolisms, and a repeat-rich compartment, which evolves rapidly containing important virulence effectors involved in pathogenicity (5-10). To avoid recognition by the plant immune system, some pathogen effectors such as avirulence genes were shown to be influenced or silenced by transposon elements (9, 11). Transposable elements (TEs) make up over 10% of the *M. oryzae* genome (12, 13). Recent evidence has revealed that TEs can be inserted in or around important effectors and alter the virulence spectrum of *M. oryzae*. For example, the insertion of POT3 into the promoter region of AVR-Pita in *M. oryzae* led to the acquisition of virulence towards the resistant rice cultivar Yashiro-mochi (14). TEs are also able to affect gene expression networks, and TE-dependent transcriptional regulation of some essential effectors can facilitate the pathogen's transition in its life cycle. For instance, in *Phytophthora sojae*, the avirulence genes *PsAvr1a*, *1b*, and *3a/5* were found to be transcriptionally inactive due to TE insertions in their promoter or 3' UTR regions (15).

These studies have demonstrated the crucial role of TE-mediated genomic variations in pathogen adaptation. However, previous investigations have mainly focused on the impact of TEs on specific effectors or secreted proteins (13, 16-20). To gain a better

83   understanding on the functions of TEs in the complex *M. oryzae*-rice pathosystem, we

84   conducted a comprehensive analysis of TE insertion polymorphisms in 275 *M. oryzae*

85   isolates (176 rice isolates and 99 non-rice isolates; **Table S1**), and systematically

86   investigated the roles of TEs in the regulation of gene expression and population

87   divergence in the *M. oryzae* rice population.

RESULTS

Recent large-scale TE bursts in the *M. oryzae* genome

To assess the activity of transposable elements (TEs) on the M. oryzae genome, we

estimated the insertion time of long terminal repeat (LTR)-retrotransposons (LTR-RTs)

by measuring the genetic distance between their 5' and 3' LTRs, which were identical

at the time of TE insertion and gradually accumulated mutations over time. Using a de

novo method that is based on the structure of LTR-RTs, we identified 1,129 intact

LTR-RTs in seven near chromosomal-level *M. oryzae* rice isolates, including 70-15,

Guy11, FJ81278, FJ98099, FJ72ZC7-77, AV1-1-1, and Sar-2-20-1 (12, 13, 21).

Surprisingly, 91.7% (1,036/1,129) of the LTR-RTs showed extremely low levels of

divergence between their LTR pairs (over 99% identity), and 69.4% (784/1,129) of

them possessed identical LTR pairs, indicating that these LTR-RTs were recently

inserted and could be still active in the *M. oryzae* genomes.

Then, we analyzed 11 TE families that are most abundant in the genomes of *M.*

oryzae rice isolates, including six LTR-RTs (RETRO5, RETRO6, RETRO7, Maggy,

MGLR-3, and Pyret), two non-LTR retrotransposons (MGL and Mg-SINE), and three

DNA transposons (POT2, POT3, and Occan) (**Fig S1**) (12, 22-25). To assess the

activity of these TE families, we calculated the Kimura 2-Parameter genetic distance

(k-value) to measure the divergence between TE sequences and their associated

consensus sequences (26). Low k-values indicate that the TE fragments were

generated through recent insertion events, while high k-values indicate that the TE

fragments are divergent copies generated through ancient transposition events (27).

Our analysis revealed that all 11 TE families, especially sequences of MGL, Mg-SINE,

Maggy, and a subset of POT2, POT3, and Occan, exhibit very low k-values, and more

than half of all TE contents consisted of newly emerged TEs (k-values less than 5)

(**Fig. 1a and b**), indicating a recent, large-scale burst of TEs in the genome of *M.*

oryzae rice isolates. Notably, we observed two or more k-value peaks in POT2,

Mg-SINE, and Pyret, indicating that these TEs families have undergone multiple

rounds of amplification.

Furthermore, we investigated these TE families in 275 genomes of *M. oryzae* isolates

(**Table S1**) (28, 29). We found that TEs only accounted for less than five percent of

the genomes of non-rice isolates, which is dramatically lower than that found in *M.*

oryzae rice isolates (**Table S1 and S2**). Interestingly, the k-values of TEs were much

larger and the proportion of newly emerged TEs was also much lower in *M. oryzae*

non-rice isolates, indicating that the TEs were generated by more ancient insertion

events and were inactive in *M. oryzae* non-rice isolates (**Fig. 1b and c**). Notably, the

M. oryzae non-rice isolates contained only a few copies of fragmented Maggy and

Mg-SINE, which were very abundant and possessed very low k-values in *M. oryzae*

rice isolates, suggesting that the two TEs specifically amplified in *M. oryzae* rice

isolates. In summary, our analysis demonstrated that the TEs were recently and

specifically expanded in the genome of *M. oryzae* rice isolates and maintained high

activity.

Whole genome landscape of TE dynamics in *M. oryzae* population

To examine the dynamics of transposable elements (TEs) in the *M. oryzae* rice

population, we conducted a genome-wide analysis of TE insertion sites in 90 rice

isolates that had previously been published (30), with two *M. oryzae Setaria viridis*

isolates as an outgroup (31). Using paired-end read mapping to the reference genome

method, we identified a total of 11,163 TE insertion sites, with an average of 1,312

sites per isolate. To verify these insertion sites, 17 insertion sites randomly selected

from Guy11 or FJ81278 isolates were proved to be presented as predicted through

PCR-based genotyping or PacBio (**Table S3,S4**). The number (1,312 versus 739) and

location of TE insertions differed dramatically between the rice and *S. viridis* isolates

(**Fig. S2**), reflecting the evolutionary divergence of these subspecies and their

corresponding TEs, which is consistent with the finding that these two subspecies

diverged ~10,000 years ago (31). More than half (6,040/11,163) of the TE insertion

sites were singletons specific to individual isolates (**Fig. 2**), indicating frequent TE

transposition events. Notably, the number of POT2 insertion sites was substantially

higher than those of other TE families, suggesting a higher activity and variability of

POT2 in the *M. oryzae* rice population.

Furthermore, we conducted a comprehensive analysis of the genomic distribution of

TE insertion sites in the *M. oryzae* rice population. Of the 11,163 TE insertion sites

identified, 77% (8,582/11,163) was found to be located within 1 kb of the flanking

regions of genes or intragenic regions, and over 40% of the genes (5,259/12,991) were

embedded by these TE insertions. Our enrichment analysis showed that the

distribution of the 11 TE families is non-random in the *M. oryzae* rice isolates, and

each family displayed a distinct preference for specific genomic regions. For instance,

Maggy, MGLR-3, RETRO5, RETRO7, Pyret, POT3, and Occan were predominantly

distributed in intergenic regions, while POT2 displayed a preference for the gene

flanking regions. Additionally, SINE, MGL, and RETRO6 were found to

preferentially target intragenic regions (adjusted p-value < 0.01, **Table S5**). These

findings provide valuable insights into the mechanisms underlying TE insertions and

their potential impact on gene regulation in *M. oryzae*.

Higher frequency of POT2 and POT3 insertions in promoter of secreted proteins

We observed that genes encoding secreted proteins were more closely associated with

TE junctions (**Fig. 3a**), and the proportion of genes encoding secreted proteins with

TE insertion within 1-kb flanking regions was significantly (p=7.6e-4) higher than in

those of non-secreted proteins (**Table 1**). Moreover, enrichment analysis for TEs

associated with genes encoding secreted proteins showed that POT2 and POT3 were

overrepresented in promoters of genes encoding secreted protein (adjusted p-value <

0.01, **Table S6**), implying that genes encoding secreted proteins are more prone to

disruption by POT2 and POT3.

Previous studies have shown that genes with presence/absence variation (PAV) tended

to be located near transposable elements (TEs) in fungal pathogen genomes, while

core genes were located further away from the TE-rich compartments (32-34). In this

study, we compared the genomic distribution of core genes and PAV genes and found

that PAV genes tended to be located closer to TE insertion sites than core genes (**Fig.**

3b). Our results were consistent with previous findings and suggested that PAV genes

may be more susceptible to TE-mediated disruption, potentially contributing to their

faster evolution in the context of host-pathogen interactions. Together, these findings

suggested that TEs play a role in the evolution of pathogen effectors and contribute to

the dynamic nature of host-pathogen interactions.

Association of TEs with *M. oryzae* rice population divergence

Considering that the *M. oryzae* rice population diverged within only one thousand

years (30) and that the large-scale TE burst happened recently, we thus raised the

question of whether the population divergence of *M. oryzae* is associated with recent

TE amplification events. To characterize correlations between the 90 isolates, we

estimated the distance for each of two isolates by calculating the identity of the TE

insertion sites. The pairwise TE insertion identities varied from 17.6% between

YN126441 and FJ12JN-084-3 to 87.2% between TW-PT3-1 and TW-PT6-1, with an

average of only 38.7%, thereby strongly implicating the high variability of TE

junctions between the different isolates. However, when we grouped these isolates

based on the pairwise TE insertion identities, we discovered two distinct clusters (**Fig.**

4a) matching the Clade2 and Clade3 isolates that we had previously defined based on

genome-wide SNPs (30). We noticed that the remaining isolates out of the two

clusters were also able to match the Clade1 isolates even though they showed a

relatively low pairwise TE insertion identity, which can be attributed to an earlier time

of divergence of Clade1 isolates from the *M. oryzae* rice population when compared

to the other two clade isolates. Furthermore, we found that the pairwise TE insertion

identity between intra-clade isolates was higher than that between inter-clade isolates

(**Table S7**). Surprisingly, the hierarchical tree constructed using TE insertion sites

showed a high degree of similarity to the phylogenetic tree constructed based on

whole-genome SNPs (**Fig. 4b**) (30) , indicating that the recent TE amplification event

has been a major force driving population divergence of *M. oryzae*. Consistently,

principal component analysis of these TE insertion sites displayed a similar pattern

(**Fig. 4c**). Considering that the TEs were largely amplified recently in the *M. oryzae*

genome (**Fig. 1**), we presumed that the recent high activity of TEs has been one of the

major forces driving population divergence of *M. oryzae*.

POT2 and Mg-SINE are critical for the divergence of Clade2 and Clade3 isolates

TEs are able to exert either beneficial or deleterious effects on host genomes, and the

retention or elimination of TEs is largely determined by their impact on the host.

Positive selection has been shown to drive the frequency of a TE locus to increase or

decrease dramatically during a population bottleneck or in response to a new

environment (35). We hypothesized that a portion of clade-specific TE insertion sites

had been fixed in the intra-clade isolates, contributing to the adaptive evolution of the

clade isolates. To identify the clade-specific TE insertion sites, we empirically

screened out those that were present in at least 80% of the intra-clade isolates and

absent in more than 80% of the other two clade isolates. A total of 11 Clade1-specific,

212 Clade2-specific, and 168 Clade3-specific TE insertion sites were identified, with

the number of Clade1-specific TE insertion sites being too small for subsequent

analysis. Enrichment analysis revealed that POT2 and Mg-SINE TE families were

overrepresented in both Clade2 and Clade3 datasets, indicating that the retention of

insertion of these two TE families in subpopulations of *M. oryzae* rice isolates was

tightly associated with the divergence of the rice-infection population (**Fig. 5a and b**).

We investigated the influence of clade-specific TE junctions on genes, which, for this

purpose, we referred to as clade-specific TE-associated genes (*CSTs*). We identified a

total of 238 Clade2-specific and 173 Clade3-specific TE-associated genes.

Interestingly, less than 10% of these genes overlapped, suggesting that the Clade2-

and Clade3-specific TEs have distinct targeted genes (**Fig. S2**). Gene ontology (GO)

enrichment analysis revealed that the Clade2- and Clade3-specific TE-associated

genes were enriched under similar GO terms (adjusted p-value < 0.01, **Fig. 5c, d**). Of

note, the top enriched term in both datasets was 'GO:0004497, monooxygenase

activity,' which is correlated to cytochrome P450s on the fungal genome. We further

validated the GO enrichment results by performing enrichment analysis for these

genes using the Pfam database (**Table S8**). Several previous studies have shown that

fungal P450s possess detoxifying functions towards compounds produced by host

plants during pathogen infection, thereby enhancing the fitness of the pathogenic

fungus to specific host genotypes (31, 36-40). Therefore, we suggest that TE-induced

variations in different members of P450s partially contribute to the differential

pathogenicity of the two clade isolates.

Genes associated with TE are significantly lower expressed

Previous studies have demonstrated that transposable elements (TEs) were able to

affect gene expression by inserting into gene promoters or intragenic regions (41-43).

Therefore, we investigated whether TE insertion polymorphisms could shape the gene

expression networks in *M. oryzae* rice populations. To this end, we selected 16

isolates and extracted total RNA for sequencing (**Table S9**). We identified 2,282 genes

targeted by 4,236 TE insertion sites that were polymorphic between the 16 isolates.

We then grouped the genes based on whether they contained TE insertion sites

(TE-present or TE-absent) and compared the expression levels between these two

groups. We observed that the TE-present gene group displayed significantly

(p=2.17e-26) lower expression levels than the TE-absent gene group (**Fig. 6a**),

suggesting that TEs have negative effects on the expression of their target genes. We

have identified 131 genes that exhibited clade-specific expression patterns.

Furthermore, based on the expression levels of these genes, we were able to divide 16

isolates into three distinct clusters using principal component analysis. These findings

suggest that transposable element-mediated gene regulation had a profound impact on

population divergence (**Fig. 6b and c**). Interestingly, we found that only genes with

insertion polymorphisms of POT2 and Mg-SINE displayed transcriptional

suppression, while other TE families displayed little impact on the expression of their

target genes. Our results suggest that the insertion polymorphisms of POT2 and

Mg-SINE were able to shape the gene expression network of *M. oryzae* by inducing

transcriptional suppression.

CST6* and *CST10* were required for the pathogenicity of *M. oryzae

To further investigate the impact of clade-specific TE in the virulence divergence

within *M. oryzae* rice population, we selected 15 of the clade-specific TE -associated

genes (CSTs) for functional analyses (**Fig. 7a and Table S10**). We defined CSTs as

genes that have TE insertion exclusively present in one clade and absent in another

clade. Among them, CST1-7 are Clade1-specific, and are only transcriptionally active

in Clade 2 isolates, while CST8-15 are Clade2-specific, and are only transcriptionally

active in Clade 1 isolates. We amplified CST1-7 from a Clade 2 isolate, 95085, and

ectopically overexpressed them in a Clade 1 isolate FJ81278. Conversely, CST8-15

were amplified from a Clade 1 isolate, FJ81278, and ectopically overexpressed in a

Clade 2 isolate 95085 (CST8-11) or transiently expressed in tobacco leaves

(CST12-15). The functional analyses revealed a role of CST6 and CST10 in the

pathogenicity of *M. oryzae* by leaf punch inoculation assays on the two Japonica

cultivars (NPB and TP309) and the two Indica cultivar (CO39 and MH63) (**Fig. 7b**

and c). While the CST6-OE strain produced smaller lesions and reduced fungal

biomass compared with the wild type, suggesting a potential role as a negative

regulator of virulence (**Fig. 7b and c**). The CST10-OE strain was more aggressive

and produced larger lesions and increased fungal biomass compared with the wild

type, suggesting its function as a positive regulator of virulence (**Fig. 7d and e**).

These results suggest that CST can profoundly impact virulence in *M. oryzae*,
influencing its interaction with different rice species.

DISCUSSION

The high genomic plasticity and rapid evolution of the plant pathogen *M. oryzae*
presents a severe challenge for rice blast disease control (44, 45). Previous studies
have identified transposons as a major driving force for the adaptive evolution of
fungal pathogen genomes (10). Insertion polymorphisms of transposable elements
(TEs) have been shown to lead to genomic instability, increased chromosomal
recombination, and accelerated gene evolution (17, 46). However, the precise role of
TE dynamics during *M. oryzae* evolution has remained poorly understood. To address
this, we conducted a population-scaled TE analysis of *M. oryzae* to investigate how
TE insertion polymorphisms contributed in shaping population structure and
divergence.

Previous study estimated that the *M. oryzae* rice population diverged 175 to 2,700
years ago (45). Here, we employed two methods to assess the activity of transposable
elements (TEs) in the genome of *M. oryzae*. First, we estimated the insertion time of
LTR-RTs and secondly, we calculated the Kimura 2-parameter genetic distance for the
11 most abundant TE families. Both analyses indicate that TEs have undergone recent
amplification in the *M. oryzae* genome and have remained highly active. Previous
studies have suggested that the insertion of TEs having caused gene presence/absence
variations to be the main evolutionary mechanism driving the divergence of
host-specific *M. oryzae* lineages (16). In this study, we investigated the abundance
and activity of 11 TEs in pathogens from wheat and grass infecting lineages, and
found that these 11 TEs have undergone specific expansion in *M. oryzae* rice isolates
while remain inactive in non-rice isolates. Based on these results, we propose that the
recent burst of TE activity could be the primary factor responsible for the divergence
of *M. oryzae* from other host infecting lineages and rice subspecies.

To elucidate the role of TEs in genome evolution and divergence, we conducted a

genome-wide survey of TE insertion sites in 90 *M. oryzae* rice isolates (30). We

utilized PoPoolationTE2, which is a validated method for estimating TE insertion

frequency in populations (47). The TE insertion junctions varied widely among the 90

isolates, indicating the highly dynamic and active feature of TEs in *M. oryzae* rice

populations. Consistent with prior research, we found that the distribution of TE

junctions across the genome was not evenly distributed, and different TE families or

superfamilies displayed preferences for insertion into distinct genomic regions (7, 17,

19).

TEs have been shown to be a major contributor in causing genomic variations, and the

evolution of plant pathogens (19, 48, 49). Previous studies have reported that in the

major wheat pathogen *Zymoseptoria tritici*, recent TE bursts were associated with the

proximity to genes (48). And TE insertion repressed the expression of *REP9-1* (49) or

Avr3D1 (50), and consequently resulted in the altered virulence in different isolates of

this fungus. In the polyphagous fungal pathogen *Rhizoctonia solani*, TEs mediated the

structural variations of regions encoding pathogenicity associated genes (51).

Similarly, TE insertions in or around *Avr* genes in *M. oryzae* were able to lead to

transcriptional silencing and loss of avirulence function (14, 52-56). Our analysis

revealed that TE junctions are frequently observed in the flanking regions of genes

encoding secreted proteins (SPs), and the proportion of genes encoding SPs with TE

insertions is higher than that of non-SPs. This is consistent with previous findings that

SPs are enriched in repeat-rich regions and are prone to rapid evolution (13, 57). We

propose that the variation in SPs, induced by TE insertion polymorphisms, is able to

facilitate adaptive evolution of *M. oryzae*. Presence/absence variations in genes

resulting from TE insertions have been identified to be associated with the divergence

of host-specific *M. oryzae* lineages (16). Consistent with this, we found that genes

exhibiting high gain/loss polymorphisms in the 90 isolates were preferentially located

near TE junctions, suggesting that TE-mediated presence/absence variation constitutes

a significant mechanism underlying differentiation of host-specific or intra-species *M.*

oryzae rice isolates.

Through the comparison of TE junctions in 90 *M. oryzae* rice isolates, we observed a

clustering pattern that was similar to the three previously defined *M. oryzae* clades.

This led us to investigate the potential role of TEs in *M. oryzae* rice population

divergence. We constructed a hierarchical tree based on the TE junctions and found

that it closely resembled the phylogenetic tree constructed using genome-wide SNPs.

Principal component analysis of the TE junctions also yielded a similar clustering

pattern. These findings suggested that the transposition of TEs was strongly

associated with *M. oryzae* rice population divergence. Given that both the junction of

the majority of TEs in the *M. oryzae* rice isolate genomes and the divergence of *M.*

oryzae rice population were occurred recently, we hypothesize the recent burst of TEs

to be a driving force contributing to the *M. oryzae* rice population divergence.

TE loci that undergo positive selection during evolution will be retained and will

exhibit high frequencies in a population. Based on the hierarchical tree and PCA

results, we postulate that the intra-clade isolates have a fixed set of TE insertion sites

that are specific to each clade. These clade-specific TE insertion sites are then able to

serve as molecular markers to distinguish isolates belonging to the three clades.

Interestingly, we found that POT2 and Mg-SINE were enriched in clade-specific TE

insertion sites, suggesting that these clade-specific insertions of POT2 and Mg-SINE

were beneficial for the adaptive evolution of clade isolates. Furthermore, we noted

that cytochrome P450 proteins were overrepresented in both Clade2- and

Clade3-specific TE-targeted genes. Given the essential roles of P450s in detoxifying

phytoalexins produced by host plants, we hypothesize that the differential

pathogenicity of clade isolates may be partially due to the variations in P450s induced

by clade-specific TE insertions.

TEs integrated within or flanking genes have been shown to induce gene silencing

through the formation of heterochromatic regions (10, 43). To investigate the

correlation between TE insertion and gene expression regulation in *M. oryzae*, we

performed RNA sequencing on 16 isolates from the three clades and conducted functional studies on genes disrupted by clade-specific TEs. Our analysis revealed that genes containing a TE insertion displayed significantly lower expression compared to genes without TE insertion. Notably, only POT2 and Mg-SINE insertions led to substantial suppression of gene expression. The initial functional analysis suggests that these CSTs play a crucial role in the pathogenic process. Surprisingly, these CSTs exhibit dual functionality, acting as both negative and positive regulators of virulence, while the detailed mechanisms underlying these roles require further investigation. Therefore, we hypothesized that clade-specific insertions of POT2 and Mg-SINE contribute to the adaptive evolution of clade isolates by regulating expression of specific genes and affecting adaptive traits.

Through a comprehensive analysis of TEs in populations of the rice-infecting fungus *M. oryzae*, we have revealed the crucial roles played by recent TE insertional bursts in the adaptive evolution and diversification of this fungal lineage. We demonstrate that recent TE insertions have led to the emergence of genes with clade-specific expression patterns, contributing significantly to the divergence of *M. oryzae* rice population and their adaptation to different rice subspecies. These findings highlight the significance of TE-mediated genetic changes in the regulation of gene expression, which in turn contributes to clade divergence and allows *M. oryzae* to adapt to diverse environmental pressures, including those imposed by different rice cultivars. Our findings highlight the significance of TE-mediated genetic changes in the regulation of gene expression, which in turn contributes to clade divergence.

MATERIALS AND METHODS

RNA extraction, library generation, and sequencing

The fungal strains were cultured in liquid CM medium by incubation at 28℃ under shaking at 110 rpm for three days. The mycelium was filtered, washed with

double-deionized water, and dried before being ground in liquid nitrogen. Ground

samples were transferred into DNase/RNase-free Eppendorf tubes, suspended in 1 mL

Trizol, and vortexed vigorously. To eliminate proteins, 200 μL of chloroform was

added to the mixture, which was then shaken for 15 s. After centrifugation at 12,000

rpm for 15 min at 4℃, 400 μL of the supernatant was collected and mixed with 400

μL of cold isopropanol. The mixture was kept at -20℃ for at least two hours, then

centrifuged at 12,000 rpm for 15 min at 4℃. The supernatant was discarded, and the

precipitates were washed with 1 mL of 70% alcohol and centrifuged at 12,000 rpm for

5 min at 4℃. After air drying for 5 min at room temperature, the pellets were diluted

with 54 μL of DNase/RNase-free deionized water and treated with 2 μL of DNase I

at 37℃ for 30 min. The mixture was brought up to 800 μL with RNase-free water,

followed by the addition of an equal volume of chloroform. After gentle mixing, the

mixture was centrifuged at 12,000 rpm for 15 min at 4℃. About 500 μL of the

supernatant was collected and mixed with 500 μL of cold isopropanol. The mixture

was then kept at -20℃ for more than three hours, followed by centrifugation at

12,000 rpm for 15 min at 4℃. The precipitates were washed with 1 mL of 70%

alcohol, air-dried for 5 min at room temperature, and eluted with DNase/RNase-free

deionized water. The RNA samples were then stored at -80℃ for RNA sequencing

analysis.

Estimation of TE activity

409 LTR-Finder (58) with modified parameters of '-D 15000 -d 1000 -L 7000 -l 100 -p 20

410 -C -M 0.8', LTR-harvest (59) with modified parameters of '-similar 80 -vic 10 -seed

411 20 -seqids yes -minlenltr 100 -maxlenltr 7000 -mintsd 4 -maxtsd 6 -motif TGCA

412 -motifmis 1' and LTR-Retriever, and LTR-Retriever (60) with default parameter were

413 used for *de novo* identification of full-length LTR-RTs and estimation of insertion

time. Information on TE classification is based on previous research (12) and the conserved domains of TE consensus sequences were predicted by Conserved Domain Database (61). The Kimura 2-Parameter genetic distances (k-values) of TE fragments were calculated by RepeatMasker (62) with option '-a'.

Prediction of secreted proteins and PAV genes

To identify putative secreted proteins, several criteria were employed, including the presence of a signal peptide cleavage site, the absence of a transmembrane domain, and a protein length of less than 400 amino acids. SignalP 4.0 and TMHMM 2.0 were utilized for signal peptide and transmembrane domain prediction, respectively (63, 64). The transcript sequences of the 70-15 reference genome were aligned to the previously published assemblies of 90 isolates using NCBI-blastn with default parameters. Genes that exhibited more than 90% similarity with no gaps longer than 50 bp when compared to the assemblies were marked as "present". Genes that did not meet these criteria were marked as "absent". PAV (presence-absence variation) genes were defined as those that were absent in more than five isolates, while core genes were defined as those present in all isolates.

Identification of TE insertion sites in *M. oryzae* populations

M. oryzae 70-15 TE is annotated by RepeatMasker with 11 TEs as TE library. The pair-end reads were mapped to the *M. oryzae* 70-15 reference genome using BWA (65) with default parameters. The alignment files were exported to PoPoolation2 (66) with default parameter to identify TE insertion site for each isolate. The TE insertion sites with score less than 0.3 were filtered, and the insertion sites located within 50 bp were considered as one insertion event.

Construction of TE hierarchical tree, principal component analysis

The insertion sites that are present in at least 5 isolates were used for constructing hierarchical tree and principal component analysis (PCA). The TE insertion sites in

440 presence/absence variation format were exported to the R package 'hclust' (67) for
441 construction of TE hierarchical tree. Vcftools (68) and Plink (69) were used for PCA.

Functional enrichment analysis

443 GO annotation information of *M. oryzae* was obtained from the JGI database (70).
444 Conserved domains of *M. oryzae* protein sequences were predicted using the Pfam
445 database (71). Fisher right-tailed test was used for enrichment analysis and a cutoff
446 p-value less than 0.05 was used to define significant enrichment.

RNA-seq analysis

The clean reads were mapped to the 70-15 reference genome using hisat2 v2.2.1 (72)
with default parameters. Stingtie v2.1.4 was used for expression quantification (73).
Gene expressions were normalized with transcripts per million (TPM). The
expression of each gene in the isolates with or without TE insertion were counted,
averaged and compared. Two-tailed Wilcoxon paired test was used to estimate the
significance of expression difference.

Pathogenicity assay of the *CST*-overexpressing transformants

The coding sequence (CDS) of *CST6* and *CST10* were amplified from the Clade 2
isolate, 95085, and the Clade 1 isolate, FJ81278, respectively, and were inserted into
the *pKNT-RP27* vector. The resulting constructs were then introduced into FJ81278
(*CST6*) or 95085 (*CST10*), respectively. The PEG-mediated protoplast transformation
was performed as described previously (74). To determine the role of selected *CST*
genes in the pathogenicity of *M. oryzae*, punch inoculation was performed as
previously described (75). In brief, 10 μL spore solution (5×10^5 spores/mL in
sterilized water containing 0.02% Tween) was inoculated into wounded rice leaves.
The inoculated rice plants were placed in a greenhouse. Disease symptoms were
recorded 10 days post inoculation. DNA extracted from the diseased rice leaves was
subjected to the quantitative fluorescence analyses. The primers used in this study

were listed in Table S11.

DATA AVAILABILITY

All high-throughput sequencing data generated in this study are accessible at NCBI's Gene Expression Omnibus (GEO) via GEO Series accession number GSE205351 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE205351). The consensus TE sequnces of *M. oryzae* (https://github.com/S-t-ing/mBio-data-availablility/blob/main/Mo.TE_Consensus.fasta), Genome-wide annotation of TEs in the genomes of 275 *Magnaporthe* isolates (https://github.com/S-t-ing/mBio-data-availablility/tree/main/gff), and the TE insertion sites among the 90 *Magnaporthe* rice isolates (https://github.com/S-t-ing/mBio-data-availablility/blob/main/TE%20insertion%2090%20isolates.xlsx) were available online.

ACKNOWLEDGMENTS

This work was financially supported by the National Natural Science Foundation of China (32272513 and 32172365) and Central Guidance on Local Science and Technology Development Fund of Fujian Province (2022L3088). Open Access funding provided by Fujian Agriculture and Forestry University.

REFERENCES

1.   **Wilson RA, Talbot NJ**. 2009. Under pressure: investigating the biology of plant infection by Magnaporthe oryzae. *Nat Rev Microbiol* 7:185-95.

2.   **Asibi AE, Chai Q, Coulter JA**. 2019. Rice blast: A disease with implications for global food security. *Agronomy* 9:451.

3.   **Leach JE, Vera Cruz CM, Bai J, Leung H**. 2001. Pathogen fitness penalty as a predictor of durability of disease resistance genes. *Annu Rev Phytopathol* 39:187-224.

4.   **Wang B-h, Ebbole DJ, Wang Z-h**. 2017. The arms race between *Magnaporthe oryzae* and rice: Diversity and interaction of *Avr* and *R* genes. *Journal of Integrative Agriculture* 16:2746-2760.

5.   **Dong S, Raffaele S, Kamoun S**. 2015. The two-speed genomes of filamentous pathogens: waltz with plants. *Current opinion in genetics & development* 35:57-65.

6.   **Chuong EB, Elde NC, Feschotte C**. 2017. Regulatory activities of transposable elements: from conflicts to benefits. *Nat Rev Genet* 18:71-86.

7.   **Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvák Z, Levin HL, Macfarlan TS**. 2018. Ten things you should know about transposable elements. *Genome biology* 19:1-12.

8.   **Gout L, Kuhn ML, Vincenot L, Bernard-Samain S, Cattolico L, Barbetti M, Moreno-Rico O,**

Balesdent MH, Rouxel T. 2007. Genome structure impacts molecular evolution at the AvrLm1
avirulence locus of the plant pathogen Leptosphaeria maculans. *Environ Microbiol* 9:2978-92.

9.   **Raffaele S, Kamoun S**. 2012. Genome evolution in filamentous plant pathogens: why bigger
can be better. *Nat Rev Microbiol* 10:417-30.

10.   **Faino L, Seidl MF, Shi-Kunne X, Pauper M, van den Berg GC, Wittenberg AH, Thomma BP**.
2016. Transposons passively and actively contribute to evolution of the two-speed genome of
a fungal pathogen. *Genome Res* 26:1091-100.

11.   **Seidl MF, Thomma B**. 2017. Transposable Elements Direct The Coevolution between Plants
and Microbes. *Trends Genet* 33:842-851.

12.   **Dean RA, Talbot NJ, Ebbole DJ, Farman ML, Mitchell TK, Orbach MJ, Thon M, Kulkarni R, Xu**
JR, Pan H, Read ND, Lee YH, Carbone I, Brown D, Oh YY, Donofrio N, Jeong JS, Soanes DM,
Djonovic S, Kolomiets E, Rehmeyer C, Li W, Harding M, Kim S, Lebrun MH, Bohnert H,
Coughlan S, Butler J, Calvo S, Ma LJ, Nicol R, Purcell S, Nusbaum C, Galagan JE, Birren BW.
2005. The genome sequence of the rice blast fungus Magnaporthe grisea. *Nature* 434:980-6.

13.   **Bao J, Chen M, Zhong Z, Tang W, Lin L, Zhang X, Jiang H, Zhang D, Miao C, Tang H, Zhang J,**
Lu G, Ming R, Norvienyeku J, Wang B, Wang Z. 2017. PacBio Sequencing Reveals
Transposable Elements as a Key Contributor to Genomic Plasticity and Virulence Variation in
Magnaporthe oryzae. *Mol Plant* 10:1465-1468.

14.   **Kang S, Lebrun MH, Farrall L, Valent B**. 2001. Gain of virulence caused by insertion of a Pot3
transposon in a Magnaporthe grisea avirulence gene. *Mol Plant Microbe Interact* 14:671-4.

15.   **Whisson S, Vetukuri R, Avrova A, Dixelius C**. 2012. Can silencing of transposons contribute to
variation in effector gene expression in Phytophthora infestans? *Mobile genetic elements*
2:110-114.

16.   **Yoshida K, Saunders DG, Mitsuoka C, Natsume S, Kosugi S, Saitoh H, Inoue Y, Chuma I, Tosa**
Y, Cano LM, Kamoun S, Terauchi R. 2016. Host specialization of the blast fungus
Magnaporthe oryzae is associated with dynamic gain and loss of genes linked to transposable
elements. *BMC Genomics* 17:370.

17.   **Thon MR, Pan H, Diener S, Papalas J, Taro A, Mitchell TK, Dean RA**. 2006. The role of
transposable element clusters in genome evolution and loss of synteny in the rice blast
fungus Magnaporthe oryzae. *Genome Biol* 7:R16.

18.   **Castanera R, Lopez-Varas L, Borgognone A, LaButti K, Lapidus A, Schmutz J, Grimwood J,**
Perez G, Pisabarro AG, Grigoriev IV, Stajich JE, Ramirez L. 2016. Transposable Elements
versus the Fungal Genome: Impact on Whole-Genome Architecture and Transcriptional
Profiles. *PLoS Genet* 12:e1006108.

19.   **Muszewska A, Steczkiewicz K, Stepniewska-Dziubinska M, Ginalski K**. 2019. Transposable
elements contribute to fungal genes and impact fungal lifestyle. *Sci Rep* 9:4307.

20.   **Shirke MD, Mahesh HB, Gowda M**. 2016. Genome-Wide Comparison of Magnaporthe
Species Reveals a Host-Specific Pattern of Secretory Proteins and Transposable Elements.
PLoS One 11:e0162458.

21.   **Zhong Z, Lin L, Zheng H, Bao J, Chen M, Zhang L, Tang W, Ebbole DJ, Wang Z**. 2020.
Emergence of a hybrid PKS-NRPS secondary metabolite cluster in a clonal population of the
rice blast fungus Magnaporthe oryzae. *Environ Microbiol* doi:10.1111/1462-2920.14994.

22.   **Sanchez E, Asano K, Sone T**. 2011. Characterization of Inago1 and Inago2 retrotransposons in
Magnaporthe oryzae. *Journal of general plant pathology* 77:239-242.

23. **Kachroo P, Leong SA, Chattoo BB**. 1994. Pot2, an inverted repeat transposon from the rice
blast fungus Magnaporthe grisea. *Mol Gen Genet* 245:339-48.

24. **Farman ML, Tosa Y, Nitta N, Leong SA**. 1996. MAGGY, a retrotransposon in the genome of the
rice blast fungus Magnaporthe grisea. *Mol Gen Genet* 251:665-74.

25. **Dobinson KF, Harris RE, Hamer JE**. 1993. Grasshopper, a long terminal repeat (LTR)
retroelement in the phytopathogenic fungus Magnaporthe grisea. *Mol Plant Microbe Interact*
6:114-26.

26. **Kimura M**. 1980. A simple method for estimating evolutionary rates of base substitutions
through comparative studies of nucleotide sequences. *J Mol Evol* 16:111-20.

27. **Chalopin D, Naville M, Plard F, Galiana D, Volff JN**. 2015. Comparative analysis of
transposable elements highlights mobilome diversity and evolution in vertebrates. *Genome*
Biol Evol 7:567-80.

28. **Peng Z, Oliveira-Garcia E, Lin G, Hu Y, Dalby M, Migeon P, Tang H, Farman M, Cook D, White**
FF, Valent B, Liu S. 2019. Effector gene reshuffling involves dispensable mini-chromosomes in
the wheat blast fungus. *PLoS Genet* 15:e1008272.

29. **Gómez Luciano LB, Tsai IJ, Chuma I, Tosa Y, Chen Y-H, Li J-Y, Li M-Y, Lu M-YJ, Nakayashiki H,**
Li W-H. 2019. Blast fungal genomes show frequent chromosomal changes, gene gains and
losses, and effector gene turnover. *Molecular biology and evolution* 36:1148-1161.

30. **Zhong Z, Chen M, Lin L, Han Y, Bao J, Tang W, Lin L, Lin Y, Somai R, Lu L, Zhang W, Chen J,**
Hong Y, Chen X, Wang B, Shen W-C, Lu G, Norvienyeku J, Ebbole DJ, Wang Z. 2018.
Population genomic analysis of the rice blast fungus reveals specific events associated with
expansion of three main clades. *The ISME Journal* doi:10.1038/s41396-018-0100-6.

31. **Zhong Z, Norvienyeku J, Chen M, Bao J, Lin L, Chen L, Lin Y, Wu X, Cai Z, Zhang Q, Lin X,**
Hong Y, Huang J, Xu L, Zhang H, Chen L, Tang W, Zheng H, Chen X, Wang Y, Lian B, Zhang L,
Tang H, Lu G, Ebbole DJ, Wang B, Wang Z. 2016. Directional Selection from Host Plants Is a
Major Force Driving Host Specificity in Magnaporthe Species. *Sci Rep* 6:25591.

32. **Lorrain C, Feurtey A, Möller M, Haueisen J, Stukenbrock E**. 2021. Dynamics of transposable
elements in recently diverged fungal pathogens: lineage-specific transposable element
content and efficiency of genome defenses. *G3* 11:jkab068.

33. **Plissonneau C, Hartmann FE, Croll D**. 2018. Pangenome analyses of the wheat pathogen
Zymoseptoria tritici reveal the structural basis of a highly plastic eukaryotic genome. *BMC*
Biol 16:5.

34. **Badet T, Oggenfuss U, Abraham L, McDonald BA, Croll D**. 2020. A 19-isolate
reference-quality global pangenome for the fungal wheat pathogen Zymoseptoria tritici. *BMC*
Biol 18:12.

35. **Torres DE, Oggenfuss U, Croll D, Seidl MF**. 2020. Genome evolution in fungal plant pathogens:
looking beyond the two-speed genome model. *Fungal Biology Reviews*.

36. Schuler MA. 2015. P450s in plants, insects, and their fungal pathogens, p 409-449,
Cytochrome P450. Springer.

37. **Ide M, Ichinose H, Wariishi H**. 2012. Molecular identification and functional characterization
of cytochrome P450 monooxygenases from the brown-rot basidiomycete Postia placenta.
Arch Microbiol 194:243-53.

38. **Nazir KH, Ichinose H, Wariishi H**. 2011. Construction and application of a functional library of
cytochrome P450 monooxygenases from the filamentous fungus Aspergillus oryzae. *Appl*

Environ Microbiol 77:3147-50.

39.   **Doddapaneni H, Chakraborty R, Yadav JS**. 2005. Genome-wide structural and evolutionary
analysis of the P450 monooxygenase genes (P450ome) in the white rot fungus
Phanerochaete chrysosporium: evidence for gene duplications and extensive gene clustering.
BMC Genomics 6:92.

40.   **Zhang S, Widemann E, Bernard G, Lesot A, Pinot F, Pedrini N, Keyhani NO**. 2012. CYP52X1,
representing new cytochrome P450 subfamily, displays fatty acid hydroxylase activity and
contributes to virulence and growth on insect cuticular substrates in entomopathogenic
fungus Beauveria bassiana. *J Biol Chem* 287:13477-86.

41.   **Santana M, Queiroz M**. 2015. Transposable elements in fungi: a genomic approach. *Scientific*
J Genetics Gen Ther 1:012.

42.   **Krishnan P, Meile L, Plissonneau C, Ma X, Hartmann FE, Croll D, McDonald BA,**
Sanchez-Vallet A. 2018. Transposable element insertions shape gene regulation and melanin
production in a fungal pathogen of wheat. *BMC Biol* 16:78.

43.   **Lewis ZA, Honda S, Khlafallah TK, Jeffress JK, Freitag M, Mohn F, Schubeler D, Selker EU**.
2009. Relics of repeat-induced point mutation direct heterochromatin formation in
Neurospora crassa. *Genome Res* 19:427-37.

44.   **Gladieux P, Condon B, Ravel S, Soanes D, Maciel JLN, Nhani A, Jr., Chen L, Terauchi R, Lebrun**
MH, Tharreau D, Mitchell T, Pedley KF, Valent B, Talbot NJ, Farman M, Fournier E. 2018.
Gene flow between divergent cereal- and grass-specific lineages of the rice blast fungus
Magnaporthe oryzae. mBio 9.

45.   **Gladieux P, Ravel S, Rieux A, Cros-Arteil S, Adreit H, Milazzo J, Thierry M, Fournier E,**
Terauchi R, Tharreau D. 2018. Coexistence of multiple endemic and pandemic lineages of the
rice blast pathogen. *mBio* 9.

46.   **Chadha S, Sharma M**. 2014. Transposable elements as stress adaptive capacitors induce
genomic instability in fungal pathogen Magnaporthe oryzae. *PLoS One* 9:e94415.

47.   **Kofler R, Gomez-Sanchez D, Schlotterer C**. 2016. PoPoolationTE2: Comparative Population
Genomics of Transposable Elements Using Pool-Seq. *Mol Biol Evol* 33:2759-64.

48.   **Oggenfuss U, Croll D**. 2023. Recent transposable element bursts are associated with the
proximity to genes in a fungal plant pathogen. *PLoS Pathog* 19:e1011130.

49.   **Wang C, Milgate AW, Solomon PS, McDonald MC**. 2021. The identification of a transposon
affecting the asexual reproduction of the wheat pathogen *Zymoseptoria tritici. Mol Plant*
Pathol 22:800-816.

50.   **Fouché S, Badet T, Oggenfuss U, Plissonneau C, Francisco CS, Croll D**. 2020. Stress-driven
transposable element de-repression dynamics and virulence evolution in a fungal pathogen.
Mol Biol Evol 37:221-239.

51.   **Francis A, Ghosh S, Tyagi K, Prakasam V, Rani M, Singh NP, Pradhan A, Sundaram RM,**
Priyanka C, Laha GS, Kannan C, Prasad MS, Chattopadhyay D, Jha G. 2023. Evolution of
pathogenicity-associated genes in *Rhizoctonia solani* AG1-IA by genome duplication and
transposon-mediated gene function alterations. *BMC Biology* 21:15.

52.   **Li W, Wang B, Wu J, Lu G, Hu Y, Zhang X, Zhang Z, Zhao Q, Feng Q, Zhang H, Wang Z, Wang G,**
Han B, Wang Z, Zhou B. 2009. The Magnaporthe oryzae avirulence gene AvrPiz-t encodes a
predicted secreted protein that triggers the immunity in rice mediated by the blast resistance
gene Piz-t. *Mol Plant Microbe Interact* 22:411-20.

53. **Zhang S, Wang L, Wu W, He L, Yang X, Pan Q**. 2015. Function and evolution of Magnaporthe
oryzae avirulence gene AvrPib responding to the rice blast resistance gene Pib. *Sci Rep*
5:11642.

54. **Chen C, Chen M, Hu J, Zhang W, Zhong Z, Jia Y, Allaux L, Fournier E, Tharreau D, Wang G-L**.
2014. Sequence variation and recognition specificity of the avirulence gene AvrPiz-t in
Magnaporthe oryzae field populations. *Fungal Genomics & Biology* 4.

55. **Miki S, Matsui K, Kito H, Otsuka K, Ashizawa T, Yasuda N, Fukiya S, Sato J, Hirayae K, Fujita Y,**
Nakajima T, Tomita F, Sone T. 2009. Molecular cloning and characterization of the AVR-Pia
locus from a Japanese field isolate of Magnaporthe oryzae. *Mol Plant Pathol* 10:361-74.

56. **Hu ZJ, Huang YY, Lin XY, Feng H, Zhou SX, Xie Y, Liu XX, Liu C, Zhao RM, Zhao WS, Feng CH,**
Pu M, Ji YP, Hu XH, Li GB, Zhao JH, Zhao ZX, Wang H, Zhang JW, Fan J, Li Y, Peng YL, He M, Li
DQ, Huang F, Peng YL, Wang WM. 2022. Loss and Natural Variations of Blast Fungal
Avirulence Genes Breakdown Rice Resistance Genes in the Sichuan Basin of China. *Front Plant*
Sci 13:788876.

57. **Wu J, Kou Y, Bao J, Li Y, Tang M, Zhu X, Ponaya A, Xiao G, Li J, Li C, Song MY, Cumagun CJ,**
Deng Q, Lu G, Jeon JS, Naqvi NI, Zhou B. 2015. Comparative genomics identifies the
Magnaporthe oryzae avirulence effector AvrPi9 that triggers Pi9-mediated blast resistance in
rice. *New Phytol* 206:1463-75.

58. **Xu Z, Wang H**. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Res* 35:W265-8.

59. **Ellinghaus D, Kurtz S, Willhoeft U**. 2008. LTRharvest, an efficient and flexible software for de
novo detection of LTR retrotransposons. *BMC Bioinformatics* 9:18.

60. **Ou S, Jiang N**. 2018. LTR_retriever: A Highly Accurate and Sensitive Program for Identification
of Long Terminal Repeat Retrotransposons. *Plant Physiol* 176:1410-1422.

61. **Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI,**
Marchler GH, Song JS, Thanki N, Yamashita RA, Yang M, Zhang D, Zheng C, Lanczycki CJ,
Marchler-Bauer A. 2020. CDD/SPARCLE: the conserved domain database in 2020. *Nucleic*
Acids Res 48:D265-D268.

62. **Tempel S**. 2012. Using and understanding RepeatMasker. *Methods Mol Biol* 859:29-51.

63. **Petersen TN, Brunak S, von Heijne G, Nielsen H**. 2011. SignalP 4.0: discriminating signal
peptides from transmembrane regions. *Nat Methods* 8:785-6.

64. **Krogh A, Larsson B, von Heijne G, Sonnhammer EL**. 2001. Predicting transmembrane protein
topology with a hidden Markov model: application to complete genomes. *J Mol Biol*
305:567-80.

65. **Li H, Durbin R**. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* 25:1754-60.

66. **Kofler R, Pandey RV, Schlotterer C**. 2011. PoPoolation2: identifying differentiation between
populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics* 27:3435-6.

67. **Galili T**. 2015. dendextend: an R package for visualizing, adjusting and comparing trees of
hierarchical clustering. *Bioinformatics* 31:3718-20.

68. **Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,**
Marth GT, Sherry ST, McVean G, Durbin R, Genomes Project Analysis G. 2011. The variant
call format and VCFtools. *Bioinformatics* 27:2156-8.

69. **Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de**

677    Bakker PI, Daly MJ, Sham PC. 2007. PLINK: a tool set for whole-genome association and
678    population-based linkage analyses. *Am J Hum Genet* 81:559-75.

679 70. Nordberg H, Cantor M, Dusheyko S, Hua S, Poliakov A, Shabalov I, Smirnova T, Grigoriev IV,
680    Dubchak I. 2014. The genome portal of the Department of Energy Joint Genome Institute:
681    2014 updates. *Nucleic Acids Res* 42:D26-31.

682 71. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE,
683    Paladin L, Raj S, Richardson LJ, Finn RD, Bateman A. 2021. Pfam: The protein families
684    database in 2021. *Nucleic Acids Res* 49:D412-D419.

685 72. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
686    genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37:907-915.

687 73. Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015. StringTie
688    enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol*
689    33:290-5.

690 74. Zhang L-m, Chen S-t, Qi M, Cao X-q, Liang N, Li Q, Tang W, Lu G-d, Zhou J, Yu W-y, Wang Z-h,
691    Zheng H-k. 2021. The putative elongator complex protein Elp3 is involved in asexual
692    development and pathogenicity by regulating autophagy in the rice blast fungus. *Journal of*
693    *Integrative Agriculture* 20:2944-2956.

694 75. Han Y, Song L, Peng C, Liu X, Liu L, Zhang Y, Wang W, Zhou J, Wang S, Ebbole D, Wang Z, Lu
695    GD. 2019. A *Magnaporthe* chitinase interacts with a rice jacalin-related lectin to promote
696    host colonization. *Plant Physiol* 179:1416-1430.

Figure legends

FIG 1 The distribution of 11 TE families in the *M. oryzae* population. a) The Kimura 2-Parameter genetic distance of 11 TE families in three different *Magnaporthe* species were shown in density plot. **b, c)** The k-value proportions of 11 TE families in *M. oryzae* rice isolates (**b**) and non-rice isolates (**c**). The numbers of TE fragment were marked at the top of bars.

FIG 2 The number of TE junctions in the *M. oryzae* rice population. The circos diagram (from outer to inner) displays chromosomes, gene distribution, distribution of DNA transposon, Non-LTR-RT and LTR-RT junctions. The histogram shows the insertion site numbers of different TE families. The non-transparent and transparent colors represent the numbers of isolate-specific and isolate-shared TE junctions, respectively

FIG 3 The TEs were in proximity to secreted proteins and PAV genes. The distance of TEs to the 5' and 3' ends of secreted proteins and non-secreted proteins (**a**) or core genes and PAV genes (**b**).

FIG 4 TE insertion was associated with the divergence of *M. oryzae* rice population. a) The heatmap showing the identity of TE insertion site between each two isolates. **b)** The hierarchical tree built by using the TE insertion sites of *M. oryzae* population. Three distinct clades were marked in red, blue and green colors respectively. **c)** Principal component analysis using the TE insertion sites.

FIG 5 Characteristics of clade-specific TEs in the *M. oryzae* rice population. a, b) Enrichment analysis of 11 TE families in Clade2- (**a**) and Clade3-specific (**b**) insertion sites. Fisher's exact test was used for significance test. **c)** GO enrichment analysis for the clade2- (**c**) and clade3-specific (**d**) TE-associated genes.

FIG 6 The impact of clade-specific TE insertion on gene expression. a) Comparison of gene expressions between the isolates present or absent with TE insertion. **b)** PCA clustering of 16 isolates based on expression level of 131 clade-specific DEGs. **c)** Heatmap showing expression level of 131 clade-specific DEGs in 16 isolates.

FIG 7 CST6 and CST10 were required for the pathogenicity of *M. oryzae* rice isolates. a) Heatmap showing expression level of 15 clade-specific TE-associated genes (CSTs) in 16 isolates. **b)** Leaf punch inoculation assays were conducted to assess the impact of CST6 overexpression on two Japonica cultivars (NPB and TP309) and the two Indica cultivar (CO39 and MH63). The Clade3 isolate (FJ81278), lacking CST6 expression, was used as the wild-type control. **c)** Leaf punch inoculation assays were conducted to assess the impact of CST10 overexpression on two Japonica cultivars (NPB and TP309) and the two Indica cultivar (CO39 and MH63). The Clade3 isolate (95085), lacking CST10 expression, was used as the wild-type control. **d)** Relative fungal biomass on leaf punch inoculation assays of CST6 overexpression on two Japonica cultivars (NPB and TP309) and the two Indica cultivar (CO39 and MH63). **e)** Relative fungal biomass on leaf

punch inoculation assays of CST10 overexpression on two Japonica cultivars (NPB and TP309) and the two Indica cultivar (CO39 and MH63).

TABLE 1 The numbers of secreted protein and non-secreted protein that have TE insertion in 1-kb flanking regions or within gene.

SUPPLEMENTAL MATERIAL

FIG S1 Schematic of the 11 most abundant TE families on the genome of the *M. oryzae* rice isolate. The conserved domains in the relative location of the TE consensus were highlighted using different colors.

FIG S2 The venn chart compares the TE insertion loci of 90 *M. oryzae* rice isolates and two *S. viridis* isolates.

FIG S3 The venn chart displays the intersection of clade2-specific and clade3-specific TE-associated genes.

TABLE S1 The proportions of 11 TE families on the genomes of 275 *M. oryzae* isolates.

TABLE S2 The proportions of 11 TE families on the genomes of a *M. oryzae* wheat isolate and three grass isolates.

TABLE S3 Validation of 17 TE insertion sites by PacBio.

TABLE S4 Validation of 17 TE insertion sites by PCR amplification.

TABLE S5 Insertion preference of various TEs in different genomic regions.

TABLE S6 Enrichment analysis of TEs in the promoter of secreted proteins.

TABLE S7 Identity of TE insertion sites between intra- or inter-clade isolates.

TABLE S8 Enrichment analysis of the clade2- and clade3-specific TE-associated genes using conserved domains.

TABLE S9 GEO information of 16 RNA seq samples.

TABLE S10 Basic information about 15 clade-specific TE-associated genes (CSTs).

TABLE S11 Primers used in this study.

a

Retro5 — k=1.4
Retro6 — k=2.3
Retro7 — k=3
Maggy — k=0.2
MGLR-3 — k=2.5
Pyret — k=3.6
MGL — k=0.8
Mg-SINE — k=0.4
POT2 — k=0.2
POT3 — k=0.3
Occan — k=0.8

K-value

Rice Isolates
Grass Isolates
Wheat Isolates

b *M. oryzae* Rice Isolates

n=2,559 n=1,770 n=2,721 n=1,589 n=1,015 n=6,185 n=2,175 n=1,234 n=4,000 n=1,167 n=764 n=25,179

RETRO5 RETRO6 RETRO7 Maggy MGLR-3 Pyret MGL SINE POT2 POT3 Occan Total

LTR-RTs   Non-LTR-RTs   DNA Transposon

k-value
> 20
5 - 20
<= 5

c *M. oryzae* Non-Rice Isolates

n=716 n=865 n=775 n=58 n=579 n=3,199 n=499 n=56 n=716 n=569 n=343 n=8,375

RETRO5 RETRO6 RETRO7 Maggy MGLR-3 Pyret MGL SINE POT2 POT3 Occan Total

LTR-RTs   Non-LTR-RTs   DNA Transposon