Cryo-electron tomography reveals postsynaptic nanoblocks in excitatory synapses

for transcellular alignment

Rong Sun[1], Liana Wilson[1], Mariam Haider[1,2], Baris Alten[3,6], Zimeng Zhou[1,4], Xinyi

Wang[1,5], Qiangjun Zhou[1*].

[1] Department of Cell and Developmental Biology, Vanderbilt Brain Institute, Center for

Structural Biology, Vanderbilt Kennedy Center, Vanderbilt University, Nashville, TN,

37240, USA.

[2] Center for Structural Biology Cryo-EM Facility, Vanderbilt University, Nashville, TN,

37240, USA.

[3] Department of Pharmacology, Vanderbilt University, Nashville, TN, 37212, USA.

[4] School of Engineering, Vanderbilt University, Nashville, TN, 37212, USA.

[5] Peabody College, Vanderbilt University, Nashville, TN, 37212, USA

[6] Present address: Harvard Medical School, Boston, MA, 02115, USA and Department of

Neurology, Massachusetts General Hospital and Brigham & Women's Hospital, Boston,

MA, 02115, USA

Abstract

The nanoscale organization of proteins within synapses is critical for maintaining and
regulating synaptic transmission and plasticity. Here, we use cryogenic electron
tomography to directly visualize the *in situ* three-dimensional architecture and
supramolecular organization of transsynaptic alignment of pre-cleft-postsynaptic
components in their native cellular context in both synaptosomes from rat hippocampi
and synapses from rat primary cultured neurons. High-resolution electron microscopy
and quantitative analyses reveal that release sites align with adhesion molecules, receptor
clusters, and postsynaptic density (PSD) nanoblocks as a physical, transsynaptic, nano-
alignment. Additionally, it has been determined that the PSDs contain different-sized,
membrane-associated, subsynaptic protein nanoblocks. Furthermore, large nanoblocks
are formed by small nanoblocks positioned close enough together. Lastly, the tomograms
of synaptosomes allow us to detect the glutamate receptor-like particles by subtomogram
averaging at resolutions of 24 Å and 26 Å. The results of this study provide a more
comprehensive understanding of synaptic ultrastructure and demonstrate that PSD is
organized into subsynaptic nanoblocks to form transcellular alignment.

Introduction

Cellular function is driven by nanoscale spatial organization of multi-protein assemblies and their interactions with other molecular components and cellular organelles. Typical examples of such nanoscale structures are neuronal chemical synapses. These are the most complex cell-cell junctions, with more than 2,000 distinct synaptic proteins.[1] In particular, laboratories have been accumulating evidence that multiple presynaptic, postsynaptic, and cell-adhesion proteins in excitatory synapses are finely organized in lipid-bound nanoscale structures to facilitate extremely rapid, dynamic, efficient, and tightly regulated transmission of information between neurons[2-10]. Moreover, the activity-dependent rearrangement of these synaptic proteins, including neurotransmitter receptors and other synaptic proteins, plays a key mechanistic role in synaptic plasticity. Synaptic plasticity is essential for perception, decision making, learning, and memory formation.

Recent studies provide evidence for the high-level organization of nanoscale supramolecular assemblies spanning the synaptic cleft, giving rise to *nanocolumns* of juxtaposed pre- and postsynaptic compartments in excitatory synapses[3,4,11-16]. Specifically, presynaptic active-zone scaffold proteins, such as RIM, form nanodomains that define synaptic vesicle docking and release sites. These sites align with clustered neurotransmitter receptors such as α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (also known as AMPAR) and postsynaptic scaffolding proteins such as PSD95 via adhesion proteins in the synaptic cleft[10,17,18].

The postsynaptic density (PSD) appears in conventional electron micrographs as an electron-dense lamina just beneath the postsynaptic membran.[19]. A PSD meshwork model comprising regularly spaced vertical filaments was proposed based on electron microscopy (EM) and biochemical assays[20-24]. Two additional forms of PSD organization have been proposed: nano-domains based on super-resolution imaging[3-5,25,26], and liquid condensate based on *in vitro* PSD mixing assay[27-29]. All these studies provide more evidence for the high-level organization of nanoscale supramolecular assemblies in the synapse, indicating that synaptic proteins are not randomly distributed and instead form functional synaptic nanostructures. However, nanoscale organization of the PSD and postsynaptic receptors is largely unknown due to the technical challenges imposed by synaptic molecular complexity.

Cryo-electron tomography (cryo-ET) is a powerful technique that produces three-dimensional (3D) volume reconstructions, or *tomograms*, of macromolecular and subcellular biological structures in their native context at nanometer resolution. To prevent producing artifacts through chemical fixation, dehydration, and heavy-metal staining, biological samples are rapidly frozen to obtain a "true, instantaneous snapshot"[30-33]. In general, obtaining high-quality data requires cryo-ET samples to be thin enough for electrons to pass through (~200 nm), posing a technical challenge during sample preparation, which limits the resolution of cellular cryo-ET[34]. The resolution is also limited by the crowded cellular environment. To overcome these limitations, isolated synaptic terminals have been used for cryo-ET studies to decrease sample thickness[10,35]. A classic type of isolated synaptic terminal called a *synaptosome* is highly suitable for cryo-ET, because it retains a piece of attached postsynaptic membrane that faces the

active zone[36,37]. Synaptosomes help to improve the resolution, as the average thickness of
a synaptosome is 200-300 nm with the core of PSD in our preparations. Here, we
developed a simplified sample preparation workflow to obtain the synaptosomes from
mature rat hippocampi, achieving high-resolution synaptic nanostructures with cryo-ET
and sub-tomogram averaging. With this method, we directly visualized the overall
architecture of the transcellular alignment in excitatory synapses. Our results revealed
that PSD is formed by subsynaptic PSD nanoblocks of specific sizes. We also determined
in situ structures of two types of putative ionotropic glutamate receptor-like particles at
24-Å and 26-Å resolution. Finally, we demonstrated subsynaptic organization of PSD
nanoblocks, as well as the physical transcellular alignment between tethered synaptic
vesicles and PSD nanoblocks in intact synapses within rat primary cultured hippocampal
neurons.

Results

Physical transsynaptic alignment in synaptosomes

To study subsynaptic organization, we prepared synaptosomes via a simplified workflow.
After dissecting the hippocampi from rat brains, we performed four strokes of
homogenization and a 10-minute centrifugation at 800 *g,* followed by plunge-freezing
and cryo-ET data acquisition (Supplementary Fig. 1a). Our preparations included two
different types of isolated synaptic terminals based on their morphology: Synaptosomes
retained a patch of postsynaptic membrane with PSD (Fig. 1 and Supplementary Fig. 1b-
e, see also Supplementary Video 2), while synaptoneurosomes retained an enclosed
postsynaptic compartment (Supplementary Fig. 1f-i)[38,39]. For this study, we primarily
focused on synaptosomes as we were able to achieve higher-resolution structural
information from these relatively thin and molecularly- uncrowded samples compared to
synaptoneurosomes and intact synapses. We acquired 60 high quality tomograms of
isolated synaptic terminals (40 synaptosomes and 20 synaptoneurosomes), which enabled
us to directly visualize characteristic features of a neuronal synapse, including typical
synaptic proteins (Fig. 1a, 1b and Supplementary Fig. 1p-u). In all our preparations, we
were able to find synaptosomes and synaptoneurosomes and did not observe obvious
morphological differences in synaptosomes or synaptoneurosomes at nanoscale between
batches of sample preparations. To ensure the data consistency in PSD density analyses
and subtomogram averaging, all 60 tomograms used were from the same data collection
session and from the same batch of sample preparation. All the synaptosomes we
obtained were likely from excitatory synapses, as they have thick PSDs of more than 30
nm, which is a known feature of excitatory synapses[19,22].

To study the relationship between release sites, adhesion molecules, postsynaptic
densities, and synaptic receptors, we scrutinized the tethered synaptic vesicles that
indicate the position of synaptic release sites. We also investigated the corresponding
nanostructures in the cleft and in the postsynaptic compartment. Examining the electron
densities and measuring the gray values of synaptosome tomograms (Fig. 1c and
Supplementary Fig. 1p-s) allowed us to detect the adhesion molecule-like proteins and
membrane proteins in the synaptic cleft, as well as the PSD protein clusters, or which we
call *nanoblocks*. The tomographic slice in Fig. 1c's left panel shows protein densities in

the middle of the cleft, which we posit to be the adhesion molecules, as well as a density
layer on the postsynaptic membrane that is comprised of membrane proteins. These direct
visualizations are consistent with the density plot in Fig. 1c's right panel. Some of the
postsynaptic membrane proteins are most likely synaptic receptors, based on their
geometry and size (see Methods regarding the criteria to select particles for subtomogram
averaging). We found that synaptosomes and synaptoneurosomes have specific cleft
widths (Fig. 1d); and the clefts of synaptosomes tend to be wider than those of
synaptoneurosomes (39.7 ± 0.9 nm for synaptosomes n = 33; and 30.5 ± 0.9 nm for
synaptoneurosomes n = 20). Of note, 7 synaptosomes were not included in these
measurements as the postsynaptic membranes were not parallel to the presynaptic
membranes. This might be due to different adhesion molecules surviving the sample
isolation process, thus resulting in different widths of synaptic clefts. We observed that
tethered vesicles aligned with the adhesion-like molecules, the membrane protein-like
particles, and PSD densities (Fig. 1e-i). This direct visualization of pre-cleft-postsynaptic
alignment suggests that postsynaptic receptors, which anchored by PSD scaffold proteins,
are aligned with presynaptic release sites through adhesion molecules in the synaptic
cleft. Two examples of adhesion-receptor-PSD nanoblock alignment without a tethered
synaptic vesicle are also shown here, indicating these might be potential release sites
(Fig. 1j-k).

PSD subsynaptic organization in synaptosomes

Next, we investigated PSD densities and performed quantitative analyses on the PSD
nanoblock dimensions. A representative tomogram of a synaptosome acquired by cryo-
ET is shown in Fig. 2a, in which PSD nanoblocks (indicated by red arrows) are clearly
visualized beneath the membrane. In terms of the overall PSD electron density patterns,
this result is consistent with previous EM studies stating that the core of the PSD is
dominated by vertically oriented filaments[20,40] or spiky structures[24]. In our results, there
were roughly three types of synaptosomes based on the organization of the PSD:
synaptosomes with separated PSD nanoblocks; synaptosomes with both separated and
continuous PSD nanoblocks; and synaptosomes with continuous PSD (Supplementary
Fig. 1j-o). To delineate the PSD nanostructures of all three synaptosome types, we
applied the following quantitative analysis: First, we measured gray values of protein
densities of a 40-nm thick PSD area beneath the postsynaptic membrane (Fig. 2a-b).
Then, we combined 2D gray values across a series of continuous virtual tomographic
slices, generating a gray value density map (Fig. 2c, top panel). With normalization and
filtering processing, **d**ensity-**b**ased **s**patial **c**lustering of **a**pplications with **n**oise
(DBSCAN) results show projected PSD density nanoblocks in the 40-nm range from the
postsynaptic membrane (Fig.2c, bottom panel). DBSCAN is a density-based clustering
non-parametric algorithm[41] and it is widely used in biological data analysis[42,43]. We next
measured the area and width of each protein nanoblock (Fig. 2c-d). The area of
nanoblock was defined as its convex hull area shown by encircling red lines. The width
of nanoblock was defined as the largest x-axis distance between two pixels in each
cluster. Fig. 2d shows examples of nanoblocks with different areas and widths quantified
by the DBSCAN clustering method. Multiple peaks were detected in the area distribution

of nanoblocks from all synaptosomes (Fig. 2e), while the width distribution was fit with
two peaks (Fig. 2f). Distinct fitting peaks in the area distribution histogram indicate that
nanoblocks have distinct sizes, and that large nanoblocks may be composed of several
small nanoblocks, which is also supported by examining the morphology of some
nanoblocks with large widths (Fig. 2d). Though the geometric shape of the nanoblock
was irregular, the way we quantified the width reflected the dimension of the nanoblock
in x axis, while the dimension of nanoblock in z axis was affected by the missing wedge
effect. Overall, our analysis suggests that PSD nanoblocks can be positioned closely
enough to one another to form large nanoblocks and coordinate synaptic function.

To further analyze the size and morphology of subsynaptic nanoblocks, we measured the
heights of the protein nanoblocks obtained via DBSCAN clustering. The height was
quantified based on gray values along the nanoblock at its largest length (Fig. 1c). The
distribution of nanoblock heights reveals that most nanoblocks are in the 30~60 nm
range, while a small population is below 20 nm or above 80 nm (Fig. 2g). The large
nanoblocks shown on the right side of Fig. 2h appear to be composed of multiple small
protein nanoblocks, which were positioned close together in our clustering analysis. The
two schematics on the right side of Figure 2H show these large nanoblocks' morphology.
Their heights are also variable. Additionally, we analyzed the gap between PSD
nanoblocks. The distance between each nanoblock centroid (Fig. 2i) and the nearest
distance between each nanoblock edge (Fig. 2j) together describe the spacing between
nanoblocks, suggesting that the gaps with less protein densities segmented the PSD.
Overall, our analysis demonstrates that subsynaptic nanoblocks of different sizes form the
PSD.

***In situ* organization of receptor-like particles on the postsynaptic membrane**

To further investigate the molecular organization of nano-alignment, we employed the
subtomogram averaging method to detect glutamate receptors in our synaptosome
samples. AMPARs or n-methyl-d-aspartate receptors (NMDARs) are prominent receptors
in excitatory synapses and consist of N-terminal domains (NTD) and ligand binding
domains (LBD) within the synaptic cleft[44-46]. We manually selected 1,565 particles from
28 synaptosomes with NTD and LBD in the range from 10 nm to 14 nm for
subtomogram averaging (Fig. 3a and Supplementary Fig. 3a). Eventually, we achieved an
averaged structure of type A receptor with 391 particles (Fig. 3b-e) and an averaged
structure of type B receptor with 189 particles (Fig. 3f-i). The type A receptor is
structurally similar to the O-shape AMPAR (PDB: 5ide), while the type B receptor is
structurally similar to the Y-shape AMPAR (PDB: 6qkz). In previous studies, the
extracellular N-termini of Y-shape AMPARs were determined to be ~14 nm in height[47,48],
while the extracellular N-termini of O-shape AMPARs were found to be ~10 nm[49], and
the extracellular N-termini of NMDARs were ~11 nm in height[50,51]. Considering the
morphology similarity of N-termini between O-shape AMPAR and NMDAR at the
resolution we achieved (Supplementary Fig. 3b), it is difficult to exclude the possibility
that some NMDARs were also counted in the type A receptor population.

To study the distribution of the averaged receptors, we anchored the type A and type B
receptors to the PSD clustering results. As displayed in Fig. 3j, three examples
demonstrate that some of the averaged receptors (type A receptors and type B receptors)
are within PSD nanoblocks, while there are also receptors outside of the PSD clusters.
We also identified some particles that remained unclassified and were neither type A nor
type B receptors. Among these undefined particles, some are within, and some are outside
of the PSD nanoblocks. Previous studies show that AMPAR and NMDAR are targeted on
the postsynaptic membrane by scaffolding proteins within the PSD[52-56]. However, the
protein nanoblocks represent the core of the PSD; and it is possible that these particles
which do not fall onto the PSD nanoblocks, are responsible for scaffolding by interacting
with proteins outside of the nanoblocks. Compared to the type B receptors, the nearest
neighbor distance analysis indicates that the type A receptors are more clustered (Fig. 3k-
l). The number of type A and type B receptors varies in all synaptosomes (Fig. 3m),
showing the heterogeneity of the synaptosomes.

PSD subsynaptic organization in intact neurons

We also examined the subsynaptic organization of synapses to determine whether similar
subsynaptic nanoblocks identified in synaptosomes can also be found in intact synapses
from cultured neurons. To accomplish this, we seeded dissociated primary hippocampal
neurons on gold EM grids and examined the synaptic ultrastructure in intact neurons
using cryo-ET (Supplementary Fig. 4a-e). In reconstructed tomograms, a typical synapse
consists of a presynaptic compartment with a population of synaptic vesicles, a synaptic
cleft, and a postsynaptic compartment with PSD. We also focused on excitatory synapses
and excluded synapses with thin PSDs in cultured neurons from our analysis
(Supplementary Fig. 4f-h).

We collected 211 tilt series of synapse-like structures from the wild-type primary
hippocampal neurons and reconstructed them. 131 of the reconstructed tomograms were
synapses, and 121 synapses had thick PSDs. Three examples of excitatory synapses are
shown in Fig. 4. Tomographic slices (Fig. 4a, x, f, h, k, and m) and segmented structures
(Fig. 4b and g, see also Supplementary Video 2) display PSDs' protein densities,
suggesting that protein distribution is not homogeneous within the PSD. However, the
crowded molecular environment and overall thickness of neurons rendered a low signal-
to-noise ratio in the cellular context, making patterns of protein complexes difficult to
recognize and analyze. Using the software package IsoNet[57], we performed contrast
transfer function (CTF) deconvolution and corrected the missing wedge effect to enhance
contrast in 32 out of 121 cryo-electron tomograms of excitatory synapses. Following
IsoNet correction, tomographic slices show more obvious protein nanoblocks in the PSDs
(Fig. 4d, i, and n). Quantitative characterization of the PSDs' 3D volume (Fig. 4e, j, and
o) indicates that proteins distribute unevenly within the PSD and form subsynaptic
nanoblocks. The nanoblocks' area and distance (Fig. 4p-r) show similar distributions to
those from our synaptosome datasets. Our observations suggest that in intact synapses,
there are nanoblocks of different sizes consisting of postsynaptic proteins, similar to
those were observed in synaptosomes. Additionally, multiple tethered vesicles, docked

vesicles, or fused vesicles represent release sites (Supplementary Fig. 4i-n) and tend to
align with large PSD nanoblocks (Fig. 4i and o).

Discussion

In this study, we utilized cryo-ET to directly visualize the ultrastructure of excitatory
synapses in their near-native state within purified hippocampal synaptosomes from 10-
week-old young adult rats as well as primary hippocampal neurons cultured from
postnatal day 0 newborn rats. Importantly, we found that in our synaptosome samples, the
exposed postsynaptic membranes remain attached to the presynaptic terminals. The only
components remaining were the membrane-associated PSD core components (PSD
nanoblocks), while weakly interacting proteins (in PSD gaps) and other postsynaptic
components might be lost. This allowed us to study the core PSD components against a
relatively clean background. Meanwhile, studies of the whole synapse in intact cultured
neurons provided more physiological and complete ultrastructure features. With these
two different samples, we consistently recorded the physical transsynaptic alignment and
subsynaptic organization in the form of PSD nanoblocks. Our observations suggest that
membrane-associated PSD core components within synaptosomes are relatively stable,
and that they maintain structural integrity during the isolation process for synaptosome
preparation via our simplified protocol. This finding is consistent with previous
observations that PSD complexes and the PSD membranes are detergent-resistant and
stable enough to be purified from conventional synaptic fractions[58-60]. Thus, it is likely
that the nanoscale organization and basic ultrastructure features we observed also reflect
synaptic architecture within the brain.

An alignment between pre- and postsynaptic elements has long been proposed but was
only recently reported and named a *nanocolumn*, discovered through super-resolution
light microscopy[3]. Thus far, only labeled synaptic proteins such as PSD95, receptors, and
RIM have been shown to form nanoclusters as a part of nanocolumns. Our cryo-ET
approach allows us to examine the molecular organization of all cellular components at
molecular resolution, as well as the physical alignment of subsynaptic nanostructures[30-
32,61]. As such, this study provides direct evidence supporting the nanocolumn hypothesis
with higher-resolution structural information, particularly through direct visualization of
potential release sites indicated by tethered, docked, and fused vesicles aligning with PSD
nanoblocks both in synaptosomes and intact synapses from hippocampal cultures. We
also observed that multiple release sites aligning with a large PSD nanoblock is common
(Fig. 4). Though, it should be noted that fewer tethered, docked, or fused vesicles were
found in synaptosomes due to our isolation process. While most of these tethered vesicles
aligned with PSD nanoblocks and adhesion-like proteins, as well as a cluster of receptors
(Fig. 1e-i), there may be more undetected release sites (Fig. 1j-k). Further improvements
in cellular tomography will surely enable direct visualization of individual synaptic
proteins in cultured neurons or brain tissues. These discoveries will provide us with
ultrastructure information at a more natural state.

The PSD in the excitatory postsynaptic terminal is a dense area packed with postsynaptic

proteins, where hundreds of molecules have been identified[62,63]. Efforts have been made
to clarify the PSD's molecular organization and the synaptic functions that these PSD
proteins serve. The development of imaging techniques such as super-resolution
fluorescent microscopy has facilitated the discovery of subsynaptic protein clusters, or
nanodomains[3-5,64]. These nanodomains are ~70 to 160 nm in size, which is larger than
most protein nanoblocks we visualized within the PSDs of synaptosomes. Therefore, we
postulate that the small nanoblocks we discovered within the PSD may be a structural
unit that forms the nanodomain, which in turn forms the PSD. A nanoblock consists of
certain copies of several different scaffold proteins that form a supramolecular structure,
which links to certain copies of synaptic receptors and aligns with the presynaptic release
sites through adhesion molecules. However, in considering the diversity of PSD protein
complexes[65-68], it remains possible that the composition of different-sized PSD
nanoblocks varies.

Chemical fixation, or high-pressure freezing and freeze substitution EM sample
preparation, displays a PSD mesh model[20,24,69]. Using cryo-ET, we also observed vertical
filaments within synaptosomes and intact synapses, which were smaller in size. We
suspect that the size difference is due to the inherent differences between sample
preparation methods and imaging modalities. In the liquid condensate mode of PSD[27-29],
the liquid droplet is at micron scale. It is possible that the synaptic proteins in cells form
nanoblocks or nanoclusters following the same mechanism as in the *in vitro* PSD protein
mixing assay.

Subtomogram averaging of receptor-like particles produced a 24-Å and a 26-Å structure
from a small number of particles (Supplementary Fig.3). We believe that we failed to
achieve higher resolutions due to AMPARs having different reconfigurations secondary
to interactions between AMPARs' flexible extracellular domains and other proteins[44,47,70-
73]. The ligand proteins and neighboring proteins also contribute to the density, rendering
each receptor structurally heterogenous. Overcoming these difficulties in observing high-
resolution structures of synaptic receptors, as well as linking AMPARs' *in situ*
conformation information with their functional states, would require an enormous cryo-
ET dataset of synaptosomes or synapses with proper thickness.

Here we propose a nanoscale organization model at excitatory synapses (Fig. 4s). The
model is supported by our data from synaptosomes and cultured neurons (Fig. 1-4),
showing the commonality and diversity across synapses. Different-sized PSD nanoblocks
and their organizing model within the transcellular alignment at the synapse suggest a
potential mechanism for PSD nanoblocks to regulate synaptic strength. Evidence from
super-resolution and EM studies also suggests that synaptic activity affects the number
and size of PSD nanoclusters or densities[4,74-76]. In addition, this model may explain two
seemingly contradictory properties of the PSD: stable, yet dynamic. PSD proteins form
stable, similarly sized nanoblocks, enabling them to recruit other postsynaptic proteins to
form dispersed nanoblocks or clustered nanoblocks (Fig. 4s) in response to neuronal
activity[77].

Figure Legends

Fig. 1. A presynaptic release site correlates with adhesion molecules, membrane proteins, and PSD nanoblocks.

(a) A tomographic slice showing an excitatory synaptosome. (b) 3D segmentation of the synaptosome shown in panel A. (c) Quantification of the synaptic cleft and the PSD shown in panel A. The zoomed-in tomographic slice showing the excitatory synaptosome from panel A is on the left and its density plot showing gray values along the red solid line is on the right. (d) Cleft widths of synaptosomes and synaptoneurosomes. (e-k) Examples of transsynaptic alignment of adhesion molecules and postsynaptic membrane proteins in the cleft, as well as PSD nanoblocks in the postsynapse. Tethered synaptic vesicles also align with cleft and PSD densities. Scale bars: (a-b) 100 nm; (e-g) 20 nm. *Abbreviations PreSM: Presynaptic membrane. PostSM: Postsynaptic membrane. PostMP: Postsynaptic membrane protein.*

Fig. 2. PSD subsynaptic organization in synaptosomes.

(a) Top panel: A tomographic slice showing the subsynaptic nanoblocks of a synaptosome. Arrows and numbers indicate the seven protein clusters in the PSD. Bottom panel: 3D segmentation of the synaptosome in the top panel. (b) Density plot shows average gray values along the 40-nm blue bar in the top panel of (a). (c) 3D density analysis in the PSD of synaptosome in (a). Top panel: Density plot shows gray values of 40-nm PSD density projected to the xz plane. The blue dotted line indicates the plane of the tomographic slice shown in (a). Bottom panel: DBSCAN clustering result of 40-nm PSD density projected to the xz plane. (d) Examples of PSD nanoblocks in 2D tomographic slices and corresponding projected DBSCAN clustering results. Values in each panel indicate the width (Units: nm) of the PSD block. The blue dotted lines indicate the plane of the corresponding tomographic slices. (e) Distribution of PSD nanoblock area. (f) Distribution of PSD nanoblock width. (g) Distribution of nanoblock height. (h) Scatter plot of PSD nanoblock area to its height. Vertical lines are the peak values of gaussian fitting in (e). Schematics in four corners show corresponding nanoblock morphology in the scatter plot. (i) Distribution of nearest neighbor distance between PSD nanoblock centroids. (j) Distribution of distance between nearest pixels of PSD nanoblocks. Scale bars: (a, c) 50 nm; (d) 20 nm.

Fig. 3. Subtomogram averaging of receptor-like particles on the postsynaptic membrane.

(a) Receptor-like particles on the postsynaptic membrane. (b-i) Subtomogram averaged type A receptor and type B receptor. Models from PDB bank are fitted to the averaged electron densities. Yellow: PDB 5ide. Red: PDB 6qkz. (j) Examples of type A receptors, type B receptors, and undefined particles on the postsynaptic clusters. (k-l) Nearest neighbor distances of type A receptor (k) and type B receptor (l). (m) Type A and type B receptor numbers in each synaptosome. Scale bars: (a) top panel 100 nm, bottom panel 20 nm; (b-i) 5 nm; (j) 50 nm.

Fig. 4. PSD subsynaptic organization in intact neurons.

(a) Tomographic slice of an excitatory synapse. (b) Segmentation of the synapse in (a). (c) Zoom-in view of the synapse in (a). (d) IsoNet-corrected tomographic slice of (c). Red arrowheads indicate PSD nanoblocks. (e) Clustering results of 40-nm PSD density projected to the plane parallel to the postsynaptic membrane. Each color represents one cluster. Docked or tethered vesicle positions are marked as circles. (f) Tomographic slice of a second excitatory synapse. (g) Segmentation of the synapse in (f). (h) Zoom-in view of the synapse in (f). (i) IsoNet-corrected tomographic slice of (h). Red arrowheads indicate PSD nanoblocks. (j) Clustering results of 40-nm PSD density projected to the plane parallel to the postsynaptic membrane. Each color represents one cluster. Docked or tethered vesicle positions are marked as circles. (k) Tomographic slice of a third excitatory synapse. (l) Segmentation of the synapse in (k). (m) Zoom-in view of the synapse in (k). (n) IsoNet-corrected tomographic slice of (m). Red arrowheads indicate PSD nanoblocks. (o) Clustering results of 40-nm PSD density projected to the plane parallel to the postsynaptic membrane. Each color represents one cluster. Docked or tethered vesicle positions are marked as circles. (p) Distribution of PSD nanoblock area. (q) Distribution of nearest neighbor distance between PSD nanoblock centroids. (r) Distribution of distance between nearest pixels of PSD nanoblocks. (s) Working model of subsynaptic organizations. Parameter ranges of nanoblock edge distance, centroid distance and nanoblock width are FWHM (the full width at half maximum) of single-peak gaussian fitting. Scale bars: (a, b, f, g, k, l) 100 nm; (c-e, h-j, m-o) 50 nm.

Acknowledgments

We thank Drs. Ege T. Kavalali, Terunaga Nakagawa, William Wan, Roger J Colbran and Mark E. Bowen for stimulating discussions and critical reading of the manuscript; the NIH for support (R00 MH113764 to Q.Z.); the Vanderbilt Faculty Fellowship Endowment Fund; and "Neurodegenerative" TIPS Initiative Award for support.

EM data collections were conducted at the Center for Structural Biology Cryo-EM Facility at Vanderbilt University. We also acknowledge the use of the Glacios cryo-TEM, which was acquired by NIH grant 1 S10 OD030292-01.

Author contributions

Conceptualization, R.S. and Q.Z.; Methodology, R.S. and Q.Z.; Investigation, R.S. and Q.Z.; Experiments, R.S. and B.A., Data analysis, R.S., L.W., M.H., X.W. and ZM.Z.; Writing – Original Draft, R.S. and Q.Z.; Writing – Review & Editing, R.S., L.W., M.H., X.W., ZM.Z. and Q.Z.; Funding Acquisition, Q.Z..

Competing interests

The authors declare no competing interests.

Methods

Synaptosome isolation

10-week Sprague-Dawley rats with either sex were used for synaptosome isolation. The rats were kept in 12 hours: 12 hours dark:light cycle and provided with treats as well as cardboard enrichments before the experiments. The rats were deeply anesthetized with isoflurane. Hippocampi tissue were dissected in the ice-cold homogenization buffer containing 0.32 M sucrose, 4 mM Hepes, 1mM EDTA, at pH 7.4 Then the tissue was homogenized at ~500 rpm for 4 strokes using an overhead stirrer (Electron Microscopy Sciences 6480610) and a tissue grinder (Electron Microscopy Sciences 6479310) in the ice-cold homogenization buffer. After homogenization, the homogenates were centrifuged at 800g at 4 °C for 10 min. The supernatant was immediately taken for cryo-ET sample preparation. All animal procedures were performed in accordance with the guide for the care and use of laboratory animals and were approved by the Institutional Animal Care and Use Committee at Vanderbilt University.

Primary dissociated neuronal cultures

The protocol was modified from previous works[22,78,79]. Postnatal day 0 Sprague-Dawley rats of either sex were used for primary hippocampal cultures. Pregnant Sprague-Dawley rats were housed individually until they give birth to a litter and were kept in 12 hours: 12 hours dark:light cycle. The pregnant rats were provided with the same treats as well as cardboard enrichments. Postnatal day 0 littermates were used to prepare primary dissociated neuronal cultures. Hippocampi were dissected in ice cold 20% fetal bovine serum (FBS) containing Hanks' balanced salt solution. Tissues were then washed and treated with 10 mg/ml trypsin and 0.5 mg/ml DNase at 37 °C for 10 min. The tissues were washed again, dissociated using filtered P1000 tip and centrifuged at 1000 rpm for 10 mins at 4 °C. Pellet containing neurons was resuspended in Neurobasal Plus medium supplemented with GlutaMAX-I and B27 supplement. Neurons were plated onto Quantifoil R2/2 Au 200 EM grids coated with poly-L-lysine. Cultures were kept in humidified incubators at 37 °C and gassed with 95% air and 5% CO2. On DIV1, the neurobasal plus medium was replaced with 4 μM cytosine arabinoside containing neurobasal plus medium. On DIV4, the cytosine arabinoside concentration was dropped 2 μM by performing a half media change. Cultures were kept without any disruption until DIV14. The cultures on EM grids were plunged frozen between DIV14-18, when synapses reached maturity. All animal procedures were performed in accordance with the guide for the care and use of laboratory animals and were approved by the Institutional Animal Care and Use Committee at Vanderbilt University.

Synaptosome and neuronal culture vitrification

Synaptosome

FEI Vitrobot Mark III was used for the plunge freezing. Set the parameters of the plunge freezer as follows: humidity 100%, temperature 4 °C, blot time 3.5 s, blot total 1, blot

offset -1.5, and wait time 6 s. Supernatant after centrifugation mixed with 10-nm gold
beads solution (Aurion, cat # 25486) at 1:1 ratio was added to the mounted grid
(Quantifoil R2/2 Cu 200 EM). The grids were plunged into liquid nitrogen cooled liquid
ethane for rapid vitrification and were stored in liquid nitrogen until use.

Primary dissociated neuronal cultures

The vitrification process of neuronal cultures is similar to the synaptosome with a few
modifications. The culture dishes were taken out from the incubator on DIV 14-18. The
culture medium was replaced with a modified Tyrode's solution containing the followings
was used: (in mM): 150 NaCl, 4 KCl, 1.25 $MgCl_2$, 2 $CaCl_2$, 10 D-glucose, 10 Hepes at
pH 7.4. After setting up the plunge freezer as follows: humidity 100%, temperature
22 °C, blot time 3.5 s, blot total 1, blot offset -1.5, and wait time 6 s, the grids with
cultures were mounted. 4 μL 10-nm gold beads solution was added to the mounted grid
before plunging. After rapid vitrification, the grids were stored in liquid nitrogen until
use.

Cryo-ET data collection

Automated batch tilt series collection was done with Thermo Fisher Tomography5
software on Thermo Fisher Titan Krios G4 using the Gatan K3 camera in zero-loss mode
(slit width 20 eV). The tilt series of neuronal cultures were collected using the dose-
symmetric scheme[80] starting from 0° to ± 60° with an interval of 2°, and with the defocus
value at -7 μm, with the total electron dosage of ~150 $e^-/Å^2$ which were evenly
distributed between tilts. The final pixel size was 3.3 Å with 26000× magnification. The
tilt series of synaptosomes were collected with Voltage Phase Plate. The phase plate
dataset was collected at the pixel size 3.3 Å with 26000× magnification, using the dose-
symmetric scheme starting from 0° to ± 60°with an interval of 2°, and with the defocus
value at -2 μm. The total electron dosage of ~150 $e^-/Å^2$ was evenly distributed between
tilts. The phase plate was switched to a new position and activated to gain a phase shift of
around 0.3π prior to each tilt series. Conditioning of the phase plate was also performed
between each tilt.

Data reconstruction and subtomogram averaging

Tomograms of synaptosome at pixel size of 3.3 Å with phase plate were used for
subtomogram averaging. Synaptosome thicker than 500 nm were excluded, and 28
tomograms were selected. TOMOMAN package[81] was used to preprocess the data. The
workflow of TOMOMAN included motion correction by MotionCorr2[82], image sorting,
dose filtering and CTF estimation by Gctf [83]. Then the tilt series were aligned the
reconstructed using IMOD[84]. Fine alignment of the tilt series was performed by using the
10 nm gold beads (Aurion, cat # 25486) as fiducial markers. Both back projection and

back projection with five-iteration SIRT (simultaneous iterative reconstruction
technique)-like were performed to reconstruct each tilt series. Segmentation and 3D
rendering were done using IMOD. In the synaptosome dataset, 1565 receptor-like
particles were picked according to the size and shape. Only the particles around 10 ~14
nm on the postsynaptic membrane with LBD (ligand binding domain) and ATD (Amino
terminal domain) were included. Subtomogram averaging was done using RELION
3.0.8[85]. We applied a box size of 96 using unbinned data. First, we generated the initial
model using 3d auto-refine with the unbinned data without applying symmetry. 3d
classification resulted in two good classes. After 3d classification, we generated 3d initial
models for each class respectively with C2 symmetry and did 3d auto-refine for each
class using the corresponding initial models. Manual repicking of the particles was
performed to exclude the particles which were too close (distance < 8 nm) and final
averaging was performed using the remaining particles in 3d auto-refine.

PSD density analysis and quantification

Density of PSD was measured using following method. In the synaptosome dataset,
synaptosome with a PSD close to cryo grid carbon hole were excluded as the gray values
were affected by the hole edge. Synaptosomes thicker than 500 nm and synaptosomes
with the postsynaptic membrane not parallel to its presynaptic membrane were also
excluded. 16 tomograms were selected for PSD density clustering analysis. In the culture
neuron dataset, synapses thicker than 700 nm were excluded, and 34 tomograms were
selected for PSD density analysis. The gray values within 40 nm range from the
postsynaptic membrane were projected to xz plane (Fig. 2) using Fiji[86]. The gray value of
each pixel of the projected 2D plane was the average value of the 40 nm column. Next,
we used different strategies to process synaptosome and cultured neuron dataset. For
synaptosome data, we measured the average gray value in empty space as background
gray value. Then we chose the bottom 25% gray values (25% darkest) below the
background value for the following clustering analysis. For synapses from cultured
neurons, we analyzed the density in the IsoNet[57] corrected tomograms. The data set was
normalized after IsoNet internal process, and we chose the bottom 20% gray values (20%
darkest) for the following clustering analysis. Lastly, clustering analysis algorithm
DBSCAN (density-based spatial clustering of applications with noise) was adopted to do
the clustering of PSD density. In the synaptosome sample analysis, DBSCAN with
different parameters were performed (Supplementary Fig. 2) and DBSCAN (6, 32) was
applied as the results are consistent with the visualized electron density clusters in the
tomographic slices. In the synapse analysis, DBSCAN (7, 32) was applied. Positions of
tethered, docked, or fused vesicles were projected to the xz plane in the same coordinate
system of PSD density to analyze the relation between release sites and PSD clusters.
Positions of receptors were also projected to the xz plane in the same coordinate system
of PSD density to analyze the relation between receptors and PSD clusters. The area of a
nanoblock was defined as its convex hull area shown by red lines in Fig. 2b-c and
Supplementary Fig. 2b bottom panel. The width of a nanoblock was defined as the largest
x-axis distance between two pixels in the cluster. The widths of nanoblocks are also
marked in Figures 2C and the bottom panel of 2D. To precisely measure the height of a

nanoblock, we first averaged the tomographic slices in the range that contains the
nanoblock determined in the DBSCAN clustering results. Then we drew a line as shown
in Fig. 1c to measure the gray value of the nanoblock in y axis at its largest length. The
height of the nanoblock was defined as the distance from the farthest pixel of the
nanoblock to the postsynaptic membrane.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request.

References

Dieterich, D. C. & Kreutz, M. R. Proteomics of the Synapse--A Quantitative Approach to Neuronal Plasticity. *Molecular & cellular proteomics : MCP* **15**, 368-381 (2016). https://doi.org:10.1074/mcp.R115.051482

Perez de Arce, K. *et al.* Topographic Mapping of the Synaptic Cleft into Adhesive Nanodomains. *Neuron* **88**, 1165-1172 (2015). https://doi.org:10.1016/j.neuron.2015.11.011

Tang, A. H. *et al.* A trans-synaptic nanocolumn aligns neurotransmitter release to receptors. *Nature* **536**, 210-214 (2016). https://doi.org:10.1038/nature19058

MacGillavry, H. D., Song, Y., Raghavachari, S. & Blanpied, T. A. Nanoscale scaffolding domains within the postsynaptic density concentrate synaptic AMPA receptors. *Neuron* **78**, 615-622 (2013). https://doi.org:10.1016/j.neuron.2013.03.009

Nair, D. *et al.* Super-resolution imaging reveals that AMPA receptors inside synapses are dynamically organized in nanodomains regulated by PSD95. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 13204-13224 (2013). https://doi.org:10.1523/JNEUROSCI.2381-12.2013

Chamma, I. *et al.* Mapping the dynamics and nanoscale organization of synaptic adhesion proteins using monomeric streptavidin. *Nature communications* **7**, 10773 (2016). https://doi.org:10.1038/ncomms10773

Maschi, D. & Klyachko, V. A. Spatiotemporal Regulation of Synaptic Vesicle Fusion Sites in Central Synapses. *Neuron* **94**, 65-73 e63 (2017). https://doi.org:10.1016/j.neuron.2017.03.006

Hruska, M., Henderson, N., Le Marchand, S. J., Jafri, H. & Dalva, M. B. Synaptic nanomodules underlie the organization and plasticity of spine synapses. *Nature neuroscience* **21**, 671-682 (2018). https://doi.org:10.1038/s41593-018-0138-9

Glebov, O. O. *et al.* Nanoscale Structural Plasticity of the Active Zone Matrix Modulates Presynaptic Function. *Cell Rep* **18**, 2715-2728 (2017).

https://doi.org:10.1016/j.celrep.2017.02.064

10   Martinez-Sanchez, A. *et al.* Trans-synaptic assemblies link synaptic vesicles and
neuroreceptors. *Sci Adv* **7** (2021). https://doi.org:10.1126/sciadv.abe6204

11   Frank, R. A. & Grant, S. G. Supramolecular organization of NMDA receptors and
the postsynaptic density. *Curr Opin Neurobiol* **45**, 139-147 (2017).
https://doi.org:10.1016/j.conb.2017.05.019

12   Kellermayer, B. *et al.* Differential Nanoscale Topography and Functional Role of
GluN2-NMDA Receptor Subtypes at Glutamatergic Synapses. *Neuron* **100**, 106-
119 e107 (2018). https://doi.org:10.1016/j.neuron.2018.09.012

13   Chen, H., Tang, A. H. & Blanpied, T. A. Subsynaptic spatial organization as a
regulator of synaptic strength and plasticity. *Curr Opin Neurobiol* **51**, 147-153
(2018). https://doi.org:10.1016/j.conb.2018.05.004

14   Biederer, T., Kaeser, P. S. & Blanpied, T. A. Transcellular Nanoalignment of
Synaptic Function. *Neuron* **96**, 680-696 (2017).
https://doi.org:10.1016/j.neuron.2017.10.006

15   Guzikowski, N. J. & Kavalali, E. T. Nano-Organization at the Synapse: Segregation
of Distinct Forms of Neurotransmission. *Front Synaptic Neurosci* **13**, 796498
(2021). https://doi.org:10.3389/fnsyn.2021.796498

16   Gramlich, M. W. & Klyachko, V. A. Nanoscale Organization of Vesicle Release at
Central Synapses. *Trends Neurosci* **42**, 425-437 (2019).
https://doi.org:10.1016/j.tins.2019.03.001

17   Ramsey, A. M. *et al.* Subsynaptic positioning of AMPARs by LRRTM2 controls
synaptic strength. *Sci Adv* **7** (2021). https://doi.org:10.1126/sciadv.abf3126

18   Haas, K. T. *et al.* Pre-post synaptic alignment through neuroligin-1 tunes synaptic
transmission efficiency. *eLife* **7** (2018). https://doi.org:10.7554/eLife.31755

19   Palay, S. L. Synapses in the central nervous system. *J Biophys Biochem Cytol* **2**,
193-202 (1956). https://doi.org:10.1083/jcb.2.4.193

20   Chen, X. *et al.* Organization of the core structure of the postsynaptic density.
Proceedings of the National Academy of Sciences of the United States of America
105, 4453-4458 (2008). https://doi.org:10.1073/pnas.0800897105

21   DeGiorgis, J. A., Galbraith, J. A., Dosemeci, A., Chen, X. & Reese, T. S.
Distribution of the scaffolding proteins PSD-95, PSD-93, and SAP97 in isolated
PSDs. *Brain Cell Biol* **35**, 239-250 (2006). https://doi.org:10.1007/s11068-007-
9017-0

22   Tao, C.-L. *et al.* Differentiation and Characterization of Excitatory and Inhibitory
Synapses by Cryo-electron Tomography and Correlative Microscopy. *Journal of*
Neuroscience **38**, 1493-1510 (2018). https://doi.org:10.1523/jneurosci.1548-
17.2017

23   Chen, X. *et al.* PSD-95 is required to sustain the molecular organization of the
postsynaptic density. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **31**, 6329-6338 (2011).
https://doi.org:10.1523/JNEUROSCI.5968-10.2011

24   Burette, A. C. *et al.* Electron tomographic analysis of synaptic ultrastructure. *J*
Comp Neurol **520**, 2697-2711 (2012). https://doi.org:10.1002/cne.23067

25   Pennacchietti, F. *et al.* Nanoscale Molecular Reorganization of the Inhibitory
Postsynaptic Density Is a Determinant of GABAergic Synaptic Potentiation. *The*

Journal of neuroscience : the official journal of the Society for Neuroscience **37**, 1747-1756 (2017). https://doi.org:10.1523/JNEUROSCI.0514-16.2016

Crosby, K. C. *et al.* Nanoscale Subsynaptic Domains Underlie the Organization of the Inhibitory Synapse. *Cell Rep* **26**, 3284-3297 e3283 (2019). https://doi.org:10.1016/j.celrep.2019.02.070

Zeng, M. *et al.* Reconstituted Postsynaptic Density as a Molecular Platform for Understanding Synapse Formation and Plasticity. *Cell* **174**, 1172-1187 e1116 (2018). https://doi.org:10.1016/j.cell.2018.06.047

Zeng, M. *et al.* Phase Transition in Postsynaptic Densities Underlies Formation of Synaptic Complexes and Synaptic Plasticity. *Cell* **166**, 1163-1175 e1112 (2016). https://doi.org:10.1016/j.cell.2016.07.008

Feng, Z., Chen, X., Zeng, M. & Zhang, M. Phase separation as a mechanism for assembling dynamic postsynaptic density signalling complexes. *Curr Opin Neurobiol* **57**, 1-8 (2019). https://doi.org:10.1016/j.conb.2018.12.001

Taylor, K. A. & Glaeser, R. M. Electron diffraction of frozen, hydrated protein crystals. *Science* **186**, 1036-1037 (1974). https://doi.org:10.1126/science.186.4168.1036

Dubochet, J. *et al.* Cryo-electron microscopy of vitrified specimens. *Q Rev Biophys* **21**, 129-228 (1988). https://doi.org:10.1017/s0033583500004297

Dubochet, J. & Sartori Blanc, N. The cell in absence of aggregation artifacts. *Micron* **32**, 91-99 (2001). https://doi.org:10.1016/s0968-4328(00)00026-3

Bleck, C. K. *et al.* Comparison of different methods for thin section EM analysis of Mycobacterium smegmatis. *Journal of microscopy* **237**, 23-38 (2010). https://doi.org:10.1111/j.1365-2818.2009.03299.x

Peet, M. J., Henderson, R. & Russo, C. J. The energy dependence of contrast and damage in electron cryomicroscopy of biological molecules. *Ultramicroscopy* **203**, 125-131 (2019). https://doi.org:10.1016/j.ultramic.2019.02.007

Fernandez-Busnadiego, R. *et al.* Quantitative analysis of the native presynaptic cytomatrix by cryoelectron tomography. *The Journal of cell biology* **188**, 145-156 (2010). https://doi.org:10.1083/jcb.200908082

Gray, E. G. & Whittaker, V. P. The isolation of nerve endings from brain: an electron-microscopic study of cell fragments derived by homogenization and centrifugation. *J Anat* **96**, 79-88 (1962).

Evans, G. J. The synaptosome as a model system for studying synaptic physiology. *Cold Spring Harb Protoc* **2015**, 421-424 (2015). https://doi.org:10.1101/pdb.top074450

Hollingsworth, E. B. *et al.* Biochemical characterization of a filtered synaptoneurosome preparation from guinea pig cerebral cortex: cyclic adenosine 3':5'-monophosphate-generating systems, receptors, and enzymes. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **5**, 2240-2253 (1985). https://doi.org:10.1523/JNEUROSCI.05-08-02240.1985

Quinlan, E. M., Olstein, D. H. & Bear, M. F. Bidirectional, experience-dependent regulation of N-methyl-D-aspartate receptor subunit composition in the rat visual cortex during postnatal development. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 12876-12880 (1999). https://doi.org:10.1073/pnas.96.22.12876

Petersen, J. D. *et al.* Distribution of Postsynaptic Density (PSD)-95 and Ca^{2+}/Calmodulin-Dependent Protein Kinase II at the PSD. *The Journal of Neuroscience* **23**, 11270-11278 (2003). https://doi.org:10.1523/jneurosci.23-35-11270.2003

Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. in *Proceedings of the Second International Conference on Knowledge Discovery and Data Mining* 226–231 (AAAI Press, Portland, Oregon, 1996).

Rahbek-Clemmensen, T. *et al.* Super-resolution microscopy reveals functional organization of dopamine transporters into cholesterol and neuronal activity-dependent nanodomains. *Nat Commun* **8**, 740 (2017). https://doi.org:10.1038/s41467-017-00790-3

Natalie, J. G. a. E. T. K. Super-resolution imaging of synaptic scaffold proteins in rat hippocampal neurons. *STAR Protocols* **4**, 102080 (2023). https://doi.org:https://doi.org/10.1016/j.xpro.2023.102080

Garcia-Nafria, J., Herguedas, B., Watson, J. F. & Greger, I. H. The dynamic AMPA receptor extracellular region: a platform for synaptic protein interactions. *J Physiol* **594**, 5449-5458 (2016). https://doi.org:10.1113/JP271844

Zhao, Y., Chen, S., Swensen, A. C., Qian, W. J. & Gouaux, E. Architecture and subunit arrangement of native AMPA receptors elucidated by cryo-EM. *Science* **364**, 355-362 (2019). https://doi.org:10.1126/science.aaw8250

Greger, I. H., Watson, J. F. & Cull-Candy, S. G. Structural and Functional Architecture of AMPA-Type Glutamate Receptors and Their Auxiliary Proteins. *Neuron* **94**, 713-730 (2017). https://doi.org:10.1016/j.neuron.2017.04.009

Nakagawa, T., Cheng, Y., Sheng, M. & Walz, T. Three-dimensional structure of an AMPA receptor without associated stargazin/TARP proteins. *Biological chemistry* **387**, 179-187 (2006). https://doi.org:10.1515/BC.2006.024

Sobolevsky, A. I., Rosconi, M. P. & Gouaux, E. X-ray structure, symmetry and mechanism of an AMPA-subtype glutamate receptor. *Nature* **462**, 745-756 (2009). https://doi.org:10.1038/nature08624

Herguedas, B. *et al.* Structure and organization of heteromeric AMPA-type glutamate receptors. *Science* **352**, aad3873 (2016). https://doi.org:10.1126/science.aad3873

Lee, C. H. *et al.* NMDA receptor structures reveal subunit arrangement and pore architecture. *Nature* **511**, 191-197 (2014). https://doi.org:10.1038/nature13548

Karakas, E. & Furukawa, H. Crystal structure of a heterotetrameric NMDA receptor ion channel. *Science* **344**, 992-997 (2014). https://doi.org:10.1126/science.1251915

El-Husseini, A. E., Schnell, E., Chetkovich, D. M., Nicoll, R. A. & Bredt, D. S. PSD-95 involvement in maturation of excitatory synapses. *Science* **290**, 1364-1368 (2000).

Schnell, E. *et al.* Direct interactions between PSD-95 and stargazin control synaptic AMPA receptor number. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 13902-13907 (2002). https://doi.org:10.1073/pnas.172511199

Beique, J. C. *et al.* Synapse-specific regulation of AMPA receptor function by PSD-95. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 19535-19540 (2006). https://doi.org:10.1073/pnas.0608492103

714   55   Elias, G. M. *et al.* Synapse-specific and developmentally regulated targeting of
715   AMPA receptors by a family of MAGUK scaffolding proteins. *Neuron* **52**, 307-320
716   (2006). https://doi.org:10.1016/j.neuron.2006.09.012
717   56   Bats, C., Groc, L. & Choquet, D. The interaction between Stargazin and PSD-95
718   regulates AMPA receptor surface trafficking. *Neuron* **53**, 719-734 (2007).
719   https://doi.org:10.1016/j.neuron.2007.01.030
720   57   Liu, Y. T. *et al.* Isotropic reconstruction for electron tomography with deep learning.
721   *Nature communications* **13**, 6482 (2022). https://doi.org:10.1038/s41467-022-
722   33957-8
723   58   Frank, R. A. W., Zhu, F., Komiyama, N. H. & Grant, S. G. N. Hierarchical
724   organization and genetically separable subfamilies of PSD95 postsynaptic
725   supercomplexes. *J Neurochem* **142**, 504-511 (2017).
726   https://doi.org:10.1111/jnc.14056
727   59   Fernandez, E. *et al.* Targeted tandem affinity purification of PSD-95 recovers core
728   postsynaptic complexes and schizophrenia susceptibility proteins. *Mol Syst Biol* **5**,
729   269 (2009). https://doi.org:10.1038/msb.2009.27
730   60   Frank, R. A. *et al.* NMDA receptors are selectively partitioned into complexes and
731   supercomplexes during synapse maturation. *Nature communications* **7**, 11264
732   (2016). https://doi.org:10.1038/ncomms11264
733   61   Turk, M. & Baumeister, W. The promise and the challenges of cryo-electron
734   tomography. *FEBS letters* **594**, 3243-3261 (2020). https://doi.org:10.1002/1873-
735   3468.13948
736   62   Dosemeci, A., Weinberg, R. J., Reese, T. S. & Tao-Cheng, J. H. The Postsynaptic
737   Density: There Is More than Meets the Eye. *Front Synaptic Neurosci* **8**, 23 (2016).
738   https://doi.org:10.3389/fnsyn.2016.00023
739   63   Vyas, Y. & Montgomery, J. M. The role of postsynaptic density proteins in neural
740   degeneration and regeneration. *Neural Regen Res* **11**, 906-907 (2016).
741   https://doi.org:10.4103/1673-5374.184481
742   64   Broadhead, M. J. *et al.* PSD95 nanoclusters are postsynaptic building blocks in
743   hippocampus circuits. *Scientific reports* **6**, 24626 (2016).
744   https://doi.org:10.1038/srep24626
745   65   Sheng, M. & Hoogenraad, C. C. The postsynaptic architecture of excitatory
746   synapses: a more quantitative view. *Annual review of biochemistry* **76**, 823-847
747   (2007). https://doi.org:10.1146/annurev.biochem.76.060805.160029
748   66   Kaizuka, T. & Takumi, T. Postsynaptic density proteins and their involvement in
749   neurodevelopmental disorders. *J Biochem* **163**, 447-455 (2018).
750   https://doi.org:10.1093/jb/mvy022
751   67   Zeng, M., Ye, F., Xu, J. & Zhang, M. PDZ Ligand Binding-Induced Conformational
752   Coupling of the PDZ-SH3-GK Tandems in PSD-95 Family MAGUKs. *Journal of
753   molecular biology* **430**, 69-86 (2018). https://doi.org:10.1016/j.jmb.2017.11.003
754   68   Hassani Nia, F. *et al.* Structural deficits in key domains of Shank2 lead to alterations
755   in postsynaptic nanoclusters and to a neurodevelopmental disorder in humans.
756   *Molecular psychiatry* (2022). https://doi.org:10.1038/s41380-022-01882-3
757   69   Chen, X. *et al.* PSD-95 family MAGUKs are essential for anchoring AMPA and
758   NMDA receptor complexes at the postsynaptic density. *Proceedings of the National
759   *Academy of Sciences of the United States of America* **112**, E6983-6992 (2015).

https://doi.org:10.1073/pnas.1517045112

70 Dutta, A. *et al.* Cooperative Dynamics of Intact AMPA and NMDA Glutamate
Receptors: Similarities and Subfamily-Specific Differences. *Structure* **23**, 1692-
1704 (2015). https://doi.org:10.1016/j.str.2015.07.002

71 Krieger, J., Bahar, I. & Greger, I. H. Structure, Dynamics, and Allosteric Potential
of Ionotropic Glutamate Receptor N-Terminal Domains. *Biophysical journal* **109**,
1136-1148 (2015). https://doi.org:10.1016/j.bpj.2015.06.061

72 Durr, K. L. *et al.* Structure and dynamics of AMPA receptor GluA2 in resting, pre-
open, and desensitized states. *Cell* **158**, 778-792 (2014).
https://doi.org:10.1016/j.cell.2014.07.023

73 Meyerson, J. R. *et al.* Structural mechanism of glutamate receptor activation and
desensitization. *Nature* **514**, 328-334 (2014). https://doi.org:10.1038/nature13603

74 Geinisman, Y. *et al.* Structural synaptic correlate of long-term potentiation:
formation of axospinous synapses with multiple, completely partitioned
transmission zones. *Hippocampus* **3**, 435-445 (1993).
https://doi.org:10.1002/hipo.450030405

75 Stewart, M. G. *et al.* Chemically induced long-term potentiation increases the
number of perforated and complex postsynaptic densities but does not alter
dendritic spine volume in CA1 of adult mouse hippocampal slices. *Eur J Neurosci*
21, 3368-3378 (2005). https://doi.org:10.1111/j.1460-9568.2005.04174.x

76 Nieto-Sampedro, M., Hoff, S. F. & Cotman, C. W. Perforated postsynaptic densities:
probable intermediates in synapse turnover. *Proceedings of the National Academy*
of Sciences of the United States of America **79**, 5718-5722 (1982).
https://doi.org:10.1073/pnas.79.18.5718

77 Sturgill, J. F., Steiner, P., Czervionke, B. L. & Sabatini, B. L. Distinct Domains
within PSD-95 Mediate Synaptic Incorporation, Stabilization, and Activity-
Dependent Trafficking. *The Journal of Neuroscience* **29**, 12845-12854 (2009).
https://doi.org:10.1523/jneurosci.1841-09.2009

78 Alten, B. *et al.* Role of Aberrant Spontaneous Neurotransmission in SNAP25-
Associated Encephalopathies. *Neuron* **109**, 59-72 e55 (2021).
https://doi.org:10.1016/j.neuron.2020.10.012

79 Sun, R. *et al.* An efficient protocol of cryo-correlative light and electron microscopy
for the study of neuronal synapses. *Biophysics Reports* **5**, 111-122 (2019).
https://doi.org:10.1007/s41048-019-0092-4

80 Hagen, W. J. H., Wan, W. & Briggs, J. A. G. Implementation of a cryo-electron
tomography tilt-scheme optimized for high resolution subtomogram averaging.
Journal of structural biology **197**, 191-198 (2017).
https://doi.org:10.1016/j.jsb.2016.06.007

81 williamnwan/TOMOMAN: TOMOMAN 08042020 v. 08042020 (Zenodo, 2020).

82 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).
https://doi.org:10.1038/nmeth.4193

83 Zhang, K. Gctf: Real-time CTF determination and correction. *Journal of structural*
biology **193**, 1-12 (2016). https://doi.org:10.1016/j.jsb.2015.11.003

84 Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of
three-dimensional image data using IMOD. *Journal of structural biology* **116**, 71-

806   76 (1996). https://doi.org:10.1006/jsbi.1996.0013

807   85   Bharat, T. A. & Scheres, S. H. Resolving macromolecular structures from electron
808   cryo-tomography data using subtomogram averaging in RELION. *Nat Protoc* **11**,
809   2054-2065 (2016). https://doi.org:10.1038/nprot.2016.124

810   86   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
811   *Methods* **9**, 676-682 (2012). https://doi.org:10.1038/nmeth.2019

Fig. 1

Fig. 2

a

b

c

d Tomo-slice

Top view projection

15.8   19.8   22.4   27.7   43.6   52.8   59.4   63.4

e

f

g

h

i

j

Fig. 3

j
- Type A receptor
- Type B receptor
- Undefined particle

Fig. 4

p Relative Frequency (%) — Nanoblock area (nm²)

q Relative Frequency (%) — Nanoblock centroid distance (nm)

r Relative Frequency (%) — Nanoblock nearest pixel distance (nm)

s
Dispersed Nanoblocks   Clustered Nanoblocks

1 – PSD nanoblock width 8-35 nm
2 – Nanoblock edge distance
        3-17 nm (synaptosome)
        4-16 nm (intact synapse)
3 – Nanoblock centroid distance
        27-63 nm (synaptosome)
        25-53 nm (intact synapse)

Supplementary Fig. 1

Supplementary Fig. 2

a

b

Supplementary Fig. 3

a

Subtomogram average

1565 particles picked | 3D auto refine without symmetry

3D classification with C2 symmetry

458 particles | 219 particles

2 good classes

Remove duplicates and particles not on the membrane

391 particles | 189 particles

Post processing

?

Putative type A receptor | Putative type B receptor | Undefined particles
391 particles | 189 particles | 888 particles

b

Putative type A receptor   Final resolution = 24 Å

- Corrected
- Unmasked Maps
- Masked Maps
- Phase Randomized Masked Maps

Putative type B receptor   Final resolution = 26 Å

- Corrected
- Unmasked Maps
- Masked Maps
- Phase Randomized Masked Maps

Supplementary Fig. 4