Cryo-electron tomography reveals postsynaptic nanoblocks in excitatory synapses

Rong Sun[1,2], James P Allen[2], Liana Wilson[1,2], Mariam Haider[1,3], Baris Alten[4,6], Zimeng

Zhou[1,5], Xinyi Wang[1,6], Qiangjun Zhou[1,2*].

[1] Department of Cell and Developmental Biology, Center for Structural Biology,

Vanderbilt Kennedy Center, Vanderbilt University, Nashville, TN, 37240, USA.

[2] Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN, 37240, USA

[3] Center for Structural Biology Cryo-EM Facility, Vanderbilt University, Nashville, TN,

37240, USA.

[4] Department of Pharmacology, Vanderbilt University, Nashville, TN, 37212, USA.

[5] School of Engineering, Vanderbilt University, Nashville, TN, 37212, USA.

[6] Peabody College, Vanderbilt University, Nashville, TN, 37212, USA

[7] Present address: Harvard Medical School, Boston, MA, 02115, USA and Department of

Neurology, Massachusetts General Hospital and Brigham & Women's Hospital, Boston,

MA, 02115, USA

Abstract

The nanoscale organization of proteins within synapses is critical for maintaining and
regulating synaptic transmission and plasticity. Here, we use cryogenic electron
tomography to directly visualize the three-dimensional architecture and supramolecular
organization of pre-cleft-postsynaptic components in their near-native cellular context in
both synaptosomes from rat hippocampi and synapses from rat primary cultured neurons.
High-resolution electron microscopy and quantitative analyses revealed that postsynaptic
density (PSD) is composed of membrane-associated nanoblocks of various sizes, which
are in close relationship with potential presynaptic release sites through adhesion
molecules spanning the synaptic cleft, as well as with post-synaptic receptors.
Subtomogram averaging from synaptosomes showed two distinct types of postsynaptic
membrane proteins at resolutions of 24 Å and 26 Å respectively. Furthermore, our data
reveal that majority of potential release sites and ~50% subtomogram averaged receptor-
like particles are located within the boundary of PSD nanoblocks, while PSD nanoblocks
might be redundant for neurotransmission. The results of this study provide a more
comprehensive understanding of synaptic ultrastructure and suggest that PSD is
composed of clustering of various nanoblocks, which likely underlies the dynamic nature
of PSD to modulate synaptic strength.

Introduction

Cellular function is driven by nanoscale spatial organization of multi-protein assemblies
and their interactions with other molecular components and cellular organelles. Typical
examples of such nanoscale structures are neuronal chemical synapses. These are the
most complex cell-cell junctions, with more than 2,000 distinct synaptic proteins[1].
Compelling evidence suggests that that multiple presynaptic, postsynaptic, and cell-
adhesion proteins in excitatory synapses are finely organized in lipid-bound nanoscale
structures to facilitate extremely rapid, dynamic, efficient, and tightly regulated
transmission of information between neurons[2-10]. Moreover, the activity-dependent
rearrangement of these synaptic proteins, including neurotransmitter receptors and other
synaptic proteins, plays a key mechanistic role in synaptic plasticity, which is essential
for perception, decision making, learning, and memory formation.

Recent studies provide evidence for a high-level organization of nanoscale
supramolecular assemblies spanning the synaptic cleft, giving rise to *nanocolumns* of
juxtaposed pre- and postsynaptic compartments in excitatory synapses[3,4,11-16].
Specifically, presynaptic active-zone scaffold proteins, such as RIM, form nanodomains
that define synaptic vesicle docking and release sites. These sites align with clustered
neurotransmitter receptors such as α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic
acid receptor (AMPAR) and postsynaptic scaffolding proteins such as PSD95 via
adhesion proteins in the synaptic cleft[10,17,18].

The postsynaptic density (PSD) appears in conventional electron micrographs as an
electron-dense lamina just beneath the postsynaptic membrane[19]. A PSD meshwork
model comprising regularly spaced vertical filaments was proposed based on electron
microscopy (EM) and biochemical assays[20-24]. Two additional forms of PSD organization
have been proposed: nano-domains based on super-resolution imaging[3-5,25-27], and liquid
condensate based on *in vitro* PSD mixing assay[28-30]. All these studies provide more
evidence for the high-level organization of nanoscale supramolecular assemblies in the
synapse, indicating that synaptic proteins are not randomly distributed and instead form
functional synaptic nanostructures. However, nanoscale organization of the PSD and
postsynaptic receptors is largely unknown due to the technical challenges imposed by
synaptic molecular complexity.

Cryo-electron tomography (cryo-ET) is a powerful technique that produces three-
dimensional (3D) volume reconstructions, or *tomograms*, of macromolecular and
subcellular biological structures in their native context at nanometer resolution. To
prevent introducing artifacts through chemical fixation, dehydration, and heavy-metal
staining, biological samples are rapidly frozen to obtain a "true, instantaneous
snapshot"[31-34]. In general, obtaining high-quality data requires cryo-ET samples to be thin
enough for electrons to pass through (~200 nm), posing a technical challenge during
sample preparation, which limits the resolution of cellular cryo-ET[35]. The resolution is
also limited by the crowded cellular environment. To overcome these limitations, isolated
synaptic terminals have been used for cryo-ET studies to decrease sample thickness[10,36].

A classic type of isolated synaptic terminal called a *synaptosome* is highly suitable for
cryo-ET, because it retains a piece of attached postsynaptic membrane that faces the
active zone[37,38]. Previous studies have demonstrated that PSD complexes and PSD
membranes are resistant to detergents and stable enough to be purified from conventional
synaptic fractions fractions[39-42]. This stability suggests that the core scaffold of the native
PSD is likely maintained on the attached postsynaptic membrane[42]. Consequently,
synaptosomes are ideal for cryo-ET data collection at high resolution, as the average
thickness of a synaptosome is 200-300 nm in our preparations. Here, we developed a
simplified sample preparation workflow to obtain the synaptosomes from mature rat
hippocampi, achieving high-resolution synaptic nanostructures with cryo-ET and sub-
tomogram averaging. With this method, we directly visualized the overall architecture of
the transcellular organization in excitatory synapses. Our results revealed that PSD is
formed by subsynaptic PSD nanoblocks of specific sizes. We also determined structures
of two types of postsynaptic membrane proteins at 24-Å and 26-Å resolution.

Results

Transsynaptic organization of synaptosomes

To study subsynaptic organization, we employed a simplified workflow to prepare
synaptosomes. After dissecting the hippocampi from rat brains, we performed four
strokes of homogenization and a 10-minute centrifugation at 800 *g,* followed by plunge-
freezing and cryo-ET data acquisition (Supplementary Fig. 1a). Our preparations yielded
two different types of isolated synaptic terminals based on their morphology:
synaptosomes retained a patch of postsynaptic membrane with PSD (Fig. 1 and
Supplementary Fig. 1b-e, see also Supplementary Video 2), while synaptoneurosomes
retained an enclosed postsynaptic compartment (Supplementary Figs. 1f-i and 2g)[43,44].
For this study, we primarily focused on synaptosomes, since they can provide higher-
resolution structural information compared to synaptoneurosomes and intact synapses
due to lower samples thickness less molecular crowding. In total, we acquired 123
tomograms of isolated synaptic terminals from three preparations without protease
inhibitor or DTT in the homogenization buffer (99 synaptosomes and 24
synaptoneurosomes). This allowed us to directly visualize characteristic features of a
synapse, including typical synaptic proteins (Fig. 1a, 1b and Supplementary Fig. 1p-u). In
all our preparations, we successfully identified both synaptosomes and
synaptoneurosomes, and no noticeable morphological differences were observed at the
nanoscale across different batches of sample preparations. All the synaptosomes we
obtained were likely from excitatory synapses, as they have thick PSDs of more than 30
nm, which is a known feature of excitatory synapses[19,22].

To study the relationship between potential release sites, adhesion molecules,
postsynaptic densities, and postsynaptic receptors, we examined the tethered, docked or
fused synaptic vesicles that likely indicate the location of synaptic release sites. We also
investigated the corresponding nanostructures in the cleft and in the postsynaptic
compartment. Examining the electron densities and measuring the gray values of
synaptosome tomograms (Fig. 1c) allowed us to detect the adhesion molecule-like
proteins and membrane proteins in the synaptic cleft, as well as the PSD protein clusters,

which we refer as *nanoblocks*. The tomographic slice in Fig. 1c's left panel shows protein
densities in the middle of the cleft, which we posit to be the adhesion molecules, as well
as a density layer on the postsynaptic membrane that is comprised of membrane proteins.
These observations based on direct visualization are presented with the density plot in
Fig. 1c's right panel. We found that synaptosomes and synaptoneurosomes have specific
cleft widths (Fig. 1d); and the clefts of synaptosomes tend to be wider than those of
synaptoneurosomes (40.1±0.5 nm for synaptosomes n=74; and 31.5±0.9 nm for
synaptoneurosomes n=24). Of note, 15 synaptosomes were not included in these
measurements as the postsynaptic membranes were not parallel to the presynaptic
membranes. The disparity in cleft widths between synaptosomes and synaptoneurosomes
may result from distinct adhesion molecules that withstand the sample isolation process.
We observed that tethered vesicles are in close proximity of the adhesion-like molecules,
the membrane protein-like particles, and PSD densities (Fig. 1e-i). This direct
visualization of pre-cleft-postsynaptic structure suggests that PSD scaffold proteins,
might be specifically aligned with potential presynaptic release sites through adhesion
molecules in the synaptic cleft. Two examples of adhesion-PSD nanoblock alignment
without a tethered synaptic vesicle are also shown here, indicating these might be
potential unoccupied release sites (Fig. 1j-k). To quantify the relationship between PSD
nanoblock and potential release sites occupied by docked, tethered, or fused synaptic
vesicles, we analysed the distance from vesicle centroid to nanoblock centroids,
normalized by the distance between nanoblock centroid to nanoblock edge, and compared
with what is observed with random distribution. We found that 76.4% of potential release
sites were transsynaptically located in the boundary of a nanoblock (Fig. 1l-n). However,
considering all docked, tethered, or fused synaptic vesicles (n=106) in synaptosomes, the
alignment between measured positions of synaptic vesicles and nanoblocks did not show
significant difference from the alignment between randomized positions of synaptic
vesicles and nanoblocks (Fig. 1n), which suggests that PSD nanoblocks are redundant for
alignment with the potential release sites occupied by synaptic vesicles.

PSD subsynaptic organization in synaptosomes

Next, we investigated PSD densities and performed quantitative analyses on the PSD
nanoblock dimensions. A representative tomogram of a synaptosome acquired by cryo-
ET is shown in Fig. 2a, in which PSD nanoblocks (indicated by red arrows) are clearly
visualized beneath the membrane. In terms of the overall PSD EM density patterns, this
result is consistent with previous EM studies stating that the core of the PSD is
dominated by vertically oriented filaments[20,45] or spiky structures[24]. Our data revealed
three types of synaptosomes based on the organization of the PSD: synaptosomes with
separated PSD nanoblocks; synaptosomes with both separated and continuous PSD
nanoblocks; and synaptosomes with continuous PSD (Supplementary Fig. 1j-o).
Importantly, it is evident that the distribution of PSD densities is not uniform across these
three types of synaptosomes. The presence of PSD nanoblocks is clearly observable in
the original 3D tomograms (Supplementary Fig. 1jo). To delineate the PSD
nanostructures of all three synaptosome types, we applied the following quantitative
analysis: First, we measured gray values of protein densities of a 40-nm thick PSD area

beneath the postsynaptic membrane (Fig. 2a-b). Then, we combined 2D gray values
across a series of continuous virtual tomographic slices, generating a gray value density
map (Fig. 2c, top panel). With normalization and filtering processing, **d**ensity-**b**ased
spatial **c**lustering of **a**pplications with **n**oise (DBSCAN) results show projected PSD
density nanoblocks in the 40-nm range from the postsynaptic membrane (Fig. 2c, bottom
panel). DBSCAN is a density-based clustering non-parametric algorithm[46] and it is
widely used in biological data analysis[47,48]. In this way, 2D DBSCAN results represented
PSD densities flattened and projected to *xz-plane* (Fig. 2d). We next measured the area
and width of each protein nanoblock (Fig. 2c and 2e). The area of nanoblock was defined
as its convex hull area shown by encircling red lines. The width of nanoblock was
defined as the largest *x-axis* distance between two pixels in each cluster. Fig. 2e shows
examples of nanoblocks with different areas and widths quantified by the DBSCAN
clustering method. Multiple peaks were detected in the area distribution of nanoblocks
from all synaptosomes (Fig. 2f), while the width distribution was fit with two peaks (Fig.
2g). Distinct fitting peaks in the area statistics histogram indicate that nanoblocks have
distinct sizes, and that large nanoblocks may be composed of several small nanoblocks,
which is also supported by examining the morphology of some nanoblocks with large
widths (Fig. 2e). Though the geometric shape of the nanoblock was irregular, the way we
quantified the width reflected the dimension of the nanoblock in *x-axis*, while the
dimension of nanoblock in *z-axis* was affected by the missing wedge effect[49]. We also
tested DBSCAN with different parameters (Supplementary Fig. 2a-b) and DBSCAN (6,
32) was applied as the results were more consistent with the visualized electron density
clusters in the tomographic slices. We isolated synaptosome sample from three separate
preparations and obtained consistent cleft with and nanoblock sizes across all samples
(Supplementary Fig. 2c-d). To further rule out the influence of sample preparation on
these observed ultrastructural features, we collected an additional 83 tomograms of
isolated synaptic terminals (72 synaptosomes and 11 synaptoneurosomes) from three
preparations with protease inhibitor and DTT in the homogenization buffer. The data of
synaptosomes treated with protease inhibitors and DTT (Supplementary Fig. 2e-f), along
with the data of synaptoneurosomes (Supplementary Fig. 2g-h), collectively present
compelling evidence that supports the presence of nanoblocks across different sample
types, ruling out the possibility of isolation artifacts. Overall, our analyses suggest that
PSD nanoblocks can be positioned closely enough to one another to form larger
nanoblocks.

To further analyse the size and morphology of subsynaptic nanoblocks, we measured the
height of the protein nanoblocks obtained via DBSCAN clustering. The height was
quantified based on gray values along the nanoblock at its largest length (Fig. 1c). The
distribution of nanoblock heights revealed that most nanoblocks are in the 30~90 nm
range, while a small population is below 20 nm or above 100 nm (Fig. 2h). Additionally,
we analysed the gap between PSD nanoblocks. The distance between each nanoblock
centroid (Fig. 2i) and the nearest distance between each nanoblock edge (Fig. 2j) together
describe the spacing between nanoblocks, suggesting that the gaps with less protein
densities segmented the PSD. Overall, our analysis demonstrates that subsynaptic
nanoblocks of different sizes form the PSD.

Organization of membrane protein particles on the postsynaptic membrane

To further investigate the nanoscale organization of the PSD, we employed the subtomogram averaging method to detect postsynaptic membrane proteins in our synaptosome samples. We manually selected 1,565 protein particles from 28 synaptosomes in the range from 10 nm to 14 nm for subtomogram averaging (Fig. 3a and Supplementary Fig. 3a). Eventually, we achieved an averaged structure of type A particle with 391 particles (Fig. 3b) and an averaged structure of type B particle with 189 particles (Fig. 3c).

To study the postsynaptic distribution of the membrane particles, we superimposed the type A particles, type B particles, and all remaining undefined particles with the PSD clustering results (Fig. 3d). To quantify the relationship between these two types of membrane particles and nanoblocks, we used the same method with which we analysed potential release sites (Fig. 3e-f). This analysis was performed in 13 synaptosomes which were used both in PSD nanoblock quantification and subtomogram averaging. We found that 64.7% type A particles and 45.2% type B particles are within nanoblocks ($d/D \leqslant 1$). Additionally, we found distribution of d/D for type A and type B particles are different (two-sample Kolmogorov–Smirnov test, D=0.2551, n=207 for type A particles and n=93 for type B particles, $p=3.54\times10^{-4}$) and type A particles are closer to nanoblocks compared to type B particles (two-sample t test, n=207, mean=0.89 nm for type A particles and n=93, mean=1.34 nm for type B particles, $p=2.52\times10^{-5}$). We performed a nearest neighbor distance analysis and found type A particles are more clustered than type B particles (Fig. 3i-j, two-sample t test, mean for type A is 30.8 nm and mean for type B is 50.5 nm, p=0.00139). The number of type A and type B particles vary in all synaptosomes (Fig. 3k), showing the heterogeneity of the synaptosomes. However, the relatively low resolution achieved through subtomogram averaging renders it challenging to determine the protein identity of type A or type B particles with certainty.

PSD subsynaptic organization in intact neurons

We also examined the subsynaptic organization of synapses to determine whether similar subsynaptic nanoblocks identified in synaptosomes can also be found in intact synapses from cultured neurons. To accomplish this, we seeded dissociated primary hippocampal neurons on gold EM grids and examined the synaptic ultrastructure in intact neurons using cryo-ET (Supplementary Fig. 4a-e). In reconstructed tomograms, a typical synapse consists of a presynaptic compartment with a population of synaptic vesicles, a synaptic cleft, and a postsynaptic compartment with PSD. We also focused on excitatory synapses and excluded synapses with thin PSDs in cultured neurons from our analysis (Supplementary Fig. 4f-h).

We collected 211 tilt series of synapse-like structures from the wild-type primary hippocampal neurons and reconstructed them. 131 of the reconstructed tomograms were synapses, and 121 synapses had thick PSDs. Three examples of excitatory synapses are shown in Fig. 4. Tomographic slices (Fig. 4a, c, f, h, k, and m) and segmented structures

(Fig. 4b and g, see also Supplementary Video 2) display EM densities within the PSD
area are not uniformly distributed, suggesting that protein distribution is not
homogeneous within the PSD. However, the crowded molecular environment and overall
higher thickness of intact neurons rendered a low signal-to-noise ratio, making patterns of
protein complexes difficult to recognize and analyse. Using the software package
IsoNet[50], we performed contrast transfer function (CTF) deconvolution and corrected the
missing wedge effect to enhance contrast in 34 out of 121 cryo-electron tomograms of
excitatory synapses. Following IsoNet correction, tomographic slices show more robust
protein nanoblocks in the PSDs (Fig. 4d, i, and n). Quantitative characterization of the
PSDs' 3D volume (Fig. 4e, j, and o) indicates that proteins distribute unevenly within the
PSD and form subsynaptic nanoblocks. The size and distribution statistics of the PSD
nanoblocks (Fig. 4p-r) closely resemble those obtained from synaptosomes and
synaptoneurosomes (Fig. 2f and Supplementary Fig. 2e-h). Our observations suggest that
intact synapses also feature nanoblocks of varying sizes composed of postsynaptic
proteins, similar to those observed in synaptosomes and synaptoneurosomes.
Furthermore, we observed multiple tethered vesicles, docked vesicles, or fused vesicles
(Supplementary Fig. 4i-n) which serve as potential occupied release sites and are
transsynaptically located in the boundary of PSD nanoblocks (Fig. 4s-t). In comparison to
~1.8 potential release sites occupied by synaptic vesicles per synapse in synaptosomes,
intact synapses in cultured neuron exhibit ~4.3 potential release sites occupied by
synaptic vesicles per synapse. Considering all docked, tethered, or fused synaptic vesicles
(n=145) as potential release sites, we employed the same analysis of normalized vesicle-
nanoblock centroid distances, and found no differences between the distribution of
observed vesicle-nanoblock distances and those with randomized release sites (Fig. 1n).
Like our previous results in synaptosomes, this further indicates that in hippocampal
neuronal cultures the PSD nanoblocks are redundant for alignment with the potential
release sites.

Discussion

In this study, we utilized cryo-ET to directly visualize the ultrastructure of excitatory
synapses in their near-native state of purified hippocampal synaptosomes from 10-week-
old young adult rats as well as primary hippocampal neurons cultured from postnatal day
0 newborn rats. Importantly, we found that in our synaptosome samples, the exposed
postsynaptic membranes remain attached to the presynaptic terminals. Thus, we were
able to directly visualize membrane-associated PSD components in these preparations,
which we refer as PSD nanoblocks. However, we acknowledge that some postsynaptic
components located in the gap between PSD nanoblocks might be lost during sample
preparation. This allowed us to study the PSD components against a relatively clean
background. Meanwhile, studies of the whole synapse in intact cultured neurons provided
more physiological and complete ultrastructure features. With these two different
samples, we consistently captured a subsynaptic organization of PSD in the form of
nanoblocks, which are facing presynaptic release sites. Our observations suggest that
membrane-associated PSD components within synaptosomes are relatively stable, and
that they maintain structural integrity during the isolation process for synaptosome

preparation via our simplified protocol. This finding is consistent with previous
observations that PSD complexes and the PSD membranes are detergent-resistant and
stable enough to be purified from conventional synaptic fractions[39-42]. Thus, it is likely
that the nanoscale organization and basic ultrastructure features we observed also reflect
synaptic architecture *in vivo*.

An alignment between pre- and postsynaptic elements has been proposed but was only
recently reported and named a *nanocolumn*, discovered through super-resolution light
microscopy[3]. Thus far, only labeled synaptic proteins such as PSD95, receptors, and RIM
have been shown to form nanoclusters as a part of nanocolumns. In contrast, our cryo-ET
approach allows us to examine the molecular organization of all cellular components at
molecular resolution, nanostructures not only those can be labeled for super-resolution
light microscopy[31-33,51]. Martinez-Sanchez et al. applied cryo-ET to examine the
alignment between presynaptic vesicles and postsynaptic receptors obtained through
subtomogram averaging in synaptoneurosomes[10]. Here, we showed that the majority of
the potential release sites, indicated by tethered, docked and fused vesicles, are
transsynaptically located in the boundary of the PSD nanoblocks in both synaptosomes
and intact neurons (Figs. 1 and 4). However, our analyses showed that this alignment is
statistically indistinguishable from a random distribution of potential release sites and
PSD nanoblocks. There might be several plausible explanations to this seemingly
unreasonable result. First, the postsynaptic density might be redundant for
neurotransmission, since majority of nanoblocks have less than 20 nm distance in
between each other, encompassing the entire postsynaptic density in a near-continuous
fashion. Second, technical limitations of sample preparation might have obscured a
physiologically relevant alignment. For example, we have found fewer tethered, docked
and fused vesicles in synaptosomes compared to intact neurons, suggesting that although
the presence and distribution of postsynaptic nanoblocks are consistent throughout, the
transsynaptic alignment might be susceptible to various steps in sample preparation. We
also observed potential release sites that weren't occupied by vesicles in synaptosomes
(Fig. 1j-k), further confounding the analyses of transsynaptic alignment. Third and last,
limitations of cryo-ET might have contributed to experimental absence of a specific
transsynaptic alignment between release sites and PSD nanoblocks. For example, we
were unable to further dissect the constituents of individual PSD nanoblocks, which
might differ from one nanoblock to another, indicating distinct synaptic functions for
each nanoblock requiring a specific transsynaptic alignment. Further improvements in
cellular tomography will surely enable direct visualization of individual synaptic proteins
ex vivo without requiring additional steps of sample preparation that might alter nano-
scale ultrastructure of a synapse, providing with near native-state structures. Future
advances in cryo-ET might reveal a specific alignment for different subgroups of
postsynaptic nanoblocks with different release sites.

The PSD in the excitatory postsynaptic terminal is a dense area packed with postsynaptic
proteins, where hundreds of molecules have been identified[52,53]. Efforts have been made
to clarify the PSD's molecular organization and the synaptic functions that these PSD
proteins serve. The development of imaging techniques such as super-resolution
fluorescent microscopy has facilitated the discovery of subsynaptic protein clusters, or
nanodomains[3-5,8,54]. These nanodomains are ~70 to 160 nm in size, which is larger than

most protein nanoblocks we directly visualized within the PSDs of synaptosomes using
cryo-ET. Therefore, we postulate that the small nanoblocks we discovered within the
PSD may be a structural unit that forms the nanodomain, which in turn forms the PSD. A
nanoblock consists of certain copies of several different scaffold proteins that form a
supramolecular structure, which links to certain copies of synaptic receptors and
allineates with the presynaptic release sites through adhesion molecules. Other
postsynaptic proteins situated in the gaps between these nanoblocks may play a role in
the regulation and organization of smaller nanoblocks into larger ones. However,
considering the diversity of PSD protein complexes[55-58], it remains possible that the
composition of different-sized PSD nanoblocks varies. Further exploration of the
composition of PSD nanoblocks will help us uncover the distinctions and similarities
between nanoblocks and nanodomains.

Chemical fixation, or high-pressure freezing and freeze substitution EM sample
preparation, displays a PSD mesh model[20,24,59]. Using cryo-ET, we also observed vertical
filaments within synaptosomes and intact synapses, which were smaller in size. We
suspect that the size difference is due to the inherent differences between sample
preparation methods and imaging modalities. In the liquid condensate mode of PSD[28-30],
the liquid droplet is at micron scale. It is possible that the synaptic proteins in cells form
nanoblocks or nanoclusters following the same mechanism as in the *in vitro* PSD protein
mixing assay.

Subtomogram averaging of postsynaptic protein particles produced a 24-Å and a 26-Å
structure from a small number of particles (Supplementary Fig.3). Our inability to
achieve higher resolutions to identify protein composition is attributed to the complexity
and diversity of postsynaptic proteins and protein complexes present on the postsynaptic
membrane of brain tissue samples, which include various types of glutamate receptors[60-67].
The ligand proteins and neighboring proteins also contribute to the density, rendering
each postsynaptic protein structurally heterogenous. Moreover, distinct postsynaptic
receptors exhibit varying degrees of retention in biochemical PSD preparations[68-71].
Therefore, it is challenging to identify the protein complex based on the current
resolution of our subtomogram averaging. When we compared the geometric shape of the
average structure with existing PDB models, type A particles and type B particles seem to
resemble the O-shaped and the Y-shaped AMPA receptors, respectively (Supplementary
Fig. 3c-h). Similar to the comparison made by Martinez-Sanchez *et al.*[10], our averaged
structures don't look like other available structures of postsynaptic complexes. It is likely
the distribution of type A and type B particles shown in Fig. 3e-h reflected the
distribution of different types of AMPA receptors in synapses. This is also consistent with
studies showing AMPA receptor nano-organization does not fully align with PSD95
nano-organization[5,8,27]. Overcoming these difficulties in observing high-resolution
structures of postsynaptic receptors, as well as linking their *in situ* conformation
information with their functional states, would require an enormous cryo-ET dataset of
intact synapses from cultured neurons or even brain tissues with proper thickness.

Here we propose a nanoscale organization model at excitatory synapses (Fig. 4u). The
model is supported by our data from isolated synaptic terminals, and cultured neurons
(Figs. 1-4), showing the commonality and diversity across synapses. Evidence from

super-resolution and EM studies also suggests that synaptic activity affects the number
and size of PSD nanoclusters or densities[4,72-74]. Thus, we postulate that the nanoblocks
that we identified in both synaptosomes and intact neurons might be the fundamental unit
of postsynaptic density and might play a role in regulating the synaptic strength.

Figure Legends

Fig. 1. A potential presynaptic release site links with adhesion molecules, membrane
proteins, and PSD nanoblocks.

(a) A tomographic slice showing an excitatory synaptosome. (b) 3D segmentation of the
synaptosome shown in panel a. (c) Quantification of the synaptic cleft and the PSD
shown in panel a. The zoomed-in tomographic slice showing the excitatory synaptosome
from panel a is on the left and its density plot showing gray values along the red solid line
is on the right. (d) Cleft widths of synaptosomes and synaptoneurosomes. (e-k) Examples
of transsynaptic alignment of adhesion molecules and postsynaptic membrane proteins in
the cleft, as well as PSD nanoblocks in the postsynapse. Tethered synaptic vesicles also
align with cleft and PSD densities. Red arrowheads indicate putative adhesion molecules
in the cleft. (l) A schematic showing distance between synaptic vesicle centroid and
nanoblock centroid, and distance between nanoblock centroid and edge. (m-n)
Distribution of normalized distance between potential release site and nanoblock. $d/D \leqslant 1$
indicates that the SV centroid aligns within the PSD nanoblock. Scale bars: (a-b) 100 nm;
(e-g) 20 nm. *Abbreviations PreSM: Presynaptic membrane. PostSM: Postsynaptic*
membrane. PostMP: Postsynaptic membrane protein.

Fig. 2. PSD subsynaptic organization in synaptosomes.

(a) Top panel: A tomographic slice showing the subsynaptic nanoblocks of a
synaptosome. Arrows and numbers indicate the seven nanoblocks in the PSD. Bottom
panel: 3D segmentation of the synaptosome in the top panel. (b) Density plot shows
average gray values (black lines) along the 40-nm blue bar in the top panel of (a). Red
lines indicate the fitting curve of average gray values. (c) 3D density analysis in the PSD
of synaptosome in (a). Top panel: Density plot shows gray values of 40-nm PSD density
projected to the xz plane. The blue dotted line indicates the plane of the tomographic slice
shown in (a). Bottom panel: DBSCAN clustering result of 40-nm PSD density projected
to the xz plane. (d) A schematic showing the clustering results of PSD densities produced
from 3D synaptosome data. (e) Examples of PSD nanoblocks in 2D tomographic slices
and corresponding projected DBSCAN clustering results. Values in each panel indicate
the width (Units: nm) of the PSD block. The blue dotted lines indicate the plane of the
corresponding tomographic slices. (f) Distribution of PSD nanoblock area. (g)
Distribution of PSD nanoblock width. (h) Distribution of nanoblock height. (i)
Distribution of nearest neighbor distance between PSD nanoblock centroids. (j)
Distribution of distance between nearest pixels of PSD nanoblocks. Scale bars: (a, c) 50
nm; (e) 20 nm.

Fig. 3. Subtomogram averaging of postsynaptic membrane particles.

(a) Membrane particles on the postsynaptic membrane. (b-c) Subtomogram averaged type
A and B particles. (d) Examples of type A, type B, and undefined particles on the
postsynaptic PSD nanoblocks. (e-h) Distribution of normalized distance between type A
or type B particles and nanoblock. (i-j) Nearest neighbor distances of type A particle (j)
and type B particle (j). (k) Type A and type B particle numbers in each synaptosome.
Scale bars: (a) left panel 100 nm, right panel 20 nm; (b-c) 5 nm; (d) 50 nm.

Fig. 4. PSD subsynaptic organization in intact neurons.

(a) Tomographic slice of an excitatory synapse. (b) Segmentation of the synapse in (a).
(c) Zoom-in view of the synapse in (a). (d) IsoNet-corrected tomographic slice of (c). Red
arrowheads indicate PSD nanoblocks. (e) Clustering results of 40-nm PSD density
projected to the plane parallel to the postsynaptic membrane. Each color represents one
cluster. Docked or tethered vesicle positions are marked as circles. (f) Tomographic slice
of a second excitatory synapse. (g) Segmentation of the synapse in (f). (h) Zoom-in view
of the synapse in (f). (i) IsoNet-corrected tomographic slice of (h). Red arrowheads
indicate PSD nanoblocks. (j) Clustering results of 40-nm PSD density projected to the
plane parallel to the postsynaptic membrane. Each color represents one cluster. Docked or
tethered vesicle positions are marked as circles. (k) Tomographic slice of a third
excitatory synapse. (l) Segmentation of the synapse in (k). (m) Zoom-in view of the
synapse in (k). (n) IsoNet-corrected tomographic slice of (m). Red arrowheads indicate
PSD nanoblocks. (o) Clustering results of 40-nm PSD density projected to the plane
parallel to the postsynaptic membrane. Each color represents one cluster. Docked or
tethered vesicle positions are marked as circles. (p) Distribution of PSD nanoblock area.
(q) Distribution of nearest neighbor distance between PSD nanoblock centroids. (r)
Distribution of distance between nearest pixels of PSD nanoblocks. (s-t) Distribution of
normalized distance between potential release site and nanoblock. (u) Working model of
subsynaptic organizations. Scale bars: (a, b, f, g, k, l) 100 nm; (c-e, h-j, m-o) 50 nm.

Acknowledgments

We thank Drs. Ege T. Kavalali, Terunaga Nakagawa, William Wan, Roger J Colbran and
Mark E. Bowen for stimulating discussions and critical reading of the manuscript; the
NIH for support (R00 MH113764 to Q.Z., R01 MH132918-01A1 to Q.Z.); the Vanderbilt
Faculty Fellowship Endowment Fund; and "Neurodegenerative" TIPS Initiative Award
for support.

EM data collections were conducted at the Center for Structural Biology Cryo-EM
Facility at Vanderbilt University. We also acknowledge the use of the Glacios cryo-TEM,
which was acquired by NIH grant 1 S10 OD030292-01.

Author contributions

Conceptualization, R.S. and Q.Z.; Methodology, R.S. and Q.Z.; Investigation, R.S. and

Q.Z.; Experiments, R.S. and B.A., Data analysis, R.S., J.P.A., L.W., M.H., X.W. and
ZM.Z.; Writing – Original Draft, R.S. and Q.Z.; Writing – Review & Editing, R.S.,
J.P.A., L.W., M.H., X.W., ZM.Z. and Q.Z..; Funding Acquisition, Q.Z..

Competing interests

468 The authors declare no competing interests.

Methods

Synaptosome isolation

10-week Sprague-Dawley rats with either sex were used for synaptosome isolation. The
rats were kept in 12 hours: 12 hours dark:light cycle and provided with treats as well as
cardboard enrichments before the experiments. The rats were deeply anesthetized with
isoflurane. Hippocampi tissue were dissected in the ice-cold homogenization buffer
containing 0.32 M sucrose, 4 mM HEPES, 1mM EDTA, at pH 7.4 with or without
protease inhibitor cocktail (1 tablet in 50 ml homogenization buffer) and 20 mM DTT.
Then the tissue was homogenized at ~500 rpm for 4 strokes using an overhead stirrer
(Electron Microscopy Sciences 6480610) and a tissue grinder (Electron Microscopy
Sciences 6479310) in the ice-cold homogenization buffer. After homogenization, the
homogenates were centrifuged at 800g at 4 °C for 10 min. The supernatant was
immediately taken for cryo-ET sample preparation. All animal procedures were
performed in accordance with the guide for the care and use of laboratory animals and
were approved by the Institutional Animal Care and Use Committee at Vanderbilt
University.

Primary dissociated neuronal cultures

The protocol was modified from previous works[22,75,76]. Postnatal day 0 Sprague-Dawley
rats of either sex were used for primary hippocampal cultures. Pregnant Sprague-Dawley
rats were housed individually until they gave birth to a litter and were kept in 12 hours:
12 hours dark:light cycle. The pregnant rats were provided with the same treats as well as
cardboard enrichments. Postnatal day 0 littermates were used to prepare primary
dissociated neuronal cultures. Hippocampi were dissected in ice cold 20% fetal bovine
serum (FBS) containing Hanks' balanced salt solution. Tissues were then washed and
treated with 10 mg/ml trypsin and 0.5 mg/ml DNase at 37 °C for 10 min. The tissues
were washed again, dissociated using filtered P1000 tip and centrifuged at 1000 rpm for
10 mins at 4 °C. Pellet containing neurons was resuspended in Neurobasal Plus medium
supplemented with GlutaMAX-I and B27 supplement. Neurons were plated onto
Quantifoil R2/2 Au 200 EM grids coated with poly-L-lysine. Cultures were kept in
humidified incubators at 37 °C and gassed with 95% air and 5% CO_2. On DIV1, the
neurobasal plus medium was replaced with 4 μM cytosine arabinoside containing
neurobasal plus medium. On DIV4, the cytosine arabinoside concentration was dropped 2

μM by performing a half media change. Cultures were kept without any disruption until
DIV14. The cultures on EM grids were plunged frozen between DIV14-18, when
synapses reached maturity. All animal procedures were performed in accordance with the
guide for the care and use of laboratory animals and were approved by the Institutional
Animal Care and Use Committee at Vanderbilt University.

Synaptosome and neuronal culture vitrification

Synaptosome

FEI Vitrobot Mark III was used for plunge freezing. Set the parameters of the plunge
freezer as follows: humidity 100%, temperature 4 °C, blot time 3.5 s, blot total 1, blot
offset -1.5, and wait time 6 s. Supernatant after centrifugation mixed with 10-nm gold
beads solution (Aurion, cat # 25486) at 1:1 ratio was added to the mounted grid
(Quantifoil R2/2 Cu 200 EM). No other chemicals or stimulation were applied prior to
plunge freezing. The grids were plunged into liquid nitrogen cooled liquid ethane for
rapid vitrification and were stored in liquid nitrogen until use.

Primary dissociated neuronal cultures

The vitrification process of neuronal cultures is similar to the synaptosome with a few
modifications. The culture dishes were taken out from the incubator on DIV 14-18. The
culture medium was replaced with a modified Tyrode's solution containing the followings
was used: (in mM): 150 NaCl, 4 KCl, 1.25 $MgCl_2$, 2 $CaCl_2$, 10 D-glucose, 10 HEPES at
pH 7.4. Then the cultured neurons were kept in the modified Tyrode's solution for 10min.
After setting up the plunge freezer as follows: humidity 100%, temperature 22 °C, blot
time 3.5 s, blot total 1, blot offset -1.5, and wait time 6 s, the grids with cultures were
mounted. 4 μL 10-nm gold beads solution (in the modified Tyrode's solution) was added
to the mounted grid before plunging. No other chemicals or stimulation were applied
prior to plunge freezing. After rapid vitrification, the grids were stored in liquid nitrogen
until use.

Cryo-ET data collection

Automated batch tilt series collection was done with Thermo Fisher Tomography5
software on Thermo Fisher Titan Krios G4 using the Gatan K3 camera in zero-loss mode
(slit width 20 eV). The tilt series of neuronal cultures were collected using the dose-
symmetric scheme[77] starting from 0° to ± 60° with an interval of 2°, and with the defocus
value at -7 μm, with the total electron dosage of ~150 e⁻/Å² which were evenly
distributed between tilts. The final pixel size was 3.3 Å with 26000× magnification. The
tilt series of synaptosomes were collected with Voltage Phase Plate. The phase plate
dataset was collected at the pixel size 3.3 Å with 26000× magnification, using the dose-

symmetric scheme starting from 0° to ± 60°with an interval of 2°, and with the defocus
value at -2 μm. The total electron dosage of ~150 e⁻/Å² was evenly distributed between
tilts. The phase plate was switched to a new position and activated to gain a phase shift of
around 0.3π prior to each tilt series. Conditioning of the phase plate was also performed
between each tilt.

Data reconstruction and subtomogram averaging

Tomograms of synaptosome at pixel size of 3.3 Å with phase plate were used for
subtomogram averaging. Synaptosome thicker than 500 nm were excluded, and 28
tomograms were selected. TOMOMAN package[78] was used to preprocess the data. The
workflow of TOMOMAN included motion correction by MotionCorr2[79], image sorting,
dose filtering and CTF estimation by Gctf [80]. Then the tilt series were aligned the
reconstructed using IMOD[81]. Fine alignment of the tilt series was performed by using the
10 nm gold beads (Aurion, cat # 25486) as fiducial markers. Both back projection and
back projection with five-iteration SIRT (simultaneous iterative reconstruction
technique)-like were performed to reconstruct each tilt series. Segmentation and 3D
rendering were done using IMOD. In the synaptosome dataset, 1565 particles were
picked according to the size and shape. Only the particles around 10 ~14 nm on the
postsynaptic membrane were included. Subtomogram averaging was done using
RELION 3.0.8[82]. We applied a box size of 96 using unbinned data. First, we generated
the initial model using 3d auto-refine with the unbinned data without applying symmetry.
3d classification resulted in two good classes. After 3d classification, we generated 3d
initial models for each class respectively with C2 symmetry and did 3d auto-refine for
each class using the corresponding initial models. Manual repicking of the particles was
performed to exclude the particles which were too close (distance < 8 nm) and final
averaging was performed using the remaining particles in 3d auto-refine.

PSD density analysis and quantification

Density of PSD was measured using the following method. In the synaptosome dataset,
synaptosome with a PSD close to cryo grid carbon hole were excluded as the gray values
were affected by the hole edge. Synaptosomes thicker than 500 nm and synaptosomes
with the postsynaptic membrane not parallel to its presynaptic membrane were also
excluded. 123 tomograms (58 tomograms for synaptosome isolated without protease
inhibitor and DTT in the homogenization buffer, 41 tomograms for synaptosome isolated
with protease inhibitor and DTT in the homogenization buffer, and 24 tomograms for
synaptoneurosomes isolated with and without protease inhibitor and DTT in the
homogenization buffer) were selected for PSD density clustering analysis. In the culture
neuron dataset, synapses thicker than 700 nm were excluded, and 34 tomograms were
selected for PSD density analysis. The gray values within 40 nm range from the
postsynaptic membrane were projected to xz plane (Fig. 2) using Fiji[83]. The gray value of
each pixel of the projected 2D plane was the average value of the 40 nm column. Next,

we used different strategies to process synaptosome and cultured neuron dataset. For
synaptosome data, we measured the average gray value in empty space as background
gray value. Then we chose the bottom 25% gray values (25% darkest) below the
background value for the following clustering analysis. For synapses from cultured
neurons, we analysed the density in the IsoNet[50] corrected tomograms. The data set was
normalized after IsoNet internal process, and we chose the bottom 20% gray values (20%
darkest) for the following clustering analysis. Lastly, clustering analysis algorithm
DBSCAN (density-based spatial clustering of applications with noise) was adopted to do
the clustering of PSD density. In the synaptosome sample analysis, DBSCAN with
different parameters were performed (Supplementary Fig. 2) and DBSCAN (6, 32) was
applied as the results are consistent with the visualized electron density clusters in the
tomographic slices. In the synaptosome isolated with protease inhibitor and DTT,
synaptoneurosome and synapse analysis, DBSCAN (7, 32) was applied.

The area of a nanoblock was defined as its convex hull area shown by red lines in Fig.
2b-c and Supplementary Fig. 2b bottom panel. All area distributions were cut off to 3500
nm^2 or 6000 nm^2 to show the fitting peaks more clearly. The width of a nanoblock was
defined as the largest x-axis distance between two pixels in the cluster. The widths of
nanoblocks are also marked in Figures 2c and the bottom panel of 2d. To precisely
measure the height of a nanoblock, we first averaged the tomographic slices in the range
that contains the nanoblock determined in the DBSCAN clustering results. Then we drew
a line as shown in Fig. 1c to measure the gray value of the nanoblock in y axis at its
largest length. The height of the nanoblock was defined as the distance from the farthest
pixel of the nanoblock to the postsynaptic membrane. Since we took the weight of each
pixel in the nanoblock the same, the centroid of a nanoblock is the geometry centroid of
the nanoblock by averaging all coordinates of the pixels of a nanoblock. The distance
between the nanoblock edge and centroid of the nanoblock was defined as the distance
from the farthest pixel of the nanoblock to centroid of the nanoblock.

Positions of all tethered, docked, or fused vesicles were projected to the xz plane in the
same coordinate system of PSD density to analyse the relation between potential release
sites and PSD nanoblocks for both synaptosomes (vesicle number is 106) and synapses
(vesicle number is 145). d/D was calculated as shown in Fig. 1l and 4t. Here d was
defined as the distance from synaptic vesicle centroid to nanoblock centroid. D was
defined as the distance from farthest pixel of the nanoblock to centroid of the nanoblock.
Thus d/D is the normalized distance to from the synaptic vesicle to the nanoblock. To test
if the alignment between potential release sites and PSD nanoblock is specialized, the
position of the vesicles were randomized within the bounds of the entire PSD. We
randomized each synaptic vesicle position one hundred times, and then performed the
analysis for the randomized vesicle position to calculate d/D as we did for actual vesicles.
Positions of postsynaptic membrane proteins were also projected to the xz plane in the
same coordinate system of PSD density to analyse the relation between membrane
proteins and PSD nanoblocks. Then we calculate d/D for each protein particle. Here d
was defined as the distance from particle to nanoblock centroid. D was defined as the
distance from farthest pixel of the nanoblock to centroid of the nanoblock.

Data and code availability

All data and code reported in this paper will be shared by the lead contact upon request.

References

Dieterich, D. C. & Kreutz, M. R. Proteomics of the Synapse--A Quantitative Approach to Neuronal Plasticity. *Molecular & cellular proteomics : MCP* **15**, 368-381 (2016). https://doi.org/10.1074/mcp.R115.051482

Perez de Arce, K. *et al.* Topographic Mapping of the Synaptic Cleft into Adhesive Nanodomains. *Neuron* **88**, 1165-1172 (2015). https://doi.org/10.1016/j.neuron.2015.11.011

Tang, A. H. *et al.* A trans-synaptic nanocolumn aligns neurotransmitter release to receptors. *Nature* **536**, 210-214 (2016). https://doi.org/10.1038/nature19058

MacGillavry, H. D., Song, Y., Raghavachari, S. & Blanpied, T. A. Nanoscale scaffolding domains within the postsynaptic density concentrate synaptic AMPA receptors. *Neuron* **78**, 615-622 (2013). https://doi.org/10.1016/j.neuron.2013.03.009

Nair, D. *et al.* Super-resolution imaging reveals that AMPA receptors inside synapses are dynamically organized in nanodomains regulated by PSD95. *J Neurosci* **33**, 13204-13224 (2013). https://doi.org/10.1523/jneurosci.2381-12.2013

Chamma, I. *et al.* Mapping the dynamics and nanoscale organization of synaptic adhesion proteins using monomeric streptavidin. *Nature communications* **7**, 10773 (2016). https://doi.org/10.1038/ncomms10773

Maschi, D. & Klyachko, V. A. Spatiotemporal Regulation of Synaptic Vesicle Fusion Sites in Central Synapses. *Neuron* **94**, 65-73 e63 (2017). https://doi.org/10.1016/j.neuron.2017.03.006

Hruska, M., Henderson, N., Le Marchand, S. J., Jafri, H. & Dalva, M. B. Synaptic nanomodules underlie the organization and plasticity of spine synapses. *Nat Neurosci* **21**, 671-682 (2018). https://doi.org/10.1038/s41593-018-0138-9

Glebov, O. O. *et al.* Nanoscale Structural Plasticity of the Active Zone Matrix Modulates Presynaptic Function. *Cell Rep* **18**, 2715-2728 (2017). https://doi.org/10.1016/j.celrep.2017.02.064

Martinez-Sanchez, A. *et al.* Trans-synaptic assemblies link synaptic vesicles and neuroreceptors. *Sci Adv* **7** (2021). https://doi.org/10.1126/sciadv.abe6204

Frank, R. A. & Grant, S. G. Supramolecular organization of NMDA receptors and the postsynaptic density. *Curr Opin Neurobiol* **45**, 139-147 (2017). https://doi.org/10.1016/j.conb.2017.05.019

Kellermayer, B. *et al.* Differential Nanoscale Topography and Functional Role of GluN2-NMDA Receptor Subtypes at Glutamatergic Synapses. *Neuron* **100**, 106-119 e107 (2018). https://doi.org/10.1016/j.neuron.2018.09.012

Chen, H., Tang, A. H. & Blanpied, T. A. Subsynaptic spatial organization as a

regulator of synaptic strength and plasticity. *Curr Opin Neurobiol* **51**, 147-153
(2018). https://doi.org/10.1016/j.conb.2018.05.004
14 Biederer, T., Kaeser, P. S. & Blanpied, T. A. Transcellular Nanoalignment of
Synaptic Function. *Neuron* **96**, 680-696 (2017).
https://doi.org/10.1016/j.neuron.2017.10.006
15 Guzikowski, N. J. & Kavalali, E. T. Nano-Organization at the Synapse: Segregation
of Distinct Forms of Neurotransmission. *Front Synaptic Neurosci* **13**, 796498
(2021). https://doi.org/10.3389/fnsyn.2021.796498
16 Gramlich, M. W. & Klyachko, V. A. Nanoscale Organization of Vesicle Release at
Central Synapses. *Trends Neurosci* **42**, 425-437 (2019).
https://doi.org/10.1016/j.tins.2019.03.001
17 Ramsey, A. M. *et al.* Subsynaptic positioning of AMPARs by LRRTM2 controls
synaptic strength. *Sci Adv* **7** (2021). https://doi.org/10.1126/sciadv.abf3126
18 Haas, K. T. *et al.* Pre-post synaptic alignment through neuroligin-1 tunes synaptic
transmission efficiency. *eLife* **7** (2018). https://doi.org/10.7554/eLife.31755
19 Palay, S. L. Synapses in the central nervous system. *J Biophys Biochem Cytol* **2**,
193-202 (1956). https://doi.org/10.1083/jcb.2.4.193
20 Chen, X. *et al.* Organization of the core structure of the postsynaptic density.
Proceedings of the National Academy of Sciences of the United States of America
105, 4453-4458 (2008). https://doi.org/10.1073/pnas.0800897105
21 DeGiorgis, J. A., Galbraith, J. A., Dosemeci, A., Chen, X. & Reese, T. S.
Distribution of the scaffolding proteins PSD-95, PSD-93, and SAP97 in isolated
PSDs. *Brain Cell Biol* **35**, 239-250 (2006). https://doi.org/10.1007/s11068-007-
9017-0
22 Tao, C.-L. *et al.* Differentiation and Characterization of Excitatory and Inhibitory
Synapses by Cryo-electron Tomography and Correlative Microscopy. *Journal of*
Neuroscience **38**, 1493-1510 (2018). https://doi.org/10.1523/jneurosci.1548-
17.2017
23 Chen, X. *et al.* PSD-95 is required to sustain the molecular organization of the
postsynaptic density. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **31**, 6329-6338 (2011).
https://doi.org/10.1523/JNEUROSCI.5968-10.2011
24 Burette, A. C. *et al.* Electron tomographic analysis of synaptic ultrastructure. *J*
Comp Neurol **520**, 2697-2711 (2012). https://doi.org/10.1002/cne.23067
25 Pennacchietti, F. *et al.* Nanoscale Molecular Reorganization of the Inhibitory
Postsynaptic Density Is a Determinant of GABAergic Synaptic Potentiation. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience **37**,
1747-1756 (2017). https://doi.org/10.1523/JNEUROSCI.0514-16.2016
26 Crosby, K. C. *et al.* Nanoscale Subsynaptic Domains Underlie the Organization of
the Inhibitory Synapse. *Cell Rep* **26**, 3284-3297 e3283 (2019).
https://doi.org/10.1016/j.celrep.2019.02.070
27 Hruska, M., Cain, R. E. & Dalva, M. B. Nanoscale rules governing the organization
of glutamate receptors in spine synapses are subunit specific. *Nat Commun* **13**, 920
(2022). https://doi.org/10.1038/s41467-022-28504-4
28 Zeng, M. *et al.* Reconstituted Postsynaptic Density as a Molecular Platform for
Understanding Synapse Formation and Plasticity. *Cell* **174**, 1172-1187 e1116

(2018). https://doi.org/10.1016/j.cell.2018.06.047

29   Zeng, M. *et al.* Phase Transition in Postsynaptic Densities Underlies Formation of
Synaptic Complexes and Synaptic Plasticity. *Cell* **166**, 1163-1175 e1112 (2016).
https://doi.org/10.1016/j.cell.2016.07.008

30   Feng, Z., Chen, X., Zeng, M. & Zhang, M. Phase separation as a mechanism for
assembling dynamic postsynaptic density signalling complexes. *Curr Opin*
Neurobiol **57**, 1-8 (2019). https://doi.org/10.1016/j.conb.2018.12.001

31   Taylor, K. A. & Glaeser, R. M. Electron diffraction of frozen, hydrated protein
crystals. *Science* **186**, 1036-1037 (1974).
https://doi.org/10.1126/science.186.4168.1036

32   Dubochet, J. *et al.* Cryo-electron microscopy of vitrified specimens. *Q Rev Biophys*
21, 129-228 (1988). https://doi.org/10.1017/s0033583500004297

33   Dubochet, J. & Sartori Blanc, N. The cell in absence of aggregation artifacts.
Micron **32**, 91-99 (2001). https://doi.org/10.1016/s0968-4328(00)00026-3

34   Bleck, C. K. *et al.* Comparison of different methods for thin section EM analysis of
Mycobacterium smegmatis. *Journal of microscopy* **237**, 23-38 (2010).
https://doi.org/10.1111/j.1365-2818.2009.03299.x

35   Peet, M. J., Henderson, R. & Russo, C. J. The energy dependence of contrast and
damage in electron cryomicroscopy of biological molecules. *Ultramicroscopy* **203**,
125-131 (2019). https://doi.org/10.1016/j.ultramic.2019.02.007

36   Fernandez-Busnadiego, R. *et al.* Quantitative analysis of the native presynaptic
cytomatrix by cryoelectron tomography. *The Journal of cell biology* **188**, 145-156
(2010). https://doi.org/10.1083/jcb.200908082

37   Gray, E. G. & Whittaker, V. P. The isolation of nerve endings from brain: an
electron-microscopic study of cell fragments derived by homogenization and
centrifugation. *J Anat* **96**, 79-88 (1962).

38   Evans, G. J. The synaptosome as a model system for studying synaptic physiology.
Cold Spring Harb Protoc **2015**, 421-424 (2015).
https://doi.org/10.1101/pdb.top074450

39   Frank, R. A. W., Zhu, F., Komiyama, N. H. & Grant, S. G. N. Hierarchical
organization and genetically separable subfamilies of PSD95 postsynaptic
supercomplexes. *J Neurochem* **142**, 504-511 (2017).
https://doi.org/10.1111/jnc.14056

40   Fernandez, E. *et al.* Targeted tandem affinity purification of PSD-95 recovers core
postsynaptic complexes and schizophrenia susceptibility proteins. *Mol Syst Biol* **5**,
269 (2009). https://doi.org/10.1038/msb.2009.27

41   Frank, R. A. *et al.* NMDA receptors are selectively partitioned into complexes and
supercomplexes during synapse maturation. *Nature communications* **7**, 11264
(2016). https://doi.org/10.1038/ncomms11264

42   Jung, J. H., Chen, X. & Reese, T. S. Cryo-EM tomography and automatic
segmentation delineate modular structures in the postsynaptic density. *Front*
Synaptic Neurosci **15**, 1123564 (2023).
https://doi.org/10.3389/fnsyn.2023.1123564

43   Hollingsworth, E. B. *et al.* Biochemical characterization of a filtered
synaptoneurosome preparation from guinea pig cerebral cortex: cyclic adenosine
3':5'-monophosphate-generating systems, receptors, and enzymes. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience **5**, 2240-2253
(1985). https://doi.org/10.1523/JNEUROSCI.05-08-02240.1985

44 Quinlan, E. M., Olstein, D. H. & Bear, M. F. Bidirectional, experience-dependent
regulation of N-methyl-D-aspartate receptor subunit composition in the rat visual
cortex during postnatal development. *Proceedings of the National Academy of*
Sciences of the United States of America **96**, 12876-12880 (1999).
https://doi.org/10.1073/pnas.96.22.12876

45 Petersen, J. D. *et al.* Distribution of Postsynaptic Density (PSD)-95 and
Ca^{2+}/Calmodulin-Dependent Protein Kinase II at the PSD. *The*
Journal of Neuroscience **23**, 11270-11278 (2003).
https://doi.org/10.1523/jneurosci.23-35-11270.2003

46 Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. in *Proceedings of the Second*
International Conference on Knowledge Discovery and Data Mining   226–231
(AAAI Press, Portland, Oregon, 1996).

47 Rahbek-Clemmensen, T. *et al.* Super-resolution microscopy reveals functional
organization of dopamine transporters into cholesterol and neuronal activity-
dependent nanodomains. *Nat Commun* **8**, 740 (2017).
https://doi.org/10.1038/s41467-017-00790-3

48 Natalie, J. G. a. E. T. K. Super-resolution imaging of synaptic scaffold proteins in
rat hippocampal neurons. *STAR Protocols* **4**, 102080 (2023).
https://doi.org/https://doi.org/10.1016/j.xpro.2023.102080

49 Baumeister, W., Grimm, R. & Walz, J. Electron tomography of molecules and cells.
Trends Cell Biol **9**, 81-85 (1999). https://doi.org/10.1016/s0962-8924(98)01423-8

50 Liu, Y. T. *et al.* Isotropic reconstruction for electron tomography with deep learning.
Nature communications **13**, 6482 (2022). https://doi.org/10.1038/s41467-022-
33957-8

51 Turk, M. & Baumeister, W. The promise and the challenges of cryo-electron
tomography. *FEBS letters* **594**, 3243-3261 (2020). https://doi.org/10.1002/1873-
3468.13948

52 Dosemeci, A., Weinberg, R. J., Reese, T. S. & Tao-Cheng, J. H. The Postsynaptic
Density: There Is More than Meets the Eye. *Front Synaptic Neurosci* **8**, 23 (2016).
https://doi.org/10.3389/fnsyn.2016.00023

53 Vyas, Y. & Montgomery, J. M. The role of postsynaptic density proteins in neural
degeneration and regeneration. *Neural Regen Res* **11**, 906-907 (2016).
https://doi.org/10.4103/1673-5374.184481

54 Broadhead, M. J. *et al.* PSD95 nanoclusters are postsynaptic building blocks in
hippocampus circuits. *Scientific reports* **6**, 24626 (2016).
https://doi.org/10.1038/srep24626

55 Sheng, M. & Hoogenraad, C. C. The postsynaptic architecture of excitatory
synapses: a more quantitative view. *Annual review of biochemistry* **76**, 823-847
(2007). https://doi.org/10.1146/annurev.biochem.76.060805.160029

56 Kaizuka, T. & Takumi, T. Postsynaptic density proteins and their involvement in
neurodevelopmental disorders. *J Biochem* **163**, 447-455 (2018).
https://doi.org/10.1093/jb/mvy022

57 Zeng, M., Ye, F., Xu, J. & Zhang, M. PDZ Ligand Binding-Induced Conformational
Coupling of the PDZ-SH3-GK Tandems in PSD-95 Family MAGUKs. *Journal of*

805   *molecular biology* **430**, 69-86 (2018). https://doi.org/10.1016/j.jmb.2017.11.003

806 58 Hassani Nia, F. *et al.* Structural deficits in key domains of Shank2 lead to alterations
807   in postsynaptic nanoclusters and to a neurodevelopmental disorder in humans.
808   *Molecular psychiatry* (2022). https://doi.org/10.1038/s41380-022-01882-3

809 59 Chen, X. *et al.* PSD-95 family MAGUKs are essential for anchoring AMPA and
810   NMDA receptor complexes at the postsynaptic density. *Proceedings of the National*
811   *Academy of Sciences of the United States of America* **112**, E6983-6992 (2015).
812   https://doi.org/10.1073/pnas.1517045112

813 60 Dutta, A. *et al.* Cooperative Dynamics of Intact AMPA and NMDA Glutamate
814   Receptors: Similarities and Subfamily-Specific Differences. *Structure* **23**, 1692-
815   1704 (2015). https://doi.org/10.1016/j.str.2015.07.002

816 61 Krieger, J., Bahar, I. & Greger, I. H. Structure, Dynamics, and Allosteric Potential
817   of Ionotropic Glutamate Receptor N-Terminal Domains. *Biophysical journal* **109**,
818   1136-1148 (2015). https://doi.org/10.1016/j.bpj.2015.06.061

819 62 Garcia-Nafria, J., Herguedas, B., Watson, J. F. & Greger, I. H. The dynamic AMPA
820   receptor extracellular region: a platform for synaptic protein interactions. *J Physiol*
821   **594**, 5449-5458 (2016). https://doi.org/10.1113/JP271844

822 63 Durr, K. L. *et al.* Structure and dynamics of AMPA receptor GluA2 in resting, pre-
823   open, and desensitized states. *Cell* **158**, 778-792 (2014).
824   https://doi.org/10.1016/j.cell.2014.07.023

825 64 Meyerson, J. R. *et al.* Structural mechanism of glutamate receptor activation and
826   desensitization. *Nature* **514**, 328-334 (2014). https://doi.org/10.1038/nature13603

827 65 Nakagawa, T., Cheng, Y., Sheng, M. & Walz, T. Three-dimensional structure of an
828   AMPA receptor without associated stargazin/TARP proteins. *Biological chemistry*
829   **387**, 179-187 (2006). https://doi.org/10.1515/BC.2006.024

830 66 Traynelis, S. F. *et al.* Glutamate receptor ion channels: structure, regulation, and
831   function. *Pharmacol Rev* **62**, 405-496 (2010).
832   https://doi.org/10.1124/pr.109.002451

833 67 Sheng, M. & Kim, E. The postsynaptic organization of synapses. *Cold Spring Harb*
834   *Perspect Biol* **3** (2011). https://doi.org/10.1101/cshperspect.a005678

835 68 Husi, H., Ward, M. A., Choudhary, J. S., Blackstock, W. P. & Grant, S. G. Proteomic
836   analysis of NMDA receptor-adhesion protein signaling complexes. *Nat Neurosci* **3**,
837   661-669 (2000). https://doi.org/10.1038/76615

838 69 Vinade, L. *et al.* Affinity purification of PSD-95-containing postsynaptic
839   complexes. *J Neurochem* **87**, 1255-1261 (2003). https://doi.org/10.1046/j.1471-
840   4159.2003.02091.x

841 70 Dosemeci, A. *et al.* Composition of the synaptic PSD-95 complex. *Mol Cell*
842   *Proteomics* **6**, 1749-1760 (2007). https://doi.org/10.1074/mcp.M700040-MCP200

843 71 Husi, H. & Grant, S. G. Isolation of 2000-kDa complexes of N-methyl-D-aspartate
844   receptor and postsynaptic density 95 from mouse brain. *J Neurochem* **77**, 281-291
845   (2001). https://doi.org/10.1046/j.1471-4159.2001.t01-1-00248.x

846 72 Geinisman, Y. *et al.* Structural synaptic correlate of long-term potentiation:
847   formation of axospinous synapses with multiple, completely partitioned
848   transmission zones. *Hippocampus* **3**, 435-445 (1993).
849   https://doi.org/10.1002/hipo.450030405

850 73 Stewart, M. G. *et al.* Chemically induced long-term potentiation increases the

number of perforated and complex postsynaptic densities but does not alter dendritic spine volume in CA1 of adult mouse hippocampal slices. *Eur J Neurosci* **21**, 3368-3378 (2005). https://doi.org/10.1111/j.1460-9568.2005.04174.x

Nieto-Sampedro, M., Hoff, S. F. & Cotman, C. W. Perforated postsynaptic densities: probable intermediates in synapse turnover. *Proceedings of the National Academy of Sciences of the United States of America* **79**, 5718-5722 (1982). https://doi.org/10.1073/pnas.79.18.5718

Alten, B. *et al.* Role of Aberrant Spontaneous Neurotransmission in SNAP25-Associated Encephalopathies. *Neuron* **109**, 59-72 e55 (2021). https://doi.org/10.1016/j.neuron.2020.10.012

Sun, R. *et al.* An efficient protocol of cryo-correlative light and electron microscopy for the study of neuronal synapses. *Biophysics Reports* **5**, 111-122 (2019). https://doi.org/10.1007/s41048-019-0092-4

Hagen, W. J. H., Wan, W. & Briggs, J. A. G. Implementation of a cryo-electron tomography tilt-scheme optimized for high resolution subtomogram averaging. *Journal of structural biology* **197**, 191-198 (2017). https://doi.org/10.1016/j.jsb.2016.06.007

williamnwan/TOMOMAN: TOMOMAN 08042020 v. 08042020 (Zenodo, 2020).

Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017). https://doi.org/10.1038/nmeth.4193

Zhang, K. Gctf: Real-time CTF determination and correction. *Journal of structural biology* **193**, 1-12 (2016). https://doi.org/10.1016/j.jsb.2015.11.003

Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of three-dimensional image data using IMOD. *Journal of structural biology* **116**, 71-76 (1996). https://doi.org/10.1006/jsbi.1996.0013

Bharat, T. A. & Scheres, S. H. Resolving macromolecular structures from electron cryo-tomography data using subtomogram averaging in RELION. *Nat Protoc* **11**, 2054-2065 (2016). https://doi.org/10.1038/nprot.2016.124

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012). https://doi.org/10.1038/nmeth.2019

Fig. 1

a

b

c

Gray value

Cleft width

PSD height

PreSM
Putative adhesion
PostMP
PostSM
PSD nanoblock

Distance (nm)

d

Cleft width (nm)

Synaptosome   Synaptoneurosome

e   **f**   **g**

h   **i**   **j**   **k**

l

SV centroid

Nanoblock edge

Nanoblock centroid

d

D

m

Relative Frequency (%)

d/D

n

Cumulative Frequency (%)

Experiemntal
Random

d/D

Fig. 2

Fig. 3

Fig. 4

p 130 nm² 325 nm² 651 nm² 1161 nm²
Relative Frequency (%)
Nanoblock area (nm²)

q
Relative Frequency (%)
Nanoblock centroid distance (nm)

r
Relative Frequency (%)
Nanoblock edge distance (nm)

s
Relative Frequency (%)
d/D

t
Cumulative Frequency (%)
Experimental
Random
d/D

u
Dispersed nanoblocks
Clustered nanoblocks

Supplementary Fig. 1

Supplementary Fig. 2

Supplementary Fig. 3

a

Subtomogram average

1565 particles picked   3D auto refine without symmetry

3D classification with C2 symmetry

458 particles   219 particles

2 good classes

Remove duplicates and particles not on the membrane

391 particles   189 particles

Post processing

?

Type A particle   Type B particle   Undefined particles
391 particles   189 particles   888 particles

b

Type A particle   Final resolution = 24 Å

Fourier Shell Correlation vs Resolution (1/Å)

Corrected   Masked Maps
Unmasked Maps   Phase Randomized Masked Maps

Type B particle   Final resolution = 26 Å

Fourier Shell Correlation vs Resolution (1/Å)

Corrected   Masked Maps
Unmasked Maps   Phase Randomized Masked Maps

c   **d**   **e**

f   **g**   **h**

Supplementary Fig. 4