Serotonin modulates infraslow oscillation in the dentate gyrus during Non-REM sleep

Gergely F. Turi[1,2#*], Sasa Teng[3,4#], Xinyue Chen[4,5], Emily CY Lim[6], Carla Dias[1,2], Ruining Hu[3], Ruizhi Wang[4], Fenghua Zhen[3,7], Yueqing Peng[3,4,8*]

[1]New York State Psychiatric Institute, Division of Systems Neuroscience New York, NY 10032, USA

[2]Department of Psychiatry, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY 10032, USA

[3]Institute for Genomic Medicine, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY 10032, USA

[4]Department of Neurology, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY 10032, USA

[5]Department of Neuroscience, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY 10032, USA

[6]Columbia College, Columbia University, New York, NY 10027, USA

[7]Present address: National Institute of Allergy and Infectious Diseases, Bethesda, MD, 20894, USA

[8]Department of Pathology and Cell Biology, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY 10032, USA

[#]These authors contributed equally

[*]Correspondence to: gt2253@cumc.columbia.edu, yp2249@cumc.columbia.edu

Keywords:

Dentate gyrus, serotonin, infraslow oscillation, microarousal, sleep, calcium imaging, fiber photometry, two-photon imaging

ABSTRACT

Synchronous neuronal activity is organized into neuronal oscillations with various frequency and time domains across different brain areas and brain states. For example, hippocampal theta, gamma and sharp wave oscillations are critical for memory formation and communication between hippocampal subareas and the cortex. In this study, we investigated the neuronal activity of the dentate gyrus (DG) with electrophysiological and optical imaging tools during sleep-wake cycles. We found that the activity of major glutamatergic cell populations in the DG is organized into infraslow oscillations (0.01 – 0.03 Hz) during NREM sleep. Although the DG is considered a sparsely active network during wakefulness, we found that 50% of granule cells and about 25% of mossy cells exhibit increased activity during NREM sleep. Further experiments revealed that the infraslow oscillation in the DG is modulated by rhythmic serotonin release during sleep, which oscillates at the same frequency but in an opposite phase. Genetic manipulation of 5-HT receptors revealed that this neuromodulatory regulation is mediated by 5-HT1a receptors and the knockdown of these receptors leads to memory impairment. Together, our results provide novel mechanistic insights into how the 5-HT system can influence hippocampal activity patterns during sleep.

INTRODUCTION

Sleep is an evolutionarily conserved biological process observed in the animal kingdom. Invertebrates such as *Drosophila*, *C. elegans*, and even *Hydra* show sleep-like behavior (1-5). In mammals, sleep stages can be subdivided to rapid eye movement (REM) sleep and non-REM (NREM) phases. During NREM sleep the activity of the skeletal muscle is reduced and the EEG is dominated by slow, high amplitude synchronous oscillations while in REM sleep the EEG activity is more similar to the awake brain state. There are also smaller building blocks of sleep stages. For instance, NREM sleep is often interrupted by brief epochs, which based on their EEG and EMG signatures, show remarkable reminiscence to awake states, thus, they are often referred to as microarousals (MAs). These events can be captured as brief motor-bursts on the EMG signal and by abrupt alternations of low frequency oscillatory patterns in EEG recordings (6-8). Even though MAs are natural parts of the sleep architecture, the function of these events is largely unknown.

Sleep is a fundamental biological process that plays a crucial role in many physiological functions. Studies using various model organisms have demonstrated that severe sleep loss or total

sleep deprivation can even have fatal effect (9-11). While the exact biological function of sleep is not fully understood, research has suggested that it plays a crucial role in memory consolidation, which is the process of converting newly acquired memories into a permanent form. The supportive evidence for this hypothesis is quite rich although far from consolidated. Current models suggest that slow wave sleep, which makes up deeper stages of NREM sleep, processes declarative memory and it requires the active participation of the hippocampus through the replay of episodic memory traces captured during waking hours (12, 13). While the role of hippocampal CA1 and CA3 subareas in this process is extensively studied, less is known about the contribution of the dentate gyrus (DG) which is the first station of the classic trisynaptic loop and located upstream from CA3. The main glutamatergic cell types of the DG are the granule cells (GCs) located in the granule cell layer and the mossy cells (MCs) which populate the hilus or polymorph layer between the upper and lower blades of the GC layers (14). Experimental work has shown that GCs and MCs form a functional unit to perform pattern separation during active exploration(15-17), a neuronal mechanism by which distinct memory traces can be created even if the input pattern is highly overlapping. Furthermore, it has been also shown that in awake immobile mice the DG generates sparse, synchronized activity patterns driven by inputs from the entorhinal cortex(18).

Sleep/wake cycles are regulated by various monoaminergic and peptidergic cell groups located in the hypothalamus and brainstem. Among the monoaminergic neuromodulators, the serotonin (5-HT) system has been shown to promote wakefulness and suppress REM sleep (19, 20). It is widely accepted in the field that 5-HT release is reduced during NREM sleep and 5-HT cells become entirely silent during REM sleep (20, 21). Serotonin is also recognized as one of the primary neuromodulatory inputs to the DG and a substantial body of evidence implicates that genetic manipulation of 5-HT or 5-HT-related genes has a profound impact on a broad range of hippocampal processes, including anxiety behavior, learning and memory (22-27). Serotonin receptors are expressed throughout the dorsoventral axis of the hippocampus (28), with GCs expressing both the inhibitory 5-HT1a and excitatory 5-HT4 receptor subtypes and about quarter of MCs expressing the excitatory 5-HT2a receptors (28). Together these findings suggest that the 5-HTergic system have a strong impact on hippocampal functions.

In this study, we investigated the activity of GCs and MCs during sleep-wake cycles and their modulation by the 5-HT system. We found that both glutamatergic cell types – especially

GCs – exhibit higher activity during specific sleep stages compared to wakefulness, and the calcium activity is organized to infraslow oscillatory (ISO) cycles during NREM sleep epochs. Our further findings obtained by recording from major 5-HTergic cell populations in the Raphe and by manipulating specific 5-HT receptor subtypes suggest that the slow oscillatory activity in the DG is regulated by an ISO of the 5-HT system oscillating in the opposite phase during NREM which is also tightly coupled to MA events. Genetic knockdown of 5-HT1a receptors in the DG impairs the ISOs and contextual fear memory, suggesting a key role for this inhibitory 5-HT receptor in governing the oscillatory effect.

RESULTS

Infraslow neural oscillation of DG populations during NREM sleep

To investigate population activity during sleep and awake brain states in the DG in a cell type-specific manner, we injected a set of Dock10-Cre$^{+/-}$ mice with AAV1-FLEX-GcaMP6s to drive the expression of GCaMP specifically in GCs (29). The mice were then implanted with fiber photometry probes, EEG and EMG electrodes to facilitate brain state classification. Two weeks after recovery, we conducted chronic photometry and EEG recordings while the animals experienced natural wake/sleep cycles in a behavioral chamber. Consistent with previous studies (15, 18, 30), we observed significantly higher populational calcium activity during sleep states (NREM and REM), compared to wakefulness (**Fig 1A-1C**). Strikingly, our data also revealed a structured pattern in the calcium signal: GC activity was organized to an infraslow oscillation (ISO, 1-2 cycles/ minute, or 0.017-0.033 Hz) during NREM sleep (**Fig 1A, 1D**). Cross correlation between calcium activity and EMG/EEG revealed that the troughs of the ISO partially coincided with MA episodes (**Fig. 1E**). Quantitative analysis showed that 30% of the calcium troughs during NREM sleep were followed by a MA epoch while 59% of troughs were accompanied by the maintenance of NREM sleep (**Fig 1F**). Notably, the reduction in GC activity significantly preceded the occurrence of MA events as judged from the EMG, where the average latency between the calcium trough and the onset of the MA was 5.89 +/- 0.28 s (Fig **1G-1H**).

MCs are among the first synaptic partners of GCs, therefore we set out to record the calcium activity of MCs by injecting a new set of mice with dopamine 2 receptor-Cre (DrD2-Cre$^{+/-}$) genetic background (**Fig 2A-B**). DrD2 expression in the hilus is highly specific to MCs (31, 32).

MCs displayed similar levels of calcium activity during wake and NREM sleep, and significantly increased activity during REM sleep (**Fig 2C**). Similar to GCs, we observed the ISO in MCs calcium signal, although with slightly lower peak frequency (**Fig 2A, 2D**). Furthermore, correlation analysis between photometric and EEG/EMG signals showed similar results as with GCs, i.e. around 30% of calcium troughs in MCs lead to MAs and 60% were resulted in the maintenance of NREM (**Fig 2E-2G**). The average latency between the reduction of calcium signal and the onset of MAs was somewhat shorter than in GCs (4.89 +/- 0.40 s) (**Fig 1H**).

Overall, our data showed similar NREM-specific oscillatory activity in both GCs and MCs. The increased activity of GCs and MCs during sleep stages is consistent with previous studies (15, 18, 30, 33). However, the ISO in the calcium activity of glutamatergic neurons, to the best of our knowledge, has not been reported yet.

Ensemble activity of dentate gyrus in sleep

To study the cellular mechanisms underlying sleep-specific activity, we conducted head-fixed two-photon imaging experiments combined with EEG/EMG recordings to record neuronal ensemble activity in the DG during sleep stages (**Fig 3A**). Wild type mice were injected with AAVdj-syn-jGCaMP7b in the DG followed by optical window, EEG and EMG implantations. After a few weeks of recovery periods, we recorded the cellular activity in the granule cell layer and in the hilus (**Fig 3A-C**) while the mouse was engaged in locomotory and motionless epochs on a treadmill. To facilitate sleep under the two-photon microscope, the mice underwent mild sleep deprivation before the recordings (see Methods for details). *Post hoc* analysis of EEG/EMG signals confirmed the presence of sleep states (**Fig. 3B-C,** also **Suppl Fig 1A-B**). Comparing the calcium signals recorded during awake and sleep intervals revealed that the calcium activity was significantly upregulated during NREM in about 50% of the GC and 28% of the putative MC populations (**Fig 3D, 3E**) in the hilus. Similar to photometry experiments, we then performed MA analysis and its correlation with calcium activity. We found that up-regulated, but not down-regulated putative GCs and putative MCs displayed decreased activity during MAs in NREM sleep (**Fig 3D, 3E, Suppl Fig 1C-E**).

ISO is driven by serotonin release during NREM sleep

The prominent electrographic patterns in the DG during NREM sleep include dentate spikes and sharp-waves (34-36), but these events are generally too short-lived (30-120 ms) to account for the long-lasting active periods during ISO. Given the second-long latencies between calcium troughs and MAs, we turned our attention toward neuromodulatory systems that are thought to modulate sleep/wake periods on a slower scale and display phasic activity during sleep cycles (37). Given the recent identification by Kato et al. of a slow oscillatory activity at the population level of the dorsal Raphe 5-HT neurons, which shares a similar infraslow oscillatory frequency range (0.01-0.3Hz) with what we observed in the DG (38), we looked at the possibility whether the 5-HT system might be responsible for the modulation of the ISO in the DG. In a subsequent set of experiments, we used GRAB$_{5-HT2h}$, a recently developed and optimized genetically encoded 5-HT sensor (39) to measure 5-HT dynamics during sleep. Considering that 5-HT is exclusively synthetized by the Raphe nuclei in the adult brain (40), our initial step involved stereotactic injection of an AAV expressing the GRAB$_{5-HT2h}$ sensor into this region, followed by the implantation of an optic fiber above the injection site. Our results recapitulated the canonical view on 5-HT dynamics during sleep/wake cycles: we observed the highest 5-HT level during wakefulness, significantly lower level of 5-HT during NREM and detected the lowest 5-HT levels during REM periods (**Suppl Fig. 2A, 2E**) (19-21). We also observed a phasic pattern of 5-HT during NREM sleep (**Suppl Fig. 2A, 2F**) which is consistent with recently published studies (38, 39). Our further analyses revealed that the rhythmic 5-HT release during NREM sleep was also correlated with MA episodes (**Suppl Fig. 2B**). Correlation analysis revealed that 35% of phasic 5-HT peaks during NREM sleep were accompanied by MA events detected by the EMG bursts (**Suppl Fig. 2C**). The risetime of the phasic increase in 5-HT during NREM preceded the onset of the MA by 6.82 +/- 0.44 s (**Suppl Fig. 2D**).

To confirm that this infraslow oscillatory pattern in 5-HT is manifested in the hippocampus as well, we injected AAV-GRAB$_{5-HT2h}$ into the DG and implanted an optic fiber above the injection site. (**Fig 4A**). Similar to 5-HT signals recorded from the Raphe nuclei, we observed the highest level of 5-HT in wake, the lowest in REM sleep, and intermediate level in NREM sleep (**Fig 4B-D**). Importantly, the 5-HT dynamics during NREM sleep displayed the same oscillatory pattern in the DG that we observed in the Raphe (**Fig 4B**). Thirty-four percent of the phasic increases in 5-HT were associated with MAs (**Fig 4C**). Furthermore, the frequency of phasic 5-HT release during

NREM was 1.14 +/- 0.04 cycle/min (**Fig 4E**), closely matching the frequency of the ISO in the DG.

The tight alignments of both the calcium troughs in the DG and the phasic 5-HT peaks in the Raphe with MA events suggest a causal relationship between them. To test this hypothesis, we conducted dual site fiber photometry recordings in SERT-Cre[+/-] mice by injecting AAV1-FLEX-GCaMP6s in the Raphe and with AAV9-CaMKII-GCaMP6s in the DG (**Fig 5A**) followed by fiber implants to these areas. *Post hoc* histological analysis of AAV9-CaMKII-GCaMP6s injected brains confirmed that the expression of GCaMP is largely restricted to GCs (**Suppl Fig. 3**). Furthermore, analysis of the calcium signals from the AAV9-CaMKII-GCaMP6s-labeled population revealed a similar ISO pattern as we observed with the genetically more selective approach (**Suppl Fig. 3**). Strikingly, the phases of the photometry signals in the DG and Raphe appeared to be anticorrelated during NREM sleep, i.e. the calcium peaks in the Raphe appeared when the intensity of the calcium signal dropped in the DG (**Fig. 5B**). Indeed, Pearson's correlation analysis of the two signals resulted in a strong negative correlation between the Raphe and DG during NREM sleep (**Fig. 5C, 5D**). Interestingly, during wake periods, the DG activity was positively correlated with the Raphe activity (**Fig. 5D**). Together, these results suggest that the down-phase of the population activity during the ISO is driven by the phasic release of 5-HT during NREM sleep.

Granule cell activity is inhibited via 5-HT1a receptors during the ISO

Our results so far suggested that the periodic increase in 5-HT levels during NREM sleep inhibits GC and MC activity. To identify the molecular basis of this relationship, we conducted local genetic manipulations of the 5-HT1a receptors. We selected this subtype because of its high expression pattern in GCs (28) and its well-characterized inhibitory effect on this cell type(41). We injected a cohort of 5-HT1a[flox/flox] mice (43) with a mix of AAV1-hSyn-Cre and AAV9-CaMKII-GCaMP6s to simultaneously express a calcium sensor and to knock out 5-HT1a receptors specifically in the GCs (**Fig 6A**). A control group of 5-HT1a[flox/flox] mice was injected with AAV9-CaMKII-GcaMP6s. We then conducted fiber photometry recordings in the DG. Analysis of the calcium signals during awake and sleep period resulted in significantly decreased ISO during NREM in the 5-HT1a knock out group compared to the control (**Fig. 6B-D**) Post hoc *in situ* hybridization confirmed the lack of 5-HT1a receptors at the injection site (**Suppl Fig. 4**).

Together, our two-site recordings and genetic manipulation data indicated that the ISO in the DG is modulated through an inhibitory mechanism via 5-HT1a receptors.

5-HTergic modulation of the DG via 5-HT1a receptors is required for contextual memory

Neural oscillations during NREM sleep are thought to be involved in memory consolidation (42). We hypothesized that the 5-HT1a-dependent ISO is a required component for hippocampus-dependent memory consolidation. To test this hypothesis, we genetically knocked out 5-HT1a receptors in the DG and examined its effect on memory performance. We injected 5-HT1a$^{flox/flox}$ mice with AAV9-CaMKII-Cre-GFP (intervention group) or with AAV9-CaMKII-GFP (control group) bilaterally in the dorsal DG. Two weeks after surgery, we tested the memory performance of the mice by examining freezing behavior in a contextual fear conditioning (CFC) paradigm **(Fig. 6E)**. In accordance with our hypothesis, we observed diminished memory performance in retrieval tests in the intervention group (**Fig. 6F**) measured by the time spent with freezing (30.0%±4.63% *vs.* 49.2%±7.10%).

DISCUSSION

In this study, we demonstrated that major glutamatergic cell types of the DG, the GCs and MCs, display ISO activity (1-2 cycles/min or 0.02-0.03 Hz) during NREM sleep (**Fig. 1-2**) but not during awake or REM periods. Our 2-photon calcium imaging data shows that this oscillation is mainly driven by NREM-active glutamatergic cells (about 50% of GCs and 28% of MCs, **Fig. 3**). Furthermore, we found that the ISO is negatively correlated with phasic 5-HT release in Raphe nuclei during NREM sleep (**Fig. 4-5**). Finally, we showed that local genetic ablation of 5-HT1a receptors in GCs impaired the ISO and memory consolidation (**Fig. 6**). Taken together, our findings demonstrate that the serotonin system plays a pivotal role in modulating DG activity during NREM sleep.

MCs and GCs are two major cell populations of DG which are thought to be involved in pattern separation and other cognitive functions in awake animals (15-17, 53). The activity of GCs and MCs has been investigated with both electrophysiological and imaging tools and the available data suggests that both cell types are significantly more active during sleep (15, 18, 33) than wakefulness. To our knowledge however, ours is the first report which shows that the population activity

is dynamically rendered to ISO pattern during NREM sleep which is organized by at least one of the neuromodulatory systems, the 5-HT.

There is much less available data on the activity pattern of MCs and GCs during MA episodes. Here we showed that about 50% of putative GCs and 28% of putative MCs are upregulated during NREM and their activity is abruptly terminated by MAs through a 5-HT mediated mechanism.

We also observed that local downregulation of 5-HT1a receptors disrupts the ISO, demonstrating that the 5-HT is directly acting in the DG to gate the oscillations. 5-HT1a receptors are expressed by GCs which suggest that the inhibitory effect is mediated via this cell type. Since MCs are driven by GCs via highly efficient "detonator" type of synapses, our current working hypothesis is that MC excitation is driven by GCs during NREM. We speculate that the drop of the calcium activity in MCs during the ISO is not mediated by serotonin but rather by the lack of the excitation from GCs during the phasic 5-HT release. This hypothesis is supported by the difference in the delay preceding MAs in GCs and MCs (5.89 +/- 0.28 s in GCs *vs.* 4.89 +/- 0.40 s in MCs) which shows that GC inactivity is followed by MC inactivity.

ISOs have been observed in multiple brain regions during sleep wake cycles (44). Polysomnographic recordings have revealed a 0.02 Hz ISO pattern recorded from cortical areas (45) which frequency is on the same time scale as the ISO we recorded from the DG. The cortical ISO is dominant in the sigma (10-15 Hz) power range in both mice and humans(45) which is the most prominent frequency band that contains neural rhythms associated with the gating of sensory information during sleep. The authors did not correlate the ISO with MAs but they designed an experiment where white noise stimulus woke up the mice 38% of the time. This stimulation was the most effective when the ISO was in its ascending phase. Our data with others (38) show that 5-HT concentration slowly oscillates in the hippocampus and in the Raphe during NREM, thus it would be intriguing to conclude that peaks of the phasic 5-HT bursts act as the gating signal when the animal is more receptive to external stimulation and this can lead to awakening. However, we also need to point out that there are multiple other systems that can be active in parallel and contribute to the shaping of the sleep microarchitecture (46). Two recent publications from different laboratories have shown rhythmic release of norepinephrine (NE) (~0.03 Hz) in the medial prefrontal cortex and in the locus coeruleus of the mouse during sleep-wake cycles (46, 47). The recording techniques used in these studies are highly similar to ours and the analyses arrive at the

same conclusion, that is, about 30% of the phasic neuromodulator bursts lead to MAs while NREM sleep is maintained about 60% of the time. Qualitative assessment of the available data obtained with other novel neuromodulatory sensors suggests, that many other major neuromodulatory systems (i.e. histaminergic, cholinergic) (48, 49) also display this phasic activity during NREM sleep which raises the question of how the individual components of the ascending reticular activating system shapes sleep microarchitecture. Increase in the infraslow oscillatory frequency of NE during NREM leads to increased frequency of MAs (47, 50). However, increasing the extracellular 5-HT level by acute SSRI administration does not seem to change the frequency of these events (38). This suggests that the 5-HT system might have different neuromodulatory functions from the NE system during sleep.

Our data demonstrated that phasic release of 5-HT during sleep acts as an inhibitory input to the neuronal activity in DG via 5-HT1a receptors, thus it serves as a pacemaker signal. In addition, we speculate that there could be an excitatory input which is responsible for the elevated activity of GC and MC subpopulations during NREM. The DG is the first station of the hippocampal tri-synaptic loop receiving glutamatergic inputs from distant brain areas such as the entorhinal cortex or the supramammillary nucleus(51). Recent data have shown a similar ISO in the medial entorhinal cortex in headfixed mice while moving on a rotating wheel (52)which also transcended immobility epochs, but further data would be needed to confirm that they exist during NREM sleep as well.

A recent work from Pofahl et al. have used *in vivo* two-photon microscopy and showed that GCs display synchronous activity patterns during immobility(18) which might be related to the phenomenon we observed. However, the authors were not able to break down the immobility intervals to sleep stages due to the lack of simultaneous EEG/EMG recordings, nor did they assess the activity of the MCs in their work. Our results demonstrated dynamic activity of DG cells during sleep (**Fig. 3**). Interestingly, we did not observe the ISO at the single-cell level under two-photon microscope. One possibility is that this was due to different sleep patterns between head-restricted and freely moving animals. We hypothesize that deep sleep is needed for the ISO. Even though we habituated our mice extensively to our two-photon setup to minimize stress, headfixed posture may still result in differences in sleep behavior due to the slightly increased stress-response caused by headfixation (54). Another possibility is that the ISO mainly has dendritic origin and manifest less at the somatic level. Future dendritic imaging in sleeping mice can test this possibility.

In summary, here we showed that calcium activity in the DG is highly increased during NREM sleep periods and population activity is entrained to ISO. The activity of the serotonergic system is highly correlated with this oscillation and rhythmic bursts of 5-HT are responsible for the maintenance of the oscillatory activity, via 5-HT1a receptors expressed by GCs. Together, our study highlights the role of neuromodulation in organizing neuronal activity during sleep and sleep-dependent brain functions, such as memory.

ACKNOWLEDGEMENTS
We thank René Hen at Columbia University for providing *5-HT1a^{flox/flox}* mice. We thank Kitti Rusznak and Hayley Judice for helping with behavioral experiments. The 5-HT sensor (AAV9-hSyn-5-HT2h) was a gift from the Yulong Li's laboratory at Peking University. This work was supported by startup funds from Columbia University to Y.P., and by Columbia University Precision Medicine Initiative to Y.P. G.F.T was supported by a BBRF Young Investigator Award, a NIMH R21MH122965 and NIA R21AG079025. The AAVdj-syn-jGCaMP7b virus was packaged by the Gene Vector and Virus Core at Wu Tsai Neurosciences Institute at Stanford University.

AUTHOR CONTRIBUTIONS
G.T., S.T. and Y.P. designed the study, carried out the experiments, and analyzed data. X.C, E.L. and R.H. performed behavioral experiments. R.W. performed in-situ hybridization, C.D. and F.Z. performed histology and data analysis. Y.P. and G.T. wrote the paper.

DECLARATION OF INTERESTS
Authors declare that they have no competing interests.

FIGURE LEGENDS

Fig. 1 Infraslow calcium oscillation of granule cells during NREM sleep.
A. Representative recording session showing infraslow oscillation (ISO) during NREM sleep. From top to bottom: brain states, EEG power spectrogram (0-25 Hz), EMG amplitude, photometric

signal. **B.** Left, Schematic representation of the recording setup. Right, a fluorescence image showing the expression of GCaMP6s (green) in the granule cell layer and the optic fiber placement (dashed line) in a Dock10-Cre mouse. Scale bar, 500 μm. **C.** Quantification of calcium activity in granule cells (13 recording sessions in 5 Dock10-Cre mice, n.s. – no significance, ** - P<0.01, paired t-test) in wake (W), NREM sleep (N), and REM sleep (R). **D.** Left: oscillation peak frequency during NREM sleep and wake based on Fourier transformation of the photometry signal. Right: Quantification of calcium oscillations in GCs (13 sessions in 5 mice). **E.** A representative example showing the coincidence of calcium troughs with MAs. **F.** Percentage of outcome of brain states following calcium drops during NREM sleep. **G.** Peri-stimulus time histogram (PSTH) in one recording session showing calcium signal aligned with the onset of MAs. Bottom left: parameters of the calcium signal used for quantification. **H.** Quantification of the latency (t) and magnitude of the calcium trough (Drop) during MAs (13 sessions from 5 Dock10-Cre mice).

Fig. 2 Infraslow calcium oscillation of mossy cells during NREM sleep.

A. Representative recording session showing infraslow calcium oscillation during NREM sleep in a Drd2-Cre$^{+/-}$ mouse injected with AAV-FLEX-GCaMP6s. From top to bottom: brain states, EEG power spectrogram (0-25 Hz), EMG amplitude, photometric signal. **B.** Fluorescence images showing the expression of GCaMP6s (green) in the mossy cells in the area marked with the white rectangle. Blue, DAPI. Scale bars, 500 μm and 50 μm. **C.** Quantification of calcium activity in mossy cells during different brain states (12 recording sessions in 4 Drd2-Cre$^{+/-}$ mice; n.s. – no significance, ** - P<0.01, paired t-test). **D.** Left: oscillation peak frequency during NREM sleep based on Fourier transformation of the photometry signal. Right: quantification of calcium oscillations during NREM sleep. **E.** A representative example showing the coincidence of calcium troughs with MAs (represented with vertical green lines). **F.** Percentage of outcome of brain states following calcium troughs during NREM sleep. **G.** PSTH from one recording session showing calcium signals aligned with the onset of MAs. **H.** Quantification of the latency (t) and the magnitude of calcium troughs (Drop) during the MAs (12 sessions from 4 Drd2-Cre mice).

Fig. 3 Two-photon calcium imaging of DG activity in sleeping mice.

A. Left, schematic layout of two-photon imaging and EEG recording setup. Right, representative images showing putative granule cells (pGCs) and putative mossy cells (pMCs). Scale bars, 50

μm. **B-C** Representative two-photon recording sessions of field of views containing putative pGCs pMCs. From top to bottom: brain states (gray – awake, orange – NREM), EEG spectrogram (0-25 Hz), EMG, velocity (vel.), and calcium traces in individual cells. Red traces - upregulated cells, blue traces –downregulated cells, gray traces – non-significant cells. **D-E.** Left, Percentage of Up-, downregulated and non-significant cells from the entire recorded cell populations. Middle, quantification of calcium activity in different brain states in up-, down-regulated cells (pGCs: $P<0.05$ in up-regulated cells between NREM and MA, $P=0.77$ in down-regulated cells between NREM and MA; pMCs: $P<0.001$ in up-regulated cells between NREM and MA, $P=0.63$ in down-regulated cells between NREM and MA, paired t-test). Right, Averaged activity in up-, down-regulated pGCs and pMCs during microarousals (MA). Time 0 indicates the MA onset. Putative GCs: 369 cells from 3 C57BL/6J mice; putative MCs: 269 cells from 2 C57BL/6J mice.

Fig. 4 Phasic release of 5-HT in the DG during NREM sleep.
A. Left, schematic of experimental design. Right, expression of 5-HT sensor in the hippocampus.
B. A representative example of 5-HT signals during different brain states. From top to bottom: brain states, EEG power spectrogram (0-25 Hz), EMG signal, photometric signal. The dashed-box enlarged below in panel B. Note the coincidence of 5-HT release with MAs during NREM sleep (black arrows and vertical green lines). **C.** Outcome of brain states following 5-HT release during NREM sleep (averaged data from 6 mice). **D.** Quantification of 5-HT signals in the DG during different brain states (14 sessions from 6 C57BL/6J mice, ** - $P<0.01$, *** - $P<0.001$, paired t-test). Data were normalized to Z scores in each recording session. **E.** Quantification of oscillatory cycles of 5-HT signals in the DG (14 recording sessions from 6 C57BL/6J mice).

Fig. 5 Correlation between DG oscillation and activity of 5-HT neurons during NREM sleep.
A. Left, Schematic representation of the 2-site photometry experimental design. Right, Expression of CaMKII-GCaMP6s and fiber placement in the DG and Raphe nuclei. **B.** A representative example of concurrent recording of DG and Raphe 5-HT neurons in a Sert-Cre[+/-] mouse during sleep. From top to bottom: brain states, EEG power spectrogram (0-25 Hz), EMG amplitude, photometric calcium signals (CaMKII-G6s) in DG and in dorsal Raphe. **C.** Correlation analysis of calcium activity between DG and Raphe 5-HT neurons during NREM sleep and wakefulness in one re-

cording session. **D.** Quantification of correlation coefficient between DG activity and Raphe activity during different brain states (11 sessions from 3 Sert-Cre$^{+/-}$ mice, *** - P<0.001, paired t-test).

Fig. 6 Genetic knockdown of 5-HT1a receptors in DG impairs ISO and memory performance. A. Schematic representation of the experimental design. A mix of AAV9-CamKII-GCaMP6s and AAV1-hSyn-Cre was injected into the DG of 5-HT1a$^{flox/flox}$ mice. **B.** Representative example showing photometry and EEG recordings in the DG of a control mouse injected with AAV9-CaMKII-GCaMP6s alone. Right, Fourier transformation of calcium activity during wake (blue) NREM sleep (red). **C.** A representative example showing photometry and EEG recordings in the DG of a mouse injected with AAV9-CaMKII-GCaMP6s and AAV1-hSyn-Cre. Right, Fourier transformation of calcium activity during wake (blue) NREM sleep (red). **D.** Left, Quantification of the relative power of the calcium oscillation in the range of 1-2 cycles/min in the Cre and control groups (16 sessions from 5 mice for Cre, 16 sessions from 6 mice for control). Right, Quantification of calcium oscillation amplitudes in the Cre and control groups. Calcium signals in each mouse were normalized to Z scores. ** - P<0.01, *** - P<0.001, unpaired t-test. **E,** Schematic representation of the CFC experimental design. **F,** Left, Contextual fear recall tests showing percentage of freezing in one minute time bins for 5-HT1a$^{flox+/+}$ mice bilaterally injected with AAV9-CaMKII-Cre-GFP (Cre) or AAV9-CaMKII-GFP (GFP). Right, Quantification of freezing behavior over 5-minute interval during contextual recall tests in Cre and GFP groups (N=11 for Cre, N=12 for GFP, *, P<0.05, ***, P<0.001, un-paired t-test).

METHODS

Animals

All procedures were carried out in accordance with the US National Institute of Health (NIH) guidelines for the care and use of laboratory animals and approved by the Animal Care and Use Committees of Columbia University and New York State Psychiatric Institute. Adult (10-16 weeks of age) were used for all experiments from both sexes. The following mouse lines were used in the current study: C57BL/6J (JAX 000664), Dock10-Cre, Drd2-Cre, Sert-Cre, 5-HT1a$^{flox/flox}$. Mice were housed in 12-hour light-dark cycles (lights on at 07:00 am and off at 07:00 pm). Dock10-Cre, Drd2-Cre, Sert-Cre were bred with C57BL/6J mice, and the heterozygote offspring was used

in the experiments. Homozygote offspring of the 5-HT1a^{flox} line was used in local genetic manipulations.

Viral constructs

Name	Cat#	Source
AAV1-Syn-FLEX-GCaMP6s	100845-AAV1	Addgene
AAV9-CamKIIa-GCaMP6s	107790-AAV9	Addgene
AAVdj-syn-jGCaMP7b	NA	Custom made
AAV9-hSyn-5-HT2h	YL10097-AV9	Vigene Biosciences
AAV1-hSyn-Cre	105553-AAV1	Addgene
AAV9-CaMKII-Cre-GFP	105551-AAV9	Addgene
AAV9-CaMKII-GFP	105541-AAV9	Addgene

Surgical procedures

EEG and Fiber implants

Mice were anaesthetized with a mixture of ketamine and Xylazine (100 mg*kg-1 and 10 mg*kg-1, intraperitoneally), then placed on a stereotaxic frame with a closed-loop heating system to maintain body temperature. After asepsis, the skin was incised to expose the skull and a small craniotomy (~0.5 mm in diameter) was made on the skull above the regions of interest. A solution containing 50-200 nl viral construct was loaded into a pulled glass capillary and injected into the target region using a Nanoinjector (WPI). Optical fibers (0.2 mm diameter, 0.39 NA, Thorlabs) were implanted into the target region with the tip 0.1 mm above the virus injection site for fiber photometry recording. For EEG and EMG recordings, a reference screw was inserted into the skull on top of the cerebellum. EEG recordings were made from two screws on top of the cortex 1 mm from midline, 1.5 mm anterior to the bregma and 1.5 mm posterior to the bregma, respectively. Two EMG electrodes were bilaterally inserted into the neck musculature. EEG screws and EMG electrodes were connected to a PCB board which was soldered with a 5-position pin connector. All the implants were secured onto the skull with dental cement (Lang Dental Manufacturing). After surgery, the animals were returned to their home cages for recovery for at least two weeks before any experiment.

Virus injection

For fiber photometry, 150-200 nl AAV1-FLEx-GCaMP6s or AAV9-hSyn-5-HT2h was unilaterally injected in the DG (AP -1.9mm, ML 1.5mm, DV 1.7mm) or the dorsal Raphe (AP -4.5mm, ML 0mm, DV 3.2mm) respectively. An optical fiber was implanted 0.1mm above the injection site. The DV Coordinates listed above are relative to the pial surface.

For two-photon imaging, the 200 nl AAVdj-syn-jGCaMP7b was stereotactially injected unilaterally to the left DG (AP -1.9mm, ML 1.5mm, DV -1.7mm).

Optical cannula, EEG and EMG implant for two-photon imaging

For two-photon imaging we modified our standard surgical procedures (55) by implanting a pair of bone screws for unilateral EEG recording on the contralateral side to the cannula. Briefly, a week after the virus injection the mice were deeply anesthetized with Isoflurane, then a sagittal incision was made on the top of the skull. After removing the skin, trephination was done on the left side of the skull, by using a sterile 2mm diameter tissue punch. The overlying cortical and hippocampal tissue was removed from above of the DG, while the brain was constantly irrigated with sterile cortex buffer (125 mM NaCl, 5mM KCl,10mM glucose, 10mM HEPES, 2mM CaCl2, 2mM MgSO4. pH set to 7.4, osmolality to 305 mOsm). A sterile metal cannula (diameter: 2mm, height: 1.8 mm) with a glass coverslip glued (Norland) to the bottom was implanted above the DG and secured in place with layers of tissue adhesive (3M Vetbond) and dental acrylic. After the cannula was safely in place, we predrilled 2 holes on the right side of the skull (1.5 mm posterior of the Bregma) for implanting recording electrodes, and one hole above the cerebellum for a reference electrode. Silver wires (125μm diameter, PFA-coated) used for EEG recordings were placed into the predrilled holes and secured in place with stainless-steal skull screws (0.8 mm diam, 2.15 mm long). A pair of silver wires (125μm diameter, PFA-coated) were implanted to the neck muscle as described in the fiber photometry section to facilitate EMG recordings. Finally, a metal head bar was secured to the skull with dental acrylic and the custom PCB interface with the connected silver wires and a connector was secured to the head bar with dental acrylic as well. All exposed bone areas were covered with dental acrylic and the mice were returned to their home cage for postoperative care and recovery.

EEG recording

Mouse sleep behavior was monitored using EEG and EMG recording along with an infrared video camera at 30 frames per second. Recordings were performed for 24 hours (light on at 7:00 am and off at 7:00 pm) in a behavioral chamber inside a sound attenuating cubicle (Med Associated Inc.). Animals were habituated in the chamber for at least 4 hours before recording. EEG and EMG signals were recorded, bandpass filtered at 0.5-500 Hz, and digitized at 1017 Hz with 32-channel amplifiers (TDT, PZ5 and RZ5D or Neuralynx Digital Lynx 4S). For sleep analysis, spectral analysis was carried out using fast Fourier transform (FFT) over a 5 s sliding window, sequentially shifted by 2 s increments (bins). Brain states were semi-automatically classified into wake, NREM sleep, and REM sleep states using a custom-written MATLAB program using the following scoring criteria: wake: desynchronized EEG and high EMG activity; NREM: synchronized EEG with high-amplitude, delta frequency (0.5–4 Hz) activity and low EMG activity; REM: high power at theta frequencies (6–9 Hz) and low EMG activity. Semi-auto classification was validated manually by trained experimenters. We categorized wake bouts of <15 seconds as microarousals. To examine the DG activity during microarousals, the calcium signals were aligned to the onset of each microarousal event to generate the per-stimulus time histogram (PSTH). The time difference between the minimum of the averaged calcium signals (trough) to the MA onset was named as latency in Figure 1 and 2.

Two-photon recording

Mice were extensively habituated to the two-photon rig and to the head-fixed position before the recording sessions. They were placed in the head holder apparatus for gradually increasing amount of time (from 5 min to 1h) over the course of two weeks. Mice were head-restrained under the objective of the two-photon microscope, but otherwise able to walk on a custom-built treadmill. The mouse's behavior on the treadmill was collected with an earlier version of BehaviorMate(56). The locomotory and stationary epochs were detected with a rotary encoder attached to the axel of the treadmill wheel. The signal from the encoder was processed with BehaviorMate and saved on a PC.

A commercial *in vivo* multiphoton imaging system was used for the two-photon recordings (Bruker Ultima IV), equipped with a dual scan head. Nikon 10x (0.4 NA) or 16x (0.8 NA) objectives were used to record cellular activity. The emitted light was collected with photomultiplier

tubes (Hamamatsu, H11706-40 GaAsP). Excitation was achieved with an InSight X3 (Spectra-Physics) laser tuned to 940 nm. The light intensity was regulated with pockels cells (Model 350-80-LA-02 KD*P Series E-O Modulator, Conoptics). The collection optics consisted of a standard filter set (565 dichroic, 525/70 and 595/50 band pass) before the detectors.

Two-photon imaging sessions took 45-60 minutes. Images were collected at a 10 Hz frequency sampling rate in resonant scanning mode. Image dimensions were set to 512x512 pixel (width and height). The collected binary data was directly converted to Hierarchical Data Format (HDF) then it was deposited to our server. Image registration and cell detection was performed using Suite2p software (ver. 0.7.0) package(57). The registered and segmented images went through hand curation by an experienced investigator. During this process the ROI selection in the GC layer was restricted to cells with small soma size and sparse activity pattern while ambiguous ROIs (non-cellular structures, cells with big soma size and high activity) were discarded from the segmented data. In the hilar recordings only large, multipolar cells were accepted as putative MCs which had large amplitude calcium transients with fast onset and exponential decay. Delta F/F was calculated using the extracted raw calcium traces and neuropil signals with custom written code in Python as to the following. The unfiltered neuropil was first subtracted from the unfiltered raw trace. Two partial baselines were estimated then by filtering the residual raw trace, and the neuropil. The residual raw trace minus its partial baseline forms the numerator term of the dF/F. The denominator is the sum of the partial baselines. The partial baselines were calculated by applying a minimax filter to the respective (smoothed) traces.

To detect wake/sleep states, a Neuralynx Digital Lynx 4S system was used for EEG/EMG recordings under the 2-photon microscopy. EMG and EEG signals were acquired and filtered as described above. The EEG/EMG data was collected on a PC. Sleep states were calculated as described above. The up-regulated (i.e., more active during NREM sleep), down-regulated (i.e., less active during NREM sleep), and non-significant cells were defined by comparing each cell's activity between epochs of quiet wake (without motion) and NREM sleep (unpaired t-test).

To facilitate sleep under head-fixed condition, the mice underwent mild sleep deprivation during the night (from 5pm to 8am) before the recording session. Please note that the night is the active phase for mice. The mice were placed in a custom designed cage within a sound attenuated enclosure on a treadmill controlled by a microcontroller. The treadmill was set to 5s on and 10s off

cycles. On the morning of the two-photon recording sessions, the mice were transferred to the head-fixed apparatus from the sleep deprivation box.

Fiber Photometry

Fiber photometry recordings were performed as previously described (58). In brief, calcium dependent GCaMP fluorescence was excited by sinusoidal modulated LED light (473nm, 220 Hz; 405nm, 350 Hz, Doric lenses) and detected by a femtowatt silicon photoreceiver (New Port, 2151). Photometric signals and EEG/EMG signals were simultaneously acquired by a real-time processor (RZ5D, TDT) and synchronized with behavioral video recording. A motorized commutator (ACO32, TDT) was used to route electric wires and optical fiber. The collected data were analyzed by custom MATLAB scripts. They were first extracted and subject to a low-pass filter at 2 Hz. A least-squares linear fit was then applied to produce a fitted 405 nm signal. The DF/F was calculated as: (F-F0)/F0, where F0 was the fitted 405 nm signals. To compare activity across animals, photometric data were further normalized using Z-score calculation in each mouse. To analyze the slow oscillation of calcium signals, data were first downsampled to 1 Hz, then spectral analysis was carried out using FFT over a 2-min sliding window, sequentially shifted by 2 s increments (bins). The spectral power in the range of 0-6 cycles per min was used for statistical analysis.

Fluorescence in situ Hybridization (FISH)

Mice were deeply anaesthetized then decapitated and their brains were snap frozen on powdered dry ice. The fresh frozen brains were sectioned at 20 μm thickness using a cryostat. FISH was performed using RNAscope Multiplex Fluorescent Assay V2 (Advanced Cell Diagnostics) as per the manufacturer's recommendations. Reagents: *Htr1a in situ* hybridization probe: cat# 312301, *GFP in situ* hybridization probe: cat# 400281 (Advanced Cell Diagnostics). Images were acquired using a Zeiss 810 confocal microscope.

Histology

Viral expression and placement of optical implants were verified at the termination of the experiments using DAPI counterstaining of 100 μm coronal sections (Prolong Gold Antifade Mountant with DAPI, Invitrogen). Images were acquired using a Zeiss 810 confocal microscope. Cell numbers were counted manually in ImageJ.

Statistics

No method of randomization was used to determine how animals were allocated to experimental groups. Investigators were not blinded to group allocations. Mice in which the *post hoc* histological examination showed off target viral injection or fiber implantation were excluded from further analysis. Paired and unpaired t-tests were used and are indicated in the respective figure legends. All analyses were performed in MATLAB. Data are presented as mean ± s.e.m.

Code availability.

Custom scripts for EEG/EMG and behavioral analysis are available from the corresponding author upon reasonable request.

Data availability

All data supporting the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES:

1. C. Cirelli *et al.*, Reduced sleep in Drosophila Shaker mutants. *Nature* **434**, 1087-1092 (2005).
2. K. Koh *et al.*, Identification of SLEEPLESS, a sleep-promoting factor. *Science* **321**, 372-376 (2008).
3. D. M. Raizen *et al.*, Lethargus is a Caenorhabditis elegans sleep-like state. *Nature* **451**, 569-572 (2008).
4. F. Guo *et al.*, Circadian neuron feedback controls the Drosophila sleep--activity profile. *Nature* **536**, 292-297 (2016).
5. H. J. Kanaya *et al.*, A sleep-like state in Hydra unravels conserved sleep mechanisms during the evolutionary development of the central nervous system. *Sci Adv* **6** (2020).
6. C. C. Lo *et al.*, Common scale-invariant patterns of sleep-wake transitions across mammalian species. *Proc Natl Acad Sci U S A* **101**, 17545-17548 (2004).
7. R. P. Bartsch, A. Y. Schumann, J. W. Kantelhardt, T. Penzel, P. Ivanov, Phase transitions in physiologic coupling. *Proc Natl Acad Sci U S A* **109**, 10181-10186 (2012).
8. G. Z. Dos Santos Lima *et al.*, Hippocampal and cortical communication around micro-arousals in slow-wave sleep. *Sci Rep* **9**, 5876 (2019).
9. M. Bentivoglio, G. Grassi-Zucconi, The pioneering experimental studies on sleep deprivation. *Sleep* **20**, 570-576 (1997).
10. A. Rechtschaffen, M. A. Gilliland, B. M. Bergmann, J. B. Winter, Physiological correlates of prolonged sleep deprivation in rats. *Science* **221**, 182-184 (1983).

11. P. J. Shaw, G. Tononi, R. J. Greenspan, D. F. Robinson, Stress response genes protect against lethal effects of sleep deprivation in Drosophila. *Nature* **417**, 287-291 (2002).

12. G. Buzsáki, Two-stage model of memory trace formation: a role for "noisy" brain states. *Neuroscience* **31**, 551-570 (1989).

13. J. G. Klinzing, N. Niethard, J. Born, Mechanisms of systems memory consolidation during sleep. *Nat Neurosci* **22**, 1598-1610 (2019).

14. D. G. Amaral, A Golgi study of cell types in the hilar region of the hippocampus in the rat. *J Comp Neurol* **182**, 851-914 (1978).

15. Y. Senzai, G. Buzsaki, Physiological Properties and Behavioral Correlates of Hippocampal Granule Cells and Mossy Cells. *Neuron* **93**, 691-704 e695 (2017).

16. D. GoodSmith *et al.*, Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. *Neuron* **93**, 677-690.e675 (2017).

17. N. B. Danielson *et al.*, In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice. *Neuron* **93**, 552-559 e554 (2017).

18. M. Pofahl *et al.*, Synchronous activity patterns in the dentate gyrus during immobility. *Elife* **10**, e65786 (2021).

19. J. M. Monti, Serotonin control of sleep-wake behavior. *Sleep Med Rev* **15**, 269-281 (2011).

20. M. Jouvet, Sleep and serotonin: an unfinished story. *Neuropsychopharmacology* **21**, 24S-27S (1999).

21. S. H. Lee, Y. Dan, Neuromodulation of brain states. *Neuron* **76**, 209-222 (2012).

22. C. M. Teixeira *et al.*, Hippocampal 5-HT Input Regulates Memory Formation and Schaffer Collateral Excitation. *Neuron* **98**, 992-1004.e1004 (2018).

23. A. Luchetti *et al.*, Two functionally distinct serotonergic projections into hippocampus. *J Neurosci* **40**, 4936-4944 (2020).

24. K. Yoshida, M. R. Drew, M. Mimura, K. F. Tanaka, Serotonin-mediated inhibition of ventral hippocampus is required for sustained goal-directed behavior. *Nat Neurosci* **22**, 770-777 (2019).

25. L. Santarelli *et al.*, Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* **301**, 805-809 (2003).

26. A. Caspi, A. R. Hariri, A. Holmes, R. Uher, T. E. Moffitt, Genetic sensitivity to the environment: the case of the serotonin transporter gene and its implications for studying complex diseases and traits. *Am J Psychiatry* **167**, 509-527 (2010).

27. A. Holmes, R. J. Yang, K. P. Lesch, J. N. Crawley, D. L. Murphy, Mice lacking the serotonin transporter exhibit 5-HT(1A) receptor-mediated abnormalities in tests for anxiety-like behavior. *Neuropsychopharmacology* **28**, 2077-2088 (2003).

28. K. F. Tanaka, B. A. Samuels, R. Hen, Serotonin receptor expression along the dorsal-ventral axis of mouse hippocampus. *Philos Trans R Soc Lond B Biol Sci* **367**, 2395-2401 (2012).

29. K. Kohara *et al.*, Cell type-specific genetic and optogenetic tools reveal hippocampal CA2 circuits. *Nat Neurosci* **17**, 269-279 (2014).

30. J. Shen, H. Kudrimoti, B. McNaughton, C. Barnes, Reactivation of neuronal ensembles in hippocampal dentate gyrus during sleep after spatial experience. *Journal of Sleep Research* **7**, 6-16 (1998).

31.   E. Puighermanal *et al.*, drd2-cre:ribotag mouse line unravels the possible diversity of dopamine d2 receptor-expressing cells of the dorsal mouse hippocampus. *Hippocampus* **25**, 858-875 (2015).

32.   G. Gangarossa *et al.*, Characterization of dopamine D1 and D2 receptor-expressing neurons in the mouse hippocampus. *Hippocampus* **22**, 2199-2207 (2012).

33.   M. W. Jung, B. L. McNaughton, Spatial selectivity of unit activity in the hippocampal granular layer. *Hippocampus* **3**, 165-182 (1993).

34.   A. Bragin, G. Jando, Z. Nadasdy, M. van Landeghem, G. Buzsaki, Dentate EEG spikes and associated interneuronal population bursts in the hippocampal hilar region of the rat. *J Neurophysiol* **73**, 1691-1705 (1995).

35.   G. Buzsáki, Hippocampal sharp waves: their origin and significance. *Brain Res* **398**, 242-252 (1986).

36.   J. S. Farrell, E. Hwaun, B. Dudok, I. Soltesz, Neural and behavioural state switching during hippocampal dentate spikes. *Nature* 10.1038/s41586-024-07192-8 (2024).

37.   T. E. Scammell, E. Arrigoni, J. O. Lipton, Neural Circuitry of Wakefulness and Sleep. *Neuron* **93**, 747-765 (2017).

38.   T. Kato *et al.*, Oscillatory population-level activity of dorsal Raphe serotonergic neurons is inscribed in sleep structure. *J Neurosci* **42**, 7244-7255 (2022).

39.   J. Wan *et al.*, A genetically encoded sensor for measuring serotonin dynamics. *Nat Neurosci* **24**, 746-752 (2021).

40.   A. Muzerelle, S. Scotto-Lomassese, J. F. Bernard, M. Soiza-Reilly, P. Gaspar, Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5-B9) to the forebrain and brainstem. *Brain Struct Funct* **221**, 535-561 (2016).

41.   T. Tsetsenis, X. H. Ma, L. Lo Iacono, S. G. Beck, C. Gross, Suppression of conditioning to ambiguous cues by pharmacogenetic inhibition of the dentate gyrus. *Nat Neurosci* **10**, 896-902 (2007).

42.   G. Girardeau, V. Lopes-Dos-Santos, Brain neural patterns and the memory function of sleep. *Science* **374**, 560-564 (2021).

43.   B. A. Samuels *et al.*, 5-HT1A receptors on mature dentate gyrus granule cells are critical for the antidepressant response. *Nat Neurosci* **18**, 1606-1616 (2015).

44.   S. W. Hughes, M. L. Lőrincz, H. R. Parri, V. Crunelli, "Chapter 10 - Infraslow (<0.1Hz) oscillations in thalamic relay nuclei: basic mechanisms and significance to health and disease states" in Progress in Brain Research*, E. J. W. Van Someren, Y. D. Van Der Werf, P. R. Roelfsema, H. D. Mansvelder, F. H. Lopes Da Silva, Eds. (Elsevier, 2011), vol. 193, pp. 145-162.

45.   S. Lecci *et al.*, Coordinated infraslow neural and cardiac oscillations mark fragility and offline periods in mammalian sleep. *Sci Adv* **3**, e1602026 (2017).

46.   C. Kjaerby *et al.*, Memory-enhancing properties of sleep depend on the oscillatory amplitude of norepinephrine. *Nat Neurosci* **25**, 1059-1070 (2022).

47.   H. Antila *et al.*, A noradrenergic-hypothalamic neural substrate for stress-induced sleep disturbances. *Proc Natl Acad Sci U S A* **119**, e2123528119 (2022).

48.   H. Dong *et al.*, Genetically encoded sensors for measuring histamine release both in vitro and in vivo. *Neuron* 10.1016/j.neuron.2023.02.024 (2023).

49.   M. Jing *et al.*, An optimized acetylcholine sensor for monitoring in vivo cholinergic activity. *Nat Methods* **17**, 1139-1146 (2020).

50.	C. Kjaerby *et al.*, Memory-enhancing properties of sleep depend on the oscillatory amplitude of norepinephrine. *Nature Neuroscience* **25**, 1059-1070 (2022).
51.	J. S. Farrell *et al.*, Supramammillary regulation of locomotion and hippocampal activity. *Science* **374**, 1492-1496 (2021).
52.	S. Gonzalo Cogno *et al.*, Minute-scale oscillatory sequences in medial entorhinal cortex. *Nature* **625**, 338-344 (2024).
53.	S. Kim, D. Jung, S. Royer, Place cell maps slowly develop via competitive learning and conjunctive coding in the dentate gyrus. *Nat Commun* **11**, 4550 (2020).
54.	K. Juczewski, J. A. Koussa, A. J. Kesner, J. O. Lee, D. M. Lovinger, Stress and behavioral correlates in the head-fixed method: stress measurements, habituation dynamics, locomotion, and motor-skill learning in mice. *Sci Rep* **10**, 12245 (2020).
55.	G. F. Turi *et al.*, Vasoactive Intestinal Polypeptide-Expressing Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning. *Neuron* **101**, 1150-1165 e1158 (2019).
56.	J. C. Bowler *et al.* (2023) behaviorMate: An<i>Intranet</i>of Things Approach for Adaptable Control of Behavioral and Navigation-Based Experiments. (Cold Spring Harbor Laboratory).
57.	M. Pachitariu *et al.*, Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *bioRxiv* 10.1101/061507 %J bioRxiv, 061507 (2017).
58.	S. Teng *et al.*, Sensory regulation of absence seizures in a mouse model of Gnb1 encephalopathy. *iScience* **25**, 105488 (2022).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

A

Photometry

AAV-GRAB₅₋ₕₜ

EEG

EMG

DG

Raphe

fiber

B

| wake | NREM | REM | MA |

EEG

EMG

Photometry

1 mV

10% DF/F

3 min

2% DF/F

1 mV

30 s

C

% of outcome after 5-HT

Wake
NREM
REM
MA

4%

34%

61%

< 1%

D

Activity (Z)

Wake   NREM   REM

**

**

E

Cycle (/min)

Fig. 5

Fig. 6