2 Title: Investigating the developmental onset of regenerative potential in the annelid *Capitella teleta*

4 Authors: Alicia A. Boyd, Elaine C. Seaver

6 Whitney Laboratory for Marine Bioscience, University of Florida

8 Corresponding author: Elaine C. Seaver (seaver@whitney.ufl.edu)

10 Keywords: regeneration, cell division, larva, *vasa*, metamorphosis

Abstract:

An animal's ability to regrow lost tissues or structures can vary greatly during its life cycle. The annelid
Capitella teleta exhibits posterior, but not anterior, regeneration as juveniles and adults. In contrast,
embryos display only limited replacement of specific tissues. To investigate when during development *C.*
teleta becomes capable of regeneration, we assessed the extent to which larvae can regenerate. We
hypothesized that larvae exhibit intermediate regeneration potential and demonstrate some features of
juvenile regeneration, but do not successfully replace all lost structures. Both anterior and posterior
regeneration potential of larvae were evaluated following amputation. Wound sites were analyzed for
re-epithelialization, cell proliferation by EdU incorporation, stem cell and differentiation marker
expression by in situ hybridization, presence of neurites and muscle fibers by immunohistochemistry
and phalloidin staining respectively, and regrowth of structures. Wound healing occurred within 6 hours
of amputation for both anterior and posterior amputations. Cell proliferation at both wound sites was
observed for up to 7 days following amputation. In addition, the stem cell marker *vasa* was expressed at
anterior and posterior wound sites. However, growth of new tissue was observed only in posterior
amputations. Neurites from the ventral nerve cord were also observed at posterior wound sites. *De*
novo ash expression in the ectoderm of anterior wound sites indicated neuronal cell specification,
although the absence of *elav* expression indicated an inability to progress to neuronal differentiation. In
rare instances, cilia and eyes reformed. Both amputations induced expanded expression of the
myogenesis gene *MyoD* in pre-existing tissues. Our results indicate that amputated larvae complete
early, but not late, stages of regeneration, indicating a gradual acquisition of regenerative ability in *C.*
teleta. Furthermore, amputated larvae can metamorphose into burrowing juveniles, including those
missing brain and anterior sensory structures. To our knowledge, this is the first study to assess
regenerative potential of annelid larvae.

1. Introduction

Regeneration is the replacement of tissues or structures after injury or amputation. The ability to
regenerate lost tissues varies throughout the animal kingdom (Alvarado, 2000). For example, animals
such as planarians and hydra can completely reform any lost structure with hardly any restriction (i.e.,
whole-body regeneration) (Reddien, 2020; Reddy et al., 2020). Other animals have a more restrictive
capacity to regenerate and reform only specific structures, such as limbs in crustaceans (Alwes et al.,
2016) and amphibians (Brockes, 1997; Byrnes, 1904; Singer, 1954). Finally, some animals, like the leech,
display almost no regenerative capabilities following bisection (Özpolat & Bely, 2016). Here, we briefly
review differences in regeneration potential across animal life cycles and describe the need to more
carefully examine regeneration in pre-adult stages. We highlight annelids as a clade with exceptional
regeneration diversity and introduce *Capitella teleta* as a prime candidate to examine how regeneration
changes during the life cycle of an indirect developer.

Most published surveys of regeneration ability focus on adult stages, while few studies examine
regeneration potential at earlier stages of the life cycle. Several questions emerge when one considers
regeneration potential across the life cycle. In animals with a biphasic life cycle, does regeneration
ability differ among embryos, larvae, and adults? If so, when in the life cycle is the regeneration program
activated – during larval stages or after metamorphosis? Are there conserved trends of increasing or
decreasing regeneration abilities across the life cycle? From a developmental perspective, one might
expect a trend of decreasing regeneration abilities through the life cycle as embryonic cells become
committed to distinct cell types, undergo terminal differentiation, and lose their pluripotency. As a
result, there would be progressive loss in the ability to replace tissues (e.g., appendages, nervous
system, and heart) (Yun, 2015; Porello et al., 2011; Walters & Zuo, 2013). However, in animals with
indirect life cycles, larvae and adults often have distinct tissues, and adult tissue formation requires the
retention of progenitors or multipotent stem cells through the larval phase (Peterson et al., 1997). The
presence of stem cells in larvae suggests the possibility that animals with indirect development may
have greater regeneration potential relative to direct developing species.

The published studies that do examine regeneration in larvae are disproportionately sampled from
deuterostomes. From these examples, a complex picture emerges; a single relationship between larval
and adult regeneration ability does not explain published observations across species. For example, the
Xenopus frog (Phipps et al., 2020) has a pattern of regeneration loss similar to what is observed in direct
developers: an ability to replace specific tissues (i.e., appendages, heart, brain, lens) as larvae that is lost
with maturation (Harrison, 1898). The crinoid, *Antedon rosacea*, is very different such that the larvae
exhibit limited ability to regenerate following bisection, but adults are capable of whole-body
regeneration (Carnevali, 2006; Carnevali et al., 1993). Furthermore, sea stars exhibit whole-body
regeneration during larval, juvenile, and adult stages (Carnevali, 2006). Other animals have a more
dynamic change in regenerative ability through their life cycle. *Ciona intestinalis* lacks regenerative
abilities as a larva yet exhibits structural regeneration as adults only to progressively lose this ability with
advanced age (Jeffery, 2015). A broad understanding of how regeneration ability relates to the life cycle
is largely unexplored, and it is critical to sample pre-adult stages in animal taxa with demonstrated
regeneration abilities as adults, particularly from protostome lineages.

Annelids are segmented worms that have a wide range of regenerative abilities as adults. Some lack
the ability to regenerate following transverse amputation, others regenerate either anterior or posterior

segments only, or even undergo whole-body regeneration (Bely, 2006). A systematic review of annelid
regenerative ability revealed that anterior and posterior regeneration are widespread across the
phylum, alongside numerous examples of loss of regeneration (Bely, 2006). Furthermore, it is likely that
both anterior and posterior regeneration are ancestral for adult annelids (Zattara & Bely, 2016).
Although some studies have characterized regeneration during juvenile stages (del Olmo et al., 2021;
Kostyuchenko, 2022; Planques et al., 2019), to our knowledge, a broad assessment of larval regenerative
potential is lacking for annelids.

Capitella teleta BLAKE ET AL 2009 is an annelid capable of posterior regeneration and has an indirect
life cycle (Seaver, 2022). Previous studies hint at changes in the ability of *C. teleta* to replace lost
structures across its life cycle. Juvenile and adult worms regenerate following bisection along the main
body axis (de Jong & Seaver, 2016; Giani et al., 2011). In contrast, early-stage embryos do not regulate
following deletion of individual precursor cells (Pernet et al., 2012) (apart from eye and germline
precursor cells (Dannenberg & Seaver, 2018; Yamaguchi et al., 2016)). It is unknown when *C. teleta* gains
its regenerative ability, or to what degree larvae might be capable of regeneration.

Following *C. teleta* embryogenesis, a non-feeding lecithotrophic larva forms. Lecithotrophic larvae
often have 'abbreviated developmental times' (Vickery et al., 2001), which characterizes the *C. teleta*
larval phase of approximately 5 - 6 days in duration. Larvae are characterized by trochal ciliated bands
that divide the body into three regions along the anterior–posterior axis: the head, the segmented
trunk, and the pygidium. The prototroch separates the head from the trunk and the telotroch separates
the trunk from the pygidium (Fig 1A, B) (Seaver et al., 2005). These two ciliary bands are composed of
long cilia. A third ciliary band, the neurotroch, runs along the ventral midline of the trunk and is
composed of short cilia. Short cilia are also present in the pygidium. Larvae utilize the prototroch and
telotroch to swim in the water column, aiding in dispersal.

During larval stages, the adult centralized nervous system, musculature, digestive system, and initial
systems mature (Meyer et al., 2015). A previously described staging system uses morphological features
to identify nine distinct stages of embryonic (Stages 1-3) and larval development (Stages 4-9) (Seaver et
al., 2005). *C. teleta* has a centralized nervous system whose development is initiated in late
embryogenesis. A ladder-like ventral nerve cord (VNC) positioned along the midline of the trunk forms
and is connected to the brain by a pair of circumoral nerves (Meyer et al., 2015; Seaver et al., 2005). The
body wall musculature is organized segmentally with circular and longitudinal muscles, and the first
muscle fibers appear at the beginning of larval life (Stage 4). Two pigmented eyes appear at Stage 5 and
are located lateral and anterior to the prototroch (Meyer et al., 2015; Yamaguchi et al., 2016). Gut
organogenesis becomes visibly pronounced between Stages 6–9. The foregut consists of the mouth,
buccal cavity, pharynx (morphogenesis in Stage 7), and esophagus (Boyle & Seaver, 2008; see Fig 1A and
Fig 1B). In Stages 8–9, the midgut differentiates, becoming olive green and stereotypically coiled (Fig
1B). Stage 6 larvae have approximately 10 segments, generating one segment a day until there are
approximately 13 segments present in Stage 9 larvae (Seaver et al., 2005). Larval development
concludes when Stage 9 larvae metamorphose into burrowing juvenile worms (Cohen & Pechenik, 1999;
Seaver et al., 2005).

We sought to determine when regeneration potential is initiated during the life history of *C. teleta*.
We hypothesized that larvae may have limited regeneration potential distinct from juveniles and adults.
We anticipated that larvae would display early stages of regeneration without successfully replacing lost

tissues and structures. This would suggest that *C. teleta* acquires its regenerative ability gradually
through its life cycle. To test our hypothesis, we developed an amputation protocol for larvae to
investigate both anterior and posterior regeneration. We characterized larval responses to amputation
by making direct comparisons with regeneration events in juveniles. At the wound site, we documented
epithelial wound healing, evaluated cell proliferation by EdU incorporation, analyzed changes in
neuronal organization with antibody labeling, and visualized musculature with phalloidin staining. We
also conducted *in situ* hybridization experiments to analyze expression of the neuronal specification and
differentiation gene markers, *ash1* and *elav*, respectively, as well as examine muscle fate and
differentiation through analysis of the *MyoD* gene.

160
2.   Materials and Methods

2.1 Animal Care

As described in previous work (Grassle & Grassle, 1976), a colony of *Capitella teleta* was
maintained in the laboratory at 19°C in glass finger bowls containing filtered sea water (FSW) and a thin
layer of mud. Using forceps, larvae were dissected from the brood tubes of healthy females. Larvae from
a single brood tube develop synchronously and the number of individuals/brood averaged 100 – 250
individuals. Brood tubes were maintained in FSW until the desired age for amputation or treatment.

2.2 Amputations

Stage 6 larvae were used for amputation experiments. Larvae were developmentally staged
following a standard staging system (Seaver et al., 2005). Stage 6 is characterized by presence of yolk in
the anterior portion of the larva, swimming behavior with a positive phototactic response, and lack of
chaetae. Prior to amputation, larvae were incubated in 1:1 $MgCl_2$: FSW for 10 minutes to inhibit muscle
contraction. Individual larvae were transferred a 60 mm petri dish lid filled with a methyl cellulose
solution (0.3 g methyl cellulose in 15 ml filtered sea water) for amputation. Additional methyl cellulose
was added as necessary to achieve immobilization. Glass capillary tubes (50 µl, Becton Dickinson and
Company) were heated over a flame and pulled into thin, flexible, sharp tips to use as amputation tools.
Larvae were amputated with a single continuous, downward motion. Cut sites for anterior amputations
and eye amputations were located just posterior to the prototroch, removing the anterior third of the
animal (i.e., brain, eyes, and prototroch). Cut sites for posterior amputations were located ~2/3 of the
body length posterior to where the width of the trunk decreases and slopes to the pygidium, effectively
removing the posterior growth zone, telotroch and pygidium. Following amputation, each individual was
visually inspected for correct cut location and complete separation of the two tissue fragments.
Amputated larvae were transferred to a 60 mm petri dish containing 1:1 0.37M $MgCl_2$: FSW and
incubated for a minimum of 1 hour. Larvae were then washed into FSW containing 60 µg/ml penicillin
and 50 µg/ml streptomycin (Sigma-Aldrich Co.) (P/S) and raised at 19°C. P/S FSW was replaced daily, and
vitality and swimming behavior were monitored daily. For each amputation experiment, age matched
larvae from the same brood tube as experimental animals were raised as uncut controls. In a typical
experiment, 40 – 60 individuals from a single brood were amputated. For posterior amputations,
amputations were performed on sets of individuals from at least 10 broods, and for anterior

amputations, sets of individuals from at least 5 broods were amputated. Both anterior and posterior
amputated larvae had high rates of survivorship, ranging from 75% (n= 37/50) to 85% (n=28/33).

Two-week-old juveniles were prepared for amputation by incubation in 1:1 $MgCl_2$: FSW for 10
minutes. Individual worms were transferred to a platform of black dissecting wax (American Educational
Products, Fort Collins, CO, USA) in the lid of a 35 mm plastic dish for amputation in a drop of 1:1 $MgCl_2$:
FSW. For eye amputations, worms were amputated immediately anterior to the mouth, removing the
eyes and brain. Amputations for posterior regeneration were at the boundary between segment 10 and
11. Individual worms were visually inspected for the correct cut site and successful separation before
being transferred to a 60 mm petri dish and incubated in 1:1 $MgCl_2$: FSW for a minimum of 1 hour. The
juveniles were then washed into FSW and moved to a 60 mm petri dish containing mud and FSW and
raised at 19°C for 1 week. For analysis, juvenile worms were sifted from the mud and placed into a petri
dish of FSW. Worms were fixed with 4% paraformaldehyde (PFA) in FSW for 30 min before being washed
into phosphate-buffered saline (PBS). Fixed juveniles were scored for pigmented eye cells before being
processed for antibody labeling.

2.3 Cell Proliferation Assay

Larvae were incubated in EdU (Click-iT EdU Alexa Fluor 488 imaging kit - Invitrogen) diluted in
FSW to a final concentration of 3 uM at room temperature for 30 minutes. Larvae were then washed
into a 1:1 $MgCl_2$: FSW solution for 10 minutes and then fixed in 4% PFA for 30 minutes. Afterward, they
were rinsed into PBS and incubated for 20 minutes at room temperature in PBS + 0.05% Triton-X-100
with rocking. The specimens were incubated in the commercially provided EdU reaction buffer for 30
minutes, rocking at room temperature. Finally, larvae were washed into PBS and stored at 4°C until
further analysis.

2.4 EdU Quantification

Quantification of EdU and DAPI stained cells in uncut and amputated larvae was conducted by
importing Z-stacks generated from confocal microscopy (see Microscopy) into ImageJ software. Using
DAPI and DIC channels to establish boundaries, each image was cropped around the perimeter of the
animal. The channels were split and treated separately to optimize cell counting. The threshold for each
channel was adjusted manually to minimize background and enhance distinction of individual cells.
Afterward, a mask and watershed application were applied to each channel. The resulting image was
divided into three equal sections along the anterior-posterior axis using the Montage to Stack
application. The modified images were compiled into 20 slice projections (20 μm total) before being
counted automatically by Analyze Particles in the Analyze menu. Each 20 μm Z-stack was counted, with
the results automatically summarized via section. Manual inspection eliminated any cells that were
counted twice by the software. Results were manually compiled by channel and section for each animal,
as well as for each time point. A ratio of EdU stained cells to DAPI stained cells was calculated for each
time point from the ImageJ cell count summaries. Averages of each body section (anterior, middle, and
posterior) for each time point (uncut controls, 6 hpa, 3dpa, and 5dpa) were calculated separately. Ten
randomly selected larvae were scored for each time point and a mean value was calculated for each

body section across individuals. Each body section of uncut controls was compared with corresponding
body sections of amputated animals. One-way weighted ANOVA statistical tests were performed using
VassarStats (Lowry, 2023) to test for significance between the control and amputation group of interest.

2.5 Antibody labeling and phalloidin staining

Amputated larvae or juveniles were incubated in a 1:1 $MgCl_2$: FSW solution for 10 minutes and fixed in
4% paraformaldehyde in FSW for 30 minutes. Animals were washed several times into PBS to remove
fixative and then washed into a PBS + Triton solution (PBT) (0.1% Triton-X-100 for larvae and 0.2%
Triton-X-100 for juveniles). Animals were transferred to a 3 well glass depression plate and incubated in
a solution of 10% heat inactivated normal goat serum in PBT at room temperature with rocking for 45–
60 min. For antibody labeling, the anti-acetylated tubulin antibody (goat anti-mouse, Sigma-Aldrich Co.)
was diluted 1:400 in block solution (10% heated treated goat serum in PBT), and incubated with larvae
or juveniles at room temperature for 2 hours or overnight at 4°C. Following multiple washes in PBT (4
times in PBT for 30 minutes each), a secondary antibody (goat anti-mouse red 594, Invitrogen) was
diluted 1:400 in block solution and added to animals prior to incubation for 2–4 hours rocking at room
temperature or overnight at 4°C. The 22C10 antibody (Goat anti-mouse, Developmental Studies
Hybridoma bank; 22C10 was deposited to the DSHB by Benzer, S. / Colley, N. (DSHB Hybridoma Product
22C10)) was diluted 1:200 in block solution, added to animals and incubated overnight at 4°C. Following
multiple washes in PBT over two hours, the secondary antibody (goat anti-mouse green 488 - Invitrogen)
was diluted 1:400 in block solution, added to animals and incubated overnight at 4°C. Following
incubation in secondary antibody, larvae were rinsed 2–3 times in PBT and then washed 4 times in PBT
for 30 minutes each wash. For phalloidin staining, specimens were incubated in a 1:400 dilution of
phalloidin (Alexa Fluor 488 phalloidin, Invitrogen) in PBT. Specimens were incubated on a rocker at room
temperature for 45–60 min before being washed twice in PBT for 30 minutes each wash. Animals were
cleared in 80% glycerol in PBS with 1 µg/ml Hoechst-33342 (Molecular Probes) overnight prior to
imaging.

2.6 Whole mount in situ hybridization

Larvae collected at different time points following amputation were incubated in a 1:1 $MgCl_2$:
FSW solution for 10 minutes and then fixed in 4% paraformaldehyde overnight at 4°C. Fixative was
removed by multiple rinses in PBS and then larvae were gradually washed into 100% methanol and
stored at -20°C. Whole mount in situ hybridization was performed following a previously published
protocol (de Jong & Seaver, 2017). DIG-labeled RNA probes were generated with the MegaScript SP6 or
T7 Transcription kit (Invitrogen) according to manufacturer instructions. The following probes and their
working concentrations were used: *ash1* with 3ng/ul, *MyoD* with 0.5ng/ul, *vasa* with 1ng/ul, and *elav*
with 2ng/ul. Animals were hybridized for ~24 – 48 hours with riboprobe at 65°C. Each RNA probe was
visualized by exposure to an anti-digoxigenin-alkaline phosphatase conjugate antibody (Roche) prior to
exposure to a color reaction solution containing NBT/BCIP (nitro blue tetrazolium chloride/5-bromo-4-
chloro-3-indolyphosphate) and visually monitored for optimal development time. For each sample,
uncut control larvae were developed for the same duration as experimental, amputated larvae. Color
reactions were terminated following washes of PBT followed by a final fixation in 4% PFA in FSW for 30

min. Tissue was cleared in 80% glycerol in PBS plus 0.1 ug/ml Hoescht to stain nuclei, and then
individuals were placed on microscope slides for analysis and imaging. At least two independent
repetitions were performed for each gene.

2.7 Tissue and Cilia Measurements

To measure tissue formation in amputated larvae, confocal images were analyzed using ImageJ
software. Multiple measurements were made on each specimen. The nervous system was measured
from the cut site to the position most distal from the cut site that contained neurites. Muscle
measurements were taken to the position most distal from the cut site that contained phalloidin-
positive fibers. Changes in body width were calculated by making two measurements per specimen; the
first measurement was taken at the axial level of the mouth, and the second was just posterior to the
cut site. Gut measurements were made from the posterior end of the pigmented midgut to the
posterior edge of the posterior ectoderm. For anterior amputations, the muscle and nervous system
were measured on 6 hpa (n = 8) and 3 dpa (n = 10) larvae. For posterior amputations, the muscle,
nervous system, and changes in body width were measured for 6 hpa (n = 7) and 5 dpa (n = 10).
Additionally, the posterior edge of the midgut was used as a landmark for measurements in 5 dpa and
Stage 9 larvae at 6 hpa (n = 10). One-way weighted ANOVA statistical tests were performed using the
Good Calculators software (Good calculators, 2023) to test for significance between the uncut control
and experimental amputation group of interest.

The cilia of larvae were measured using ImageJ software on confocal image projections.
Individual labeled cilia of the prototroch, neurotroch, telotroch, and pygidium were traced. Lines
resulting from the trace were then measured using the 'measure' tool found under 'Analyze.' At least
three cilia for each ciliary band type were measured in a representative individual, and three individuals
were measured, resulting in a total of at least nine measurements for each of the prototroch, telotroch,
neurotroch and pygidium ciliary lengths. The range of cilia lengths across the nine measurements is
reported (see Results).

2.8 Metamorphosis assays

Metamorphosis is rapidly induced in *C. teleta* by exposure to vitamin B supplements (Burns et
al., 2014). A single multivitamin (Centrum Men Multivitamin) was dissolved in 200 ml FSW and stored at
room temperature. To test whether amputated larvae are capable of successful metamorphosis, uncut,
anteriorly amputated, and posteriorly amputated larvae were exposed to the multivitamin solution
containing vitamin B. Larvae were placed in a 30mm petri dish in FSW. Most of the FSW was removed
from the dish and replaced with 3 ml of the multi-vitamin solution. Animals were incubated in the
vitamin solution for 20 - 30 minutes. The multi-vitamin solution was removed by a minimum of three
washes with FSW (2 ml/wash), and then animals were individually scored for settlement and
metamorphosis. Successful settlement and metamorphosis were defined as cessation of swimming
behavior and initiation of burrowing, loss of the prototroch and telotroch ciliary bands, and elongation
of the body.

2.9 Microscopy and Imaging Analysis

Larvae were imaged with a Zeiss 710 confocal microscope using 4x bidirectional and
unidirectional scan settings for the green and red lasers. Z-stacks were compiled using ImageJ software
(Schindelin et al., 2012). Differential interference contrast microscopy images were captured on a Zeiss
AxioSkop II motplus compound microscope (Zeiss) coupled with a SPOT FLEX digital camera (Diagnostic
Instruments Inc). Multiple focal planes were merged for DIC images using Helicon Focus software
(Helicon Focus 7). Images were cropped and processed for brightness and contrast with Photoshop
(Adobe Photoshop 2020) or Illustrator (Adobe Illustrator 2020). Figures were composed in Adobe
Illustrator (Adobe Illustrator 2020).

3. Results

3. Results
3.1 Characteristics of successful regeneration in C. teleta

To characterize larval regenerative potential, we looked to the juvenile posterior regeneration
program in *C. teleta* to define features of successful regeneration. *C. teleta* juveniles maintain three
conserved sequential stages of posterior regeneration: wound healing, cell proliferation, and
differentiation of new tissues (de Jong & Seaver, 2016). Following amputation, juveniles wound heal in
4–6 hours and thereafter form a blastema (Fig 1C) (de Jong & Seaver, 2016). Neuronal projections
extend from the VNC into the wound site as early as 2 days post amputation (dpa) (Fig 1D; see also de
Jong & Seaver, 2016). In regenerating juveniles, the thickness of the connective nerve decreases
dramatically at the point of amputation, with multiple thin neural projections extending from the old
VNC tissue into the wound tissue and thereafter into the blastema/new tissue. This change in VNC
thickness indicates the position of the cut site. Localized cell proliferation is also observed at the wound
site by 2 dpa and persists for several days as new segments form (de Jong & Seaver, 2016). We observed
the density of circular and longitudinal muscle fibers decrease abruptly at the wound site at 3 dpa (Fig
1E). Circular and longitudinal muscle fibers are absent between the posterior boundary of the old tissue
and phalloidin-positive fibers at the posterior end of the animal. The decrease in muscle fibers spatially
align with the decrease in thickness of the connective nerves of the VNC. After approximately 7 days,
additional segments are formed that contain organized ganglia, peripheral nerves, circular and
longitudinal muscles, digestive tissue and other differentiated tissues (de Jong & Seaver, 2016).

3.2 Amputation responses in larvae and subsequent development

To assess the regenerative potential of *C. teleta* larvae, sites were selected for both anterior and
posterior amputations that removed approximately 1/3 of the body (Fig 2A). Initial analysis focused on
wound healing, which is identified by presence of a continuous epithelial layer covering the wound site
and results from contraction of the severed edges of the body wall. To assess wound healing in
amputated larvae, we analyzed Stage 6 larvae fixed at 2 hours post amputation (hpa), 3 hpa, 5 hpa, and
6 hpa for presence of a continuous epithelial layer covering the wound site. Following anterior
amputations, we observed consistent wound healing 6 hpa (Fig 2B) (n = 54/54). For posterior
amputations, we observed wound healing as early as 2 hpa (n = 3/15); however, in most cases successful
wound healing occurred around 6 hpa (Fig 2C) (n = 48/51). In posterior amputations performed on Stage
9 larvae, successful wound healing also occurred by 6 hpa (Fig 2D). Presence of tissue between the
posterior edge of the pigmented midgut and the posterior epidermis of most animals indicated wound
healing of the mesodermal layer by 6 hpa.

Larvae survived multiple days following amputation and continued to develop towards metamorphic
competency. Amputated larvae generally had high survivorship (~75-85%) at 5 dpa, with anteriorly
amputated animals and posteriorly amputated animals surviving up to 5 dpa and 7 dpa, respectively.
Furthermore, amputated larvae continued to develop on schedule relative to uncut animals (Fig 2E, 2F).
For example, in posteriorly amputated larvae, chaetae were present in the original tissue by 1 dpa
(comparable to Stage 7), pigmentation appeared in the midgut by 2 dpa (comparable to Stage 8), and
the midgut became coiled by 3 dpa (comparable to Stage 9). Anteriorly amputated larvae also
developed chaetae and a pigmented midgut, but the curvature of midgut—a hallmark of Stage 9 larvae
(Boyle & Seaver, 2008; Seaver et al., 2005) was delayed by one day. The growth of anterior and posterior
amputations differed. The shape of the body at the cut site in anterior amputations was frequently
observed to be blunt, similar in shape to a specimen following wound healing (6 hpa) (Fig 2F), whereas
the tissue distal to the cut site in posterior amputations had a rounded tapered shape (Fig 2E). The
tapered shape in posterior amputations is different from the shape of the wound site in aged-matched
individuals (Stage 9) fixed at 6 hpa (compare Fig 2D, E). Our observations demonstrate that larvae not
only survive removal of a third of their body, but they continue to develop with a similar time course to
uncut animals.

3.4 Cell proliferation

One key feature of regeneration is the birth of new cells in the region of the wound site. Larval
development is a dynamic growth process that also exhibits cell proliferation. Stage 6 and 7 larvae have
a characteristic pattern of EdU incorporation following a one hour exposure to EdU. Specifically, there
are numerous EdU+ cells distributed throughout the trunk, straddling the midline (Fig 3A, B). In contrast,
by Stage 9, EdU+ cells are not ordinarily observed, apart from a few cells in the midgut or hindgut and
posterior growth zone (PGZ) (Fig 3C, 3D). To delineate between growth and a response to amputation,
we compared age-matched, uncut larvae with amputated larvae.

Anterior amputations show EdU incorporation near the amputation site. The initial pattern of EdU
incorporation in anterior amputations (6 hpa) is similar to the pattern observed in age-matched, uncut
controls (Fig 3E, 3I). This pattern of EdU incorporation persists in 1 dpa animals and in age matched
controls (Fig 3B, F, J). However, by 3 dpa, EdU+ cells are localized near the wound site in the anterior
ectoderm and occasionally in the mesoderm of the amputated animals (Fig 3G). EdU+ cells near the
wound site persist through 5 dpa in anterior ectoderm and mesoderm (Fig 3H). In contrast, there is a
notable lack of EdU+ cells in the corresponding body region of unamputated animals (Fig 3C, 3D). The
ratio of EdU+ cells to nuclei is higher in amputated animals at both 3 dpa and 5 dpa at the respective
amputation sites compared with the equivalent body region in uncut controls (Table 1). There is a
significant difference between the number of EdU+ cells in the anterior body region anteriorly
amputated larvae at 3 dpa and 5 dpa when compared to age-matched, uncut controls (One-way ANOVA
test) (Fig 3M).

Posterior amputations also show EdU incorporation localized near the wound site. In both cut and
uncut larvae, the number of EdU+ cells decrease over time in the anterior and middle sections of the
animal. While the initial pattern of EdU incorporation in cells at 6 hpa appears similar to that observed in
Stage 6 uncut controls (Fig 3I), there is a progressive shift toward expression localized to the cut site in
amputated larvae. Animals at 1 dpa have fewer EdU+ cells in the anterior 2/3 of the body (Fig 3J) in
comparison with uncut Stage 7 controls. By 3 dpa, EdU incorporation is not detected through the

anterior 2/3 of the body, and EdU+ cells are restricted to the area surrounding the cut site (Fig 3K). This
localized pattern of EdU incorporation persists through 5 dpa (Fig 3L). Furthermore, cut larvae have
more EdU+ cells in the posterior region than the uncut controls (Table 2). The number of EdU+ cells in
posterior sections of 3 dpa and 5 dpa larvae is significantly higher compared to age matched, uncut
controls (One-way ANOVA test) (Fig 3N). Additionally, there is a higher ratio of EdU+ cells/nuclei
localized to the wound site in posterior amputated larvae relative to anterior amputations. Together,
these data show a marked increase in cell proliferation at the wound site following both anterior and
posterior amputations.

3.4 Expression of the stem cell marker vasa

To further characterize the larval response to amputation, expression of the stem cell marker *vasa*
was analyzed by *in situ* hybridization (Dannenberg & Seaver, 2018; Juliano et al., 2010). In uncut larvae,
vasa expression is dynamic from Stage 6 through Stage 9 (Dill & Seaver, 2008). In Stage 6 larvae, there is
expression in the foregut, the mesoderm of the trunk and the PGZ (Fig 4A). By Stage 8, expression is
restricted to the foregut, the multipotent progenitor cell cluster (MPC), and the PGZ (Fig 4B). This
pattern continues through Stage 9 (with some variability in detection of the MPC) (Fig 4C).

Vasa expression is detected at the wound site in both anterior and posterior amputations. Initially,
the *vasa* expression pattern in both amputation groups is very similar to the observed pattern in uncut
controls (6 hr post amputation) (Fig 4D). However, by 2 dpa, there is *de novo* expression in the anterior
ectoderm at the amputation site of anteriorly cut larvae (Fig 4E). By 5 dpa, expression at the wound site
is no longer detected, although expression in the MPC and PGZ persists (Fig 4F). In posterior
amputations at 2 hpa, trunk mesodermal expression is continuous from one side of the animal to the
other and wraps around the posterior end of the body (Fig 4G). We interpret this expression along the
posterior face of the animal to be the result of wound healing that pulls the two lateral trunk domains of
expression together. At 2 dpa, *vasa* is expressed in the posterior mesoderm and ectoderm near the
wound site (Fig 4H). This expression near the wound site is also detectable 3 dpa (Fig 4I) and 4 dpa (data
not shown). In summary, *vasa* expression is detected at the wound site 2–3 dpa for both anterior and
posterior amputations.

3.5 Generation of new tissue

The presence of localized cell proliferation at the wound site (Fig 3G, 3H, 3K, and 3L) and what
appears to be growth of new tissue (Fig 2E) warranted a more detailed investigation of the tissue
surrounding the wound site. To distinguish between pre-exiting tissue and new tissue, the location of
the cut site was determined using the nervous system and musculature markers. We used a framework
commonly used for identifying the cut site in juvenile worms with an anti-acetylated tubulin antibody
marking the nervous system (Fig 1B). An abrupt change in the thickness of the connective nerves was
used to indicate the position of the cute site (Fig 1D, 5B, 5B', and 5C). An abrupt decrease in the density
of muscle fibers also indicated the location of the cut site (Fig 1E, Fig 5F and 5F'). Typically, there was a
precise concordance in position between an abrupt change in muscle fiber density and thickness of
connective nerves. Therefore, we could consistently identify the location of the cut site using these two
markers.

The identification of the cut site in amputated larvae allowed us to identify new tissue and
quantify tissue distal to the amputation site. No evidence of significant growth of tissue was detected in

anterior amputations between 6 hpa and 3 dpa (see Table 3). In contrast, there was significant growth in
posterior amputations between 6 hpa and 5 dpa. Additional measurements of posterior amputations
indicated a significant change in body width (body width at the mouth compared to immediately
posterior to the cut site) between 6 hpa and 5 dpa. This information is consistent with the presence of
new growth in posterior amputations; regenerated tissue is often more narrow than the width of the
original tissue. While the position of the cut site indicated by muscle and nerves indicated new growth,
there was one measurement that did not. Measurements from the posterior end of the pigmented gut
to the posterior of the animal for both the 6 hpa stage 9 and 5 dpa larvae did not show a significant
difference between these two groups.

We next characterized the nervous system and musculature of the tissue distal to the cut site.
Similar to what was observed in 6 hpa control larvae (Fig 5D), anterior amputations in 3 dpa larvae
lacked clear evidence of neurites and muscle fibers distal to the wound site (Fig 5 E–E"). In posteriorly
amputated larvae, a mixture of results was observed. In 2 dpa posteriorly amputated larvae with a
discernable cut site and new growth, neuronal projections distal to the amputation site were observed
that were not detected in 6 hpa larvae (Fig 5B, 5F, 5F', 5F"). There were also cases in which there was
seemingly no change in the larval nervous system (Fig 5G; n = 165/395). Cases in which new growth was
not observed lend confidence to those instances when we observed processes distal to the cut site
(compare Fig 5F" and 5G). At 5 dpa, nerve processes persisted in tissue distal to the cut site (Fig 5H,
5H'). These longitudinal nerve processes are located at the midline and span the length of the new
tissue. Additionally, there were circumferential processes similar to the orientation of peripheral nerves,
although these processes were somewhat disorganized relative to peripheral nerves in uncut animals
(Fig 5H'). These neuronal processes are oriented medial to lateral, extending from the midline. The
entire region of the new tissue has a lower density of muscle fibers relative to the preexisting tissue (Fig
5B, 5H'). The muscle fibers that are present are predominately longitudinal muscle fibers. In many cases,
individual longitudinal muscle fibers appear to extend from preexisting tissue into tissue distal to the
amputation site (Fig 5H", arrow), whereas circular muscle fibers were only occasionally observed
posterior to the amputation site. These circular fibers are unlikely associated with a telotroch because
amputated larvae lack evidence of regrown ciliary bands. A similar arrangement of muscle and nerve
fibers in the tissue distal to the amputation site is present in animals 7 dpa (data not shown). While we
observed multiple tissue types distal to the cut site, we did not observe evidence of new segmentation.
Conventionally, newly formed segments are characterized by the appearance of new ganglia, regularly
spaced circumferential peripheral nerves and segmental boundaries. The nervous system distal to the
cut site did not organize into ganglia by 7 dpa, and peripheral nerves were disorganized. We did not
observe segmental boundaries in the ectoderm. In conclusion, new tissue growth was observed along
with changes in organization of the nervous system and musculature in posteriorly amputated larvae,
although there was a lack of new tissue in anteriorly amputated larvae.

3.6 Neural specification and differentiation

Next, we sought to determine if the neural and muscle cell types observed distal to the amputation
site resulted from the birth of new cells. We rationalized that expression of molecular markers of cell
fate specification and differentiation in the tissue distal to the amputation site indicated birth of new
cells. For the nervous system, gene expression patterns of neural specification (*ash1*) and neural
differentiation markers (*elav*) were analyzed near the wound site (Meyer & Seaver, 2009). In *C. teleta*,
ash1 is a marker of neural specification during initial neurogenesis of the brain and VNC (Meyer &

Seaver, 2009; Sur et al., 2017). In Stage 6 larvae, *ash1* is expressed in the brain, foregut, ventral neural
ectoderm, PGZ, and the mesoderm in the pygidium (Fig 6A). By Stage 9, *ash1* expression is restricted to
the foregut and PGZ (Fig 6B). In anterior amputations, the pattern observed at 6 hpa for *ash1* expression
mirrors that of uncut Stage 6 larvae, with the notable exception of missing brain expression due to the
removal of the head (Fig 6C). In 3 dpa specimens, *ash1* is expressed in both the anterior ectoderm and
pharynx (Fig 6D). Expression in the ectoderm can be regarded as *de novo ash1* anterior expression since
age-matched, uncut controls lack anterior ectoderm expression (compare panels Fig 6B and 6D, E) (n
=32/39). The novel anterior ectodermal expression of *ash1* at 3 dpa was present in a punctate pattern
similar to the pattern observed in the anterior ectoderm during *C. teleta* brain development. In contrast,
expression in the pharynx corresponds with expression in age matched uncut controls, albeit with fewer
cells.

After observing novel anterior neural specification following amputation, the next step of
neurogenesis was analyzed by characterizing the expression pattern of *elav*, a marker of differentiating
neurons. Between Stages 6 and 9 in uncut controls, *elav* is expressed in the brain and the VNC (Fig 6F &
G). In 6 hpa anterior amputations, *elav* is expressed only in the VNC (Fig 6H). This expression pattern
persists through 3 dpa, with no new expression domains observed in amputated animals (n =21/22) (Fig
6I). In the one differing case of the 22 scored animals, there were a few *elav*-expressing cells in the
anterior ectodermanteriorly located cells. These cells are part of the pharynx that abnormally everted
during morphogenesis.

3.7 Amputation induces expansion of MyoD expression

We also examined the expression of *MyoD*, a gene known for its role in muscle specification and
differentiation (Zammit, 2017). In uncut larvae, *MyoD* is expressed in the mesoderm of the posterior
trunk segments at Stage 6 (Fig 7A) and Stage 9 (Fig 7B). There is a developmental progression from
anterior to posterior in the larval trunk, with the more mature segments having an anterior position
(Seaver et al., 2005). Thus, *MyoD* is expressed in the younger and newly forming segments. Six hours
after both anterior and posterior amputations (6 hpa), expression is restricted to the posterior portion
of the trunk (Fig 7 C & E), similar to what is observed in uncut controls. At 7 dpa in anterior amputees
(Fig 7D) and at 3 dpa in posterior amputees (Fig 7F), there is a drastic expansion of *MyoD* mesodermal
expression along the entire length of the trunk. This expansion in *MyoD* expression is in pre-existing
tissue in older segments and includes tissue distant from the wound site in the case of posterior
amputations.

3.8 Rare regeneration of differentiated cell types

Following amputations, a few differentiated cell types occasionally regenerated, namely the
larval eyes and cilia. Larvae have two cerebral eyes, with each eye consisting of three cells: a pigment
cell (Fig 8A), a sensory cell (Fig 8C), and a support cell (Yamaguchi & Seaver, 2013). The pigment cell is
readily visible by its orange pigment granules and the sensory cell can be visualized by immunolabeling
with the antibody 22C10 (Yamaguchi & Seaver, 2013). Currently, we are unable to assess the presence
of the support cell. In uncut Stage 9 larvae, the pigment cells are located slightly anterior, if not inline,
with the prototroch, in a lateral position (Fig 8A). The sensory cell in uncut Stage 9 larvae (Fig 8C) is
located immediately adjacent and lateral to the pigment cell and has an axonal projection extending
from the cell body to the external surface of the ectodermal epithelium. The pigment and sensory cells
were scored as independent characters. Following head amputation, a few pigment cells regenerated,

and they are variable in location and number (n = 28/420; 3-5 dpa). Additionally, the pigment cells in
amputated larvae are ectopically positioned. For example, as seen in Fig 8B, there is a pigment cell in
proximity to the midgut. Regenerated sensory cells were also observed in amputated larvae and also
exhibited variability in location (n= 8/220, 3 –7 dpa). Axon projections in regenerated sensory cells
extended to the outer ectodermal surface of the animals (Fig 8D). There was a higher percentage of
cases with regenerated pigment cells (7%) compared with sensory cells (4%) in anterior amputations (Fig
8G). Summation of the total number of pigment cells and sensory cells indicates that at least one cell of
the eye regenerated in 6% of cases amputated (n = 36/640).

We were surprised by the appearance of new larval eye cells in amputated *C. teleta* larvae since
regeneration of anterior structures is not reported in adults (Yamaguchi & Seaver, 2013). Therefore, we
conducted a similar experiment with juveniles. Four-week-old juveniles were amputated immediately
anterior to the mouth, removing the eye cells. After screening for successful wound healing and visual
inspection that pigment cells were removed, a subset of the amputated juveniles was fixed at 4 hpa to
confirm the removal of the sensory cells (n =21/21). No pigment cells were observed in amputated
juveniles at 7 dpa (n = 0/43). However, a small proportion of cases with eye sensory cells were observed
(Fig 8E) (n = 6/43; 14%). Regenerated sensory cells were ectopically located (compare Fig 8E and 8F).
Five of the six juveniles with regenerated sensory cells had a small portion of brain present in the
anterior of the animal; only one case lacking a brain had a sensory cell. In summary, pigment cells did
not regenerate in juveniles, although sensory cells were detected in a small fraction of cases. These rare
instances of eye sensory cell regeneration represented the most dramatic phenotype observed following
juvenile anterior amputations.

There are four external ciliated structures in *C. teleta* larvae: prototroch, neurotroch, telotroch,
and pygidial cilia (Fig 9A, B). The prototroch and telotroch are composed of relatively long cilia
(prototroch cilia length ~25-30 microns and telotroch cilia length ~20-28 microns), and the neurotroch
and pygidium have short cilia (neurotroch cilia length ~ 8-10 microns, pygidium length ~5-7 microns). At
6 hpa, larvae lacked long cilia at the cut site (Fig 9 C and 9D), due to amputation of the prototroch and
telotroch in anterior and posterior amputations, respectively. At 5 dpa, short cilia (approximately 8
microns in length) were observed in rare cases (n = 12/161; 7%) at the anterior end of anteriorly cut
animals, distal to the amputation site (compare Fig 9E and F). Long cilia were not observed following
anterior amputations (n = 0/161). While rare (n = 22/370; 6%), both long (~25 microns) and short (~9
microns) cilia appeared distal to the cut site 3–7 days after posterior amputations (compare Fig 9G with
9H and 9H').

3.9 Metamorphosis following amputation

C. teleta is competent and can be induced to undergo metamorphosis into juveniles as Stage 9
larvae (Burns et al., 2014). Following amputation, larvae were screened for stereotypical signs of
successful settlement and metamorphosis following the definition of Cohen and Pechenik (1999). These
changes include a loss of swimming and initiation of burrowing behavior, loss of both ciliary bands
(telotroch and prototroch), and the appearance of an elongated, worm-like body shape. Uncut,
uninduced control larvae swam with their prototroch and telotroch bands intact for the duration of the
experiment (Fig 10A) (2 hr, n = 532/534). In contrast, virtually all larvae in the induced, unamputated
control group successfully underwent metamorphosis within two hours of exposure to the metamorphic
cue, settling on the bottom of the dish, dropping their cilia, adopting burrowing behavior, and extending

in length along the anterior-posterior axis (Fig 10B) (n = 203/207). Both sets of amputated larvae
followed a similar pattern to what was observed for unamputated larvae. Specifically, one day after
anterior amputation, larvae either continued to swim in the water column in the absence of a
metamorphic cue (Fig. 10C) (n=65/66) or settled to the bottom of the dish and underwent
metamorphosis within two hours of exposure to a metamorphic cue (Fig. 10D) (n=135/136). Similarly,
individuals whose posterior ends had been removed retained their prototrochal bands and continued to
swim in the absence of a metamorphic cue (Fig. 10E, arrow) (n=65/65), yet underwent rapid settlement
and metamorphosis within two hours of exposure (Fig. 10F) (n=143/144). In the absence of an induction
cue, amputated larvae continued to swim in the water column and retained their ciliary bands for
several days, demonstrating that amputation itself does not induce metamorphosis (Fig 9E and 9F).
Together, these results demonstrate *C. teleta* larvae can detect a metamorphic cue and successfully
execute metamorphosis in the absence of either anterior or posterior structures.

4. Discussion

4.1 Limited regeneration in C. teleta larvae

This study demonstrates that the initial events in response to amputation appear to be quite
similar between *C. teleta* larvae and juveniles. Following both anterior and posterior amputation, larvae
wound heal in a similar time frame as previously reported for posterior amputation of juveniles (i.e.,
within 6 hpa) (de Jong & Seaver, 2016). Following wound healing, EdU+ cells appear localized at the
wound site at 2 dpa in juvenile posterior regeneration, and the number of EdU+ cells increase over time
(de Jong & Seaver, 2016). EdU incorporation in anterior-facing wound sites in juveniles of *C. teleta* has
not been formally characterized. In both anterior and posterior larval amputations, EdU is incorporated
near the wound site by 3 dpa and this persists through 5 dpa, comparable to the pattern of EdU
incorporation at the wound site in regenerating juveniles. There is a difference in the number of EdU+
cells between anterior and posterior amputations in larvae, with posterior amputations exhibiting more
EdU+ cells. One difference in cell proliferation patterns between juveniles and larvae is a change in
position of the localized EdU+ cells. During posterior regeneration in juveniles, localization of EdU+ cells
progressively shifts anteriorly to a subterminal position, a shift that corresponds with the re-
establishment of the juvenile PGZ. We did not observe such a shift in larvae. It is interesting to note that
in adults of the annelid *Paranais litoralis*, BrdU is incorporated near the anterior wound site even though
these animals have lost the ability to regenerate their heads (Bely & Sikes, 2010). This finding, along with
our results in *C. teleta* larvae, emphasize that a block in regeneration can occur subsequent to the
initiation of cell division at the wound site, and cell division at the wound site is not a reliable predictor
of successful regeneration.

Interestingly, expression of the stem cell marker *vasa* at the wound site was observed in both
anteriorly and posteriorly amputated larvae. In successful posteriorly regenerating juveniles, *vasa* is
expressed locally at the wound site at 2 dpa and persists for several days (de Jong & Seaver, 2017). In
larvae, the duration of *vasa* expression and proportion of cases is different between anterior and
posterior amputations. In anterior amputations, *vasa* is expressed in the anterior ectoderm in
approximately 1/3 of cases and is transient, detectable at 2 dpa but not 5 dpa. Conversely, localized *vasa*
expression is observed at 2 dpa and persists through 5 dpa in posterior amputations. We interpret the
localized expression of *vasa* at the posterior wound site as either the result of *de novo* expression or
expression maintained from an earlier stage (i.e., 6 hpa). The spatial and temporal patterns of EdU

incorporation and *vasa* expression are very similar in posterior amputations. In contrast, *vasa* expression
has a shorter duration than EdU incorporation near the wound site in anterior amputations. We
hypothesize that these differences between anterior and posterior amputations is related to the new
tissue growth in posterior amputations that is absent following anterior amputations.

Additional features of juvenile posterior regeneration are the presence of neural projections
into the wound site and muscle reorganization distal to the amputation site. In juveniles, this
phenomenon is observed approximately at 2 dpa (de Jong & Seaver, 2016). In larval posterior
amputations, neurites are present distal to the cut site by 2 dpa. However, these neurites were not
observed in larval anterior amputations. With respect to muscle fiber organization, there is an abrupt
transition between the presence and absence of muscle fibers distal to the cut site in both juveniles and
larval posterior amputations that was not observed in larval anterior amputations. We were unable to
determine whether fibers present distal to the amputation site in posterior larval amputations resulted
from birth of new neural or muscle cells or from the extension of pre-existing cells.

To better understand the characteristics and development of the tissue distal to the cut site in
larvae, we characterized the specification and differentiation of cell types (i.e., neural cell types and
cilia) in amputated larvae. The timing of neural specification or differentiation in juveniles has not been
investigated in detail, but differentiated tissues are present by 7 dpa (de Jong & Seaver, 2016). Analysis
of *de novo ash1* expression indicates the specification of neural fates in larval anterior amputations.
Surprisingly, subsequent differentiation of neural tissues was not detectable, as illustrated by the
absence of *elav* expression in larval amputations. Anterior amputation of larvae results in incomplete
tissue and structure regeneration—an observation consistent with the response in juveniles to anterior
amputations. Despite a general lack of anterior structural differentiation in amputated larvae, we
surprisingly observed rare cases in which differentiated eye cells and cilia reappeared within 3–7 dpa.
Additionally, sensory eye cells (but not pigment cells) regenerated in juveniles at a higher proportion
than in larvae (14% in juveniles, 4% in larvae). While rare, the absence of eyes immediately following
amputation and the ectopic location of the observed eye cells made us confident that these cells are
regenerated eye cells in amputated larvae and juveniles. One possible explanation for the variation in
cilia and eye cell regeneration across individuals is variance in maternal nutritional output.

In summary, the beginning stages of regeneration (i.e., wound healing, localized cellular
proliferation, and expression of stem cell markers, etc.) occur in amputated larvae but there is a lack
complete replacement of lost tissue or of cell or tissue differentiation. These data indicate that *C. teleta*
larvae exhibit limited regeneration abilities relative to the successful posterior regeneration observed in
juvenile and adult stages. Additionally, detection of initial stages of regeneration in larvae suggests that
C. teleta has a gradual transition in regenerative potential, rather than a stark change in potential that
corresponds with the metamorphoric transition.

4.2 Amputation-induced expression in preexisting tissue

MyoD functions in muscle development and its expression in muscle progenitor cells is an indication
of myogenesis (Zammit, 2017; Rawls & Olson, 1997). In response to amputation, *MyoD* expression
expands across the length of the larval body in *C. teleta*, far from the wound site. This expanded
expression may result in an increase in muscle cell number to compensate for a decrease in body length
following amputation, although we did not observe an obvious increase in the number of muscle fibers 3
– 5 dpa. Alternatively, expanded expression of *MyoD* may be an injury-specific response that leads to

cell dedifferentiation and return to a precursor state. Such a precursor cell population may contribute to
regenerating tissue. It is notable that the induction of expanded *MyoD* expression occurs in the absence
of a complete regeneration response. Regardless, future functional work is needed to better understand
the implications of this change in *MyoD* expression in response to amputation. Changes in gene
expression in response to amputation has previously been documented in *C. teleta* and other annelids
(de Jong & Seaver, 2016; Kozin et al., 2017; Pfeifer et al., 2012; Takeo et al., 2008). However, most
examples documented to date are changes in expression of positional identity genes, and their new
expression likely reflects repatterning of the body fragment in response to amputation.

4.3 Metamorphosis following amputation

Amputated *C. teleta* larvae undergo successful metamorphosis. We were surprised to observe
metamorphosis following anterior amputations. In other spiralians, it has been hypothesized that the
brain and other anterior sensory organs function in the detection and response to metamorphic cues
(Chartier et al., 2018; Conzelmann et al., 2013; Hadfield, 2010; Hadfield et al., 2000). In *C. teleta*, chemo-
sensitive ciliated cells in the head ectoderm were proposed to relay metamorphosis signals directly to
the cerebral ganglia (Biggers & Laufer, 1999). More recently, a study hypothesized that the inductive
signal for metamorphosis is detected by sensory neurons that innervate the dorsal pharyngeal pad of
the pharynx (Biggers et al., 2012). If the metamorphic cue was detected by anterior neurons, we would
expect that removing the head would prevent metamorphosis. However, our results demonstrate that
the brain is not required for detecting the metamorphic cue or for undergoing metamorphosis in *C.
teleta*, supporting the hypothesis that detection of metamorphic cues in *C. teleta* likely occur in the
pharynx. It was similarly demonstrated in the polychaete *Hydroides elegans* that apical sensory neurons
are not necessary for successful metamorphosis (Nedved et al., 2021), suggesting that the brain may not
be necessary for metamorphosis in some annelids.

It is yet to be determined whether the amputations performed during larval stages have lasting
effects on survival and growth following metamorphosis. In addition, it remains to be seen whether
juveniles induced from posteriorly amputated larvae redevelop posterior growth zones, can generate
additional segments, or are capable of posterior regeneration. Future studies could examine the
regeneration potential of juveniles that result from posteriorly amputated larvae. Additionally, it could
be interesting to determine if juveniles induced from anterior amputated larvae are capable of posterior
regeneration in the absence of a brain.

4.4 Larval regeneration in marine invertebrates

The regenerative potential of an animal can vary significantly across its life stages. In the literature,
larvae with reduced regenerative ability relative to adults are described as having "attempted" or
"limited" regeneration (Vickery et al., 2001). Using this convention, we categorize *C. teleta* larvae as
having limited regeneration. Here, we show that *C. teleta* gains regeneration ability over its life cycle and
exhibits different regeneration abilities between larval and juvenile stages. To our knowledge, published
studies characterizing larval regeneration abilities in other annelid larvae are scarce.

Other bilaterian taxa suffer from a similar dearth in sampling larval and immature stages for their
regenerative abilities. Within Spiralia, larval regeneration has been examined in one nemertean species.
Similar to annelids, nemertean adult regeneration is variable (Zattara et al., 2019; Zattara & Fernández-
Alvarez, 2022). *Maculaura alaskensis* appears to have high regenerative potential as early stage larvae

that progressively decreases with maturation until it stabilizes as an adult capable of posterior
regeneration (Hiebert & Maslakova, 2015). Younger *M. alaskensis* larvae are five times more likely to
regenerate than older larvae following amputation (Moss, 2017). In Ecdoysozoa, the other protostome
clade, regeneration ability is generally restricted to appendages, rather than along the main body axis
(Brenneis et al., 2023). However, recent work shows that immature instars of the sea spider
Pycnogonum littorale can regenerate the midgut and gonads as well as appendages following bisection.
Adult counterparts of *P. littorale* cannot regenerate and do not even survive the amputations,
suggesting that regeneration ability is restricted to the immature instar stages (Brenneis et al., 2023).

Deuterostome larvae have been better surveyed for regeneration ability and there are numerous
documented examples. The hemichordate *Ptychodera flava* has the ability to repeatedly regenerate all
body parts when cut transversely or decapitated as an adult (Humphreys et al., 2022). Although *P. flava*
larvae can regenerate along its main body axis, it does not regenerate as well as the adults. In addition,
regeneration ability differs across the larval body. Specifically, ventral regions of *P. flava* larvae
regenerate better than dorsal regions and posterior segments regenerate better than anterior segments
(Luttrell et al., 2018). The larval tail of the cephalochordate *Branchistoma floridae* regenerates multiple
tissues faster than adults following similar amputations (Zou et al., 2021). Unlike other clades,
echinoderm larvae have been widely surveyed for regenerative potential in all classes with documented
adult whole body regeneration abilities (Cary et al., 2019; Vickery et al., 2001). Asteroids and echinoid
larvae, such as *Patiria miniate,* can regenerate completely, regardless of developmental stage,
demonstrating a stable regenerative ability throughout life (Oulhen, 2016; Carnevali, 2006; Vickery et
al., 2002). In crinoids, there appears a trend of increasing regenerative potential over developmental
time. For example, crinoid larvae display limited ability to regenerate body parts, with early-stage
Antedon rosacea larvae regenerate more poorly than late-stage larvae (Carnevali, 2006). In contrast,
adults can replace lost structures even when reduced to 1/5 of the original body size (Carnevali, 2006).
In summary, no clear trends of increasing or decreasing regeneration abilities across the life cycle
emerge from regeneration studies either within or across taxa, suggesting changes in regeneration
ability across the life cycle is a variable trait.

Although we cannot currently identify trends reflecting how regeneration abilities change between
larval and adult stages from a phylogenetic perspective, we can consider the impact of life history
factors. Factors such as feeding (de Jong & Seaver, 2016; Özpolat et al., 2016; Zattara & Bely, 2013),
duration of larval stage, and sublethal predation (Schoeman and Simon, 2023; Stewart, 1996) may
better correlate with the regenerative abilities observed in an individual species. Unlike *C. teleta*, *M.
alaskensis*, *P. flava*, and *P. miniate* larvae feed and display an ability to replace complex structures
following amputation. *M. alaskensis* and *P. miniate* can swim in the water column for weeks or months
(Moss, 2017; Cary et al., 2019) and *P. flava* for up to 300 days (Luttrell et al., 2018). In contrast, larvae of
C. teleta and the crinoid *A. rosacea* are non-feeding (Vickery et al., 2001) and have a short larval period
before undergoing metamorphosis. *C. teleta* larvae swim in the water column for hours to a few days,
and some crinoids for up to 10 days before settlement (Nakano et al., 2003). It has been previously
proposed that the longer an animal is in the water column and vulnerable to predation, the greater risk
there is of sublethal injury (Vickery & McClintock, 1998). Sublethal damage could act as an evolutionary
pressure to select for a robust regenerative ability. To that end, there may not be a strong selective
advantage for regeneration in non-feeding, short lived, brood dwelling larvae like *C. teleta*.

4.5 Future directions

This study demonstrates differences in regenerative ability across developmental stages of
Capitella teleta. Future studies could investigate physiological, cellular, and molecular differences
between larvae and juveniles in response to amputation. One such difference may relate to nutritional
status. It has been demonstrated that the extent of regeneration in *C. teleta* juveniles and other
annelids can fluctuate depending on nutrition intake (de Jong & Seaver, 2016; Zattara & Bely, 2013).
Because *C. teleta* larvae are non-feeding, limited energy resources may contribute to their regeneration
limitations. This nutrition hypothesis could be tested by two separate approaches. In the first, cells with
high yolk content can be deleted in early-stage embryos to reduce maternally contributed nutrition.
Previous work demonstrated that such deletions result in morphologically normal, albeit smaller larvae
(Pernet et al., 2012). These larvae would have a reduced nutritional supply and may exhibit a further
reduction in regenerative potential to what was observed in this study. The second approach would
involve investigating regeneration abilities in a different annelid with a feeding larval form. A feeding
larva with more nutritional resources may exhibit full regenerative abilities following amputation.
Additionally, an annelid with a longer larval period would have more time to completely regenerate lost
tissue following amputation during the early larval period, while also continually feeding.

The timing of cell maturation may also limit larval regeneration and explain differences between
larval and juvenile regenerative abilities in *C. teleta*. An example of one such developmental constraint
may be in the timing of stem cell maturation. A previous study proposed that stem cells travel from the
multipotent progenitor cell (MPC) cluster to the wound site during juvenile regeneration (de Jong &
Seaver, 2017). MPC cells in larvae may be immature, thereby preventing these stem cells from positively
contributing to successful regeneration. Specifically, nascent stem cells may not be capable of self-
renewal or migration, or of contributing to the blastema (Juliano et al., 2010). The number of cells in the
MPC cluster does not increase during larval development (de Jong & Seaver, 2017), suggesting these
cells may be quiescent during this time. A heterotopic transplant of a mature MPC cluster from juveniles
into larvae would test this hypothesis, and an increased regenerative ability in larvae following
transplantation would be expected. Such experiments in *C. teleta* regeneration might lead to a better
understanding of the role that development has on regeneration.

Future studies can characterize the regeneration ability of other annelid larvae, particularly
those whose adults have different regenerative potential than that of *C. teleta*. A correlation between
adult and larval regenerative potential within individual species may exist. For example, there may be
anterior regenerative ability in larvae in a species capable of anterior regeneration as an adult. If this
correlation holds, the larvae of *Chaetopterous pergamentaceus* (Cuvier & Latreille, 1829) (formally
variopedatus) would be predicted to regenerate both anterior and posterior structures. Adult *C.*
pergamentaceus can regenerate anteriorly following amputations at segment 15 or more anterior and
can regenerate posteriorly following amputation at any segment (Berrill, 1928; Cuvier & Latreille, 1829).
Extensive additional sampling is needed before we can meaningfully determine whether the pattern
observed in *C. teleta* is representative of other annelids. Additional sampling across annelids will also
provide insight as to the extent of variation in larval regeneration abilities, and whether larval
regenerative ability is as common as adult regenerative ability.

Acknowledgements

We thank Brent Foster, Lauren Kunselman, and Katie Feerst for critical reading of the manuscript.

Author Contributions

Conceived and designed the experiments: AAB ECS. Performed the experiments: AAB. Analyzed the
data: AAB ECS. Contributed reagents/materials/analysis tools: ECS AAB. Wrote the paper: AAB ECS.

References

Alvarado, A. S. (2000). Regeneration in the metazoans: why does it happen? *BioEssays*.
https://doi.org/10.1002/(SICI)1521-1878(200006)22:6

Bely, A. E. (2006). Distribution of segment regeneration ability in the *Annelida. Integrative and Comparative Biology*, *46*(4), 508–518. https://doi.org/10.1093/icb/icj051

Bely, A. E., & Sikes, J. M. (2010). Latent regeneration abilities persist following recent evolutionary loss in asexual annelids. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(4), 1464–1469. https://doi.org/10.1073/pnas.0907931107

Berrill, N.J. (1928). Regeneration in the polychaet *Chaetopterus variopedatus. Journal of Marine Biological Association of the United Kingdom*, 15(1), 151-158.
https://doi.org/10.1017/S0025315400055594

Biggers, W. J., & Laufer, H. (1999). Settlement and metamorphosis of *Capitella* larvae induced by juvenile hormone-active compounds is mediated by protein kinase C and ion channels. 196(2), 187–198. https://doi.org/10.2307/1542564

Biggers, W. J., Pires, A., Pechenik, J. A., Johns, E., Patel, P., Polson, T., & Polson, J. (2012). Inhibitors of nitric oxide synthase induce larval settlement and metamorphosis of the polychaete annelid *Capitella teleta. Invertebrate Reproduction and Development,* 56(1), 1–13.
https://doi.org/10.1080/07924259.2011.588006

Blake, J.A., Grassle, J.P., & Eckelbarger, K.J. (2009). *Capitella teleta*, a new species designation for the opportunistic and experimental *Capitella* sp. I, with a review of the literature for confirmed records. *Zoosymposia*, 2(1), 25– 53. DOI:10.11646/zoosymposia.2.1.6

Boyle, M. J., & Seaver, E. C. (2008). Developmental expression of *foxA* and *gata* genes during gut formation in the polychaete annelid, *Capitella* sp. I. *Evolution & Development*, 10(1), 89–105.
https://doi.org/10.1111/J.1525-142X.2007.00216.X

Brenneis, G., Frankowski, K., MaaB, L., & Scholtz, G. (2023) The sea spider *Pycnogonum littorale* overturns the paradigm of the absence of axial regeneration in molting animals. *PNAS*, 120(5).
https://doi.org/10.1073/pnas.221727120

Brockes, J. P. (1997). Amphibian limb regeneration: rebuilding a complex structure. *Science*, 276(5309), 81–87. https://doi.org/10.1126/SCIENCE.276.5309.81

Burns, R. T., Pechenik, J. A., Biggers, W. J., Scavo, G., & Lehman, C. (2014). The B vitamins nicotinamide (B3) and riboflavin (B2) stimulate metamorphosis in larvae of the deposit-feeding polychaete capitella teleta: implications for a sensory ligand-gated ion channel. *PLoS ONE*, 9(11).
https://doi.org/10.1371/journal.pone.0109535

Byrnes, E. F. (1904). Regeneration of the anterior limbs in the tadpoles of frogs. *Archiv Für Entwicklungsmechanik Der Organismen*, *18*(2), 171–177. https://doi.org/10.1007/BF02163652

Carnevali, C. (2006). Regeneration in echinoderms: repair, regrowth, cloning. *Invertebrate Survival Journal.* 3(1), 64-76.

Carnevali, M. D. C., Lucca, E., & Bonasoro, F. (1993). Mechanisms of arm regeneration in the feather star *Antedon mediterranea*: healing of wound and early stages of development. *Journal of Experimental*

Zoology, 267(3), 299–317. https://doi.org/10.1002/JEZ.1402670308

Cary, G. A., Wolff, A., Zueva, O., Pattinato, J., & Hinman, V. F. (2019). Analysis of sea star larval
regeneration reveals conserved processes of whole-body regeneration across the metazoa. *BMC*
Biology, 17(1), 16. https://doi.org/10.1186/s12915-019-0633-9

Chartier, T. F., Deschamps, J., Dürichen, W., Jékely, G., & Arendt, D. (2018). Whole-head recording of
chemosensory activity in the marine annelid *Platynereis dumerilii. Open Biology, 8*(10).
https://doi.org/10.1098/RSOB.180139

Cohen, R. A., & Pechenik, J. A. (1999). Relationship between sediment organic content, metamorphosis,
and postlarval performance in the deposit-feeding polychaete *Capitella* sp. I *Journal of*
Experimental Marine Biology and Ecology, 240, 1–18. https://doi.org/10.1016/S0022-
0981(99)00047-7

Conzelmann, M., Williams, E. A., Tunaru, S., Randel, N., Shahidi, R., Asadulina, A., Berger, J., Offermanns,
S., & Jékely, G. (2013). Conserved MIP receptor-ligand pair regulates *Platynereis* larval settlement.
Proceedings of the National Academy of Sciences of the United States of America, 110(20), 8224–
8229. https://doi.org/10.1073/PNAS.1220285110/-/DCSUPPLEMENTAL

Cuvier, G., & Latreille, P. A. (1829). Le règne animal distribué d'après son organisation, pour servir de
base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Chez Déterville.
https://doi.org/10.5962/bhl.title.49223

Dannenberg, L. C., & Seaver, E. C. (2018). Regeneration of the germline in the annelid *Capitella teleta*.
Developmental Biology, 440(2), 74–87. https://doi.org/10.1016/j.ydbio.2018.05.004

de Jong, D. M., & Seaver, E. C. (2016). A stable thoracic hox code and epimorphosis characterize
posterior regeneration in *Capitella teleta. PLOS ONE, 11*(2), e0149724.
https://doi.org/10.1371/journal.pone.0149724

de Jong, D. M., & Seaver, E. C. (2017). Investigation into the cellular origins of posterior regeneration in
the annelid *Capitella teleta. Regeneration (Oxford, England), 5*(1), 61–77.
https://doi.org/10.1002/reg2.94

del Olmo, I., Verdes, A., & Alvarez-Campos, P. (2022) Distinct patterns of gene expression during
regeneration and asexual reproduction in the annelid *Pristina leidyi. Journal of experimental*
zoology. doi: 10.1002/jez.b.23143

Dill, K. K., & Seaver, E. C. (2008). *Vasa* and *nanos* are coexpressed in somatic and germ line tissue from
early embryonic cleavage stages through adulthood in the polychaete *Capitella* sp. I. *Development*
Genes and Evolution, 218(9), 453–463. https://doi.org/10.1007/s00427-008-0236-x

Giani, V. C., Yamaguchi, E., Boyle, M. J., & Seaver, E. C. (2011). Somatic and germline expression of *piwi*
during development and regeneration in the marine polychaete annelid *Capitella teleta. EvoDevo,*
2(1). https://doi.org/10.1186/2041-9139-2-10

c. Good Calculators. [online software]. https://goodcalculators.com/one-way-anova-calculator/

Grassle, J., & Grassle, J. F. (1976). Sibling species in the marine pollution indicator *Capitella* (polychaeta).
Science 192(4239),567– 569. DOI: 10.1126/science.1257794

Hadfield, M. G. (2010). Biofilms and marine invertebrate larvae: What bacteria produce that larvae use

to choose settlement sites. *Annual review of marine science, 3*, 453–470.
https://doi.org/10.1146/ANNUREV-MARINE-120709-142753

Hadfield, M. G., Meleshkevitch, E. A., & Boudko, D. Y. (2000). The apical sensory organ of a gastropod
veliger is a receptor for settlement cues. *Biological Bulletin*, 198(1):67-76. doi: 10.2307/1542804

Harrison, R. G. (1898). The growth and regeneration of the tail of the frog larva - Studied with the aid of
Born's method of grafting. *Archiv Für Entwickelungsmechanik Der Organismen, 7*(2–3), 430–485.
https://doi.org/10.1007/BF02161494

Hiebert, T. C., & Maslakova, S. (2015). Integrative taxonomy of the *Micrura alaskensis* Coe, 1901 species
complex (Nemertea: Heteronemertea), with descriptions of a new genus maculaura gen. nov. and
four new species from the NE pacific. *Zoological Science, 32*(6), 615–637.
https://doi.org/10.2108/ZS150011

Humphreys, T., Weiser, K., Arimoto, A., Sasaki, A., Uenishi, G., Fujimoto, B., Kawashima, T., Taparra, K.,
Molnar, J., Satoh, N., Marikawa, Y., & Tagawa, K. (2022) Ancestral stem cell reprogramming genes
active in hemichordate regeneration. *Frontiers in Ecology and Evolution*.
doi:10.3389/fevo.2022.769433

Jeffery, W. R. (2015). Regeneration, stem cells, and aging in the tunicate *Ciona*: insights from the oral
siphon. *International Review of Cell and Molecular Biology, 319*, 255–282.
https://doi.org/10.1016/bs.ircmb.2015.06.005

Juliano, C. E., Swartz, S. Z., & Wessel, G. M. (2010). A conserved germline multipotency program.
Development, 137(24), 4113–4126. https://doi.org/10.1242/DEV.047969

Kostyunchenko, R.P. (2022) *Nanos* is expressed in somatic and germline tissue during larval and post-
larval development of the annelid *Alitta virens. Genes.*13, 270.
http://doi.org/10.3390/genes13020270

Lowry, R. (2023) vassarstats [online software]. http://vassarstats.net/anova1u.html

Luttrell, S. M., Su, Y. H., & Swalla, B. J. (2018). Getting a head with *Ptychodera flava* larval regeneration.
Biological Bulletin, 234(3), 152–164. https://doi.org/10.1086/698510

Meyer, N. P., Boyle, M. J., Martindale, M. Q., & Seaver, E. C. (2010). A comprehensive fate map by
intracellular injection of identified blastomeres in the marine polychaete *Capitella teleta. EvoDevo,*
1(1). https://doi.org/10.1186/2041-9139-1-8

Meyer, N. P., Carrillo-Baltodano, A., Moore, R. E., & Seaver, E. C. (2015). Nervous system development in
lecithotrophic larval and juvenile stages of the annelid *Capitella teleta. Frontiers in Zoology*, 12(1),
1–27. https://doi.org/10.1186/s12983-015-0108-y

Meyer, N. P., & Seaver, E. C. (2009). Neurogenesis in an annelid: Characterization of brain neural
precursors in the polychaete *Capitella* sp. I. *Developmental Biology, 335*(1), 237–252.
https://doi.org/10.1016/J.YDBIO.2009.06.017

Nakano, H., Hibino, T., Oji, T., Hara, Y., & Amemiya, S. (2003) Larval stages of a living sea lily (stalked
crinoid echinoderm). Nature 421, 158–160. https://doi.org/10.1038/nature01236

Nedved, B. T., Freckelton, M. L., & Hadfield, M. G. (2021). Laser ablation of the apical sensory organ of
Hydroides elegans (Polychaeta) does not inhibit detection of metamorphic cues. *The Journal of*

Experimental Biology, 224(20). https://doi.org/10.1242/JEB.242300

Oulhen, N., Heyland, A., Carrier, T. J., Zazueta-Novoa, V., Fresques, T., Laird, J., Onorato, T. M., Janies, D.,
& Wessel, G. (2016). Regeneration in bipinnaria larvae of the bat star *Patiria miniata* induces rapid
and broad new gene expression. *Mechanisms of Development, 142*, 10–21.
https://doi.org/10.1016/j.mod.2016.08.003

Özpolat, B. D., & Bely, A. E. (2016). Developmental and molecular biology of annelid regeneration: a
comparative review of recent studies. *Current Opinion in Genetics & Development, 40*, 144–153.
https://doi.org/10.1016/j.gde.2016.07.010

Özpolat, B.D., Sloane, E.S., Zattara, E.E., and Bely, A.E. (2016) Plasticity and regeneration of gonads in the
annelid *Pristina leidyi*. *EvoDevo* 7 (22). https://doi.org/10.1186/s13227-016-0059-1

Pernet, B., Amiel, A., & Seaver, E. C. (2012). Effects of maternal investment on larvae and juveniles of the
annelid *Capitella teleta* determined by experimental reduction of embryo energy content.
Invertebrate Biology, 131(2), 82–95. https://doi.org/10.1111/j.1744-7410.2012.00263.x

Phipps, L. S., Marshall, L., Dorey, K., & Amaya, E. (2020). Model systems for regeneration: *Xenopus*.
Development (Cambridge), 147(6). https://doi.org/10.1242/DEV.180844/223048

Planques, A., Malem, J., Parapar, J., Vervoot, M., & Gazave, E. (2019) Morphological, cellular and
molecular characterization of posterior regeneration in the marine annelid *Playnereis dummerilli*.
Developmental Biology 445(189-210). http://doi.org/10.10106/j.ydbio.2018.11.004

Rawls, A., & Olson, E. N. (1997). *MyoD* meets its maker. *Cell, 89*(1), 5–8. https://doi.org/10.1016/S0092-
8674(00)80175-0

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C.,
Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., &
Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods
2012 9:7, 9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schoeman, S., & Simon, C.A. (2023). Live to die another day: regeneration in *Diopatra aciculata* Knox
and Cameron, 1971 (Annelida: Onuphidae) collected as bait in Knysna Estuary, South Africa.
Biology. 12(3), 483. https://doi.org/10.3390/biology12030483

Seaver, E. C. (2022). Sifting through the mud: a tale of building the annelid *Capitella teleta* for EvoDevo
studies. *Current Topics in Developmental Biology, 147*, 401–432.
https://doi.org/10.1016/BS.CTDB.2021.12.018

Seaver, E. C., Thamm, K., & Hill, S. D. (2005). Growth patterns during segmentation in the two
polychaete annelids, *Capitella* sp. I and *Hydroides elegans*: comparisons at distinct life history
stages. *Evolution Development, 7*(4), 312–326. https://doi.org/10.1111/j.1525-142X.2005.05037.x

Singer, M. (1954). Induction of regeneration of the forelimb of the postmetamorphic frog by
augmentation of the nerve supply. *Journal of Experimental Zoology, 126*(3), 419–471.
https://doi.org/10.1002/jez.1401260304

Stewart, B. (1998). Sub-lethal predation and rate of regeneration in the euryalinid snake star
Astrobrachion constrictum (Echinodermata, Ophiuroidea) in a New Zealand fiord. *Journal of
Experimental Marine Biology and Ecology*. 199(2), 269-283. https://doi.org/10.1016/0022-
0981(95)00177-8

Sur, A., Magie, C. R., Seaver, E. C., & Meyer, N. P. (2017). Spatiotemporal regulation of nervous system
development in the annelid *Capitella teleta*. *8*, 13. https://doi.org/10.1186/s13227-017-0076-8

Vickery, M.S., McClinktock, J.B. (1998). Regeneration in metazoan larvae. *Nature*. 394, 140.
https://doi.org/10.1038/28086

Vickery, M. C. L., Vickery, M. S., Amsler, C. D., & McClintock, J. B. (2001). Regeneration in echinoderm
larvae. *Microscopy Research and Technique*, *55*(6), 464–473. https://doi.org/10.1002/jemt.1191

Vickery, Minako S., Vickery, M. C. L., & McClintock, J. B. (2002). Morphogenesis and organogenesis in the
regenerating planktotrophic larvae of asteroids and echinoids. *Biological Bulletin*, *203*(2), 121–133.
https://doi.org/10.2307/1543381

Wilhelm, E., Bückmann, D., & Tomaschko, K.H. (1997). Life cycle and population dynamics of
Pycnogonum litorale (Pycnogonida) in a natural habitat. Marine Biology 129, 601–606.
https://doi.org/10.1007/s002270050202

Yamaguchi, E., Dannenberg, L. C., Amiel, A. R., & Seaver, E. C. (2016). Regulative capacity for eye
formation by first quartet micromeres of the polychaete *Capitella teleta*. *Developmental Biology*,
410(1), 119–130. https://doi.org/10.1016/j.ydbio.2015.12.009

Yamaguchi, E., & Seaver, E. C. (2013). The importance of larval eyes in the polychaete *Capitella teleta*:
effects of larval eye deletion on formation of the adult eye. *Invertebrate Biology*, *132*(4), 352–367.
https://doi.org/10.1111/ivb.12034

Zammit, P. S. (2017). Function of the myogenic regulatory factors *Myf5*, *MyoD*, *Myogenin* and *MRF4* in
skeletal muscle, satellite cells and regenerative myogenesis. *Seminars in Cell & Developmental
Biology*, *72*, 19–32. https://doi.org/10.1016/J.SEMCDB.2017.11.011

Zattara, E. E., & Bely, A. E. (2013). Investment choices in post-embryonic development: quantifying
interactions among growth, regeneration, and asexual reproduction in the annelid *Pristina leidyi*.
Journal of Experimental Zoology Part B: Molecular and Developmental Evolution, *320*(8), 471–488.
https://doi.org/10.1002/jez.b.22523

Zattara, E. E., & Bely, A. E. (2016). Phylogenetic distribution of regeneration and asexual reproduction in
Annelida: regeneration is ancestral and fission evolves in regenerative clades. *Invertebrate Biology*,
135(4), 400–414. https://doi.org/10.1111/IVB.12151

Zattara, E.E. & Fernández-Alvarez, F.A. (2022) Collecting and culturing *Lineus sanguineus* to study
Nemertea WBR. In: Blanchoud S, Galliot B, editors. Whole-Body Regeneration: Methods and
Protocols [Internet]. New York (NY) Chapter 12. doi: 10.1007/978-1-0716-2172-1_12

Zattara E.E., Fernández-Álvarez F. A., Hiebert T. C., Bely A. E. & Norenburg J. L. (2019) A phylum-wide
survey reveals multiple independent gains of head regeneration in *Nemertea*. The Royal Society.
https://doi.org/10.1098/rspb.2018.2524

Zou, J., Wu, X., Shi, C., Zhong, Y., Yan, Q., Su, L., and Li, G. (2021) A potential method for rapid screening
of amphioxus founder harboring germline mutation and transgene. *Front. Cell Dev. Biol.* 9 (702290)
doi: 10.3389/fcell.2021.702290

Figure Legends

Figure 1. Larvae and regenerating juveniles of *C. teleta*. All images are oriented in ventral view with
anterior to the left. **A** Stage 6 larva. **B** Stage 9 larva. **C**, **D**, and **E** are images of the posterior end of a
regenerating juvenile 3 days post-amputation (dpa) and are from the same specimen. **C** Differential
interference contrast (DIC) image showing regeneration of new tissue in juveniles. **D** Nervous system
visualized with anti-acetylated tubulin antibody labeling. **E** Phalloidin staining shows muscle fibers with
few fibers in the regenerating tissue. The white dotted lines indicate amputation sites. The asterisk
marks the position of the mouth. N denotes the number of animals scored that resemble the image
shown. Cm, circular muscle; ey, eyes; g, gut; mg, midgut; np, neural projection; pt, prototroch; py,
pygidium; tt, telotroch.

Figure 2. Amputated larvae undergo successful wound healing. All images are oriented in ventral view
with anterior to the left. Arrows indicate the continuous layer of epithelium, indicative of successful
wound healing. **A** Stage 6 uncut larva showing experiment design for amputations. The dotted lines
indicate cut sites: (aa) anterior amputation and (pa) posterior amputation. **B** Stage 6 larva 6 hpa anterior
amputation. **C** Stage 6 larva 6 hpa posterior amputation. **D** Stage 9 larva 6 hpa posterior amputation. **E**
Stage 9 larva 2 dpa anterior amputation. **F** Larva 3 dpa anterior amputation. The black bracket
represents the tissue between the gut and ectoderm. Amputation site indicated by the black dotted line.
N denotes the number of animals scored that resembled the image in the panel. Scale bar is 50 μm.

Figure 3. Localized cell proliferation near the amputation site in larvae. All images are oriented in ventral
view with anterior to the left. Animals were exposed to EdU to label dividing cells. EdU+ cells are green.
A–D are uncut controls: **A** Stage 6, **B** Stage 7, **C** Stage 9, and **D** Stage 9 larvae. **E–H** are anterior
amputations: **E** 6 hpa, **F** 1 dpa, **G** 3 dpa, and **H** 5 dpa larvae. **I–L** are posterior amputations: **I** 6 hpa, **J** 1
dpa, **K** 3 dpa, and **L** 5 dpa larvae. **M**, **N** Quantification of EdU incorporation in anterior (M) and posterior
(N) amputations and comparison with uncut controls. Time after amputation is on the X-axis and the
ratio of EdU/DAPI cells is on the Y-axis. The white arrows indicate EdU+ cells near the wound site. The
developmental stage or time post-amputation is indicated in the bottom left corner of each panel. N
denotes the number of animals scored that resemble the image shown. Dark blue bars in the graphs
represent uncut controls and light blue bars represent amputated animals. Asterisk denotes significance
with $p < 0.05$. Ant amp, anterior amputation; Post amp, posterior amputation. Scale bar is 50 μm.

Figure 4. *Vasa* is expressed at the wound site in amputated larvae. All images are oriented in ventral
view with anterior to the left. The purple coloration indicates localization of *vasa* transcripts by in situ
hybridization. **A**, **B**, and **C** *Vasa* expression patterns in uncut larvae at Stage 6, Stage 8, and Stage 9. **D**, **E**,
and **F** are anterior larval amputations. **G**, **H**, and **I** are posterior larval amputations. **D** and **G** are
amputated larvae 6 hpa (D) and 2 hpa (**G**). **E** *De novo* expression is observed in anterior amputations at 2
dpa, as indicated by the arrow and is no longer detectable by 5 dpa (**F**). In posterior amputations,
expression near the wound site is observed at both 2 dpa (**H**) and 3 dpa (**I**), as indicated by arrows. The
time post-amputation is indicated in the bottom left corner of each panel in D–I. N denotes the number
of animals scored that resemble the image shown. Ant amp, anterior amputation; Fg, foregut; mpc,
multi-progenitor cell cluster; Post amp, posterior amputation; pgz, posterior growth zone. Scale bar is 50
μm.

Figure 5. Tissue response to amputation. **A–C** and **F–I"** are oriented in ventral view. **C–D"** are oriented in
lateral view. In all images, anterior is to the left. **A** uncut Stage 6 larva labeled with anti-acetylated
tubulin to visualize the nervous system and ciliary bands. Larvae are labeled with anti-acetylated tubulin
(red) and phalloidin (green) showing the nervous system and musculature for all panels except A, C and
G. Larvae shown in panels A, C and G were labeled with anti-acetylated tubulin. **B** and **B'** are the same
specimen 6 hpa. **C** neural fibers are present distal to the cut site in 2 dpa posterior amputee (arrow). **D**
larva 6 hpa after anterior amputation. **E, E'** and **E"** are the same specimen, 3 dpa anterior amputation. **E'**
magnified view of muscle fibers as marked by phalloidin. **E"** magnified view of anti-acetylated tubulin
labeling. **F, F',** and **F"** are the same larva 2 dpa. **F** Nervous system and musculature in a larva 2 dpa. **F'**
magnified view of posterior wound site showing few muscle fibers distal to cut site. **F"** magnified view of
posterior wound site showing fibers in the tissue distal to the amputation site (arrows). **G** shows a
specimen lacking neural fibers distal to the cut site. **H, H',** and **H"** are the same individual at 5 dpa
posterior amputation. **H'** neurites distal to the cut site (arrow). **H"** longitudinal and circular muscles are
present distal to the cut site (bracket). White brackets in H, **H'** and **H"** indicate tissue distal to the
amputation site. N denotes the number of animals scored that resemble the representative image.
Amputation site is indicated by a white dotted line. vnc, ventral nerve cord; pn, peripheral nerves. Scale
bar is 50 µm.

Figure 6. Amputation induces *ash1* but not *elav* expression at the anterior wound site. All images are
oriented with anterior to the left. **A-E** are ventral views and **F-I** are lateral views. The purple coloration
indicates expression of either *ash1* (**A-E**) or *elav* (**F-I**) transcripts. **C-E** and **H-I** are anterior amputations. **A**
Stage 6 uncut control. **B** Stage 9 uncut control. **C** 6 hpa. **D** and **E** are the same 3 dpa individual. Vertical
arrows indicate novel *ash1* ectodermal expression. The boundary between ectodermal and mesodermal
layers is indicated by a black dotted line. **F** Stage 6 and **G** Stage 9 uncut controls. **H** 6 hpa. **I** 3 dpa. N
denotes the number of animals scored that resemble the image indicated. Ant amp, anterior
amputation; br, brain; fg, foregut; vnc, ventral nerve cord. Scale bar is 50 µm.

Figure 7. Amputation induces expansion of *MyoD* expression. All images are oriented in ventral view
with anterior to the left. Localization of *MyoD* transcripts is visualized by the purple coloration. **A** Stage 6
and **B** Stage 9 uncut controls. *MyoD* is expressed in the posterior trunk mesoderm. **C** and **D** are anterior
amputations (ant amp). **C** 6 hpa. **D** anterior shift of anterior boundary of *MyoD* expression is observed at
7 dpa. **E** 6 hpa posterior amputation (post amp). Spots of staining anterior to the bracket is nonspecific
trapping in the chaetae. **F** *MyoD* is expressed along the length of the body in 3 dpa animals (compare
with E). Brackets indicate the regions of detectable *MyoD* expression along the anterior-posterior axis.
Amputation sites are indicated by black dotted lines. N denotes the number of animals scored that
resemble the representative image. Ant amp, anterior amputation; Post amp, posterior amputation.
Scale bar is 50 µm.

Figure 8. Rare eye reformation following anterior amputation. All images are oriented with anterior to
the left. **A – D** are ventral views. **A** DIC image of the anterior end of a Stage 9 uncut larva. **B** DIC image of
the anterior end of a larva 5 days after anterior amputation. The image shown is a merge of multiple
focal planes. The pigmented cells (pc) of the eye are indicated by arrows. **C–F** eye sensory cells (sc, white
arrows), labeled by the 22C10 antibody (green). **C** Uncut Stage 9 control. **D** anterior amputated larva 5
dpa. **E** and **F** are juvenile worms in lateral view. **E** uncut 2 week-old juvenile and **F** juvenile 1 week post-
amputation of the head. **G** Table displaying the number of pigment cells, sensory cells, or either cell

(total) present in anteriorly amputated larvae at 6 hpa and 3 dpa or later. The dotted line indicates the
outline of the animal. Juv, juvenile; mg, midgut; pc, pigment cells; sc, sensory cells.

Figure 9. Rare reformation of cilia occurs in amputated larvae. Animals are labelled with anti-acetylated
tubulin (white). All images are oriented in with anterior to the left. **A, C, E, B,** and **F** are oriented in
lateral view. **B, D, G, H,** and **H'** are oriented in ventral view. **A** Stage 6 uncut larva. **D** Stage 9 uncut larva.
B and **E** are anterior amputations. **B** larva 6 hpa. **E** larva 5 dpa with short cilia visible distal to the cut site
(arrow). **D, H,** and **H'** are posterior amputations in ventral view. **D** larva 6 hpa. **H** and **H'** larva 5 dpa with
both long cilia (LC) and short cilia (SC) distal to the cut site (arrows). **H'** is a magnification of **H**. N denotes
the number of animals scored that are similar to the image represented. Amputation site is indicated by
the white dotted line. Ant amp, anterior amputation; lc, long cilia; nt, neurotroch; post amp, posterior
amputation; pt, prototroch; pyg, pygidium; sc, short cilia; tt, telotroch. Scale bar is 50 μm.

Figure 10. Induction of metamorphosis in amputated larvae. All images are oriented in ventral view with
anterior to the left. **A** Uncut Stage 9 larva. **B** Uncut animal 2 hr after induction with metamorphic cue. **C**
anteriorly amputated larva not exposed to metamorphic cue. **D** anteriorly amputated animal 2 hr after
exposure to metamorphic cue. **E** posteriorly amputated larva not exposed to metamorphic cue. **F**
posteriorly amputated animal 2 hr after exposure to metamorphic cue. N denotes the number of
animals scored that are similar to the image shown. Ant amp, anterior amputation; Br, brain; ph,
pharynx; post amp, posterior amputation; pt, prototroch; tt, telotroch. All images have the same
magnification; scale bar, 50 μm.

1067

1068

A — aa ― pa

B — n = 54/54 — ant amp

C — n = 48/51 — post amp

D — 6 hpa — n = 10/10

E — 2 dpa — n = 10/10

F — 3 dpa — n = 10/10

ash1

elav

Larval pigment cell	Larval sensory cell	Juv sensory cell
A Cont — pc, pc — St 9	**C** sc — St 9	**E** sc — 2 wk
B Amp — pc — 5 dpa	**D** sc — 7 dpa	**F** sc — 1 wpa

G

	Pigment cell	Sensory cell	Total
6 hpa	0/46	0/31	0/57
3 dpa +	28/420	8/220	36/640
	7%	4%	6%

Uncut controls | Ant amp | Post amp

A — Uninduced — n=532/534 — br, ph, pt, tt

C — n=65/66 — ph, tt

E — n=65/65 — br, ph, pt

B — Induced — n=203/207 — br, ph

D — n=135/136 — ph

F — n=143/144 — br, ph

Table 1. Mean values of EdU+ cells in anterior amputations

Anterior Amp		Anterior		Middle		Posterior	
		Uncut	Cut	Uncut	Cut	Uncut	Cut
6 hpa	# EdU	133	102	466	235	387	136
	# Nuclei	418	392	1216	417	551	250
	%	32%	26%	38%	56%	70%	54%
3 dpa	# EdU	3	29	18	2	14	24
	# Nuclei	447	363	856	457	413	185
	%	1%	8%	2%	0%	3%	13%
5dpa	# EdU	3	28	18	5	14	10
	# Nuclei	147	359	856	516	413	219
	%	2%	8%	2%	1%	3%	5%

Table 2. Mean of EdU+ cells in posterior amputations

Posterior Amp		Anterior		Middle		Posterior	
		Uncut	Cut	Uncut	Cut	Uncut	Cut
6 hpa	**# EdU**	133	70	466	217	387	225
	# Nuclei	418	155	1216	556	551	463
	%	32%	45%	38%	39%	70%	49%
3 dpa	**# EdU**	3	2	18	6	14	78
	# Nuclei	447	306	856	702	413	525
	%	1%	1%	2%	1%	3%	15%
5dpa	**# EdU**	3	0	18	0	14	51
	# Nuclei	147	331	856	547	413	513
	%	2%	0%	2%	0%	3%	10%

Table 3. Measurement of tissue distal to cut site

		Muscle	Nervous System	Gut	Width 2 /Width 1
Ant Amp	**6 hpa**	19 μm	18 μm	N/A	N/A
	3 dpa	22 μm	15 μm	N/A	N/A
Post Amp	**6 hpa**	* 17 μm	* 12 μm	N/A	0.675
	5 dpa	* 62 μm	* 67 μm	* 45 μm	0.453
	St 9 6 hpa	N/A	N/A	* 37 μm	N/A

* = significance, p < .05