2   Title: Investigating the developmental onset of regenerative potential in the annelid *Capitella teleta*

4   Authors: Alicia A. Boyd, Elaine C. Seaver

6   Whitney Laboratory for Marine Bioscience, University of Florida

8   Corresponding author: Elaine C. Seaver (seaver@whitney.ufl.edu)

10   Keywords: regeneration, cell division, larva, *vasa*, metamorphosis

Abstract:

An animal's ability to regrow lost tissues or structures can vary greatly during its life cycle. The annelid
Capitella teleta has robust posterior, but not anterior, regeneration as a juvenile and an adult. In
contrast, embryos display only limited replacement of specific tissues. To investigate when during
development *C. teleta* becomes capable of regeneration, we assessed the extent to which larvae can
regenerate. We hypothesized that larvae exhibit intermediate regeneration potential and demonstrate
some features of juvenile regeneration, but do not successfully replace all lost structures. Both anterior
and posterior regeneration potential of larvae were evaluated following amputation. Responses were
analyzed by EdU incorporation, immunohistochemistry, phalloidin staining, and in situ hybridization.
Wound sites were evaluated for re-epithelialization, cell proliferation, stem cell marker expression, the
presence of neurites and regrowth of structures. Wound healing occurred within 6 hours of amputation.
Cell proliferation at the wound site was observed for up to 7 days following amputation. In addition, the
stem cell marker *vasa* was expressed at anterior and posterior wound sites. However, growth of new
tissue was observed only in posterior amputations. Neurites from the ventral nerve cord were also
observed at posterior wound sites. *De novo ash* expression in the ectoderm of anterior wound sites
indicated neuronal cell specification, although the absence of *elav* expression indicated an inability to
progress to neuronal differentiation. In rare instances, cilia and eyes reformed. Both amputations
induced ectopic expression of the myogenesis gene *myoD*. Our results indicate that amputated larvae
complete early, but not late, stages of regeneration, indicating a gradual acquisition of regenerative
ability in *C. teleta.* Furthermore, amputated larvae can metamorphose into burrowing juveniles,
including those missing brain and anterior sensory structures. To our knowledge, this is the first study to
assess regenerative potential of annelid larvae.

1. Introduction

Regeneration is the replacement of tissues or structures after injury or amputation. The ability
to regenerate lost tissues varies throughout the animal kingdom (Alvarado, 2000). Animals such as
echinoderms can to completely reform any lost structure (i.e., whole-body regeneration) with hardly
any restriction. Other animals have a more restrictive capacity to regenerate and reform only specific
structures; for instance, limbs in amphibians (Brockes, 1997; Byrnes, 1904; Singer, 1954) or the heart in
spiny mouse (Gaire et al., 2021). Finally, there are animals like the leech that display almost no
regenerative capability (Özpolat & Bely, 2016).

The capacity to regenerate can also vary throughout an animal's life cycle. Some animals lose
regenerative potential as they mature. For example, the *Xenopus* frog has robust regenerative ability to
replace tissues (i.e., appendages, nervous system, and heart) as a tadpole (Harrison, 1898), but with age
this regenerative ability is progressively lost (Phipps et al., 2020). Other animals gain regenerative
potential with age. Crinoids, such as *Antedon rosacea,* have a limited ability to regenerate as larvae, but
are capable of whole-body regeneration as adults (C. Carnevali, 2006; M. D. C. Carnevali et al., 1993).
Ascidians such as *Ciona* have a more dynamic change in regenerative ability through their life cycle.
Ciona intestinalis lacks regenerative ability as larvae yet exhibit robust structural regeneration as adults
only to progressively lose this ability with advanced age (Jeffery, 2015). Other animals exhibit robust
regenerative potential throughout their entire life cycle. For example, sea stars exhibit whole-body
regeneration at larval, juvenile, and adult stages (C. Carnevali, 2006). Most published surveys of
regeneration ability focus on adult stages, while few studies examine regeneration potential throughout
the life cycle. Furthermore, the mechanisms that control changes in regeneration abilities across the life
cycle are unknown. This historically narrow focus may be expanded by appraising larval regeneration in
a range of indirect developing animals.

Annelids are segmented worms that, as a phylum, have a wide range of documented
regenerative abilities as adults. There are worms that lack the ability to regenerate following transverse
amputation, have partial regeneration (e.g., anterior or posterior only), or undergo whole-body
regeneration (A. E. Bely, 2006). A systematic review of regenerative ability revealed that anterior and
posterior regeneration are widespread across the phylum, alongside numerous examples of loss of
regeneration (A. E. Bely, 2006). Furthermore, it is likely that both anterior and posterior regeneration
are ancestral for adult annelids (Zattara & Bely, 2016). It is not known if such a range of regeneration
potential exists at other stages of the annelid life cycle. To our knowledge, a broad assessment of larval
regenerative potential is lacking for annelids.

Several features of *C. teleta* biology make it an excellent choice for contributing to research of
regeneration potential across the lifespan. First, previous studies indicate there is a change in the
regeneration ability across the life cycle of *C. teleta*. Juvenile and adult worms are capable of robust
posterior regeneration (de Jong & Seaver, 2016; Giani et al., 2011). In contrast, early-stage embryos
generally do not regulate following the laser deletion of individual cells (Pernet et al., 2012) (apart from
eye and germline precursor cells (Dannenberg & Seaver, 2018; Yamaguchi et al., 2016)). It is not
currently known when *C. teleta* gains its structural regenerative ability, or to what degree larvae might
be capable of regeneration. Second, worms can be maintained in the laboratory and bred throughout
the year, providing consistent production of embryos, larvae, and juveniles for experimentation. Since
the sequencing of the *C. teleta* genome (Simakov et al., 2013), many tools are now available for
experimental manipulation in this animal. Methods for analyzing spatiotemporal gene expression such

as in situ hybridization, antibody labeling, and cell fate maps are well-established and readily available
(Meyer et al., 2010).
111
We sought to determine when regeneration potential is initiated during the life history of *C.*
teleta. We hypothesized that *C. teleta* larvae may have limited regeneration potential distinct from
juveniles and adults. We anticipated that limited potential would manifest itself as displaying early
stages of regeneration without successfully replacing lost tissues and structures. Such a scenario would
suggest that *C. teleta* acquires its regenerative ability gradually through its life cycle. Alternatively, the
onset of regenerative potential may coincide with metamorphosis. Such a rapid shift would be indicated
by a complete lack of regenerative features in larvae. To test our hypotheses, we developed an
amputation protocol for larvae to investigate both anterior and posterior larval regeneration. We
characterized responses of *C. teleta* larvae to amputation by making direct comparisons with
regeneration events in juveniles. At the wound site, we documented epithelial wound healing, utilized
EdU incorporation to evaluate cell proliferation, antibody labeling to analyze changes in neuronal
organization, and phalloidin staining to visualize the musculature. We also conducted in situ
hybridization to analyze the expression of *ash1* and *elav*, neuronal specification and differentiation
genes, respectively, as well as *MyoD*, a muscle differentiation gene.

3. Results

3.1 Capitella teleta larval development

Capitella teleta is an annelid with an indirect life cycle that is an emerging model for
development and regeneration studies (Seaver, 2022)(Seaver & De Jong, 2021). It is commonly found in
muddy estuaries (Cohen & Pechenik, 1999). During embryonic and larval stages, the adult centralized
nervous system, musculature, digestive system, and initial segments develop (Meyer et al., 2015). The
non-feeding larvae utilize their larval-specific cilia to swim, aiding in dispersal. Competent larvae
metamorphose into burrowing juvenile worms, which reach sexual maturity after approximately 8
weeks.

C. teleta juveniles successfully regenerate in the posterior direction (de Jong & Seaver, 2016).
There are three conserved sequential stages of regeneration: wound healing, cell proliferation, and
differentiation of new tissues. Following amputation, juveniles wound heal in 4–6 hours and thereafter
form a blastema (Fig 1A) (de Jong & Seaver, 2016). Neuronal projections extend from the ventral nerve
cord (VNC) into the wound site as early as 2 days post amputation (dpa) in amputated juveniles (Fig 1B),
consistent with previously published research (de Jong & Seaver, 2016). In these regenerating juveniles,
the thickness of the connective nerve decreases dramatically at the point of amputation, with multiple
thin neural projections extending from the old VNC tissue into the wound tissue and thereafter into the
blastema/new tissue. This change in the thickness of the VNC can be used as an indicator of the position
of the cut site. A localized area of cell proliferation is also observed at the wound site 2 dpa and persists
for several days as new segments are formed (de Jong & Seaver, 2016). The musculature, as visualized
by phalloidin staining, shows an abrupt change in density of the circular and longitudinal muscle fibers at
3 dpa (Fig 1C); this change in density aligns with the cut site (designated by the acetylated tubulin
labelling of the VNC). Specifically, there is a lack of circular muscle and few longitudinal muscle fibers
between the old tissue and phalloidin-positive fibers at the posterior end of the animal. Distal to this
area of reduced muscles, there is a circular muscle associated with the anus in the new tissue (Fig 1 C).

After approximately 7 days, a new pygidium and additional segments are formed with organized ganglia,
peripheral nerves, muscles, gut, and other differentiated tissues (de Jong & Seaver, 2016). Although
juveniles are capable of robust posterior regeneration, it is currently unclear to what extent the larvae
are capable of regenerating.

The development of *C. teleta* is indirect and begins as eggs are laid in a maternal brood tube.
Following embryogenesis, non-feeding larvae develop and remain in the brood tube until they are
competent to undergo metamorphosis (Cohen & Pechenik, 1999; Dannenberg & Seaver, 2018). A
previously described staging system uses morphological features to identify 9 distinct stages (Seaver et
al., 2005). Stages 1–3 describe embryonic development, namely fertilization, cleavage, and gastrulation.
Stages 4–9 describe larval development. The transition to a larva in Stage 4 is denoted by the emergence
of trochal ciliated bands that divide the body into three regions along the anterior–posterior axis: the
head, the segmented trunk, and the pygidium. The prototroch separates the head from the trunk and
the telotroch separates the trunk from the pygidium (Fig 1D, E) (Seaver et al., 2005). These two ciliary
bands are composed of long cilia. A third ciliary band, the neurotroch, runs along the ventral midline of
the trunk and is composed of short cilia. Short cilia are also present in the pygidium. Larvae utilize these
ciliary bands to swim in the water column.

As larvae develop (Stages 6 – 9), tissue types differentiate. *C. teleta* has a centralized nervous
system with a ladder-like ventral nerve cord (VNC) positioned along the midline of the trunk (Meyer et
al., 2015; Seaver et al., 2005). The VNC connects with the brain in the anterior of the head (Meyer et al.,
2015). During these stages, musculature is organized segmentally with circular and longitudinal muscles.
Two pigmented eyes are located laterally, just anterior to the prototroch (Meyer et al., 2015; Yamaguchi
et al., 2016). Gut organogenesis is one of the most visibly pronounced set of changes that occurs
between Stage 6 and Stage 9. The foregut is comprised of the mouth (located posterior to the
prototroch, along the ventral midline), buccal cavity, pharynx (developed in Stage 7), and esophagus
(Boyle & Seaver, 2008). In Stages 8–9, the mid- and hindgut differentiate, becoming olive green and
stereotypically coiled by Stage 9 (Fig 1E). In addition, Stage 6 and Stage 9 larvae differ significantly in
shape; Stage 6 larvae are elliptical (Fig 1D) and become elongated over time into a rod-like form by
Stage 9 (Fig 1E). Stage 9 concludes larval development when larvae become competent to undergo
metamorphosis (Seaver et al., 2005).

3.2 Larval amputations

To assess the regenerative potential of *C. teleta* larvae, amputation sites were selected on Stage
6 larvae for both anterior and posterior amputations (Fig 2A). Cut sites for anterior amputations were
located just posterior to the prototroch, removing the head and anterior third of the animal (i.e., brain,
eyes, and prototroch). Cut sites for posterior amputations were located posterior to where the width of
the trunk decreases and slopes to the pygidium, effectively removing the posterior third of the animal,
including the posterior growth zone, telotroch and pygidium.

Successful wound healing was identified as the closure of the wound by contraction of the
severed edges of the body wall resulting in a continuous epithelial layer at the wound site. To assess
wound healing in amputated larvae, we analyzed larvae fixed at 2 hours post amputation (hpa), 3 hpa, 5
hpa, and 6 hpa for an epithelial layer covering the wound site. For anterior amputations, we observed
consistent wound healing 6 hpa (Fig 2B) (n = 54/54). For posterior amputations, we observed wound

healing as early as 2 hpa (n = 3/15); however, in most cases successful wound healing occurred around 6
hpa (Fig 2C) (n = 48/51).

Larvae survived multiple days following amputation (i.e., up to 5 dpa in anteriorly amputated
animals and up to 7 dpa in posteriorly amputated animals). Furthermore, posteriorly amputated larvae
continued to develop on schedule relative to uncut animals (Fig 2D, 2E). For example, chaetae were
present by 1 dpa (comparable to Stage 7), pigmented gut appeared by 2 dpa (comparable to Stage 8),
and the midgut became coiled by 3 dpa (comparable to Stage 9). Anteriorly amputated larvae also
developed chaetae and a pigmented midgut, but the curvature of midgut—a hallmark of Stage 9 larvae
(Boyle & Seaver, 2008; Seaver et al., 2005) was delayed by a day. Another difference between anterior
and posterior amputations was a difference in shape at the cut site: a blunt end was frequently
observed in anterior amputations (Fig 2D) whereas a rounded tapering of tissue was observed distal to
the cut site in posterior amputations (Fig 2E). The tapered shape in posterior amputations is different
from the shape of the wound site in aged-matched individuals (Stage 9) fixed only 6 hpa. In addition,
there was a clear difference in the distance between the posterior edge of the pigmented midgut and
the posterior end of most animals 2 dpa–7 dpa due to the presence of mesodermal tissue (compare Fig
2E, F).

3.3 Cell proliferation

One key feature of regeneration is the birth of new cells in the region of the wound site. We
used age-matched larvae as controls to distinguish between normal growth and an amputation
response. Stage 6 larvae have a characteristic pattern of EdU incorporation with limited dividing cells in
the anterior ectoderm and anterior mesoderm surrounding the mouth and foregut as well as in the
telotroch and pygidium. There are many EdU+ cells in the trunk straddling the midline (Fig 3A). A similar
pattern is observed in Stage 7 larvae (Fig 3B). By Stage 9, EdU+ cells are not ordinarily observed, apart
from a few cells in the gut and posterior growth zone (PGZ) (Fig 3C, 3D).

Anterior amputations show EdU incorporation near the amputation site. The initial pattern of EdU
incorporation in anterior amputations (6 hpa) is similar to the pattern observed in age-matched, uncut
controls (Fig 3E, 3I). However, by 3 dpa, EdU+ cells are localized near the wound site of the amputated
animals in the anterior ectoderm and occasionally in the mesoderm (Fig 3G). EdU+ cells near the wound
site persisted through 5 dpa in anterior ectoderm and mesoderm (Fig 3H). In contrast, there is a notable
lack of EdU+ cells in the corresponding body region of unamputated animals (Fig 3C, 3D). The ratio of
EdU+ cells to nuclei is higher in amputated animals at 3 dpa and 5 dpa at the respective amputation sites
compared to uncut controls (Table 1). There is a significant difference between the number of EdU+ cells
in the anterior section of 3 dpa and 5 dpa larvae when compared to age-matched, uncut controls (One-
way ANOVA test) (Fig 3M).

Posterior amputations also show EdU incorporation localized near the wound site. While the pattern
of EdU incorporated cells at 6 hpa appears similar to that observed in Stage 6 uncut controls (Fig 3I),
there is a progressive shift toward localized expression in amputated larvae. Animals at 1 dpa have
fewer EdU+ cells in the anterior 2/3 of the body (Fig 3J) in comparison with uncut Stage 7 controls. By 3
dpa, EdU incorporation is not detected throughout the anterior 2/3 of the body, and EdU+ cells are
restricted to the area surrounding the cut site (Fig 3K). This localized pattern of EdU incorporation
persists through 5 dpa (Fig 3L). In both cut and uncut larvae, the number of EdU+ cells decrease over

time in the anterior and middle sections of the animal. However, cut larvae have more EdU+ cells in the
posterior region than the uncut controls (Table 2). The number of EdU+ cells in posterior sections of 3
dpa and 5 dpa larvae is significantly higher compared to age matched, uncut controls (One-way ANOVA
test) (Fig 3N). Additionally, there is a higher ratio of localized EdU+ cells/nuclei in posterior amputated
larvae than in anterior amputations. Together, this data shows that there is a marked increase in cell
proliferation at the wound site following both anterior and posterior amputations.

3.4 Expression of the stem cell marker vasa

To further characterize the larval response to amputation, expression of the stem cell marker *vasa*
was analyzed (Dannenberg & Seaver, 2018; Juliano et al., 2010) by in situ hybridization. In uncut larvae,
vasa is expressed in a dynamic pattern from Stage 6 through Stage 9 (Dill & Seaver, 2008). In Stage 6
larvae, there is expression in the foregut, the PGZ, and in the mesoderm of the trunk (Fig 4 A). By Stage
8, expression is restricted to the foregut, the multi-potent cell cluster (MPC), and the PGZ (Fig 4B). This
pattern continues through Stage 9 (with some variability in detection of the MPC) (Fig 4C).

Six hours after anterior amputation, the *vasa* expression pattern in both amputation groups is very
similar to the observed pattern in uncut controls (Fig 4D). By 2 dpa, there is *de novo* expression in the
anterior ectoderm at the amputation site of anteriorly cut larvae (Fig 4E). By 5 dpa, expression at the
wound site is no longer detected, although expression in the MPC and PGZ persists (Fig 4F). In posterior
amputations at 2 hpa, trunk mesodermal expression is continuous from one side of the animal to the
other and wraps around the posterior end of the body (Fig 4G). We interpret this expression along the
posterior face of the animal to be the result of wound healing that pulls the two lateral trunk domains of
expression together. At 2 dpa, *vasa* is expressed in the posterior mesoderm and ectoderm near the
wound site (Fig 4H). This expression near the wound site is also detectable 3 dpa (Fig 4I) and 4 dpa (data
not shown). In summary, we detect *vasa* expression at the wound site 2–3 dpa for both anterior and
posterior amputations.

3.5 Generation of new tissue

With the presence of localized cell proliferation at the wound site (Fig 3G, 3H, 3K, and 3L) and
what appears to the be growth of new tissue (Fig 2E), the tissue surrounding the wound site was
investigated in more detail. To distinguish between pre-exiting tissue and new tissue, the location of the
cut site was determined using the nervous system and musculature as markers. We used a framework
commonly used for identifying the cut site in juvenile worms with anti-acetylated tubulin marking the
nervous system (Fig 1B). An abrupt change in the thickness of the connective nerves, multiple thin
extensions from the VNC to the posterior end of the body, and/or unpaired neuronal fibers were
indications of a cut site (Fig 1B, 5B, and 5B'). An abrupt decrease in the density of muscle fibers also
indicated the location of the cut site (Fig 1C, 5B and 5B"). In many cases, there was a precise
concordance in position between an abrupt change in muscle fiber density and thickness of connective
nerves. Therefore, using these two markers, we could consistently identify the location of the cut site.

The identification of the cut site allowed us to identify and quantify the presence of new tissue
distal to the amputation site in amputated larvae. No evidence of significant growth of tissue in anterior
amputations between 6 hpa and 3 dpa was detected (see Table 3). In contrast, there was significant
growth in posterior amputations between 6 hpa and 5 dpa. Additional measurements of posterior
amputations indicated a significant change in body width (body width at the mouth compared to

immediately posterior to the cut site) between 6 hpa and 5 dpa. This information is consistent with the
presence of new growth in posterior amputations. While the position of the cut site indicated by muscle
and nerves indicated new growth, there was one measurement that did not. Measurements from the
posterior end of the pigmented gut to the posterior of the animal for both the 6 hpa stage 9 and 5 dpa
larvae did not show a significant difference between these two groups.

We next characterized the nervous system and musculature of the tissue distal to the cut site.
Similar to that observed in 6 hpa control larvae (Fig 5C), anterior amputations in 3 dpa larvae lacked
clear evidence of neurites and muscle fibers distal to the wound site (Fig 5 D–D"). In posteriorly
amputated larvae, there were also cases in which there was seemingly no change in the larval nervous
system (Fig 5G; n = 165/395). Cases in which new growth was not observed lend confidence to those
instances when we observed processes distal to the cut site (compare Fig 5F and 5G). In 2 dpa
posteriorly amputated larvae with a discernable cut site and new growth, neuronal projections distal to
the amputation site were observed that were not detected in 6 hpa larvae (Fig 5E, E', F). At 5 dpa, nerve
processes persist in tissue distal to the cut site (Fig 5I, 5I"). These longitudinal nerve processes are
located at the midline and span the length of the new tissue. Additionally, there were circumferential
processes that are similar to the orientation of peripheral nerves (Fig 5I"). These processes are oriented
medial to lateral, extending from the midline. The entire region of the new tissue has lower density of
muscle relative to the preexisting tissue (Fig 5H, 5I'). The fibers that are present are predominately
longitudinal muscle fibers. In many cases, individual longitudinal muscle fibers appeared to be
continuous with fibers in the preexisting tissue that extend through the amputation site (Fig 5I', arrow).
A phalloidin positive fiber with a circular orientation muscle was occasionally observed posterior to the
amputation site. This phalloidin positive fiber is reminiscent of a serotonin positive fiber beneath the
telotroch in uncut animals (Meyer et al., 2015); however, a ciliary band was not seen in amputated
larvae. A similar arrangement of muscle and nerve fibers in the tissue distal to the amputation site
persists through 7 dpa (data not shown). In conclusion, new growth was not observed in anteriorly
amputated larvae, but in posteriorly amputated larvae, new growth along with changes in the nervous
system and musculature was observed.

3.6 Neural specification and differentiation

Next, we sought to determine if the neural and muscle cell types we observed distal to the
amputation site were the result of birth of new cells. We rationalized that the expression of molecular
markers of cell fate specification and differentiation in the tissue distal to the amputation site would
indicate birth of new cells. For the nervous system, gene expression patterns of neural specification
(*ash1*) and neural differentiation markers (*elav*) were analyzed (Meyer & Seaver, 2009). In *C. teleta*, *ash1*
is a marker of neural specification during initial neurogenesis of the brain and VNC (Meyer & Seaver,
2009; Sur et al., 2017). In Stage 6 larvae, *ash1* is expressed in the brain, foregut, PGZ, and the pygidium
(Fig 6A). By Stage 9, *ash1* expression is restricted to the foregut and PGZ (Fig 6B). In anterior
amputations, the pattern observed at 6 hpa for *ash1* expression mirrors that of uncut Stage 6 larvae,
with the notable exception of missing brain expression due to the removal of the head (Fig 6C). In 3 dpa
specimens, *ash1* was expressed in both the anterior ectoderm and pharynx (Fig 6D). Expression in the
ectoderm can be regarded as *de novo ash1* anterior expression since age-matched, uncut controls lack
anterior ectoderm expression (compare panels Fig 6B and 6D, E) (n =32/39). The novel anterior
ectodermal expression of *ash1* at 3 dpa was present in a punctate pattern similar to what is observed

during *C. teleta* brain development. In contrast, expression in the pharynx corresponds with expression
in age matched uncut controls, albeit with fewer cells.

After observing novel anterior neural specification following amputation, the next step of
neurogenesis was analyzed by characterizing the pattern of *elav*, a marker of differentiating neurons.
Between Stages 6 and 9 in uncut controls, *elav* expression is restricted to the brain and the VNC (Fig 6F
& G). In 6 hpa anterior amputations, *elav* is expressed only in the VNC (Fig 6H). This expression pattern
persists through 3 dpa, with no new expression domains observed in amputated animals (n =21/21) (Fig
6I). In the one differing case of the 22 scored, we were not able to definitively score it.

3.7 Amputation induces ectopic MyoD expression

We also examined the expression of *MyoD*, a gene known for a role in muscle specification and
differentiation (Zammit, 2017). In uncut larvae, *MyoD* is expressed in the mesoderm of the posterior
trunk segments at Stage 6 (Fig 7A); this pattern persists through Stage 9 (Fig 7B). There is a
developmental progression from anterior to posterior in the larval trunk, with the more mature
segments having an anterior position (Seaver et al., 2005). Thus, *MyoD* is expressed in the younger and
newly forming segments. Six hours after both anterior and posterior amputations (6 hpa), expression is
restricted to the posterior portion of the trunk (Fig 7 C & E), similar to what is observed in uncut
controls. At 7 dpa and 3 dpa, there is a drastic expansion of *MyoD* mesodermal expression along the
entire length of the trunk in anterior (Fig 7D) and posterior amputees (Fig 7F), respectively. This ectopic
expression is in pre-existing tissue in older segments and includes tissue distant from the wound site in
the case of posterior amputations.

3.8 Rare regeneration of differentiated cell types

Following amputations, a few cell types occasionally regenerated, namely the larval eyes and
cilia. The larvae have two eyes, with each eye consisting of three cells: a pigment cell (Fig 8A), a sensory
cell (Fig 8C), and a support cell (not pictured) (Yamaguchi & Seaver, 2013). The pigment cell is readily
visible by its orange pigment granules and the sensory cell can be visualized by immunolabeling with the
antibody 22C10 (Yamaguchi & Seaver, 2013). Currently, we are unable to assess the presence of the
support cell. In uncut Stage 9 larvae, the pigment cells are located slightly anterior, if not inline, with the
prototroch, in a lateral position (Fig 8A). The sensory cell in uncut Stage 9 larvae (Fig 8C) is located
immediately adjacent and lateral to the pigment cell and has an axonal projection extending from the
cell body to the external surface of the ectodermal epithelium. The pigment and sensory cells were
scored as independent characters. Following head amputation, the few regenerated pigment cells
observed were variable in location and number (n = 28/420; 3-5 dpa). Additionally, as seen in Fig 8B,
pigment cells in amputated larvae were ectopically positioned and in proximity to tissues normally not
observed in uncut controls (i.e., midgut). Regenerated sensory cells were also observed in amputated
larvae and also exhibited variability in location (n= 8/220, 3 –7 dpa). Axon projections in regenerated
sensory cells extended to the outer ectodermal surface of the animals (Fig 8D). There was a higher
percentage of cases with regenerated pigment cells (7%) compared with sensory cells (4%) in anterior
amputations (Fig 8G). Summation of the total number of pigment cells and sensory cells indicates that at
least one cell of the eye regenerated in 6% of cases amputated (n = 36/640).

We were surprised by the appearance of new larval eye cells in *C. teleta* larvae since
regeneration of anterior structures is not reported in adults (Yamaguchi & Seaver, 2013). Therefore, we

conducted a similar experiment with juveniles. Four-week-old juveniles were amputated immediately
anterior to the mouth, removing the eye cells. After screening for successful wound healing and visual
inspection that pigment cells were removed, a portion of the amputated juveniles were fixed at 4 hpa
and used to confirm the removal of the sensory cells (n =21/21). No pigment cells were observed in
amputated juveniles at 7 dpa (n = 0/43). Using 22C10 antibody labelling, a small proportion of cases with
eye sensory cells were observed (Fig 8E) (n = 6/43; 14%). Regenerated sensory cells were ectopically
located compared to their position in uncut controls (Fig 8E, 8F). Five of the six juveniles with
regenerated sensory cells had a small portion of brain present in the anterior of the animal; only one
case lacking a brain had a sensory cell. In summary, pigment cells did not regenerate in juveniles,
although sensory cells were detected in a small fraction of cases.

There are four external ciliated structures in *C. teleta* larvae: prototroch, neurotroch, telotroch,
and pygidium. The prototroch and telotroch are composed of relatively long cilia (prototroch cilia length
~25-30 microns and telotroch cilia length ~20-28 microns), and the neurotroch and pygidium have short
cilia (neurotroch cilia length ~ 8-10 microns, pygidium length ~5-7 microns) (Fig 9A). At 6 hpa, larvae
lacked long cilia at the cut site (Fig 9 B & C), due to amputation of the prototroch and telotroch in
anterior and posterior amputations, respectively. In rare cases (n = 12/161), short cilia were observed at
the anterior end of anteriorly cut animals at 5 dpa, distal to the amputation site (approximately 8
microns in length) (Fig 9E). Long cilia were not observed following anterior amputations (n = 0/161).
While rare (n = 22/370), both long (~25 microns) and short (~9 microns) cilia were observed following
posterior amputations between 3–7 dpa (Fig 9F).

3.9 Metamorphosis following amputation

C. teleta is typically competent to undergo metamorphosis into juveniles as Stage 9 larvae.
Metamorphosis is rapidly induced by exposure to vitamin B supplements dissolved in salt water (Burns
et al., 2014). To test whether amputated larvae are capable of successfully undergoing metamorphosis,
uncut, anteriorly amputated, and posteriorly amputated larvae were exposed to a solution containing
vitamin B. Animals were screened for stereotypical signs of successful settlement and metamorphosis
following the definition of Cohen and Pechenik (1999). These changes include a loss of swimming
behavior and initiation of burrowing behavior, loss of both ciliary bands (telotroch and prototroch), and
the appearance of an elongated, worm-like body shape. Uncut, uninduced control larvae swam with
their prototroch and telotroch bands intact for the duration of the experiment (Fig. 10A) (2 hr, n =
532/534). In contrast, virtually all larvae in the induced, unamputated control group successfully
underwent metamorphosis within two hours of exposure, settling on the bottom of the dish, dropping
their cilia, adopting burrowing behavior, and extending in length along their anterior-posterior axis (Fig
10B) (n = 203/207). Both sets of amputated larvae followed a similar pattern to what was observed for
unamputated larvae. Specifically, one day after anterior amputation, larvae either continued to swim in
the water column in the absence of a metamorphic cue (Fig. 10C) (n=65/66) or settled to the bottom of
the dish and underwent metamorphosis within two hours of exposure to a metamorphic cue (Fig. 10D)
(n=135/136). Similarly, individuals whose posterior ends had been removed retained their prototrochal
bands and continued to swim in the absence of a metamorphic cue (Fig. 10E, arrow) (n=65/65), yet,
underwent rapid settlement and metamorphosis within two hours of exposure (Fig. 10F) (n=143/144).
Without an induction cue, amputated laravae continue to swim in the water column and retain their
ciliary bands for several days, demonstrating that amputation itself does not induce metamorphosis (Fig

9E, F). Together, these results demonstrate *C. teleta* larvae can detect a metamorphic cue and
successfully execute metamorphosis in the absence of either anterior or posterior structures.

4. Discussion

4.1 Limited regeneration in C. teleta larvae

This study demonstrates that the initial events following amputation appear to be quite similar
between *C. teleta* larvae and juveniles. Following both anterior and posterior amputation, larvae wound
heal in a similar time frame as previously reported for posterior amputation of juveniles (i.e., within 6
hpa) (de Jong & Seaver, 2016). Following wound healing, EdU+ cells appear localized at the wound site
at 2 dpa in juvenile posterior regeneration, and the number of EdU+ cells increases over time (de Jong &
Seaver, 2016). EdU incorporation in anterior-facing wound sites in juveniles of *C. teleta* has not been
formally characterized. In both anterior and posterior larval amputations, EdU is incorporated near the
wound site 3 dpa and this persists through 5 dpa, comparable to the pattern of EdU incorporation at the
wound site in regenerating juveniles. There is a difference in the number of EdU+ cells between anterior
and posterior amputations in larvae, with posterior amputations exhibiting more EdU+ cells. One
difference in cell proliferation patterns between juveniles and larvae is a change in position of the
localized EdU+ cells. During posterior regeneration in juveniles, localization of EdU+ cells progressively
shifts anteriorly to a subterminal position, a shift that corresponds with the re-establishment of the
juvenile PGZ. We did not observe such a shift in larvae. It is interesting to note that in the annelid
Paranais litoralis, BrdU is incorporated near the anterior wound site even though these animals have
lost the ability to regenerate their heads (Bely and Sikes, 2010). This finding, along with our results in *C.*
teleta larvae, emphasize that a block in regeneration can occur subsequent to the initiation of cell
division at the wound site, and cell division at the wound site is not a reliable predictor of successful
regeneration.

Interestingly, expression of the stem cell marker *vasa* at the wound site was observed in both
sets of amputated larvae. In successful posteriorly regenerating juveniles, *vasa* is expressed locally at the
wound site at 2 dpa and persists for several days (de Jong & Seaver, 2017). In larvae, *vasa* expression has
a different duration between anterior and posterior amputations. In anterior amputations *vasa*
expression is transient; it is detectable at 2 dpa but not 5 dpa. Conversely, localized *vasa* expression is
observed at 2 dpa and persists through 5 dpa in posterior amputations. We interpret the localized
expression of *vasa* at the posterior wound site as either the result of *de novo* expression or expression
maintained from an earlier stage (i.e., 6 hpa). The spatial and temporal patterns of EdU incorporation
and *vasa* expression are very similar in posterior amputations. In contrast, near the wound site in
anterior amputations, *vasa* expression is of a shorter duration than EdU incorporation. We hypothesize
that this difference between anterior and posterior amputations is related to the new tissue growth in
posterior amputations yet absent in anterior amputations.

Additional features of juvenile posterior regeneration are the presence of neural projections
into the wound site and muscle reorganization distal to the amputation site. In juveniles, this
phenomenon is observed approximately at 2 dpa (de Jong & Seaver, 2016). In larval posterior
amputations, neurites were present distal to the cut site by 2 dpa. However, these neurites were not
observed in larval anterior amputations. With respect to muscle fiber organization, there is an abrupt
transition between the presence and absence of muscle fibers distal to the cut site in both juveniles and
larval posterior amputations that was not observed in larval anterior amputations. We were unable to

determine whether fibers present distal to the amputation site in posterior larval amputations resulted
from birth of new neural or muscle cells or from the extension of pre-existing cells.

443   To understand the development of the tissue distal to the cut site in larvae, we characterized
the specification and differentiation of cell types (i.e., neural cell types and cilia) in amputated larvae.
The timing of neural specification or differentiation in juveniles has not been investigated in detail, but
differentiated tissues are present by 7 dpa (de Jong & Seaver, 2016). Analysis of *de novo ash1* expression
indicates the specification of neural fates in larval anterior amputations. Surprisingly, subsequent
differentiation of neural tissues was not detectable, as illustrated by the absence of *elav* expression in
larval amputations. Anterior amputation of larvae results in incomplete tissue and structure
regeneration—an observation consistent with the response in juveniles to anterior amputations.
Despite a general lack of anterior structural differentiation in amputated larvae, we surprisingly
observed rare cases in which differentiated eye cells and cilia reappeared within 3–7 dpa. Additionally,
sensory eye cells (but not pigment cells) regenerated in juveniles at a higher proportion than in larvae.
While rare, we are confident that these cells are regenerated differentiated eye cells in amputated
larvae and juveniles. In summary, the beginning stages of regeneration occur in amputated larvae but
there is a lack of cell or tissue differentiation. These data indicate that *C. teleta* larvae exhibit limited
regeneration abilities relative to the successful posterior regeneration observed in juvenile and adult
stages. Additionally, detection of initial stages of regeneration in larvae suggests that *C. teleta* has a
gradual transition in regenerative potential, rather than a stark change in potential that corresponds
with the onset of metamorphosis*.*

4.2 Ectopic expression in preexisting tissue

MyoD functions in muscle development and its expression in muscle progenitor cells is an indication
of myogenesis (Zammit, 2017) (Rawls & Olson, 1997). In response to amputation, *MyoD* becomes
ectopically expressed across the length of the larval body in *C. teleta*. This ectopic expression may result
in an increase in muscle cell number to compensate for a decrease in body length following amputation,
although we did not observe an obvious increase in the number of muscle fibers 3 – 5 dpa. Alternatively,
ectopic expression of *MyoD* may be an injury-specific response that leads to cell dedifferentiation and
return to a precursor state. Such a precursor cell population may contribute to regenerating tissue. It is
notable that the induction of ectopic *MyoD* expression occurs in the absence of a complete regeneration
response. Regardless, future functional work is needed to better understand the implications of this
novel *MyoD* expression in amputated larvae. Ectopic gene expression in response to amputation has
previously been documented in *C. teleta* and other annelids (de Jong & Seaver, 2016; Kozin et al., 2017;
Pfeifer et al., 2012; Takeo et al., 2008). However, in most cases, these are positional identity genes and
their ectopic expression likely reflects repatterning of the body fragment in response to amputation.

4.3 Metamorphosis following amputation

476   Amputated larvae can undergo successful metamorphosis. We were surprised to observe
metamorphosis following anterior amputations. In other spiralians, it has been hypothesized that the
brain and other anterior sensory organs function in the detection and response to metamorphic cues
(Chartier et al., n.d., 2018; Conzelmann et al., 2013; Hadfield, 2010; Hadfield et al., 2000). In *C. teleta*, it
has been hypothesized that chemo-sensitive ciliated cells in the head ectoderm relay metamorphosis
signaling directly to the cerebral ganglia (Biggers & Laufer, 1999). Therefore, we expected that by
removing the head, anterior amputees would not undergo metamorphosis. Our results demonstrate

that the brain is not required for detecting the metamorphic cue, nor for undergoing metamorphosis in
C. teleta. Without the sensory structures to detect the metamorphic cue, we would expect cilia to be
retained. A recent study in the polychaete *Hydroides elegans* similarly demonstrated that its apical
sensory neurons are not necessary for successful metamorphosis (Nedved et al., 2021). Furthermore,
our results are consistent with a previous study hypothesizing that the inductive metamorphosis signal is
detected by sensory neurons that innervate the dorsal pharyngeal pad of the pharynx (Biggers et al.,
2012). In our experiments, the pharynx remained intact in both anterior and posterior amputated
larvae. Successful metamorphosis from amputated larvae supports the hypothesis that detection of the
metamorphic cue in *C. teleta* can occur in the pharynx.

4.4 Larval regeneration in marine invertebrates

Larval regeneration studies are overshadowed by numerous studies of adult and juvenile
regeneration. Of the taxa where larval regeneration is studied, there is a disproportionate emphasis on
echinoderms (C. Carnevali, 2006; G. Cary & Hinman, 2017; Oulhen et al., 2016; M. C. L. Vickery et al.,
2001; M. S. Vickery & McClintock, 1998; Minako S. Vickery et al., 2002). Larval regeneration has been
documented in all classes of echinoderms examined (G. A. Cary et al., 2019; M. C. L. Vickery et al., 2001).
Regenerative potential can vary significantly across life stages and age. In some animals, larval tissues
and organs have similar regenerative abilities as adults. Asteroids and echinoid larvae have
demonstrated the ability to regenerate completely, regardless of developmental stage (C. Carnevali,
2006; Minako S. Vickery et al., 2002). In others, like in crinoids, larvae display a limited ability to
regenerate. Echinoderms in which larvae have a reduced regenerative ability relative to that of the adult
are described as having "attempted" or "limited" regeneration. When regeneration has been recorded
in crinoid larvae, it has been described as partial regeneration of body parts in both sessile and
swimming larval stages (C. Carnevali, 2006). In contrast, adults are capable of robust structural
regeneration even when reduced to 1/5 of the original body size (C. Carnevali, 2006). In the crinoid
Antedon rosacea, amputated early-stage larvae differ in their regeneration ability compared to late-
stage. For early stage amputated larvae, both anterior and posterior halves demonstrate a reduced
regenerative capacity. In late-stage larvae, the anterior portions developed typical skeletons, while
posterior halves lack stalk development, but have numerous duplicated organs and vestibules (M. C. L.
Vickery et al., 2001). Similar to *Antedon rosacea* early-stage larvae*,* amputation of 2-day old *Holothuria*
poli into right and left halves resulted in "attempted" regeneration (M. C. L. Vickery et al., 2001). In
amputated *Holothuria*, processes similar to development were observed in an amputated coelom in the
absence of tissue replacement (M. C. L. Vickery et al., 2001).

These cases of limited or attempted regeneration in echinoderms have remarkable parallels with
what we documented in *C. teleta* larvae relative to features characteristic of successful regeneration in
C. teleta juveniles and adults. Amputated larvae (both anterior and posterior amputations) resulted in
completed wound healing, localized cell proliferation, and innervation. However, in most cases, we did
not observe neural and muscle differentiation, nor the reformation of ciliary bands or other structures.
While larvae appear to have the rare capacity to regenerate cilia and eyes, we conclude that complete
regeneration is not the typical response to amputation. The initiation of the regeneration response in
larvae, but a lack of complete replacement of lost structures is consistent with our hypothesis that
larvae have limited regeneration potential. Furthermore, our data do not support an abrupt onset of
regenerative potential coincident with metamorphosis in *C. teleta*, although differences between larval

and juvenile regenerative potential demonstrate that there is an increase in regenerative potential
following metamorphosis.

Outside of echinoderms, there are far fewer examples of larval regeneration studies. In the
hemichordate *Ptychodera flava*, larvae display a more restricted regeneration ability than adults.
Specifically, ventral regions of larvae regenerate better than dorsal regions and posterior segments
regenerate better than anterior segments (Luttrell et al., 2018). Another example comes from the
nemertean, *Maculaura alaskensis*, which as an adult is capable of posterior regeneration (Hiebert &
Maslakova, 2015). During its life cycle, *Maculaura alaskensis* forms a feeding larva that eventually
undergoes a catastrophic metamorphosis. Amputations in larvae revealed that younger larvae were five
times more likely to regenerate than older larvae (Moss, 2017). This study substantially contributes to
addressing that have been, until now, predominately restricted to deuterostomes.

4.5 Future directions

This study demonstrates differences in regenerative ability across developmental stages of
Capitella teleta. Future studies could investigate differences in physiological, cellular, and molecular
responses to amputation between larvae and juveniles. It has been demonstrated that the extent of
regeneration in *C. teleta* juveniles and other annelids can fluctuate depending on nutrition intake (de
Jong & Seaver, 2016; Zattara & Bely, 2013). Because *C. teleta* larvae are non-feeding, limited energy
resources may contribute to their regeneration limitations. This nutrition hypothesis could be tested by
two separate approaches. In the first, cells with high yolk content can be targeted for deletion in early-
stage embryos to reduce maternally contributed nutrition. Previous work demonstrated that such
deletions result in morphologically normal, albeit smaller larvae (Pernet et al., 2012). These larvae would
have a reduced nutritional supply. After performing amputations on the smaller larvae, there may be a
further reduction in regenerative potential to what was observed in this study. The second approach
would involve investigating the regeneration abilities in a different annelid with a feeding larval form. A
feeding larva with more nutritional resources may exhibit full regenerative abilities following
amputation. Additionally, an annelid with a longer larval period would have more time to complete
regeneration following amputation during the early larval period, while also continually feeding.

Timing of developmental maturation may also limit regeneration ability and explain differences
between larval and juvenile regenerative abilities in *C. teleta*. An example of one such developmental
constraint may be in the maturity of stem cells. A previous study proposed that stem cells travel from
the MPC cluster to the wound site during juvenile regeneration (de Jong & Seaver, 2017). MPC cells in
larvae may be immature, thereby preventing these stem cells from positively contributing to successful
regeneration. Specifically, nascent stem cells may not be capable of self-renewal or migration (Juliano
et al., 2010), or of contributing to the blastema. The number of cells in the MPC cluster does not
increase during larval development (de Jong & Seaver, 2017), suggesting these cells may be quiescent
during this time. A heterotopic transplant of a mature MPC cluster from juveniles into larvae would test
this hypothesis. We would expect to see an increased regenerative ability in larvae following
transplantation.

It is yet to be determined whether the amputations performed during larval stages have lasting
effects on survival and growth following metamorphosis. It remains to be seen whether juveniles
induced from posteriorly amputated larvae redevelop posterior growth zones, can generate additional
segments or are capable of posterior regeneration. Future studies could examine the regeneration

potential of juveniles that result from amputated larvae. Additionally, it could be interesting to
determine if the juveniles induced from anterior amputated larvae are capable of posterior regeneration
in the absence of a brain.

Previous studies have identified a wide range of regeneration abilities in adult annelids. Adult
annelids also display differences in regenerative ability with respect to cut site location. For example, in
Chaetopterous pergamentaceus (Cuvier et al., 1829) (formally *variopedatus*), posterior regeneration
occurs following amputation at any segment; in contrast, anterior regeneration does not occur following
amputations posterior to segment 15 (Berrill, 1928) (Cuvier et al., 1829). Regenerative success appears
to be influenced by species-specific differences, developmental timing, nutritional status, amputation
location, or a combination of these. Future studies can characterize the regeneration ability of other
annelid larvae, particularly those whose adults have different regenerative potential than that of *C.
teleta*. There may a correlation between adult and larval regenerative potential within individual
species. For example, there may be anterior regenerative ability in larvae in a species capable of anterior
regeneration as an adult. We are intrigued to learn whether larval regenerative ability is as common as
adult regenerative ability in annelids. Comparison of *C. teleta* regeneration abilities to other annelids
might lead to a better understanding of the role that development has on regeneration.

2. **Materials and Methods**

2.1 Animal Care

As described in previous work, a colony of *Capitella teleta* was maintained in laboratory at 19°C
in glass finger bowls filled with mud and filtered sea water (FSW). Using forceps, larvae were dissected
from the brood tubes of healthy females. Broods were maintained individually in FSW until the desired
age for amputation or treatment.

2.2 Amputations

Amputations were conducted on a 60 mm petri dish lid covered in a methyl cellulose solution
(0.3 g methyl cellulose to 15 ml 1:1 solution of 0.37 M $MgCl_2$: FSW. Glass pipettes pulled into thin,
flexible, sharp tips capable of slicing were used as amputation tools. Individual broods were isolated and
selected when identified as containing Stage 6 larvae following a standard staging system (Seaver et al.,
2005). Stage 6 was characterized by yolk in the anterior portion of the larvae, swimming behavior with a
positive phototactic response, and lack of chaetae.

Larvae were incubated in 1:1 $MgCl_2$: FSW for 10 minutes. Individual larvae were transferred to
an amputation stage (a platform of black dissecting wax (American Educational Products, Fort Collins,
CO, USA in the lid of a 35 mm plastic dish) in a 6 % methyl cellulose in a 1:1 solution of 0.37 M $MgCl_2$:
FSW, adding additional methyl cellulose as necessary to achieve immobilization. Larvae were amputated
with a pulled glass micro-needle in a continuous, downward motion. The positions of amputations were
~1/3 of the body length for anterior amputations and ~2/3 of the body length for posterior amputations.
With each amputation, complete separation of body sections was visually noted. Amputated larvae
were transferred to a 60 mm petri dish and incubated in 1:1 $MgCl_2$: FSW for a minimum of 1 hour.

Larvae were then washed into FSW containing 60 µg/ml penicillin and 50 µg/ml streptomycin (Sigma-
Aldrich Co.)(P/S) and incubated at 19°Celsius until the designated time point. P/S FSW was replaced
daily, as necessary, while also monitoring for vitality and swimming behavior.

For eye amputations, Stage 6 larvae were amputated posterior to the prototroch as described
above, and each individual was visually inspected for correct cut site and complete separation of the
two tissue fragments. Two-week-old juveniles were prepared for amputation by incubation in 1:1 $MgCl_2$:
FSW for 10 minutes. Individual worms were transferred to a wax amputation stage in a drop of 1:1
$MgCl_2$: FSW. The worms were amputated immediately anterior to the mouth, removing the eyes and
brain. Individual worms were visually inspected for the correct cut site and successful separation before
being transferred to a 60 mm petri dish and incubated in 1:1 $MgCl_2$: FSW for a minimum of 1 hour. The
juveniles were then washed into FSW and moved to a 60 mm petri dish containing mud and FSW for 1
week. For analysis, juvenile worms were sifted from the mud and placed into a petri dish of FSW. Worms
were fixed with 4% paraformaldehyde (PFA) in FSW for 30 min before being washed into phosphate-
buffered saline (PBS). Fixed juveniles were scored for pigmented eye cells before being processed for
antibody labeling.

2.3 Cell Proliferation Assay

Sibling larvae dissected from a single brood tube were incubated in EdU (Click-iT EdU Alexa Fluor
488 imaging kit - Invitrogen) diluted in FSW to a final concentration of 3 uM at room temperature for 30
minutes. Larvae were then washed into a 1:1 $MgCl_2$: FSW solution for 10 minutes and then fixed in 4%
PFA for 30 minutes. Afterward, they were rinsed into PBS and incubated for 20 minutes at room
temperature in PBT (PBS + 0.05% Triton-X-100) with rocking. The specimens were incubated in EdU
reaction buffer for 30 minutes, rocking at room temperature. Finally, they were washed into PBS and
stored at 4 degrees celsius until needed.

2.4 EdU Quantification

Quantification of EdU and DAPI stained cells was conducted by importing Z-stacks generated
from confocal microscopy (see Microscopy) into ImageJ software. Quantification of anterior and
posterior amputations were analyzed using the same methodology. Using DAPI and DIC channels to
establish boundaries, each image was cropped around the perimeter of the animal. The channels were
split and treated separately to optimize cell counting. The threshold for each channel was adjusted
manually to minimize background and enhance distinction of individual cells. Afterward, a mask and
watershed application were applied to each channel. The resulting image was divided into three equal
sections along the anterior-posterior axis using the Montage to Stack application. The modified images
were compiled into 20 slice projections (20 µm total) before being counted automatically by the Analyze
Particles in the Analyze menu. Each 20 µm Z-stack was counted, with the results automatically
summarized via section. Manual inspection eliminated any cells that were counted twice by the
software. Results were manually compiled by channel and section for each animal, as well as for each
time point. A ratio of EdU stained cells to DAPI stained cells was calculated for each time point from the
ImageJ cell count summaries. Averages of each section (anterior, middle, and posterior) for each time

point (uncut controls, 6 hpa, 3dpa, and 5dpa) was calculated separately. Each body section of uncut
controls was compared with corresponding body sections of amputated animals. One-way weighted
ANOVA statistical tests were performed using VassarStats to test for significance between the control
and amputation group of interest.

2.5 Antibody labeling and phalloidin staining

Amputated larvae or juveniles were incubated in a 1:1 $MgCl_2$: FSW solution for 10 minutes and fixed in
4% paraformaldehyde in FSW for 30 minutes. Animals were washed into PBS and then a PBS+ Triton
solution (0.1% Triton-X-100 for larvae and 0.2% Triton-X-100 for juveniles), rocking at room temperature
for 45–60 min. For antibody labeling, the anti-acetylated tubulin antibody (goat anti-mouse, Sigma-
Aldrich Co.) was diluted 1:400 in block solution (10% heated treated goat serum in PBT), and incubated
at room temperature for 2 hours or overnight at 4°C. Following multiple washes in PBT (4 times in PBT
for 30 minutes each), a secondary antibody (goat anti-mouse red 594, Invitrogen) was diluted 1:400 in
block solution and incubated 2–4 hours rocking at room temperature or overnight at 4°C. The 22C10
antibody (Goat anti-mouse, Developmental Studies Hybridoma bank) (22C10 was deposited to the DSHB
by Benzer, S. / Colley, N. (DSHB Hybridoma Product 22C10)) was diluted 1:200 in block solution, and
incubated overnight at 4°C. Following multiple washes in PBT over two hours, the secondary antibody
(goat anti-mouse green 488 - Invitrogen) was diluted 1:400 in block solution and incubated overnight at
4°C. Following incubation in secondary antibody, larvae were rinsed 2–3 times in PBT and then washed 4
times in PBT for 30 minutes each wash. For phalloidin staining, specimens were transferred to a 3 well
glass depression plate and incubated in a 1:400 dilution of phalloidin (Alexa Fluor 488 phalloidin,
Invitrogen) in PBT. Specimens were incubated on a rocker at room temperature for 45–60 min before
being washed twice in PBT for 30 minutes each wash. Animals were cleared in 80% glycerol in PBS with
Hoechst overnight prior to imaging.

2.6 Whole mount in situ hybridization

Larvae collected at different time points following amputation were incubated in a 1:1 $MgCl_2$:
FSW solution for 10 minutes and then fixed in 4% paraformaldehyde overnight. Fixative was removed by
multiple rinses in PBS and then larvae were gradually washed into 100% methanol and stored at -20°C.
Following a previously described protocol (de Jong & Seaver, 2017), whole mount in situ hybridization
was performed. DIG-labeled RNA probes were generated with the MegaScript SP6 or T7 Transcription kit
(Invitrogen). The following working concentrations and probes were used: *ash1* with 3ng/ul, *MyoD* with
0.5ng/ul, *vasa* with 1ng/ul, and *elav* with 2ng/ul. Animals were hybridized for ~24 – 48 hours at 65°C.
Each RNA probe was visualized by color reaction with NBT/BCIP and visually monitored. For each
sample, uncut control larvae were developed alongside experimental, amputated larvae. The reaction
was terminated following washes of PBT followed by a final fixation in 4% PFA in FSW for 30 min. Nuclei
were stained with 0.1 ug/ml Hoescht in PBS and 80% glycerol before individuals were placed on
microscope slides for analysis and imaging.

2.7 Tissue and Cilia Measurements

To measure tissue formation in amputated larvae, confocal images were analyzed using ImageJ
software. Multiple measurements were made on each specimen for the applicable traits. Nervous
system measurements were measured from the cut site to the ectoderm distal to the cut site. The
muscle measurements were taken from the cut site to the ectoderm distal to the cut site. Changes in
body width were calculated by making two measurements per specimen; the first measurement was
taken at the axial level of the mouth, and the second was taken just posterior to the cut site. Gut
measurements were made from the posterior end of the pigmented gut to the edge of the posterior
ectoderm. For anterior amputations the muscle and nervous system were measured in 6 hpa (n = 8) and
3 dpa (n = 10) larvae. For posterior amputations, the muscle, nervous system, and changes in body width
were measured for 6 dpa (n = 7) and 5 dpa (n = 10). Additionally, the gut was used as a measurement
landmark for 5 dpa and Stage 9 larvae at 6 hpa (n = 10). One-way weighted ANOVA statistical tests were
performed using *goodcalculators* to test for significance between the control and amputation group of
interest.

The cilia of larvae were measured using ImageJ software on confocal images. Using the channel
in which the cilia are labelled with anti-acetylated tubulin, we traced individual labeled cilia of the
prototroch, neurotroch, telotroch, and pygidium. Lines resulting from the trace were then measured
using the 'measure' tool found under 'Analyze.' At least three cilia per ciliary band were measured in a
representative specimen of a given experimental group.

2.8 Metamorphosis assays

A single multivitamin (Centrum Men Multivitamin) was dissolved in 200 ml FSW. Larvae were
placed in a 30mm petri dish in FSW. Most of the FSW was removed from the dish and replaced with 3 ml
of the multi-vitamin solution. Animals were left in the vitamin solution for 20 - 30 minutes. Animals were
then washed rapidly with FSW (2 ml/wash), a minimum of three times. Animals were individually scored
for successful settlement and metamorphosis. Successful settlement and metamorphosis were defined
as cessation of swimming behavior and initiation of burrowing, loss of the prototroch and/or telotroch
ciliary bands and elongation of the body.

2.9 Microscopy and Imaging

Larvae were imaged with a Zeiss 710 confocal microscope using 4x bidirectional and
unidirectional settings for green and red channels. Z-stacks were compiled using ImageJ software
(Schindelin et al., 2012). Differential interference contrast microscopy images were captured on a Zeiss
AxioSkop II motplus compound microscope (Zeiss) coupled with a SPOT FLEX digital camera (Diagnostic
Instruments Inc). Multiple focal planes were merged for DIC images using Helicon Focus software
(Helicon Focus 7). Images were processed for brightness, cropping, and contrast with Illustrator (Adobe
Illustrator 2020) or or Photoshop (Adobe Photoshop 2020). Figures were composed using Adobe
Illustrator (Adobe Illustrator 2020).

Acknowledgements

We thank Brent Foster, Lauren Kunselman, and Katie Feerst for critical reading of the manuscript.

Author Contributions

Conceived and designed the experiments: AAB ECS. Performed the experiments: AAB. Analyzed the
data: AAB ECS. Contributed reagents/materials/analysis tools: ECS AAB. Wrote the paper: AAB ECS.

References

Alvarado, A. S. (2000). Regeneration in the metazoans: why does it happen? *BioEssays*.
https://doi.org/10.1002/(SICI)1521-1878(200006)22:6

Bely, A. E. (2006). Distribution of segment regeneration ability in the Annelida. *Integrative and*
Comparative Biology, *46*(4), 508–518. https://doi.org/10.1093/icb/icj051

Bely, Alexandra E, & Sikes, J. M. (2010). Latent regeneration abilities persist following recent
evolutionary loss in asexual annelids. *Proceedings of the National Academy of Sciences of the*
United States of America, *107*(4), 1464–1469. https://doi.org/10.1073/pnas.0907931107

Berrill, N. J. (1928). *Regeneration in the polychret Choetopterus Ohcetopterus*. 151–158.

Biggers, W. J., & Laufer, H. (1999). Settlement and Metamorphosis of Capitella Larvae Induced by
Juvenile Hormone-Active Compounds Is Mediated by Protein Kinase C and Ion Channels.
Https://Doi.Org/10.2307/1542564, *196*(2), 187–198. https://doi.org/10.2307/1542564

Biggers, W. J., Pires, A., Pechenik, J. A., Johns, E., Patel, P., Polson, T., & Polson, J. (2012). Inhibitors of
nitric oxide synthase induce larval settlement and metamorphosis of the polychaete annelid
Capitella teleta. *Invertebrate Reproduction and Development*, *56*(1), 1–13.
https://doi.org/10.1080/07924259.2011.588006

Boyle, M. J., & Seaver, E. C. (2008). Developmental expression of foxA and gata genes during gut
formation in the polychaete annelid, Capitella sp. I. *Evolution & Development*, *10*(1), 89–105.
https://doi.org/10.1111/J.1525-142X.2007.00216.X

Brockes, J. P. (1997). Amphibian Limb Regeneration: Rebuilding a Complex Structure. *Science*, *276*(5309),
81–87. https://doi.org/10.1126/SCIENCE.276.5309.81

Burns, R. T., Pechenik, J. A., Biggers, W. J., Scavo, G., & Lehman, C. (2014). The b vitamins nicotinamide
(b>3) and riboflavin (b2) stimulate metamorphosis in larvae of the deposit- Feeding polychaete
capitella teleta: Implications for a sensory ligand-gated ion channel. *PLoS ONE*, *9*(11).
https://doi.org/10.1371/journal.pone. 0109535

Byrnes, E. F. (1904). Regeneration of the anterior limbs in the tadpoles of frogs. *Archiv Für*
Entwicklungsmechanik Der Organismen, *18*(2), 171–177. https://doi.org/10.1007/BF02163652

Carnevali, C. (2006). Regeneration in Echinoderms: repair, regrowth, cloning. In *ISJ* (Vol. 3).

Carnevali, M. D. C., Lucca, E., & Bonasoro, F. (1993). Mechanisms of arm regeneration in the feather star
Antedon mediterranea: Healing of wound and early stages of development. *Journal of*
Experimental Zoology, *267*(3), 299–317. https://doi.org/10.1002/JEZ.1402670308

Cary, G. A., Wolff, A., Zueva, O., Pattinato, J., & Hinman, V. F. (2019). Analysis of sea star larval
regeneration reveals conserved processes of whole-body regeneration across the metazoa. *BMC*
Biology, *17*(1), 16. https://doi.org/10.1186/s12915-019-0633-9

Cary, G., & Hinman, V. (2017). Conserved processes of metazoan whole-body regeneration identified in
sea star larvae. *BioRxiv*, 118232. https://doi.org/10.1101/118232

Chartier, T. F., Arendt, D., Deschamps, J., Dürichen, W., & Jékely, G. (n.d.). *Whole-head recording of*
chemosensory activity in the marine annelid Platynereis dumerilii.
https://doi.org/10.1098/rsob.180139

Chartier, T. F., Deschamps, J., Dürichen, W., Jékely, G., & Arendt, D. (2018). Whole-head recording of chemosensory activity in the marine annelid Platynereis dumerilii. *Open Biology*, *8*(10). https://doi.org/10.1098/RSOB.180139

Cohen, R. A., & Pechenik, J. A. (1999). Relationship between sediment organic content, metamorphosis, and postlarval performance in the deposit-feeding polychaete Capitella sp. *Journal of Experimental Marine Biology and Ecology*, *240*, 1–18.

Conzelmann, M., Williams, E. A., Tunaru, S., Randel, N., Shahidi, R., Asadulina, A., Berger, J., Offermanns, S., & Jékely, G. (2013). Conserved MIP receptor-ligand pair regulates Platynereis larval settlement. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(20), 8224–8229. https://doi.org/10.1073/PNAS.1220285110/-/DCSUPPLEMENTAL

Cuvier, G., Cuvier, G., & Latreille, P. A. (1829). Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. In *Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée*. Chez Déterville. https://doi.org/10.5962/bhl.title.49223

Dannenberg, L. C., & Seaver, E. C. (2018). Regeneration of the germline in the annelid Capitella teleta. *Developmental Biology*, *440*(2), 74–87. https://doi.org/10.1016/j.ydbio.2018.05.004

de Jong, D. M., & Seaver, E. C. (2016). A Stable Thoracic Hox Code and Epimorphosis Characterize Posterior Regeneration in Capitella teleta. *PLOS ONE*, *11*(2), e0149724. https://doi.org/10.1371/journal.pone.0149724

de Jong, D. M., & Seaver, E. C. (2017). Investigation into the cellular origins of posterior regeneration in the annelid Capitella teleta. *Regeneration (Oxford, England)*, *5*(1), 61–77. https://doi.org/10.1002/reg2.94

Dill, K. K., & Seaver, E. C. (2008). Vasa and nanos are coexpressed in somatic and germ line tissue from early embryonic cleavage stages through adulthood in the polychaete Capitella sp. I. *Development Genes and Evolution*, *218*(9), 453–463. https://doi.org/10.1007/s00427-008-0236-x

Gaire, J., Varholick, J. A., Rana, S., Sunshine, M. D., Doré, S., Barbazuk, W. B., Fuller, D. D., Maden, M., & Simmons, C. S. (2021). Spiny mouse (Acomys): an emerging research organism for regenerative medicine with applications beyond the skin. *Npj Regenerative Medicine 2021 6:1*, *6*(1), 1–6. https://doi.org/10.1038/s41536-020-00111-1

Giani, V. C., Yamaguchi, E., Boyle, M. J., & Seaver, E. C. (2011). Somatic and germline expression of piwi during development and regeneration in the marine polychaete annelid Capitella teleta. *EvoDevo*, *2*(1). https://doi.org/10.1186/2041-9139-2-10

Hadfield, M. G. (2010). Biofilms and Marine Invertebrate Larvae: What Bacteria Produce That Larvae Use to Choose Settlement Sites. *Https://Doi.Org/10.1146/Annurev-Marine-120709-142753*, *3*, 453–470. https://doi.org/10.1146/ANNUREV-MARINE-120709-142753

Hadfield, M. G., Meleshkevitch, E. A., & Boudko, D. Y. (2000). The apical sensory organ of a gastropod veliger is a receptor for settlement cues. *Https://Doi.Org/10.2307/1542804*, *198*(1), 67–76. https://doi.org/10.2307/1542804

Harrison, R. G. (1898). The growth and regeneration of the tail of the frog larva - Studied with the aid of Born's method of grafting. *Archiv Für Entwickelungsmechanik Der Organismen*, *7*(2–3), 430–485.

https://doi.org/10.1007/BF02161494

Hiebert, T. C., & Maslakova, S. (2015). Integrative taxonomy of the Micrura alaskensis Coe, 1901 species
complex (Nemertea: Heteronemertea), with descriptions of a new genus maculaura gen. nov. and
four new species from the NE pacific. *Zoological Science*, *32*(6), 615–637.
https://doi.org/10.2108/ZS150011

Jeffery, W. R. (2015). Regeneration, Stem Cells, and Aging in the Tunicate Ciona: Insights from the Oral
Siphon. *International Review of Cell and Molecular Biology*, *319*, 255–282.
https://doi.org/10.1016/bs.ircmb.2015.06.005

Juliano, C. E., Swartz, S. Z., & Wessel, G. M. (2010). A conserved germline multipotency program.
Development, *137*(24), 4113–4126. https://doi.org/10.1242/DEV.047969

Luttrell, S. M., Su, Y. H., & Swalla, B. J. (2018). Getting a head with ptychodera flava larval regeneration.
Biological Bulletin, *234*(3), 152–164. https://doi.org/10.1086/698510

Meyer, N. P., Boyle, M. J., Martindale, M. Q., & Seaver, E. C. (2010). A comprehensive fate map by
intracellular injection of identified blastomeres in the marine polychaete Capitella teleta. *EvoDevo*,
1(1). https://doi.org/10.1186/2041-9139-1-8

Meyer, N. P., Carrillo-Baltodano, A., Moore, R. E., & Seaver, E. C. (2015). Nervous system development in
lecithotrophic larval and juvenile stages of the annelid Capitella teleta. *Frontiers in Zoology*, *12*(1),
1–27. https://doi.org/10.1186/s12983-015-0108-y

Meyer, N. P., & Seaver, E. C. (2009). Neurogenesis in an annelid: Characterization of brain neural
precursors in the polychaete Capitella sp. I. *Developmental Biology*, *335*(1), 237–252.
https://doi.org/10.1016/J.YDBIO.2009.06.017

Nedved, B. T., Freckelton, M. L., & Hadfield, M. G. (2021). Laser ablation of the apical sensory organ of
Hydroides elegans (Polychaeta) does not inhibit detection of metamorphic cues. *The Journal of
Experimental Biology*, *224*(20). https://doi.org/10.1242/JEB.242300

Oulhen, N., Heyland, A., Carrier, T. J., Zazueta-Novoa, V., Fresques, T., Laird, J., Onorato, T. M., Janies, D.,
& Wessel, G. (2016). Regeneration in bipinnaria larvae of the bat star Patiria miniata induces rapid
and broad new gene expression. *Mechanisms of Development*, *142*, 10–21.
https://doi.org/10.1016/j.mod.2016.08.003

Özpolat, B. D., & Bely, A. E. (2016). Developmental and molecular biology of annelid regeneration: a
comparative review of recent studies. *Current Opinion in Genetics & Development*, *40*, 144–153.
https://doi.org/10.1016/j.gde.2016.07.010

Pernet, B., Amiel, A., & Seaver, E. C. (2012). Effects of maternal investment on larvae and juveniles of the
annelid Capitella teleta determined by experimental reduction of embryo energy content.
Invertebrate Biology, *131*(2), 82–95. https://doi.org/10.1111/j.1744-7410.2012.00263.x

Phipps, L. S., Marshall, L., Dorey, K., & Amaya, E. (2020). Model systems for regeneration: Xenopus.
Development (Cambridge), *147*(6). https://doi.org/10.1242/DEV.180844/223048

Rawls, A., & Olson, E. N. (1997). MyoD meets its maker. *Cell*, *89*(1), 5–8. https://doi.org/10.1016/S0092-
8674(00)80175-0

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C.,

Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., &
Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*
2012 9:7, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Seaver, E. C. (2022). Sifting through the mud: A tale of building the annelid Capitella teleta for EvoDevo
studies. *Current Topics in Developmental Biology*, *147*, 401–432.
https://doi.org/10.1016/BS.CTDB.2021.12.018

Seaver, E. C., & De Jong, D. M. (2021). Regeneration in the Segmented Annelid Capitella teleta. *Genes*,
12. https://doi.org/10.3390/genes12111769

Seaver, E. C., Thamm, K., & Hill, S. D. (2005). Growth patterns during segmentation in the two
polychaete annelids, Capitella sp. I and Hydroides elegans: comparisons at distinct life history
stages. *Evolution Development*, *7*(4), 312–326. https://doi.org/10.1111/j.1525-142X.2005.05037.x

Simakov, O., Marletaz, F., Cho, S. J., Edsinger-Gonzales, E., Havlak, P., Hellsten, U., Kuo, D. H., Larsson, T.,
Lv, J., Arendt, D., Savage, R., Osoegawa, K., De Jong, P., Grimwood, J., Chapman, J. A., Shapiro, H.,
Aerts, A., Otillar, R. P., Terry, A. Y., … Rokhsar, D. S. (2013). Insights into bilaterian evolution from
three spiralian genomes. In *Nature* (Vol. 493, Issue 7433, pp. 526–531). Nature Publishing Group.
https://doi.org/10.1038/nature11696

Singer, M. (1954). Induction of regeneration of the forelimb of the postmetamorphic frog by
augmentation of the nerve supply. *Journal of Experimental Zoology*, *126*(3), 419–471.
https://doi.org/10.1002/jez.1401260304

Sur, A., Magie, C. R., Seaver, E. C., & Meyer, N. P. (2017). *Spatiotemporal regulation of nervous system*
development in the annelid Capitella teleta. *8*, 13. https://doi.org/10.1186/s13227-017-0076-8

Vickery, M. C. L., Vickery, M. S., Amsler, C. D., & Mcclintock, J. B. (2001). Regeneration in Echinoderm
Larvae. *Microscopy Research and Technique*, *55*(6), 464–473. https://doi.org/10.1002/jemt.1191

Vickery, M. S., & McClintock, J. B. (1998). Regeneration in metazoan larvae [6]. In *Nature* (Vol. 394, Issue
6689, p. 140). Nature Publishing Group. https://doi.org/10.1038/28086

Vickery, Minako S., Vickery, M. C. L., & McClintock, J. B. (2002). Morphogenesis and organogenesis in the
regenerating planktotrophic larvae of asteroids and echinoids. *Biological Bulletin*, *203*(2), 121–133.
https://doi.org/10.2307/1543381

Yamaguchi, E., Dannenberg, L. C., Amiel, A. R., & Seaver, E. C. (2016). Regulative capacity for eye
formation by first quartet micromeres of the polychaete Capitella teleta. *Developmental Biology*,
410(1), 119–130. https://doi.org/10.1016/j.ydbio.2015.12.009

Yamaguchi, E., & Seaver, E. C. (2013). The importance of larval eyes in the polychaete Capitella teleta:
Effects of larval eye deletion on formation of the adult eye. *Invertebrate Biology*, *132*(4), 352–367.
https://doi.org/10.1111/ivb.12034

Zammit, P. S. (2017). Function of the myogenic regulatory factors Myf5, MyoD, Myogenin and MRF4 in
skeletal muscle, satellite cells and regenerative myogenesis. *Seminars in Cell & Developmental*
Biology, *72*, 19–32. https://doi.org/10.1016/J.SEMCDB.2017.11.011

Zattara, E. E., & Bely, A. E. (2013). Investment choices in post-embryonic development: Quantifying
interactions among growth, regeneration, and asexual reproduction in the annelid *Pristina leidyi*.
Journal of Experimental Zoology Part B: Molecular and Developmental Evolution, *320*(8), 471–488.

https://doi.org/10.1002/jez.b.22523

Zattara, E. E., & Bely, A. E. (2016). Phylogenetic distribution of regeneration and asexual reproduction in
Annelida: regeneration is ancestral and fission evolves in regenerative clades. *Invertebrate Biology*,
135(4), 400–414. https://doi.org/10.1111/IVB.12151

Figure Legends

Figure 1. Regenerating juveniles and larvae of *C. teleta*. All images are oriented in ventral view with
anterior to the left. **A**, **B**, and **C** are images of the posterior end of a regenerating juvenile 3 days post-
amputation (dpa) and are from the same specimen. **A** Differential interference contrast (DIC) image
showing regeneration of new tissue in juveniles. **B** Nervous system visualized with anti-acetylated
tubulin antibody labeling. **C** Phalloidin staining shows muscle fibers with few in regenerating tissue . **D**
Stage 6 larva. **E** Stage 9 larva. (G) gut, (N) neural projection, (CM) circular muscle, (Ey) eye, (Pt)
prototroch, (Tt) telotroch, (Py) pygidium. The white dotted lines indicate amputation sites. The asterisk
marks the position of the mouth. N denotes the number of animals scored that resemble the image
shown.

Figure 2. Amputated larvae undergo successful wound healing. All images are oriented in ventral view
with anterior to the left. Arrows indicate the continuous layer of epithelium, indicative of successful
would healing. **A** Stage 6 uncut larva showing experiment design for amputations. The dotted lines
indicate cut sites: (AA) anterior amputation and (PA) posterior amputation. **B** Stage 6 larva 6 hpa
anterior amputation. **C** Stage 6 larva 6 hpa posterior amputation. **D** Larva 3 dpa anterior amputation. **E**
Stage 9 larva 2 dpa anterior amputation. **F** Stage 9 larva 6hpa posterior amputation. The black bracket
represents the tissue between the gut and ectoderm. Amputation site is indicated by the black dotted
line. N denotes the number of animals scored that resembled the image in the panel. Scale bar is 50 µm.

Figure 3. Localized cell proliferation near the amputation site in larvae. All images are oriented in ventral
view with anterior to the left. Animals were exposed to EdU to label dividing cells. EdU+ cells are in
green. **A–D** are uncut controls: **A** Stage 6, **B** Stage 7, **C** Stage 8, and **D** Stage 9. **E–H** are anterior
amputations: **E** 6 hpa, **F** 1 dpa, **G** 3 dpa, and **H** 5 dpa. **I–L** are posterior amputations: **I** 6 hpa, **J** 1 dpa, **K** 3
dpa, and **L** 5 dpa. **M**, **N** Quantification of EdU incorporation in anterior (M) and posterior (N)
amputations and comparison with uncut controls. Time after amputation is on the X-axis and the ratio of
EdU/DAPI cells is on the Y-axis. The white arrows indicate EdU+ cells near the wound site.
Developmental stage or time post-amputation is indicated in the bottom left corner of each panel. N
denotes the number of animals scored that resemble the image shown. Dark blue represents uncut
controls and light blue represents amputated animals. Asterisk denotes significance with $p < 0.05$. Scale
bar is 50 µm.

Figure 4. *Vasa* is expressed at the wound site in amputated larvae. All images are oriented in ventral
view with anterior to the left. The purple coloration indicates localization of *vasa* transcripts by in situ
hybridization. **A**, **B**, and **C** *Vasa* expression patterns in uncut larvae at Stage 6, Stage 8, and Stage 9. **D**, **E**,
and **F** are anterior larval amputations. **G**, **H**, and **I** are posterior larval amputations. **D** and **G** are
amputated larvae 6 hpa (D) and 2 hpa (**G**). **E** De novo expression is observed in anterior amputations at 2
dpa, as indicated by the arrow and is no longer detectable by 5 dpa (**F)**. In posterior amputations,
expression near the wound site is observed at both 2 dpa (**H**) and 3 dpa (**I**), as indicated by arrows. The

time post-amputation is indicated in the bottom left corner of each panel in D–I. N denotes the number
of animals scored that resemble the image shown. (FG) foregut, (MPC) multi-progenitor cell cluster,
(PGZ) posterior growth zone. Scale bar is 50 μm.

Figure 5. Tissue response to amputation. **A–B”** and **E–I”** are oriented in ventral view. **C–D”** are oriented
in lateral view. In all images, anterior is to the left. **A** uncut Stage 6 larva labeled with anti-acetylated
tubulin to visualize the nervous system and ciliary bands. **B, C, D, H, I** larvae are labeled with anti-
acetylated tubulin (red) and phalloidin (green) showing the nervous system and musculature,
respectively. **B**, **B’**, and **B”** are the same specimen. **B** Nervous system and musculature in a larva 2 dpa.
B’ magnified view of posterior wound site showing few muscle fibers distal to cut site. **B”** magnified view
of posterior wound site showing fibers in the tissue distal to the amputation site (arrows). **C** larva 6 hpa
after anterior amputation. **D**, **D’** and **D”** are the same specimen, 3 dpa anterior amputation. **D’** magnified
view of muscle fibers as marked by phalloidin. **D”** magnified view of anti-acetylated tubulin labeling. **E**
and **E’** are the same larva 6 hpa labeled with anti-acetylated tubulin; **E’** as a magnification of **E**. **F** and **G**
are separate 2 dpa posterior amputated larvae. **F** neural fibers are present distal to the cut site (arrow).
G shows a specimen lacking neural fibers distal to the cut site. **H** 6 hpa posterior amputation. **I, I’,** and **I”**
are the same individual at 5 dpa posterior amputation. **I’** longitudinal and circular muscles are present
distal to the cut site (arrow). **I”** neurites distal to the cut site (arrow). White brackets in **I’** and **I”** indicate
tissue distal to the amputation site. N denotes the number of animals scored that resemble the
representative image. Amputation site is indicated by a white dotted line. (VNC) ventral nerve cord, (PN)
peripheral nerves. Scale bar is 50 μm.

Figure 6. Amputation induces *ash1* but not *elav* expression at the anterior wound site. All images are
oriented with anterior to the left. **A-E** are ventral views and **F-I** are lateral views. The purple coloration
indicates expression of either *ash1* (**A-E**) or *elav* (**F-I**) transcripts by in situ hybridization. **C-E** and **H-I** are
anterior amputations. **A** Stage 6 uncut control. **B** Stage 9 uncut control. **C** 6 hpa. **D** and **E** are the same 3
dpa individual. Vertical arrows indicate novel *ash1* ectodermal expression. The boundary between
ectodermal and mesodermal layers is indicated by a black dotted line. **F** Stage 6 and **G** Stage 9 uncut
controls. **H** 6 hpa. **I** 3 dpa. (Br) brain, (FG) foregut, (VNC) ventral nerve cord. N denotes the number of
animals scored that resemble the image indicated. Scale bar is 50 μm.

Figure 7. Amputation induces ectopic expression of the myogenesis marker *MyoD*. All images are
oriented in ventral view with anterior to the left. *MyoD* expression is visualized by the purple coloration
via in situ hybridization. **A** Stage 6 and **B** Stage 9 uncut controls. *MyoD* is expressed in the posterior
trunk mesoderm. **C** and **D** are anterior amputations (ant amp). **C** 6 hpa. **D** anterior shift in *MyoD*
expression is observed at 7 dpa. **E** 6 hpa posterior amputation (post amp). Spots of staining anterior to
the bracket is nonspecific trapping in the chaetae. **F** *MyoD* is expressed along the length of the body in 3
dpa animals (compare with E). Brackets indicate the boundaries of detectable *MyoD* expression along
the anterior-posterior axis. Amputation sites are indicated by black dotted lines. N denotes the number
of animals scored that resemble the representative image. Scale bar is 50 μm.

Figure 8. Eye reformation following anterior amputation. All images are oriented with anterior to the
left. **A** – **D** are ventral views. **A** DIC image of the anterior end of a Stage 9 uncut larva. **B** DIC image of the
anterior end of a larva 5 days after anterior amputation. Image shown is a merge of multiple focal
planes. The pigmented cells (PC) of the eye are indicated by arrows. **C**–**F** eye sensory cells (SC, white
arrows), labeled by the 22C10 antibody (green). **C** Uncut Stage 9 control. **D** anterior amputated larva 5

dpa. **E** and **F** are juvenile worms in lateral view. **E** uncut 2 week-old juvenile and **F** juvenile 1 week post-
amputation of the head. **G** Table displaying the number of pigment cells, sensory cells, or either cell
(total) present in anteriorly amputated larvae at 6 hpa and 3 dpa or later. Dotted line indicates the
outline of the animal. (MG) midgut.

Figure 9. Rare reformation of cilia occurs in amputated larvae. Animals are labelled with anti-acetylated
tubulin (white). All images are oriented in with anterior to the left. **A** and **D** are uncut controls, oriented
in ventral view. **A** Stage 6 uncut larva. **D** Stage 9 uncut larva. **B** and **E** are anterior amputations oriented
in lateral view, with anterior to the left. **B** larva 6 hpa. **E** larva 5 dpa with short cilia visible distal to the
cut site (arrow). **C** and **F** are posterior amputations in ventral view. **C** larva 6 hpa. **F** larva 5 dpa with both
long cilia (LC) and short cilia (SC) distal to the cut site (arrows). (LC) long cilia, (SC) short cilia, (NT)
neurotroch, (PT) prototroch, (PYG) pygidium, (TT) telotroch. N denotes the number of animals scored
that are similar to the image represented. Amputation site is indicated by the white dotted line. Scale
bar is 50 µm.

Figure 10. Induction of metamorphosis in amputated larvae. All images are oriented in ventral view with
anterior to the left. **A** Uncut Stage 9 larva not exposed to the metamorphic cue. **B** Uncut Stage 9 larva 2
hr after inducement with metamorphic cue. **C** anteriorly amputated larva not exposed to metamorphic
cue. **D** anteriorly amputated larva 2 hr after exposure to metamorphic cue. **E** posteriorly amputated
larva not exposed to metamorphic cue. **F** posteriorly amputated larva 2 hr after exposure to
metamorphic cue. (Br) brain, (Ph) pharynx, (Pt) prototroch, and (Tt) telotroch. N denotes the number of
animals scored that are similar to the image shown. All images have the same magnification; scale bar
50 µm.

A — AA, PA

B — N = 54/54 — Ant Amp

C — N = 48/51 — Post Amp

D — N = 10/10 — 3 dpa

E — N = 10/10 — 2 dpa

F — N = 10/10 — 6 hpa

Ash1

Elav

A

Uncut Cont

St 6

B

St 9

C

Ant Amp

6 hpa

D

7 dpa

E

Post Amp

6 hpa

F

3 dpa

	Larval Pigment Cell	Larval Sensory Cell	Juv. Sensory Cell
Cont	A St 9	C St 9	E 2 wk
Amp	B 5 dpa	D 7 dpa	F 1 wpa

G

	Pigment Cell	Sensory Cell	Total
6 hpa	0/46	0/31	0/57
3 dpa +	28/420	8/220	36/640
	7%	4%	6%

| Uncut | Ant Amp | Post Amp |

	Uncut Controls	Ant Amp	Post Amp
Not Induced	A — N = 532/534	C — N = 65/66	E — N = 65/65
Induced	B — N = 203/207	D — N = 135/136	F — N = 143/144

Labels: Br, Ph, Pt, Tt

Table 1. Quantification of EdU+ cells in anteriorly amputated larvae

Anterior Amp		Anterior		Middle		Posterior	
		Uncut	Cut	Uncut	Cut	Uncut	Cut
6 hpa	# Nuclei	418	392	1216	417	551	250
	# EdU	133	102	466	235	387	136
	%	32%	26%	38%	56%	70%	54%
3 dpa	# Nuclei	447	363	856	457	413	185
	# EdU	3	29	18	2	14	24
	%	1%	8%	2%	0%	3%	13%
5dpa	# Nuclei	147	359	856	516	413	219
	# EdU	3	28	18	5	14	10
	%	2%	8%	2%	1%	3%	5%

Table 2. Quantification of EdU+ cells in posteriorly amputated larvae

Posterior Amp		Anterior		Middle		Posterior	
		Uncut	Cut	Uncut	Cut	Uncut	Cut
6 hpa	# Nuclei	418	155	1216	556	551	463
	# EdU	133	70	466	217	387	225
	%	32%	45%	38%	39%	70%	49%
3 dpa	# Nuclei	447	306	856	702	413	525
	# EdU	3	2	18	6	14	78
	%	1%	1%	2%	1%	3%	15%
5dpa	# Nuclei	147	331	856	547	413	513
	# EdU	3	0	18	0	14	51
	%	2%	0%	2%	0%	3%	10%

Table 3. Amputated larvae measured for tissue distal to cut site

		Muscle	Nervous System	Gut	Width 2 /Width 1
Ant Amp	**6 hpa**	19μm	18μm	N/A	N/A
	3 dpa	22μm	15μm	N/A	N/A
Post Amp	**6 hpa**	* 17μm	* 12μm	N/A	* 0.675
	5 dpa	* 62μm	* 67μm	45μm	* 0.453
	St 9 6 hpa	N/A	N/A	37μm	N/A

* = significance, p < .05