Multiplex PCR with Nanopore Sequencing for Sequence-Based Detection of Four Tilapia Pathogens

Jérôme Delamare-Deboutteville[1#], Watcharachai Meemetta[2#], Khaettareeya Pimsannil[2], Han Ming Gan[3], Laura Khor Li Imm[1], Chadag Vishnumurthy Mohan[1], Ha Thanh Dong[4], Saengchan Senapin[2,5]

[1]WorldFish, Penang, Malaysia
[2]Fish Health Platform, Center of Excellence for Shrimp Molecular Biology and Biotechnology (Centex Shrimp), Faculty of Science, Mahidol University, Bangkok, Thailand
[3]Patriot Biotech Sdn Bhd, Bandar Sunway, 47500, Selangor, Malaysia
[4]School of Environment, Resources and Development, Asian Institute of Technology, Pathum Thani, 12120, Thailand
[5]National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), Pathum Thani, Thailand

Corresponding authors:
Jérôme Delamare-Deboutteville
Jalan Batu Maung, Batu Maung, 11960 Bayan Lepas, Penang, Malaysia
Email address: j.delamare@cgiar.org

Saengchan Senapin
272 Rama VI Road, Ratchathewi, Bangkok 10400, THAILAND
Email address: saengchan@biotec.or.th
# These authors contributed equally to this work.

Abstract

Background. Tilapia aquaculture faces significant threats posed by four prominent pathogens: tilapia lake virus (TiLV), infectious spleen and kidney necrosis virus (ISKNV), *Francisella orientalis*, and *Streptococcus agalactiae*. Currently, employed molecular diagnostic methods for these pathogens rely on multiple singleplex PCR reactions, which are time-consuming and expensive.

Methods. In this study, we present an approach utilizing a multiplex PCR (mPCR) assay, coupled with rapid Nanopore sequencing, enabling the one-tube simultaneous detection and one-reaction Nanopore sequencing-based validation of four pathogens.

Results. Our one-tube multiplex assay exhibits a detection limit of 1,000 copies per reaction for TiLV, ISKNV, and *S. agalactiae*, while for *F. orientalis*, the detection limit is 10,000 copies per reaction. This sensitivity is sufficient for diagnosing infections and co-infections in clinical samples from sick fish, enabling rapid confirmation of the presence of pathogens. Integrating multiplex PCR and Nanopore sequencing provides an alternative approach platform for fast and precise diagnostics of major tilapia pathogens in clinically sick animals, adding to the available toolbox for disease diagnostics.

Introduction

Tilapia (*Oreochromis* spp.) is one of the most widely farmed freshwater fish species globally due to its high adaptability, fast growth, and excellent meat quality. Global production is estimated at 6,100,719 tonnes in 2020 (FAO, 2022). By providing sustenance, employment opportunities, and domestic and export revenues, this valuable species supports large populations worldwide (Wang & Lu, 2016). In the past decade, tilapia production has almost doubled (FAO, 2020), attributed mainly to its ease of cultivation, market demand, and stable pricing (Wang & Lu, 2016).

However, the production of tilapia has been threatened by several viral and bacterial diseases that can cause significant economic losses (Debnath et al., 2023). Among the most important tilapia pathogens are tilapia lake virus (TiLV), infectious spleen and kidney necrosis virus (ISKNV), *Francisella orientalis*, and *Streptococcus agalactiae* (SAG) also called group B *streptococcus* (GBS) (Kawasaki et al., 2018; Machimbirike et al., 2019; Mabrok et al., 2021; Haenen et al., 2023; Alathari et al., 2023). Effective diagnosis and monitoring of these pathogens are crucial for disease management and control (Dong et al., 2023). Traditional diagnostic techniques such as virus and bacterial isolation, immunofluorescence assay (IFA), and enzyme-linked immunosorbent assay (ELISA) are time-consuming, labor-intensive, and require specialized equipment and expertise (Dong et al., 2023). In contrast, molecular methods such as polymerase chain reaction (PCR) and quantitative PCR (qPCR) have gained widespread use in pathogen detection due to their high sensitivity and specificity, rapid turnaround time, and ability to detect multiple pathogens in a single reaction (Soto et al., 2010; Kralik & Ricchi, 2017; Liamnimitr et al., 2018; Waiyamitra et al., 2018; López Porras et al., 2019; Ramírez-Paredes et al., 2021; Taengphu et al., 2022; Dong et al., 2023).

In current real-world situations, it is uncommon to perform subsequent sequencing of the positive PCR or qPCR products to validate the pathogen identity due to the time and resources involved in conventional Sanger sequencing. Sequence-based verification is particularly valuable in multiplex PCR since non-specific amplification is more likely to occur in a multiplex PCR reaction due to multiple primer combinations. Fortunately, this is now possible with Oxford nanopore technology (ONT) that allows on-site amplicon sequencing (Delamare-Deboutteville et al., 2021). Therefore, proper integration of multiplex PCR and ONT-based genotyping may present a promising tool for the efficient and precise detection and characterization of pathogens in tilapia aquaculture.

We present a multiplex PCR assay with a detection level suitable for simultaneously identifying TiLV,
ISKNV, *F. orientalis*, and *S. agalactiae* in sick tilapia samples. Using the portable nanopore sequencing
platform, we confirmed the presence of these pathogens at the sequence level and gathered genetic
information from the amplicons. This assay offers a rapid and practical solution for detecting and
genetically characterizing these pathogens, with the potential to complement existing diagnostic tools and
inform targeted surveillance and control strategies.

Materials & Methods

Ethics declarations
93
94 The authors confirm that the journal's ethical policies, as noted on the journal's author guidelines page,
95 have been adhered to. No ethical approval was required as no animals were used in this study. Virus
96 sequences were generated from archived samples.
97

98 **Clinical samples and nucleic acid extraction**

99
We utilized archival clinical samples of fry, fingerling, juvenile and adult Nile tilapia, red tilapia, and Asian
sea bass from challenge experiments or from various disease outbreaks between 2015 and 2020. Our
investigation included samples that were either confirmed to be caused by a single pathogen using PCR
diagnosis or suspected to have resulted from co-infections. All sample details are comprehensively listed
in Table 1. To extract the nucleic acid from the samples, some were processed using the PathoGen-spin
DNA/RNA extraction kit from iNtRON Biotechnology, while others were archival RNA samples extracted
using Trizol reagent from Invitrogen, and DNA samples extracted using the conventional
phenol/chloroform ethanol precipitation method as described by (Meemetta et al., 2020). If done close to
the farm, the diagnostic workflow from the point of sample collection to the final data analysis can take
less than 12 hours. The processes include nucleic acid extraction, multiplex PCR, library preparation,
Nanopore sequencing, and data analysis, as illustrated in Figure 1.
111

Primers used in this study
We obtained the primers for the four selected target pathogens from Bio Basic (Canada), and the primer
sequences are summarized in Supplemental Table 1. Primers for TiLV, ISKNV, *F. orientalis*, and *S.*
agalactiae were reported in previous studies (Yang et al., 2013; Dong et al., 2016; Paria et al., 2016;
Leigh et al., 2018; Kawato et al., 2021; Taengphu et al., 2022). The expected amplicon sizes were 137,
190, 203, and 351 bp, respectively. The specificity of each primer pair was assessed *in silico* using the
Primer-BLAST program (https://www.ncbi.nlm.nih.gov/tools/primer-blast/).
119

Plasmid positive controls and the analytical sensitivity assay
All plasmid-positive controls except for *S. agalactiae* (constructed in this study) were obtained from our
previous studies (Supplemental Table 2). Positive plasmid control containing *S. agalactiae groEL* partial
gene was obtained by cloning a 351 bp-*groEL* amplified fragment purified before being ligated into
pGEM-T easy vector (Promega). A recombinant plasmid was subjected to DNA sequencing (Macrogen).
The copy number of each plasmid was calculated based on its size in base pair (bp) and amount in
nanogram (ng) using a web tool at http://cels.uri.edu/gsc/cndna.html. A combination of four recombinant
plasmids was mixed for the multiplex PCR sensitivity assay. This mixture was then subjected to 10-fold
serial dilution, resulting in a 1 to 10^6 copies/µl range. Subsequently, 4 µl of the diluted series was used in
the multiplex PCR reaction. To simulate a clinical sample from a fish, each PCR reaction was spiked with
50 ng of RNA extracted from a healthy tilapia.
131

Multiplex PCR condition optimization and the detection of clinical samples
Previous studies claimed that qPCR reagents usually contain PCR additives and enhancers that can
improve amplification efficiency (Karunanathie et al., 2022). To optimize our multiplex PCR, we used
KAPA SYBR FAST One-Step qRT-PCR master mix (Roche), known to contain such additives. Each
reaction was prepared in a 25 µl volume, comprising 1X master mix, 4 µl of template, and 80-240 nM of
each primer pair (Supplemental Table 3). We determined the optimal annealing temperature (Ta) using

gradient PCR with Ta ranging between 55-65 °C. To identify the best combination of PCR components,
we added ammonium sulfate, BSA, dNTPs, and MgCl2 in different proportions. The final cycling
conditions comprised a reverse transcription step at 42 °C for 5 min, followed by inactivation at 95 °C for
3 min, and 40 cycles of 95°C for 10 s and 60°C for 30 sec, with a final extension step at 60 °C for 5 min
(Supplemental Table 3). We then analyzed 10 µl of each product by electrophoresis on a 3.5% agarose
gel stained with ethidium bromide. The newly optimized multiplex PCR assay was subsequently used to
detect the presence of TiLV, ISKNV, *F. orientalis*, and *S. agalactiae* in archival clinical samples (Table 1).
Most of these samples were previously tested for a single pathogen using PCR or qPCR assays.

Quantification of pathogens by qPCR assays
A total of 15 clinical samples, whose mPCR products underwent Nanopore sequencing, were analyzed to
determine the presence and quantity of each pathogen. The established protocols (Leigh et al., 2018;
Kawato et al., 2021; Taengphu et al., 2022) were followed to detect TiLV, ISKNV, and SAG, as described
in Supplemental Table 4. For FnO qPCR, the same primers used in mPCR were used in this study's
SYBR Green-based qPCR assay.

Nanopore sequencing
The present study employed amplicons generated from both single and multiplex PCR reactions (Table 1,
samples highlighted in grey) as templates for library preparation using the ligation sequencing kit (SQK-
LSK109) and the native barcoding expansion 1-12 kit (EXP-NBD104) according to the standard protocols
of Oxford Nanopore Technologies (ONT) adapted for the Flongle flow cell. Three sequencing runs (r1-r3)
were conducted, with 250 ng of PCR product per sample and a unique native barcode (BC) assigned to
each sample. Run 1 (r1) involved sequencing amplicons obtained from individual PCR reactions and a
combination of these amplicons (Table 1). Run 2 (r2) focused on sequencing multiplex PCR (mPCR)
products derived from clinical samples. Run 3 (r3) employed mPCR products from the remaining clinical
samples. Additionally, different concentrations of single PCR products were included in this run. The
library of pooled barcoded samples was subjected to a Short Fragment Buffer (SFB) wash before the final
elution step of the protocol. The DNA concentration was quantified at every step using the Qubit assay.
Subsequently, the prepared library was loaded onto a Flongle flow cell (FLO-FLG106), following the
Nanopore standard protocol, and each Flongle flow cell was fitted to a Flongle adapter (FLGIntSP) for
MinION.

Reads filtering and read abundance calculation
Basecalled reads in fastq format were primer-trimmed with cutadapt v4.3 (Martin, 2011), and reads that
have been trimmed with length ranging from 75-400 bp will be retained for the subsequent analysis,
leaving out overly short or long reads without intact primer sequence on both ends. The filtered reads
were aligned to the four pathogen gene segments using Minimap2 v2.17 (Li, 2018). Reads that aligned
uniquely (only a single hit reported) with more than 80% coverage to the target region were used for
abundance calculation and variant calling (consensus generation). Calculation of raw, trimmed, and
aligned read statistics used seqkit v2.20 (Shen et al., 2016). Reads failing to align at this stringent level
were extracted and re-aligned using the default Minimap2 setting, followed by a less stringent blastN
alignment. The host genome was included in the reference sequence to assess the fraction of reads
mapping to the host.

Generation of consensus and variant sequences
Only reads with unique alignments were used as the template for Minimap2 and medaka variant calling
based on the ARTIC pipeline ("Core Pipeline - artic pipeline"; "sars-cov-2-ont-artic-variant-calling/COVID-
19-ARTIC-ONT"). The reads were mapped to the reference gene segments of four pathogens, followed
by variant calling using the medaka variant model r941_min_g507. Variant filtering was then performed
based on mapping quality and read depth. The number of ambiguous bases in the consensus sequences
was calculated using QUAST v5 (Gurevich et al., 2013). Only sequences without any ambiguous base
were selected for subsequent variant sequence generation. Variants were generated using cd-hit v.4.8.1
(Huang et al., 2010), whereby sequences with 100% nucleotide identity and exact sequence length were
clustered into the same genotype. The variant sequences were compared against the NCBI Blast nt
database (accessed on 25 April 2023) to identify the top 5 BLAST hits for each sequence.

Code availability
The Linux scripts used to generate initial FastQ files, assembled amplicons (public and from this study)
are publicly available in the Zenodo.org dataset https://doi.org/10.5281/zenodo.7866295.

Results

199
Primer testing by single PCR
Single PCR reactions were performed using each primer pair with its respective target template to
validate the selected primer pairs for each pathogen. The amplification of the target regions was
confirmed by amplicons of the expected sizes, as illustrated in Supplemental Figure 1. To simulate the
expected products from multiplex PCR (mPCR), an equal amount of each individual PCR product was
loaded into lane C of the Supplemental Figure 1. Gel electrophoresis analysis revealed distinct and
separable bands, demonstrating the potential for further utilization in the multiplex PCR analysis. Note
that the products generated from this step were subsequently employed in Nanopore sequence run 1 (r1).

208
Multiplex PCR detection of four pathogens among clinical samples
We have successfully optimized a multiplex PCR assay to simultaneously detect four important tilapia
pathogens, TiLV, ISKNV, *F. orientalis*, and *S. agalactiae*, in a single reaction. The detection sensitivity of
the new assay in the presence of spiked host RNA was 10^3 copies/reaction for TiLV, ISKNV, and *S.*
agalactiae. In the presence of 10^4 copies of each template, the assay could detect *F. orientalis* and three
other pathogens (Supplemental Figure 2). The assay could detect each pathogen efficiently in tilapia
clinical samples, as confirmed by distinct PCR bands (Figure 2). The assay was able to detect TiLV in
heavily infected samples (3-7) both on the gel and by nanopore but failed to detect it on the gel for
samples (8-12) with low levels of TiLV infection. Similarly, the highest number of reads for ISKNV were
found in sample #25 (twice more than in sample #27), yet it had a much fainter band on the gel (Table 1).
There is a good correlation between read numbers and bands intensity for *F. orientalis* in four samples
(#30-33) but only a weak band for *S. agalactiae* in one sample (#39) compared to number of amplicons
(Figure 2, 3, Supplemental Table 5). In some samples, we detected dual infections with more than one
pathogen, as evidenced by the presence of multiple bands, including a possible co-infection with *F.*
orientalis and TiLV in samples 3-7. However, differentiation of *F. orientalis* and ISKNV was challenging as
their respective bands have similar sizes (Figure 2).

225
Successful recovery of pathogen sequence variants from infected samples
A total of 246,756 reads were successfully demultiplexed from three separate Flongle runs (r1-r3), each
containing different sample types (Table 1, Supplemental Table 5). Following primer and length filtering, a
substantial reduction in read count was observed, resulting in an average of 64% of reads being removed
(range: 46%-87%) (Supplemental Table 5). An additional 10% of reads were removed after alignment
filtering. After these filtering steps, 75,203 reads remained, providing an average of 2,500 filtered reads
per sample for subsequent read abundance and consensus sequence generation (Supplemental Table
5).

234
Given the small sampling size and possibly highly conserved nature of some of the pathogen gene
segments, one sequence variant was generated per pathogen gene segment except for the TiLV gene
segment, whereby two sequence variants were recovered (Table 2). All recovered sequence variants
exhibited 100% nucleotide identity to at least one publicly available sequence in the NCBI database. In
addition, samples derived from the single PCR product inputs and their combined mixture were clustered
in their respective sequence variants. In the case of the TiLV amplicons, we observed two unique
sequence variants (Var 0 and Var 1) for TiLV gene segment 9, one of which displayed 100% nucleotide
identity to the gene region of a relatively divergent TiLV strain from Vietnam (Table 2). This sequence
variant was detected in samples r2_NT_F1 to r2_NT_F5, collected from the same sampling site
simultaneously (Table 2).

245
246
247

Effect of template input on data distribution and pathogen detection specificity on the Nanopore platform

In the first run (r1), using well-defined pathogen nucleic acids as templates for the detection assay, confident alignments were observed for each sample to their expected pathogen gene segments, except for sample r1_TILV_1 which consisted purely of TiLV amplicons and did not sequence well enough to enable subsequent variant calling (Figure 3 and Table 2). For sample 4PAT, which consisted of a combination of products from 4 pathogens, the read distribution was broadly similar, as was the case for sample 4PAT_DIL, which used a ten-fold diluted version of the template (Figure 3). In the second run (r2), actual clinical samples with PCR-verified infections were used, and over 50% and up to 99% of reads were primarily mapped to the primary suspected pathogen gene fragments, with the remaining reads mapping to the other three pathogens. Interestingly, in the third run (r3), a significant percentage (>5%) of reads aligned to other pathogens were observed among samples spiked with different amounts of single amplified products (Figure 3). For example, among the TiLV samples (TILV05, 15, 30, 50), the lowest amount of PCR product template (TILV05) resulted in a low number of TiLV reads and a higher percentage of reads belonging to non-TiLV pathogens. A similar trend was observed for pure SAG samples, with the most diluted SAG sample (SAG05) having the highest percentage of non-SAG reads. However, in contrast to the pure synthetic TiLV samples, the percentage of non-specific reads decreased to less than 2% in the SAG50 sample with the highest read count, while it remained around 10% in the TILV50 sample.

Reads that failed stringent alignment to the pathogen gene panel were host-derived or partial pathogen sequences with lower nucleotide identity

Upon investigation of reads that failed to align, a significant proportion was found to map not only to the four pathogen gene segments but also to the host genome when a more lenient alignment configuration was employed (Figure 4). Samples from run 2 (r2), predominantly derived from fish tissues, exhibited a higher percentage of reads mapping to the host genome than other samples. As expected, given the use of input PCR products as the template for all samples in run 1 (r1), no reads were found to align to the host genome when they were aligned with Minimap2 (default setting). On the contrary, a small portion of reads belonging to the host genome was found among different concentrations of single PCR products from run 3 (r3) (Figure 4 and Supplemental Table 5), which also consists of samples derived from tilapia organs (Table 1).

Discussion

The intensification of tilapia farming has led to a surge in emerging and re-emerging infectious diseases, resulting in substantial losses in the aquaculture industry (Machimbirike et al., 2019; Haenen et al., 2023; Shinn et al., 2023). These diseases continue to evolve and spread under intense selective pressures, particularly in regions with high-density tilapia farming. To address this issue, we built upon our previous study (Delamare-Deboutteville et al., 2021), and improved the detection and genotyping of multiple pathogens by switching to multiplex PCR coupled with Nanopore sequencing. Our updated approach enabled the simultaneous and sequence-based detection and verification of four major tilapia pathogens: ISKNV, TiLV, *S. agalactiae*, and *F. orientalis.*

The requirement for simultaneous diagnosis of multiple pathogens is particularly relevant to high mortalities observed in larger grow-out tilapia. These mortalities are likely exacerbated by co-infections, as indicated by (Ramírez-Paredes et al., 2021) previous research findings, which revealed the active co-infection of ISKNV-positive fish with Streptococcus agalactiae and other bacterial pathogens.

The discrepancies between gel-based interpretation of multiplex PCR products and qPCR or direct amplicon sequencing approaches are not surprising, given the subjective nature of gel visualization and the labor-intensive process of running multiple samples. The accuracy of pathogen detection can be significantly impacted by faint or similar-sized bands, making it challenging to standardize results. To streamline the multiplex PCR approach and increase throughput, the gel visualization step can be eliminated in favor of direct Nanopore sequencing. With high-degree sample-based multiplexing, it is now possible to perform high throughput sequence-based detection of multiple tilapia pathogens using

302 Nanopore sequencing. The utilization of this application is paramount for the aquaculture industry, given
303 that disease outbreaks are frequently linked with multiple infections (Abdel☐Latif & Khafaga, 2020; Huang
304 et al., 2020; Liu et al., 2020; Basri et al., 2020). Furthermore, as aquaculture farms are often situated in
305 remote locations far from diagnostic reference laboratories, the ability to mobilize diagnostic testing with
306 rapid results, high accuracy, and less complicated equipment near the farm is preferred.
307
Although our approach is scalable, there are limitations due to the old version of MinKNOW used during
the study. Short amplicons are sequenced less efficiently and have lower read accuracy. We addressed
this issue using a stringent read filtering and alignment approach, which led to a reduced read recovery in
the final abundance calculation. Some synthetic samples analyzed contained reads from other
pathogens, which was unexpected and suggests cross-ligation of native barcodes during library
preparation. The availability of unbound adapters is higher among samples with low DNA input, leading to
higher crosstalk levels and reads from other samples. New improvements in the current Nanopore
technology and library preparation protocols (as of 14 September 2024) will mitigate these issues. This
includes (1) Switching to a PCR-barcoding kit to only enrich for amplicon with Nanopore partial adapter,
(2) the use of a new Q20+ LSK114 kit with improved read accuracy, and (3) the use of high accuracy
Dorado base-calling model (https://github.com/nanoporetech/dorado).
319
By enforcing a stringent alignment setting, the reads aligning to the pathogen gene segment are directly
suitable for subsequent reference-based variant calling, producing highly accurate variants useful for
biological interpretation. This is exemplified by the consistent recovery of a TiLV segment 9 variant
previously only found in a Vietnamese TiLV strain in all Thai tilapia samples from the same sampling
batch. Considering the proximity of Thailand and Vietnam, this may suggest a possible spillover event,
highlighting the need for more thorough sampling and sequencing to elucidate the degree of TiLV
diversity in both regions.
327
For field diagnostics, some of the equipment used in the present study can be replaced with portable
items that are small enough to fit in a bag for use in remote settings. An example of the field application of
this diagnostic workflow is the Lab-in-a-backpack concept developed by WorldFish (WorldFish, 2020;
Huso et al., 2020; Cagua et al., 2021; The University of Queensland, 2021; Chadag et al., 2021; Barnes
et al., 2021). It includes all the necessary sampling, extraction, and sequencing kits, pipettes,
and consumables, as well as the use of the miniPCR thermocycler and blueGel electrophoretic system
from miniPCR bio™, a minicentrifuge, a magnetic rack, a fluorometer, a Nanopore MinION sequencer,
and flow cells for sequencing. Additionally, a laptop with minimum requirements to run the MinKNOW
software is required. Similar PCR-sequencing approaches using portable equipment have been employed
for the rapid identification of a diverse range of biological specimens, including plants, insects, reptiles
(Pomerantz et al., 2022), and viruses such as COVID-19, Ebola, and Zika (González-González et al.,
2019, 2020).
340
Improvements in the remote and low-resource applications of the current methodology will broaden its
use for capacity building and implementations in low-income countries, where high mortalities due to
unknown causes are prevalent in tilapia farming. Accurate and rapid genomic detection of fish pathogens
through our amplicon-based approach will allow health professionals to promptly provide advice and take
necessary actions for producers. Moreover, it can facilitate informed decisions regarding additional
investigations, such as whole-genome sequencing (WGS), on isolates stored in biobanks. The sequence
data obtained from WGS offer crucial epidemiological insights, which can be utilized to develop
customized multivalent autogenous vaccines using local pathogens (Barnes et al., 2022). These vaccines
can then be administered to broodstock and seeds before distribution among grow-out farmers for
restocking purposes.
351

Conclusions

353 In summary, our study presents an attractive approach for detecting and verifying four tilapia pathogens
354 in clinically sick fish, which can also be applied to pathogens in other livestock, fish, or crustaceans if the
355 genetic information of the pathogen is publicly available for primer design and reference mapping.

Although there are limitations to the current pipeline version used at the time of this study, we are optimistic that the current improvements in Nanopore technology will further enhance the accuracy and scalability of our approach.

Acknowledgements

We want to extend our utmost gratitude and profound recognition to the late Dr. Pattanapon Kayansamruaj for his valuable contributions during the early stages of conceiving this research endeavor.

References

Abdel□Latif HMR, Khafaga AF. 2020. Natural co-infection of cultured Nile tilapia Oreochromis niloticus with Aeromonas hydrophila and Gyrodactylus cichlidarum experiencing high mortality during summer. *Aquaculture Research* 51:1880–1892. DOI: 10.1111/are.14538.

Alathari S, Chaput DL, Bolaños LM, Joseph A, Jackson VLN, Verner-Jeffreys D, Paley R, Tyler CR, Temperton B. 2023. A Multiplexed, Tiled PCR Method for Rapid Whole-Genome Sequencing of Infectious Spleen and Kidney Necrosis Virus (ISKNV) in Tilapia. *Viruses* 15:965. DOI: 10.3390/v15040965.

Barnes A, Das S, Rudenko O, Wilkinson S, Cagua F, Delamare-Deboutteville J. 2021. Diagnosis in a fish farmer's backpack. In: *Food & Nutrition Security – The Biosecurity, Health, Trade Nexus. https://www.crawfordfund.org/events/other-events/2021-conference/speakers-chairs/dr-andrew-barnes/.* Crawford Fund,.

Barnes AC, Silayeva O, Landos M, Dong HT, Lusiastuti A, Phuoc LH, Delamare-Deboutteville J. 2022. Autogenous vaccination in aquaculture: A locally enabled solution towards reduction of the global antimicrobial resistance problem. *Reviews in Aquaculture* 14:907–918. DOI: 10.1111/raq.12633.

Basri L, Nor RM, Salleh A, Md. Yasin IS, Saad MZ, Abd. Rahaman NY, Barkham T, Amal MNA. 2020. Co-Infections of Tilapia Lake Virus, Aeromonas hydrophila and Streptococcus agalactiae in Farmed Red Hybrid Tilapia. *Animals* 10:2141. DOI: 10.3390/ani10112141.

Cagua F, Wilkinson S, Barnes A, Delamare-Deboutteville J. 2021.Lab in a Backpack: Rapid and accurate diagnosis for the control and prevention of diseases in aquatic animals. *Available at https://labinabackpack.com/* (accessed June 15, 2023).

Chadag V, Delamare-Deboutteville J, Beveridge M, Marwaha N. 2021. Reducing disease risks in fish through better detection, management and prevention. *WorldFish. https://worldfishcenter.org/publication/reducing-disease-risks-fish-through-better-detection-management-and-prevention.*

Core Pipeline - artic pipeline. *Available at https://artic.readthedocs.io/en/latest/minion/* (accessed June 15, 2023).

Debnath SC, McMurtrie J, Temperton B, Delamare-Deboutteville J, Mohan CV, Tyler CR. 2023. Tilapia aquaculture, emerging diseases, and the roles of the skin microbiomes in health and disease. *Aquaculture International*. DOI: 10.1007/s10499-023-01117-4.

Delamare-Deboutteville J, Taengphu S, Gan HM, Kayansamruaj P, Debnath PP, Barnes A, Wilkinson S, Kawasaki M, Vishnumurthy Mohan C, Senapin S, Dong HT. 2021. Rapid genotyping of tilapia lake virus (TiLV) using Nanopore sequencing. *Journal of Fish Diseases* 44:1491–1502. DOI: 10.1111/jfd.13467.

Dong HT, Chaijarasphong T, Barnes AC, Delamare-Deboutteville J, Lee PA, Senapin S, Mohan
CV, Tang KFJ, McGladdery SE, Bondad-Reantaso MG. 2023. From the basics to
emerging diagnostic technologies: What is on the horizon for tilapia disease diagnostics?
Reviews in Aquaculture 15:186–212. DOI: 10.1111/raq.12734.
Dong HT, Gangnonngiw W, Phiwsaiya K, Charoensapsri W, Nguyen VV, Nilsen P, Pradeep PJ,
Withyachumnarnkul B, Senapin S, Rodkhum C. 2016. Duplex PCR assay and in situ
hybridization for detection of Francisella spp. and Francisella noatunensis subsp.
orientalis in red tilapia. *Diseases of Aquatic Organisms* 120:39–47. DOI:
10.3354/dao03021.
FAO. 2020. *The State of World Fisheries and Aquaculture 2020: Sustainability in action*. Rome,
Italy: FAO, Rome. DOI: 10.4060/ca9229en.
FAO. 2022. FishStatJ: universal software for fishery statistical time series: aquaculture
production 1950–2020. FAO, Rome.
González-González E, Mendoza-Ramos JL, Pedroza SC, Cuellar-Monterrubio AA, Márquez-
Ipiña AR, Lira-Serhan D, Santiago GT, Alvarez MM. 2019. Validation of use of the
miniPCR thermocycler for Ebola and Zika virus detection. *PLOS ONE* 14:e0215642.
DOI: 10.1371/journal.pone.0215642.
González-González E, Santiago GT, Lara-Mayorga IM, Martínez-Chapa SO, Alvarez MM.
2020. Portable and accurate diagnostics for COVID-19: Combined use of the miniPCR
thermocycler and a well-plate reader for SARS-CoV-2 virus detection. *PLOS ONE*
15:e0237418. DOI: 10.1371/journal.pone.0237418.
Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome
assemblies. *Bioinformatics* 29:1072–1075. DOI: 10.1093/bioinformatics/btt086.
Haenen OLM, Dong HT, Hoai TD, Crumlish M, Karunasagar I, Barkham T, Chen SL, Zadoks
R, Kiermeier A, Wang B, Gamarro EG, Takeuchi M, Azmai MNA, Fouz B, Pakingking
Jr. R, Wei ZW, Bondad-Reantaso MG. 2023. Bacterial diseases of tilapia, their zoonotic
potential and risk of antimicrobial resistance. *Reviews in Aquaculture* 15:154–185. DOI:
10.1111/raq.12743.
Huang Y, Cai S, Jian J, Liu G, Xu L. 2020. Co-infection of infectious spleen and kidney necrosis
virus and Francisella sp. in farmed pearl gentian grouper (♀Epinephelus fuscoguttatus
×♂E. lanceolatus) in China - A case report. *Aquaculture* 526:735409. DOI:
https://doi.org/10.1016/j.aquaculture.2020.735409.
Huang Y, Niu B, Gao Y, Fu L, Li W. 2010. CD-HIT Suite: a web server for clustering and
comparing biological sequences. *Bioinformatics (Oxford, England)* 26:680–682. DOI:
10.1093/bioinformatics/btq003.
Huso D, Wilkinson S, Barnes A, Delamare-Deboutteville J. 2020.Lab-in-a-backpack: Rapid
Genomic Detection to revolutionize control of disease outbreaks in fish farming.
Available at https://fish.cgiar.org/impact/stories-of-change/lab-backpack-rapid-genomic-
detection-revolutionize-control-disease (accessed June 11, 2021).
Karunanathie H, Kee PS, Ng SF, Kennedy MA, Chua EW. 2022. PCR enhancers: Types,
mechanisms, and applications in long-range PCR. *Biochimie* 197:130–143. DOI:
10.1016/j.biochi.2022.02.009.
Kawasaki M, Delamare-Deboutteville J, Bowater RO, Walker MJ, Beatson S, Ben Zakour NL,
Barnes AC. 2018. Microevolution of Streptococcus agalactiae ST-261 from Australia
Indicates Dissemination via Imported Tilapia and Ongoing Adaptation to Marine Hosts

445   or Environment. *Applied and Environmental Microbiology* 84. DOI:
446   10.1128/AEM.00859-18.
447 Kawato Y, Cummins DM, Valdeter S, Mohr PG, Ito T, Mizuno K, Kawakami H, Williams LM,
448   Crane MSJ, Moody NJG. 2021. Development of New Real-time PCR Assays for
449   Detecting *Megalocytivirus* Across Multiple Genotypes. *Fish Pathology* 56:177–186.
450   DOI: 10.3147/jsfp.56.177.
451 Kralik P, Ricchi M. 2017. A Basic Guide to Real Time PCR in Microbial Diagnostics:
452   Definitions, Parameters, and Everything. *Frontiers in Microbiology* 8.
453 Leigh WJ, Zadoks RN, Jaglarz A, Costa JZ, Foster G, Thompson KD. 2018. Evaluation of PCR
454   primers targeting the groEL gene for the specific detection of Streptococcus agalactiae in
455   the context of aquaculture. *Journal of Applied Microbiology* 125:666–674. DOI:
456   10.1111/jam.13925.
457 Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–
458   3100. DOI: 10.1093/bioinformatics/bty191.
459 Liamnimitr P, Thammatorn W, U-thoomporn S, Tattiyapong P, Surachetpong W. 2018. Non-
460   lethal sampling for Tilapia Lake Virus detection by RT-qPCR and cell culture.
461   *Aquaculture* 486:75–80. DOI: https://doi.org/10.1016/j.aquaculture.2017.12.015.
462 Liu X, Sun W, Zhang Y, Zhou Y, Xu J, Gao X, Zhang S, Zhang X. 2020. Impact of Aeromonas
463   hydrophila and infectious spleen and kidney necrosis virus infections on susceptibility
464   and host immune response in Chinese perch (Siniperca chuatsi). *Fish & Shellfish
465   Immunology* 105:117–125. DOI: 10.1016/j.fsi.2020.07.012.
466 López￼Porras A, Elizondo C, Chaves AJ, Camus AC, Griffin MJ, Kenelty K, Barum S,
467   Barquero￼Calvo E, Soto E. 2019. Application of multiplex quantitative polymerase
468   chain reaction methods to detect common bacterial fish pathogens in Nile tilapia,
469   Oreochromis niloticus, hatcheries in Costa Rica. *Journal of the World Aquaculture
470   Society* 50:645–658. DOI: 10.1111/jwas.12576.
471 Mabrok M, Elayaraja S, Chokmangmeepisarn P, Jaroenram W, Arunrut N, Kiatpathomchai W,
472   Debnath PP, Delamare-Deboutteville J, Mohan CV, Fawzy A, Rodkhum C. 2021. Rapid
473   visualization in the specific detection of Flavobacterium columnare, a causative agent of
474   freshwater columnaris using a novel recombinase polymerase amplification (RPA)
475   combined with lateral flow dipstick (LFD) assay. *Aquaculture* 531:735780. DOI:
476   10.1016/j.aquaculture.2020.735780.
477 Machimbirike VI, Jansen MD, Senapin S, Khunrae P, Rattanarojpong T, Dong HT. 2019. Viral
478   infections in tilapines: More than just tilapia lake virus. *Aquaculture* 503:508–518. DOI:
479   10.1016/j.aquaculture.2019.01.036.
480 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
481   *EMBnet.journal* 17:10–12. DOI: 10.14806/ej.17.1.200.
482 Meemetta W, Domingos JA, Dong HT, Senapin S. 2020. Development of a SYBR Green
483   quantitative PCR assay for detection of Lates calcarifer herpesvirus (LCHV) in farmed
484   barramundi. *Journal of Virological Methods* 285:113920. DOI:
485   10.1016/j.jviromet.2020.113920.
486 Paria A, Dong J, Padinhate Purayil SB, Marappan M, Chaudhari A, Thirunavukkarasu AR,
487   Purushothaman C, Kv R. 2016. Evaluation of candidate reference genes for quantitative
488   expression studies in Asian seabass (Lates calcarifer) during ontogenesis and in tissues of
489   healthy and infected fishes. *Indian journal of experimental biology* 54:597–605.

Pomerantz A, Sahlin K, Vasiljevic N, Seah A, Lim M, Humble E, Kennedy S, Krehenwinkel H,
Winter S, Ogden R, Prost S. 2022. Rapid in situ identification of biological specimens via
DNA amplicon sequencing using miniaturized laboratory equipment. *Nature Protocols*
17:1415–1443. DOI: 10.1038/s41596-022-00682-x.
Ramírez-Paredes JG, Paley RK, Hunt W, Feist SW, Stone DM, Field TR, Haydon DJ, Ziddah
PA, Nkansa M, Guilder J, Gray J, Duodu S, Pecku EK, Awuni JA, Wallis TS, Verner-
Jeffreys DW. 2021. First detection of infectious spleen and kidney necrosis virus
(ISKNV) associated with massive mortalities in farmed tilapia in Africa. *Transboundary*
and Emerging Diseases 68:1550–1563. DOI: 10.1111/tbed.13825.
sars-cov-2-ont-artic-variant-calling/COVID-19-ARTIC-ONT. *Available at*
https://workflowhub.eu/workflows/111 (accessed June 15, 2023).
Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q
File Manipulation. *PLOS ONE* 11:e0163962. DOI: 10.1371/journal.pone.0163962.
Shinn AP, Avenant-Oldewage A, Bondad-Reantaso MG, Cruz-Laufer AJ, García-Vásquez A,
Hernández-Orts JS, Kuchta R, Longshaw M, Metselaar M, Pariselle A, Pérez-Ponce de
León G, Pradhan PK, Rubio-Godoy M, Sood N, Vanhove MPM, Deveney MR. 2023. A
global review of problematic and pathogenic parasites of farmed tilapia. *Reviews in*
Aquaculture 15:92–153. DOI: 10.1111/raq.12742.
Soto E, Bowles K, Fernandez D, Hawke JP. 2010. Development of a real-time PCR assay for
identification and quantification of the fish pathogen Francisella noatunensis subsp.
orientalis. *Diseases of Aquatic Organisms* 89:199–207. DOI: 10.3354/dao02204.
Taengphu S, Kayansamruaj P, Kawato Y, Delamare-Deboutteville J, Mohan CV, Dong HT,
Senapin S. 2022. Concentration and quantification of Tilapia tilapinevirus from water
using a simple iron flocculation coupled with probe-based RT-qPCR. *PeerJ* 10:e13157.
DOI: 10.7717/peerj.13157.
The University of Queensland. 2021. Diagnosis in a fish farmer's backpack. YouTube.
https://www.youtube.com/watch?v=pUqmgQ-AuXM.
Waiyamitra P, Tattiyapong P, Sirikanchana K, Mongkolsuk S, Nicholson P, Surachetpong W.
2018. A TaqMan RT-qPCR assay for tilapia lake virus (TiLV) detection in tilapia.
Aquaculture 497:184–188. DOI: https://doi.org/10.1016/j.aquaculture.2018.07.060.
Wang M, Lu M. 2016. Tilapia polyculture: a global review. *Aquaculture Research* 47:2363–
2374. DOI: 10.1111/are.12708.
WorldFish. 2020. A New Concept for Rapid Genomic Detection of Fish Disease. YouTube.
https://www.youtube.com/watch?v=iFxbodO6Fos.
Yang CG, Wang XL, Tian J, Liu W, Wu F, Jiang M, Wen H. 2013. Evaluation of reference
genes for quantitative real-time RT-PCR analysis of gene expression in Nile tilapia
(Oreochromis niloticus). *Gene* 527:183–192. DOI: 10.1016/j.gene.2013.06.013.
527

① Fish sampling for disease diagnostics	② Nucleic acid extraction	③ Multiplex PCR	④ Gel electrophoresis & DNA clean up
30 min	30 min	2 h	30 min

⑧ Disease control in aquaculture.	⑦ Mapping to reference	⑥ MinION sequencing	⑤ Nanopore multiplex Library preparation
	15 min	30 min	90 min

Cq values of the qPCR assay

TiLV	13	11	12	10	12	ud	ud	ud	34	ud	35	35	33	32	32
ISKNV	ud	ud	ud	ud	38	30	33	14	ud	ud	ud	ud	ud	ud	ud
FnO	ud	ud	ud	ud	35	ud	ud	ud	23	14	18	13	ud	ud	ud
SAG	ud	ud	ud	ud	ud	ud	ud	ud	ud	ud	ud	ud	24	25	33

Table 1. Sources of clinical tilapia samples and PCR detection results

Code	Sample_name	Life stage	Country	Year Isolation	Previous Diagnosis sPCR	mPCR	Refs/comments
1	Healthy NT 1	juvenile (NT)	Thailand	2020	-	-	[16]
2	Healthy NT 2	juvenile (NT)	Thailand	2020	-	-	[16]
3	r2_NT F1	fingerling (NT)	Thailand	2020	TiLV (++)	TiLV*	[16]
4	r2_NT F2	fingerling (NT)	Thailand	2020	TiLV (++)	TiLV*	[16]
5	r2_NT F3	fingerling (NT)	Thailand	2020	TiLV (++)	TiLV*	[16]
6	r2_NT F4	fingerling (NT)	Thailand	2020	TiLV (++)	TiLV*	[16]
7	r2_NT F5	fingerling (NT)	Thailand	2020	TiLV (++)	TiLV*	[16]
8	NT fingerling pond C1	fingerling (NT)	Thailand	2020	TiLV (+)	-	[16]
9	NT fingerling pond C2	fingerling (NT)	Thailand		TiLV (+)	-	[16]
10	NT female brood	adult (NT)	Thailand		TiLV (+)	-	[16]
11	NT male brood	adult (NT)	Thailand		TiLV (+)	-	[16]
12	NT juvenile 4	juvenile (NT)	Thailand		TiLV (+)	-	[16]
13	RT F1	fingerling (RT)	Thailand	2015	FnO (+)	-	[20]
14	RT F2	fingerling (RT)	Thailand		FnO (+)	-	[20]
15	RT F3	fingerling (RT)	Thailand		FnO (+)	-	[20]
16	RT F4	fingerling (RT)	Thailand		FnO (+)	-	[20]
17	NT 1.2	fingerling (NT)	Thailand	2015	TiLV (+)	-	[21]
18	NT 1.3	fingerling (NT)	Thailand		TiLV (+)	-	[21]
19	NT 2.1	fingerling (NT)	Thailand		TiLV (+)	-	[21]
20	NT 2.3	fingerling (NT)	Thailand		TiLV (+)	-	[21]
21	NT 3.1	fingerling (NT)	Thailand		TiLV (+)	-	[21]
22	NT 3.3	fingerling (NT)	Thailand		TiLV (+)	-	[21]
23	NT Gh 21	juvenile (NT)	West Africa	2019	ISKNV (++)	ISKNV*	-
24	NT Gh 22	juvenile (NT)	West Africa		ISKNV (++)	ISKNV*	-
25	r2_NT Gh 23	juvenile (NT)	West Africa		ISKNV (++)	ISKNV*	-
26	r2_NT Gh 24	adult (NT)	West Africa		ISKNV (++)	ISKNV*	-
27	r2_NT Gh 25	juvenile (NT)	West Africa		ISKNV (++)	ISKNV*	-
28	RT EX 01	fry (RT)	Thailand	2019	FnO (++)	FnO#	[22]
29	RT EX02	fry (RT)	Thailand		FnO (++)	FnO#	[22]
30	r2_RT EX03	fry (RT)	Thailand		FnO (++)	FnO#	[22]
31	r2_RT FM1a	adult (RT)	Thailand		FnO (++)	FnO#	[22]
32	r2_RT M1	adult (RT)	Thailand		FnO (++)	FnO#	[22]
33	r2_RT FM1b	adult (RT)	Thailand		FnO (++)	FnO#	[22]
34	NT PV F8	adult (NT)	Thailand	2020	Not done	*	-
35	NT PV F9	adult (NT)	Thailand		Not done	#	-
36	NT PV F10	adult (NT)	Thailand		Not done	#	-
37	r3_NT Group B	fry (NT)	Thailand	2019	SAG (++)	SAG	-
38	r3_NT Zone A	fry (NT)	Thailand		SAG (++)	SAG*	-
39	r3_NT Zone B	fry (NT)	Thailand		SAG (++)	SAG	-
40	r1_TILV_1	fingerling (RT)	Thailand	2018-07	TiLV	TiLV	[23]
41	r1_ISKNV_1	fry (AS)	Thailand	2018-11	ISKNV	ISKNV	[24]
42	r1_FNO_1	fingerling (RT)	Thailand	2015	FnO	FnO	[20]
43	r1_SAG_1	fry (NT)	Thailand	2019-07	SAG	SAG	
44	r3_TILV50	fingerling (RT)	Thailand	2019-07	TiLV	TiLV	TiLV PCR amplicon of 100 ng
45	r3_TILV30	fingerling (RT)	Thailand	2019-07	TiLV	TiLV	TiLV PCR amplicon of 75 ng
46	r3_TILV15	fingerling (RT)	Thailand	2019-07	TiLV	TiLV	TiLV PCR amplicon of 50 ng
47	r3_TILV05	fingerling (RT)	Thailand	2019-07	TiLV	TiLV	TiLV PCR amplicon of 25 ng
48	r3_SAG50	fry (NT)	Thailand	2019-07	SAG	SAG	SAG PCR amplicon of 100 ng
49	r3_SAG30	fry (NT)	Thailand	2019-07	SAG	SAG	SAG PCR amplicon of 75 ng
50	r3_SAG15	fry (NT)	Thailand	2019-07	SAG	SAG	SAG PCR amplicon of 50 ng
51	r3_SAG05	fry (NT)	Thailand	2019-07	SAG	SAG	SAG PCR amplicon of 25 ng
52	r1_4PAT_1	N/A	Thailand	N/A	4PAT	4PAT	PCR amplicons 4 PAT (combined1)
53	r1_4PAT_DIL_1	N/A	Thailand	N/A	4PAT	4PAT	PCR amplicons 4 PAT (combined2)
54	r3_4PAT_DIL_3	N/A	Thailand	N/A	4PAT	4PAT	PCR amplicons 4 PAT (combined3)

Notes and abbreviations: sPCR, singleplex PCR; mPCR, multiplex PCR; NT, Nile tilapia; RT, red tilapia; AS, Asian sea bass; -, negative test; ++, high pathogen load; +, low pathogen load; *, suspected dual infections with other pathogen(s); #, probably non-specific products; TiLV, tilapia lake virus; ISKNV, infectious spleen and kidney necrosis virus; FnO, *Francisella noatunensis* subsp. *orientalis*; SAG, *Streptococcus agalactiae*; 4PAT: mixture of PCR amplicons 4 pathogens (TiLV, ISKNV, FnO, SAG). Samples sequenced are italicized and highlighted in grey, and the designation "r1" to "r3" indicates the run number in Nanopore.

Table 2. List of variants for the pathogen gene region, along with the inferred variants and their top 5 NCBI BLAST hits. Pat: Pathogen; Var, Variants; %ID, % nucleotide identity. The samples were grouped based on the sequencing run from which they were generated.

Pat	Samples	Var	Top 5 BLAST hits	%ID	Accession
FnO	r1_4PAT_1,r1_4PAT_DIL_1,r1_FNO_1,r1_SAG_1 r2_NT_F1,r2_NT_F2,r2_NT_F3,r2_NT_F4,r2_NT_F5,r2_NT_Gh23,r2_NT_Gh24,r2_NT_Gh25,r2_RT_EX03,r2_RT_FM1A,r2_RT_FM1B,r2_RT_M1 r3_4PAT_DIL_3	0	*Francisella noatunensis* subsp. *orientalis* strain LPM2-AR2019	100	MN385384.1
			Francisella noatunensis subsp. *orientalis* strain FO371	100	CP022953.1
			Francisella noatunensis subsp. *orientalis* strain FNO364	100	CP022952.1
			Francisella noatunensis subsp. *orientalis* strain FNO222	100	CP022951.1
			Francisella noatunensis subsp. *orientalis* strain FNO215	100	CP022950.1
ISKNV	r1_4PAT_1,r1_4PAT_DIL_1,r1_ISKNV_1 r2_NT_F2,r2_NT_Gh23,r2_NT_Gh24,r2_NT_Gh25	0	ISKNV isolate ISKNV/10	100	MT178422.1
			ISKNV isolate ISKNV/48	100	MT178418.1
			Angelfish iridovirus AFIV-16	100	MK689685.1
			ISKNV isolate M6	100	MK084827.1
			ISKNV isolate SB04	100	KY440040.1
SAG	r1_SAG_1,r1_4PAT_1,r1_4PAT_DIL_1,r1_SAG_1 r3_4PAT_DIL_3,r3_NT_GroupB,r3_NT_ZoneA,r3_NT_ZoneB,r3_SAG05,r3_SAG15,r3_SAG30,r3_SAG50	0	*Streptococcus agalactiae* strain 01173	100	CP053027.1
			Streptococcus agalactiae strain Sag153	100	CP036376.1
			Streptococcus agalactiae strain ZQ0910	100	CP049938.1
			Streptococcus agalactiae strain NJ1606	100	CP026084.1
			Streptococcus agalactiae strain BSE009	100	CP020387.1
TiLV	r1_4PAT_1,r1_4PAT_DIL_1 r3_4PAT_DIL_3,r3_TILV15,r3_TILV30,r3_TILV50	0	TiLV isolate WVL19054 segment 9	100	MN193531.1
			TiLV isolate WVL19031-01A segment 9	100	MN193521.1
			TiLV isolate EC-2012 segment 9	100	MK392380.1
			TiLV isolate Til-4-2011 segment 9	100	KU751822.1
			TiLV AD-2016 Contig 20	100	KU552140.1
	r2_NT_F1,r2_NT_F2,r2_NT_F3,r2_NT_F4,r2_NT_F5	1	TiLV isolate RIA2-VN-2019 segment 9	100	ON376590.1
			TiLV isolate WVL19054 segment 9	98.91	MN193531.1
			TiLV isolate WVL19031-01A segment 9	98.91	MN193521.1
			TiLV strain EC-2012 segment 9	98.91	MK392380.1
			TiLV isolate Til-4-2011 segment 9	98.91	KU751822.1

Abbreviations: FnO, *Francisella noatunensis* subsp. *orientalis*; ISKNV, infectious spleen and kidney necrosis virus; SAG, *Streptococcus agalactiae;* TiLV, tilapia lake virus