Allochthonous groundwater microorganisms affect coastal seawater microbial abundance, activity and diversity

Keren Yanuka-Golub[1,4,5], Natalia Belkin[2], Nurit Weber[1], Meor Mayyani[1], Yehuda Levy[1,3], Itay J. Reznik[3], Maxim Rubin-Blum[2], Eyal Rahav[2†], Yael Kiro[1]

[1]Department of Earth and Planetary Sciences, Weizmann Institute of Science, Rehovot 7610001, Israel

[2]Israel Oceanographic and Limnological Research, National Institute on Oceanography, Haifa, 310220, Israel

[3]Geological Survey of Israel, Jerusalem 96921, Israel

[4]The Institute of Applied Research, The Galilee Society, ShefaAmr 2020000, Israel

[5]The Institute of Evolution (IoE), Haifa University, Haifa, Israel

† Corresponding author: Keren Yanuka-Golub (keren@gal-soc.org) and Eyal Rahav (eyal.rahav@ocean.org.il)

*Current address:

Key Points:

- Groundwater discharge into the coastal zone delivers both nutrients and allochthonous microbes.

- Groundwater microbes interact with seawater populations, by which affecting the delicate autotroph-heterotroph balance.

- Subterranean microbial processes are key drivers of food webs, potentially affecting biogenic carbon fluxes in the ocean and climate.

Abstract

Submarine groundwater discharge (SGD) is a globally important process supplying nutrients and

trace elements to the coastal environment, thus playing a pivotal role in sustaining marine primary

productivity. Along with nutrients, groundwater also contains allochthonous microbes that are

discharged from the terrestrial subsurface into the sea. Currently, little is known about the

interactions between groundwater-borne and coastal seawater microbial populations, and their role

upon introduction to coastal seawater populations. Here, we investigated seawater microbial

abundance, activity and diversity in a site strongly influenced by SGD (*in-situ* observations), and

through laboratory-controlled bottle incubations mimicking different mixing scenarios between

SGD (either ambient or filtered through 0.1 µm/0.22 µm) and seawater. Our results demonstrate

that the addition of <0.1 µm SGD stimulated heterotrophic activity and increased microbial

abundance compared to control, whereas <0.22 µm filtration treatments induced primary

productivity rates and *Synechococcus* growth. Amplicon sequencing of the 16S rRNA gene

showed a strong shift from a SAR11-rich community in the reference SGD-unaffected coastal

samples to a *Rhodobacteraceae*-dominated one in the <0.1 µm treatment, in agreement with their

in-situ enrichment in the SGD field site. These results suggest that despite the significant nutrient

input, microbes delivered by SGD may affect the abundance, activity and diversity of intrinsic

microbes in coastal seawater. Our results highlight the cryptic interplay between groundwater and

seawater microbes in coastal environments, which has important implications for carbon cycling

and climate.

1 Introduction

Submarine groundwater discharge (SGD) is a globally important process, mixing the recirculated seawater inside the sediment and terrestrial groundwater flowing across the seabed with coastal waters (Burnett et al., 2003). This process of water exchange occurs at the subsurface land–ocean interface, termed subterranean estuary, STE (Moore, 1999; Rocha et al., 2021). Various dynamic driving forces, such as terrestrial hydraulic gradients, tides, waves, seasonal and interannual sea level may impact the chemical solute composition of SGD flux into the coastal ocean (Robinson et al., 2018). Thus, SGD acts as a major pathway for delivering terrestrial solutes across the land-ocean interface, and the STE is an important hotspot for biogeochemical reactions (Moore, 2010). This is especially true for ultra-oligotrophic coastal regions such as the southeastern Mediterranean Sea (SEMS), where the effects of the discharge are potentially most prominent (Rahav et al., 2020; Rodellas et al., 2015).

The terrestrial subsurface is one of the largest habitats for microorganisms on Earth (Griebler and Lueders, 2009), where biogeochemical processes, including the turnover of carbon and other nutrients, mineral cycling or pollutant degradation are primarily driven by diverse microbial populations (Probst et al., 2018). Generally, natural groundwater microorganisms live under completely dark conditions, with limited bioavailable organic carbon and phosphorous (Hofmann and Griebler, 2018); thus, they are forced to develop diverse metabolic and physical strategies to survive (Griebler and Lueders, 2009). Recently, *in situ* dark inorganic carbon fixation rates were reported in a carbonate aquifer (Overholt et al., 2022). Estimations were similar to those measured in oligotrophic marine surface waters, indicating chemolithoautotrophs play an important role in subsurface ecosystems processes. Contrary to the relatively stable conditions maintained in inland aquifers, coastal aquifers entail rapid fluctuations and steep physiochemical gradients to which STE microbes need to quickly adapt. This results in a highly diverse microbial community composition along the hydrological continuum of SGD sites, reflecting a mixture of seawater and groundwater-associated taxa (Adyasari et al., 2019; Chen et al., 2020; Degenhardt et al., 2020), as well as microbial functional groups that follow redox gradients (McAllister et al., 2015; Purkamo et al., 2022). Therefore, ubiquity of diverse prokaryotic groups in coastal groundwater ecosystems that are metabolically-active (Miyoshi et al., 2005), or dormant due to scarcity of nutrients and carbon sources in the subsurface habitat (Hofmann and Griebler, 2018) could potentially affect

bacterioplankton communities upon discharge into the marine environment (Lee et al., 2017; Ruiz-
González et al., 2021).

Bacterioplankton community variation across estuarine gradients were shown to be driven
primarily by salinity (Fortunato and Crump, 2011; Herlemann et al., 2011). Adyasari et al. (2020)
also found that salinity was the most decisive variable that shaped the microbial community
composition across surface water samples, while dissolved nitrogen and phosphorus were the
major predictor of community shift within subsurface water samples. At the same study site, higher
microbial diversity was found in coastal porewater in comparison to surface water samples,
consisting of a diverse mixture of fresh water and marine microbes. The presence of
microorganisms transported from external sources that are hydraulically connected show a
bidirectional dispersal of groundwater and marine species (Ruiz-González et al., 2021). It still
remains unknown, however, whether the transported communities tolerate such wide ranges of
conditions to remain active. Nevertheless, at a different study (Adyasari et al., 2019), brackish
porewater samples were more similar to freshwater samples than to saline samples, comprising
mainly taxonomic groups associated with nitrogen transformation (nitrification and
denitrification) in natural water systems. Given that nitrogen cycling is a key pathway in the
subterranean estuary (Erler et al., 2014; Hays and Ullman, 2007; Slomp and Van Cappellen, 2004),
these findings indicate the possibility of active biological transformation of dissolved nitrogen
along the land-water interface.

SGD often exceeds the Redfield ratio and alleviate co-limitation of N and Si in marine
environments (Santos et al., 2021). The examined effect is mostly attributed to the delivery of
dissolved nutrients stimulating the growth of bloom-forming phytoplankton (Garcés et al., 2011;
Lecher et al., 2017, 2015). However, supplements of groundwater to seawater is not always
straightforward and behave in a dose-depended response. For example, Lecher et al., (2015)
observed through incubation experiments that 10% groundwater addition rather than 20-50%
(volumetric) yields the highest positive effect on chlorophyll a concentrations and phytoplankton
abundance, despite the fewer nutrients introduced. Contrary, lower addition volume ratios (1-10%)
did respond at a dose-dependent manner (Lecher et al., 2015). The authors concluded that these
responses were attributed to the elevated nutrients supplied by the SGD, but this was achieved
only to a certain level. This implies a direct biological effect of SGD through the introduction of
subsurface bacterial cells into the coastal environment, which could potentially affect the native

coastal microbiome. Recently it has been reported that STEs are continuously subjected to a bidirectional inoculation of microbial taxa from the marine and fresh groundwater endmembers (Adyasari et al., 2019). However, investigations mainly focused on health implications of the marine environment due to the transport of pathogenic or fecal indicator bacteria through coastal aquifers or sediments, neglecting the ecological implications. Currently, the role of coastal aquifers as sources of microbial diversity and the responses of marine microbial communities to groundwater-derived microorganisms is still unknown.

The occurrence of specific heterotrophic bacterial groups associated with cyanobacteria is dynamic and highly influenced by biotic and chemical factors. For example, Garcés et al., (2011) found in the Mediterranean Sea that the addition of groundwater to marine communities resulted in *Synechococcus* fast growth, due to the nutrient addition. But, this observation was not consistent in other coastal settings, where other phytoplankton groups dominated (Chamberlain et al., 2014). Marine *Synechococcus* are ubiquitous cyanobacteria in the global ocean and is abundant in both estuarine and coastal waters (Dufresne et al., 2008; Huang et al., 2012). Synergistic interactions between *Synechococcus* and alphaproteobacteria heterotrophic bacteria (*Rhodobacteraceae*), gammaproteobacteria (*Alteromonadaceae*) (Dang and Lovell, 2016; Wang et al., 2021; Zheng et al., 2018), and the Cytophaga-Flavobacteria-Bacteroides (CFB) group are fundamental in the marine food web (Buchan et al., 2014). In oligotrophic surface oceans, the three bacterial groups *Prochlorococcus*, marine *Synechococcus*, and heterotrophs in the SAR11 clade often represent more than 50% of free-living bacterial cells (Becker et al., 2019). The abundant SAR11 is a strict oligotrophic heterotroph (Kearney et al., 2021), thus other heterotrophic groups that are adapted to higher organic carbon concentrations such as those associated with phytoplankton blooms are often favored (Buchan et al., 2014; Tada et al., 2011; Teeling et al., 2012). Previous studies showed that the *Roseobacter/Rhodobacter* group of bacteria may respond as generalists because they remain active regardless of large fluctuations in organic matter, whereas the *Alteromonas* group are defined as specialists for using organic matter and became abundant only in the presence of phytoplankton blooms (Tada et al., 2011). Given the delicate autotrophic-heterotrophic metabolic balance in coastal marine ecosystems, SGD is an important factor that should be accounted for with respect to its chemical and biological effects on the ocean.

We recently compared a site strongly influenced by SGD (Achziv, northern Israel) and a nearby nutrients-poor reference site at the oligotrophic Israeli shallow rocky coast (Rahav et al.,

2020). At the SGD site (Achziv), we showed that SGD contributes to high concentrations of dissolved nitrate and silica in comparison to the reference coastal seawater, resulting in elevated *in-situ* phytoplankton biomass and primary productivity. The main objective of this study is to further elucidate the response of marine heterotrophic bacterial taxonomic and functional responses to SGD by addressing the following questions: i) Do coastal microbial communities respond to the nutrient-rich source of SGD in a dose-dependent manner (i.e. higher SGD-enriched mixtures correspond linearly with higher microbial activity and abundance)? ii) How do discharged groundwater microbes shape the activity and composition of the coastal microbiota? To explore these questions, three complementary incubation experiments were designed to illustrate the taxonomic and physiological response upon introducing groundwater and coastal communities. Additionally, porewater samples during three field campaigns were collected to follow composition, abundance and activity of the natural microbial communities at the Achziv SGD site. Our results show that groundwater prokaryotes transported through SGD influence coastal microbial activity and diversity, resulting in different heterotrophic composition patterns. We were able to mimic a similar pattern in bottle incubation experiments by filtering the groundwater prior mixing with coastal seawater. This points at potential competitive interactions for available nutrients in oligotrophic marine environments.

2 Materials and Methods

2.1 Field site, sampling procedures, and sample analyses

Field site: Groundwater and seawater samples were collected from the shore sediments and surface seawater at Achziv (Lat. 33º3'52 N, Lon. 35º6'14.94 E), located along the Northern Israeli Mediterranean shoreline (Figure 1). Achziv coastal aquifer structure is comprised of Pleistocene calcareous sandstone, beach rocks, and carbonate sand, providing a complex groundwater flow system. About five km east of the coastline, the carbonate Galilee Mountains (Judea Group) are elevated. The Quaternary permeable filling of the coastal aquifer is recharged mainly from the east, by precipitation on these carbonate mountains (Kafri and Kessler, 2001; Paldor et al., 2020). Previous studies show fresh SGD (or a fresh SGD component) both shallow along the shoreline (Weinstein et al., 2006) and in deeper parts, through preferential flow in the Judea Group aquifer (Paldor et al., 2020, 2019). The exact SGD sampling locations were selected based on the density

166 and electric conductivity to identify the fresh groundwater and fresh-saline transition zone and to

167 track changes in the salinity distribution compared to previous seasons.

Figure 1: **(a)** Map showing the location of the two sampling areas at the Israeli coastline in the SE Mediterranean Sea: Achziv, the SGD-site (33° 3′52 N, 35° 6′14.94 E) and Shikmona (32° 49′34 N, 34° 57′20 E) within the Haifa Bay (sampled as the reference non-SGD site during the incubation experiments). **(b)** Schematic representation of the field site in Achziv showing the location of the piezometers (brown bars, numbers indicate piezometer ID). Piezometer samples are referred to in the text as either "Low or High salinity porewater" samples (n = 9 and 13, respectively) and are distinguished from the surface seawater samples (n = 11). Accordingly, red dots indicate high salinity (~30 ppt) porewater and blue dots indicate low salinity porewater samples (~17 ppt). **(c)** aerial photography with the piezometers setup for Achziv sampling campaigns. The numbers indicate piezometer ID along a perpendicular (1, 5-6) and parallel (2-4) to the shoreline cross sections.

Sample collection: The fieldwork included three campaigns at Achziv Beach, Israel (August 2020, February and July 2021). Each field campaign lasted 2-5 days and covered at least 2 tidal cycles. Physical parameters were measured, and porewater samples were collected on the shoreline using piezometers (AMS piezometers that reach depths of <2 meters) and a portable peristaltic pump

(Masterflex®, Cole-Parmer, Germany). The density (g cm^{-3}), electric conductivity (mS/cm),

temperature (°C) and pH, of surface seawater, porewater and groundwater were measured on-site

at the time of the sampling. Based on average temperature and water density measurements,

salinity (ppt) was calculated.

Samples for microbial analysis were collected from the piezometers and divided to aliquots:

First, for community analysis, samples were immediately filtered through polycarbonate 0.2 μm

pore size filters (Merck, Israel), which were kept on ice and transported to the laboratory on the

same day. Filter samples were stored frozen (-20°C) until DNA extraction. Second, for Pico-/nano-

phytoplankton and heterotrophic prokaryotic abundance, non-filtered samples were chilled on ice

and transported to the laboratory on the same day.

2.2 Bottle incubation experiments

Three incubation experiments were conducted to determine the contribution of SGD to the

coastal microbial community. The experiments were conducted with five different treatments

(including no-mixing control of reference coastal seawater, i.e., ambient seawater that is not

exposed to SGD) in triplicates as described in Table 1. The first experiment (Exp. 1) was designed

to test the relative contribution of brackish discharged groundwater (salinity = 7.93 ppt) on the

microbial productivity and abundance of reference coastal seawater by mixing different ratios (1,

5, 10 and 20%) of discharged groundwater to reference coastal seawater. Discharged groundwater

was collected into acid-cleaned containers on the day the experiment initiated near Achziv

National Park (33° 3′52 N, 35° 6′14.94 E). At this sampling site, a significant groundwater

discharge was spotted (Rahav et al., 2020; Weinstein et al., 2006). The second and third

experiments (Exp. 2; Exp.3) were designed to extend Exp. 1 and aimed to specifically investigate

how groundwater-derived microorganisms affect the activity and abundance of marine organisms

once discharged into the sea. For these experiments, fresh groundwater (FGW) was collected from

drilling wells and pumped into 20 L acid-cleaned sample-rinsed carboys the same day the

experiment was initiated. At the laboratory, fresh groundwater was either filtered through a 0.1

μm polycarbonate filter (Exp. 2) or serially filtered through 0.22 and 0.1 μm polycarbonate filters

(Exp. 3). The filtrate was added to seawater in different mixing scenarios as described in Table 1.

Table 1: Summary of the three-bottle incubation experimental designs

	Experiment aim	Groundwater source	Control groups No addition		Treatment groups Mixing ratios with ambient seawater (v:v)*			
Exp. 1	Investigate dose-dependent responses	Brackish discharged groundwater (Achziv site)	Reference ambient seawater		1%	5%	10%	20%
Exp. 2	Compare filtered fraction supply of dissolved inorganic nutrients, to ambient groundwater	Inland fresh groundwater	Reference ambient seawater	Fresh groundwater	5%	5% of 0.1 µm filtered FGW		
Exp. 3	Investigate sequentially filtered groundwater and dilution effect (DDW)	Inland fresh groundwater	Reference ambient seawater		5%	5% DDW	5% 0.1 & 0.22µm filtered FGW	

* For all incubation experiments, the % ratio indicates adding groundwater to ambient seawater, unless stated otherwise.

Ambient coastal seawater was collected by pumping at the Israel Oceanographic and Limnological Research Institute (IOLR) into acid-cleaned carboys, and mixed with either brackish groundwater (Exp.1) or fresh groundwater (Exp. 2, Exp. 3) at the desired ratios and filtration size (Table 1). Then, each mixture was randomly distributed into acid-washed transparent polycarbonate Nalgene bottles (4.5 L for Exp. 1 and 2; 250 ml bottles for Exp 3).

The duration of the experiments was 3-5 days, and samples were taken for the following analyses: chlorophyll *a* (Exp. 1 & 2, every 24Hr.), dissolved nutrient concentrations (Exp. 2 & 3 T zero and T final), flow cytometry (bacterial and phytoplankton abundance, every 24Hr.), primary and heterotrophic production rates (Exp. 1 & 2, every 24Hr.; Exp. 3 T zero and T final), methods are specified below. All results were normalized to the dilution factor (for example, in the 20% treatment group, all measurement values were multiplied by 1.2). All bottles were incubated in a flow-through tank at the IOLR through which ambient Eastern Mediterranean water was continuously pumped to maintain surface ocean temperature. The incubation tanks were covered with an illumination net to maintain natural light (representing a full-day cycle).

2.3 Heterotrophic prokaryotic and Pico-/nano-phytoplankton abundance

Samples (1.8 mL) were fixed with glutaraldehyde (final concentration 0.02 % v:v, Sigma-Aldrich G7651), frozen in liquid nitrogen, and later stored at −80°C until analysis. The abundance of autotrophic pico- and nano-eukaryotes, *Synechococcus* and *Prochlorococcus*, and other

heterotrophic prokaryotes (bacteria and archaea) was determined using an Attune® Acoustic

Focusing Flow Cytometer (Applied Biosystems) equipped with a syringe based fluidic system and

488 and 405 nm lasers. To measure heterotrophic prokaryote abundance (hereafter refer to as

heterotrophic abundance), a sample aliquot was stained with SYBR Green (Applied Biosystems).

2.4 Heterotrophic productivity

Prokaryotic (bacteria and archaea) heterotrophic production was estimated using the ^{3}H-

leucine incorporation method (Perkin Elmer, specific activity $100\,\mathrm{Ci\,mmol^{-1}}$). Water samples

(1.7 mL each) were incubated in the dark (wrapped in aluminum foil) with ~ 10 nmol hot leucine

$\mathrm{L^{-1}}$ for 4 h (Rahav et al., 2019). For control treatments, an additional sample was immediately

added with $100\,\mu\mathrm{L}$ of 100 % trichloroacetic acid (TCA, 4°C) along with ^{3}H-leucine, and was

carried out in triplicates. The incubations were terminated with TCA and were later processed

following the micro-centrifugation technique (Smith and Azam, 1992) and added with 1 mL of

scintillation cocktail (Ultima-Gold). The samples were counted using a TRI-CARB 2100 TR

(Packard) liquid scintillation counter.

2.5 Primary productivity

Photosynthetic carbon fixation rates were estimated using the ^{14}C incorporation method

(Nielsen, 1952). Briefly, collected water samples (50 mL) were immediately spiked with 5 µCi of

NaH^{14}CO3 (Perkin Elmer, specific activity $56\,\mathrm{mCi\,mmol^{-1}}$). The samples were incubated for 24 h

under in situ natural illumination. The incubations were terminated by filtering the spiked seawater

through GF/F filters (Whatman, 0.7 µm pore size) at low pressure (~ 50 mmHg). Measurements

for dark controls were also performed and were removed from the sample reads. The filters were

placed overnight in 5 mL scintillation vials containing 50 µL of 32 % hydrochloric acid to remove

excess ^{14}C, after which 5 mL of scintillation cocktail (Ultima-Gold) was added. Radioactivity was

measured using a TRI-CARB 2100 TR (Packard) liquid scintillation counter.

2.6 Inorganic nutrients determination

Samples from the field were immediately filtered through 0.2 µm pore size filters into acid-

washed plastic vials, chilled on ice and transported to the laboratory on the same day. Samples

were stored frozen until analysis. Experimental samples at time-zero and final time point were

filtered through 0.2 μm pore size filters into acid-washed plastic scintillation vials, immediately

frozen (− 20°C) and kept frozen until analysis. Dissolved nutrients ($NO_2 + NO_3$ = NOx, PO_4, and

$Si(OH)_4$) were analyzed in the Interuniversity Institute for Marine Sciences (IUI, Eilat) using

QuikChem 8000 flow injection analyzer (Lachat Instruments, Milwaukee, USA). The

measurement is based on a color reaction created by each of the nutrients with its unique reagent

to create a color complex with a wavelength in the visible light range, which is absorbed by the

device's spectrophotometer.

2.7 DNA extraction, amplicon sequencing, and analysis of bacterial community

Seawater or groundwater (4-5 L) were filtered using a peristaltic pump onto polycarbonate

membrane filters (0.1 or 0.2 μm, 47 mm, Merck, Israel) and stored at −20°C until extraction (n=1

per sample except in selected samples where n=4). After thawing, each filter was cut into small

pieces using a sterile scalpel blade, which was placed immediately into PowerSoil DNA bead tubes

and extracted with the dNeasy PowerSoil Kit (Qiagen, USA) following the standard protocol. To

disrupt the cells, the FastPrep-24™ Classic (MP Biomedicals, USA) bead-beating was used (2

cycles at 5.5 m sec–1, with a 5 min interval).

To generate 16S rRNA gene libraries, the V3–V4 hypervariable region of the 16S gene was

amplified (∼ 465 bp) using the 341F/806R primers containing CS1 and CS2 linkers –

ACACTGACGACATGGTTCTACACCTACGGGNGGCWGCAG   and

TACGGTAGCAGAGACTTGGTCTGG ACTACHVGGGTWTCTAAT. PCR conditions were as

follows: initial denaturation at 95°C for 5 min, 28 cycles of denaturation (95°C for 30 sec),

annealing (50°C for 30 sec) and extension (72°C for 60 s). Sequences were obtained on the

Illumina MiSeq platform in a 2×250 bp paired-end run (HyLabs Israel). Demultiplexing and

assignment of the amplicon sequence variants (ASVs) were performed using the Qiime2 (v.

2020.2) (Bolyen et al., 2019) platform, and DADA 2 was used as a quality control for chimeric

reads (Callahan et al., 2016). The two 250-bp paired-end sequences were merged to obtain a single

read (approximately 416.27 bp, mean length). Taxonomic assignment of the ASVs was done with

q2_feature_classifier (Bokulich et al., 2018) using the Silva database (release 138-99) for 16S

rRNA gene sequencing at 99% identity (Quast et al., 2012).

Alpha diversity and variance calculations were performed using Mothur (v.1.48.0). ASV data table

was filtered (removing ASVs that appear in total <10 times) and repeatedly subsampled to the

lowest number of reads among all samples (12,538 reads x 100 times). Diversity indices included

Invsimpson, Shannon and Chao. Beta diversity was analyzed using Bray–Curtis metrics of

community dissimilarity and plotted using pCoA ordination with PAST v4.09 (Hammer et al.,

2001). Analysis of similarity (ANOSIM) (9999 permutations) was used to make pairwise

comparisons between the different samples' types (Field: Low and high-salinity porewater, Achziv

surface seawater and experimental treatment groups: control ambient coastal seawater (SGD-non

influenced), non-filtered, and 5% filtered 0.1μm pore size) treatments.

3. Data Analysis

3.1 Statistical analyses

Statistical analysis was performed in GraphPad Prism 5.03 (GraphPad, La Jolla, CA, USA).

One-way analysis of variance (ANOVA) was used to determine statistical differences between the

control and different mixing treatments for the three-bottle incubation experiments, and the three

field campaign sample types. Multiple pairwise comparisons were achieved using Tukey 95%

confidence intervals.

3.2 Accession numbers: Raw data from Illumina MiSeq sequencing of 16S rRNA gene

amplicons have been deposited in the Sequence Read Archive (SRA) of NCBI under accession

numbers.

4 Results and Discussion

4.1 Groundwater discharge enhances coastal microbial abundance but constrains activity

Three bottle incubation experiments were designed to independently study the microbial response of groundwater additions (brackish or fresh) to ambient coastal seawater. Specifically, the aim of the first experiment (hereafter, Exp. 1) was to investigate the response of ambient coastal microorganisms (unexposed to SGD and corresponding microbes) to SGD by additions of discharged groundwater at pre-determined volumetric percentages (1%, 5%, 10%, 20%) relative to ambient seawater (Figure 2). The chosen volumetric range aimed to demonstrate naturally occurring conditions that could occur in an SGD site concerning point-source discharge flux. Temporal measurements indicated that a ~40-Hr incubation period yielded maximal production rates, while longer times resulted in saturation of heterotrophic and primary production rates (Supplementary data, S1). Thus, the following results are presented for the 42-Hr incubation time point. Results showed that heterotrophic prokaryotic abundance increased in a dose-dependent manner, ranging from $4.8 \times 10^5 \pm 0.1 \times 10^5$ cells ml^{-1} in the control to $19 \times 10^5 \pm 1.4 \times 10^5$ cells ml^{-1} in the 20% GW treatment (Figure 2A). Contrary, heterotrophic production rates reached a plateau already at 1% GW enrichment; 11.8 ± 2.4 µg C L^{-1} day^{-1}, while the control values were ~6 fold lower (Figure 2B). Primary production rates (Figure 2C) and chlorophyll *a* concentrations (Supplementary data, S1), reached a plateau at 5% treatment, in agreement with an earlier study from the intertidal area affected by SGD in Monterey Bay (Lecher et al., 2015).

Figure 2: Scatter dot plots of surface seawater (control) prokaryote abundance (a), heterotrophic production rate (b) and primary production rate (c) following dilution with discharged brackish groundwater (1-20% v:v) or un-amended seawater (control). Results are shown for the 2nd time point of the experiment (42hr) as described in the text. Lowercase letters indicate significant differences between treatments (using ANOVA followed by Tukey posthoc tests, $p \leq 0.05$). The temporal variability of all time points is shown in Supplementary Information (S1). The dilution factor was calculated for each treatment to account for the volume of groundwater added to ambient seawater.

335 These observations suggest a further interaction between the groundwater microbes and the
336 seawater microbes, which limits activity, but not abundance, despite the addition of nutrients.
337

4.2 Groundwater microbes transported through SGD may limit coastal production rates
To assess the microbial contribution of SGD to heterotrophic production, a second experiment was
designed (hereafter, Exp. 2, Table 1), in which filtered (<0.1 μm) and non-filtered (5 and 20%)
groundwater samples were mixed with the reference ambient coastal seawater. Amendment of
filtered groundwater presented primarily the addition of dissolved nutrients, whereas non-filtered
groundwater supplied nutrients along with endogenous groundwater microbes. Temporal
measurements indicated that a ~67-Hr incubation period yielded maximal production rates
(Supplementary data, S2). Thus, the following results are presented for the 67-Hr incubation time
point (Figure 3, Supplementary Table S1). Natural SGD-affected coastal measurements from
Achziv were also evaluated (samples were collected at two different seasons: February 2021 and
June 2022, Table 2). Compared to non-filtered groundwater and the control treatments, the filtered
treatment showed the highest change for prokaryote abundance (Figure 3A) and heterotrophic
production rates (Figure 3B). While prokaryote abundance of Achziv surface seawater was similar
to the filtered treatment, heterotrophic production was significantly lower (similar to non-filtered
treatments and control). Contrary, no changes in primary production rates were observed between
the different groundwater amendment treatments (Figure 3C). These rates were significantly
higher than the control (ANOVA and Tukey posthoc tests, $p \leq 0.05$), but slightly lower than Achziv
surface seawater. These findings lie in agreement with the results of experiment 1, where
prokaryote abundance is dose dependent and activity reaches a stable threshold, supported by the
natural Achziv measurements (Figure 3). Furthermore, changes in concentration of phosphate
(PO_4), silica ($Si(OH)_4$), nitrate (NO_3), and ammonium (NH_4) over the incubation experiment
period indicated that mainly PO_4 and $Si(OH)_4$ were uptaken by the microbial community provided
by groundwater amendments (Supplementary data, S3). The 20% non-filtered treatment utilized
significantly more of these nutrients than the other filtered and non-filtered treatments
(Supplementary data, S3), as observed in a previous study (Lecher et al., 2015).
The higher increase in prokaryote abundance or heterotrophic production in the filtered vs. the
non-filtered addition suggests that groundwater microbes affect the seawater communities and

potentially outcompete them for the added nutrients, highlighting the intricate interactions between

autotrophic and heterotrophic microbes in oligotrophic realms such as the Mediterranean coast.

Figure 3: Scatter dot plots of surface seawater prokaryote abundance (a), heterotrophic production (b) and primary

production (c) following dilution with fresh groundwater (5% and 5% filtered 0.1 μm v:v), un-amended seawater

(control SW) or fresh groundwater (Control GW) during Experiment 2. Results are shown for the 3rd time point of the

experiment (67hr) as described in the text. Surface seawater measurements from Achziv (SGD-site) were also added

to this analysis. Lowercase letters indicate significant differences between treatments (using ANOVA followed by

Tukey posthoc tests, $p \leq 0.05$). The temporal variability of all time points is shown in the Supplementary Information

(S2). The dilution factor was calculated for each treatment to account for the volume of groundwater added to ambient

seawater.

The control fresh groundwater (collected from an inland aquifer), incubated without mixing,

showed the lowest values of abundance and productivity (Figure 3). While lower phototrophic

abundance and activity are negligible, as expected in subsurface environments, the lower

prokaryotic abundance ($8.6\pm2.4\times10^4$ cells/ml) and heterotrophic productivity (6.74 ± 3.0 μg C L^{-1}

day^{-1}) are in agreement with values reported for pristine inland fresh groundwater. These

observations show that a large fraction of communities from inland aquifers are locally inactive

(Griebler and Lueders, 2009). Despite their low numbers and activity rates, incubation experiments

of groundwater communities showed that groundwater bacteria can respond rapidly to

environmental changes as implied from our data as well, and grow even when mixed with marine

waters (Adyasari et al., 2019). This could ultimately shape coastal microbial diversity and activity.

4.4 Groundwater microbes transported through SGD may shape coastal community composition

Three sampling campaigns (August 2020, February 2021 and July 2021) were conducted at a field site, highly influenced by SGD (Achziv, northern Israel). Porewater samples were collected on the shoreline using piezometers at different depths (0.25-1m, Figure 1). According to the recorded porewater density values, the porewater samples were divided into two groups: low and high salinity. In addition, we sampled surface seawater (Table 2). The measured parameters of these samples were compared to values obtained for both Achziv and the reference site (Shikmona) during the routine monitoring program by Israel Oceanographic and Limnological Research, IOLR (Table 2).

Table 2: Summary of environmental parameters averaged by sample groups

Sample source	Group	# samples (n)	Salinity (ppt)	pH	N:P (μM)	Si (μM)
Environmental **Achziv study site.** Samples from 3 sampling campaigns.	Low (porewater)	12	17.3 *a*	7.320±0.1407 *a*	270.9±103.8 *a*	193.9±88.12 *a*
	High (porewater)	16	29.2 *b*	7.473±0.1919 *b*	225.6±70.80 *a*	122.5±75.47 *b*
	Achziv surface seawater	4	41.2 *b*	7.970 ± 0.1186 *c*	49.62±38.73 *a*	10.00±1.00 *c*
Fresh groundwater inland well	GW	4	0.6 *c*	7.251 ± 0.1774 *a*	238.9±286.0 *a*	254.5±44.4 *a*
National monitoring, SEMS, 2020-2021 (surface seawater)	Reference site (Shikmona)	24	39.16 *b*	8.120±0.0438 *c*	23.10±27.29 *b*	1.547±0.8062 *c*
	Achziv	24	38.97 *b*	8.052±0.0345 *c*	281.8±223.2 *a*	2.744±1.142 *c*

Lowercase letters indicate significant differences between groups (using ANOVA followed by Tukey post-hoc tests, $p \le 0.05$).

The measured parameters reported for Achziv surface seawater during the three field campaigns were comparable to the ongoing monitoring results obtained during the years 2020-2021 (salinity, pH, and $Si(OH)_4$, ANOVA, $p > 0.05$) with no significant seasonal or annual differences. As expected, the low-salinity porewater samples showed significantly lower salinity/pH values compared to high-salinity porewater and surface water (Table 2), further supporting the contribution of freshwater inputs. The mean N:P ratio of both high- and low-salinity porewater samples were comparable to values obtained during routine monitoring of the Achziv site for the last two years. The N:P values obtained at all Achziv samples were significantly higher than in the Shikmona reference site, signifying the influence of SGD as an important nutrient source. Notably,

Si(OH)$_4$, PO$_4$ and NO$_3$ concentrations of porewater samples exceeded the surface seawater

samples by an order of magnitude (Table 2), and lie on a conservative mixing line with water

salinity (Supplementary data, S4). Higher PO$_4$ concentrations were observed in all Achziv

porewater samples compared to the reference site, ranging from 1.0–3.7 µM, but generally peaked

at the lower-salinity samples (2.9±0.6 µM).

Sequencing of the 16S rRNA gene resulted in 1,333,846 quality sequences, which clustered into

2235 ASVs, including field campaigns and bottle incubation experiments. The number of ASVs

per sample varied between 36 and 493. Microbial community diversity of samples collected in the

Achziv field site (low and high-salinity porewater and surface seawater) was compared with those

obtained from the experimental treatments, namely control seawater (ambient seawater, not

influenced by SGD), non-filtered treatments (1%, 5%, 20%), filtered (5% 0.1 µm pore size) and

fresh groundwater from inland wells. Among these, the low-salinity porewater group was the most

diverse for the three alpha diversity indices investigated (Figure 4), indicating a high heterogeneity

of the low-salinity microbial community, and a relatively even community inferred from the higher

inverse Simpson index (Figure 4c).

Figure 4: Box-Whisker plots showing the interquartile range (25th to 75th percentile) and the median value

(horizontal line within the box) of the microbial communities' alpha diversity grouped according to field and

experimental samples assessed by the Chao richness estimating the number of ASVs (a), Shannon (b) and inverse

Simpson indices (c). Lowercase letters indicate significant differences between treatments (using ANOVA followed

by Tukey posthoc tests, p ≤ 0.05).

PCoA ordination analyses performed on the entire microbial community data (based on the Bray-Curtis distance) showed that samples separated into four groups according to salinity and experimental treatments (Figure 5; ANOSIM $p < 0.001$), where some dispersal appeared between sampling days (for the field campaigns). This implies that the microbial community in Achziv porewater is relatively stable annually and diversity is primarily influenced by water salinity. The low-salinity porewater ($n = 9$) was the most distinctive compared to the other groups of the high-salinity porewater ($n = 13$), Achziv surface water ($n = 11$), and the experimental treatments: non-filtered experimental treatment ($n = 11$) and seawater control group ($n = 4$). The exceptions were the 5% 0.1 μm filtered treatment ($n = 3$; $p > 0.05$) and fresh groundwater samples ($n = 3$; $p > 0.05$). Overall, the PCoA plot was divided into four main quarters along both axes, each representing different water characteristics (Figure 5). While, the left half of the PCoA2 axis contained samples originating from fresh groundwater and low-salinity porewater, the right half was dominated by high-salinity porewater and surface seawater, including samples from Achziv and the experimental treatments. During the field campaign in February 2021, two surface seawater samples were collected at different distances from the shore (100m and 600m). It is apparent from Figure 5 that the high-salinity porewater community composition in the aquifer was more similar to that of the near-shore (100m) Achziv surface seawater than to that of the open seawater (600m from the shore). These communities possibly affect each other, creating a new environment across the seawater-aquifer transect. For the same field sampling campaign, we measured surface seawater and porewater microbial activity (primary and heterotrophic production) and prokaryotic abundance. We observed a substantial decrease in all parameters between the conservative mixing line and seawater (Supplementary data, S5). These findings illustrate the change in the environment between open seawater and groundwater, which both prevents day light and introduces new communities, which may limit productivity.

Figure 5: Principal coordinate analysis (PCoA) of prokaryote composition (Bray-Curtis distances) based on 16S-amplicon sequencing of samples collected during three sampling campaigns (different colors as indicated) at Achziv site, grouped according to porewater salinity (Low/High) and Surface seawater. The Achziv surface seawater samples collected in February 2021 were divided according to distance from shore (100 and 600m). Additionally, experimental treatment samples (control seawater, non-filtered, 5% filtered and fresh groundwater) are plotted.

Achziv bacterial community composition (surface seawater) was co-dominated by three marine family-level taxonomic groups (relative abundance > 40%), including the autotrophic *Synechococcus* and heterotrophs (Figure 6, Supplementary Table S1). The three heterotrophic bacterial families were *Rhodobacteraceae*, *Actinomarinaceae* and SAR11 Clade. While *Actinomarinaceae* was the most abundant bacterial plankton in all surface seawater samples, including the experimental control seawater (ambient seawater, not influenced by SGD), Achziv surface seawater and high-salinity porewater, the SAR11 clade was primarily abundant in the control seawater and non-filtered experimental treatments, Figure 6a &b, respectively. On the other hand, the lowest relative abundance of *Rhodobacteraceae* was detected in experimental control seawater, comparable with fresh groundwater samples. Thus, from ambient coastal seawater to SGD-coastal samples (Achziv), the community composition shifted from a SAR11-dominated to a *Rhodobacteraceae*-dominated community. Notably, the relative abundance of

Rhodobacteraceae and *Alteromonadaceae* were the highest in the filtered treatment (Figure 6c & d, respectively). Previous studies have linked the presence of these marine microbial taxa in coastal aquifers as potential indicators of seawater intrusion, indicating an active land-ocean interface (Chen et al., 2019; Unno et al., 2015).

Figure 6: Scatter dot plots of the four dominant bacterial groups (relative abundances): (a) *Actinomarinaceae* (b) SAR11 (c) *Rhodobacteraceae* (d) *Alteromonadaceae* (e) the autotrophic bacterium *Synechococcus* in Achziv field samples (samples grouped according to water-salinity) and incubation experiments (samples grouped according to treatment). (f) *Synechococcus* relative abundances in incubation samples, differentiating between the different treatments. Lowercase letters indicate significant differences between treatments (using ANOVA followed by Tukey posthoc tests, $p \leq 0.05$). * indicates significant difference between two treatments ($p < 0.05$), where the rest are non-significant. Data derived from 16S rDNA analyses.

Based on the SAR11 to *Rhodobacteraceae* shift and the substantially higher relative
abundance of *Rhodobacteraceae* in the filtered treatment, we suggest that applying the filtration
resulted in a unique scenario, where the typical-oligotrophic coastal microbial community,
growing under constant nutrient-limiting conditions, was exposed to a nutrient-rich solution almost
without potential competitors.

Our results suggest that mixing non-filtered groundwater with ambient coastal seawater
induced competitive interactions between the two different communities, eventually affecting
primary and heterotrophic production rates. This competitive effect was essentially eliminated
through filtration because most microbes were removed from the water, relieving the nutrient
limitations of the oligotrophic coastal seawater. Although the transport of microbial diversity into
the coastal ocean through SGD has been addressed mostly with regard to pathogenic or fecal
indicator bacteria (Vollberg et al., 2019; Yau et al., 2014), our results emphasize that marine
heterotrophic bacterioplankton respond taxonomically and functionally to SGD-derived microbes.
Typically, low-salinity porewater and fresh groundwater samples were similar to each other in
terms of low relative abundance of the marine bacteria found in the other samples (Figure 6a, b, c)
and β-diversity distance (ANOSIM, $p > 0.05$). Moreover, the higher alpha diversity values in the
low-salinity porewater samples (Figure 4) are explained by a much more heterogeneous microbial
community, which consisted of a diverse mixture of freshwater and marine microbes. These
include the prevalent groundwater clades Patescibacteria and Woesearchaeales (Herrmann et al.,
2019; Tian et al., 2020). Patescibacteria was found in low abundances in Achziv surface seawater,
and absent in the control ambient samples (Supplementary data, Table S2). This hints at the
potential role of connectivity between terrestrial and marine hydrospheres driven by SGD in the
Achziv site.
Synechococcus was the dominant cyanobacterium detected in all surface seawater and high-
salinity porewater samples (Figure 6e). The lowest abundance was associated with the low-salinity
porewater and groundwater samples, probably due to the different environmental conditions,
primarily the lack of light in the aquifer. Moreover, the highest groundwater addition treatment
(20%) had the lowest *Synechococcus* relative abundance when compared to the other lower ratios
of non-filtered treatments (Figure 6f), suggesting larger volumes of groundwater inhibit
Synechococcus growth. However, this was less significant based on cell number counts
(Supplementary Table S1).

The third experiment (hereafter Exp. 3) was designed to extend on the previous two experiments

and further examine the response of autotrophic communities to different microbial cell size

fractions by mixing sequentially filtered groundwater (non-filtered, 0.22 & 0.1 μm pore size) with

the control ambient seawater, Figure 7. Also here, heterotrophic production and prokaryote

abundance increased when 0.1 μm filtered groundwater was mixed with the control seawater.

However, the 0.22 μm filtration treatment did not exert such a response on prokaryote abundance

and heterotrophic production, rather induced primary productivity rates, specifically associated

with *Synechococcus* abundance, after a 40-hour incubation (Figure 7a &b). Cell abundances

dropped for all treatments after a 65-Hr incubation period, but even the delayed production rates

measured only for the last time period (65Hr) were still showing a significant response, which

supported the abundance trend observed after a 40-Hr incubation period. The DDW treatment was

included to demonstrate that groundwater mixing effects occur as a result of terrestrial-coastal

microbial interactions, and not due to simple volumetric dilution of the seawater with fresh water.

4.5 SGD affects the interplay between *Synechococcus* and heterotrophic bacteria

In our two investigated sites (SGD and control), *Synechococcus* was the dominant cyanobacterium

based on 16S data. Although *Synechococcus* relative abundance was similar for both sites, we

observed marked differences in the composition of heterotrophic bacteria, specifically SAR11 and

Rhodobacteraceae groups (Figure 7c). The strict oligotrophic SAR11 clade are highly abundant

in the open sea, and were found to associate with Cyanobacteria primarily under poor dissolved

organic carbon and nutrient conditions (Kearney et al., 2021). Alternatively, *Rhodobacteraceae* is

much more metabolically flexible (Xia et al., 2021), allowing this group to outcompete and

dominate other heterotrophs upon groundwater discharge.

Therefore, heterotrophic bacterioplankton largely determine carbon biogeochemical cycling in the

ocean through respiration and biomass production, and their composition appears to be affected

by SGD (Kaile'a and Wiegner, 2016). These results suggest that *Synechococcus* abundance can

increase similarly in the two sites (SGD-enriched and oligotrophic), but primary production

activity is limited, maintaining a balanced ecological ecosystem (perhaps as an intrinsic control of

harmful-algal blooming). Currently, the mechanistic role SGD plays in maintaining ecosystem

stability remains unknown, and how the groundwater microbiota drives this effect, particularly

concerning different microbial cell size fractions.

Figure 7: a) Scatter dot plots of heterotrophic and primary production rates following dilution with fresh groundwater (5%, 5% filtered 0.22 μm and 5% filtered 0.1 μm v:v), un-amended seawater (SW) or diluted with 5% distilled water (5% DDW) during Experiment 3. Results are shown for the final time point of the experiment (only initial and final measurements were conducted for this experiment). Lowercase letters indicate significant differences between treatments (using ANOVA followed by Tukey posthoc tests, $p \leq 0.05$). b) Temporal variability of prokaryote and *Synechococcus* abundance detected (data derived through flow cytometry); */** indicate significant difference between two treatments ($p < 0.05$ or $p < 0.01$, respectively), where the rest are non-significant. c) Bar plot showing relative sequence abundance of dominant microbial taxa at order level for both coastal sites

4.6 potential effect on coastal heterotrophic autotrophic balance and carbon cycling in the ocean

Although freshwater SGD was found to be minor compared to freshwater river fluxes to the ocean (Luijendijk et al., 2020; Zhou et al., 2019), saline and brackish water fluxes are much more significant (e.g. Beck et al., 2013; Kwon et al., 2014; Santos et al., 2021). Our results show how SGD shape the heterotrophic community diversity and regulate activity through discharged microbes. These observations have significant implications for the coastal and ocean carbon budget. Carbon fixation by autotrophs and heterotroph respiration shape the carbon budget in the ocean, affecting directly atmospheric CO_2 through the ability to bury organic carbon (Kirchman, 2012). Estimating chemolithoautotroph fixation rates in STE ecosystems are also important given the notable inorganic carbon fixation rates recently reported in a carbonate aquifer (Overholt et al., 2022). Climate change, causing global warming and ocean acidification have shown to enhance heterotrophic activity (Duarte et al., 2013), which may affect the balance between autotrophic carbon fixation and heterotrophic recycling, thus the biogenic carbon fluxes in the ocean. Our

study suggests that this relationship may be more complex through changes in SGD fluxes due to sea level changes. An increase in sea level will reduce SGD fluxes (Kiro et al., 2008), and thus affecting the heterotrophic community in seawater, as concluded in this study. This may have direct implications on the delicate balance between heterotrophs and autotrophs, and the biogenic carbon fluxes.

5 Conclusions

SGD is an important source of nutrients in coastal environments, especially in oligotrophic environments. In this study we shed light on the complex contribution of SGD on the ecology of coastal microbial communities through the transport of groundwater-borne prokaryotes. Diverse microbial communities, naturally inhabiting the subterranean estuary, shape coastal microbial abundance, taxonomic diversity and activity. This ultimately affects photoautotrophic-heterotrophic specific interactions, marine food webs and biogeochemical cycles of the entire coastal ecological system. In our current changing climate world, these interactions are crucial for understanding the possible effects and feedback mechanisms on carbon cycling and climate. Further research is needed to integrate the global chemical-biological effect of SGD on marine environments, focusing, for example, on STE sediment microbial dynamics along redox gradients at the terrestrial–marine interface.

Acknowledgments

We thank to the Sustainability and Energy Research Initiative (SAERI) grant, the Helen Kimmel Center for Planetary Science grant and the De Botton center for Marine Sciences grant for their funding supporting this study. This study was supported by the, Zuckerman Faculty Scholars program, a research grant from the Center for Scientific Excellence, a research grant from the Raymond Lapon Fund, and the Estate of David Levinson, a research grant from Paul and Tina Gardner, and a research grant from the Center for New Scientists at the Weizmann Institute of Science. We thank H. Michael and I. Halevy for fruitful discussions and suggestions.

References

Adyasari, D., Hassenrück, C., Montiel, D., Dimova, N., 2020. Microbial community

composition across a coastal hydrological system affected by submarine groundwater

discharge (SGD). PLoS One 15, e0235235.

Adyasari, D., Hassenrück, C., Oehler, T., Sabdaningsih, A., Moosdorf, N., 2019. Microbial

community structure associated with submarine groundwater discharge in northern Java

(Indonesia). Sci. Total Environ. 689, 590–601.

https://doi.org/10.1016/j.scitotenv.2019.06.193

Beck, A.J., Charette, M. a., Cochran, J.K., Gonneea, M.E., Peucker-Ehrenbrink, B., 2013.

Dissolved strontium in the subterranean estuary – Implications for the marine strontium

isotope budget. Geochim. Cosmochim. Acta 117, 33–52.

https://doi.org/10.1016/j.gca.2013.03.021

Becker, J.W., Hogle, S.L., Rosendo, K., Chisholm, S.W., 2019. Co-culture and biogeography of

Prochlorococcus and SAR11. ISME J. 13, 1506–1519.

Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., Huttley, G.A.,

Gregory Caporaso, J., 2018. Optimizing taxonomic classification of marker-gene amplicon

sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6, 1–17.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A.,

Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., 2019. Reproducible, interactive,

scalable and extensible microbiome data science using QIIME 2. Nat. Biotechnol. 37, 852–

857.

Buchan, A., LeCleir, G.R., Gulvik, C.A., González, J.M., 2014. Master recyclers: features and

functions of bacteria associated with phytoplankton blooms. Nat. Rev. Microbiol. 12, 686–

698.

Burnett, W.C., Bokuniewicz, H., Huettel, M., Moore, W.S., Taniguchi, M., 2003. Groundwater and pore water inputs to the coastal zone. Biogeochemistry 66, 3–33. https://doi.org/10.1023/B:BIOG.0000006066.21240.53

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods 13, 581–583.

Chamberlain, S.D., Kaplan, K.A., Modanu, M., Sirianni, K.M., Annandale, S., Hewson, I., 2014. Biogeography of planktonic and benthic cyanobacteria in coastal waters of the Big Island, Hawai'i. FEMS Microbiol. Ecol. 89, 80–88.

Chen, L., Hu, B.X., Dai, H., Zhang, X., Xia, C.-A., Zhang, J., 2019. Characterizing microbial diversity and community composition of groundwater in a salt-freshwater transition zone. Sci. Total Environ. 678, 574–584.

Chen, X., Ye, Q., Sanders, C.J., Du, J., Zhang, J., 2020. Bacterial-derived nutrient and carbon source-sink behaviors in a sandy beach subterranean estuary. Mar. Pollut. Bull. 160, 111570.

Dang, H., Lovell, C.R., 2016. Microbial surface colonization and biofilm development in marine environments. Microbiol. Mol. Biol. Rev. 80, 91–138.

Degenhardt, J., Dlugosch, L., Ahrens, J., Beck, M., Waska, H., Engelen, B., 2020. Seasonal dynamics of microbial diversity at a sandy high energy beach reveal a resilient core community. Front. Mar. Sci. 7, 573570.

Duarte, C.M., Regaudie-de-Gioux, A., Arrieta, J.M., Delgado-Huertas, A., Agustí, S., 2013. The oligotrophic ocean is heterotrophic. Ann. Rev. Mar. Sci. 5, 551–569.

Dufresne, A., Ostrowski, M., Scanlan, D.J., Garczarek, L., Mazard, S., Palenik, B.P., Paulsen,

I.T., de Marsac, N.T., Wincker, P., Dossat, C., 2008. Unraveling the genomic mosaic of a

ubiquitous genus of marine cyanobacteria. Genome Biol. 9, 1–16.

Erler, D. V, Santos, I.R., Zhang, Y., Tait, D.R., Befus, K.M., Hidden, A., Li, L., Eyre, B.D.,

2014. Nitrogen transformations within a tropical subterranean estuary. Mar. Chem. 164, 38–

47.

Fortunato, C.S., Crump, B.C., 2011. Bacterioplankton community variation across river to ocean

environmental gradients. Microb. Ecol. 62, 374–382.

Garcés, E., Basterretxea, G., Tovar-Sánchez, A., 2011. Changes in microbial communities in

response to submarine groundwater input. Mar. Ecol. Prog. Ser. 438, 47–58.

https://doi.org/10.3354/meps09311

Griebler, C., Lueders, T., 2009. Microbial biodiversity in groundwater ecosystems. Freshw. Biol.

https://doi.org/10.1111/j.1365-2427.2008.02013.x

Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological statistics software

package for education and data analysis.

Hays, R.L., Ullman, W.J., 2007. Dissolved nutrient fluxes through a sandy estuarine beachface

(Cape Henlopen, Delaware, USA): Contributions from fresh groundwater discharge,

seawater recycling, and diagenesis. Estuaries and Coasts 30, 710–724.

Herlemann, D.P.R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J., Andersson, A.F.,

2011. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic

Sea. ISME J. 5, 1571–1579.

Herrmann, M., Wegner, C.-E., Taubert, M., Geesink, P., Lehmann, K., Yan, L., Lehmann, R.,

Totsche, K.U., Küsel, K., 2019. Predominance of Cand. Patescibacteria in groundwater is

668  caused by their preferential mobilization from soils and flourishing under oligotrophic

669  conditions. Front. Microbiol. 10, 1407.

670 Hofmann, R., Griebler, C., 2018. DOM and bacterial growth efficiency in oligotrophic

671  groundwater: absence of priming and co-limitation by organic carbon and phosphorus.

672  Aquat. Microb. Ecol. 81, 55–71.

673 Huang, S., Wilhelm, S.W., Harvey, H.R., Taylor, K., Jiao, N., Chen, F., 2012. Novel lineages of

674  Prochlorococcus and Synechococcus in the global oceans. ISME J. 6, 285–297.

675 Kafri, U., Kessler, A., 2001. Hydrogeological plan for the development, production and

676  exploitation of the Kabri Basin. Geol. Surv. Isr. Jerusalem, Isr.

677 Kaile'a, M.C., Wiegner, T.N., 2016. Effects of submarine groundwater discharge on bacterial

678  growth efficiency in coastal Hawaiian waters. Aquat. Microb. Ecol. 77, 167–181.

679 Kearney, S.M., Thomas, E., Coe, A., Chisholm, S.W., 2021. Microbial diversity of co-occurring

680  heterotrophs in cultures of marine picocyanobacteria. Environ. Microbiome 16, 1–15.

681 Kirchman, D.L., 2012. Degradation of organic material. Process. Microb. Ecol. Oxford Univ.

682  Press Inc., New York, NY 79–98.

683 Kiro, Y., Yechieli, Y., Lyakhovsky, V., Shalev, E., Starinsky, A., 2008. Time response of the

684  water table and saltwater transition zone to a base level drop. Water Resour. Res. 44.

685 Kwon, E.Y., Kim, G., Primeau, F., Moore, W.S., Cho, H.M., Devries, T., Sarmiento, J.L.,

686  Charette, M.A., Cho, Y.K., 2014. Global estimate of submarine groundwater discharge

687  based on an observationally constrained radium isotope model. Geophys. Res. Lett. 41,

688  8438–8444. https://doi.org/10.1002/2014GL061574

689 Lecher, A.L., Mackey, K., Kudela, R., Ryan, J., Fisher, A., Murray, J., Paytan, A., 2015. Nutrient

690  loading through submarine groundwater discharge and phytoplankton growth in Monterey

Bay, CA. Environ. Sci. Technol. 49, 6665–6673.

Lecher, A.L., Mackey, K.R.M., Paytan, A., 2017. River and submarine groundwater discharge

effects on diatom phytoplankton abundance in the Gulf of Alaska. Hydrology 4, 61.

Lee, E., Shin, D., Hyun, S.P., Ko, K.S., Moon, H.S., Koh, D.C., Ha, K., Kim, B.Y., 2017.

Periodic change in coastal microbial community structure associated with submarine

groundwater discharge and tidal fluctuation. Limnol. Oceanogr. 62, 437–451.

https://doi.org/10.1002/lno.10433

Luijendijk, E., Gleeson, T., Moosdorf, N., 2020. Fresh groundwater discharge insignificant for

the world's oceans but important for coastal ecosystems. Nat. Commun. 11, 1–12.

McAllister, S.M., Barnett, J.M., Heiss, J.W., Findlay, A.J., MacDonald, D.J., Dow, C.L., Luther,

G.W., Michael, H.A., Chan, C.S., 2015. Dynamic hydrologic and biogeochemical processes

drive microbially enhanced iron and sulfur cycling within the intertidal mixing zone of a

beach aquifer. Limnol. Oceanogr. 60, 329–345. https://doi.org/10.1002/lno.10029

Miyoshi, T., Iwatsuki, T., Naganuma, T., 2005. Phylogenetic characterization of 16S rRNA gene

clones from deep-groundwater microorganisms that pass through 0.2-micrometer-pore-size

filters. Appl. Environ. Microbiol. 71, 1084–1088.

Moore, W.S., 2010. The Effect of Submarine Groundwater Discharge on the Ocean. Ann. Rev.

Mar. Sci. https://doi.org/10.1146/annurev-marine-120308-081019

Moore, W.S., 1999. The subterranean estuary: A reaction zone of ground water and sea water, in:

Marine Chemistry. pp. 111–125. https://doi.org/10.1016/S0304-4203(99)00014-6

Nielsen, E.S., 1952. The use of radio-active carbon (c14) for measuring organic production in the

sea. ICES J. Mar. Sci. 18, 117–140. https://doi.org/10.1093/icesjms/18.2.117

Overholt, W.A., Trumbore, S., Xu, X., Bornemann, T.L. V, Probst, A.J., Krüger, M., Herrmann,

M., Thamdrup, B., Bristow, L.A., Taubert, M., 2022. Carbon fixation rates in groundwater

similar to those in oligotrophic marine systems. Nat. Geosci. 15, 561–567.

Paldor, A., Katz, O., Aharonov, E., Weinstein, Y., Roditi-Elasar, M., Lazar, A., Lazar, B., 2020.

Deep submarine groundwater discharge–Evidence from Achziv Submarine Canyon at the

exposure of the Judea Group confined aquifer, eastern Mediterranean. J. Geophys. Res.

Ocean.

Paldor, A., Shalev, E., Katz, O., Aharonov, E., 2019. Dynamics of saltwater intrusion and

submarine groundwater discharge in confined coastal aquifers: a case study in northern

Israel. Hydrogeol. J. 1–15.

Probst, A.J., Ladd, B., Jarett, J.K., Geller-Mcgrath, D.E., Sieber, C.M.K., Emerson, J.B.,

Anantharaman, K., Thomas, B.C., Malmstrom, R.R., Stieglmeier, M., Klingl, A., Woyke,

T., Ryan, M.C., Banfield, J.F., 2018. Differential depth distribution of microbial function

and putative symbionts through sediment-hosted aquifers in the deep terrestrial subsurface.

Nat. Microbiol. 3, 328–336. https://doi.org/10.1038/s41564-017-0098-y

Purkamo, L., von Ahn, C.M.E., Jilbert, T., Muniruzzaman, M., Bange, H.W., Jenner, A.-K.,

Böttcher, M.E., Virtasalo, J.J., 2022. Impact of submarine groundwater discharge on

biogeochemistry and microbial communities in pockmarks. Geochim. Cosmochim. Acta

334, 14–44.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,

2012. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Res. 41, D590–D596.

Rahav, E., Raveh, O., Yanuka-Golub, K., Belkin, N., Astrahan, P., Maayani, M., Tsumi, N.,

Kiro, Y., Herut, B., Silverman, J., 2020. Nitrate-Enrichment Structures Phytoplankton

Communities in the Shallow Eastern Mediterranean Coastal Waters. Front. Mar. Sci.

Rahav, E., Silverman, J., Raveh, O., Hazan, O., Rubin-Blum, M., Zeri, C., Gogou, A., Kralj, M., Pavlidou, A., Kress, N., 2019. The deep water of Eastern Mediterranean Sea is a hotspot for bacterial activity. Deep Sea Res. Part II Top. Stud. Oceanogr. https://doi.org/10.1016/j.dsr2.2019.03.004

Robinson, C.E., Xin, P., Santos, I.R., Charette, M.A., Li, L., Barry, D.A., 2018. Groundwater dynamics in subterranean estuaries of coastal unconfined aquifers: Controls on submarine groundwater discharge and chemical inputs to the ocean. Adv. Water Resour. 115, 315–331. https://doi.org/10.1016/j.advwatres.2017.10.041

Rocha, C., Robinson, C.E., Santos, I.R., Waska, H., Michael, H.A., Bokuniewicz, H.J., 2021. A place for subterranean estuaries in the coastal zone. Estuar. Coast. Shelf Sci. 250, 107167.

Rodellas, V., Garcia-Orellana, J., Masqué, P., Feldman, M., Weinstein, Y., Boyle, E.A., 2015. Submarine groundwater discharge as a major source of nutrients to the Mediterranean Sea. Proc. Natl. Acad. Sci. 112, 3926–3930. https://doi.org/10.1073/pnas.1419049112

Ruiz-González, C., Rodellas, V., Garcia-Orellana, J., 2021. The microbial dimension of submarine groundwater discharge: current challenges and future directions. FEMS Microbiol. Rev.

Santos, I.R., Chen, X., Lecher, A.L., Sawyer, A.H., Moosdorf, N., Rodellas, V., Tamborski, J., Cho, H.-M., Dimova, N., Sugimoto, R., 2021. Submarine groundwater discharge impacts on coastal nutrient biogeochemistry. Nat. Rev. Earth Environ. 2, 307–323.

Slomp, C.P., Van Cappellen, P., 2004. Nutrient inputs to the coastal ocean through submarine groundwater discharge: controls and potential impact. J. Hydrol. 295, 64–86. https://doi.org/10.1016/j.jhydrol.2004.02.018

Smith, D.C., Azam, F., 1992. A simple, economical method for measuring bacterial protein

synthesis rates in seawater using 3H-leucine. Mar. Microb. food webs 6, 107–114.

Tada, Y., Taniguchi, A., Nagao, I., Miki, T., Uematsu, M., Tsuda, A., Hamasaki, K., 2011.

Differing growth responses of major phylogenetic groups of marine bacteria to natural

phytoplankton blooms in the western North Pacific Ocean. Appl. Environ. Microbiol. 77,

4055–4065.

Teeling, H., Fuchs, B.M., Becher, D., Klockow, C., Gardebrecht, A., Bennke, C.M., Kassabgy,

M., Huang, S., Mann, A.J., Waldmann, J., 2012. Substrate-controlled succession of marine

bacterioplankton populations induced by a phytoplankton bloom. Science (80-.). 336, 608–

611.

Tian, R., Ning, D., He, Z., Zhang, P., Spencer, S.J., Gao, S., Shi, W., Wu, L., Zhang, Y., Yang,

Y., 2020. Small and mighty: adaptation of superphylum Patescibacteria to groundwater

environment drives their genome simplicity. Microbiome 8, 1–15.

Unno, T., Kim, J., Kim, Y., Nguyen, S.G., Guevarra, R.B., Kim, G.P., Lee, J.-H., Sadowsky,

M.J., 2015. Influence of seawater intrusion on microbial communities in groundwater. Sci.

Total Environ. 532, 337–343.

Vollberg, F., Walther, M., Gärdes, A., Moosdorf, N., 2019. Modeling the potential of submarine

groundwater discharge to facilitate growth of Vibrio cholerae bacteria. Hydrology 6, 39.

Wang, X., Ye, Y., Xu, F.-F., Duan, Y.-H., Xie, P.-F., Yang, Q., Zhang, X.-L., 2021.

Maritimibacter alexandrii sp. nov., a New Member of Rhodobacteraceae Isolated from

Marine Phycosphere. Curr. Microbiol. 78, 3996–4003.

Weinstein, Y., Less, G., Kafri, U., Herut, B., 2006. Submarine groundwater discharge in the

southeastern Mediterranean (Israel). Radioact. Environ. 8, 360–372.

https://doi.org/10.1016/S1569-4860(05)08029-0

Xia, X., Zheng, Q., Leung, S.K., Wang, Y., Lee, P.Y., Jing, H., Jiao, N., Liu, H., 2021. Distinct

metabolic strategies of the dominant heterotrophic bacterial groups associated with marine

Synechococcus. Sci. Total Environ. 798, 149208.

Yau, V.M., Schiff, K.C., Arnold, B.F., Griffith, J.F., Gruber, J.S., Wright, C.C., Wade, T.J.,

Burns, S., Hayes, J.M., McGee, C., 2014. Effect of submarine groundwater discharge on

bacterial indicators and swimmer health at Avalon Beach, CA, USA. water Res. 59, 23–36.

Zheng, Q., Wang, Y., Xie, R., Lang, A.S., Liu, Y., Lu, J., Zhang, X., Sun, J., Suttle, C.A., Jiao,

N., 2018. Dynamics of heterotrophic bacterial assemblages within Synechococcus cultures.

Appl. Environ. Microbiol. 84, e01517-17.

Zhou, Y., Sawyer, A.H., David, C.H., Famiglietti, J.S., 2019. Fresh submarine groundwater

discharge to the near-global coast. Geophys. Res. Lett. 46, 5855–5863.

795