1

Original article
UV-induced feather color change reflects its porphyrin content

Masaru Hasegawa[1], Emi Arai[2], Shosuke Ito[3], & Kazumasa Wakamatsu[3]

[1]Department of Environmental Science, Ishikawa Prefectural University, Nonoichi, Ishikawa 921-8836, Japan

[2] Research Institute for Humanity and Nature, 457-4 Motoyama, Kamigamo, Kita-ku, Kyoto, 603-8047, Japan

[3] Institute for Melanin Chemistry, Fujita Health University, 1-98 Dengakugakubo, Kutsukake-cho, Toyoake, Aichi, 470-1192, Japan

Number of tables 6

Number of figures 6

Number of supplementary materials 6

Author for corresponding

Dr Masaru Hasegawa

E-mail: perorobomusadiobe@gmail.com

Tel: (+81)- 46-858-1562

AUTHOR CONTRIBUTIONS

MH conceived the ideas; MH and EA designed methodology; MH conducted field survey, collected the data of color measurements, and analyzed the data, and led the writing of the manuscript; EA prepared samples for HPLC analyses; SI and KW conducted HPLC analyses and collected the data of pigmentation. All authors contributed to final draft and gave final approval for publication.

ACKNOWLEDGMENTS

We are grateful to the residents of Tsurugi Town and Miyazaki City for their kind support and assistance. We also thank Dr. Ichiro Tayasu, and Dr. Shumpei Kitamura and his laboratory members in Ishikawa Prefectural University. MH was supported by the Research Fellowship of the Japan Society for the Promotion of Science (JSPS, 22J40066).

DATA AVAILABILITY STATEMENT

The datasets supporting this article have been uploaded as part of electronic supplementary material, table S1, which will be uploaded to osf.io after acceptance.

CONFLICT OF INTEREST

We declare no conflict of interest.

Abstract

1. Pigmentary coloration is widespread in animals. Its evolutionary and ecological

features are often attributed to the property of predominant pigments; therefore, most

research has focused on predominant pigments such as carotenoids in carotenoid-

based coloration. However, coloration results from predominant pigments and many

other minority pigments, and the importance of the latter is overlooked.

2. Here, we focused on porphyrin, an "uncommon" pigment found in bird feathers, and

investigate its importance in the context of feather color changes in the barn swallow

Hirundo rustica. By irradiating UV light on feathers, we investigated feather color

change and its relationship with porphyrin pigmentation in the barn swallow.

3. We found that the "pheomelanin-based coloration" of the barn swallow faded after

the irradiation of UV light, and this effect was particularly strong in the feathers of

young swallows (nestlings and fledglings, here). We also found that it is not the

predominant pigment, pheomelanin, but protoporphyrin IX pigment that showed the

same pattern of depigmentation after the irradiation of UV light, particularly in the

feathers of young swallows. In fact, the abovementioned age-dependent feather color

change was statistically explained by the amount of porphyrin in the feathers.

4. The current study demonstrates that a minority pigment, porphyrin, explains within-

season dynamic color change, an ecological feature of feather coloration. The

porphyrin-mediated rapid color change would benefit young birds, in which feather

coloration affects the parental food allocation during a few weeks before

independence, but not later. Future studies should not ignore these minor but

essential pigments and their evolutionary and ecological functions.

KEYWORDS

eumelanin, pheomelanin, plumage ornaments, protoporphyrin IX, seasonal color change

1 | INTRODUCTION

Pigmentary coloration is widely distributed in animals and serves many functions including camouflage, interspecific signaling (e.g., warning coloration), and intraspecific signaling (e.g., mate attraction; reviewed in Andersson, 1994; Espmark et al., 2000; Hill & McGraw, 2006). Pigmentary coloration is often subdivided by predominant pigments, such as carotenoid-based coloration, eumelanin-based coloration, and pheomelanin-based coloration (e.g., see McGraw, 2006a,b on bird coloration). It is then not surprising that the evolutionary and ecological features of animal coloration are often attributed to the chemical properties of predominant pigments. For example, carotenoid-based coloration can serve as an honest signal of quality to signal receivers, because carotenoids consumed as antioxidants cannot simultaneously serve as feather colorants (reviewed in McGraw, 2006a; also see McGraw, 2005 for other major pigments). While such a simple explanation is appealing, coloration results from predominant pigments and several other minority pigments. The evolutionary and ecological importance of the latter is ignored, solely because they are less abundant than the former.

Porphyrin is a class of endogenously synthesized colorants that give a reddish, rusty, or brown appearance (McGraw, 2006c). Many animals manufacture porphyrins, as evident from the fact that a well-known blood pigment, hemoglobin, is synthesized from natural porphyrin via the addition of iron (i.e., iron-protoporphyrin IX; McGraw, 2006c). Nevertheless, besides some exceptions (e.g., the European hedgehog *Erinasceus europaeus*), birds are the only animals that can deposit porphyrin in integumentary organs such as skin and feathers (Galván et al., 2016). Although feather porphyrin is well-known in certain non-passerine birds, such as owls and nightjars, passerines also deposit porphyrin into their feathers (e.g., Ozaki & Imamura, 2019). The function of feather porphyrin has not been known (reviewed in McGraw, 2006c), but it might function as an "ephemeral" signal because porphyrin is easily photodegraded (Galván et al., 2016). By depositing porphyrin rather than more stable alternatives, such as pheomelanin pigments,

which also provide a rusty or brown appearance (McGraw, 2006b,c), birds might display

conspicuous coloration for only a short period and may even directly signal the freshness

of the feathers (Galván et al., 2016). However, the effect of porphyrin on feather color

change, particularly in its importance relative to predominant feather pigments (e.g.,

pheomelanin), remains to be clarified.

In the current study, we investigated porphyrin pigmentation in the context of

feather color change using the throat feathers of the barn swallow *Hirundo rustica*. Throat

feather coloration of the barn swallow is reported to be used in mate choice (e.g., Ninni,

2003; Safran et al., 2005), intrasexual contests (e.g., Hasegawa et al., 2014; Wilkins et al.,

2015), and parental preferential provisioning of their young (e.g., Romano et al., 2016;

Arai et al., unpublished data). Throat feather coloration mainly consists of pheomelanin

and eumelanin pigments (e.g., McGraw et al., 2005), and this coloration changes over

time (e.g., Hasegawa et al., 2008; Safran et al., 2010; Arai et al., 2015). Moreover, the

color change is reported to be associated with the quality of birds (e.g., Safran et al., 2010),

and, within-season changes in feather coloration affect the response of signal receivers

(e.g., Safran et al., 2005). In this species, feather color change is particularly notable in

juveniles, because their initially pinkish feathers fade to whitish before the next molt so

that old and newly molted feathers are visually distinguishable (e.g., see Fig. S1 for an

example). Previous studies often assume that pheomelanin and eumelanin pigments are

the main determinants of feather coloration and its evolutionary and ecological features,

including feather color change (e.g., Arai et al., 2015, 2017, 2019). However, this

"pheomelanin-based" coloration might be affected by other pigments, such as porphyrin,

which has not previously been examined but is a candidate pigment for feather color

change, particularly in the young, in which the feather color change seems striking (see

above).

Here, by irradiating UV light (peak: 365 nm) on feathers for 30 days, we

investigated feather color change and its relationship with porphyrin pigmentation in the

barn swallow. We predicted that, if porphyrin, which can be easily photodegraded,

contributes to feather color change in this study system, feather coloration would be

affected by UV treatment, particularly in the young (see above). The effect of UV light

would be particularly noticeable when examining the feather color change before and

after the experiment. The amount of porphyrin in the feathers would show a pattern

similar to that of feather coloration. Then, this pigment (but not eumelanin and

pheomelanin pigments) would explain the variation of the feather color change. Lastly, to

clarify whether or not age-difference in porphyrin content in feathers, if any, is caused by

the time elapsed after feather emergence, we examined porphyrin content in the feathers

collected from molting adults during the non-breeding season. This was then compared

with those collected from adults and young during the breeding season. If the amount of

porphyrin is simply determined by the time elapsed after feather emergence, the feathers

of molting adults would contain a relatively large amount of porphyrin similar to those of

young. Based on previous studies (e.g., McGraw, 2006c; Galván et al., 2016, 2018), we

focused on coproporphyrin III and protoporphyrin IX as candidate porphyrin pigments.

We discuss the observed patterns from the evolutionary and ecological perspectives.

2 | MATERIALS AND METHODS

2.1 | Field survey

The field survey was conducted during the breeding season (1 April–3 July) in 2022 in a

residential area of Tsurugi Town, Ishikawa Prefecture, Japan (36°26′N, 136°37′E). We

inspected nests every third day to record breeding events such as laying dates, clutch size,

and hatching dates. Adults were captured in sweep nets while roosting at night. Each bird

was provided with a numbered aluminum ring and a unique combination of half-sized

colored rings. The sex of each individual was determined by the tail shape and by the

presence (female) or absence (male) of an incubation patch (Turner, 2006). During the

field survey, we also captured nestlings by hand from breeding nests. Upon capture, throat

feathers (approximately 20–40 feathers per bird) were plucked from each bird. In the

current study, we used throat feathers of adult females, simply because the capture of

females was relatively easier (e.g., Safran & McGraw, 2004), and to exclude the potential

confounding effect of sex difference in pigmentation (e.g., see Arai et al., 2019 for

Japanese swallows). All adult females completed their molt. We found that nestling throat

feathers were too small to accurately measure coloration when they were young (see Arai

et al., 2017 for detailed feather development in Japanese barn swallows), and thus,

concerning nestlings, we plucked throat feathers from well-grown, 16–18-day-old

nestlings. Nestling sex was not identified, because the sex difference in throat coloration

is at best small (Arai et al., 2018). In this population, we collected samples from seven

different females and seven different nestlings. The seven nestlings were captured from

seven different breeding nests (i.e., samples were collected from one nestling per nest).

To increase sample size, feather samples were also collected from a breeding

population in Miyazaki City, Miyazaki Prefecture, Japan (31°54′N, 131°25′E). As in the

Tsurugi population, adults were captured in sweep nets in this population. In addition to

adult females, we also captured fledglings, instead of nestlings, in sweep nests, because

most nestlings fledged before the study period (5–8 July, 2022). In this population,

samples were collected from three different females and three different fledglings. The

three fledglings were captured at different nests and roosting sites.

Some barn swallows remain year-round in the Miyazaki population (Hasegawa,

2020), and thus newly molted feathers can be collected from molting swallows, in which

old and newly molted feathers can be easily distinguished (Fig. S1). For this reason,

during 12–17 February 2023, wintering swallows were captured there to collect newly

molted throat feathers. We collected samples from three different molting adults, although

we could not identify the sex of these individuals (because wintering swallows do not

develop brood patches, thereby preventing successful sex identification, particularly

when their feathers are still growing).

2.2 | Color measurements

In the laboratory, we piled five throat feathers collected from each individual on a piece of white paper so that the perimeters of the feathers coincided. From each bird, two feather samples were taken, one for the control and the other for the UV treatment group (see below). The feather samples were scanned at 800 dpi resolution using a scanner (GT X970; Epson, Tokyo, Japan), and the images obtained were imported into Photoshop Elements 15 (Adobe Systems, San Jose, CA). We measured the mean red-green-blue (RGB) values for a 30×30 pixel square near the distal end of the feather sample. The mean RGB values were converted into hue-saturation-brightness (HSB) values, using the algorithm described by Foley and van Dam (1984). Throat feathers exhibit few ultraviolet reflectance, so this method captured most of the variation in throat coloration (e.g., Safran & McGraw, 2004). Because previous studies in this study system also used HSB values (e.g., McGraw et al., 2005; Hasegawa et al., 2008; Arai et al., 2015), this method enables us to directly compare our results with those of previous studies. Details of the methodology are described elsewhere (Hasegawa et al., 2008).

2.3 | Experimental UV irradiation

The experiment was conducted from 27 October to 28 November 2022 with a short pause to check the progress of the experiment. Before the start of the experiment, we measured the coloration of all 40 feather samples (see above). Two feather samples taken from the same birds resembled each other, demonstrating high repeatability of HSB values between samples (Hue: repeatability = 0.92, $F_{19,20} = 25.34$, $p < 0.0001$; Saturation: repeatability = 0.96, $F_{19,20} = 55.56$, $p < 0.0001$; Brightness: repeatability = 0.93, $F_{19,20} = 26.84$, $p < 0.0001$). Thus, between-individual variation was much larger than within-individual variation in our measurements. Two feather samples each were taken from 20 different individuals, one for the control and the other for the UV treatment group (see

above), and thus control and UV treatment groups each had 20 feather samples. Regarding

the UV treatment group, feather samples were placed in a closed cardboard box (size: 37

cm × 25 cm × 14 cm), in which they were continuously exposed to UV light (FL365-SD,

27W; peak spectrum: 365 nm, approximate range: 350–400 nm, intensity: 1300 µW/cm^2

from a distance of 15 cm) at room temperature for 30 days. The intensity of the UV light

was within the range of natural UV radiation in Japan (e.g., the intensity of UV-A radiation

in open fields between 900 and 1600 in August was 1000–2500 µW/cm^2 in Akashi City,

Hyogo Prefecture in Japan: Ishiuchi et al., 2012). Regarding the control group, samples

were placed in the same-sized cardboard box without UV light (i.e., they were held in

complete darkness) at room temperature for 30 days. The two treatment groups

experienced the same amount of time in the boxes, because the two treatments started and

ended at the same time. The two boxes were held in the same room. After the end of the

experiment, the coloration of all feather samples was measured again, and the pigment

content of each feather sample was evaluated using high-performance liquid

chromatography (HPLC; see below).

2.4 | Pigment quantification via HPLC

We followed the extraction protocols of porphyrins from feathers given in preceding

studies (Galván et al., 2016, 2018; Loules et al., 2022). In short, five feathers (0.2–0.3

mg) were extracted in 300 µL 6 M HCOOH plus acetonitrile (1:1), by homogenizing them

with a Tenbroeck tissue grinder. Notably, unlike in the previous studies (e.g., Arai et al.

2017), we did not cut off the feathers' basal part to effectively extract protoporphyrin

pigments. After centrifugation in a glass conical tube, the supernatants (50 µL) were

subjected to the following HPLC analyses.

We used the precipitates from the feather extraction to analyze eumelanin (pyrrole-

2,3,5-tricarboxylic acid; PTCA) and pheomelanin (thiazole-2,4,5-tricarboxylic acid;

TTCA). Precipitates (and homogenizer) were washed once with 1 mL water by

centrifugation, dried in a desiccator, and subjected to alkaline hydrogen peroxide

oxidation (AHPO) as described in previous research (Ito et al., 2011, 2020) using 40%

volumes of reagents (Arai et al., 2018, 2019).

The following HPLC conditions were applied. First, A Waters (Milford, MA, USA)

Spherisorb ODS 2 (5 µm particle size, 4.6 mm × 250 mm) HPLC column was used. Then,

protoporphyrin IX (PPIX) and coproporphyrin III (CPIII) (standards: 1 µg/mL 6 M

HCOOH plus acetonitrile (1:1)) were analyzed by isocratic elution under different

methanol concentrations. PPIX was analyzed by methanol and 1 M ammonium acetate =

90:10 (v/v). We used high concentrations of ammonium acetate to ensure high buffering

capacity. The column temperature was 70°C (we use the high temperature to make the

PPIX peak sharp and high). VIS detection was performed with a UV/VIS detector at 402

nm (Galván et al., 2016). PPIX appeared at 7.5 min. CPIII was analyzed separately by

methanol and 1M ammonium acetate = 60:40 (v/v) at a column temperature of 50 °C.

CPIII appeared at 6.9 min.

2.5 | Statistical procedures

A linear mixed-effects model (lmer function in the R package lme4: Bates et al., 2015)

was used to test the effects of treatment ("Control" and "UV treatment" were denoted as

0 and 1, respectively, and "adults" and "young" were denoted as 0 and 1, respectively)

and the interaction between treatment and age on each dependent variable (i.e., hue,

saturation, and brightness for color measurements, their changes before and after the

experiment, and the amount of each pigment). As before (e.g., Arai et al., 2017), the

amount of pigmentation was log-transformed before analysis to fit a normal distribution

(Shapiro-Wilk test; before: $0.66 < W < 0.96$; after: $W > 0.93$). We also examined plumage

coloration in relation to pigmentation to test the effect of pigmentation on coloration. In

these models, bird identity was included as a random factor to control for

pseudoreplication. We did not include the identity of populations (i.e., Tsurugi/Miyazaki)

as another fixed factor to avoid complicating the model structure (note that we could not

include the study population as a random factor due to the small number of random-effect

levels, e.g., Bolker et al., 2009 suggests ">5–6 random-effect levels per random effect"

for properly estimating among-group variance). We also conducted analyses of "residual"

components of feather coloration controlling for pigment contents, in which, from

observed feather coloration, the component explained by the amount of each pigment

content was subtracted using the coefficients of the abovementioned models. If the effect

of UV exposure on feather coloration reflects on particular pigment content (e.g.,

protoporphyrin IX), the residual component controlling for the pigment content (but not

those for others) would no longer be explained by UV treatment and its interaction with

age. Additionally, to further test the effect of pigmentation on feather coloration, we

examined difference in the feather coloration between two samples from the same

individuals (i.e., one used for control and the other used for UV treatment) in relation to

the corresponding difference in pigmentation. In this analysis with a small sample size (n

= 18), we used a robust linear model (function lmrob in the R package robustbase) to

minimize the influence of outliers, as before (Arai et al., 2017). Lastly, we compared the

pigment contents of feathers collected from molting adults and those collected from adults

and young during the breeding season, to test the effect of time elapsed after feather

emergence. All data analyses were performed using the R statistical package (version

4.1.0; R Core Team, 2021).

2.6 | Replication statement

The level of interest is among feather samples. The factors of interest are age and

treatment (Control and UV group) and the unit of replication is individual, hence, they

have ten replicate individuals of each age/treatment combination (note that we used a

"paired" design, i.e., two feather samples were taken from each individual, one for the

control and the other for the UV treatment group; see above).

3 | RESULTS

3.1 | Pigmentation per feather

We could quantify PTCA, i.e., an index of eumelanin, TTCA, i.e., an index of pheomelanin, and protoporphyrin IX contents of 38 feather samples. In these samples, the throat feathers of barn swallows contained 13.65 ± 6.96 ng/feather (mean $\pm$ SD) PTCA (range: 3.5–26.9 ng/feather), and 13.44 ± 11.22 ng/feather (mean $\pm$ SD) TTCA (range: 2.3–56.5 ng/feather). We also detected protoporphyrin IX pigments (mean $\pm$ SD = 67.10 $\pm$ 97.42 pg/feather, range: 0–403 pg/feather), and its amount was on average 734 times less than TTCA (range: 14–3786 times, excluding two individuals with no detectable protoporphyrin IX content). We could not estimate the amount of protoporphyrin IX pigments in two feather samples from two different individuals, and thus the following analyses of pigmentation consisted of 36 samples from the remaining 18 individuals (note that, as we adopted a "paired" experimental design, each pair of samples should consist of one in control and the other in the UV treatment group). Although we also attempted to analyze coproporphyrin III pigment, substantial amounts of it were not detected in the current samples (i.e., only traces of this pigment were detected in HPLC analysis), preventing further analyses of this pigment.

3.2 | UV, coloration, and pigmentation

The interaction term between age and treatment explained the coloration of throat feathers, as measured by hue, saturation, and brightness (Table 1). Feathers exposed to UV light showed significantly higher hue (i.e., less red), saturation, and brightness than control feathers even in adults, as shown by the coefficients of "Treatment" in Table 1, but the effect was particularly strong in young (Fig. 1), as shown by the significant positive interaction terms in Table 1. A similar pattern was found when we analyzed feather color change before and after the experiment (Table 2; Fig. 2).

Likewise, the interaction term between age and treatment explained the

protoporphyrin IX content of throat feathers (Table 3). Feathers exposed to UV light

contained significantly lower amounts of protoporphyrin IX pigments than control

feathers, even in adults (ca. 1/3, deduced from Table 3), but again, this effect was

particularly strong in young (ca. 1/20, deduced from Table 3; Fig. 3). In contrast, the

interaction term between age and treatment was far from significant in PTCA, an index

of eumelanin, and TTCA, an index of pheomelanin (Table 3). In these models, treatment

remained nonsignificant after the nonsignificant interaction terms were excluded from the

models ($p > 0.43$; details not shown).

When we used three pigments as predictor variables in mixed-effects models,

protoporphyrin IX explained the variation of the color measurements (Table 4). All three

measurements, i.e., hue, saturation, and brightness, decreased with increasing

protoporphyrin IX pigments (Fig. 4). Although we also found the significant effect of

TTCA, an index of pheomelanin (Table 4), a close inspection of Figure 4 indicates that

this was due to between-age differences rather than within-age variation (Fig. 4). In fact,

after controlling for the age difference in coloration and pigmentation by subtracting the

coefficients of "Age" from measurements of young (see Tables 1 & 3), protoporphyrin

IX, but not TTCA, remained significant (protoporphyrin IX: all: $p < 0.0001$; PTCA: all:

$p > 0.36$; TTCA: all: $p > 0.16$; details not shown). When we examined feather color

change before and after the experiment in relation to the three pigments, protoporphyrin

IX but not PTCA or TTCA, explained the color change (Table 5). The more color changed

before and after the experiment, the fewer protoporphyrin IX remained in the feathers

after the experiment (Fig. 5).

After statistically controlling for protoporphyrin IX content, the interaction term

between age and treatment on feather coloration and its color change did not remain

significant (Tables S3 & S4). This was not the case when we statistically controlled for

PTCA or TTCA (i.e., interaction terms remained significant: $p < 0.05$; details not shown).

An alternative approach to study the linkage between coloration and

pigmentation is to examine the difference in pigmentation between the two treatments

collected from the same individuals (i.e., one in the UV treatment group and the other in

the Control; see Methods) in relation to the corresponding difference in coloration

between them, assuming that the effect of UV treatment on coloration reflects that on its

pigment content. As predicted, we found that the difference in protoporphyrin IX

consistently explained the difference in each of the three coloration measurements,

although the analysis of hue difference was marginal (i.e., $p = 0.05$; Table S2). The

difference in the color measurements between the two treatments increased with the

increasing reduction of protoporphyrin IX pigments in the UV treatment group compared

with controls (Fig. S2).

3.3 | Newly molted feathers

When comparing protoporphyrin IX pigments of feathers collected from molting adults

in winter (Fig. S1) with those collected from adults and young during the breeding season,

we found that the amount of protoporphyrin IX content of throat feathers collected from

molting adults was comparable to that of throat feathers collected from young during the

breeding season but was significantly higher than that of throat feathers collected from

adults during the breeding season (ca. five times, deduced from Table 6; Fig. 6). There

were no detectable differences in the amount of eumelanin and pheomelanin content of

throat feathers between molting adults and the other two groups, although the difference

in the amount of pheomelanin content between throat feathers collected from young and

molting adults was marginal (Table 6; Fig. 6).

4 | DISCUSSION

The main finding of the current study is that protoporphyrin IX pigments, seemingly an

"uncommon pigment" (McGraw, 2006c), explained the pheomelanin-based feather

coloration of barn swallows and its age-dependent color change under the exposure of UV light. This finding is understandable due to the wide range of protoporphyrin IX content of the feathers but is surprising because the protoporphyrin IX content of throat feathers was, on average, >500 times less than eumelanin and pheomelanin contents in the barn swallow. The observed porphyrin content is low even when compared with other species, such as the barn owl *Tyto alba*, in which protoporphyrin IX content was one tenth of pheomelanin pigments (Roulin et al., 2008). Previous studies often focused on the predominant pigments such as carotenoids and melanins, assuming that the evolutionary and ecological features of integument coloration can be attributed to the chemical properties of these predominant pigments (e.g., see McGraw, 2006b, Ducrest et al., 2008, Roulin, 2016 for reviews on melanin-based feather coloration). However, as was shown here, feather color change, an ecological feature of feather coloration, depends on a minority pigment, and thus the assumption that predominant pigments determine the evolutionary and ecological features of animal coloration is no longer valid. Because feather coloration changed even in adults (Tables 1 & 2), porphyrin would affect feather coloration of adults as well, although the feather color change of adults was smaller than that of young (Tables 1 & 2).

A fundamental question about porphyrin is why birds (and some other animals) use porphyrins as colorants in addition to other pigments (e.g., pheomelanin, here). A possible explanation is that, when the use of coloration is confined to a short period after feather emergence, birds use porphyrin to temporally "boost" feather coloration (Galván et al., 2016, 2018). In the barn swallow, nestling feather coloration affects parental food allocation (Romano et al. 2016; Arai et al. unpublished data), which lasts for only a few weeks before the independence of offspring (Turner, 2006). Then, for young birds, pigments that can easily be photodegraded (i.e., porphyrin) are preferable to relatively stable pigments (i.e., pheomelanin, here), because retaining conspicuous coloration after independence would incur viability costs (e.g., via increased predatory attack).

A relatively large amount of protoporphyrin IX pigments found in young, in

combination with a relatively low amount of pheomelanin pigments (Table 3), at first

seem consistent with this explanation. Young swallows might use many porphyrin

pigments to rapidly fade their coloration after its use. Susceptibility to UV light, however,

provides an alternative and simpler explanation for this pattern. In contrast with adults,

which molt after the breeding season (see Methods), nestlings grow their feathers in the

breeding nest just before fledging; therefore, much more protoporphyrin pigments would

remain in the feathers at sampling. In fact, even in adults, we found a relatively large

amount of protoporphyrin IX in their newly molted feathers (Table 6; Fig. 6), supporting

this perspective. Thus, the deposition of a large amount of protoporphyrin in feathers

would not be unique to nestlings but be shared with adults. For this reason, we cannot

exclude the possibility that the supposedly "adaptive" function of porphyrin in young (see

above) might be an exaptation (i.e., secondary use of already evolved traits for other

functions: Bergstrom & Dugatkin, 2016). Rather, as we found a substantial amount of

protoporphyrin IX in breeding adults, adults (rather than young) might evolve unique

mechanisms to slow down the photodegradation of porphyrin, which might be beneficial

for them to retain their coloration until the end of the breeding period. Although age-

dependent depigmentation of protoporphyrin IX (Table 3) is consistent with this

perspective, we have no data to test this possibility in the current data set. The ecological

function of porphyrin and its evolutionary importance remains to be clarified in future

studies, by focusing not only on pigment production but also on its maintenance.

A caveat of the current study is that we focused only on feather color change

under ultraviolet light, particularly, what is called, UV-A light (i.e., wavelength: 320–400

nm). We chose UV-A, simply because the stratospheric ozone absorbs most UV-B

radiation and thus UV-A light is much more abundant in the environment (e.g., see Japan

Meteorological Agency, 2010), and because UV-A light penetrates deeper integument

layers than UV-B light (and thus we speculated that this could explain the uniform

depigmentation of pre-molt feathers). However, feather color change can also occur for

other reasons, such as photodegradation via other wavelengths of sunlight including UV-

B, feather abrasion, and accumulation of stains (reviewed in Montgomerie, 2006). In fact,

using the same study species, Arai et al. (2015) showed that, in the wild, the amount of

pheomelanin content in throat feathers decreased at a rate of 15–25% per month, and thus

these components should also be involved in the wild, together with the intense

photodegradation of porphyrin pigments we demonstrated here. In addition, because the

amount of protoporphyrin IX could not explain all the feather color changes (Table S4),

even under the current experimental condition, not only pigmentation but also other

factors, such as preen oil, feather microstructures, and their interactions with

pigmentation, might be involved in feather color change, which should be studied in the

future to understand the whole picture of feather color change.

In summary, we demonstrated that an uncommon pigment, protoporphyrin IX,

affected pheomelanin-based coloration and its dynamic color change under UV light.

Feather color change, however, would not be the sole ecological feature of feather

porphyrin. For example, protoporphyrin IX is a well-known colorant of eggs (reviewed

in McGraw, 2006c), and its expression is known to correlate with hatching success and

female feather coloration, at least in the barn swallow (Corti et al., 2018). Then, it is likely

that the porphyrin content of feathers can partially explain the linkage, serving feather

coloration as an honest signal of the reproductive value of their clutches. It remains to be

clarified how these minority pigments affect the overall evolution and ecology of animal

coloration in future studies. In particular, the co-occurrence of porphyrin and pheomelanin,

i.e., two similar-colored pigments with different chemical properties (McGraw, 2006c),

and their inter- and intraspecific variation, would provide a unique study system to

understand how particular pigments and their combinations (or possibly displacement)

evolved to produce the same kind of coloration.

AUTHOR CONTRIBUTIONS

Please see title page

ACKNOWLEDGMENTS

Please see title page

DATA AVAILABILITY STATEMENT

Please see title page

REFERENCES

Andersson, M. (1994). *Sexual selection*. Princeton University Press.

Arai, E., Hasegawa, M., Nakamura, M., & Wakamatsu, K. (2015). Male pheomelanin
pigmentation and the breeding onset in Barn Swallows *Hirundo rustica gutturalis*.
Journal of Ornithology, 156, 419–427.

Arai, E., Hasegawa, M., Makino, T., Hagino, A., Sakai, Y., Ohtsuki, H., Wakamatsu, K.,
& Kawata, M. (2017). Physiological conditions and genetic controls of phaeomelanin
pigmentation in nestling barn swallows. *Behavioral Ecology*, 28, 706–716.

Arai, E., Hasegawa, M., Wakamatsu, K., & Ito, S. (2018). Males with more pheomelanin
have a lower oxidative balance in Asian barn swallows (*Hirundo rustica gutturalis*).
Zoological Science, 36, 505–513.

Arai, E., Hasegawa, M., Sato, M., Sakai, H., Ito, S., & Wakamatsu, K. (2019). Eumelanin
levels in rufous feathers explain plasma testosterone levels and survival in swallows.
Ecology and Evolution, 9, 2755–2764.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects
models using lme4. *Journal of Statistical Software*, 67, 1–48.

Bergstrom, C.T., & Dugatkin, L.A. (2016). *Evolution*. WW Norton & Co Inc.

Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H.,

& White, J.-S.S. (2009). Generalized linear mixed models: a practical guide for

ecology and evolution. *Trends in Ecology and Evolution*, 24, 127–135.

Corti, M., Romano, A., Constanzo, A., Bentz, A.B., Navara, K.J., Parolini, M., Saino, N.,

& Rubolini, D. (2018). Protoporphyrin-based eggshell pigmentation predicts

hatching success and offspring sex ratio in the barn swallow. *Journal of Avian*

Biology, 2018, e01642.

Ducrest, A.-L., Keller, L., & Roulin, A. (2008). Pleiotropy in the melanocortin system,

coloration and behavioural syndromes. *Trends in Ecology and Evolution*, 23, 502–

510.

Espmark, Y., Amundsen, T., & Rosenqvist, G. (2000). *Animal signals: signalling and*

signal design in animal communication. Tapir Academic Press.

Foley, J.D., & van Dam, A. (1984). Intensity and color. In: *Fundamentals of interactive*

computer graphics. pp. 593–622. Addison-Wesley.

Galván, I., Camarero, P.R., Mateo, R., & Negro, J.J. (2016). Porphyrins produce uniquely

ephemeral animal colouration: a possible signal of virginity. *Scientific Reports*, 6,

39210.

Galván, I., del Mar Delgado, M., Camarero, P.R., Mateo, R., Lourenço, R., & Penteriani,

V. (2018). Feather content of porphyrins in Eurasian eagle owl (*Bubo bubo*)

fledglings depends on body condition and breeding site quality. *Integrative Zoology*,

13, 569–578.

Hadfield, J.D. (2010). MCMC methods for multi-response generalized linear mixed

models: the MCMCglmm R package. *Journal of Statistical Software*, 33, 1–22.

Hasegawa, M. (2020). *Tsubame no himitsu (in Japanese)*. Midorishobo, Tokyo.

Hasegawa, M., Arai, E., Watanabe, M., & Nakamura, M. (2008). Methods for correcting

plumage color fading in the Barn Swallow. *Ornithological Science*, 7, 117–122.

Hasegawa, M., Arai, E., Watanabe, M., & Nakamura, M. (2014). Colourful males hold

high quality territories but exhibit reduced paternal care in barn swallows. *Behaviour*,

151, 591–612.

Hill, G.E., & McGraw, K.J. (2006). *Bird coloration, vol II. Function and evolution.*

Harvard University Press.

Ishiuchi, T., Koyanagi, T., & Kuwahara, Y. (2012). A Study on Ground Arrival Ultra

Violet Radiation in Outdoor Space. *Papers on Environmental Information. Science,*

26, 225–230.

Ito, S., Nakanishi, Y., Valenzuela, R.K., Brilliant, M.H., Kolbe, L., & Wakamatsu, K.

(2011). Usefulness of alkaline hydrogen peroxide oxidation to analyze eumelanin and

pheomelanin in various tissue samples: Application to chemical analysis of human

hair melanins. *Pigment Cell & Melanoma Research,* 24, 605–613.

Ito, S., Del Bino, S., Hirobe, T., & Wakamatsu, K. (2020). Improved HPLC conditions to

determine eumelanin and pheomelanin contents in biological samples using an ion

pair reagent. *International Journal of Molecular Sciences,* 21, 2134.

Japan Meteorological Agency (2010). Annual report of ozone layer monitoring 2010.

Retrieved from https://www.data.jma.go.jp/gmd/env/ozonehp/o3report2010.html

Louleb, M., Galván, I., Latrous, L., Justyn, N.M., Hill, G.E., Ríos, Á., & Zougagh, M.

(2022). Detection of porphyrins in hair using capillary liquid chromatography-mass

spectrometry. *International Journal of Molecular Sciences,* 23, 6230.

McGraw, K.J. (2005). The antioxidant function of many animal pigments: are there

consistent health benefits of sexually selected colourants? *Animal Behaviour,* 69,

757–764.

McGraw, K.J. (2006a). Mechanics of melanin-based coloration. In: Hill, G.E., & McGraw,

K.J., editors. *Bird coloration, vol.1: mechanisms and measurements.* Harvard

University Press, p. 243–294.

McGraw, K.J. (2006b). Mechanics of carotenoid-based coloration. In: Hill, G.E., &

McGraw, K.J., editors. *Bird coloration, vol.1: mechanisms and measurements.*

Harvard University Press, p. 177–242.

McGraw, K.J. (2006c). Mechanics of uncommon colors: Pterins, Porphyrins, and

Psittacofulvins. In: Hill, G.E., & McGraw, K.J., editors. *Bird coloration, vol.1:*

mechanisms and measurements. Harvard University Press, p. 354–398.

McGraw, K.J., Safran, R.J., & Wakamatsu, K. (2005). How feather color reflects its

melanin content. *Functional Ecology*, 19, 816–821.

Montgomerie, R. (2006). Cosmetic and adventitious colors. In: Hill, G.E., & McGraw,

K.J., editors. *Bird coloration. II. Function and evolution*. Harvard University Press,

p. 399–427.

Ninni, P. (2003). *Carotenoid Signals in Barn Swallows*. PhD thesis, Université Pierre et

Marie Curie, Paris, France.

Ozaki, T., & Imamura, S. (2019). Distribution of protoporphyrin IX in bird feathers.

International Journal of Analytical Bio-Science, 7, 41–48.

R Core Team (2021). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/.

Romano, A., Bazzi, G., Caprioli, M., Corti, M., Costanzo, A., Rubolini, D., & Saino, N.

(2016). Nestling sex and plumage color predict food allocation by barn swallow

parents. *Behavioral Ecology*, 27, 1198-1205.

Roulin, A. (2016). Condition-dependence, pleiotropy and the handicap principle of sexual

selection in melanin-based colouration. *Biological Reviews*, 91, 328–348.

Roulin, A., Almasi, B., Rossi-Pedruzzi, P., Ducrest, A.-L., Wakamatsu, K., Miksik, I.,

Blount, J.D., Jenni-Eiermann, S., & Jenni, L. (2008). Corticosterone mediates the

condition-dependent component of melanin-based coloration. *Animal Behaviour*, 75,

1351–1358.

Safran, R.J., & McGraw, K.J. (2004). Plumage coloration, not length or symmetry of tail-

streamers, is a sexually selected trait in North American barn swallows. *Behavioral*

Ecology, 15, 455–461.

Safran, R.J., Neuman, C.R., McGraw, K.J., & Lovette, I.J. (2005). Dynamic paternity

allocation as a function of male plumage color in barn swallows. *Science*, 309, 2210–

2212.

Safran, R.J., McGraw, K.J., Wilkins, M.R., Hubbard, J.K., & Marling, J. (2010). Positive

carotenoid balance correlates with greater reproductive performance in a wild bird.

PLoS ONE, 5, e9420.

Turner, A.K. (2006). *The barn swallow*. T & AD Poyser.

Wilkins, M.R., Shizuka, D., Joseph, M.B., Hubbard, J.K., & Safran, R.J. (2015).

Multimodal signalling in the North American barn swallow: a phenotype network

approach. *Proceedings of the Royal Society of London B*, 282, 20151574.

TABLE 1   Linear mixed-effects model explaining coloration of the throat feathers in

relation to age and treatment in the barn swallow ($n_{adult,control} = 10$, $n_{young,control} = 10$, $n_{adult,UV}$

$= 10$, $n_{young,UV} = 10$, $n_{total} = 40$).

Variables	Coefficient ± SE	t	p
Hue			
Age	2.65 ± 0.67	3.96	<0.001
Treatment	1.69 ± 0.27	6.24	<0.0001
Interaction	1.18 ± 0.38	3.07	<0.01
	Variance explained by a random factor, ID = 1.87		
Saturation			
Age	30.08 ± 2.94	-10.23	<0.0001
Treatment	8.17 ± 1.35	6.03	<0.0001
Interaction	6.58 ± 1.92	3.43	<0.01
	Variance explained by a random factor, ID = 34.04		
Brightness			
Age	19.71 ± 7.21	2.73	0.01
Treatment	9.83 ± 2.73	3.61	<0.01
Interaction	16.41 ± 3.85	4.26	<0.001
	Variance explained by a random factor, ID = 222.78		

576 Bird identity (ID) was included as a random factor in each model.

580 TABLE 2   Linear mixed-effects model explaining plumage color change of the throat

581 feathers in relation to age and treatment in the barn swallow ($n_{adult,control}$ = 10, $n_{young,control}$

582 = 10, $n_{adult,UV}$ = 10, $n_{young,UV}$ = 10, n_{total} = 40).

Variables	Coefficient ± SE	t	p
Hue changes			
Age	-0.14 ± 0.20	-0.70	0.49
Treatment	1.78 ± 0.15	11.91	<0.0001
Interaction	0.99 ± 0.21	4.69	<0.001
	Variance explained by a random factor, ID = 0.08		
Saturation changes			
Age	0.45 ± 1.36	0.33	0.74
Treatment	7.90 ± 1.05	7.54	<0.0001
Interaction	6.77 ± 1.48	4.57	<0.001
	Variance explained by a random factor, ID = 3.81		
Brightness changes			
Age	-1.52 ± 1.90	-0.80	0.43
Treatment	11.55 ± 1.52	7.61	<0.0001
Interaction	12.88 ± 2.15	6.00	<0.0001
	Variance explained by a random factor, ID = 6.57		

583 Bird identity (ID) was included as a random factor in each model.

585 TABLE 3 Linear mixed-effects model explaining pigmentation of the throat feathers in

586 relation to age and treatment in the barn swallow ($n_{adult,control} = 9$, $n_{young,control} = 9$, $n_{adult,UV}$

587 $= 9$, $n_{young,UV} = 9$, $n_{total} = 36$).

Variables	Coefficient ± SE	t	p
Protoporphyrin IX			
Age	0.95 ± 0.40	2.35	0.03
Treatment	-1.22 ± 0.30	-4.02	<0.001
Interaction	-1.80 ± 0.43	-4.20	<0.001
	Variance explained by a random factor, ID = 0.32		
PTCA			
Age	-0.10 ± 0.26	-0.38	0.71
Treatment	-0.01 ± 0.19	-0.04	0.97
Interaction	-0.03 ± 0.27	-0.11	0.91
	Variance explained by a random factor, ID = 0.14		
TTCA			
Age	-1.00 ± 0.21	-4.66	<0.0001
Treatment	0.18 ± 0.18	0.97	0.35
Interaction	-0.15 ± 0.26	-0.59	0.57
	Variance explained by a random factor, ID = 0.06		

PITCA and TTCA are indices of eumelanin and pheomelanin, respectively.

Bird identity (ID) was included as a random factor in each model.

Dependent variables were log-transformed before each analysis.

Note that sample sizes were reduced in this analysis because of missing information (see

Methods).

TABLE 4 Linear mixed-effects model explaining coloration of the throat feathers in

relation to pigmentation in the barn swallow ($n_{adult,control} = 9$, $n_{young,control} = 9$, $n_{adult,UV} = 9$,

$n_{young,UV} = 9$, $n_{total} = 36$).

Variables	Coefficient $\pm$ SE	t	p
Hue			
Protoporphyrin IX	-0.83 $\pm$ 0.14	-6.04	<0.0001
PTCA	0.24 $\pm$ 0.57	0.42	0.68
TTCA	-1.87 $\pm$ 0.48	-3.84	<0.001
	Variance explained by a random factor, ID = 1.42		
Saturation			
Protoporphyrin IX	-4.31 $\pm$ 0.57	-7.53	<0.0001
PTCA	-5.18 $\pm$ 2.99	-1.73	0.094
TTCA	12.47 $\pm$ 2.80	4.45	<0.0001
	Variance explained by a random factor, ID = 88.83		
Brightness			
Protoporphyrin IX	-7.58 $\pm$ 1.03	-7.37	<0.0001
PTCA	2.96 $\pm$ 4.97	0.59	0.56
TTCA	-16.69 $\pm$ 4.51	-3.70	<0.001
	Variance explained by a random factor, ID = 180.37		

PITCA and TTCA are indices of eumelanin and pheomelanin, respectively.

Bird identity (ID) was included as a random factor in each model.

Independent variables were log-transformed before each analysis.

Note that sample sizes were reduced in this analysis because of missing information (see

Methods).

TABLE 5 Linear mixed-effects model explaining color change of the throat feathers

before and after the experiment in relation to pigmentation in the barn swallow ($n_{adult,control}$

$= 9$, $n_{young,control} = 9$, $n_{adult,UV} = 9$, $n_{young,UV} = 9$, $n_{total} = 36$).

Variables	Coefficient ± SE	t	p
Hue change			
Protoporphyrin IX	-0.72 ± 0.09	-7.76	<0.0001
PTCA	-0.03 ± 0.27	-0.12	0.91
TTCA	-0.31 ± 0.21	-1.49	0.15
	Variance explained by a random factor, ID < 0.001		
Saturation change			
Protoporphyrin IX	-4.08 ± 0.45	-9.11	<0.0001
PTCA	-1.18 ± 1.80	-0.66	0.52
TTCA	-1.30 ± 1.52	-0.86	0.40
	Variance explained by a random factor, ID = 13.10		
Brightness change			
Protoporphyrin IX	-7.16 ± 0.66	-10.91	<0.0001
PTCA	-1.59 ± 2.74	-0.58	0.57
TTCA	-1.04 ± 2.34	-0.45	0.66
	Variance explained by a random factor, ID = 33.88		

PITCA and TTCA are indices of eumelanin and pheomelanin, respectively.

Bird identity (ID) was included as a random factor in each model.

Independent variables were log-transformed before each analysis.

Note that sample sizes were reduced in this analysis because of missing information (see

Methods).

A fitted model for hue change is near singular, and thus, based on the recommendation

(Bates et al. 2015), we conducted a Bayesian approach using the function MCMCglmm

in the package MCMCglmm (Hadfield, 2010) with a normal error distribution and found

that qualitatively similar results were obtained (protoporphyrin IX: coefficient = -0.72,

95%CI = -0.92, -0.55, p_{MCMC} < 0.0001; PTCA: coefficient = -0.03, 95%CI = -0.65, 0.46,

p_{MCMC} = 0.92; TTCA: coefficient = -0.30, 95%CI = -0.73, 0.12, p_{MCMC} = 0.18; no. iteration

= 130000, burnin = 30000, thinning interval = 100).

TABLE 6 Linear mixed-effects model explaining pigmentation of the throat feathers in
the barn swallow including newly molted adults ($n_{post\text{-}molting\ adult} = 3$, $n_{breeding\ adult} = 10$,
$n_{young} = 9$, $n_{total} = 22$)

Variables	Coefficient ± SE	t	p
Protoporphyrin IX			
Difference to breeding adults	-1.54 ± 0.49	-3.16	<0.01
Difference to young	-0.57 ± 0.49	-1.16	0.26
PTCA			
Difference to breeding adults	-0.08 ± 0.39	-0.20	0.84
Difference to young	-0.08 ± 0.40	-0.20	0.85
TTCA			
Difference to breeding adults	0.24 ± 0.38	0.62	0.54
Difference to young	-0.69 ± 0.39	-1.79	0.090

623 PITCA and TTCA are indices of eumelanin and pheomelanin, respectively.

624 Dependent variables were log-transformed before each analysis.

FIGURE 1   Boxplots of coloration of the throat feathers in relation to age and treatment in the barn swallow: hue (upper panel); saturation (middle panel); brightness (lower panel). The horizontal bar in each boxplot indicates the median, and the box shows the first and third quartiles of data. The whiskers range from the lowest to the highest data points within the 1.5 × interquartile range of the lower and upper quartiles, respectively. Circles indicate each data point. See Table 1 for detailed statistics.

FIGURE 2   Boxplots of color changes (value after experiment minus value before experiment) of the throat feathers in relation to age and treatment in the barn swallow: hue change (upper panel); saturation change (middle panel); brightness change (lower panel). The horizontal bar in each boxplot indicates the median, and the box shows the first and third quartiles of data. The whiskers range from the lowest to the highest data points within the 1.5 × interquartile range of the lower and upper quartiles, respectively. Circles indicate each data point. See Table 2 for detailed statistics.

FIGURE 3   Boxplots of pigmentation of the throat feathers in relation to age and

treatment in the barn swallow: protoporphyrin IX (upper panel); PTCA (middle panel);

TTCA (lower panel). PITCA and TTCA are indices of eumelanin and pheomelanin,

respectively. The horizontal bar in each boxplot indicates the median, and the box shows

the first and third quartiles of data. The whiskers range from the lowest to the highest data

points within the 1.5 × interquartile range of the lower and upper quartiles, respectively.

Circles indicate each data point. See Table 3 for detailed statistics.

FIGURE 4　　　Relationships between color measurements of the throat feathers and pigmentation in the barn swallow: protoporphyrin IX (left column); PTCA (middle column); TTCA (right column). PITCA and TTCA are indices of eumelanin and pheomelanin, respectively. The upper, middle, and bottom rows show hue, saturation, and brightness, respectively. Red and pink circles (dark and light gray, in print) indicate adults and young, respectively. Simple regression lines are depicted for illustrative purpose (See Table 4 for detailed statistics).

FIGURE 5   Relationships between color change and pigmentation of the throat feathers in the barn swallow: protoporphyrin IX (left column); PTCA (middle column); TTCA (right column). The upper, middle, and bottom rows show hue, saturation, and brightness, respectively. Red and pink circles (dark and light gray, in print) indicate adults and young, respectively. Simple regression lines are depicted for illustrative purpose (See Table 5 for detailed statistics).

FIGURE 6 Boxplots of pigmentation of the throat feathers in relation to molting adults and adults and young captured during the breeding season in the barn swallow: protoporphyrin IX (upper column); PTCA (middle column); TTCA (lower column). PITCA and TTCA are indices of eumelanin and pheomelanin, respectively. The horizontal bar in each boxplot indicates the median, and the box shows the first and third quartiles of data. The whiskers range from the lowest to the highest data points within the 1.5 × interquartile range of the lower and upper quartiles, respectively. See Table 6 for detailed statistics.

794   TABLE S1 Dataset of the current study.

ID	Age	Treatment	post-H	post-S	post-V	pre-H	pre-S	pre-V	log(Porphyrin +1)	log(PTCA +1)	log(TTCA +1)
1	Adult	Control	21.45	133.38	101.56	21.88	130.54	98.92	3.78	1.63	2.22
2	Adult	Control	25.67	127.32	142.10	25.72	129.37	137.13	4.11	2.15	2.32
3	Adult	Control	26.24	130.55	142.55	26.36	130.09	138.23	5.29	2.07	2.10
4	Adult	Control	21.98	129.67	101.75	22.32	129.69	101.28	5.02	2.38	2.82
5	Young	Control	28.09	86.57	157.29	28.61	85.41	152.95	4.96	1.65	1.53
6	Young	Control	28.27	91.78	162.11	28.22	84.86	163.22	4.98	2.85	1.46
7	Adult	Control	26.07	125.70	142.63	25.75	126.21	138.40	2.83	3.06	2.64
8	Adult	Control	22.91	129.90	105.40	23.30	127.58	101.60	3.71	3.00	3.62
9	Young	Control	28.35	90.24	146.83	28.67	88.27	143.80	5.97	2.53	1.89
10	Young	Control	25.93	100.68	126.87	26.66	100.07	130.54	4.49	1.50	1.74
11	Adult	Control	24.06	134.87	112.31	24.15	132.09	109.56	4.03	3.33	4.05
12	Young	Control	27.22	93.63	139.98	27.74	89.94	139.10	4.36	3.01	2.44
13	Young	Control	25.91	97.38	122.24	26.23	95.69	119.07	6.00	3.18	2.86
14	Young	Control	26.89	95.81	141.78	27.46	94.60	138.96	5.53	2.30	1.93
15	Young	Control	28.58	102.06	165.18	28.28	102.55	164.67	3.87	2.73	1.19
16	Adult	Control	26.40	120.59	127.68	26.82	118.49	125.34	4.37	2.77	3.06
17	Adult	Control	26.91	119.79	135.90	26.43	121.57	130.49	3.14	2.31	2.52
18	Young	Control	25.29	113.32	118.05	25.02	110.84	118.18	4.30	3.14	2.07
19	Adult	Control	24.47	125.46	113.64	24.26	121.70	116.75	3.43	2.73	2.94
20	Young	Control	28.10	104.93	142.27	27.94	109.75	139.50	NA	2.42	2.15
1	Adult	UV	24.83	142.39	118.82	22.87	129.36	104.02	NA	2.17	3.29
2	Adult	UV	27.24	136.98	145.59	25.45	126.09	129.17	3.40	2.45	2.91
3	Adult	UV	26.84	135.62	138.31	25.40	130.64	126.04	3.22	2.85	2.86
4	Adult	UV	23.64	133.39	110.36	22.74	124.00	101.18	3.22	2.09	2.75
5	Young	UV	31.88	96.97	181.36	29.02	82.87	155.37	3.22	1.57	1.76
6	Young	UV	30.13	111.26	170.38	27.49	92.95	149.82	3.04	3.12	1.84
7	Adult	UV	29.04	129.23	163.54	26.12	125.06	146.18	3.14	2.44	2.67
8	Adult	UV	25.34	131.75	111.60	24.50	121.74	103.98	2.94	3.03	3.56
9	Young	UV	31.37	104.00	180.39	28.21	88.14	143.39	2.48	2.84	1.79
10	Young	UV	28.70	111.91	143.03	26.74	94.31	129.81	2.20	1.61	1.59
11	Adult	UV	25.93	140.78	125.20	24.24	130.69	110.87	2.64	2.60	3.12
12	Young	UV	30.59	110.00	174.26	27.98	92.49	145.98	0.00	2.76	1.82
13	Young	UV	29.75	111.20	163.61	27.91	90.09	133.14	2.48	1.81	1.93
14	Young	UV	29.59	114.53	165.88	27.21	95.26	134.51	1.95	2.31	1.89
15	Young	UV	30.96	115.04	184.86	28.33	108.84	162.76	1.95	3.21	2.08
16	Adult	UV	27.44	132.93	135.18	25.63	124.43	117.39	2.40	2.04	3.34
17	Adult	UV	27.13	137.89	146.28	25.07	130.29	127.84	1.95	3.13	3.15
18	Young	UV	28.21	125.57	149.22	25.57	108.46	121.39	0.00	3.32	2.64
19	Adult	UV	25.64	137.93	128.97	24.10	127.72	113.86	2.08	3.11	3.31
20	Young	UV	30.11	123.35	172.02	27.38	109.28	151.95	1.95	2.48	1.96

"Pre-" indicates measurement before the experiment, whereas "post-" indicates measurement after the experiment.
"NA" indicates "Not Available".

796   TABLE S2 Linear mixed-effects model explaining the color difference of the throat

797   feathers between treatments in relation to pigmentation difference between treatments in

798   the barn swallow ($n_{adult} = 9$, $n_{young} = 9$, $n_{total} = 18$).

Variables	Coefficient ± SE	t	p
Hue difference			
Protoporphyrin IX difference	-0.32 ± 0.15	-2.10	0.0545
PTCA difference	-0.69 ± 0.92	-0.75	0.47
TTCA difference	-0.24 ± 0.92	-0.26	0.80
Saturation			
Protoporphyrin IX difference	-2.56 ± 0.93	-2.74	0.016
PTCA difference	0.96 ± 2.88	0.33	0.74
TTCA difference	1.72 ± 2.56	0.67	0.52
Brightness			
Protoporphyrin IX difference	-6.27 ± 2.01	-3.12	<0.01
PTCA difference	-5.92 ± 12.29	-0.48	0.64
TTCA difference	-1.33 ±9.09	-0.15	0.89

799   PITCA and TTCA are indices of eumelanin and pheomelanin, respectively.

800   Note that, due to the small sample sizes in these analyses, in which small deviation from

801   normal distribution affects results, we here used robust linear models (function lmrob in

802   the R package robustbase) as before (Arai et al., 2017).

803   Independent variables were log-transformed before each analysis.

TABLE S3   Linear mixed-effects model explaining coloration of the throat feathers in relation to age and treatment with and without controlling for protoporphyrin IX in the barn swallow ($n_{adult,control} = 9$, $n_{young,control} = 9$, $n_{adult,UV} = 9$, $n_{young,UV} = 9$, $n_{total} = 36$).

Variables	Without controlling for protoporphyrin IX			With controlling for protoporphyrin IX		
	Coefficient ± SE	t	p	Coefficient ± SE	t	p
Hue						
Age	2.20 ± 0.68	3.25	<0.01	2.99 ± 0.77	3.87	<0.001
Treatment	1.50 ± 0.25	5.94	<0.0001	0.49 ± 0.39	1.27	0.22
Interaction	1.46 ± 0.36	4.08	<0.001	-0.03 ± 0.55	-0.07	0.95
	Variance explained by a random factor, ID = 1.78			Variance explained by a random factor, ID = 2.01		
Saturation						
Age	-30.26 ± 2.98	-10.15	<0.0001	-26.18 ± 2.84	-9.21	<0.0001
Treatment	8.07 ± 1.48	5.47	<0.0001	2.83 ± 1.78	1.59	0.13
Interaction	6.26 ± 2.09	3.00	<0.01	-1.49 ± 2.52	-0.59	0.56
	Variance explained by a random factor, ID = 30.20			Variance explained by a random factor, ID = 22.12		
Brightness						
Age	17.37 ± 7.75	2.24	0.037	24.54 ± 8.38	2.93	<0.01
Treatment	9.01 ± 2.96	3.04	<0.01	-0.21 ± 3.16	-0.67	0.95

Interaction	16.84 ± 4.19	4.02	<0.001	3.23 ± 4.47	0.72	0.48
	Variance explained by a random factor, ID = 230.86			Variance explained by a random factor, ID = 271.14		

Bird identity (ID) was included as a random factor in each model.

Note that sample sizes were reduced in this analysis because we lacked measurements in two samples (one in the adult UV treatment group and the other in the young control group). For the paired experimental design (see Methods), we had to exclude "unpaired" samples as well.

For analyses controlling for protoporphyrin IX pigmentation, we subtracted the estimated contribution of this pigment on color measurements (i.e., hue, saturation, brightness) using the coefficient of protoporphyrin IX and intercept using the models in Table 4. For analyses without controlling for protoporphyrin IX pigmentation, we subtracted the intercept alone (note that, although this model was quite similar to that in Table 1, we presented this model to compare models with and without controlling for protoporphyrin IX pigments using the same sample size).

TABLE S4   Linear mixed-effects model explaining plumage color change of the throat feathers in relation to age and treatment in the barn swallow with and without controlling for protoporphyrin IX ($n_{adult,control} = 9$, $n_{young,control} = 9$, $n_{adult,UV} = 9$, $n_{young,UV} = 9$, $n_{total} = 36$).

Variables	Without controlling for protoporphyrin IX			With controlling for protoporphyrin IX		
	Coefficient ± SE	t	p	Coefficient ± SE	t	p
Hue change						
Age	-0.22 ± 0.21	-1.03	0.31	0.46 ± 0.32	1.45	0.16
Treatment	1.71 ± 0.16	10.91	<0.0001	0.83 ± 0.29	2.88	0.01
Interaction	1.08 ± 0.22	4.88	<0.001	-0.21 ± 0.41	-0.52	0.61
	Variance explained by a random factor, ID = 0.09			Variance explained by a random factor, ID = 0.08		
Saturation change						
Age	1.35 ± 1.35	1.00	0.33	5.21 ± 2.11	2.48	0.021
Treatment	7.65 ± 1.09	7.00	<0.0001	2.68 ± 1.38	1.95	0.07
Interaction	6.56 ± 1.55	4.24	<0.001	-0.77 ± 1.95	-0.40	0.70
	Variance explained by a random factor, ID = 2.86			Variance explained by a random factor, ID = 11.40		
Brightness change						
Age	-1.70 ± 2.06	-0.83	0.42	5.08 ± 3.43	1.48	0.15
Treatment	11.48 ± 1.58	7.29	<0.0001	2.77 ± 2.00	1.38	0.19

Interaction	13.74 ± 2.23	6.17	<0.0001	0.86 ± 2.83	0.31	0.76
	Variance explained by a random factor, ID = 7.99			Variance explained by a random factor, ID = 34.77		

Bird identity (ID) was included as a random factor in each model.

Note that sample sizes were reduced in this analysis because we lacked measurements in two samples (one in the adult UV treatment group and the other in the young control group). For the paired experimental design (see Methods), we had to exclude "unpaired" samples as well.

For analyses controlling for protoporphyrin IX pigmentation, we subtracted the estimated contribution of this pigment on feather color change (i.e., hue change, saturation change, and brightness change) using the coefficient of protoporphyrin IX and intercept using the models in Table 5. For analyses without controlling for protoporphyrin IX pigmentation, we subtracted the intercept alone (note that, although this model was quite similar to that in Table 2, we presented this model to compare models with and without controlling for protoporphyrin IX pigments using the same sample size).

FIGURE S1   Reddish throat patch of a molting barn swallow. Note the color difference between newly molted (darker) and previously molted, faded (paler) feathers.

FIGURE S2   Relationships between color and pigmentation differences between treatments in the barn swallow: protoporphyrin IX (left column); PTCA (middle column); TTCA (right column). PITCA and TTCA are indices of eumelanin and pheomelanin, respectively. The upper, middle, and bottom rows show hue change, saturation change, and brightness change, respectively. Red and pink circles (dark and light gray, in print) indicate adults and young, respectively. Simple regression lines are depicted (see Table S4 for detailed statistics).