Deep-learning quantified cell-type-specific nuclear morphology predicts genomic

instability and prognosis in multiple cancer types

John Abel[1,*], Suyog Jain[1,#], Deepta Rajan[1,#], Harshith Padigela[1], Kenneth Leidal[1,#],

Aaditya Prakash[1,#], Jake Conway[1], Michael Nercessian[1], Christian Kirkup[1], Syed Ashar

Javed[1], Raymond Biju[1], Natalia Harguindeguy[1], Daniel Shenker[1], Nicholas Indorf[1],

Darpan Sanghavi[1], Robert Egger[1], Benjamin Trotter[1], Ylaine Gerardin[1], Jacqueline A.

Brosnan-Cashman[1], Aditya Dhoot[1], Michael C. Montalto[1#], Chintan Parmar[1], Ilan

Wapinski[1,#], Archit Khosla[1,#], Michael G. Drage[1,#], Limin Yu[1], Amaro Taylor-Weiner[1,#]

[1]PathAI; Boston, Massachusetts, United States

[#]Employee of PathAI at time of study

* **Correspondence to:**

John Abel, Ph.D.

PathAI

1325 Boylston Street, Suite 10000

Boston, Massachusetts 02215

Phone: 617-500-8457

Email: john.abel@pathai.com

Declaration of Competing interests: The authors declare the following competing

interests: J.A., S.J., D.R., H.P., K.L., A.P., J.C., M.N., C.K., S.A.J., R.B., N.H., D.Sh,

N.I., D.Sa., R.E., B.T., Y.G., J.A.B., A.D., M.C.M., C.P., I.W., A.K., M.G.D., L.Y., and

A.T.W. are currently, or were formerly, employed by and receive stock options from

PathAI, Inc., a company that builds artificial intelligence tools for pathology. In addition,

S.J. declares employment and stock from Meta, D.R. declares employment and stock

from Microsoft, and A.P. declares employment and stock from Spring Discovery.

ABSTRACT

While alterations in nucleus size, shape, and color are ubiquitous in cancer,

comprehensive quantification of nuclear morphology across a whole-slide histologic

image remains a challenge. Here, we describe the development of a pan-tissue, deep

learning-based digital pathology pipeline for exhaustive nucleus detection,

segmentation, and classification and the utility of this pipeline for nuclear morphologic

biomarker discovery. Manually-collected nucleus annotations were used to train an

object detection and segmentation model for identifying nuclei, which was deployed to

segment nuclei in H&E-stained slides from the BRCA, LUAD, and PRAD TCGA cohorts.

Interpretable features describing the shape, size, color, and texture of each nucleus

were extracted from segmented nuclei and compared to measurements of genomic

instability, gene expression, and prognosis. The nuclear segmentation and classification

model trained herein performed comparably to previously reported models. Features

extracted from the model revealed differences sufficient to distinguish between BRCA,

LUAD, and PRAD. Furthermore, cancer cell nuclear area was associated with increased

aneuploidy score and homologous recombination deficiency. In BRCA, increased

fibroblast nuclear area was indicative of poor progression-free and overall survival and

was associated with gene expression signatures related to extracellular matrix

remodeling and anti-tumor immunity. Thus, we developed a powerful pan-tissue

approach for nucleus segmentation and featurization, enabling the construction of

predictive models and the identification of features linking nuclear morphology with

clinically-relevant prognostic biomarkers across multiple cancer types.

INTRODUCTION

Histological assessment of tissue is central to the diagnosis and classification of malignancy, and critically informs patient management. Pathologists routinely report visible alterations in nuclear morphology. Altered nuclear features are ubiquitous in cancer, and changes in nuclear size, shape, coloration, texture, nucleoli, and nuclear-cytoplasmic ratio, as well as their intratumoral variance, are important features of histologic grade, which has prognostic relevance independent of disease stage[1,2]. The enumeration and morphologic features of mitoses also informs pathologist assessment of malignancy[3]. Furthermore, nuclear morphology can be important diagnostic features of certain cancers, such as nuclear clearing ("Ophan Annie Eyes") and pseudoinclusions of papillary thyroid carcinoma[4].

A complex interplay exists between nuclear morphology and the genetic, epigenetic, and transcriptomic milieu of cancer cells, reflecting the importance of the nucleus to the process of oncogenic transformation. Distorted nuclei can indicate dysregulated replication processes, aneuploidy, genomic instability, and genetic mutations that affect stability and function of the nuclear envelope[5]. Indeed, many cancers have altered expression of nuclear envelope components, resulting in nuclear rupture and micronuclei formation, further increasing genomic instability[5,6]. In addition, components of the nuclear envelope are known to bind to both chromatin and transcription factors, providing a spatial regulation to gene transcription and expression[5,7]. Therefore, the visual appearance of cancer cell nuclei has the potential to reveal key information about the biology of a tumor.

The quantitation of nuclear morphology has been a long sought-after goal[8]. Early studies used semi-quantitative approaches to enumerate features such as nuclear size and shape; these works revealed relationships between increased nuclear area and altered nuclear shape with poor prognosis and advanced disease in breast cancer and prostate cancer, respectively[9–11]. The use of computational approaches in pathology image analysis to identify and quantify nuclear changes has gained traction as modern computer vision methods have allowed for rapid, reproducible and cost-effective quantification of nuclear morphology. Using these methods, nuclear morphometric

features have been shown to correlate with relevant clinical and pathological metrics,

such as oligodendroglioma component in glioblastoma[12], as well as stage[13], disease

aggressiveness[14], recurrence[15–17], and outcome[18] in other cancer subtypes. In addition,

increased nuclear size has been correlated with whole genome duplication[19,20], and

nuclear morphometric features have allowed for the prediction of relevant molecular

information, such as ER status[21] and Oncotype DX risk scores[22,23] in breast cancer.

Most recently, Nimgaonkar et al. described an AI-derived histologic signature, the main

component of which was variance in nuclear morphology in cancer cells, that predicted

response to gemcitabine in patients with pancreatic adenocarcinoma[24].

96   Digitized whole slide images (WSIs) have enhanced the degree to which nuclear

morphology can be studied in histological specimens[12,13]. However, the large size of

WSIs - up to billions of pixels and containing thousands of nuclei - makes exhaustive

manual annotation infeasible; thus studies have relied on manually-selected subregions

of interest rather than entire slides[20,25]. Automated methods are, therefore, needed to

fully quantify nuclear features in WSIs. We recently described a cell- and tissue-level

computational pathology pipeline using WSIs for the automated computation of human

interpretable features (HIFs), distinctive features with tangible methods for validation[26].

This pipeline allows the use of HIFs to predict treatment-relevant molecular phenotypes

and allows for integration with current pathological methods. Given that morphological

analysis of histology features is central to pathology workflows, we sought to extend this

work to identify nuclear human interpretable features (nuHIFs) in multiple cancer types.

108   In this paper, we present a multi-tissue model for the exhaustive detection,

109   segmentation, and classification of nuclei from entire hematoxylin and eosin (H&E)-

110   stained WSIs, allowing for the exhaustive analysis of slide-level descriptors of nuclear

111   size, shape, texture, and staining intensity. Furthermore, we demonstrate that these

112   nuHIFs are predictive of clinically relevant information in multiple cancer types.

MATERIALS AND METHODS

Study Design

Manually collected annotations were used to train and validate an object detection and segmentation model to detect and segment nuclei from H&E stained tissue slides. Training data variation and number of annotations were selected to exceed previously used standards in the field[27] and exhibit wide variation in tissue morphology as subjectively assessed by study pathologists (MGD and LY). This model was deployed on whole-slide H&E images from The Cancer Genome Atlas (TCGA) to extract features from each nucleus in each slide, and the resulting features were used to analyze the relationship between nuclear morphology and underlying molecular markers of cancer, and patient outcomes. Inclusion of TCGA slides was performed in accordance with literature norms (e.g. as by Saltz et al.[28]): TCGA slides were selected to be the DX1 (primary diagnostic) slide for each case in TCGA and no outlier exclusion was performed, to conservatively reflect real-world conditions where same-case replicates may not be available. Where multiple hypotheses were tested, all reported statistics were corrected to control false discovery rate as described below.

Dataset Description and Annotation Collection

Over 29,000 manual annotations of cell nuclei were collected from H&E images from 21 tumor types at 40x and 20x magnification from TCGA[29], as well as a proprietary set of H&E-stained tissue biopsies of skin, liver non-alcoholic steatohepatitis, colon inflammatory bowel disease, and kidney lupus. Board-certified pathologists (MGD and LY) selected 1000 x 1000 pixel patches that were exemplary of varied tissue and nuclear morphology from the training slides and trained collaborators to perform exhaustive manual annotation of nuclei in the patches. Annotations were checked for quality, adjusted, and confirmed by MGD and LY. This process resulted in 67 WSI patches exhaustively annotated for nuclei. These patches were split into training, validation, and held-out test data sets to ensure distribution of tissue types (Table 1).

Following model training and initial testing, an additional two data sources were used to

collect additional annotations for model testing. H&E-stained slides of ulcerative colitis

were obtained from BioIVT (Westbury, NY), and H&E-stained breast cancer slides were

generously provided by Cleveland Clinic Foundation (Cleveland, OH). An additional 14

512 x 512 pixel patches were identified from these data sources (seven patches from

each source), and an additional 2,647 manual, exhaustive nucleus annotations were

collected for model evaluation (Table 2).

Nuclear Segmentation Model Architecture

A Mask-RCNN-style architecture was selected for nuclear segmentation. A ResNet50

backbone pretrained on the ImageNet dataset was used to produce the feature pyramid

network. The first two of five modules that comprise ResNet50 were frozen during

training to preserve the pretrained weights of early layers. Model development was

performed using the PyTorch library[30].

Nuclear Segmentation Model Training

The manually-collected annotations were used to train the model for detecting and

segmenting cellular nuclei (Figure 1A). During training, the annotated patches were

augmented by crops, flips, rotations, and affine deformations.

Cell Classification

Following nuclear segmentation, the cell class of each nucleus was assigned using

PathExplore™ (PathAI, Boston, MA)[31] models specific to breast cancer (BRCA), lung

adenocarcinoma (LUAD), and prostate adenocarcinoma (PRAD); PathExplore is for

research use only and is not for use in diagnostic procedures. Cancer epithelial cells,

fibroblasts, macrophages, lymphocytes and plasma cells were predicted for all three

cancer types, while additional cell classes were predicted for LUAD (granulocytes and

normal cells) and PRAD (smooth muscle cells, endothelial cells, and normal epithelial

cells). Model performance for the prediction of cell types was assessed by comparing

model predictions to pathologist annotations in nested pairwise fashion[32]. Model

performance metrics for BRCA, LUAD, and PRAD are shown in Supplementary Figures

S1-S3, respectively, and Supplementary Tables S1 and S2. Example prediction results

are shown in Figure 3. The five pan-indication cell classes (cancer epithelial cells,

fibroblasts, macrophages, lymphocytes, and plasma cells) were used for analyses

assessing the biological implications of nuclear feature differences in BRCA, LUAD, and

PRAD.

<u>Deployment Dataset and Feature Extraction</u>

The nuclear segmentation model was deployed on publicly available images of H&E

slides from the BRCA (N=886), PRAD (N=392), and LUAD (N=426) TCGA cohorts; a

summary of clinicopathologic features of each cohort is shown in Table 3. Model

performance was qualitatively assessed by board-certified pathologists and determined

to be consistent with performance on the held-out test dataset. The features computed

for each individual nucleus were: area, circularity, eccentricity, major and minor axis

length, perimeter, solidity, and the mean and standard deviation of pixel grayscale

intensity, pixel saturation, and pixel A and B channels in LAB colorspace. The mean and

standard deviation of each feature from each nucleus class on the slides were used to

summarize the nuclear morphology on each slide. This yielded 30 slide-level nuHIFs for

each cell type, e.g. the mean area of cancer nuclei, the standard deviation of fibroblast

nuclear eccentricity, or the mean pixel grayscale intensity of lymphocyte nuclei.

Attributes and features described by nuHIFs are included in Figure 1B. Thus, the total

number of features summarizing the morphology on each slide was 30 times the

number of cell classes.

<u>Exploring Cancer Type and Nuclear Morphology</u>

To compare the nuHIFs quantifying cancer cell, fibroblast, and lymphocyte morphology,

uniform manifold approximation and projection (UMAP) analysis was performed.

Nuclear HIFs were z-scored across all cancer types for standardization. UMAP was

parameterized with 100 neighbors, an embedding dimension of 2, and the Euclidean

distance metric. Features characteristic of each cancer type were evaluated by

averaging each feature across the samples of each cancer type and z-scoring for

visualization; hierarchical clustering (using Euclidean distance with average linkage)

identified features that varied across cancer types.

<u>Classifying Cancer Type from Nuclear Morphology</u>

Random forest (RF) binary classification models were trained and applied to each cell-

type-specific nuHIF set to differentiate between pairs of cancer types. RF classification

models were trained using 5-fold stratified cross-validation with balanced class

weighting. The performance of each model was assessed using the area under the

receiver operating characteristic curve (AUROC) on each held-out validation split. The

mean AUROC on the held-out validation splits is reported. RF model training was

performed in scikit-learn with default hyperparameters (100 trees)[33].

<u>Classifying Breast Cancer Subtype from Nuclear Morphology</u>

Characteristics of breast cancer molecular subtypes (luminal A, N=457; luminal B,

N=159; HER-2, N=66; normal-like, N=31; basal-like, N=161) were obtained from a prior

study by Berger and colleagues[34]. Random forest (RF) binary classification models

were trained and applied to each cell-type-specific nuHIF set to differentiate between

subtypes in a one-vs.-all manner. RF classification models and cross-validation

schemes were identical to cancer type classification.

<u>Statistical Analysis</u>

Spearman (rank-based) correlation was used to find the association between variation

in cancer nuclear morphology and metrics of genomic instability. Variation in size was

captured by the nuHIF "standard deviation of cancer cell nuclear area" for each slide.

For metrics of genomic instability, previously published metrics were selected:

aneuploidy score[35] and homologous recombination deficiency (HRD) score[36]. RF binary

classification models were trained in scikit-learn with default hyperparameters[33] using 5-

fold stratified cross-validation with balanced class weighting, and applied to the cancer

nuHIF set from each cancer type to predict binarized whole-genome doubling (WGD; 1-

2 doublings = 1; no doublings = 0). The performance of each model was evaluated

using AUROC on each held-out validation split, and the mean AUROC is reported. The

mean RF Gini importance (also called the mean decrease in impurity) of the top five

features for each cancer type across the five splits are reported. Cox proportional

hazard models were utilized to explore the relationship between BRCA fibroblast

nuHIFs and overall and progression-free survival (OS and PFS, respectively). Ordinal

tumor stage (1-4) and patient age were included as clinical covariates; 17 subjects

missing tumor stage and the one missing survival data were excluded. Robust z-scoring

(i.e. using the median and scaled interquartile range) of each nuHIF before modeling

was performed for simple interpretation of the hazard ratios (HRs). The p-values

associated with each nuHIF were corrected for false discovery rate (FDR) by the

Benjamini-Hochberg procedure. Survival analyses were performed using the Lifelines

library[37]. Gene expression data was acquired from the Genomic Data Commons (GDC)-

processed TCGA BRCA cohort (release 18.0) from the UCSC Xena data portal[38]. Gene

expression samples were paired to case-matched slides in our dataset, yielding 868

expression-nuHIF pairs. Spearman (rank-based) correlation was used to quantify the

association between bulk RNAseq expression and the mean fibroblast nucleus area

nuHIF for each gene and corrected for FDR via Benjamini-Hochberg procedure. Genes

with corrected $p < 0.05$ and Spearman correlation greater than 0.15 or less than -0.15

were selected to comprise the significant positively and negatively associated gene

sets, respectively, for gene set enrichment analysis (GSEA). GSEA[39] was performed

using the Molecular Signatures Database (MSigDB)[40] and the REACTOME pathway

database[41], and the ten most significant pathway overlaps, with FDR-corrected $p <$

0.05, are reported.

RESULTS

<u>Model development, performance, and nuclear feature extraction</u>

We collected annotations and trained a machine learning (ML) model to detect and segment nuclei in H&E-stained WSIs as described in the Methods and shown in Figure 1. The model is not limited to sampling regions of interest from tissue samples, but rather can be utilized to exhaustively annotate WSIs. Application of the model to our held-out test data, including held-out tissue and disease types, demonstrated performance (mean Dice score=0.818, aggregated Jaccard index (AJI) =0.619) comparable to models reported previously in the literature[27,42]. Importantly, model speed was adequate to apply to multi-gigabyte WSIs at full resolution (approximately 0.25 μm/pixel; roughly 30 minutes per slide). Examples of model performance in mesothelioma, head and neck squamous cell carcinoma, and stomach adenocarcinoma are shown in Figure 2.

We selected clinical samples from two additional datasets, designated OOD-Test-1 and OOD-Test-2, characterized in Table 2. We collected additional annotations on these datasets and characterized model performance. We found performance numerically comparable or superior to our initial held-out test data despite different sample origin, and one dataset containing non-cancer tissue samples (OOD-Test-1 mean Dice = 0.818, AJI = 0.628; OOD-Test-2 mean Dice = 0.826, AJI = 0.649).

Having evaluated our model's performance, we deployed the resulting model on primary diagnostic (DX1) H&E slides from the breast cancer (BRCA; N=892), prostate adenocarcinoma (PRAD; N=392), and lung adenocarcinoma (LUAD; N=426) TCGA cohorts (Figure 3); model performance was visually assessed to be consistent with test data. The distribution of pixel sizes (microns per pixel; MPP) of these three cohorts are shown in Supplementary Figure 4. The median MPPs were 0.248, 0.252, and 0.252 for BRCA, LUAD, and PRAD datasets respectively. We extracted interpretable features describing the shape, size, staining intensity, and texture of every nucleus on each WSI (Supplementary Table S3). We performed further analysis on nuHIFs specific to cancer

cells, fibroblasts, and lymphocytes, as these three cell classes are common across all

cancer types and have been implicated in clinical outcomes.

nuHIFs show within- and between-cancer type variation

To assess whether nuHIFs differ between cancer types, we performed UMAP to

compare the nuHIFs from cancer cells (Figure 4A), fibroblasts (Figure 4B), and

lymphocytes (Figure 4C) in BRCA, LUAD, and PRAD datasets. We observed notable

inter- and intra-dataset variation in nuHIFs. For cancer cells, nuclear morphology was

distinct between PRAD and LUAD datasets, while BRCA dataset cancer cells showed

nuclear features similar to both PRAD and LUAD (Figure 4A). Unsupervised hierarchical

clustering of z-scored features revealed specific nuHIFs differentially exhibited in these

three cancer subtypes (Figure 4B). For example, features associated with nuclear size

were higher in LUAD cancer nuclei relative to PRAD. Assessment of the distribution of

three size-related features in BRCA, LUAD, and PRAD confirmed these observations –

cancer nuclei in PRAD were smaller in area and major axis length than cancer nuclei in

BRCA and LUAD (Supplementary Figure 5A), while fibroblast area and major axis

length is larger in BRCA than in LUAD and PRAD (Supplementary Figure 5B). Minute

variation in minor axis length was observed between the three cancer types for cancer

cells and fibroblasts (Supplementary Figure 5C). In contrast, lymphocyte nucleus size

parameters did not appear to differ between cancer types (Supplementary Figure 5). In

addition to size features, features associated with nuclear staining were observed to

differ between cancer types. In particular, notable differences in features relating to

nucleus stain intensity, color, and shape between PRAD and LUAD were observed

(Figure 4). The clearest distinction between cancer subtypes was discerned through

nuHIFs of fibroblasts in BRCA, LUAD, and PRAD (Figure 4B,D). Unsupervised

hierarchical clustering revealed specific features enriched in fibroblasts from these

cancer subtypes. Interestingly, lymphocyte nuHIFs also differed between cancer types.

To ensure that the observed differences in nuclear features between cancer types were

not biased by scanned image pixel size, we measured the Pearson correlation between

nuclear size (using major axis length as a representative feature) and MPP for each cell

type within BRCA, LUAD, and PRAD datasets individually, to remove the potential effect

of possible between-cancer-type variation in nuclear size. The within-cancer-type

variation in mean nuclear major axis length between slides at the same MPP is large for

cancer cells (Supplementary Figure 6A), fibroblasts (Supplementary Figure 6B), and

lymphocytes (Supplementary Figure 6C). In addition, the magnitude of the within-

cancer-type Pearson correlations is low, although some rise to the level of significance,

perhaps due to the high power of the large dataset. The within-cancer-type Pearson

correlations also show an inconsistent sign, ranging from 0.206 to -0.151. Generally,

these results suggest that there is an inconsistent directional effect of MPP on nuclear

size, and other factors are likely driving the observed differences.

Because of the apparent association between nuclear morphology and cancer type, we

hypothesized that nuHIF-quantified nuclear morphology could be a distinguishing

feature of cancer types. To test this, we constructed a simple random forest binary

classification model for differentiating between each pair of cancer types (BRCA, PRAD,

LUAD) using cancer, fibroblast, or lymphocyte nuclear HIFs. We performed five-fold

cross validation to estimate the extent to which cancer types may be differentiated by

nuclear morphology. We found consistently strong performance for differentiating

between cancer types using nuclear morphology (Figure 4D). Although lymphocyte

nuclear morphology was less distinct between cancer types when visualized with

UMAP, supervised analysis indicated that lymphocyte morphology differed between

cancer types.

Cancer nuclear morphology is associated with metrics of genomic instability in multiple

cancer types

Cancer nuclear atypia is used clinically as a marker of malignancy. We therefore

hypothesized that underlying levels of genomic instability may partially explain the

observed heterogeneity in cancer nuclear morphology within cancer subtypes, as well

as between cancer types with known differences in malignancy. We tested this

hypothesis by assessing the relationship between cancer nuclear morphology and

genomic instability in LUAD, BRCA, and PRAD cohorts using aneuploidy score and

homologous recombination deficiency (HRD) score as metrics of genomic instability.

Indeed, using the standard deviation of cancer nuclear area as a metric of nuclear

atypia, we detected significant correlation between this nuHIF and both aneuploidy

score (Figure 5A) and HRD score (Figure 5B). When assessed in a pan-cancer manner,

the overall correlation increased, and the pattern observed in cancer nuHIF UMAP

analysis persisted: PRAD displayed a lower level of genomic instability across both

metrics compared to LUAD, while BRCA showed a wide range of genomic instability,

with similarities to both PRAD and LUAD. These results confirm that cancer nuclear

morphology, especially variability in nuclear size, is associated with the level of genomic

instability.

Because aneuploidy score was correlated to variation in cancer nuclear area, we

posited that cancer nuclear morphology was predictive of whole genome doubling. To

address this hypothesis, we trained random forest models for predicting binarized

whole-genome doubling using cancer nuHIFs from each of the BRCA, LUAD, and

PRAD cancer types. We found that cancer nuclear morphology was predictive of WGD

for each cancer type, with strongest predictive power in BRCA, and more variation in

performance expected for PRAD, where WGD occurs less frequently (Figure 5C). The

mean RF importance across the five splits is reported for the top five features for each

cancer type in Supplementary Table S4. Briefly, variation in cancer nuclear dimensions

were most important for predicting WGD in BRCA, mean cancer nuclear dimensions

were most important for predicting WGD in LUAD, and a mix of color and shape

features were found to be most important for PRAD.

<u>Nuclear morphology enables prediction of breast cancer molecular subtype</u>

We hypothesized that nuclear morphology would differ in subtle but meaningful ways

between molecular subtypes of breast cancer, and that these differences might enable

classification of molecular subtypes of breast cancer from H&E images. To test this

theory, we trained nuHIF-based classification models for predicting breast cancer

subtype in a one-vs.-all manner (Figure 6). Briefly, we found that cell-type-specific

nuclear morphology enabled classification of some but not all breast cancer molecular

subtypes. Interestingly, the ability to predict subtype varied by subtype as well as by cell

type being used to make the inference. Cancer nuclear morphology (Figure 6A) or

lymphocyte nuclear morphology (Figure 6C) enabled moderate prediction (AUROC >

0.7) of luminal A and basal-like breast cancer subtypes. Cancer nuclear morphology but

not lymphocyte or fibroblast nuclear morphology enabled moderate prediction of HER-2

breast cancer subtype. Interestingly, fibroblast nuclear morphology alone was a poor

predictor of molecular subtype (Figure 6B). When aggregating cell types (Figure 6D),

luminal A and basal-like prediction AUROC increased further to ≥ 0.80. These results

suggest that altered nuclear morphology is a possible histological presentation of breast

cancer molecular subtypes.

<u>Fibroblast nuclear morphology is associated with survival and gene expression patterns</u>
<u>in breast cancer</u>
The interplay between fibroblasts and cancer cells is complex and prognostically

relevant, as associations between cancer-associated fibroblasts (CAFs) and cancer

progression have been recently described[43–46]. Notably, in breast cancer, CAFs have

been shown to contribute to prognosis[47], while CAF subset heterogeneity correlates

with metastasis[48]. We therefore hypothesized that fibroblast nuHIFs in BRCA would be

clinically prognostic, independent of further molecular testing. We sought to identify

fibroblast nuHIFs that are associated with progression-free (PFS) and/or overall survival

(OS). We performed regression between each fibroblast nuHIF and PFS and OS using

Cox proportional hazards models with patient age and ordinal cancer stage as

regression covariates. After FDR correction, multiple fibroblast nuHIFs were significantly

prognostic of PFS (Supplementary Table S5) and OS (Supplementary Table S6).

Features quantifying the same general attribute, *e.g.* nuclear area and nuclear axis

length as measures of size, were indeed found to be correlated with one another (mean

pairwise Pearson r = 0.90 for fibroblast nuclear area, major axis length, minor axis

length, and perimeter). We selected the mean fibroblast nucleus area

("MEAN[FIBROBLAST_NUCLEUS_AREA]_H & E") for further evaluation, and show

Kaplan-Meier survival curves for PFS and OS for the population binarized by this

feature median value. High nuclear area was prognostic of worse outcomes (Figure 7,

PFS HR = 1.81, 95% CI [1.32-2.48], p = 0.0002; OS HR = 1.77, 95% CI [1.22, 2.56], p =

0.002).

Having identified this relationship between fibroblast nuclear size and prognosis, we

sought to assess whether mean fibroblast nuclear area was associated with differences

in bulk gene expression in breast cancer. We computed the rank-based (Spearman)

correlation between fibroblast mean nuclear area and each gene in TCGA bulk gene

expression to identify genes associated with this nuHIF (see Methods for details).

Fibroblast nuclear area was significantly, albeit weakly (absolute r > 0.15), associated

with expression of numerous individual genes (Supplementary Table S7). In contrast to

the weak associations observed at the individual gene level, gene set enrichment

analysis performed on the genes associated with morphology revealed significant

relationships between fibroblast nuclear size and levels of several previously identified

expression pathways. Notably, larger fibroblast nuclear size showed positive

association with gene expression in pathways associated with degradation and

remodeling of the extracellular matrix (Supplementary Table S8) indicating higher

fibroblast activity. Meanwhile, larger fibroblast nuclear size showed negative association

with the expression of genes in pathways relating to immune response to the tumor,

such as B cell receptor signaling and lymphoid cell interactions with non-lymphoid cells

(Supplementary Table S9). Taken together, these results suggest that fibroblast nuclear

morphology is indicative of underlying patterns of gene expression and is thus

biologically grounded.

DISCUSSION

In this study, we have presented a pan-tissue approach for nucleus segmentation, classification, and featurization on entire whole-slide pathology images. This method enabled the construction of predictive models and the identification of features linking nuclear morphology with quantitative biomarkers across BRCA, PRAD, and LUAD. These results highlight the potential of ML-enabled quantification of nuclear morphometry as a prognostic feature of many cancer types and a potential biomarker to be used by pathologists. Furthermore, this approach enables the quantitative testing of hypotheses and numerical quantification of histological relationships proposed by pathologists (*e.g.*, by establishing a numerical relationship between nuclear atypia and disease metrics). In addition, our approach enables the data-driven identification of sub-visual changes that may be clinically meaningful.

One particular strength of our approach is the ability to not only measure morphologic features associated with nuclei in a cancer specimen, but to assign a cell class to each nucleus, as well. To our knowledge, this work provides the first characterization of nuclear morphologies of specific cell types in different cancers at scale. As such, we were not only able to assess the associations of cancer cell nuclear morphology with clinically relevant metrics, but we were also able to examine these relationships using nuclear features of fibroblasts and lymphocytes. For example, fibroblast nuHIFs provided a clear separation of cancer types in both unsupervised and supervised analyses, indicating that the nuclear morphologies of fibroblasts differ in breast, lung, and prostate cancers. Given recent observations that CAFs can be classified into multiple functional subtypes based on gene expression[49], the distinctive nuclear morphologies seen in fibroblasts of breast, lung, and prostate cancers suggests that fibroblasts may contribute to cancer progression differently in these three cancer types. Importantly, we cannot distinguish between the multiple known subtypes of intratumoral fibroblasts using the approach described herein. This caveat is particularly relevant to the associations of fibroblast nuclear morphology with gene expression in breast cancer. Increased nuclear size was positively associated with an extracellular matrix

remodeling gene expression profile and negatively associated with the expression of

genes relating to anti-tumor immune response (Supplementary Tables S4, S5).

Interestingly, single-cell analysis of fibroblasts in breast cancer has revealed several

disparate populations, including an immunosuppressive population characterized by the

expression of genes involved in collagen production and extracellular matrix remodeling

and a separate class with an inflammatory gene expression profile[50]. While our model

cannot directly predict the presence of these fibroblast sub-populations, given the

prognostic associations of nuclear morphology in our dataset and sc-RNAseq

expression[50], it will be of interest to test whether specific nuclear features of CAFs

associate with functional subtypes.

Furthermore, nuclear features derived from our model were associated with PFS and

OS in breast cancer. It is worth noting that this analysis, while incorporating patient age

and clinical stage as regression covariates, was conducted on a large cohort of patients

across study sites for whom relevant clinical information (e.g., treatment history) was

not readily available. Therefore, while our result linking fibroblast nuclear morphology to

prognosis in breast cancer is intriguing, further study in more controlled patient cohorts

is needed to confirm this observation.

Herein, we observed that nuclear morphology differed between cancers as assessed

using nucleus segmentation models. This result was observed not only for cancer

epithelial cells and fibroblasts, but also, surprisingly, for lymphocytes. However, caution

is warranted in interpretation – it is plausible that batch effects between slides from

different tumor groups could drive variation in nuclear presentation, especially due to

differences in pre-analytic variables such as slide preparation and staining. However, it

is also plausible that this finding reflects the differences in genetic and epigenetic

landscapes between tumor types, levels of genomic instability, and overall differences in

cancer evolution between these cancer types that may manifest as disparate nuclear

morphologies.

The observed relationship between greater variation in cancer nuclear area and

genomic instability was consistent across cancer types, indicating a quantitative link

between nuclear pleomorphism and genomic instability pertinent to numerous cancer

histologies. Prior analyses have noted an association between increased variation in

nuclear size and whole genome doubling, suggesting a direct link between variation in

nuclear size and genomic instability[19,20]. Additional work has noted a correlation

between nuclear morphology and HDR in luminal and triple-negative breast cancer[51].

Given that nuclear size reflects DNA content, variation in nuclear size features between

cells may be linked to underlying genomic instability. Similarly, recent work identified a

histologic signature based on variability in nuclear morphology in pancreatic cancer

cells that was associated with improved response to gemcitabine but was not

associated with a previously defined gene expression-based disease subtype[24].

Pancreatic cancer patients with BRCA1/2 mutations, associated with increased genomic

instability, are known to respond more favorably to therapy regimens involving

gemcitabine [52]; thus, our result that nuclear variation is associated with genomic

instability may explain this recent finding. To this end, our observation that variability in

nuclear size (measured here by standard deviation of cancer cell nuclear area) is

consistent with these prior hypotheses and allows for them to be tested on a larger

scale for each case (all cells for each cell type in the WSI). While the biological result

linking nuclear morphology with genomic instability is not novel, the observation of this

expected result through the analyses of our novel model-derived nuclear features

indicates that our approach supports the technical robustness and biological

applicability of our approach.

One mitigation to potential batch effects is to analyze nuclear morphology within a

single cancer type, and additionally to focus on size and shape features that are more

likely to be robust to tissue preparation variabilities. For example, in breast cancer, we

observed a clear relationship between fibroblast nuclear size, prognosis, and gene

expression patterns. In breast cancer, increased fibroblast nuclear area was positively

correlated with gene expression in extracellular matrix remodeling pathways and

negatively correlated with genes in anti-tumor immune response pathways. The CAF

subtypes present in a breast cancer sample may impact the tumor immune

microenvironment[49]. While it would be interesting to posit that fibroblast nuclear

morphology could reflect these subtypes, the ability to explore this is precluded by the

use of bulk RNAseq data, since fibroblast nuclear features and the bulk expression

profiling reflect a summarization of a whole slide. However, because nuclear

morphology is quantified at single-cell resolution, this approach could be tied directly to

single-cell expression analysis. Further work is necessary to delineate the functional

relevance of nuclear morphology changes in fibroblasts in cancer.

As noted, batch effects have the potential to influence the interpretation of model

outputs due to data that are aggregated across different sites, sources, and preparation

laboratories. Pixel size variability, due to slide scanning with different MPP resolution, is

one aspect of how these differences may manifest, but there are others to consider as

well: differences in stain reagents, sample preparation, sample storage, or other pre-

analytical variables. For the analyses described herein, the median MPP values were

highly similar across the three indications, with the BRCA MPP slightly lower than that

of LUAD and PRAD (Supplementary Figure S4). That said, to further ensure against

differences in pixel dimension contributing to bias, the size-related features of the nuclei

are reported here in units of microns or square microns, which is created by multiplying

the size of the mask by the appropriate MPP conversion factor. Thus, differences in the

MPP should not propagate into length-features, and the slide scan characteristics

should not bias the features. Furthermore, we measured the Pearson correlation

between nuclear size (using major axis length as a representative feature) and MPP for

each cell type within BRCA, LUAD, and PRAD datasets individually to eliminate the

potential effect of possible inter-cancer-type variation in nuclear size (Supplementary

Figure S6). While the within-cancer-type variation in mean nuclear major axis length

between slides at the same MPP is large, the magnitude of the within-cancer-type

Pearson correlations is low, although some rise to the level of significance (likely due to

the high power of the large datasets). Lastly, it is worth noting that cancer-type

differences in nuclear size appear to be an outlier of relatively larger magnitude than

expected if MPP bias was the primary driving factor (Supplementary Figure S5). Thus,

we are confident that the observations noted in this study regarding nuclear size

features are not biased by scan-specific metrics.

The approach that we undertook for nucleus segmentation and morphometry analysis in

this paper has several key strengths. First, the ability to compute human-interpretable

nuclear features at scale enables testing quantitative biological hypotheses, rather than

relying on by-eye estimation of parameters such as variation in nuclear morphology.

The ability to perform these analyses on WSIs of H&E-stained cancer tissue additionally

obviates the need to hand-select regions of interest, which may contribute to biased

analyses. In addition, we were able to train and deploy our model on tissues from

diverse cancer types, suggesting that the model can be readily deployed on samples

from varied cancer indications[53].

A particular strength of this approach is the interpretability of the predictions made.

While HIF-based predictive clinical models are inherently less flexible than end-to-end

black-box approaches (and, thus, can yield lower performance), they benefit from the

lower dimensionality of features as a method of regularization, as the HIFs used herein

directly map to low-dimensional representations of the tissue image. Furthermore, HIF-

based models allow researchers and clinicians to learn from the features and generate

novel hypotheses without discarding the wealth of known biology.

Although our results point to the potential of nuclear segmentation, classification, and

feature analysis as a clinical screening tool, our study is limited in that our biomarker

analysis was focused on academically curated datasets. These datasets were selected

due to their size, completeness, and rich genomic and transcriptomic profiling data.

Construction and validation of generalizable predictive machine-learning models

requires the inclusion of a broad range of training and validation data, and future efforts

should focus on validating these hypotheses in additional cohorts. The technical

approaches we describe here have been validated by their application to other clinical

data sets, showing their generalizability of this methodology and robustness of these

models (data not shown).

In sum, this work highlights the power of ML-driven quantitative nuclear morphometry in multiple cancer types. The models and resulting features described herein have the potential not only to aid pathologists and research teams in discerning novel biomarkers but to provide meaningful prognostic information for cancer patients. The ability to measure these features robustly and consistently at scale may enable the development of improved clinical tools for advancing precision medicine.

ACKNOWLEDGEMENTS

The authors would like to thank the software engineering and machine learning operations teams at PathAI for developing the systems and pipelines used for model development and feature extraction. The authors also thank SciStories, LLC and GCI Health for assistance with figure design.

FUNDING

This work was funded by PathAI, Inc.

AUTHOR CONTRIBUTIONS

Conceptualization: J.A., S.J., D.J., A.P., M.G.D., L.Y., A.T.W.

Methodology: J.A., S.J., D.R., H.P., K.L., A.P., J.C., M.N., C.K., S.A.J., N.I., D.Sa., R.E., B.T., Y.G., A.D., C.P., I.W., A.K., M.G.D., L.Y., A.T.W.

Investigation: J.A., S.J., D.R., H.P., K.L., A.P., J.C., M.N., C.K., S.A.J., R.B., N.H., D.Sh., D.Sa., R.E., B.T., Y.G., A.D., C.P., M.G.D., L.Y., A.T.W.

Visualization: J.A., Y.G., J.A.B., I.W., A.T.W.

Funding acquisition: N/A

Project administration: J.A., D.Sa, C.P., I.W., A.T.W.

Supervision: M.C.M., I.W., A.K., L.Y., A.T.W.

Writing – original draft: J.A., Y.G., J.A.B.

Writing – review & editing: J.A., S.J., D.R., H.P., K.L., A.P., J.C., M.N., C.K., S.A.J.,

R.B., N.H., D.Sh., N.I., D.Sa., R.E., B.T., Y.G., J.A.B., A.D., M.C.M., C.P., I.W., A.K.,

M.G.D., L.Y., A.T.W.

DATA AND MATERIALS AVAILABILITY

Histopathology images from the Cancer Genome Atlas dataset are available at

https://www.cancer.gov/about-nci/organization/ccg/research/structural-genomics/tcga.

Images and annotations of nuclei from the following datasets will be made available on

GitHub prior to publication: training set, validation set, test set, OOD-2 dataset, and

TCGA datasets. Images and annotations of nuclei from the OOD-1 dataset will be

shared upon written request. Access to feature tables, cell-, tissue-, and nuclei-type

heatmaps, as well as usage of cell- and tissue-type classification models, are available

upon reasonable request to academic investigators without relevant conflicts of interest

for non-commercial use who agree not to distribute the data. Access requests can be

made to publications@pathai.com.

CODE AVAILABILITY

Model parameters for nuclei, cell, and tissue models, and codes model training,

inference, and feature extractions are not disclosed. Access requests for such a code

will not be considered to safeguard PathAI's intellectual property. All source code for

reproducing correlational analyses and molecular predictions, will be deposited to

GitHub prior to publication, and the link will be provided at that time.

FIGURE LEGENDS

Figure 1. Machine learning model annotation collection, training, and application.
(A) Model workflow. Briefly, pathologists trained expert annotators to perform
exhaustive annotations of nuclei on H&E slide patches from diverse tissue sources.
These were used to train a pan-H&E nucleus detection and segmentation model, which
was subsequently evaluated on held-out patches and applied to exhaustively segment
nuclei in three WSI datasets. **(B)** Features extracted from the model. Mean and
standard deviation values were calculated for these features at the whole-slide level for
cancer cells, lymphocytes, and fibroblasts.

Figure 2. Example of model performance. Representative WSI patches from
mesothelioma, head and neck squamous cell carcinoma (HNSCC), and stomach
adenocarcinoma stained with H&E are shown in the left-most panel. Ground truth nuclei
identified manually and nuclei predicted by the model are shown in the middle and right-
most panels, respectively. Each color represents a nucleus instance.

**Figure 3. Nuclear segmentation and cell type identification in multiple cancer
types.** Representative H&E images of **(A)** breast cancer (TCGA BRCA), **(B)** lung
adenocarcinoma (TCGA LUAD), and **(C)** prostate adenocarcinoma (TCGA PRAD) are
shown at 40X magnification. **(D-F)** Nuclear segmentation and cell type identification
masks are overlaid onto H&E images shown in panels A-C. **(H-I)** High-magnification
images of BRCA **(G)**, LUAD **(H)**, and PRAD **(I)**. Magnified regions are indicated by
dashed boxes in panels D-F. Scale bars indicate a distance of 50 μm.

Figure 4. nuHIFs show variation within and between cancer types. Uniform
manifold approximation and projection (UMAP) visualization of BRCA, LUAD, and
PRAD defined by nuclear human interpretable feature (HIF) for **(A)** cancer cells, **(B)**
fibroblasts, and **(C)** tumor-infiltrating lymphocytes. Clustered heatmaps of median Z-
scores for all 30 nuHIFs are shown for each cell type. **(D)** Receiver operating
characteristic (ROC) curves for binary classification between paired cancer types using

nuHIFs from each of cancer, fibroblast, or lymphocyte nuclei. ROCs are shown for the

five held-out validation splits and mean area under ROC (AUROC) is shown for each

classification problem. In particular, fibroblast and lymphocyte nuclear features are

highly able to differentiate between cancer types. Mean AUROC is shown for each class

of nuclear HIF.

Figure 5. Variation in cancer nuclear size correlates with metrics of genomic

instability. Standard deviation of cancer cell nuclear area was compared to **(A)**

aneuploidy score and **(B)** homologous recombination deficiency (HRD) score for BRCA,

LUAD, and PRAD. **(C)** Receiver operating characteristic (ROC) curves for prediction of

whole-genome doublings in BRCA, LUAD, and PRAD. ROCs are shown for the five

held-out validation splits; mean AUROC is shown for each cancer type.

Figure 6. Cell-type-specific nuclear morphology enables classification of breast

cancer molecular subtypes. One-vs-all binary classification of breast cancer

molecular subtypes (luminal A, luminal B, HER2-like, basal-like, and normal-like)[34] was

performed using random forest classification on nuHIFs derived from A) cancer cells, B)

fibroblasts, C) lymphocytes, and D) aggregated cell types. Five-fold stratified cross-

validation was used, and mean AUROC for each of the iteratively held-out test sets is

reported here.

Figure 7. Association between fibroblast nuclear area and survival in breast

cancer. Increased fibroblast nuclear area ($\geq$ 50th percentile) corresponds to poor PFS

(HR = 1.8163, 95% CI [1.3119-2.4823], p = 0.0002) and OS (HR = 1.7753, 95% CI

[1.2206, 2.5620], p = 0.0022).

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure S1. Performance of cell classification model in breast cancer. A) Comparison of model-predicted cell types to average pathologist annotations. The bar graph at the top left depicts the breakdown of average pathologist annotations for each class of model prediction (precision). The bar graph at the bottom right shows the breakdown of model predictions for each class of average pathologist annotation (recall). "Other Than Specified" refers to predictions of classes other than those listed, or background class. **B)** Agreement between model-derived cell counts and pathologist consensus counts . **C**) Precision, Recall, and F1 scores of model predictions compared to pathologists' annotations in nested pairwise fashion[32]. **D**) Difference in the nested pairwise metric comparing mean difference between model and individual pathologist performance. Positive values indicate that the model out-performed pathologists when evaluated against held-out pathologists, while negative values indicate the model under-performed pathologists. Confidence intervals were obtained by bootstrapping.

Supplementary Figure S2. Performance of cell classification model in NSCLC. A) Comparison of model-predicted cell types to average pathologist annotations. The bar graph at the top left depicts the breakdown of average pathologist annotations for each class of model prediction (precision). The bar graph at the bottom right shows the breakdown of model predictions for each class of average pathologist annotation (recall). "Other Than Specified" refers to predictions of classes other than those listed, or background class. **B)** Agreement between model-derived cell counts and pathologist consensus counts . **C**) Precision, Recall, and F1 scores of model predictions compared to pathologists' annotations in nested pairwise fashion[32]. **D**) Difference in the nested pairwise metric comparing mean difference between model and individual pathologist performance. Positive values indicate that the model out-performed pathologists when evaluated against held-out pathologists, while negative values indicate the model under-performed pathologists. Confidence intervals were obtained by bootstrapping.

Supplementary Figure S3. Performance of cell classification model in prostate

cancer. A) Comparison of model-predicted cell types to average pathologist

annotations. The bar graph at the top left depicts the breakdown of average pathologist

annotations for each class of model prediction (precision). The bar graph at the bottom

right shows the breakdown of model predictions for each class of average pathologist

annotation (recall). "Other Than Specified" refers to predictions of classes other than

those listed, or background class. **B)** Agreement between model-derived cell counts and

pathologist consensus counts . **C**) Precision, Recall, and F1 scores of model predictions

compared to pathologists' annotations in nested pairwise fashion[32]. **D**) Difference in the

nested pairwise metric comparing mean difference between model and individual

pathologist performance. Positive values indicate that the model out-performed

pathologists when evaluated against held-out pathologists, while negative values

indicate the model under-performed pathologists. Confidence intervals were obtained by

bootstrapping.

Supplementary Figure S4. Distribution of pixel sizes (in MPP) across the three TCGA datasets (BRCA, LUAD, PRAD) used in this study.

Supplementary Figure S5. Distributions of nuclear size features in BRCA, LUAD, and PRAD datasets. A) Distribution of area. B) Distribution of major axis length. C) Distribution of minor axis length.

Supplementary Figure S6. Pearson correlation between nuclear size (using major axis length as a representative feature) and MPP for each cell type within BRCA, LUAD, and PRAD datasets. A) Correlation between major axis length and MPP in cancer epithelial cells. B) Correlation between major axis length and MPP in fibroblasts. C) Correlation between major axis length and MPP in lymphocytes.

REFERENCES

1. Zink, D., Fischer, A. H. & Nickerson, J. A. Nuclear structure in cancer cells. *Nat. Rev. Cancer* **4**, 677–687 (2004).

2. Fischer, E. G. Nuclear Morphology and the Biology of Cancer Cells. *Acta Cytol.* **64**, 511–519 (2020).

3. Cree, I. A. *et al.* Counting mitoses: SI(ze) matters! *Mod. Pathol.* **34**, 1651–1657 (2021).

4. Hapke, M. R. & Dehner, L. P. The optically clear nucleus. A reliable sign of papillary carcinoma of the thyroid? *Am. J. Surg. Pathol.* **3**, 31–38 (1979).

5. Chow, K.-H., Factor, R. E. & Ullman, K. S. The nuclear envelope environment and its cancer connections. *Nat. Rev. Cancer* **12**, 196–209 (2012).

6. Shah, P., Wolf, K. & Lammerding, J. Bursting the bubble--nuclear envelope rupture as a path to genomic instability? *Trends Cell Biol.* **27**, 546–555 (2017).

7. Zuleger, N., Robson, M. I. & Schirmer, E. C. The nuclear envelope as a chromatin organizer. *Nucleus* **2**, 339–349 (2011).

8. Stenkvist, B. O. *et al.* Computerized Nuclear Morphometry as an Objective Method for Characterizing Human Cancer Cell Populations1.

9. Baak, J. P. A., Van Dop, P. H., Kurver, t. P. E. T. H. & Hermans, J. O. The Value of Morphometry to Classic Prognosticators in Breast Cancer. doi:10.1002/1097-0142(19850715)56:2.

10. Pienta, K. J. & Coffey, D. S. Correlation of Nuclear Morphometry With Progression of Breast Cancer. doi:10.1002/1097-0142(19911101)68:9.

11. Partin, A. W. *et al.* Use of Nuclear Morphometry, Gleason Histologic Scoring,

Clinical Stage, and Age to Predict Disease-Free Survival Among Patients With

Prostate Cancer. doi:10.1002/1097-0142(19920701)70:1.

12. Kong, J. *et al.* Machine-based morphologic analysis of glioblastoma using whole-

slide pathology images uncovers clinically relevant molecular correlates. *PLoS One*

8, e81049 (2013).

13. Kumar, N. *et al.* Computer-extracted features of nuclear morphology in hematoxylin

and eosin images distinguish stage II and IV colon tumors. *J. Pathol.* **257**, 17–28

(2022).

14. Lewis, J. S., Jr, Ali, S., Luo, J., Thorstad, W. L. & Madabhushi, A. A quantitative

histomorphometric classifier (QuHbIC) identifies aggressive versus indolent p16-

positive oropharyngeal squamous cell carcinoma. *Am. J. Surg. Pathol.* **38**, 128–137

(2014).

15. Tokuyama, N. *et al.* Prediction of non-muscle invasive bladder cancer recurrence

using machine learning of quantitative nuclear features. *Mod. Pathol.* (2021)

doi:10.1038/s41379-021-00955-y.

16. Ji, M.-Y. *et al.* Nuclear shape, architecture and orientation features from H&E

images are able to predict recurrence in node-negative gastric adenocarcinoma. *J.*

Transl. Med. **17**, 92 (2019).

17. Lee, G. *et al.* Nuclear Shape and Architecture in Benign Fields Predict Biochemical

Recurrence in Prostate Cancer Patients Following Radical Prostatectomy:

Preliminary Findings. *Eur Urol Focus* **3**, 457–466 (2017).

18. Yu, K.-H. *et al.* Predicting non-small cell lung cancer prognosis by fully automated

microscopic pathology image features. *Nat. Commun.* **7**, 12474 (2016).

19. Cancer Genome Atlas Research Network. Comprehensive and Integrated Genomic

Characterization of Adult Soft Tissue Sarcomas. *Cell* **171**, 950–965.e28 (2017).

20. Fu, Y. *et al.* Pan-cancer computational histopathology reveals mutations, tumor

composition and prognosis. *Nature Cancer* **1**, 800–810 (2020).

21. Rawat, R. R., Ruderman, D., Macklin, P., Rimm, D. L. & Agus, D. B. Correlating

nuclear morphometric patterns with estrogen receptor status in breast cancer

pathologic specimens. *NPJ Breast Cancer* **4**, 32 (2018).

22. Whitney, J. *et al.* Quantitative nuclear histomorphometry predicts oncotype DX risk

categories for early stage ER+ breast cancer. *BMC Cancer* **18**, 610 (2018).

23. Li, H. *et al.* Quantitative nuclear histomorphometric features are predictive of

Oncotype DX risk categories in ductal carcinoma in situ: preliminary findings.

Breast Cancer Res. **21**, 114 (2019).

24. Nimgaonkar, V. *et al.* Development of an artificial intelligence-derived histologic

signature associated with adjuvant gemcitabine treatment outcomes in pancreatic

cancer. *Cell Rep Med* **0**, (2023).

25. Joshi, R. P. *et al.* Imaging-based histological features are predictive of MET

alterations in Non-Small Cell Lung Cancer. *arXiv [cs.CV]* (2022).

26. Diao, J. A. *et al.* Human-interpretable image features derived from densely mapped

cancer pathology slides predict diverse molecular phenotypes. *Nat. Commun.* **12**,

1613 (2021).

27. Kumar, N. *et al.* A Multi-Organ Nucleus Segmentation Challenge. *IEEE Trans. Med.*

Imaging **39**, 1380–1391 (2020).

28. Saltz, J. *et al.* Spatial Organization and Molecular Correlation of Tumor-Infiltrating

Lymphocytes Using Deep Learning on Pathology Images. *Cell Rep.* **23**, 181–

193.e7 (2018).

29. Gutman, D. A. *et al.* Cancer Digital Slide Archive: an informatics resource to

support integrated in silico analysis of TCGA pathology data. *J. Am. Med. Inform.*

Assoc. **20**, 1091–1098 (2013).

30. Paszke, A. *et al.* PyTorch: An imperative style, high-performance deep learning

library. *arXiv [cs.LG]* (2019).

31. Markey, M. *et al.* Abstract B010: Spatially-resolved prediction of gene expression

signatures in H&E whole slide images using additive multiple instance learning

models. *Mol. Cancer Ther.* **22**, B010–B010 (2023).

32. Gerardin, Y. *et al.* Improved statistical benchmarking of digital pathology models

using pairwise frames evaluation. *arXiv [cs.CV]* (2023)

doi:10.48550/ARXIV.2306.04709.

33. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *arXiv [cs.LG]* 2825–

2830 (2012).

34. Berger, A. C. *et al.* A Comprehensive Pan-Cancer Molecular Study of Gynecologic

and Breast Cancers. *Cancer Cell* **33**, 690–705.e9 (2018).

35. Taylor, A. M. *et al.* Genomic and Functional Approaches to Understanding Cancer

Aneuploidy. *Cancer Cell* **33**, 676–689.e3 (2018).

36. Marquard, A. M. *et al.* Pan-cancer analysis of genomic scar signatures associated

with homologous recombination deficiency suggests novel indications for existing

cancer drugs. *Biomark Res* **3**, 9 (2015).

37. Davidson-Pilon, C. lifelines: survival analysis in Python. *J. Open Source Softw.* **4**,

853  1317 (2019).

854 38. Goldman, M. J. *et al.* Visualizing and interpreting cancer genomics data via the

855  Xena platform. *Nat. Biotechnol.* **38**, 675–678 (2020).

856 39. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based

857  approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.*

858  *U. S. A.* **102**, 15545–15550 (2005).

859 40. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,

860  1739–1740 (2011).

861 41. Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids*

862  *Res.* **50**, D687–D692 (2022).

863 42. Graham, S. *et al.* Hover-Net: Simultaneous segmentation and classification of

864  nuclei in multi-tissue histology images. *Med. Image Anal.* **58**, 101563 (2019).

865 43. Friedman, G. *et al.* Cancer-associated fibroblast compositions change with breast

866  cancer progression linking the ratio of S100A4+ and PDPN+ CAFs to clinical

867  outcome. *Nat Cancer* **1**, 692–708 (2020).

868 44. Becker, L. M. *et al.* Epigenetic Reprogramming of Cancer-Associated Fibroblasts

869  Deregulates Glucose Metabolism and Facilitates Progression of Breast Cancer.

870  *Cell Reports* vol. 31 107701 Preprint at

871  https://doi.org/10.1016/j.celrep.2020.107701 (2020).

872 45. Suh, J., Kim, D., Lee, Y., Jang, J. & Surh, Y. Fibroblast growth factor-2, derived

873  from cancer-associated fibroblasts, stimulates growth and progression of human

874  breast cancer cells via FGFR1 signaling. *Molecular Carcinogenesis* vol. 59 1028–

875  1040 Preprint at https://doi.org/10.1002/mc.23233 (2020).

46. Houthuijzen, J. M. & Jonkers, J. Cancer-associated fibroblasts as key regulators of

the breast cancer tumor microenvironment. *Cancer Metastasis Rev.* **37**, 577–597

(2018).

47. Ren, J. *et al.* Cancer-associated fibroblast-derived Gremlin 1 promotes breast

cancer progression. *Breast Cancer Res.* **21**, 109 (2019).

48. Pelon, F. *et al.* Cancer-associated fibroblast heterogeneity in axillary lymph nodes

drives metastases in breast cancer through complementary mechanisms. *Nature*

Communications vol. 11 Preprint at https://doi.org/10.1038/s41467-019-14134-w

(2020).

49. Costa, A. *et al.* Fibroblast Heterogeneity and Immunosuppressive Environment in

Human Breast Cancer. *Cancer Cell* **33**, 463–479.e10 (2018).

50. Kieffer, Y. *et al.* Single-Cell Analysis Reveals Fibroblast Clusters Linked to

Immunotherapy Resistance in Cancer. *Cancer Discov.* **10**, 1330–1351 (2020).

51. Lazard, T. *et al.* Deep learning identifies morphological patterns of homologous

recombination deficiency in luminal breast cancers from whole slide images. *Cell*

Rep Med **3**, 100872 (2022).

52. Rosen, M. N., Goodwin, R. A. & Vickers, M. M. BRCA mutated pancreatic cancer:

A change is coming. *World J. Gastroenterol.* **27**, 1943–1958 (2021).

53. Michener, C. M. *et al.* 593P AI-powered analysis of nuclear morphology associated

with prognosis in high-grade serous carcinoma. *Ann. Oncol.* **33**, S818 (2022).

900 **TABLES**

902 **Table 1. Samples used for training and evaluating the segmentation model.**

Source	# Frames	Magnification	Microns Per Pixel (MPP)	Indication	Dataset
TCGA-A2-A3XS-01Z-00-DX1	3	40X	0.2456	Breast, Infiltrating Ductal Carcinoma	Train
TCGA-XF-AAMJ-01Z-00-DX1	2	40X	0.2527	Bladder, Urothelial Carcinoma	Train
TCGA-UW-A72Q-01Z-00-DX1	3	40X	0.2472	Liver, Hepatocellular Carcinoma	Train
TCGA-G3-A7M6-01Z-00-DX1	2	40X	0.2527	Liver, Hepatocellular Carcinoma	Train
TCGA-KM-	1	40X	0.2525	Kidney, Renal Cell Carcinoma	Train

A7QL-01Z-00-DX1					
TCGA-CV-A6JD-01Z-00-DX1	1	40X	0.2527	Head and Neck Squamous Cell Carcinoma	Train
TCGA-02-0009-01Z-00-DX1	1	20X	0.5015	Brain, Glioblastoma	Train
TCGA-LB-A8F3-01Z-00-DX1	3	40X	0.2527	Pancreas, Pancreatic Adenocarcinoma	Train
TCGA-36-2547-01A-01-TS1	1	20X	0.5015	Ovary, Ovarian Serous Cystadenocarcinoma	Train
TCGA-W5-AA30-01Z-00-DX1	3	40X	0.2529	Bile Duct, Cholangiocarcinoma	Train

TCGA-86-7701-01Z-00-DX1	2	40X	0.252	Lung, Lung Adenocarcinoma	Train
TCGA-B0-4823-01Z-00-DX1	3	40X	0.252	Kidney, Renal Clear Cell Carcinoma	Train
TCGA-02-0009-01Z-00-DX1	1	20X	0.5015	Brain, Glioblastoma multiforme	Train
TCGA-2G-AAKO-05Z-00-DX1	1	40X	0.2277	Testicle, Acute Myeloid Leukemia	Train
PathAI	3	40X	0.2511	Liver, Hepatitis B	Train
PathAI	3	40X	0.2522	Liver, Non-alcoholic steatohepatitis	Train
PathAI	3	40X	0.2522	Kidney, Lupus	Train
PathAI	1	40X	0.2522	Colon, Inflammatory Bowel Disease	Train

PathAI	2	20X	0.5023	Small Intestine, Carcinoid	Train
PathAI	1	40X	0.2522	Prostate, Prostate Adenocarcinoma	Train
TCGA-EJ-5495-01Z-00-DX1	2	40X	0.252	Prostate, Prostate Adenocarcinoma	Validation
TCGA-IG-A3YA-01Z-00-DX1	2	40X	0.2465	Esophagus, Esophageal Carcinoma	Validation
TCGA-GS-A9U3-01Z-00-DX1	2	40X	0.2525	Lymph, Diffuse Large B Cell Lymphoma	Validation
TCGA-LL-A5YM-01Z-00-DX1	2	40X	0.2456	Breast, Invasive Ductal Carcinoma	Validation
TCGA-P4-AAVO-01Z-00-	3	40X	0.2526	Kidney, Renal Papillary Cell Carcinoma	Validation

DX1					
PathAI	2	40X	0.2522	Liver, Non-alcoholic steatohepatitis	Validation
PathAI	1	20X	0.5023	Colon, Human Papillomavirus	Validation
PathAI	2	40X	0.2511	Brain, Glioma	Validation
PathAI	2	40X	0.2522	Skin, Normal	Validation
TCGA-VQ-A8DV-01Z-00-DX1	1	40X	0.2525	Stomach, Stomach Adenocarcinoma	Test
TCGA-CV-6934-01Z-00-DX1	2	40X	0.2525	Head and Neck Squamous Cell Carcinoma	Test
TCGA-MQ-A6BR-01Z-00-DX1	1	40X	0.2465	Lung, Mesothelioma	Test
TCGA-OR-	1	40X	0.2527	Adrenal, Adrenal Cortical Carcinoma	Test

A5J8-01Z-00-DX1					
PathAI	1	40X	0.2522	Colon, Inflammatory Bowel Disease	Test
PathAI	1	20X	0.5023	Breast, Invasive Ductal Carcinoma	Test
PathAI	2	40X	0.2522	Prostate, Prostate Adenocarcinoma	Test

Table 2: Samples used for out-of-distribution (OOD) evaluation of model

performance.

Slide Identifier	# Frames	Magnification	Microns Per Pixel (MPP)	Indication	Dataset
581805	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test 1
581720	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test-1
581810	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test-1
581875	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test-1
591979	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test-1
581916	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test-1
581956	1	40X	0.2518	Colon, Ulcerative Colitis	OOD-Test-1
591275	1	40X	0.2521	Breast, Breast Cancer	OOD-Test-2
591351	1	40X	0.262385	Breast, Breast Cancer	OOD-Test-2
591347	1	40X	0.262385	Breast, Breast Cancer	OOD-Test-2

592506	1	40X	0.2521	Breast, Breast Cancer	OOD-Test-2
592559	1	40X	0.262125	Breast, Breast Cancer	OOD-Test-2
597204	1	40X	0.262125	Breast, Breast Cancer	OOD-Test-2
597271	1	40X	0.262125	Breast, Breast Cancer	OOD-Test-2

Table 3. Characteristics of patients in TCGA cohorts.

	TCGA BRCA N=886	TCGA LUAD N=426	TCGA PRAD N=392
Age at initial pathologic diagnosis (median, range)	58 (26-90)	66 (33-88)	61 (41-77)
Sex (male), n (%)	10 (1.1%)	191 (44.8%)	392 (100%)
American Joint Committee on Cancer (AJCC) Tumor Stage (n)	Stage I: 77 Stage IA: 76 Stage IB: 5 Stage II: 6 Stage IIA: 286 Stage IIB: 207 Stage III: 2 Stage IIIA: 128 Stage IIIB: 18 Stage IIIC: 51 Stage IV: 13 Unknown: 17	Stage I: 5 Stage IA: 122 Stage IB: 108 Stage II: 1 Stage IIA: 47 Stage IIB: 58 Stage IIIA: 50 Stage IIIB: 5 Stage IV: 22 Unknown: 8	Unknown: 392 (100%)
Whole genome doublings (WGD)	0: 435, 56.8% 1: 298, 38.5% 2: 37, 4.8% Unknown: 116	0: 168, 41.9% 1: 193, 48.1% 2: 40, 10.0% Unknown: 25	0: 337, 92.1% 1: 29, 7.9% Unknown: 26

Progression-free survival (PFS, median)	122.3 mo	29.3 mo	116.7 mo
Overall survival (OS, median)	131.4 mo	48.5 mo	Not Reached

Figure 1

A

B

Attribute	Feature
Size	Area Major Axis Length Minor Axis Length Perimeter Length
Shape	Circularity Eccentricity
Texture	Solidity Standard deviation of the grayscale intensity Standard deviation of the saturation value in HSV color space Standard deviation of the A value in LAB color space Standard deviation of the B value in LAB color space
Stain Intensity	Mean greyscale intensity Mean saturation value in HSV color space Mean A value in LAB color space Mean B value in LAB color space

Figure 2

Figure 3

TCGA BRCA **TCGA LUAD** **TCGA PRAD**

Segmentation Color Scheme:

Cancer epithelial cell Lymphocyte Fibroblast Macrophage

Plasma cell Granulocyte Other cell

Figure 4

Figure 5

Figure 6

Figure 7

< 50th Percentile:						
At risk	434	133	48	4	1	0
Censored	1	259	330	365	367	368
Events	0	43	57	66	67	67

≥ 50th Percentile:						
At risk	433	110	23	5	3	3
Censored	1	259	325	335	337	337
Events	0	65	86	94	94	94

< 50th Percentile:						
At risk	434	146	55	7	3	0
Censored	1	263	336	378	382	384
Events	0	26	44	50	50	51

≥ 50th Percentile:						
At risk	433	116	28	9	6	4
Censored	1	276	348	358	361	362
Events	0	42	58	67	67	68