Surface exclusion of IncC conjugative plasmids and their relatives

Nicolas Rivard[1][#a], Malika Humbert[1][¶], Kévin T. Huguet[1][#b][¶], Aurélien Fauconnier[1][¶], César Pérez

Bucio[2], Eve Quirion[1], Vincent Burrus[1][*]

[1]Département de biologie, Université de Sherbrooke, Sherbrooke, Quebec, Canada

[2]Instituto Tecnológico y de Estudios Superiores de Monterrey, Monterrey, Nuevo León, Mexico

[#a]Current address: Laboratoire d'Ingénierie des Systèmes Macromoléculaires, UMR7255,

Institut de Microbiologie de la Méditerranée, Aix-Marseille Univ, CNRS, Marseille, France.

[#b]Current address: CNRS, LCPME, UMR 7564, Université de Lorraine, Villers-lès-Nancy,

France

Short title: Surface exclusion of IncC conjugative plasmids

*Corresponding author

E-mail: Vincent.Burrus@USherbrooke.ca

[¶]These authors contributed equally to this work.

Abstract

The phenomenon of exclusion allows conjugative plasmids to selectively impede the entry of

identical or related elements into their host cell to prevent the resulting instability. Entry

exclusion blocks DNA translocation into the recipient cell, whereas surface exclusion

destabilizes the mating pair. IncC conjugative plasmids largely contribute to the dissemination of

antibiotic-resistance genes in Gammaproteobacteria. IncC plasmids are known to exert

exclusion against their relatives, including IncC and IncA plasmids, yet the entry exclusion factor

eexC alone does not account for the totality of the exclusion phenotype. In this study, a

transposon-directed insertion sequencing approach identified *sfx* as necessary and sufficient for

the remaining exclusion phenotype. Sfx is an exclusion factor unrelated to the ones described to

date. A cell fractionation assay localized Sfx in the outer membrane. Reverse transcription PCR

and beta-galactosidase experiments showed that *sfx* is expressed constitutively at a higher level

than *eexC*. A search in Gammaproteobacteria genomes identified Sfx homologs encoded by

IncC, IncA and related, untyped conjugative plasmids and an uncharacterized family of

integrative and mobilizable elements that likely rely on IncC plasmids for their mobility. Mating

assays demonstrated that *sfx* is not required in the donor for exclusion, ruling out Sfx as the

exclusion target. Instead, complementation assays revealed that the putative adhesin TraN in

the donor mediates the specificity of surface exclusion. Mating assays with TraN homologs from

related untyped plasmids from *Aeromonas* spp. and *Photobacterium damselae* identified two

surface exclusion groups, with each Sfx being specific of TraN homologs from the same group.

Together, these results allow us to better understand the apparent incompatibility between IncA

and IncC plasmids and to propose a mechanistic model for surface exclusion mediated by Sfx in

IncC plasmids and related elements, with implications for the rampant dissemination of antibiotic

resistance.

Author summary

Bacterial conjugation plays a pivotal role in the evolution of bacterial populations. The circulation of drug resistance genes bolsters the emergence of multidrug-resistant pathogens, with which contemporary medicine struggles to cope. Exclusion is a natural process preventing the redundant acquisition of a plasmid via conjugation by a host harbouring an identical or similar plasmid. Although exclusion has been known for the past half-century, the mechanisms involved remain poorly understood. This study describes an exclusion factor, Sfx, encoded by IncC, IncA and related conjugative plasmids and by unrelated integrative and mobilizable elements. We report that Sfx is a lipoprotein of the recipient that selectively inhibits conjugation based on the adhesin TraN expressed at the surface of the donor. We propose a mechanistic model for Sfx-mediated exclusion. Ultimately, a better understanding of exclusion could facilitate the design of conjugation inhibitors targeting mating pair formation to curb the circulation of drug-resistance genes in healthcare settings, agriculture, animal husbandry and food and drug production.

Introduction

IncC plasmids are large conjugative plasmids frequently associated with multidrug resistance phenotypes in a broad range of Gammaproteobacteria species [1]. IncC plasmids' ability to mobilize non-autonomous, unrelated integrative and mobilizable elements (IMEs) of the *Salmonella* Genomic Island 1 (SGI1) and MGI*Vch*Hai6 families exacerbates their importance in antibiotic resistance dissemination [2–5]. SGI1 and its multiple variants are important vehicles of antibiotic resistance genes and are frequently found in *Salmonella enterica* and a broad range of Gammaproteobacteria [6]. MGI*Vch*Hai6 conferred multidrug resistance to *Vibrio cholerae* non-O1/non O139 infecting cholera patients during the 2010 cholera outbreak in Haiti [4]. IncC, IncA and untyped Inc<u>A</u>/<u>C</u>-like (ACL) plasmids (e.g., pAsa4c, pAhD4-1, pAQU1) share a syntenic set of genes involved in conjugation, DNA repair, and regulation. All seem regulated by closely related homologs of the master activator of transfer AcaCD [7–9]. IncC plasmid-encoded AcaCD promotes the excision and mobilization of MGI*Vch*Hai6-like IMEs [4,5]. AcaCD also stimulates the excision and replication of SGI1 and the expression of its *traN*, *traG* and *traH* genes, which modify the mating apparatus of IncC plasmids to enhance its dissemination [7,10,11].

Large conjugative plasmids are autonomous replicons that usually remain at a low-copy number. The entry into the same cell of a plasmid sharing similar replication or partitioning determinants promotes plasmid destabilization and loss, a phenomenon known as incompatibility [12]. Exclusion refers to the mechanisms employed by conjugative elements to preclude instability resulting from incompatibility by hindering redundant transfer into the cells they reside [13]. While the definite nature of exclusion mechanisms remains elusive, they can be categorized as entry and surface exclusion by whether they interfere with DNA transfer or mating pair stabilization [13,14]. Exclusion seems to be a staple of the conjugative plasmid lifestyle. Exclusion factors have been identified in the IncF, IncHI1, IncI, IncN, IncP, IncW and ColE1

families, plasmids of the pKPC_UVA01 replicon type and the enterococcal conjugative plasmid

pCF10 [14–22]. Integrative and conjugative elements (ICEs) of the SXT/R391 family and

ICE*Bs*1 of *Bacillus subtilis* also encode exclusion determinants [23,24]. Most exclusion factors

identified to date are associated with the cytoplasmic membrane, where they would exert entry

exclusion, with only the IncF TraT firmly established as a surface exclusion factor [13,14]. Like

TraT, the IncHI1 EexB localizes in the outer membrane and is thought to engage in surface

exclusion [15]. Likewise, the surface-exposed PrgA protease of pCF10 reduces aggregation and

transfer [22].

Exclusion is not specific to the transferred DNA [25], and proteins in the donor have been

identified as targets of exclusion factors in several systems. EexA and EexB of IncHI1 plasmids

and TrbK of pKPK_UVA01 appear to act concurrently as exclusion factors and exclusion targets,

as they are necessary in both the donor and recipient for exclusion to take place [15,21]. Other

exclusion factors target different plasmid-encoded proteins. The IncI ExcA targets TraY, which

is found in the cytoplasmic membrane of the donor [16]. Similarly, the IncF TraS and SXT/R391

Eex of the recipient seem to interact directly with the VirB6 homolog TraG of the donor, and the

regions involved in both proteins are cytoplasmic [23,26]. In the cases above, the genes

encoding the exclusion factor and its target are adjacent. In IncP, IncW and IncN plasmids, for

which the exclusion target remains unknown, the exclusion gene abuts a gene coding for a

VirB6 homolog, suggesting a widespread usage of this target beyond the IncF and SXT/R391

families [13].

The surface exclusion factor TraT of IncF plasmids reduces the percentage of mating

aggregates, and five contiguous residues determine specificity between exclusion groups

[14,27]. Harrison *et al*. proposed that TraT destabilizes the mating pair by interacting with the

pilus tip; however, it was eventually shown that the pilin does not define exclusion specificity

[27,28]. The discovery of interactions between TraT and the outer membrane protein OmpA in

the recipient and between OmpA in the recipient and the putative adhesin TraN in the donor

suggested surface exclusion could instead result from TraT preventing docking of TraN to

OmpA, thus hindering mating pair stabilization [29–31]. The identification of TraN receptor

specificity groups across IncF plasmids makes this hypothesis extremely appealing, yet *traN*

substitutions failed to flip exclusion specificity between F and R100-1 [30,32,33].

IncC plasmids share with IncA plasmids the same entry exclusion system, *eexC*/*traG*, and

inhibit each other's entry into the cell in which they reside [9,34]. During the characterization of

eexC/*traG*, we reported that knocking-out *eexC* of the IncC plasmid pVCR94 in the recipient

only reduced the exclusion phenotype in mating assays using an IncC$^+$ donor. Likewise, the

deletion of a 45.6-kb fragment encompassing 33 open reading frames located between *traN* and

traF led to a statistically significant reduction of exclusion. Hence, another mechanism likely

prevents redundant transfer between cells containing identical plasmids.

In this report, we investigate the cause of the incomplete abolition of exclusion by a recipient

strain bearing an IncC plasmid lacking the entry exclusion gene *eexC*. Using a transposon-

directed insertion sequencing (TraDIS) approach, we identified IncC-borne candidate genes

whose disruption in recipient cells facilitated the acquisition of an incoming mobilizable plasmid

from an IncC$^+$ donor. Further analyses revealed that a single open reading frame, hereafter

named *sfx*, flanked by *traN* and *acaB*, accounts for the residual exclusion phenotype. Deletion

of *sfx* with *eexC* completely abolished exclusion. Signal peptide prediction and a subcellular

localization assay suggest the Sfx protein is an outer-membrane lipoprotein and support a role

as a surface exclusion factor. Sfx belongs to an uncharacterized family of broadly distributed

surface exclusion proteins encoded by IncC plasmids, their relatives and several putative IMEs.

Surface exclusion assays designed to test *sfx*/*traN* gene pairs from ACL plasmids revealed that

sfx and *traN* are part of a surface exclusion system, enabling us to propose a model of surface

exclusion of ACL plasmids.

Results

vcrx085 impedes conjugative transfer between IncC$^+$ cells

To identify additional factors responsible for the inhibition of transfer between IncC$^+$ cells, we used transposon-directed insertion sequencing (TraDIS) aimed at mapping Tn*5* insertions that enhance the entry of the mobilizable plasmid pClo transferred from an IncC$^+$ donor strain into a Tn*5*$^+$ IncC$^+$ recipient library (Fig 1A). pClo was used as a proxy for conjugative transfer to bypass IncC plasmid incompatibility. Briefly, we constructed a high-density mini-Tn*5* (Sp^R) insertion library in *E. coli* GG56 carrying pVCR94^{Kn} Δ*eexC*::*cat*, a kanamycin-resistant variant of pVCR94 unable to exert entry exclusion in recipient cells [9]. This set of mutants formed the input library containing Tn*5* insertions in the chromosome or pVCR94 that allowed plasmid replication and maintenance. The input library was then used as recipient in a mating assay with a donor strain carrying pVCR94^{Kn} and the mobilizable plasmid pClo. GG56 transconjugant colonies that acquired pClo made up the output library. We expected that genes or sequences of pVCR94^{Kn} Δ*eexC*::*cat* more frequently disrupted in the output than in the input (high insertion index ratio) were those impeding the mobilization of pClo between IncC$^+$ cells. Since our assays monitored pClo entry only, we did not consider the role of chromosomal genes in the recipient.

Fig 1. Identification of an additional IncC exclusion factor. (A) Overview of the TraDIS workflow to identify exclusion factors in pVCR94^{Kn} Δ*eexC*::*cat*. (B) Mini-Tn5 reads near *vcrx085*. The tracks plot the number of reads from two independent replicates as a function of position in pVCR94^{Kn} Δ*eexC*::*cat* for both the input and output libraries. ORFs with similar functions are colour-coded as indicated in the panel. This panel was created using the UCSC Genome Browser (http://genome.ucsc.edu). (C) Nine loci of unknown function show a mini-Tn*5* insertion index ratio greater than 5 (dashed line), calculated as the ratio of insertion counts between the output and input libraries. ORFs with similar functions are colour-coded as indicated in Fig S1.

(D) *sfx* (*vcrx085*) is the only functional exclusion factor among candidates identified in (B) and

suffices alongside *eexC* (*eex*) to account for the totality of IncC exclusion. *E. coli* VB112 (Rf^r)

containing pVCR94^{Kn} and pClo (Ap^r) served as the donor strain, and *E. coli* GG56 (Nx^r)

containing the specified derivatives of pVCR94^{Sp} served as the recipient strains. When indicated,

eexC or *sfx* were expressed in the recipients from single-copy, chromosomally integrated

pAH56 (Kn^r) in the presence of IPTG or together from pBeloBAC11 (Cm^r) under the control of

their respective native promoters. Transconjugants containing pClo were selected as the Nx^r Ap^r

colonies. Crossbars show the mean and standard error of the mean of three independent

experiments. One-way ANOVA (p=1.6e-25) with a Tukey-Kramer post-test was used on the

log$_{10}$-transformed values to compare the means. Statistical significance (S2 File) is shown as a

compact letter display for pairwise comparisons where means grouped under identical letters

are not statistically different. Exclusion indices, shown at the bottom of each crossbar, are

calculated as the frequency of transfer into the empty recipient divided by the frequency of

transfer into the indicated mutant.

We identified nine candidate genes in pVCR94^{Kn} Δ*eexC*::*cat* with insertion index ratios above

the arbitrary threshold of 5 (Table 1, Fig 1B and 1C, S1 Fig). Among these genes, only *acr2*

(*vcrx150*) is a known repressor of conjugative transfer [7]. To validate the ability of the

candidates to inhibit incoming conjugative transfer, we tested mobilization of pClo into the

corresponding deletion mutants of pVCR94^{Sp} used as recipients. All but one had no impact on

transfer of pClo (Fig 1D). The deletion of *vcrx085* (hereafter referred to as *sfx* for surface

exclusion) improved conjugation to the same level similar as the deletion of the entry exclusion

factor *eexC*. Like *eexC*, we could complement the *sfx* deletion in *trans* (Fig 1D, pAH56). Both

mutations combined completely abolished exclusion, allowing conjugation to a level comparable

to an empty recipient. Conversely, the standalone expression of *sfx* and *eexC* from their native

promoter in the absence of an IncC plasmid in the recipient matched the full exclusion

phenotype of the wildtype IncC plasmid (Fig 1D, pBeloBAC11).

Table 1. Candidate genes identified by TraDIS.

ORF	Fold change	Predicted size (aa)	Predicted product[a]
vcrx018	6.39	119	Hypothetical protein
vcrx019	5.17	148	Fe^{3+}-siderophore ABC transporter permease
vcrx062	8.67	194	Hypothetical protein
vcrx085 (sfx)	6.22	286	Lipoprotein (Sec/SPII)
vcrx107	6.07	114	Hypothetical protein
vcrx108	6.08	100	Hypothetical protein
vcrx111	6.83	184	Hypothetical protein
vcrx137	6.63	163	Hypothetical protein
acr2	6.50	139	Transcriptional repressor

[a]As predicted by NCBI blastp or SignalP 6.0.

Deletion of *eexC*, *sfx*, or both did not significantly impact the fitness of the cells (similar CFU

counts/ml) when compared to wild-type pVCR94^{Sp} or empty cells (S2 Fig). Conversely, the

overexpression of *eexC* or *sfx* from pBAD30 slightly decreased the fitness.

Sfx is a constitutively expressed outer membrane protein

The SignalP 6.0 server predicts a lipoprotein signal peptide in the translation product of *sfx* (S3

Fig), suggesting Sfx is displayed on the cell surface [35]. A C-terminally 3xFLAG-tagged version

of *sfx* was cloned under the control of the inducible promoter P_{tac}. The Sfx^{3xFLAG} polypeptide has

a predicted size of 33.4 kDa before signal peptide cleavage and 31.7 kDa afterwards. A cell

fractionation assay confirmed the presence of Sfx^{3xFLAG} in the inner and outer membrane

fractions (Fig 2A). In contrast, the known outer membrane protein OmpC, used as a control,

was only detected in the outer membrane fraction. This cellular localization remains consistent

with the role of Sfx in surface exclusion. The accumulation of Sfx^{3xFLAG} in the inner membrane

suggests that the high expression level of *sfx*^{3xFLAG} from P_{tac} on a multi-copy vector saturates the

Lpt trafficking pathway. We could not undoubtedly assign the processed and unprocessed forms

in the inner membrane fraction, and two additional products appeared between 20 and 25 kDa

on the western blot. These weaker bands are presumably degradation products of an unknown

nature.

Fig 2. Expression of IncC surface exclusion is independent of the master activator of

transfer AcaCD. (A) Cell fractionation assay. The soluble (S), inner membrane (IM), and outer

membrane (OM) cell fractions of *E. coli* DH5α expressing 3xFLAG-tagged Sfx$_{94}$ or 3xFLAG-

tagged AcaC from pAH56 under the control of the IPTG-inducible P_{tac} promoter with (+) and

without (-) IPTG were analyzed by SDS-PAGE and immunodetected using rabbit anti-FLAG or

anti-OmpC antibodies. Molecular weight markers (M) are indicated on the left. A vertical white

line separates non-adjacent lanes from a single blot. AcaC-3xFLAG (22 kDa), a known

cytoplasmic protein [7] and OmpC (36 kDa), a known outer membrane protein [36], were used

as controls in the fractionation. (B) Schematic representation of the *acaB-sfx* locus. The relative

positions of reverse transcription primer 85RT and primers used to amplify *sfx* and *acaB* are

shown. A dotted line shows the reverse transcription product. Stem loops represent predicted

Rho-independent terminators. The translational *lacZ* fusion used for enzymatic assays was

introduced after the third codon of *sfx*. (C) *sfx* is expressed constitutively and independently of

the upstream AcaCD-dependent P_{acaB} promoter. A 2% agarose gel of the PCR products from an

sfx-derived reverse transcribed RNA. Untreated RNA (RNA) and genomic DNA from *E. coli*

GG56 (Nx^r) containing pVCR94^{Kn} Δ*acaCD* (gDNA) were used as negative and positive controls,

respectively. (D) β-galactosidase activity of GG56 (Nx^r) containing pVCR94^{Kn} *eexC'-'lacZ*,

pVCR94^{Kn} *sfx'-'lacZ*, or pVCR94^{Kn} *traN'-'lacZ* translational fusions with or without p*acaCD* and

arabinose. Crossbars show the mean and standard error of the mean of three independent

experiments. One-way ANOVA with a Tukey-Kramer post-test was used to compare the means.

sfx is located downstream of the transfer activator *acaB*, which is expressed from an AcaCD-

inducible promoter (Fig 2B) [37].. Hence, we wondered whether *sfx* expression was controlled

by AcaCD. To test this hypothesis, we constructed a translational fusion between the third

codon of *sfx* and the eighth codon of *lacZ* in pVCR94^{Kn}. Expression of AcaCD did not impact the

β-galactosidase activity of the *sfx'- 'lacZ* fusion. A similarly constructed *eexC'- 'lacZ* fusion yielded

a ~150-fold lower β-galactosidase activity, regardless of the presence of AcaCD (Fig 2D). We

used a similar *traN'-'lacZ* fusion as a control and observed a very low level of expression in

IncC$^+$ cells, which increased ~260 fold upon AcaCD overexpression. Furthermore, to test

whether *sfx* and *acaB* are part of the same transcript, we attempted to amplify both loci from a

cDNA synthesized from the 3' end of *sfx*. While *sfx* could be amplified regardless of AcaCD,

acaB remained undetectable, suggesting that these two genes are expressed as independent

mRNA transcripts (Fig 2C). A Rho-independent terminator predicted between *sfx* and *acaB*

supports this idea (Fig 2B, S3B Fig). Altogether, our results indicate that *sfx* is constitutively

expressed independently of AcaCD at a much higher level than *eexC*.

Genes encoding homologs of Sfx$_{94}$ are found in diverse genomic backgrounds

To explore the diversity of exclusion factors, we first used NCBI tblastn to identify Sfx$_{94}$

homologs in the ACL plasmids pAsa4c of *Aeromonas salmonicida*, pAhD4-1 of *Aeromonas*

hydrophila, and pAQU1 of *Photobacterium damselae* that are known untyped relatives of IncA

and IncC plasmids [8,9]. For pAsa4c and pAhD4-1, closely related homologs were found

adjacent to the gene *traN*. For pAQU1, the search resulted in a single hit (*215*, hereafter

referred to as *sfx*$_{AQU1}$) whose product shares only 39% identity over 273 amino acid residues

with Sfx$_{94}$. *sfx*$_{AQU1}$ is surrounded by genes of unknown function. Sfx$_{AQU1}$ was manually added to

the dataset for downstream analyses. We next searched the GenBank database for homologs

of Sfx_{94} and extracted the resulting sequences for further analyses. 130 unique proteins were

identified (S3 File), most encoded by ACL plasmids (Fig 3A). 78 Sfx homologs share more than

90% identity with Sfx_{94}, including nine variants encoded by IncC plasmids. All of the Sfx

homologs encoded by IncC plasmids are inside this cluster. Sfx_{94} belongs to an identical protein

group of 193 representatives, all encoded by IncC or putative IncC plasmids. Sfx_{RA1} from the

IncA plasmid pRA1 belongs to the Sfx_{94} cluster as they share 92% identity over 286 amino acid

residues. All plasmids outside the Sfx_{94} cluster are ACL plasmids, including pAsa4c and pAhD4-

1. In most ACL plasmids and with the rare exception of pAQU1, *sfx* lies downstream of the

dsbC-traC-trhF-traWU operon between *traN* and the transfer activator gene *acaB* (Fig 3C).

Fig 3. Sfx homologs are found across a wide range of Gammaproteobacteria. Maximum

likelihood phylogenetic analyses of Sfx (A) and TraN (B) homologs. Trees with the highest

likelihood (9,621.78 and 28,204.23 for Sfx and TraN, respectively) are shown. Bootstrap

supports are indicated as percentages at the branching points only when >80%. Branch lengths

represent the number of substitutions per site over 269 and 813 amino acid positions for Sfx

and TraN, respectively. Only one representative per cluster of similar proteins (>90%) is shown

in each tree, except for $TraN_{Asa4c}$ and $TraN_{AhD4-1}$, which share 95% identity over the aligned

sequences. S3 File provides additional details on host strains. (C) Schematic representation of

the genetic context of *sfx* homologs in a selection of putative IMEs and ACL plasmids. Predicted

attL and *attR* attachment sites flanking the IMEs are represented by blue bars. pAQU1 encodes

homologs of *acaB*, *vcrx087-088-089-090*, *bet* and *exo* clustered in a distinct locus not shown

here. ORFs with similar functions are colour-coded as indicated in the panel.

The remaining Sfx proteins are encoded by putative IMEs integrated at four distinct

chromosomal locations. First, ten of these IMEs belong to the MGI*Vch*Hai6 family and are

integrated at the 5' end of *dusA* (tRNA-dihydrouridine synthase A) in environmental and clinical

Aeromonas strains [5]. Represented by IE*Ave*Tha1 (Fig 3C), all possess MGI*Vch*Hai6-like *mobI*

and *oriT* and predicted AcaCD-responsive promoters (S2 Table). A second group represented

by IE*Vch*Aut1 comprises six IMEs integrated at the 3' end of *yicC* (putative RNase adaptor

protein YicC) in environmental and clinical *Vibrio* strains. In addition to a putative recombination

directionality factor related to RdfM of MGI*Vfl*Ind1 [38], these IMEs carry a set of conserved

genes, including a predicted type IV coupling protein (*traD*) and a putative relaxase of the MOB_F

family (*traI*). Finally, two IMEs coding for an identical Sfx are integrated at the 3' end of *tRNA-*

TRP and the 5' end of *rlmD* (23S rRNA uracil methyltransferase) in environmental isolates of

Shewanella chilikensis and *Shewanella putrefaciens*, respectively. Despite sharing a conserved

core set of genes with IE*Vch*Aut1, the Sfx protein of these two IMEs resembles and clusters

with Sfx_{94} (Fig 3A). Although we can predict AcaCD binding sites across these IMEs (S2 Table),

no other conserved features of the SGI1 and MGI*Vch*Hai6 families was found, suggesting they

belong to an unrelated IME family mobilizable by IncC plasmids.

Surface exclusion specificity is determined by TraN

To test the role of Sfx in the donor, we monitored the transfer of pClo from donors carrying the

Δ*eexC* and Δ*sfx* deletion mutants of pVCR94^{Sp} toward an IncC$^+$ strain (pVCR94^{Kn}). No

alleviation of exclusion, i.e., an increase in transfer frequency, was observed (Fig 4A). On the

contrary, a modest yet significant decrease was observed for the Δ*sfx* and Δ*eexC* Δ*sfx* mutants,

suggesting that augmented redundant transfer between donors was detrimental to transfer

toward recipients. Hence, *sfx* is not self-targeting to promote surface exclusion.

Next, we focused on TraN as a potential target for surface exclusion, as the gene *traN* encodes

a putative adhesin thought to stabilize the mating pair, and it is adjacent to *sfx* in most

conjugative plasmids found but pAQU1 [30,31,39]. A search for homologs of $TraN_{94}$ yielded 412

homologous proteins (S3 File), most encoded by ACL plasmids. Others belonged to IncC-

mobilizable IMEs of the SGI1 family integrated into *trmE* or *dusA*, with no clear relationship

between TraN phylogeny and insertion site (Fig 3B). The cluster represented by pAQU1 differs

significantly from the close ACL plasmids as the *dsbC-traC-trhF-traWUN* and *acaB*/*bet-exo*

regions are not adjacent, and *sfx* lies elsewhere.

Fig 4. TraN directs surface exclusion specificity. (A) The absence of *eexC* and *sfx* in the

donor decreases the frequency of transconjugant formation. pClo (Ap^r) was mobilized from *E.*

coli VB112 (Rf^r) bearing the indicated deletion mutants of pVCR94^{Sp} (donor, D) into *E. coli*

GG56 (Nx^r) bearing pVCR94^{Kn} (recipient). Transconjugants containing pClo were selected as

the Nx^r Ap^r colonies. Crossbars show the mean and standard error of the mean of three

independent experiments. One-way ANOVA (p=0.016) with a Tukey-Kramer post-test was used

on the log$_{10}$-transformed values to compare the means. Statistical significance (S2 File) is

shown as a compact letter display for pairwise comparisons where means grouped under

identical letters are not statistically different. (B) *sfx*$_{94}$ is ineffective against the *traN*$_{94}$ distant

homolog *traN*$_{AQU1}$. Pairwise t-tests were used on the log$_{10}$-transformed values to compare the

means. Exclusion indices are calculated by dividing the frequency of transfer of pVCR94^{Sp}

Δ*traN* to the empty recipient divided by the transfer frequency to the recipient expressing *sfx*$_{94}$.

(C) Heatmap of surface exclusion indices of *sfx* and *traN* pairs from a diverse set of ACL

plasmids. Exclusion indices are calculated by dividing the frequency of transfer of pVCR94^{Sp}

Δ*traN* to the empty recipient divided by the transfer frequency to the recipient expressing the

indicated *sfx* variant. In (B) and (C), *E. coli* GG56 (Nx^r) containing pVCR94^{Sp} Δ*traN* and

expressing the indicated *traN* variant from pBAD30 (Ap^r) (donor, D) was mated with *E. coli*

CAG18439 (Tc^r) expressing the indicated *sfx* variant from a single-copy, chromosomally

integrated pAH56 (Kn^r) (recipient, R). Transconjugants containing pVCR94^{Sp} Δ*traN* were

selected as the Tc^r Sp^r colonies.

To investigate the involvement of TraN in Sfx-dependent exclusion, we used in mating assays a

traN deletion mutant of pVCR94^{Sp} complemented ectopically with either its native *traN$_{94}$* or the

distantly related homolog *traN$_{AQU1}$* [40]. Despite sharing only 66% identity with TraN$_{94}$, TraN$_{AQU1}$

fully complemented the Δ*traN$_{94}$* mutation (Fig 4B). Furthermore, whereas the expression of *sfx$_{94}$*

in the recipient strongly inhibited conjugation supported by *traN$_{94}$*, it did not impair transfer

supported by *traN$_{AQU1}$* (Fig 4B). This result implies that TraN is, if not a direct interactant of Sfx,

at least a key player in Sfx-mediated exclusion and the determinant of exclusion specificity. To

further investigate the diversity of exclusion specificity, we tested the *traN*/*sfx* pairs of pAsa4c

and pAhD4-1, two plasmids belonging to distinct entry exclusion groups, and pAQU1 [9,41,42].

The corresponding genes were tested in pairs in all possible combinations to determine

exclusion indices. Two separate exclusion groups emerged (Fig 4C and S5 Fig). *sfx* homologs

from pVCR94, pAsa4c and pAhD4-1 specifically exerted exclusion against *traN* homologs from

these same plasmids. *sfx$_{AQU1}$* only excluded *traN$_{AQU1}$*. In addition, we found that *sfx$_{Aut1}$* from the

putative IME IE*Vch*Aut1 is functional. *sfx$_{Aut1}$* exhibits a surface exclusion pattern resembling

sfx$_{Asa4c}$, excluding transfer mediated by *traN* of pVCR94, pAsa4c and pAhD4-1 but not pAQU1.

Discussion

All known conjugative plasmids encode at least one exclusion factor [13]. An entry exclusion

system involving the entry exclusion factor EexC and the VirB6-homolog TraG was previously

identified in IncA and IncC conjugative plasmids [9]. Here, we identified a second exclusion

factor within a region suspected by Humbert *et al*. to be responsible for additional exclusion

activity [9]. A TraDIS experiment pointed to nine candidate genes likely to enhance incoming

transfer once disrupted. *sfx* was the only one whose activity in exclusion could be

experimentally validated. Together, *eexC* and *sfx* account for the totality of the exclusion

phenotype (Fig 1D), akin to the exclusion mediated by *traS* and *traT* of the F plasmid [43]. We

could not detect any impact of abolishing exclusion on cell viability (S2 Fig). This result strongly

contrasts with reports on F plasmid exclusion for which double mutants of *traS* and *traT* could

not be isolated [43]. This discrepancy could be due to the strong repression of the conjugative

machinery of IncC plasmids (Fig 2D), whereas F transfer is constitutive due to the inactivation of

F *finO* by IS*3* [7,44].

The translation product of *sfx* is a predicted lipoprotein, and we detected Sfx in the outer

membrane (Fig 2A), a trait shared with the surface exclusion factor TraT of the F plasmid. In

addition, we showed that exclusion mediated by *sfx* is inhibited or abolished by the substitution

of the putative adhesin TraN (Fig 4C), suggesting that exclusion specificity is determined by

sfx/*traN* pairs. Together, these observations support a mechanism of surface exclusion [14,45].

Our search in the Genbank database using Sfx_{94} failed to reveal any known entry or surface

exclusion factor, including F plasmid TraS or TraT, establishing Sfx of IncA and IncC plasmids

as an unrelated surface exclusion factor (S3 File). As the ACL plasmids pAsa4c and pAhD4-1

belong to the same surface exclusion group as pVCR94 despite their divergence (Fig 4C), IncA

and IncC plasmids are expected to exert surface exclusion against each other, hence

contributing alongside entry exclusion to the previously observed exclusion phenotype between

the two incompatibility groups [9,34]. Among the Sfx homologs tested in this study, only Sfx_{AQU1},

which failed detection by the NCBI blastp search due to low similarity with Sfx_{94}, belongs to a

separate surface exclusion group (Fig 4C). A pairwise comparison illustrates that, like Sfx_{AQU1},

the putative surface exclusion proteins WP_035235290.1 and WP_121638668.1 share minimal

homology with all other representatives (Fig 3A and S4 Fig). Therefore, we speculate that

neither belongs to the two surface exclusion groups described here and that all other IncC, IncA

and close ACL plasmids belong to the Sfx_{94} surface exclusion group. This hypothesis is further

supported by the exclusion pattern of sfx_{Aut1} (IE*Vch*Aut1), virtually identical to sfx_{Asa4c}. From our

observations with ACL plasmids, surface exclusion strikingly contrasts with entry exclusion,

which seems much more specific. Indeed, pAsa4c and pAhD4-1 each belong to separate entry

exclusion groups (EexE and EexD, respectively) distinct from the one encompassing IncC and

IncA plasmids (EexC) [9]; yet, all belong to the same surface exclusion group. These

observations suggest IncC, IncA and ACL plasmids form a two-tier community, exerting surface

and entry exclusion against closely related peers but only surface exclusion against a broader

range of distantly related conjugative elements.

This work has identified orphan surface exclusion factors borne by IMEs. IE*Ave*Tha1-like

elements belong to the MGI*Vch*Hai6 family of IncC-mobilizable IMEs [4,5], whereas IE*Vch*Aut1-

like elements constitute a new family of IMEs likely to be mobilizable by ACL plasmids (Fig 3C).

We showed that sfx_{Aut1} of IE*Vch*Aut1 is functional (Fig 4D). Since these IMEs lack a *traN* gene,

we propose they engage in surface exclusion as a defensive mechanism, forcing their host to

favour mating partners that could enhance both their stability and transmissibility. Hence, these

IMEs could modulate the circulation of conjugative elements in bacterial populations without

affecting whether surface exclusion from a potential recipient will target transfer mediated by

their helper element. Such behaviour would contrast with the way SGI1 manipulates the

exclusion system of its helper IncC plasmid. SGI1, which is not known to exert exclusion,

evades IncC entry exclusion by substituting TraG in the mating apparatus with a distant VirB6

homolog [11]. Manipulation of exclusion by mobilizable elements is not well known. While

homologous ColE1 mobilizable plasmids were reported to inhibit each other's transfer mildly,

this phenotype was attributable to the accessory mob gene *mbeD* with no identified target in the

donor [20,46,47]. It thus remains unclear whether exclusion is at play in that case. The putative

IMEs we reported here may be the first instance of exclusion exerted by mobilizable elements.

The close relationship between Sfx proteins encoded by IE*Ave*Tha1 and IE*Vch*Aut1 with those

tested in this study (Sfx$_{94}$, Sfx$_{AhD4-1}$, and Sfx$_{Asa4c}$) suggests they belong to the same exclusion

group (Fig 3, S4 Fig). This activity has been corroborated by experimental evidence for Sfx$_{Aut1}$

(Fig 4C, S5 Fig), establishing surface exclusion as an actual barrier to the circulation of mobile

genetic elements on par with other bacterial defense mechanisms instead of being limited to

preventing redundant transfer and, potentially, mediating mating pair separation [13,39].

Surprisingly, none of the IMEs identified by the TraN search, including SGI1, appears in the Sfx

tree, suggesting they do not encode a surface exclusion factor related to Sfx$_{94}$. Conversely,

IMEs identified by the Sfx search are ostensibly absent from the TraN tree as they lack the

components of the conjugative apparatus. This observation suggests different survival

strategies. Whereas SGI1 maximizes its own dissemination by remodelling the mating

apparatus to improve transfer and bypass entry exclusion [11], IE*Ave*Tha1 and IE*Vch*Aut1 may

ensure persistence by hindering via surface exclusion the entry of destabilizing, excision-

inducing ACL plasmids. Although the mobilization mechanism of IE*Vch*Aut1 remains unknown,

its gene content and the presence of AcaCD binding motifs suggest mechanistic parallels with

that of SGI1, albeit using totally unrelated mobilization factors. In ACL^{+} cells, AcaCD likely

promotes IE*Vch*Aut1's excision through the expression of *rdfM* [38]. A predicted homolog of the

replication protein P from bacteriophage lambda suggests the excised IME is replicative [48].

IME DNA transfer seems to rely on the IME-encoded TraI, a predicted relaxase of the MOB$_{F}$

family [49]. While SGI1 encodes its own replication and DNA processing functions [10,50,51],

IE*Vch*Aut1 additionally encodes a type IV coupling protein, presumably making it independent

from the IncC-encoded TraD and DtrJ (also known as TraJ) [5,52].

IncHI1 EexB and IncF TraT are two other known surface exclusion factors. While EexB seems

to have a dual role as an exclusion factor and exclusion target, TraT's mechanism remains

elusive as exclusion specificity could not be attributed to direct interaction with pilin or indirect

interaction with the putative adhesin TraN through OmpA [15,27,28,30]. Our results

demonstrate that Sfx is only required in the recipient to enable exclusion (Fig 4A), ruling out the

self-targeting of Sfx. Instead, we found that Sfx exclusion can be neutralized by substituting

TraN in the donor with TraN from another surface exclusion group (Fig 4B and 4C). Distinct

exclusion groups within a given family were previously reported in IncI and IncF plasmids and

SXT/R391 ICEs [16,23,53–55].

Although a direct interaction between Eex and TraG has been suggested in SXT/R391 ICEs

and likely extends to other VirB6-targeting entry exclusion factors [26], direct contact between

Sfx and TraN is unlikely for surface exclusion. In fact, the binding of outer membrane proteins

on donor and recipient cells is hard to reconcile with the idea that surface exclusion physically

prevents the formation or disrupts mating aggregates [14]. Instead, surface exclusion factors

have been proposed to act by concealing outer membrane receptors involved in mating

aggregation [29]. According to Riede and Eschbach, TraT of F directly interacts with the major

outer membrane protein OmpA as TraT inhibits the OmpA-specific phage K3 [29]. Yet, despite

ample evidence that $TraN_F$ interacts with OmpA during conjugation, it does not seem to

determine exclusion specificity as $TraN_F$ substitution with $TraN_{R100-1}$ does not counter exclusion

by TraT of F [30–32].

We found that sfx_{94} is constitutively expressed independently of AcaCD at a much higher level

than the entry exclusion factor $eexC$ or the adhesin gene $traN$ when donor cells are in pure

culture (Fig 2C and 2D). The high expression of sfx relative to $eexC$, despite similar levels of

exclusion exerted by the two genes (Fig 1D, Fig 2D), is consistent with an interaction with a

highly abundant surface protein. While few EexC proteins might suffice to target TraG in one

mating pore during conjugation, an abundance of Sfx proteins is likely required to conceal most,

if not all, TraN receptors on the recipient surface at any time. In pure cultures, expression of

traN remained low, suggesting that the level of Sfx could be sufficient to quench spurious donor-

donor mating (Fig 2D). Unlike TraT of F, we have identified TraN as the specificity determinant

for Sfx-mediated exclusion between ACL plasmids. Thus, we propose that $TraN_{94}$ and $TraN_{AQU1}$

recognize distinct receptors on the recipient surface and that Sfx prevents mating pair

stabilization by specific binding to the receptor of its cognate TraN. Further investigation will be

required to identify the receptors involved.

Materials and methods

Bacterial strains and media

The bacterial strains and plasmids used in this study are described in Table 2. Bacterial strains were grown at 37°C on Luria-Bertani (LB) agar or with agitation in LB broth and maintained at −75°C in LB broth containing 20% (vol/vol) glycerol. Antibiotics were used at the following concentrations: ampicillin (Ap), 100 µg/ml; kanamycin (Kn), 50 µg/ml; spectinomycin (Sp), 50 µg/ml; rifampicin (Rf), 50 µg/ml; chloramphenicol (Cm), 20 µg/ml and nalidixic acid (Nx), 40 µg/ml. Media were supplemented with 0.02% L-arabinose or 0.1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) to induce gene expression from pBAD30 and pAH56 constructs, respectively.

Table 2. Strains and plasmids used in this study

Strain, plasmid or element	Relevant genotype or phenotype	Reference
Escherichia coli		
DH5α	F$^-$ φ80*lacZ*ΔM15 Δ(*lacZYA-argF*)U169 *recA1 endA1 hsdR17*(r$_K^-$, m$_K^+$) *phoA supE44* λ$^-$ *thi*-1 *gyrA96 relA1*	[56]
GG56	Nx^r derivative of BW25113 (Nx)	[57]
KH95	Rf^r derivative of BW25113 (Rf)	[10]
MFD*pir*	MG1655 RP4-2-Tc::[ΔMu1::*aac(3)IV*-Δ*aphA*-Δ*nic35*-ΔMu2::*zeo*] Δ*dapA*::(*erm-pir*) Δ*recA*	[58]
VB112	Rf^r derivative of MG1655 (Rf)	[59]
CAG18439	MG1655 *lacZU118 lacI42*::Tn*10* (Tc)	[60]
Vibrio cholerae		
A12JL5W90	non-O1/non-O139 isolated from a large Austrian lake, 2012	[61]
Plasmids		
pVCR94	IncC conjugative plasmid, *V. cholerae* O1 El Tor (Su Tm Cm Ap Tc Sm)	[62]
pVCR94^{Sp}	Sp^r derivative of pVCR94 (pVCR94ΔX2) (Su Sp)	[11]
pVCR94^{Kn}	Kn^r derivative of pVCR94 (pVCR94ΔX3) (Su Kn)	[11]
pVCR94^{Kn} Δ*eexC*::*cat*	*eex* deletion mutant of pVCR94^{Kn} (Su Kn Cm)	This study
pFG036	*ori$_{pMB1}$*, *cI857* (Ts) repressor, *tetM* (Tc)	Addgene #137996
pFG051	*ori$_{R6K}$*, Tn*5 tnp* under λP$_L$ promoter, *oriT$_{RP4}$*, Tn*5d-aadA7* (Sp)	Addgene #137997
pVCR94^{Sp} Δ*018-019*	*vcrx018-vcrx019* deletion mutant of pVCR94^{Sp} (Su Sp)	This study

pVCR94^{Sp} Δ062	$vcrx062$ deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ107-108	$vcrx107$-$vcrx108$ deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ111	$vcrx111$ deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ137	$vcrx137$ deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ$acr2$	$vcrx150$ deletion mutant of pVCR94^{Sp} (Su Sp)	[7]
pVCR94^{Sp} Δ$eexC$	$vcrx145$ deletion mutant of pVCR94^{Sp} (Su Sp)	[9]
pVCR94^{Sp} Δsfx	$vcrx085$ deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ$eexC$Δsfx	$vcrx145$-$vcrx085$ double deletion mutant of pVCR94^{Sp} (Su Sp Cm)	This study
pVCR94^{Kn} Δ$acaCD$	$acaCD$ deletion mutant of pVCR94^{Kn} (Su Kn)	This study
pClo	CloDF13::TnAΔEcoRV (pSU4628) (Ap)	[63]
p$acaCD$	pBAD30::$acaCD$ (Ap)	[7]
pVCR94^{Sp} Δ$traN$	$traN$ deletion mutant of pVCR94^{Sp} (Su Sp)	[11]
pVCR94^{Kn} Δ(087-$traG$)	$vcrx87$-$traG$ deletion mutant of pVCR94^{Kn} (Su Kn)	This study
pAsa4c	Cm^r conjugative plasmid from *A. salmonicida* JF2267 (Cm)	[41]
p$traN_{94}$	pBAD30::$traN_{pVCR94}$ (Ap)	[11]
p$traN_{Asa4c}$	pBAD30::$traN_{pAsa4c}$ (Ap)	This study
p$traN_{AhD4-1}$	pBAD30::$traN_{pAhD4-1}$ (Ap)	This study
p$traN_{AQU1}$	pBAD30::$traN_{pAQU1}$ (Ap)	This study
psfx_{94}	pAH56::sfx_{pVCR94} (Kn)	This study
psfx_{Asa4c}	pAH56::sfx_{pAsa4c} (Kn)	This study
psfx_{AhD4-1}	pAH56::$sfx_{pAhD4-1}$ (Kn)	This study
psfx_{AQU1}	pAH56::sfx_{pAQU1} (Kn)	This study
psfx_{Aut1}	pAH56::$sfx_{IEVchAut1}$ (Kn)	This study
p$eexC$	pAH56::eex_{pVCR94} (Kn)	This study
p$sfx^{3×Flag}$	pAH56::$sfx_{pVCR94}{}^{3×Flag}$ (Kn)	This study
p$acaDC^{3×Flag}$	pAH56::$acaDC^{3×Flag}$ (Kn)	[7]
pVCR94^{Kn} $eexC$'-'$lacZ$	Translational fusion of the 8th codon of $lacZ$ at the 3rd codon of $vcrx145$ (Su Kn)	This study
pVCR94^{Kn} sfx'-'$lacZ$	Translational fusion of the 8th codon of $lacZ$ at the 3rd codon of $vcrx085$ (Su Kn)	This study
pVCR94^{Kn} $traN$'-'$lacZ$	Translational fusion of the 8th codon of $lacZ$ at the 3rd codon $traN$ (Su Kn)	This study
pVCR94^{KnΔCD} $eexC$'-'$lacZ$	$acaCD$ deletion mutant of pVCR94^{Kn} $eexC$'-'$lacZ$ (Su Kn Cm)	This study
pVCR94^{KnΔCD} sfx'-'$lacZ$	$acaCD$ deletion mutant of pVCR94^{Kn} sfx'-'$lacZ$ (Su Kn Cm)	This study
pVCR94^{KnΔCD} $traN$'-'$lacZ$	$acaCD$ deletion mutant of pVCR94^{Kn} $traN$'-'$lacZ$ (Su Kn Cm)	This study
p$eexC$-sfx	pBeloBAC11::(sfx-$eexC$) from pVCR94^{Kn} Δ(087-$traG$) (Cm)	This study
pBeloBAC11	Single-copy vector derived from the F plasmid (Cm)	New England Biolabs
pAH56	$oriV_{R6Kγ}$ $attP_λ$ $lacI$ P_{tac} (Kn)	[64]
pBAD30	ori_{p15A} bla $araC$ P_{BAD} (Ap)	[65]

pINT-ts	ori_{R101} $cl857$ λp_R-int_λ (ts, Ap)	[64]
pVI42b	pVI36 BamHI::P_{lac}-$lacZ$ (Sp)	[66]
pKD3	*cat* template for one-step chromosomal gene inactivation (Cm)	[67]
pKD4	*aph* template for one-step chromosomal gene inactivation (Kn)	[67]
pCP20	Flp recombinase thermo-inducible encoding plasmid (ts, Ap Cm)	[68]
pSIM6	λ Red recombination thermo-inducible encoding plasmid (ts, Ap)	[69]

Ap, ampicillin; Cm, chloramphenicol; Kn, kanamycin; Nx, nalidixic acid; Rf, rifampicin; Sm, streptomycin; Sp, Spectinomycin; Su, sulfamethoxazole; Tc, tetracycline; Tm, trimethoprim; ts, thermosensitive

Plasmid and strain construction

Plasmid DNA was prepared using the QIAprep Spin Miniprep kit (Qiagen), and genomic DNA

was isolated with the QIAamp DNA mini kit (Qiagen) as recommended by the manufacturer.

PCR products were purified using the PCR purification kit (Qiagen). All molecular biology

manipulations were carried out by standard procedures following the Current Protocols in

Molecular Biology [70]. The oligonucleotides used in this study are described in S1 Table.

Deletions were constructed using the one-step chromosomal gene inactivation technique (λ

Red-mediated mutagenesis) with pKD3 and pKD4 as templates for antibiotic resistance

cassettes [67]. The deletions of *vcrx018-vcrx019*, *vcrx062*, *vcrx085*, *vcrx107-vcrx108*, *vcrx111*,

vcrx137 and *eexC* in pVCR94^{Sp} or pVCR94^{Kn} were done using primer pairs

94delvcrx018.for/94delvcrx019.rev, 94delvcrx062.for/94delvcrx062.rev,

94delvcrx085.for/94delvcrx085.rev, 94delvcrx107.for/94delvcrx108.rev,

94delvcrx111.for/94delvcrx111.rev, 94delvcrx137.for/94delvcrx137.rev and

94del145.for/94del145.rev, respectively, and pKD3 or pKD4 as the template. Deletion of *vcrx085*

in pVCR94^{Sp} Δ*eexC* was done using primer pair 94delsfx.for/94delsfx.rev and pKD3 as the

template. The λRed recombination system was expressed using pSIM6. When appropriate,

resistance cassettes were excised from the resulting constructions using the Flp-encoding

plasmids pCP20. All deletions were validated by antibiotic profiling and PCR.

The *eexC'-'lacZ*, *sfx'-'lacZ*, and *traN'-'lacZ* translational fusions were introduced into pVCR94^{Kn}

using primer pairs 94eex-lacZ.f/94eex-lacZ.r, 94sfx-lacZ.f/94sfx-lacZ.r, and 94traN-lacZ.f/94traN-

lacZ.r, and pVI42b as the template. In all cases, the third codon of the gene of interest was fused

to the eighth codon of *lacZ*. The deletion of *acaCD* in pVCR94^{Kn}, pVCR94^{Kn} *eexC'-'lacZ*,

pVCR94^{Kn} *sfx'-'lacZ*, and pVCR94^{Kn} *traN'-'lacZ* was done using primer pair

94DelacaD.for/94DelacaC.rev and pKD3 as the template.

pVCR94 and pAsa4c served as templates to amplify their respective genes of interest. *sfx*$_{AhD4-1}$,

sfx$_{AQU1}$, *traN*$_{AhD4-1}$ and *traN*$_{AQU1}$ were chemically synthesized (Bio Basic, Markham). *sfx*$_{94}$, *sfx*$_{Asa4c}$,

sfx$_{AhD4-1}$, *sfx*$_{AQU1}$ and *eexC* were amplified using primer pairs 94sfxNdeI.f/94sfxSalI.r,

Asa4sfxNdeI.f/Asa4sfxSalI.r, AhD4sfxNdeI.f/AhD4sfxSalI.r, AQU1sfxNdeI.f/AQU1sfxSalI.r, and

eexCNdeI.f/eexCSalI.r, respectively. *sfx*$_{Aut1}$ was amplified using primer pair

Aut1sfxNdeI.f/Aut1sfxSalI.r and the genomic DNA of *V. cholerae* A12JL5W90 as the template.

Amplicons were digested with NdeI and SalI and cloned into NdeI/SalI-digested pAH56. p*sfx*^{3xFlag}

was constructed from p*sfx*$_{94}$ using primer pair pAH56_3xFlag.F/94sfx_3xFlag.R and the Q5 Site-

Directed Mutagenesis Kit (New England Biolabs) according to the manufacturer's instructions.

The resulting constructs were single-copy integrated into the *attB*$_\lambda$ chromosomal site of

CAG18439 using pINT-ts.

To assemble a DNA fragment encompassing *sfx*$_{94}$ and *eexC* alongside their native promoters,

the region spanning from *acaB* to *traG* in pVCR94^{Kn} was deleted using the one-step

chromosomal gene inactivation technique with primer pair 94del86acaB.for/94del144traG.rev

and pKD3 as the template. After the excision of the resistance cassette using pCP20, the region

of interest was amplified using primer pair 94eexBamHI.for/94sfxHindIII.rev. The amplicon was

digested with BamHI and HindIII and cloned into a BamHI/HindIII-digested pBeloBAC11.

$traN_{Asa4c}$, $traN_{AhD4-1}$, $traN_{AQU1}$ were amplified using primer pairs Asa4traNEcoRI.f/Asa4traNSalI.r,

AhD4traNEcoRI.for/AhD4traNEcoRI.rev and AQU1traNEcoRI.for/AQU1traNEcoRI.rev,

respectively. Amplicons were digested with EcoRI or EcoRI/SalI and cloned into EcoRI- or

EcoRI/SalI-digested pBAD30.

All constructs were verified by PCR and DNA sequencing at the Plateforme de Séquençage et

de Génotypage du Centre de Recherche du CHUL (Québec, QC, Canada).

Bacterial conjugation and exclusion assays

Bacteria were grown for 16 h in LB broth with the appropriate antibiotics. Mating assays were

carried out by mixing 100 µl of donor and recipient cells. Cells were pelleted by centrifugation,

then washed once in 1 volume of LB broth and resuspended in 1/20 volume of LB broth.

Bacterial mixtures were incubated for 6 h on LB agar plates at 37°C to allow conjugation. Serial

dilutions were then plated on selective LB-agar plates with appropriate antibiotics to discriminate

between donor, recipient and transconjugant CFUs. Transfer frequencies were calculated by

dividing the number of transconjugant CFUs by the number of donor CFUs. Exclusion indices

were calculated as the ratio of an element's transfer frequency toward an empty recipient to its

transfer frequency toward the tested recipient.

Transposon-directed insertion sequencing (TraDIS)

A conjugation-assisted random transposon mutagenesis experiment was performed on *E. coli*

GG56 bearing pVCR94^{Kn} Δ*eexC*::*cat*. The transposition system was composed of *E. coli* MFD*pir*

carrying pFG036 (a plasmid coding for the thermosensitive cI857 transcriptional repressor) and

pFG051 (a Pi-dependent RP4-mobilizable plasmid coding for the Tn*5* transposition machinery

repressed by cI857, and carrying a mini-Tn*5* (Sp) transposon). This diaminopimelate (DAP)-

auxotrophic strain contains a chromosomal RP4 conjugative machinery and expresses the Pi

protein required for pFG051 replication. The TraDIS experiment was performed in several

successive steps. First, pFG051 was transferred by conjugation from MFD*pir* to GG56 bearing

pVCR94^{Kn} Δ*eexC*::*cat* in a 2-h mating experiment at 30°C on LB agar plates supplemented with

DAP in duplicates. Once in the recipient strain that lacks cI, the constitutively expressed Tn*5*

machinery of pFG051 mediates random mini-Tn*5* (Sp) insertions in the genome. The mating

mixture was then entirely spread onto 40 large LB agar plates (150 mm) supplemented with Cm,

Kn, Nx, and Sp. Plates were incubated until near confluence (40k to 60k CFUs) to select clones

carrying mini-Tn*5* (Sp) insertions. After overnight incubation at 37°C, Cm Nx Kn Sp-resistant

colonies were collected using a cell scraper and resuspended in LB broth. The collected sample,

designated as the 'input library' was washed, then resuspended in 4.5 ml of LB broth and

cryopreserved. The total DNA of a 1.5 ml aliquot of the input library was extracted for

sequencing. Another 1 ml aliquot of input library was used to inoculate 50 ml of LB broth

supplemented with Cm, Kn, Nx and Sp, which was then grown overnight at 37°C. The resulting

culture was used as the recipient in a mating assay, mixed in equal volumes with *E. coli* KH95

carrying pVCR94^{Kn} and pClo used as the donor. After 2-h incubation at 37°C, mating mixtures

were spread onto 20 large LB agar plates (150 mm) supplemented with Nx, Kn, Sp, Cm, and Ap

to select for mini-Tn*5* (Sp) insertions that allowed the entry and replication of pClo into the

recipients. After overnight incubation at 37°C, Nx, Kn, Sp, Cm, and Ap-resistant colonies were

collected and subsequently resuspended in LB broth, washed and resuspended in 4.5 ml of LB

broth and cryopreserved. These samples were designated as 'output libraries'. The total DNA of

a 1.5 ml aliquot of the output libraries was extracted and used for sequencing.

Preparation of TraDIS libraries and Illumina sequencing

For each library, a 1.5 ml frozen stock aliquot was thawed on ice for 15 min and used to prepare

sequencing libraries as described previously [71]. Mutant libraries were then pooled and

sequenced by Illumina using the NextSeq® 500/550 High Output Kit v2 at the RNomics platform

of the Laboratoire de Génomique Fonctionnelle de l'Université de Sherbrooke

(https://rnomics.med.usherbrooke.ca) (Sherbrooke, QC, Canada). The transposon data analysis

was carried out as described previously [71].

Fractionation of cellular proteins

Cell fractionation was carried out using a protocol derived from Sandrini *et al.* [72]. Briefly, *E. coli*

DH5α λ*pir* bearing p*acaDC*^{3xFlag} or p*sfx*^{3xFlag} was grown overnight with or without 0.1 mM IPTG.

The cells were pelleted at 3,900 *g* for 20 min at 4°C. Cell pellets were washed twice with 10 mM

Tris buffer (pH 7.5), resuspended in 10 ml of the same buffer, and frozen for 2 h at -80°C.

Samples were lysed by ultrasonication (Qsonica Q125) in an ice bath using five cycles of 30 s

each at 70% amplitude, followed by 30 s of cooling. The lysates were treated with DNAse I (1

mg/ml). Cell debris were removed by centrifugation (3,900 *g* for 30 min at 4°C). The proteins in

the supernatant were separated by ultracentrifugation (185,000 *g* for 30 min at 4°C). The

supernatant (cytoplasmic proteins) was stored at -20°C. The pellet (total membrane proteins)

was washed three times with 10 mM Tris buffer (pH 7.5), resuspended in 10 mM Tris buffer with

2% (v/v) Triton X-100, and incubated for 30 min at room temperature. The mixture was

centrifuged at 185,000 *g* for 30 min at 4°C. The supernatant (inner membrane proteins) was

concentrated using Pierce™ Protein Concentrators PES 3K, following the manufacturer's

instructions (Thermo Scientific, cat.# 88515) and stored at -20°C. The pellet (outer membrane

proteins) was washed three times with 10 mM Tris buffer (pH 7.5) and stored at -20°C.

SDS-page and Immunoblot analysis

Bradford quantification was used to determine the quantity of fractionated proteins and

normalize protein samples. Each normalized fraction was resuspended in NuPAGE LDS Sample

Buffer (Invitrogen) supplemented with NuPAGE Sample Reducing agent (Invitrogen), incubated

at 70°C for 10 min, and separated by 12% SDS-PAGE (120 mV, 2 h). Proteins were detected by

silver staining (Pierce Silver Stain Kit) following the manufacturer's instructions (Thermo

Scientific, cat.# 24612) or transferred onto a polyvinylidene fluoride (PVDF) membrane. Western

blotting was performed according to standard procedures. The membranes were probed using

1:300 rabbit OMPC polyclonal primary antibody (ThermoFisher, cat # BS-20213R) or 1:3000

rabbit anti-FLAG primary antibody (Sigma-Aldrich, cat. # F7425), then with 1:3000 HRP goat

anti-rabbit IgG HRP-conjugated secondary antibody (Invitrogen, cat. # G-21234). Detection was

carried out with the Clarity western ECL substrate (BioRad, cat.# 170-5061) using the

ChemiDoc™ MP imaging system (BioRad).

Molecular biology

Plasmid DNA was extracted using the EZ-10 Spin Column Plasmid DNA Minipreps Kit (Bio

Basic) following the manufacturer's instructions. Enzymes used in this study were purchased

from New England Biolabs. PCR assays were performed with primers listed in Table S1. PCR

conditions were as follows: (i) 3 min at 94°C; (ii) 30 cycles of 30 s at 94°C, 30 s at the

appropriate annealing temperature, and 1 minute/kb at 68°C; and (iii) 5 min at 68°C. When

required, the resulting products were purified using the EZ-10 Spin Column PCR Products

Purification Kit (Bio Basic) following the manufacturer's instructions. *E. coli* strains were

transformed by electroporation as described previously [73] in a Bio-Rad Gene Pulser Xcell

device set at 25 µF, 200 V and 1.8 kV using 1-mm gap electroporation cuvettes.

RNA extraction and cDNA synthesis

RNA extractions were performed as follows. *E. coli* GG56 containing pVCR94^{Kn} with or without

p*acaCD* was grown at 37°C for 16 h in LB broth containing the appropriate antibiotics. The

cultures were diluted 1:200 in fresh medium containing the appropriate antibiotics and grown to

an OD_{600} of 0.2 before being diluted 1:10 again in fresh medium containing the appropriate

antibiotics and supplemented with 0.02% arabinose when needed. After a 2-h incubation period,

4 ml of the cultures were used for total RNA extraction using the Direct-zol RNA extraction kit

(Zymo Research) and TRI Reagent (Sigma-Aldrich) according to the manufacturer's instructions.

Once purified, the RNA samples were treated using 2 units of DNase I (New England Biolabs)

according to the manufacturer's instructions to eliminate residual gDNA. cDNA was synthesized

from 0.5 ng of RNA and 1 pmol of gene-specific primer 85RT (Integrated DNA Technologies)

using the reverse transcriptase SuperScript IV (Invitrogen), according to the manufacturer's

instructions. Control reactions without reverse transcriptase treatment ('RNA') were performed

for each sample. PCR reactions aiming at amplifying sfx_{94} and *acaD* were carried out using 50

pg of RNA or the corresponding amount of cDNA, or 0.5 ng of gDNA as the template, and primer

pairs sfxF/sfxR and 086F2/086R2, respectively.

β-galactosidase assays

Quantitative β-galactosidase assays were performed with 2-nitrophenyl-β-D-galactopyranoside

(ONPG) according to a protocol adapted from Miller [74]. After overnight incubation at 37°C with

appropriate antibiotics, cultures were diluted 1:100 in 50ml LB broth supplemented with 50 µg/ml

ampicillin and grown until an OD_{600} of 0.2 was reached. Three series of 1/10 dilutions were then

prepared in total volumes of 5 ml LB broth supplemented with 50 µg/ml ampicillin with or without

0.2% arabinose and incubated for 2 hours at 37°C.

Statistical analyses

Numerical data presented in graphs are available in S1 File. All analyses were performed using

R Statistical Software (v4.2.1) and RStudio (v2022.12.0.252) [75]. We assumed a normal

distribution of the frequencies of transconjugant formation and tested the homogeneity of

variance of the log10-transformed frequencies using Bartlett's tests. Dunnett and Tukey-Kramer

tests and compact letter display were performed via the multcomp R package (v1.4-20) [76].

Details of statistical tests are provided in S2 File. Graphics were rendered via the ggplot2 R

package (v3.3.6) [77]. All figures were prepared using Inkscape 0.92 (https://inkscape.org/).

Phylogenetic analyses

The primary sequence of Sfx_{94} and $TraN_{94}$ homologs were obtained using the NCBI blastp

algorithm [78] against the nr/nt database restricted to Gammaproteobacteria (taxid: 1236).

Primary sequences sharing less than 45% identity and under 85% minimum coverage were

filtered out of subsequent analyses. The distantly related Sfx_{AQU1} from pAQU1 was added

manually to the Sfx dataset as an outgroup. The datasets were clustered with CD-HIT [79] to the

best cluster that met the 0.90 identity cut-off before alignment. Where applicable, representative

sequences ("seeds") were manually substituted with sequences of interest within the same

clusters for downstream analyses. The resulting datasets were then aligned with MUSCLE [80],

and poorly aligned regions were discarded from the resulting amino acid alignments using

TrimAl v1.3 software with the automated heuristic approach [81]. Evolutionary analyses were

performed within MEGA11 (v 11.0.13) [82] using the Maximum Likelihood method (PhyML) [83]

and the Le and Gascuel matrix-based model [84] with gamma distribution (LG + G). Initial tree(s)

for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ

algorithms to a matrix of pairwise distances estimated using a JTT model and then selecting the

topology with superior log likelihood value. Percent identity values were retrieved from the

MUSCLE output [80].

Bioinformatic predictions

639   The prediction of signal peptides and cleavage sites in Sfx_{94} and $TraN_{94}$ was performed using

640   SignalP 6.0 with the slow model mode against other organisms [85]. Terminators were predicted

641   on both strands using ARNold [86,87]. *oriT* loci and mobilization proteins in putative IMEs were

642   predicted using oriTfinder [88].

Data availability

Raw sequencing data of TraDIS experiments were submitted to Genbank Sequence Read

Archive (SRA) under BioProject accession number PRJNA857050 with the following BioSample

accession numbers: SAMN32907975 and SAMN32907976 for *E. coli* GG56 bearing pVCR94^{Kn}

Δ*eexC*::*cat* as the recipient strain (input); SAMN32907978 and SAMN32907979 for *E. coli* GG56

transconjugants bearing pVCR94^{Kn} Δ*eexC*::*cat* and pSU4628 (output). Complete data from

aligned reads can also be visualized using the UCSC genome browser [89] at

https://genome.ucsc.edu/cgi-

bin/hgHubConnect?hgHub_do_redirect=on&hgHubConnect.remakeTrackHub=on&hgHub_do_fir

stDb=1&hubUrl=https://datahub-101-cw2.p.genap.ca/Rivard_2023/pVCR94dX3deexC.hub.txt.

Acknowledgement

We are grateful to Emilie Boulerne, Colin Poirier, Simon Jeanneau, and Frédéric Grenier for

technical assistance. We thank Dr. Alexander Kirschner, Medical University of Vienna, for the

kind gift of *V. cholerae* A12JL5W90.

References

1. Harmer CJ, Hall RM. The A to Z of A/C plasmids. Plasmid. 2015;80: 63–82. doi:10.1016/j.plasmid.2015.04.003

2. Doublet B, Boyd D, Mulvey MR, Cloeckaert A. The *Salmonella* genomic island 1 is an integrative mobilizable element. Molecular Microbiology. 2005;55: 1911–1924. doi:10.1111/j.1365-2958.2005.04520.x

3. Douard G, Praud K, Cloeckaert A, Doublet B. The *Salmonella* Genomic Island 1 Is Specifically Mobilized *In Trans* by the IncA/C Multidrug Resistance Plasmid Family. PLoS One. 2010;5. doi:10.1371/journal.pone.0015302

4. Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V. IncA/C Conjugative Plasmids Mobilize a New Family of Multidrug Resistance Islands in Clinical *Vibrio cholerae* Non-O1/Non-O139 Isolates from Haiti. mBio. 2016;7: e00509-16. doi:10.1128/mBio.00509-16

5. Rivard N, Colwell RR, Burrus V. Antibiotic Resistance in *Vibrio cholerae*: Mechanistic Insights from IncC Plasmid-Mediated Dissemination of a Novel Family of Genomic Islands Inserted at *trmE*. mSphere. 2020;5. doi:10.1128/mSphere.00748-20

6. Siebor E, Neuwirth C. Overview of *Salmonella* Genomic Island 1-Related Elements Among Gamma-Proteobacteria Reveals Their Wide Distribution Among Environmental Species. Front Microbiol. 2022;13: 857492. doi:10.3389/fmicb.2022.857492

7. Carraro N, Matteau D, Luo P, Rodrigue S, Burrus V. The Master Activator of IncA/C Conjugative Plasmids Stimulates Genomic Islands and Multidrug Resistance Dissemination. PLoS Genet. 2014;10. doi:10.1371/journal.pgen.1004714

8. Poulin-Laprade D, Carraro N, Burrus V. The extended regulatory networks of SXT/R391 integrative and conjugative elements and IncA/C conjugative plasmids. Front Microbiol. 2015;6. doi:10.3389/fmicb.2015.00837

9. Humbert M, Huguet KT, Coulombe F, Burrus V. Entry Exclusion of Conjugative Plasmids of the IncA, IncC, and Related Untyped Incompatibility Groups. J Bacteriol. 2019;201: e00731-18. doi:10.1128/JB.00731-18

10. Huguet KT, Rivard N, Garneau D, Palanee J, Burrus V. Replication of the *Salmonella* Genomic Island 1 (SGI1) triggered by helper IncC conjugative plasmids promotes incompatibility and plasmid loss. PLoS Genet. 2020;16. doi:10.1371/journal.pgen.1008965

11. Carraro N, Durand R, Rivard N, Anquetil C, Barrette C, Humbert M, et al. *Salmonella* genomic island 1 (SGI1) reshapes the mating apparatus of IncC conjugative plasmids to promote self-propagation. PLoS Genet. 2017;13. doi:10.1371/journal.pgen.1006705

12. Novick RP. Plasmid incompatibility. Microbiol Rev. 1987;51: 381–395.

13. Garcillán-Barcia MP, de la Cruz F. Why is entry exclusion an essential feature of conjugative plasmids? Plasmid. 2008;60: 1–18. doi:10.1016/j.plasmid.2008.03.002

14. Achtman M, Kennedy N, Skurray R. Cell-cell interactions in conjugating *Escherichia coli*:
role of *traT* protein in surface exclusion. Proc Natl Acad Sci U S A. 1977;74: 5104–5108.

15. Gunton JE, Ussher JER, Rooker MM, Wetsch NM, Alonso G, Taylor DE. Entry exclusion in
the IncHI1 plasmid R27 is mediated by EexA and EexB. Plasmid. 2008;59: 86–101.
doi:10.1016/j.plasmid.2007.11.004

16. Sakuma T, Tazumi S, Furuya N, Komano T. ExcA proteins of IncI1 plasmid R64 and Incγ
plasmid R621a recognize different segments of their cognate TraY proteins in entry
exclusion. Plasmid. 2013;69: 138–145. doi:10.1016/j.plasmid.2012.11.004

17. Pohlman RF, Genetti HD, Winans SC. Entry Exclusion of the IncN Plasmid pKM101 Is
Mediated by a Single Hydrophilic Protein Containing a Lipid Attachment Motif. Plasmid.
1994;31: 158–165. doi:10.1006/plas.1994.1017

18. Haase J, Kalkum M, Lanka E. TrbK, a small cytoplasmic membrane lipoprotein, functions in
entry exclusion of the IncP alpha plasmid RP4. J Bacteriol. 1996;178: 6720–6729.

19. Bolland S, Llosa M, Avila P, de la Cruz F. General organization of the conjugal transfer
genes of the IncW plasmid R388 and interactions between R388 and IncN and IncP
plasmids. J Bacteriol. 1990;172: 5795–5802.

20. Yamada Y, Yamada M, Nakazawa A. A ColE1-encoded gene directs entry exclusion of the
plasmid. J Bacteriol. 1995;177: 6064–6068.

21. Kamruzzaman M, Mathers AJ, Iredell JR. A Novel Plasmid Entry Exclusion System in
pKPC_UVA01, a Promiscuous Conjugative Plasmid Carrying the *bla*$_{KPC}$ Carbapenemase
Gene. Antimicrob Agents Chemother. 66: e02322-21. doi:10.1128/aac.02322-21

22. Schmitt A, Hirt H, Järvå MA, Sun W-S, ter Beek J, Dunny GM, et al. Enterococcal PrgA
Extends Far Outside the Cell and Provides Surface Exclusion to Protect against Unwanted
Conjugation. Journal of Molecular Biology. 2020;432: 5681–5695.
doi:10.1016/j.jmb.2020.08.018

23. Marrero J, Waldor MK. The SXT/R391 Family of Integrative Conjugative Elements Is
Composed of Two Exclusion Groups. J Bacteriol. 2007;189: 3302–3305.
doi:10.1128/JB.01902-06

24. Avello M, Davis KP, Grossman AD. Identification, characterization, and benefits of an
exclusion system in an integrative and conjugative element of *Bacillus subtilis*. Mol Microbiol.
2019;112: 1066–1082. doi:10.1111/mmi.14359

25. Meynell E, Ewins A. Effect on Exclusion of Alterations to the Sex Pilus. J Bacteriol.
1973;113: 71–75.

26. Marrero J, Waldor MK. Determinants of Entry Exclusion within Eex and TraG Are
Cytoplasmic. J Bacteriol. 2007;189: 6469–6473. doi:10.1128/JB.00522-07

27.   Harrison JL, Taylor IM, Platt K, O'Connor CD. Surface exclusion specificity of the TraT
730   lipoprotein is determined by single alterations in a five-amino-acid region of the protein.
731   Molecular Microbiology. 1992;6: 2825–2832. doi:10.1111/j.1365-2958.1992.tb01462.x

732   28.   Anthony KG, Sherburne C, Sherburne R, Frost LS. The role of the pilus in recipient cell
733   recognition during bacterial conjugation mediated by F-like plasmids. Molecular
734   Microbiology. 1994;13: 939–953. doi:10.1111/j.1365-2958.1994.tb00486.x

735   29.   Riede I, Eschbach M-L. Evidence that TraT interacts with OmpA of *Escherichia coli*. FEBS
736   Letters. 1986;205: 241–245. doi:10.1016/0014-5793(86)80905-X

737   30.   Klimke WA, Frost LS. Genetic Analysis of the Role of the Transfer Gene, *traN*, of the F and
738   R100-1 Plasmids in Mating Pair Stabilization during Conjugation. J Bacteriol. 1998;180:
739   4036–4043.

740   31.   Klimke WA, Rypien CD, Klinger B, Kennedy RA, Rodriguez-Maillard JM, Frost LS. The
741   mating pair stabilization protein, TraN, of the F plasmid is an outer-membrane protein with
742   two regions that are important for its function in conjugation. Microbiology. 2005;151: 3527–
743   3540. doi:10.1099/mic.0.28025-0

744   32.   Low WW, Wong JLC, Beltran LC, Seddon C, David S, Kwong H-S, et al. Mating pair
745   stabilization mediates bacterial conjugation species specificity. Nat Microbiol. 2022;7: 1016–
746   1027. doi:10.1038/s41564-022-01146-4

747   33.   Low WW, Seddon C, Beis K, Frankel G. The Interaction of the F-Like Plasmid-Encoded
748   TraN Isoforms with Their Cognate Outer Membrane Receptors. Journal of Bacteriology.
749   2023;205: e00061-23. doi:10.1128/jb.00061-23

750   34.   Ambrose SJ, Harmer CJ, Hall RM. Compatibility and entry exclusion of IncA and IncC
751   plasmids revisited: IncA and IncC plasmids are compatible. Plasmid. 2018.
752   doi:10.1016/j.plasmid.2018.02.002

753   35.   Nivaskumar M, Francetic O. Type II secretion system: A magic beanstalk or a protein
754   escalator. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 2014;1843:
755   1568–1577. doi:10.1016/j.bbamcr.2013.12.020

756   36.   Rocque WJ, McGroarty EJ. Isolation and preliminary characterization of wild-type OmpC
757   porin dimers from *Escherichia coli* K-12. Biochemistry. 1989;28: 3738–3743.
758   doi:10.1021/bi00435a017

759   37.   Hancock SJ, Phan M-D, Luo Z, Lo AW, Peters KM, Nhu NTK, et al. Comprehensive
760   analysis of IncC plasmid conjugation identifies a crucial role for the transcriptional regulator
761   AcaB. Nature Microbiology. 2020; 1–9. doi:10.1038/s41564-020-0775-0

762   38.   Daccord A, Mursell M, Poulin-Laprade D, Burrus V. Dynamics of the SetCD-Regulated
763   Integration and Excision of Genomic Islands Mobilized by Integrating Conjugative Elements
764   of the SXT/R391 Family. J Bacteriol. 2012;194: 5794–5802. doi:10.1128/JB.01093-12

39. Lawley TD, Klimke WA, Gubbins MJ, Frost LS. F factor conjugation is a true type IV
secretion system. FEMS Microbiology Letters. 2003;224: 1–15. doi:10.1016/S0378-
1097(03)00430-0

40. Nonaka L, Maruyama F, Miyamoto M, Miyakoshi M, Kurokawa K, Masuda M. Novel
Conjugative Transferable Multiple Drug Resistance Plasmid pAQU1 from *Photobacterium*
damselae subsp. *damselae* Isolated from Marine Aquaculture Environment. Microbes
Environ. 2012;27: 263–272. doi:10.1264/jsme2.ME11338

41. Tanaka KH, Vincent AT, Trudel MV, Paquet VE, Frenette M, Charette SJ. The mosaic
architecture of *Aeromonas salmonicida* subsp. *salmonicida* pAsa4 plasmid and its
consequences on antibiotic resistance. PeerJ. 2016;4: e2595. doi:10.7717/peerj.2595

42. Zhu L, Zheng J-S, Wang W-M, Luo Y. Complete Genome Sequence of Highly Virulent
Aeromonas hydrophila Strain D4, Isolated from a Diseased Blunt-Snout Bream in China.
Microbiol Resour Announc. 2019;8: e01035-18. doi:10.1128/MRA.01035-18

43. Achtman M, Manning PA, Kusecek B, Schwuchow S, Neil W. Genetic analysis of F sex
factor cistrons needed for surface exclusion in *Escherichia coli*. Journal of Molecular Biology.
1980;138: 779–795. doi:10.1016/0022-2836(80)90065-0

44. Yoshioka Y, Ohtsubo H, Ohtsubo E. Repressor gene *finO* in plasmids R100 and F:
constitutive transfer of plasmid F is caused by insertion of IS*3* into F *finO*. J Bacteriol.
1987;169: 619–623. doi:10.1128/jb.169.2.619-623.1987

45. Perumal NB, Minkley EG. The product of the F sex factor *traT* surface exclusion gene is a
lipoprotein. Journal of Biological Chemistry. 1984;259: 5357–5360. doi:10.1016/S0021-
9258(18)91011-2

46. Inselburg J. Incompatibility Exhibited by Colicin Plasmids E1, E2, and E3 in *Escherichia coli*.
J Bacteriol. 1974;119: 478–483.

47. Chan PT, Ohmori H, Tomizawa J, Lebowitz J. Nucleotide sequence and gene organization
of ColE1 DNA. Journal of Biological Chemistry. 1985;260: 8925–8935. doi:10.1016/S0021-
9258(17)39438-3

48. Friedman DI, Olson ER, Georgopoulos C, Tilly K, Herskowitz I, Banuett F. Interactions of
bacteriophage and host macromolecules in the growth of bacteriophage lambda. Microbiol
Rev. 1984;48: 299–325.

49. Garcillán-Barcia MP, Francia MV, De La Cruz F. The diversity of conjugative relaxases and
its application in plasmid classification. FEMS Microbiology Reviews. 2009;33: 657–687.
doi:10.1111/j.1574-6976.2009.00168.x

50. Szabó M, Murányi G, Kiss J. IncC helper dependent plasmid-like replication of *Salmonella*
Genomic Island 1. Nucleic Acids Res. 2021;49: 832–846. doi:10.1093/nar/gkaa1257

51. Kiss J, Szabó M, Hegyi A, Douard G, Praud K, Nagy I, et al. Identification and
Characterization of *oriT* and Two Mobilization Genes Required for Conjugative Transfer of
Salmonella Genomic Island 1. Front Microbiol. 2019;10. doi:10.3389/fmicb.2019.00457

52. Hancock SJ, Phan M-D, Roberts LW, Vu TNM, Harris PNA, Beatson SA, et al.
Characterization of DtrJ as an IncC plasmid conjugative DNA transfer component. Molecular
Microbiology. 2021;116: 154–167. doi:https://doi.org/10.1111/mmi.14697

53. Hedges RW, Datta N. Plasmids Determining I Pili Constitute a Compatibility Complex.
Microbiology. 1973;77: 19–25. doi:10.1099/00221287-77-1-19

54. Willetts N, Maule J. Interactions between the surface exclusion systems of some F-like
plasmids. Genetics Research. 1974;24: 81–89. doi:10.1017/S0016672300015093

55. Willetts N, Maule J. Specificities of IncF plasmid conjugation genes. Genet Res. 1986;47: 1–
11. doi:10.1017/s0016672300024447

56. Woodcock DM, Crowther PJ, Doherty J, Jefferson S, DeCruz E, Noyer-Weidner M, et al.
Quantitative evaluation of *Escherichia coli* host strains for tolerance to cytosine methylation
in plasmid and phage recombinants. Nucl Acids Res. 1989;17: 3469–3478.
doi:10.1093/nar/17.9.3469

57. Grenier F, Matteau D, Baby V, Rodrigue S. Complete Genome Sequence of *Escherichia coli*
BW25113. Genome Announc. 2014;2: e01038-14. doi:10.1128/genomeA.01038-14

58. Ferrières L, Hémery G, Nham T, Guérout A-M, Mazel D, Beloin C, et al. Silent mischief:
bacteriophage Mu insertions contaminate products of *Escherichia coli* random mutagenesis
performed using suicidal transposon delivery plasmids mobilized by broad-host-range RP4
conjugative machinery. J Bacteriol. 2010;192: 6418–6427. doi:10.1128/JB.00621-10

59. Ceccarelli D, Daccord A, René M, Burrus V. Identification of the Origin of Transfer (*oriT*) and
a New Gene Required for Mobilization of the SXT/R391 Family of Integrating Conjugative
Elements. J Bacteriol. 2008;190: 5328–5338. doi:10.1128/JB.00150-08

60. Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, et al. A collection of
strains containing genetically linked alternating antibiotic resistance elements for genetic
mapping of *Escherichia coli*. Microbiol Rev. 1989;53: 1–24.

61. Lepuschitz S, Baron S, Larvor E, Granier SA, Pretzer C, Mach RL, et al. Phenotypic and
Genotypic Antimicrobial Resistance Traits of *Vibrio cholerae* Non-O1/Non-O139 Isolated
From a Large Austrian Lake Frequently Associated With Cases of Human Infection. Front
Microbiol. 2019;10: 2600. doi:10.3389/fmicb.2019.02600

62. Carraro N, Sauvé M, Matteau D, Lauzon G, Rodrigue S, Burrus V. Development of
pVCR94ΔX from *Vibrio cholerae*, a prototype for studying multidrug resistant IncA/C
conjugative plasmids. Front Microbiol. 2014;5. doi:10.3389/fmicb.2014.00044

63. Cabezón E, Sastre JI, de la Cruz F. Genetic evidence of a coupling role for the TraG protein
family in bacterial conjugation. Mol Gen Genet. 1997;254: 400–406.

64. Haldimann A, Wanner BL. Conditional-Replication, Integration, Excision, and Retrieval
Plasmid-Host Systems for Gene Structure-Function Studies of Bacteria. J Bacteriol.
2001;183: 6384–6393. doi:10.1128/JB.183.21.6384-6393.2001

65. Guzman LM, Belin D, Carson MJ, Beckwith J. Tight regulation, modulation, and high-level
expression by vectors containing the arabinose P_{BAD} promoter. Journal of Bacteriology.
1995;177: 4121–4130. doi:10.1128/jb.177.14.4121-4130.1995

66. Garriss G, Waldor MK, Burrus V. Mobile Antibiotic Resistance Encoding Elements Promote
Their Own Diversity. PLoS Genet. 2009;5. doi:10.1371/journal.pgen.1000775

67. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli*
K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97: 6640–6645.

68. Cherepanov PP, Wackernagel W. Gene disruption in *Escherichia coli*: Tc^R and Km^R
cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant.
Gene. 1995;158: 9–14. doi:10.1016/0378-1119(95)00193-A

69. Datta S, Costantino N, Court DL. A set of recombineering plasmids for gram-negative
bacteria. Gene. 2006;379: 109–115. doi:10.1016/j.gene.2006.04.018

70. Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, et al. Current
Protocols in Molecular Biology. Hoboken: John Wiley & Sons; 2003.

71. Roy D, Huguet KT, Grenier F, Burrus V. IncC conjugative plasmids and SXT/R391 elements
repair double-strand breaks caused by CRISPR-Cas during conjugation. Nucleic Acids Res.
2020;48: 8815–8827. doi:10.1093/nar/gkaa518

72. Sandrini S, Haigh R, Freestone P. Fractionation by Ultracentrifugation of Gram Negative
Cytoplasmic and Membrane Proteins. BIO-PROTOCOL. 2014;4.
doi:10.21769/BioProtoc.1287

73. Dower WJ, Miller JF, Ragsdale CW. High efficiency transformation of *E. coli* by high voltage
electroporation. Nucleic Acids Res. 1988;16: 6127–6145. doi:10.1093/nar/16.13.6127

74. Miller JF. A short course in bacterial genetics. Plainview, NY: Cold Spring Harbor Laboratory
Press; 1992.

75. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing; 2021. Available: https://www.R-project.org/

76. Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric models. Biom
J. 2008;50: 346–363. doi:10.1002/bimj.200810425

77. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York, NY: Springer-Verlag;
2016.

78. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

79. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. Bioinformatics. 2006;22: 1658–1659.
doi:10.1093/bioinformatics/btl158

80. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. 2004;32: 1792–1797. doi:10.1093/nar/gkh340

81. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25: 1972–1973.
doi:10.1093/bioinformatics/btp348

82. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. Mol Biol Evol. 2021;38: 3022–3027. doi:10.1093/molbev/msab120

83. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst Biol. 2010;59: 307–321. doi:10.1093/sysbio/syq010

84. Le SQ, Gascuel O. An Improved General Amino Acid Replacement Matrix. Molecular
Biology and Evolution. 2008;25: 1307–1320. doi:10.1093/molbev/msn067

85. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, et al.
SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat
Biotechnol. 2022;40: 1023–1025. doi:10.1038/s41587-021-01156-3

86. Macke TJ, Ecker DJ, Gutell RR, Gautheret D, Case DA, Sampath R. RNAMotif, an RNA
secondary structure definition and search algorithm. Nucleic Acids Research. 2001;29:
4724–4735. doi:10.1093/nar/29.22.4724

87. Gautheret D, Lambert A. Direct RNA motif definition and identification from multiple
sequence alignments using secondary structure profiles. Journal of Molecular Biology.
2001;313: 1003–1011. doi:10.1006/jmbi.2001.5102

88. Li X, Xie Y, Liu M, Tai C, Sun J, Deng Z, et al. oriTfinder: a web-based tool for the
identification of origin of transfers in DNA sequences of bacterial mobile genetic elements.
Nucleic Acids Res. 2018;46: W229–W234. doi:10.1093/nar/gky352

89. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The Human
Genome Browser at UCSC. Genome Res. 2002;12: 996–1006. doi:10.1101/gr.229102

Supporting information

S1 Fig. Identification of an additional IncC exclusion factor by TraDIS. The tracks plot the number of reads from two independent replicates as a function of position in pVCR94^{Kn} Δ$eexC$::cat for both the input and output libraries. ORFs with similar functions are colour-coded as indicated in the figure. This figure was created using the UCSC Genome Browser (http://genome.ucsc.edu).

S2 Fig. Impact of exclusion on the fitness of IncC$^+$ cells. Colony forming units (CFUs) were obtained from mating assays conducted in Fig 1D. Crossbars show the mean and standard error of the mean of three independent experiments. One-way ANOVA (p=3.9e-6) with a Tukey-Kramer post-test was used on the log$_{10}$-transformed values to compare the means. Statistical significance (S2 File) is shown as a compact letter display for pairwise comparisons where means grouped under identical letters are not statistically different.

S3 Fig. Sfx is a predicted lipoprotein. (A) Schematic representation of the predicted translation products of sfx_{94} and $traN_{94}$ and their predicted signal peptide sequences. (B) Predicted Rho-independent terminators in the proximity of sfx in the IncC plasmid pVCR94. Terminator sequences are provided with the stems and loops in blue and red, respectively. Their position is represented by hairpins on the schematic representation of the region. Predicted ΔG values are indicated in kcal/mol.

S4 Fig. Sequence identity heatmap showing pairwise percentage identity between 18 selected Sfx homologs. Panels are shaded as indicated in the Figure according to the identity calculated for each pair.

S5 Fig. TraN directs surface exclusion specificity. *E. coli* GG56 (Nx^r) containing pVCR94^{Sp} Δ$traN$ and expressing the indicated $traN$ variants from pBAD30 (Ap^r) served as the donor, and *E. coli* CAG18439 (Tc^r) expressing the indicated sfx variants from a single-copy, chromosomally

integrated pAH56 (Kn^r) was used as the recipient. Transconjugants containing pVCR94^{Sp} Δ*traN*

were selected as the Tc^r Sp^r colonies. Crossbars show the mean and standard error of the

mean of three independent experiments. One-way ANOVA (*traN* pVCR94, p=2.21e-8; *traN*

pAsa4c, p=4.84e-10; *traN* pAhD4-1, p=2.20e-8; *traN* pAQU1, p=1.81e-9) with a Tukey-Kramer

post-test was used on the log$_{10}$-transformed values to compare the means. Exclusion indices,

shown at the bottom of each crossbar, are calculated by dividing the frequency of transfer of

pVCR94^{Sp} Δ*traN* to the empty recipient divided by the transfer frequency to the recipient

expressing the indicated *sfx* variant. Statistical significance (S2 File) is shown as a compact

letter display for pairwise comparisons where means grouped under identical letters are not

statistically different.

S1 Table. Primers used in this study.

S2 Table. Predicted AcaCD binding sites.

S1 File. Complete data sets.

S2 File. Statistical analyses.

S3 File. Sfx and TraN homologs.

A Mobilization of mini-Tn5 into *E. coli* pVCR94^Kn Δ*eexC*::*cat* → Selection of Nx^r Kn^r Cm^r Sp^r cells (input library) → Mating with *E. coli* pVCR94^Kn pClo → Selection of Nx^r Kn^r Cm^r Sp^r Ap^r cells (output library) → High-throughput sequencing of mini-Tn5 insertion sites and read mapping on reference genome

mini-Tn5 (Sp)

pVCR94^Kn Δ*eexC*::*cat*

pVCR94^Kn

pClo (Ap)

B

pVCR94^Kn Δ*eexC*::*cat*

dsbC *traC* *trhF* *traW* *traU* *traN* (*085*) *089* *090* *ssb* *bet* *exo* kb

sfx *acaB*

Input: 20,000 — 0

Output: 20,000 — 0

■ Homologous recombination ■ Host defense evasion ■ Mating pore ■ Surface exclusion ■ Transcriptional activator □ Unknown function

C

Insertion index ratio (output/input) vs Relative position in pVCR94^Kn Δ*eexC*::*cat*

018, *019*, *062*, *085*, *107*, *108*, *111*, *137*, *acr2*

D

Log₁₀(transconjugants/donor)

		a								c		c		a	
			b	b	b	b	b	b	b		b		b		b
1	287	334	252	275	318	222	292	15	262	9.2	155	0.6	270

pVCR94^Sp	–	WT	*018-019*	*062*	*107-108*	*111*	*137*	*acr2*	*sfx*	*sfx*	*eex*	*eex*	*eex sfx*	–
pAH56	–	–	–	–	–	–	–	–	–	*sfx*	–	*eex*	–	–
pBelo BAC11	–	–	–	–	–	–	–	–	–	–	–	–	–	*eex sfx*

A Sfx

- WP_187168763.1 *Eco*
- pVCR94 (**Sfx₉₄**) *Vch*
- pYULZMPS10 *Sen*
- WP_121926689.1 *Eho*
- IE*Vch*Aut1 *Vch*
- pEI-2234-3 *Eic*
- pKAZ5 *Kpn*
- WP_199478716.1 *Msp*
- IE*Ave*Tha1 *Ave*
- pAhD4-1 (**Sfx$_{AhD4-1}$**) *Ahy*
- pAsa4c (**Sfx$_{Asa4c}$**) *Asa*
- WP_234576833.1 *Asa*
- pHDA2C.2 *Hde*
- WP_214053513.1 *Pca*
- pEP-MS-18-199 *Epi*
- WP_035235290.1 *Ana*
- WP_121638668.1 *Pag*
- pAQU1 (**Sfx$_{AQU1}$**) *Pda*

Scale: 0.20

B TraN

- pYULZMPS10 *Sen*
- pRA1 *Ahy*
- NVK74585.1 *Osp*
- pVCR94 (**TraN₉₄**) *Vch*
- pHDA2C.2 *Hde*
- pBML2537 *Pst*
- GI*Vch*O27-1 *Vch*
- EGR4451686.1 *Vch*
- WP_214053512.1 *Pca*
- IME*Vch*USA3 *Vch*
- pEP-MS-18-199 *Epi*
- WP_223939723.1 *Aca*
- pAsa4c (**TraN$_{Asa4c}$**) *Asa*
- pAhD4-1 (**TraN$_{AhD4-1}$**) *Ahy*
- KXJ48173.1 *Msp*
- WP_232902095.1 *Msp*
- SGI1 *Sen*
- pAQU1 (**TraN$_{AQU1}$**) *Pda*
- WP_224067490.1 *Vpe*

Scale: 0.10

Legend:
- □ Close ACL plasmids
- ▨ IE*Vch*Aut1-like (*yicC*)
- ▨ IE*Ave*Tha1-like (*dusA*)
- ▨ SGI1-like (*trmE*)
- ▨ SGI1-like (*dusA*)
- ▨ Distant ACL plasmids
- ▨ pAQU1-like

Aca	*Aeromonas caviae*	
Ahy	*Aeromonas hydrophila*	
Ana	*Alcanivorax nanhaiticus*	
Asa	*Aeromonas salmonicida*	
Eco	*Escherichia coli*	
Eho	*Enterobacter hormaechei*	
Eic	*Edwardsiella ictaluri*	
Epi	*Edwardsiella piscicida*	
Hde	*Hamiltonella defensa*	
Kpn	*Klebsiella pneumoniae*	
Msp	*Marinobacter* sp.	
Osp	*Oceanospirillaceae* sp.	
Pag	*Pseudoalteromonas agarivorans*	
Pca	*Photorhabdus caribbeanensis*	
Pda	*Photobacterium damselae*	
Pst	*Providencia stuartii*	
Sen	*Salmonella enterica*	
Vch	*Vibrio cholerae*	
Vpe	*Vibrio penaeicida*	

C

Gene maps for pVCR94, pAsa4c, pAhD4-1, pAQU1, IE*Ave*Tha1, IE*Spu*Chn1, IE*Sch*Chn1, IE*Vch*Aut1.

Legend (scale 1kb):
- ATP synthase/ATPase
- DNA cytosine methyltransferase
- DNA processing
- DUF927
- Helix-turn-helix
- Homologous recombination
- Host defense evasion
- Integrase
- Mating pore
- MobI
- Recombination directionality factor
- Replication
- Serine protease
- Surface exclusion
- Transcriptional activator
- tRNA/RNA modification
- Type I restriction-modification
- Unknown function
- ▷ AcaCD binding site
- ★ *oriT*