Surface exclusion of IncC conjugative plasmids and their relatives

Nicolas Rivard[1], Malika Humbert[1¶], Kévin T. Huguet[1#a¶], César Pérez Bucio[2], Eve Quirion[1],

Vincent Burrus[1*]

[1]Département de biologie, Université de Sherbrooke, Sherbrooke, Quebec, Canada

[2]Instituto Tecnológico y de Estudios Superiores de Monterrey, Monterrey, Nuevo León, Mexico

[#a]Current address: Centre international de recherche en infectiologie, Lyon, France

Short title: Surface exclusion of IncC conjugative plasmids

*Corresponding author

E-mail: Vincent.Burrus@USherbrooke.ca

¶These authors contributed equally to this work.

Abstract

The phenomenon of exclusion allows conjugative plasmids to selectively impede the entry of identical or related elements into their host cell to prevent the resulting instability. Entry exclusion blocks DNA translocation into the recipient cell, whereas surface exclusion destabilizes the mating pair. IncC conjugative plasmids largely contribute to the dissemination of antibiotic-resistance genes in Gammaproteobacteria. IncC plasmids are known to exert exclusion against their relatives, including IncC and IncA plasmids, yet the entry exclusion factor *eexC* alone does not account for the totality of the exclusion phenotype. In this study, a transposon-directed insertion sequencing approach identified *sfx* as necessary and sufficient for the remaining exclusion phenotype. Sfx is a novel exclusion factor unrelated to the ones described to date. A cell fractionation assay localized Sfx in the outer membrane. Reverse transcription PCR and beta-galactosidase experiments showed that *sfx* is expressed constitutively at a higher level than *eexC*. A search in Gammaproteobacteria genomes identified Sfx homologs encoded by IncC, IncA and related, untyped conjugative plasmids and a novel family of integrative and mobilizable elements that likely rely on IncC plasmids for their mobility. Mating assays demonstrated that *sfx* is not required in the donor for exclusion, ruling out Sfx as the exclusion target. Instead, complementation assays revealed that the putative adhesin TraN in the donor is the target of surface exclusion. Mating assays with TraN homologs from related untyped plasmids from *Aeromonas* spp. and *Photobacterium damselae* identified two surface exclusion groups, with each Sfx targeting TraN homologs from the same group. Together, these results allow us to understand better the apparent incompatibility between IncA and IncC plasmids and to propose a mechanistic model for surface exclusion mediated by Sfx in IncC plasmids and related elements, with implications for the rampant dissemination of antibiotic resistance.

Author summary

Bacterial conjugation plays a pivotal role in the evolution of bacterial populations. The circulation of drug resistance genes bolsters the emergence of multidrug-resistant pathogens, with which contemporary medicine struggles to cope. Exclusion is a natural process preventing the redundant acquisition of a plasmid via conjugation by a host harbouring an identical or similar plasmid. Although exclusion has been known for the past half-century, the mechanisms involved remain poorly understood. This study describes a novel exclusion factor, Sfx, encoded by IncC, IncA and related conjugative plasmids and by unrelated integrative and mobilizable elements. We report that Sfx is a lipoprotein of the recipient that selectively inhibits conjugation based on the adhesin TraN expressed at the surface of the donor. We propose a mechanistic model for Sfx-mediated exclusion. Ultimately, a better understanding of exclusion could facilitate the design of novel conjugation inhibitors targeting mating pair formation to curb the circulation of drug-resistance genes in healthcare settings, agriculture, animal husbandry and food and drug production.

Introduction

IncC plasmids are large conjugative plasmids frequently associated with multidrug resistance phenotypes in a broad range of Gammaproteobacteria species [1]. IncC plasmids' ability to mobilize non-autonomous, unrelated integrative and mobilizable elements (IMEs) of the SGI1 and MGI*Vch*Hai6 families exacerbates their importance in antibiotic resistance dissemination [2–5]. IncC, IncA and untyped Inc<u>A</u>/<u>C</u>-<u>l</u>ike (ACL) plasmids (e.g., pAsa4c, pAhD4-1, pAQU1) share a syntenic set of genes involved in conjugation, DNA repair, and regulation. All seem regulated by closely related homologs of the master activator of transfer AcaCD [6–8]. IncC plasmid-encoded AcaCD promotes the excision and mobilization of MGI*Vch*Hai6-like IMEs [4,5]. AcaCD also stimulates the excision and replication of SGI1 and the expression of its *traN*, *traG* and *traH* genes, which modify the mating apparatus of IncC plasmids to enhance its dissemination [6,9,10].

Large conjugative plasmids are autonomous replicons that usually remain at a low-copy number. The entry into the same cell of a plasmid sharing similar replication or partitioning determinants promotes plasmid destabilization and loss, a phenomenon known as incompatibility [11]. Exclusion refers to the mechanisms employed by conjugative elements to preclude instability resulting from incompatibility by hindering redundant transfer into the cells they reside [12]. While the definite nature of exclusion mechanisms remains elusive, they can be categorized as entry and surface exclusion by whether they interfere with DNA transfer or mating pair stabilization [12,13]. Exclusion seems to be a staple of the conjugative plasmid lifestyle. Exclusion factors have been identified in the IncF, IncHI1, IncI, IncN, IncP, IncW and ColE1 families, plasmids of the pKPC_UVA01 replicon type and the enterococcal conjugative plasmid pCF10 [13–21]. Integrative and conjugative elements (ICEs) of the SXT/R391 family and ICE*Bs*1 of *Bacillus subtilis* also encode exclusion determinants [22,23]. Most exclusion

factors identified to date are associated with the cytoplasmic membrane, where they would exert

entry exclusion, with only the IncF TraT firmly established as a surface exclusion factor [12,13].

Like TraT, the IncHI1 EexB localizes in the outer membrane and is thought to engage in surface

exclusion [14]. Likewise, the surface-exposed PrgA protease of pCF10 reduces aggregation and

transfer [21].

Exclusion is not specific to the transferred DNA [24], and proteins in the donor have been

identified as targets of exclusion factors in several systems. EexA and EexB of IncHI1 plasmids

and TrbK of pKPK_UVA01 appear to act concurrently as exclusion factors and exclusion

targets, as they are necessary in both the donor and recipient for exclusion to take place

[14,20]. Other exclusion factors target different plasmid-encoded proteins. The IncI ExcA targets

TraY, which is found in the cytoplasmic membrane of the donor [15]. Similarly, the IncF TraS

and SXT/R391 Eex of the recipient seem to interact directly with the VirB6 homolog TraG of the

donor, and the regions involved in both proteins are cytoplasmic [22,25]. In the cases above, the

genes encoding the exclusion factor and its target are adjacent. In IncP, IncW and IncN

plasmids, for which the exclusion target remains unknown, the exclusion gene abuts a gene

coding for a VirB6 homolog, suggesting a widespread usage of this target beyond the IncF and

SXT/R391 families [12].

The surface exclusion factor TraT of IncF plasmids reduces the percentage of mating

aggregates, and five contiguous residues determine specificity between exclusion groups

[13,26]. Harrison *et al*. proposed that TraT destabilizes the mating pair by interacting with the

pilus tip; however, it was eventually shown that the pilin does not define exclusion specificity

[26,27]. The discovery of interactions between TraT and the outer membrane protein OmpA in

the recipient and between OmpA in the recipient and the putative adhesin TraN in the donor

suggested surface exclusion could instead result from TraT preventing docking of TraN to

OmpA, thus hindering mating pair stabilization [28–30]. The identification of TraN receptor

specificity groups across IncF plasmids makes this hypothesis extremely appealing, yet *traN*

substitutions failed to flip exclusion specificity between F and R100-1 [29,31,32].

IncC plasmids share with IncA plasmids the same entry exclusion system, *eexC*/*traG*, and

inhibit each other's entry into the cell in which they reside [8,33]. During the characterization of

eexC/*traG*, we reported that knocking-out *eexC* of the IncC plasmid pVCR94 in the recipient

only reduced the exclusion phenotype in mating assays using an IncC[+] donor. Likewise, the

deletion of a 45.6-kb fragment encompassing 33 open reading frames located between *traN* and

traF led to a statistically significant reduction of exclusion. Hence, another mechanism likely

prevents redundant transfer between cells containing identical plasmids.

In this report, we investigate the cause of the incomplete abolition of exclusion by a recipient

strain bearing an IncC plasmid lacking the entry exclusion gene *eexC*. Using a transposon-

directed insertion sequencing (TraDIS) approach, we identified IncC-borne candidate genes

whose disruption in recipient cells facilitated the acquisition of an incoming mobilizable plasmid

from an IncC[+] donor. Further analyses revealed that a single open reading frame, hereafter

named *sfx*, flanked by *traN* and *acaB*, accounts for the residual exclusion phenotype. Deletion

of *sfx* with *eexC* completely abolished exclusion. Signal peptide prediction and a subcellular

localization assay suggest the Sfx protein is an outer-membrane lipoprotein and support a role

as a surface exclusion factor. Sfx belongs to a novel family of broadly distributed surface

exclusion proteins encoded by IncC plasmids, their relatives and several putative IMEs. Surface

exclusion assays designed to test *sfx*/*traN* gene pairs from ACL plasmids revealed that *sfx* and

traN are part of a surface exclusion system, enabling us to propose a model of surface

exclusion of ACL plasmids.

Results

vcrx085 impedes conjugative transfer between IncC[+] cells

To identify additional factors responsible for the inhibition of transfer between IncC[+] cells, we constructed a high-density Tn*5* insertion library in *E. coli* GG56 carrying pVCR94[Kn] Δ*eexC*::*cat*, a kanamycin-resistant variant of pVCR94 unable to exert entry exclusion in recipient cells [8]. This set of mutants formed the input library enriched in Tn*5* insertions that allowed plasmid replication and maintenance. The input library became the recipient in a mating assay with a donor strain carrying pVCR94[Kn] and the mobilizable plasmid pClo (Fig 1A). GG56 transconjugant colonies that acquired pClo made up the output library. We expected that genes or sequences of pVCR94[Kn] Δ*eexC*::*cat* more frequently disrupted in the output than in the input (high insertion index ratio) were those impeding the mobilization of pClo between IncC[+] cells. Of nine candidate genes with insertion index ratios above the arbitrary threshold of 5 (Table 1, Fig 1C, S1 Fig), only *acr2* (*vcrx150*) is a known repressor of conjugative transfer [6]. To validate the ability of the candidates to inhibit incoming conjugative transfer, we tested corresponding deletion mutants of pVCR94[Sp] used as recipients in mating assays. All but one had no impact on transfer (Fig 1D). The deletion of *vcrx085* (hereafter referred to as *sfx* for surface exclusion) improved conjugation to a level similar to the deletion of the entry exclusion factor *eexC*. Like *eexC*, we could complement the *sfx* deletion in *trans*. Both mutations combined completely abolished exclusion, allowing conjugation to a level comparable to an empty recipient.

Table 1. Candidate genes identified by TraDIS.

ORF	Fold change	Predicted size (aa)	Predicted product[a]
vcrx018	6.39	119	Hypothetical protein
vcrx019	5.17	148	Fe3[+]-siderophore ABC transporter permease
vcrx062	8.67	194	Hypothetical protein
vcrx085 (*sfx*)	6.22	286	Lipoprotein (Sec/SPII)

vcrx107	6.07	114	Hypothetical protein
vcrx108	6.08	100	Hypothetical protein
vcrx111	6.83	184	Hypothetical protein
vcrx137	6.63	163	Hypothetical protein
acr2	6.50	139	Transcriptional repressor

152   [a]As predicted by NCBI blastp or SignalP 6.0.

Sfx is a constitutively expressed outer membrane protein

The predicted translation product of *sfx* contains a lipoprotein signal peptide (S2 Fig),

suggesting it is displayed on the cell surface [34]. A C-terminally 3xFLAG-tagged version of *sfx*

was cloned under the control of the inducible promoter P_{tac}. The Sfx-3xFLAG polypeptide has a

predicted size of 33.4 kDa before signal peptide cleavage and 31.7 kDa afterwards. A cell

fractionation assay confirmed the presence of Sfx-3xFLAG in the outer membrane fraction (Fig

2AB). This cellular location is consistent with the role of Sfx in surface exclusion. We could not

undoubtedly assign the processed and unprocessed forms in the total cell extract, and an

additional product appeared between 25 and 30 kDa on the western blot. This weaker band is

presumably a degradation product of unknown nature.

sfx is located downstream of the transfer activator *acaB*, which is expressed from an AcaCD-

inducible promoter. Hence, we wondered whether *sfx* expression was controlled by AcaCD (Fig

2C) [35]. To test this hypothesis, we constructed a translational fusion between the third codon

of *sfx* and the eighth codon of *lacZ* in pVCR94[Kn]. Expression of AcaCD did not impact the β-

galactosidase activity of the *sfx'-'lacZ* fusion. A similarly constructed *eexC'-'lacZ* fusion yielded a

~150-fold lower β-galactosidase activity (Fig 2EF). Furthermore, to test whether *sfx* and *acaB*

are part of the same transcript, we attempted to amplify both loci from a cDNA synthesized from

the 3' end of *sfx*. While *sfx* could be amplified regardless of AcaCD, *acaB* remained

undetectable, suggesting that these two genes are expressed as independent mRNA transcripts

(Fig 2D). A Rho-independent terminator predicted between *sfx* and *acaB* supports this idea (Fig

2C, S2 Fig B). Altogether, our results indicate that *sfx* is constitutively expressed independently

of AcaCD at a much higher level than *eexC*.

Genes encoding homologs of Sfx$_{94}$ are found in diverse genomic backgrounds

To explore the diversity of exclusion factors, we first used NCBI tblastn to identify Sfx$_{94}$

homologs in the ACL plasmids pAsa4c of *Aeromonas salmonicida* and pAhD4-1 of *Aeromonas*

hydrophila, and in the distant ACL plasmid pAQU1 of *Photobacterium damselae* (*traN*$_{AQU1}$) that

are known untyped relatives of IncA and IncC plasmids [7,8]. For pAsa4c and pAhD4-1, closely

related homologs were found adjacent to the gene *traN*. For pAQU1, the search resulted in a

single hit (*215*, hereafter referred to as *sfx*$_{AQU1}$) whose product shares only 39% identity over

273 amino acid residues with Sfx$_{94}$. *sfx*$_{AQU1}$ is surrounded by genes of unknown function. Sfx$_{AQU1}$

was manually added to the dataset for downstream analyses. We next searched the GenBank

database for homologs of Sfx$_{94}$ and extracted the resulting sequences for further analyses. 130

unique proteins were identified (S3 File), most encoded by ACL plasmids (Fig 3A). 78 Sfx

homologs share more than 90% identity with Sfx$_{94}$, including nine variants encoded by IncC

plasmids. All of the Sfx homologs encoded by IncC plasmids are inside this cluster. Sfx$_{94}$

belongs to an identical protein group of 193 representatives, all encoded by IncC or putative

IncC plasmids. Sfx$_{RA1}$ from the IncA plasmid pRA1 belongs to the Sfx$_{94}$ cluster as they share

92% identity over 286 amino acid residues. All plasmids outside the Sfx$_{94}$ cluster are ACL

plasmids, including pAsa4c and pAhD4-1. In most ACL plasmids and with the rare exception of

pAQU1, *sfx* lies downstream of the *dsbC-traC-trhF-traWU* operon between *traN* and the transfer

activator gene *acaB* (Fig 3C).

The remaining Sfx proteins are encoded by putative IMEs integrated at four distinct

chromosomal locations. First, ten of these IMEs belong to the MGI*Vch*Hai6 family and are

integrated at the 5' end of *dusA* in environmental and clinical *Aeromonas* strains [5].

Represented by IE*Ave*Tha1 (Fig 3C), all possess MGI*Vch*Hai6-like *mobI* and *oriT* and predicted

AcaCD-responsive promoters. A second group represented by IE*Vch*Aut1 comprises six IMEs

integrated at the 3' end of *yicC* in environmental and clinical *Vibrio* strains. In addition to a

putative recombination directionality factor related to RdfM of MGI*Vfl*Ind1 [36], these IMEs carry

a set of conserved genes, including a predicted type IV coupling protein (*traD*) and a putative

relaxase of the MOB_F family (*traI*). Finally, two IMEs sharing an identical Sfx protein are

integrated at the 3' end of *tRNA-TRP* and the 5' end of *rlmD* in environmental isolates of

Shewanella chilikensis and *Shewanella putrefaciens*, respectively. Despite sharing a conserved

core set of genes with IE*Vch*Aut1, the Sfx protein of these two IMEs resembles and clusters

with Sfx_{94} (Fig 3A). Although we can predict AcaCD binding sites across these IMEs, they do

not share any other conserved features of the SGI1 and MGI*Vch*Hai6 families, suggesting they

belong to a novel unrelated IME family mobilizable by IncC plasmids.

Surface exclusion specificity is determined by TraN

To test the role of Sfx in the donor, we monitored the transfer of pClo from donors carrying the

Δ*eexC* and Δ*sfx* deletion mutants of pVCR94^{Sp} toward an IncC$^+$ strain (pVCR94^{Kn}). No

alleviation of exclusion, i.e., an increase in transfer frequency, was observed (Fig 4A). On the

contrary, a modest yet significant decrease was observed for the Δ*eexC* Δ*sfx* mutant,

suggesting that augmented redundant transfer between donors was detrimental to transfer

toward recipients. Hence, *sfx* is not self-targeting to promote surface exclusion.

Next, we focused on TraN as a potential target for surface exclusion, as the gene *traN* encodes

a putative adhesin thought to stabilize the mating pair, and it is adjacent to *sfx* in most

conjugative plasmids found but pAQU1 [29,30,37]. A search for homologs of $TraN_{94}$ yielded 412

homologous proteins (S3 File), most encoded by ACL plasmids. Others belonged to IncC-

mobilizable IMEs of the SGI1 family integrated into *trmE* or *dusA*, with no clear relationship

between TraN phylogeny and insertion site (Fig 3B). The cluster represented by pAQU1 differs

significantly from the close ACL plasmids as the *dsbC-traC-trhF-traWUN* and *acaB*/*bet-exo*

regions are not adjacent, and *sfx* lies elsewhere.

To investigate the involvement of TraN in Sfx-dependent exclusion, we used in mating assays a

traN deletion mutant of pVCR94^{Sp} Δ*traN* complemented ectopically with either its native *traN*$_{94}$

or the distantly related homolog *traN*$_{AQU1}$ [38]. Whereas the expression of *sfx*$_{94}$ in the recipient

strongly inhibited conjugation supported by *traN*$_{94}$, it did not impair transfer supported by

traN$_{AQU1}$ (Fig 4B). This result implies TraN is, if not a direct interactant of Sfx, a key player in

Sfx-mediated exclusion and the determinant of exclusion specificity. To further investigate the

diversity of exclusion specificity, we tested the *traN*/*sfx* pairs of pAsa4c and pAhD4-1, two

plasmids that belong to distinct entry exclusion groups, as well as pAQU1 [8,39,40]. The

corresponding genes were tested in pairs in all possible combinations to determine exclusion

indices. Two separate exclusion groups emerged (Fig 4C). *sfx* homologs from pVCR94, pAsa4c

and pAhD4-1 specifically exerted exclusion against *traN* homologs from these same plasmids.

sfx$_{AQU1}$ only excluded *traN*$_{AQU1}$.

Discussion

All conjugative plasmids encode at least one exclusion factor [12]. An entry exclusion system

involving the entry exclusion factor EexC and the VirB6-homolog TraG was previously identified

in IncA and IncC conjugative plasmids [8]. Here, we identified a second exclusion factor within a

region suspected by Humbert *et al*. as responsible for additional exclusion activity [8]. A TraDIS

experiment pointed to nine candidate genes likely to enhance incoming transfer once disrupted.

sfx was the only one whose activity could be experimentally validated. Together, *eexC* and *sfx*

account for the totality of the exclusion phenotype (Fig 1D), akin to the exclusion mediated by

traS and *traT* of the F plasmid [41].

The translation product of *sfx* is a predicted lipoprotein. Here, we confirmed that Sfx is present

in the outer membrane (Fig 2B), a trait shared with the surface exclusion factor TraT of the F

plasmid and compatible with the proposed mechanism of surface exclusion [13,42]. We found

that *sfx* is constitutively expressed independently of AcaCD at a much higher level than *eexC*

(Fig 2DEF).

Our search in the Genbank database using Sfx_{94} failed to reveal any known entry or surface

exclusion factor, including F plasmid TraS or TraT, establishing Sfx as a novel surface exclusion

factor (S3 File). As pAsa4c and pAhD4-1 belong to the same surface exclusion group as

pVCR94 despite their divergence (Fig 4C), IncA and IncC plasmids are expected to exert

surface exclusion against each other, hence contributing alongside entry exclusion to the

previously observed exclusion phenotype between the two incompatibility groups [8,33]. Among

the Sfx homologs tested in this study, only Sfx_{AQU1}, which failed detection by the NCBI blastp

search due to low similarity with Sfx_{94}, belongs to a separate surface exclusion group (Fig 4C).

A pairwise comparison illustrates that similar to Sfx_{AQU1}, the putative surface exclusion proteins

WP_035235290.1 and WP_121638668.1 share minimal homology with all other representatives

(Fig 3A and S3 Fig). Therefore, we speculate that neither belongs to the two surface exclusion

groups described here and that all other IncC, IncA and close ACL plasmids belong to the Sfx_{94}

surface exclusion group. From our observations with ACL plasmids, surface exclusion strikingly

contrasts with entry exclusion, which seems much more specific. Indeed, pAsa4c and pAhD4-1

each belong to separate entry exclusion groups (EexE and EexD, respectively) distinct from the

one encompassing IncC and IncA plasmids (EexC); yet, all belong to the same surface

exclusion group [8]. These observations suggest IncC, IncA and ACL plasmids form a two-tier

community, exerting surface and entry exclusion against closely related peers but only surface

exclusion against a broader range of distantly related conjugative elements.

This work has also identified orphan surface exclusion factors borne by IMEs. IE*Ave*Tha1-like

elements belong to the MGI*Vch*Hai6 family of IncC-mobilizable IMEs [4,5], whereas IE*Vch*Aut1-

like elements constitute a new family of IMEs likely to be mobilizable by ACL plasmids (Fig 3C).

Since these IMEs lack a *traN* gene, we propose they engage in surface exclusion as a

defensive mechanism, forcing their host to favour mating partners that could enhance both their

stability and transmissibility. Hence, these IMEs could modulate the circulation of conjugative

elements in bacterial populations without affecting whether surface exclusion from a potential

recipient will target transfer mediated by their helper element. Such behaviour would contrast

with the way SGI1 manipulates the exclusion system of its helper IncC plasmid. SGI1, which is

not known to exert exclusion, evades IncC entry exclusion by substituting TraG in the mating

apparatus with a distant VirB6 homolog [10]. Manipulation of exclusion by mobilizable elements

is not well known. While homologous ColE1 mobilizable plasmids are known to inhibit each

other's transfer mildly, this phenotype was attributable to the accessory mob gene *mbeD* with no

identified target in the donor [19,43,44]. It thus remains unclear whether exclusion is at play in

that case. The putative IMEs we reported here may be the first instance of exclusion exerted by

mobilizable elements. The close relationship between Sfx proteins encoded by IE*Ave*Tha1 and

IE*Vch*Aut1 with those tested in this study (Sfx$_{94}$, Sfx$_{AhD4-1}$, and Sfx$_{Asa4c}$) suggests they belong to

the same exclusion group (Fig 3, S3 Fig). Should this activity be corroborated by experimental

evidence, it would establish exclusion as an actual barrier to the circulation of mobile genetic

elements on par with other bacterial defense mechanisms instead of being limited to preventing

redundant transfer and, potentially, mediating mating pair separation [12,37]. Surprisingly, none

of the IMEs identified by the TraN search, including SGI1, appears in the Sfx tree, suggesting

they do not encode a surface exclusion factor related to Sfx$_{94}$. Conversely, IMEs identified by

the Sfx search are ostensibly absent from the TraN tree as they lack the components of a

conjugative apparatus. This observation could indicate different survival strategies. Whereas

SGI1 maximizes its own dissemination by remodelling the mating apparatus to improve transfer

and bypass entry exclusion [10], IE*Ave*Tha1 and IE*Vch*Aut1 may ensure persistence by

hindering the entry of destabilizing, excision-inducing ACL plasmids via surface exclusion.

Although the mobilization mechanism of IE*Vch*Aut1 remains unknown, its gene content and the

presence of AcaCD binding motifs suggest mechanistic parallels with that of SGI1, albeit using

totally unrelated factors. In ACL+ cells, AcaCD likely promotes IE*Vch*Aut1's excision through the

expression of *rdfM* [36]. A predicted homolog of the replication protein P from bacteriophage

lambda suggests the excised IME is replicative [45]. IME DNA transfer seems to rely on the

IME-encoded TraI, a predicted relaxase of the MOB_F family [46]. While SGI1 encodes its own

replication and DNA processing functions [9,47,48], IE*Vch*Aut1 additionally encodes a type IV

coupling protein, presumably making it independent from the IncC-encoded TraD and DtrJ (also

known as TraJ) [5,49].

IncHI1 EexB and IncF TraT are two other known surface exclusion factors. While EexB seems

to have a dual role as an exclusion factor and exclusion target, TraT's mechanism remains

elusive as exclusion specificity could not be attributed to direct interaction with pilin or indirect

interaction with the putative adhesin TraN through OmpA [14,26,27,29]. Our results

demonstrate that Sfx is only required in the recipient to enable exclusion (Fig 4A), ruling out the

self-targeting of Sfx. Instead, we found that Sfx exclusion can be neutralized by substituting

TraN in the donor with TraN from another surface exclusion group (Fig 4BC). Distinct exclusion

groups within a given family were previously reported in IncI and IncF plasmids and SXT/R391

ICEs [15,22,50–52].

Although a direct interaction between Eex and TraG has been suggested in SXT/R391 ICEs

and likely extends to other VirB6-targeting entry exclusion factors [25], direct contact between

Sfx and TraN is unlikely for surface exclusion. In fact, the binding of outer membrane proteins

on donor and recipient cells is hard to reconcile with the idea that surface exclusion physically

prevents the formation or disrupts mating aggregates [13].

Instead, surface exclusion factors have been proposed to act by concealing outer membrane

receptors involved in mating aggregation [28]. According to Riede and Eschbach, TraT of F

directly interacts with the major outer membrane protein OmpA as TraT inhibits the OmpA-

specific phage K3 [28]. Yet, despite ample evidence that $TraN_F$ interacts with OmpA during

conjugation, it does not seem to determine exclusion specificity as $TraN_F$ substitution with

$TraN_{R100-1}$ does not counter exclusion by TraT of F [29–31].

Our finding that *sfx* is expressed at a much higher level than *eexC* despite exerting a similar

level of exclusion (Fig 1D, Fig 2EF) is consistent with an interaction with a highly abundant

surface protein. While few EexC proteins might suffice to target TraG in one mating pore during

conjugation, an abundance of Sfx proteins is likely required to conceal most, if not all, TraN

receptors on the recipient surface at all times. Unlike TraT of F, we have identified TraN as the

specificity determinant for Sfx-mediated exclusion between ACL plasmids. Thus, we propose

that $TraN_{94}$ and $TraN_{AQU1}$ recognize distinct receptors on the recipient surface and that Sfx

prevents mating pair stabilization by specific binding to the receptor of its cognate TraN. Further

investigation will be required to identify the receptors involved.

Materials and methods

Bacterial strains and media

The bacterial strains and plasmids used in this study are described in Table 2. Bacterial strains

were grown at 37°C on Luria-Bertani (LB) agar or with agitation in LB broth and maintained at

−75°C in LB broth containing 20% (vol/vol) glycerol. Antibiotics were used at the following

concentrations: ampicillin (Ap), 100 μg/ml; kanamycin (Kn), 50 μg/ml; spectinomycin (Sp), 50

μg/ml; rifampicin (Rf), 50 μg/ml; chloramphenicol (Cm), 20 μg/ml and nalidixic acid (Nx), 40

μg/ml. Media were supplemented with 0.02% L-arabinose or 0.1 mM isopropyl β-D-1-

thiogalactopyranoside (IPTG) to induce gene expression from pBAD30 and pAH56 constructs,

respectively.

Table 2. Strains and plasmids used in this study

Strain, plasmid or element	Relevant genotype or phenotype	Reference
Escherichia coli		
GG56	Nx^r derivative of BW25113 (Nx)	[53]
KH95	Rf^r derivative of BW25113 (Rf)	[9]
MFD*pir*	MG1655 RP4-2-Tc::[ΔMu1::*aac(3)IV*-Δ*aphA*-Δ*nic35*-ΔMu2::*zeo*] Δ*dapA*::(*erm-pir*) Δ*recA*	[54]
VB112	Rf^r derivative of MG1655 (Rf)	[55]
CAG18439	MG1655 *lacZU118 lacI42*::Tn*10* (Tc)	[56]
Plasmids		
pVCR94	IncC conjugative plasmid, *V. cholerae* O1 El Tor (Su Tm Cm Ap Tc Sm)	[57]
pVCR94^{Sp}	Sp^r derivative of pVCR94 (pVCR94ΔX2) (Su Sp)	[10]
pVCR94^{Kn}	Kn^r derivative of pVCR94 (pVCR94ΔX3) (Su Kn)	[10]
pVCR94^{Kn} Δ*eexC*::*cat*	*eex* deletion mutant of pVCR94^{Kn} (Su Kn Cm)	This study
pFG036	*ori*$_{pMB1}$, *cI857* (Ts) repressor, *tetM* (Tc)	Addgene #137996
pFG051	*ori*$_{R6K}$, Tn*5 tnp* under λP$_L$ promoter, *oriT*$_{RP4}$, Tn*5d-aadA7* (Sp)	Addgene #137997
pVCR94^{Sp} Δ*018-019*	*vcrx018-vcrx019* deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ*062*	*vcrx062* deletion mutant of pVCR94^{Sp} (Su Sp)	This study

pVCR94^{Sp} Δ*107-108*	*vcrx107-vcrx108* deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ*111*	*vcrx111* deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ*137*	*vcrx137* deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ*acr2*	*vcrx150* deletion mutant of pVCR94^{Sp} (Su Sp)	[6]
pVCR94^{Sp} Δ*eexC*	*vcrx145* deletion mutant of pVCR94^{Sp} (Su Sp)	[8]
pVCR94^{Sp} Δ*sfx*	*vcrx085* deletion mutant of pVCR94^{Sp} (Su Sp)	This study
pVCR94^{Sp} Δ*eexC*Δ*sfx*	*vcrx145-vcrx085* double deletion mutant of pVCR94^{Sp} (Su Sp Cm)	This study
pVCR94^{Kn} Δ*acaCD*	*acaCD* deletion mutant of pVCR94^{Kn} (Su Kn)	This study
pClo	CloDF13::TnAΔEcoRV (pSU4628) (Ap)	[58]
p*acaCD*	pBAD30::*acaCD* (Ap)	[6]
pVCR94^{Sp} Δ*traN*	*traN* deletion mutant of pVCR94^{Sp} (Su Sp)	[10]
pAsa4c	Cm^r conjugative plasmid from *A. salmonicida* JF2267 (Cm)	[39]
p*traN*$_{94}$	pBAD30::*traN*$_{pVCR94}$ (Ap)	[10]
p*traN*$_{Asa4c}$	pBAD30::*traN*$_{pAsa4c}$ (Ap)	This study
p*traN*$_{AhD4-1}$	pBAD30::*traN*$_{pAhD4-1}$ (Ap)	This study
p*traN*$_{AQU1}$	pBAD30::*traN*$_{pAQU1}$ (Ap)	This study
p*sfx*$_{94}$	pAH56::*sfx*$_{pVCR94}$ (Ap)	This study
p*sfx*$_{Asa4c}$	pAH56::*sfx*$_{pAsa4c}$ (Ap)	This study
p*sfx*$_{AhD4-1}$	pAH56::*sfx*$_{pAhD4-1}$ (Ap)	This study
p*sfx*$_{AQU1}$	pAH56::*sfx*$_{pAQU1}$ (Ap)	This study
p*eexC*	pAH56::*eex*$_{pVCR94}$ (Ap)	This study
p*sfx*$_{94}$-*3xFlag*	pAH56::*sfx*$_{pVCR94}$-*3xFlag* (Ap)	This study
pVCR94^{Kn} *eexC'*-*'lacZ*	Translational fusion of *lacZ* at the 8th codon of *vcrx145* (Su Kn)	This study
pVCR94^{Kn} *sfx'*-*'lacZ*	Translational fusion of *lacZ* at the 8th codon of *vcrx085* (Su Kn)	This study
pVCR94^{Kn} *eexC'*-*'lacZ* Δ*acaCD*	*acaCD* deletion mutant of pVCR94^{Kn} *eexC'*-*'lacZ* (Su Kn Cm)	This study
pVCR94^{Kn} *sfx'*-*'lacZ* Δ*acaCD*	*acaCD* deletion mutant of pVCR94^{Kn} *sfx'*-*'lacZ* (Su Kn Cm)	This study
pAH56	*oriV*$_{R6Kγ}$ *attP*$_{λ}$ *lacI P*$_{tac}$ (Kn)	[59]
pBAD30	*ori*$_{p15A}$ *bla araC P*$_{BAD}$ (Ap)	[60]
pINT-ts	*ori*$_{R101}$ *cI857 λp*$_{R}$-*int*$_{λ}$ (ts, Ap)	[59]
pVI42b	pVI36 BamHI::*P*$_{lac}$-*lacZ* (Sp)	[61]
pKD3	*cat* template for one-step chromosomal gene inactivation (Cm)	[62]
pKD4	*aph* template for one-step chromosomal gene inactivation (Kn)	[62]
pCP20	Flp recombinase thermo-inducible encoding plasmid (ts, Ap Cm)	[63]
pSIM6	λ Red recombination thermo-inducible encoding plasmid (ts, Ap)	[64]

Ap, ampicillin; Cm, chloramphenicol; Kn, kanamycin; Nx, nalidixic acid; Rf, rifampicin; Sm, streptomycin; Sp, Spectinomycin; Su, sulfamethoxazole; Tc, tetracycline; Tm, trimethoprim; ts, thermosensitive

Plasmids and strains construction

Plasmid DNA was prepared using the QIAprep Spin Miniprep kit (Qiagen), and genomic DNA

was isolated with the QIAamp DNA mini kit (Qiagen) as recommended by the manufacturer.

PCR products were purified using the PCR purification kit (Qiagen). All molecular biology

manipulations were carried out by standard procedures following the Current Protocols in

Molecular Biology [65]. Transformation by electroporation was done using a Biorad Gene Pulser

Xcell apparatus (BioRad) in 1 mm cuvettes under the following conditions: 200 Ω, 25 µF and

either 18 kV/cm.

The oligonucleotides used in this study are described in Supplemental Table 1. Gene deletions

were constructed using the one-step chromosomal gene inactivation technique (λ Red-mediated

mutagenesis) with pKD3 and pKD4 as templates for antibiotic resistance cassettes [62].

The deletions of *vcrx018-vcrx019*, *vcrx062*, *vcrx085*, *vcrx107-vcrx108*, *vcrx111*, *vcrx137* and

eexC in pVCR94^{Sp} or pVCR94^{Kn} were done using primer pairs

94delvcrx018.for/94delvcrx019.rev, 94delvcrx062.for/94delvcrx062.rev,

94delvcrx085.for/94delvcrx085.rev, 94delvcrx107.for/94delvcrx108.rev,

94delvcrx111.for/94delvcrx111.rev, 94delvcrx137.for/94delvcrx137.rev and

94del145.for/94del145.rev, respectively, and pKD3 or pKD4 as the template. Deletion of

vcrx085 in pVCR94^{Sp} Δ*eexC* was done using primer pair 94delsfx.for/94delsfx.rev and pKD3 as

the template. The λRed recombination system was expressed using pSIM6. When appropriate,

resistance cassettes were excised from the resulting constructions using the Flp-encoding

plasmids pCP20. All deletions were validated by antibiotic profiling and PCR.

The *eexC'-'lacZ* and *sfx'-'lacZ* fusions were introduced into pVCR94^{Kn} using primer pairs 94eex-

lacZ.f/94eex-lacZ.r and 94sfx-lacZ.f/94sfx-lacZ.r and pVI42b as the template. The deletion of

acaCD in pVCR94^{Kn}, pVCR94^{Kn} *eexC'-'lacZ* and pVCR94^{Kn} *sfx'-'lacZ* was done using primer

pair 94DelacaD.for/94DelacaC.rev and pKD3 as the template.

pVCR94 and pAsa4c served as templates to amplify their respective genes of interest. *sfx*$_{AhD4-1}$,

sfx$_{AQU1}$, *traN*$_{AhD4-1}$ and *traN*$_{AQU1}$ were chemically synthesized and obtained from Bio Basic.

sfx$_{94}$, *sfx*$_{Asa4c}$, *sfx*$_{AhD4-1}$, *sfx*$_{AQU1}$ and *eexC* were amplified using primer pairs

94sfxNdeI.f/94sfxSalI.r, Asa4sfxNdeI.f/Asa4sfxSalI.r, AhD4sfxNdeI.f/AhD4sfxSalI.r,

AQU1sfxNdeI.f/AQU1sfxSalI.r and eexCNdeI.f/eexCSalI.r, respectively. Amplicons were

digested with NdeI and SalI and cloned into NdeI/SalI-digested pAH56. p*sfx*$_{94}$-*3xFlag* was

constructed from p*sfx*$_{94}$ using primer pair pAH56_3xFlag.F/94sfx_3xFlag.R and the Q5 Site-

Directed Mutagenesis Kit (New England Biolabs) according to the manufacturer's instructions.

The resulting constructs were single-copy integrated into the *attB*$_\lambda$ chromosomal site of

BW25113 using pINT-ts.

traN$_{Asa4c}$, *traN*$_{AhD4-1}$, *traN*$_{AQU1}$ were amplified using primer pairs Asa4traNEcoRI.f/Asa4traNSalI.r,

AhD4traNEcoRI.for/AhD4traNEcoRI.rev and AQU1traNEcoRI.for/AQU1traNEcoRI.rev,

respectively. Amplicons were digested with EcoRI or EcoRI/SalI and cloned into EcoRI- or

EcoRI/SalI-digested pBAD30.

All constructs were verified by PCR and DNA sequencing at the Plateforme de Séquençage et

de Génotypage du Centre de Recherche du CHUL (Québec, QC, Canada).

Conjugation assays

Bacteria were grown for 16 h in LB broth with the appropriate antibiotics. Mating assays were

carried out by mixing 100 µl of donor and recipient cells. Cells were pelleted by centrifugation,

then washed once in 1 volume of LB broth and resuspended in 1/20 volume of LB broth.

Bacterial mixtures were incubated for 6 h on LB agar plates at 37°C to allow conjugation. Serial

dilutions were then plated on selective LB-agar plates with appropriate antibiotics to discriminate

between donor, recipient and transconjugant CFUs. Transfer frequencies were calculated by

dividing the number of transconjugant CFUs by the number of donor CFUs. Exclusion indices

were calculated as the ratio of an element's transfer frequency toward an empty recipient to its

transfer frequency toward the tested recipient.

Transposon-directed insertion sequencing (TraDIS)

A conjugation-assisted random transposon mutagenesis experiment was performed on *E. coli*

GG56 bearing pVCR94^{Kn} Δ*eexC*::*cat*. The transposition system was composed of *E. coli*

MFD*pir* carrying pFG036 (a plasmid coding for a thermosensitive cI transcriptional repressor)

and pFG051 (a Pi-dependent RP4-mobilizable plasmid coding for the Tn*5* transposition

machinery repressed by cI, and carrying a mini-Tn*5* (Sp) transposon). This diaminopimelate

(DAP)-auxotrophic strain contains a chromosomal RP4 conjugative machinery and expresses

the Pi protein required for pFG051 replication. The TnSeq experiment was performed in several

successive steps. First, pFG051 was transferred by conjugation from MFD*pir* to GG56 bearing

pVCR94^{Kn} Δ*eexC*::*cat* in a 2-h mating experiment at 30°C on LB agar plates supplemented with

DAP in duplicates. Once in the recipient strain that lacks cI, the constitutively expressed Tn*5*

machinery of pFG051 mediates random mini-Tn*5* (Sp) insertions in the genome. The mating

mixture was then entirely spread onto 40 large LB agar plates (150 mm) supplemented with Cm,

Kn, Nx, and Sp. Plates were incubated until near confluence (40k to 60k CFUs) to select clones

carrying mini-Tn*5* (Sp) insertions. After overnight incubation at 37°C, Cm Nx Kn Sp-resistant

colonies were collected using a cell scraper and resuspended in LB broth. The collected

sample, designated as the 'input library' was washed, then resuspended in 4.5 ml of LB broth

and cryopreserved. The total DNA of a 1.5 ml aliquot of the input library was extracted for

sequencing. Another 1 ml aliquot of input library was used to inoculate 50 ml of LB broth

supplemented with Cm, Kn, Nx and Sp, which was then grown overnight at 37°C. The resulting

culture was used as the recipient in a mating assay, mixed in equal volumes with *E. coli* KH95

carrying pVCR94^{Kn} and pClo. After a 2-h incubation at 37°C, mating mixtures were spread onto

20 large LB agar plates (150 mm) supplemented with Nx, Kn, Sp, Cm, and Ap to select for mini-

Tn*5* (Sp) insertions in pVCR94^{Kn} Δ*eexC*::*cat* that allowed transmission and replication of pClo

into the recipients. After overnight incubation at 37°C, Nx, Kn, Sp, Cm, and Ap-resistant

colonies were collected and subsequently resuspended in LB broth, washed and resuspended

in 4.5 ml of LB broth and cryopreserved. These samples were designated as 'output libraries'.

The total DNA of a 1.5 ml aliquot of the output libraries was extracted and used for sequencing.

Preparation of TnSeq libraries and Illumina sequencing

For each library, a 1.5 ml frozen stock aliquot was thawed on ice for 15 minutes and used to

prepare sequencing libraries as described previously [66]. Mutant libraries were then pooled

and sequenced by Illumina using the NextSeq® 500/550 High Output Kit v2 at the RNomics

platform of the Laboratoire de Génomique Fonctionnelle de l'Université de Sherbrooke

(https://rnomics.med.usherbrooke.ca) (Sherbrooke, QC, Canada). The transposon data analysis

was carried out as described previously [66].

Protein purification and analysis

E. coli BL21 containing a single-copy, chromosomally integrated p*sfx*$_{94}$-*3xFlag* was grown to an

OD$_{600}$ = 0.8 in the presence of 0.1 mM IPTG. Cells were subjected to a lysozyme/EDTA lysis

followed by selective detergent treatment to isolate the outer membrane fraction [67]. The lysate

(total cell extract) and the pellet containing the outer membrane fraction (final volume: 225 µl)

resuspended in NuPAGE LDS Sample Buffer (Invitrogen) and supplemented with NuPAGE

Sample Reducing agent (Invitrogen) were incubated at 70°C for 10 min before loading on a 12%

SDS-PAGE gel. Proteins were then either stained with Coomassie brilliant blue R-250 (Bio-Rad)

or transferred onto a polyvinylidene fluoride (PVDF) membrane. Western blotting was performed

according to standard procedures. Sfx_{94}-3xFLAG was detected using a rabbit anti-FLAG primary

antibody (Sigma-Aldrich, cat. # F7425) and an HRP goat anti-rabbit IgG HRP-conjugated

secondary antibody (Invitrogen, cat. # G-21234).

Molecular biology

Plasmid DNA was extracted using the EZ-10 Spin Column Plasmid DNA Minipreps Kit (Bio

Basic) following the manufacturer's instructions. Enzymes used in this study were purchased

from New England Biolabs. PCR assays were performed with primers listed in Table S1. PCR

conditions were as follows: (i) 3 min at 94°C; (ii) 30 cycles of 30 s at 94°C, 30 s at the

appropriate annealing temperature, and 1 minute/kb at 68°C; and (iii) 5 min at 68°C. When

required, the resulting products were purified using the EZ-10 Spin Column PCR Products

Purification Kit (Bio Basic) following the manufacturer's instructions. E. coli strains were

transformed by electroporation as described previously [68] in a Bio-Rad Gene Pulser Xcell

device set at 25 µF, 200 V and 1.8 kV using 1-mm gap electroporation cuvettes.

RNA extraction and cDNA synthesis

RNA extractions were performed as follows. *E. coli* GG56 containing $pVCR94^{Kn}$ with or without

p*acaCD* was grown at 37°C for 16 h in LB broth containing the appropriate antibiotics. The

cultures were diluted 1:200 in fresh medium containing the appropriate antibiotics and grown to

an OD_{600} of 0.2 before being diluted 1:10 again in fresh medium containing the appropriate

antibiotics and supplemented with 0.02% arabinose when needed. After a 2-h incubation period,

4 ml of the cultures were used for total RNA extraction using the Direct-zol RNA extraction kit

(Zymo Research) and TRI Reagent (Sigma-Aldrich) according to the manufacturer's

instructions. Once purified, the RNA samples were treated using 2 units of DNase I (New

England Biolabs) according to the manufacturer's instructions to eliminate residual gDNA. cDNA

was synthesized from 0.5 ng of RNA and 1 pmol of gene-specific primer 85RT (Integrated DNA

Technologies) using the reverse transcriptase SuperScript IV (Invitrogen), according to the

manufacturer's instructions. Control reactions without reverse transcriptase treatment ('RNA')

were performed for each sample. PCR reactions aiming at amplifying sfx_{94} and *acaD* were

carried out using 50 pg of RNA or the corresponding amount of cDNA, or 0.5 ng of gDNA as the

template, and primer pairs sfxF/sfxR and 086F2/086R2, respectively.

β-galactosidase assays

Qualitative assays were performed by depositing 10 μl aliquots of overnight cultures with

appropriate antibiotics on solid agar supplemented with 5-bromo-4-chloro-3-indolyl-β-D-

galactopyranoside (X-gal) with or without 0.02% arabinose. The plates were observed after

overnight incubation at 37°C.

Quantitative assays were performed with 2-nitrophenyl-β-D-galactopyranoside (ONPG)

according to a protocol adapted from Miller [69]. After overnight incubation at 37°C with

appropriate antibiotics, cultures were diluted 1:100 in 50ml LB broth supplemented with 50

μg/ml ampicillin and grown until an OD_{600} of 0.2 was reached. Three series of 1/10 dilutions

were then prepared in total volumes of 5 ml LB broth supplemented with 50 μg/ml ampicillin with

or without 0.2% arabinose and incubated for 2 hours at 37°C.

Statistical analyses for conjugation assays

Numerical data presented in graphs are available in S3 Dataset. All analyses were performed

using R Statistical Software (v4.2.1) and RStudio (v2022.12.0.252) [70]. We assumed a normal

distribution of the frequencies of transconjugant formation and tested the homogeneity of

variance of the log10-transformed frequencies using Bartlett's tests. Dunnett and Tukey-Kramer

tests and compact letter display were performed via the multcomp R package (v1.4-20) [71].

Details regarding statistical tests are provided in S2 File. Graphics were rendered via the

ggplot2 R package (v3.3.6) [72]. All figures were prepared using Inkscape 0.92

(https://inkscape.org/).

Phylogenetic analyses

The primary sequence of homologs of Sfx_{94} and $TraN_{94}$ were obtained using the NCBI blastp

algorithm [73] against the nr/nt database restricted to Gammaproteobacteria (taxid: 1236).

Primary sequences sharing less than 45% identity and under 85% minimum coverage were

filtered out of subsequent analyses. The distantly related Sfx_{AQU1} from pAQU1 was added

manually to the Sfx dataset as an outgroup. The datasets were clustered with CD-HIT [74] to

the best cluster that met the 0.90 identity cut-off before alignment. Where applicable,

representative sequences ("seeds") were manually substituted with sequences of interest within

the same clusters for downstream analyses. The resulting datasets were then aligned with

MUSCLE [75], and poorly aligned regions were discarded from the resulting amino acid

alignments using TrimAl v1.3 software with the automated heuristic approach [76]. Evolutionary

analyses were performed within MEGA11 (v 11.0.13) [77] using the Maximum Likelihood

method (PhyML) [78] and the Le and Gascuel matrix-based model [79] with gamma distribution

(LG + G). Initial tree(s) for the heuristic search were obtained automatically by applying

Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using a JTT

model and then selecting the topology with superior log likelihood value. Percent identity values

were retrieved from the MUSCLE output [75].

Bioinformatic predictions

The prediction of signal peptides and cleavage sites in Sfx_{94} and $TraN_{94}$ was performed using SignalP 6.0 with the slow model mode against other organisms [80]. Terminators were predicted on both strands using ARNold [81,82]. *oriT* loci and mobilization proteins in putative IMEs were predicted by oriTfinder [83].

Data availability

Raw sequencing data of TraDIS experiments were submitted to Genbank Sequence Read Archive (SRA) under BioProject accession number PRJNA857050 with the following BioSample accession numbers: SAMN32907975 and SAMN32907976 for *E. coli* BW25113 Nx bearing pVCR94^{Kn} Δ*eexC*::*cat* as the recipient strain (input); SAMN32907978 and SAMN32907979 for *E. coli* BW25113 Nx transconjugants bearing pVCR94^{Kn} Δ*eexC*::*cat* and pSU4628 (output). Complete data from aligned reads can also be visualized using the UCSC genome browser [84] at https://genome.ucsc.edu/cgi-bin/hgHubConnect?hgHub_do_redirect=on&hgHubConnect.remakeTrackHub=on&hgHub_do_firstDb=1&hubUrl=https://datahub-101-cw2.p.genap.ca/Rivard_2023/pVCR94dX3deexC.hub.txt.

Acknowledgement

We are grateful to Emilie Boulerne, Colin Poirier, Simon Jeanneau, and Frédéric Grenier for technical assistance.

References

1. Harmer CJ, Hall RM. The A to Z of A/C plasmids. Plasmid. 2015;80: 63–82.
doi:10.1016/j.plasmid.2015.04.003

2. Doublet B, Boyd D, Mulvey MR, Cloeckaert A. The *Salmonella* genomic island 1 is an
integrative mobilizable element. Molecular Microbiology. 2005;55: 1911–1924.
doi:10.1111/j.1365-2958.2005.04520.x

3. Douard G, Praud K, Cloeckaert A, Doublet B. The *Salmonella* Genomic Island 1 Is
Specifically Mobilized *In Trans* by the IncA/C Multidrug Resistance Plasmid Family. PLoS
One. 2010;5. doi:10.1371/journal.pone.0015302

4. Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V. IncA/C Conjugative Plasmids
Mobilize a New Family of Multidrug Resistance Islands in Clinical *Vibrio cholerae* Non-
O1/Non-O139 Isolates from Haiti. mBio. 2016;7: e00509-16. doi:10.1128/mBio.00509-16

5. Rivard N, Colwell RR, Burrus V. Antibiotic Resistance in *Vibrio cholerae*: Mechanistic
Insights from IncC Plasmid-Mediated Dissemination of a Novel Family of Genomic Islands
Inserted at *trmE*. mSphere. 2020;5. doi:10.1128/mSphere.00748-20

6. Carraro N, Matteau D, Luo P, Rodrigue S, Burrus V. The Master Activator of IncA/C
Conjugative Plasmids Stimulates Genomic Islands and Multidrug Resistance Dissemination.
PLoS Genet. 2014;10. doi:10.1371/journal.pgen.1004714

7. Poulin-Laprade D, Carraro N, Burrus V. The extended regulatory networks of SXT/R391
integrative and conjugative elements and IncA/C conjugative plasmids. Front Microbiol.
2015;6. doi:10.3389/fmicb.2015.00837

8. Humbert M, Huguet KT, Coulombe F, Burrus V. Entry Exclusion of Conjugative Plasmids of
the IncA, IncC, and Related Untyped Incompatibility Groups. J Bacteriol. 2019;201: e00731-
18. doi:10.1128/JB.00731-18

9. Huguet KT, Rivard N, Garneau D, Palanee J, Burrus V. Replication of the *Salmonella*
Genomic Island 1 (SGI1) triggered by helper IncC conjugative plasmids promotes
incompatibility and plasmid loss. PLoS Genet. 2020;16. doi:10.1371/journal.pgen.1008965

10. Carraro N, Durand R, Rivard N, Anquetil C, Barrette C, Humbert M, et al. *Salmonella*
genomic island 1 (SGI1) reshapes the mating apparatus of IncC conjugative plasmids to
promote self-propagation. PLoS Genet. 2017;13. doi:10.1371/journal.pgen.1006705

11. Novick RP. Plasmid incompatibility. Microbiol Rev. 1987;51: 381–395.

12. Garcillán-Barcia MP, de la Cruz F. Why is entry exclusion an essential feature of
conjugative plasmids? Plasmid. 2008;60: 1–18. doi:10.1016/j.plasmid.2008.03.002

13. Achtman M, Kennedy N, Skurray R. Cell-cell interactions in conjugating *Escherichia coli*:
role of *traT* protein in surface exclusion. Proc Natl Acad Sci U S A. 1977;74: 5104–5108.

14. Gunton JE, Ussher JER, Rooker MM, Wetsch NM, Alonso G, Taylor DE. Entry exclusion in
the IncHI1 plasmid R27 is mediated by EexA and EexB. Plasmid. 2008;59: 86–101.
doi:10.1016/j.plasmid.2007.11.004

15. Sakuma T, Tazumi S, Furuya N, Komano T. ExcA proteins of IncI1 plasmid R64 and Inclγ
plasmid R621a recognize different segments of their cognate TraY proteins in entry
exclusion. Plasmid. 2013;69: 138–145. doi:10.1016/j.plasmid.2012.11.004

16. Pohlman RF, Genetti HD, Winans SC. Entry Exclusion of the IncN Plasmid pKM101 Is
Mediated by a Single Hydrophilic Protein Containing a Lipid Attachment Motif. Plasmid.
1994;31: 158–165. doi:10.1006/plas.1994.1017

17. Haase J, Kalkum M, Lanka E. TrbK, a small cytoplasmic membrane lipoprotein, functions in
entry exclusion of the IncP alpha plasmid RP4. J Bacteriol. 1996;178: 6720–6729.

18. Bolland S, Llosa M, Avila P, de la Cruz F. General organization of the conjugal transfer
genes of the IncW plasmid R388 and interactions between R388 and IncN and IncP
plasmids. J Bacteriol. 1990;172: 5795–5802.

19. Yamada Y, Yamada M, Nakazawa A. A ColE1-encoded gene directs entry exclusion of the
plasmid. J Bacteriol. 1995;177: 6064–6068.

20. Kamruzzaman M, Mathers AJ, Iredell JR. A Novel Plasmid Entry Exclusion System in
pKPC_UVA01, a Promiscuous Conjugative Plasmid Carrying the bla_{KPC} Carbapenemase
Gene. Antimicrob Agents Chemother. 66: e02322-21. doi:10.1128/aac.02322-21

21. Schmitt A, Hirt H, Järvå MA, Sun W-S, ter Beek J, Dunny GM, et al. Enterococcal PrgA
Extends Far Outside the Cell and Provides Surface Exclusion to Protect against Unwanted
Conjugation. Journal of Molecular Biology. 2020;432: 5681–5695.
doi:10.1016/j.jmb.2020.08.018

22. Marrero J, Waldor MK. The SXT/R391 Family of Integrative Conjugative Elements Is
Composed of Two Exclusion Groups. J Bacteriol. 2007;189: 3302–3305.
doi:10.1128/JB.01902-06

23. Avello M, Davis KP, Grossman AD. Identification, characterization, and benefits of an
exclusion system in an integrative and conjugative element of *Bacillus subtilis*. Mol
Microbiol. 2019;112: 1066–1082. doi:10.1111/mmi.14359

24. Meynell E, Ewins A. Effect on Exclusion of Alterations to the Sex Pilus. J Bacteriol.
1973;113: 71–75.

25. Marrero J, Waldor MK. Determinants of Entry Exclusion within Eex and TraG Are
Cytoplasmic. J Bacteriol. 2007;189: 6469–6473. doi:10.1128/JB.00522-07

26. Harrison JL, Taylor IM, Platt K, O'Connor CD. Surface exclusion specificity of the TraT
lipoprotein is determined by single alterations in a five-amino-acid region of the protein.
Molecular Microbiology. 1992;6: 2825–2832. doi:10.1111/j.1365-2958.1992.tb01462.x

599 27. Anthony KG, Sherburne C, Sherburne R, Frost LS. The role of the pilus in recipient cell
600   recognition during bacterial conjugation mediated by F-like plasmids. Molecular
601   Microbiology. 1994;13: 939–953. doi:10.1111/j.1365-2958.1994.tb00486.x

602 28. Riede I, Eschbach M-L. Evidence that TraT interacts with OmpA of *Escherichia coli*. FEBS
603   Letters. 1986;205: 241–245. doi:10.1016/0014-5793(86)80905-X

604 29. Klimke WA, Frost LS. Genetic Analysis of the Role of the Transfer Gene, *traN*, of the F and
605   R100-1 Plasmids in Mating Pair Stabilization during Conjugation. J Bacteriol. 1998;180:
606   4036–4043.

607 30. Klimke WA, Rypien CD, Klinger B, Kennedy RA, Rodriguez-Maillard JM, Frost LS. The
608   mating pair stabilization protein, TraN, of the F plasmid is an outer-membrane protein with
609   two regions that are important for its function in conjugation. Microbiology. 2005;151: 3527–
610   3540. doi:10.1099/mic.0.28025-0

611 31. Low WW, Wong JLC, Beltran LC, Seddon C, David S, Kwong H-S, et al. Mating pair
612   stabilization mediates bacterial conjugation species specificity. Nat Microbiol. 2022;7: 1016–
613   1027. doi:10.1038/s41564-022-01146-4

614 32. Low WW, Seddon C, Beis K, Frankel G. The Interaction of the F-Like Plasmid-Encoded
615   TraN Isoforms with Their Cognate Outer Membrane Receptors. Journal of Bacteriology.
616   2023;205: e00061-23. doi:10.1128/jb.00061-23

617 33. Ambrose SJ, Harmer CJ, Hall RM. Compatibility and entry exclusion of IncA and IncC
618   plasmids revisited: IncA and IncC plasmids are compatible. Plasmid.
619   doi:10.1016/j.plasmid.2018.02.002

620 34. Nivaskumar M, Francetic O. Type II secretion system: A magic beanstalk or a protein
621   escalator. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 2014;1843:
622   1568–1577. doi:10.1016/j.bbamcr.2013.12.020

623 35. Hancock SJ, Phan M-D, Luo Z, Lo AW, Peters KM, Nhu NTK, et al. Comprehensive
624   analysis of IncC plasmid conjugation identifies a crucial role for the transcriptional regulator
625   AcaB. Nature Microbiology. 2020; 1–9. doi:10.1038/s41564-020-0775-0

626 36. Daccord A, Mursell M, Poulin-Laprade D, Burrus V. Dynamics of the SetCD-Regulated
627   Integration and Excision of Genomic Islands Mobilized by Integrating Conjugative Elements
628   of the SXT/R391 Family. J Bacteriol. 2012;194: 5794–5802. doi:10.1128/JB.01093-12

629 37. Lawley TD, Klimke WA, Gubbins MJ, Frost LS. F factor conjugation is a true type IV
630   secretion system. FEMS Microbiology Letters. 2003;224: 1–15. doi:10.1016/S0378-
631   1097(03)00430-0

632 38. Nonaka L, Maruyama F, Miyamoto M, Miyakoshi M, Kurokawa K, Masuda M. Novel
633   Conjugative Transferable Multiple Drug Resistance Plasmid pAQU1 from *Photobacterium*
634   *damselae* subsp. *damselae* Isolated from Marine Aquaculture Environment. Microbes
635   Environ. 2012;27: 263–272. doi:10.1264/jsme2.ME11338

39. Tanaka KH, Vincent AT, Trudel MV, Paquet VE, Frenette M, Charette SJ. The mosaic
architecture of *Aeromonas salmonicida* subsp. *salmonicida* pAsa4 plasmid and its
consequences on antibiotic resistance. PeerJ. 2016;4: e2595. doi:10.7717/peerj.2595

40. Zhu L, Zheng J-S, Wang W-M, Luo Y. Complete Genome Sequence of Highly Virulent
Aeromonas hydrophila Strain D4, Isolated from a Diseased Blunt-Snout Bream in China.
Microbiol Resour Announc. 2019;8: e01035-18. doi:10.1128/MRA.01035-18

41. Achtman M, Manning PA, Kusecek B, Schwuchow S, Neil W. Genetic analysis of F sex
factor cistrons needed for surface exclusion in *Escherichia coli*. Journal of Molecular
Biology. 1980;138: 779–795. doi:10.1016/0022-2836(80)90065-0

42. Perumal NB, Minkley EG. The product of the F sex factor *traT* surface exclusion gene is a
lipoprotein. Journal of Biological Chemistry. 1984;259: 5357–5360. doi:10.1016/S0021-
9258(18)91011-2

43. Inselburg J. Incompatibility Exhibited by Colicin Plasmids E1, E2, and E3 in *Escherichia coli*.
J Bacteriol. 1974;119: 478–483.

44. Chan PT, Ohmori H, Tomizawa J, Lebowitz J. Nucleotide sequence and gene organization
of ColE1 DNA. Journal of Biological Chemistry. 1985;260: 8925–8935. doi:10.1016/S0021-
9258(17)39438-3

45. Friedman DI, Olson ER, Georgopoulos C, Tilly K, Herskowitz I, Banuett F. Interactions of
bacteriophage and host macromolecules in the growth of bacteriophage lambda. Microbiol
Rev. 1984;48: 299–325.

46. Garcillán-Barcia MP, Francia MV, De La Cruz F. The diversity of conjugative relaxases and
its application in plasmid classification. FEMS Microbiology Reviews. 2009;33: 657–687.
doi:10.1111/j.1574-6976.2009.00168.x

47. Szabó M, Murányi G, Kiss J. IncC helper dependent plasmid-like replication of *Salmonella*
Genomic Island 1. Nucleic Acids Res. 2021;49: 832–846. doi:10.1093/nar/gkaa1257

48. Kiss J, Szabó M, Hegyi A, Douard G, Praud K, Nagy I, et al. Identification and
Characterization of *oriT* and Two Mobilization Genes Required for Conjugative Transfer of
Salmonella Genomic Island 1. Front Microbiol. 2019;10. doi:10.3389/fmicb.2019.00457

49. Hancock SJ, Phan M-D, Roberts LW, Vu TNM, Harris PNA, Beatson SA, et al.
Characterization of DtrJ as an IncC plasmid conjugative DNA transfer component. Molecular
Microbiology. 2021;116: 154–167. doi:https://doi.org/10.1111/mmi.14697

50. Hedges RW, Datta N. Plasmids Determining I Pili Constitute a Compatibility Complex.
Microbiology. 1973;77: 19–25. doi:10.1099/00221287-77-1-19

51. Willetts N, Maule J. Interactions between the surface exclusion systems of some F-like
plasmids. Genetics Research. 1974;24: 81–89. doi:10.1017/S0016672300015093

52. Willetts N, Maule J. Specificities of IncF plasmid conjugation genes. Genet Res. 1986;47: 1–
11. doi:10.1017/s0016672300024447

53. Grenier F, Matteau D, Baby V, Rodrigue S. Complete Genome Sequence of *Escherichia coli*
BW25113. Genome Announc. 2014;2: e01038-14. doi:10.1128/genomeA.01038-14

54. Ferrières L, Hémery G, Nham T, Guérout A-M, Mazel D, Beloin C, et al. Silent Mischief:
Bacteriophage Mu Insertions Contaminate Products of Escherichia coli Random
Mutagenesis Performed Using Suicidal Transposon Delivery Plasmids Mobilized by Broad-
Host-Range RP4 Conjugative Machinery. J Bacteriol. 2010;192: 6418–6427.
doi:10.1128/JB.00621-10

55. Ceccarelli D, Daccord A, René M, Burrus V. Identification of the Origin of Transfer (*oriT*) and
a New Gene Required for Mobilization of the SXT/R391 Family of Integrating Conjugative
Elements. J Bacteriol. 2008;190: 5328–5338. doi:10.1128/JB.00150-08

56. Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, et al. A collection of
strains containing genetically linked alternating antibiotic resistance elements for genetic
mapping of *Escherichia coli*. Microbiol Rev. 1989;53: 1–24.

57. Carraro N, Sauvé M, Matteau D, Lauzon G, Rodrigue S, Burrus V. Development of
pVCR94ΔX from *Vibrio cholerae*, a prototype for studying multidrug resistant IncA/C
conjugative plasmids. Front Microbiol. 2014;5. doi:10.3389/fmicb.2014.00044

58. Cabezón E, Sastre JI, de la Cruz F. Genetic evidence of a coupling role for the TraG protein
family in bacterial conjugation. Mol Gen Genet. 1997;254: 400–406.

59. Haldimann A, Wanner BL. Conditional-Replication, Integration, Excision, and Retrieval
Plasmid-Host Systems for Gene Structure-Function Studies of Bacteria. J Bacteriol.
2001;183: 6384–6393. doi:10.1128/JB.183.21.6384-6393.2001

60. Guzman LM, Belin D, Carson MJ, Beckwith J. Tight regulation, modulation, and high-level
expression by vectors containing the arabinose P_{BAD} promoter. Journal of Bacteriology.
1995;177: 4121–4130. doi:10.1128/jb.177.14.4121-4130.1995

61. Garriss G, Waldor MK, Burrus V. Mobile Antibiotic Resistance Encoding Elements Promote
Their Own Diversity. PLoS Genet. 2009;5. doi:10.1371/journal.pgen.1000775

62. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli*
K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97: 6640–6645.

63. Cherepanov PP, Wackernagel W. Gene disruption in *Escherichia coli*: Tc[R] and Km[R]
cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant.
Gene. 1995;158: 9–14. doi:10.1016/0378-1119(95)00193-A

64. Datta S, Costantino N, Court DL. A set of recombineering plasmids for gram-negative
bacteria. Gene. 2006;379: 109–115. doi:10.1016/j.gene.2006.04.018

65. Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, et al. Current
Protocols in Molecular Biology. Hoboken: John Wiley & Sons; 2003.

66. Roy D, Huguet KT, Grenier F, Burrus V. IncC conjugative plasmids and SXT/R391 elements
repair double-strand breaks caused by CRISPR-Cas during conjugation. Nucleic Acids Res.
2020;48: 8815–8827. doi:10.1093/nar/gkaa518

67. Thein M, Sauer G, Paramasivam N, Grin I, Linke D. Efficient Subfractionation of Gram-
Negative Bacteria for Proteomics Studies. J Proteome Res. 2010;9: 6135–6147.
doi:10.1021/pr1002438

68. Dower WJ, Miller JF, Ragsdale CW. High efficiency transformation of *E. coli* by high voltage
electroporation. Nucleic Acids Res. 1988;16: 6127–6145. doi:10.1093/nar/16.13.6127

69. Miller JF. A short course in bacterial genetics. Plainview, NY: Cold Spring Harbor Laboratory
Press; 1992.

70. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing; 2021. Available: https://www.R-project.org/

71. Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric models. Biom
J. 2008;50: 346–363. doi:10.1002/bimj.200810425

72. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York, NY: Springer-Verlag;
2016.

73. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

74. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. Bioinformatics. 2006;22: 1658–1659.
doi:10.1093/bioinformatics/btl158

75. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. 2004;32: 1792–1797. doi:10.1093/nar/gkh340

76. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25: 1972–1973.
doi:10.1093/bioinformatics/btp348

77. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. Mol Biol Evol. 2021;38: 3022–3027. doi:10.1093/molbev/msab120

78. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst Biol. 2010;59: 307–321. doi:10.1093/sysbio/syq010

79. Le SQ, Gascuel O. An Improved General Amino Acid Replacement Matrix. Molecular
Biology and Evolution. 2008;25: 1307–1320. doi:10.1093/molbev/msn067

80. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, et al.
SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat
Biotechnol. 2022;40: 1023–1025. doi:10.1038/s41587-021-01156-3

81. Macke TJ, Ecker DJ, Gutell RR, Gautheret D, Case DA, Sampath R. RNAMotif, an RNA
secondary structure definition and search algorithm. Nucleic Acids Research. 2001;29:
4724–4735. doi:10.1093/nar/29.22.4724

82. Gautheret D, Lambert A. Direct RNA motif definition and identification from multiple
sequence alignments using secondary structure profiles. Journal of Molecular Biology.
2001;313: 1003–1011. doi:10.1006/jmbi.2001.5102

83. Li X, Xie Y, Liu M, Tai C, Sun J, Deng Z, et al. oriTfinder: a web-based tool for the
identification of origin of transfers in DNA sequences of bacterial mobile genetic elements.
Nucleic Acids Res. 2018;46: W229–W234. doi:10.1093/nar/gky352

84. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The Human
Genome Browser at UCSC. Genome Res. 2002;12: 996–1006. doi:10.1101/gr.229102

Fig 1. Identification of an additional IncC exclusion factor. (A) Overview of the TnSeq

workflow to identify exclusion factors in pVCR94^{Kn} Δ*eexC*::*cat*. (B) Mini-Tn5 reads in the vicinity

of *vcrx085*. The tracks plot the number of reads from two independent replicates as a function of

position in pVCR94^{Kn} Δ*eexC*::*cat* for both the input and output libraries. ORFs with similar

functions are colour-coded as indicated in the panel. (C) Nine loci of unknown function show a

mini-Tn*5* insertion index ratio greater than 5 (dashed line), calculated as the ratio of insertion

counts between the output and input libraries. ORFs with similar functions are colour-coded as

indicated in Fig S1. (D) *sfx* (*vcrx085*) is the only functional exclusion factor among candidates

identified in (B) and suffices alongside *eexC* (*eex*) to account for the totality of IncC exclusion.

E. coli VB112 containing pVCR94^{Kn} and pClo served as the donor strain, and *E. coli* GG56

containing the specified derivatives of pVCR94^{Sp} served as the recipient strains. When

indicated, *eexC* or *sfx* were expressed from single-copy, chromosomally integrated pAH56 in

the recipients in the presence of IPTG. Transconjugants containing pClo were selected as the

Nx^r Ap^r colonies. This figure was created using the UCSC Genome Browser

(http://genome.ucsc.edu).

Fig 2. Expression of IncC surface exclusion is independent of the master activator of

transfer AcaCD. (A) and (B) Cell fractionation assay. The total cell extract (T) and outer

membrane fraction (OM) from *E. coli* BL21 expressing a 3xFLAG-tagged Sfx$_{94}$ were analyzed

by SDS-PAGE and stained with Coomassie blue (A) or immunodetected using an anti-FLAG

antibody (B). Molecular weight markers (M) are indicated on the left. A vertical white line

separates non-adjacent lanes from a single gel or blot. (C) Schematic representation of the

acaB-*sfx* locus. The relative positions of reverse transcription primer 85RT and primers used to

amplify *sfx* and *acaB* are shown. A dotted line shows the reverse transcription product. Stem

loops represent predicted Rho-independent terminators. (D) *sfx* is expressed constitutively and

independently of the upstream AcaCD-dependent P_{acaB} promoter. A 2% agarose gel of the PCR

products from an *sfx*-derived reverse transcribed RNA. Untreated RNA (RNA) and genomic

DNA from *E. coli* GG56 containing pVCR94^{Kn} Δ*acaCD* (gDNA) were used as negative and

positive controls, respectively. β-galactosidase activity of GG56 containing pVCR94^{Kn} *eexC'-*

'lacZ or pVCR94^{Kn} *sfx'-'lacZ* transcriptional fusions with or without p*acaCD* and arabinose on

agar medium (E) or in quantitative enzymatic assays (F).

Fig 3. Sfx homologs are found across a wide range of Gammaproteobacteria. Maximum

likelihood phylogenetic analyses of Sfx (A) and TraN (B) homologs. Trees with the highest

likelihood (9,621.78 and 28,204.23 for Sfx and TraN, respectively) are shown. Bootstrap

supports are indicated as percentages at the branching points only when >80%. Branch lengths

represent the number of substitutions per site over 269 and 813 amino acid positions for Sfx

and TraN, respectively. Only one representative per cluster of similar proteins (>90%) is shown

in each tree, except for TraN$_{Asa4c}$ and TraN$_{AhD4-1}$, which share 95% identity over the aligned

sequences. S5 Table provides additional details on host strains. (C) Schematic representation

of the genetic context of *sfx* homologs in a selection of putative IMEs and ACL plasmids.

Predicted *attL* and *attR* attachment sites flanking the IMEs are represented by blue bars.

pAQU1 encodes homologs of *acaB*, *vcrx087-088-089-090*, *bet* and *exo* clustered in a distinct

locus not shown here. ORFs with similar functions are colour-coded as indicated in the panel.

Fig 4. TraN directs surface exclusion specificity. (A) The presence of *eexC* and *sfx* in the

donor decreases the frequency of transconjugant formation. pClo was mobilized from *E. coli*

VB112 bearing the indicated deletion mutants of pVCR94^{Sp} (donor) into *E. coli* GG56 bearing

pVCR94^{Kn} (recipient). Transconjugants containing pClo were selected as the Nx^r Ap^r colonies.

(B) *sfx*$_{94}$ is ineffective against the *traN*$_{94}$ distant homolog *traN*$_{AQU1}$. (C) Heatmap of surface

exclusion indices of *sfx* and *traN* pairs from a diverse set of ACL plasmids. In (B) and (C), *E. coli*

GG56 containing pVCR94^{Sp} Δ*traN* and expressing the indicated *traN* variant from pBAD30

served as the donor, and *E. coli* CAG18439 expressing the indicated *sfx* variant from a single-

copy, chromosomally integrated pAH56 was used as the recipient. Transconjugants containing

pVCR94^{Sp} Δ*traN* were selected as the Tc^r Sp^r colonies.

Supporting information

S1 Fig. Identification of an additional IncC exclusion factor by TraDIS. The tracks plot the

number of reads from two independent replicates as a function of position in pVCR94^{Kn}

Δ*eexC*::*cat* for both the input and output libraries. ORFs with similar functions are colour-coded

as indicated in the figure. This figure was created using the UCSC Genome Browser

(http://genome.ucsc.edu).

S2 Fig. Sfx is a predicted lipoprotein. (A) Schematic representation of the predicted

translation products of *sfx*$_{94}$ and *traN*$_{94}$ and their predicted signal peptide sequences. (B)

Predicted Rho-independent terminators in the proximity of *sfx* in the IncC plasmid pVCR94.

Terminator sequences are provided with the stems and loops in blue and red, respectively.

Their position is represented by hairpins on the schematic representation of the region.

Predicted ΔG values are indicated in kcal/mol.

S3 Fig. Sequence identity heatmap showing pairwise percentage identity between 18

selected Sfx homologs. Panels are shaded as indicated in the Figure according to the identity

calculated for each pair.

S4 Fig. TraN directs surface exclusion specificity. *E. coli* GG56 containing pVCR94^{Sp} Δ*traN*

and expressing the indicated *traN* variants from pBAD30 served as the donor, and *E. coli*

CAG18439 expressing the indicated *sfx* variants from a single-copy, chromosomally integrated

pAH56 was used as the recipient. Transconjugants containing pVCR94^{Sp} Δ*traN* were selected

as the Tc^r Sp^r colonies.

S1 Table. Primers used in this study.

S2 Table. Strains and plasmids used in this study.

S3 Table. Predicted AcaCD binding sites.

S1 File. Complete data sets.

S2 File. Statistical analyses.

S3 File. Sfx and TraN homologs.

Figure 1

Figure 3

Figure 4

Figure 2