Interaction of amisulpride with GLUT1 at the blood-brain barrier. Relevance to Alzheimer's disease

Sevda T. Boyanova[1], Ethlyn Lloyd-Morris[1], Christopher Corpe[2], K. Miraz Rahman[1], Doaa B. Farag[3], Lee K. Page[1], Hao Wang[1], Alice L. Fleckney[1], Ariana Gatt[4], Claire Troakes[5], Gema Vizcay-Barrena[6], Roland Fleck[6], Suzanne J. Reeves[7], and Sarah A. Thomas[1] *

[1] King's College London, Institute of Pharmaceutical Science, 150 Stamford St, SE1 9NH, London, UK; [2] King's College London, Department of Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences and Medicine, [3]Faculty of Pharmacy, Misr International University, 11431, Cairo, Egypt; [4] King's College London, Wolfson Centre for Age Related Disease, 20 Newcomen St, SE1 1YR, London, UK; [5] King's College London, London Neurodegenerative Diseases Brain Bank, IoPPN, De Crespigny Park, SE5 8AF, London, UK; [6] King's College London, Centre for Ultrastructural Imaging, New Hunt's House, Guy's Campus, SE1 1UL, London, UK; [7]Faculty of Brain Sciences, University College London, 149 Tottenham Court Road, W1T 7NF, London, UK

Running title or short title: Interaction of amisulpride with GLUT1 at the blood-brain barrier.

*Correspondence:

sarah.thomas@kcl.ac.uk[1] King's College London, Institute of Pharmaceutical Science, 150 Stamford St, London SE1 9NH, UK

Abstract

Blood-brain barrier (BBB) dysfunction may be involved in the increased sensitivity of Alzheimer's disease (AD) patients to antipsychotics, including amisulpride. Studies indicate that antipsychotics interact with facilitated glucose transporters (GLUT), including GLUT1, and that GLUT1 BBB expression decreases in AD. We tested the hypotheses that amisulpride (charge: +1) interacts with GLUT1, and that BBB transport of amisulpride is compromised in AD.

GLUT1 substrates and inhibitors, and GLUT-interacting antipsychotics were identified by literature review and their physicochemical characteristics summarised. Interactions between amisulpride, and GLUT1 were studied using *in silico* approaches and the human cerebral endothelial cell line, hCMEC/D3. Brain distribution of [^{3}H]amisulpride was determined using *in situ* perfusion in wild type (WT) and 5xFamilial AD (5xFAD) mice. With transmission electron microscopy (TEM) we investigated brain capillary degeneration in WT and 5xFAD mice, and human samples. Western blots determined BBB transporter expression in mouse and human.

Literature review revealed that, although D-glucose has no charge, charged molecules can interact with GLUT1. GLUT1 substrates are smaller (184.95±6.45g/mol) than inhibitors (325.50±14.40g/mol), and GLUT-interacting antipsychotics (369.38±16.04). Molecular docking showed beta-D-glucose (free energy binding: -15.39kcal/mol) and amisulpride (-29.04kcal/mol) interact with GLUT1. Amisulpride did not affect [^{14}C]D-glucose accumulation in hCMEC/D3. 5xFAD mice showed increased brain [^{3}H]amisulpride uptake, and no cerebrovascular space changes compared to WT. TEM revealed brain capillary degeneration in human AD. There was no significant effect of

AD on mouse GLUT1 and P-gp BBB expression, and in human GLUT1 expression. In

contrast, caudate P-glycoprotein expression was decreased in human AD capillaries

versus controls.

This study provides new details about the BBB transport of amisulpride, evidence that

amisulpride interacts with GLUT1, and that BBB transporter expression is altered in

AD. This suggests that antipsychotics exacerbate the cerebral hypometabolism in AD.

Further research into the mechanism of amisulpride transport by GLUT1 is important

for improving antipsychotics safety.

Keywords: blood-brain barrier; amisulpride; adverse events; transporters; GLUT1.

Introduction

Psychosis, which most commonly presents as delusions, is highly prevalent (~50%) in people with Alzheimer's disease (AD) (1,2), and is associated with poorer quality of life, a more rapid speed of cognitive and functional decline (2) and greater risk of institutionalisation (3). Safe and effective prescribing of antipsychotic drugs is challenging in older people, particularly those with AD, due to their heightened susceptibility to the side effects of these drugs (4), including sedation, postural hypotension, parkinsonism, and an increased risk of stroke and death (5,6). As a result, antipsychotic use is restricted to those with severe psychosis and/or aggression that have not responded to non-pharmacological approaches. In the UK, the National Institute for Health and Care Excellence guidance advocates use of 'the lowest possible dose for the shortest possible time' (7). There is a lack of guidance on the minimum clinically effective dose for individual drugs, or the factors that predict differing response and side effects, although a recent publication, based on clinical data on risperidone use in AD, suggests that increased dementia severity is an independent risk factor for emergent parkinsonism (7).

Amisulpride is a second-generation antipsychotic – a substituted benzamide derivative, which is a highly selective dopamine D2 receptor antagonist (8). It can be prescribed off-label in very late onset (> 60 years) schizophrenia-like psychosis (9), and in patients with AD psychosis (10–12). Therapeutic drug monitoring studies in adults predominantly aged under 65 years have shown that therapeutic striatal D2/3 receptor occupancies between 40-70% are achieved at blood drug concentrations of 100-319

ng/ml; equivalent to 400-800 mg/day (13–15). Older patients with AD psychosis (aged

69-92 years) showed a clinical response to amisulpride and parkinsonian side effects at

lower doses (25-75 mg/day) with high D2/3 receptor occupancies in the caudate (41-

83%) and lower blood drug concentrations (40-100 ng/ml) than expected (11,12).

These findings suggest that age and/or AD-related changes in central pharmacokinetics

contribute to antipsychotic drug sensitivity and implicate the blood-brain barrier (BBB)

which controls the entry of drugs to the brain through selective transport pathways.

Evidence from our animal studies further supports this hypothesis (16).   Since

amisulpride is predominantly positively (17) charged at physiological pH, observed

changes in the expression and/or function of BBB transporters for organic cations may

help explain the increased amisulpride sensitivity in AD patients. In particular, the

organic cation transporter 1 (OCT1; SLC22A1) and the plasma membrane monoamine

transporter (PMAT; SLC29A4) (16).

Compromised function and expression of other SLC transporters at the BBB, such as

the glucose transporter, GLUT1 (SLC2A1), has also been observed in AD (18–20).

Consequently, AD brains suffer chronic shortages of energy-rich metabolites (21,22).

Importantly, antipsychotic drugs, including risperidone and clozapine, inhibit glucose

uptake by transporters, it has been suggested that this could be through direct

interaction with GLUT transporters (23,24). In addition, the use of clozapine and

another antipsychotic drug, olanzapine, has been associated with the development of

type 2 diabetes. One of the proposed mechanisms is through the inhibition of glucose

transporters inducing hyperglycaemia (25–27). However, detailed studies of the drugs-

transporter interaction, including *in silico* molecular docking studies on the specific

molecular interactions of antipsychotics with GLUT1 are rare.

Thus, considering the changes in GLUT1 expression at the BBB and resulting impact

on brain metabolism in AD, and the increased sensitivity of AD patients to

antipsychotic drugs including risperidone and amisulpride, we wanted to explore the

interaction between GLUT1 and amisulpride. To do this we tested the hypotheses that

amisulpride interacts with GLUT1 at the BBB, and that expression of BBB transporters

and the transport of amisulpride and glucose into the brain will be affected by AD. We

utilised a combination of literature review, physicochemical properties analysis,

molecular docking approaches, cell culture BBB studies, studies in wild type mice and

in an animal model of AD and assessment of tissue from human cases with and without

AD. An overview of the methods deployed are presented in Supplementary Fig 1, S1

File. Abstracts of this work have been presented (28).

Materials and Methods

Radiolabelled and non-labelled chemicals

Radiolabelled [^{14}C]D-glucose (#NEC042X050UC, Lot: 2389266, specific activity 275

mCi/ mmol) and [^{3}H]mannitol (#NET101001MC, Lot: 3632303, specific activity 12.3

Ci/ mmol) were purchased from Perkin Elmer. [O-methyl-^{3}H]amisulpride (MW 374.8

g/mol; specific activity 77 Ci/mmol; 97% radiochemical purity) was custom tritiated

(#TRQ41291 Quotient, UK). [^{14}C(U)]Sucrose (MW 359.48 g/mol; specific activity 536

mCi/mmol; 99% radiochemical purity: # MC266) was purchased from Moravek

Biochemicals, USA. Non-labelled amisulpride (MW 369.5 g/mol, >⬜98% purity) was

purchased from Cayman Chemicals, UK (#71675-85-9), and non-labelled D-glucose

(#10117) was purchased from BDH.

Literature review

141   Three different groups of molecules, which interacted with GLUT, were identified by

142   literature review. These groups were GLUT1 substrates, GLUT1 inhibitors and

143   GLUT-interacting antipsychotics. The inclusion criteria are explained below.

Identification of GLUT1 substrates and inhibitors

To identify established substrates and inhibitors of GLUT1 we performed a Pubmed

literature search using the parameters ((GLUT1 substrate) OR (GLUT1 inhibitor))

AND (review [Publication Type])) AND (("1985" [Date - Publication]:"2021" [Date -

Publication])) (29) and tabulated the results. In the substrates group we included

molecules for which there is *in vitro* evidence that they are transported by GLUT1 and

that their uptake into cells is inhibited by established GLUT1 inhibitors, such as

cytochalasin B. In the inhibitors group we included molecules that were shown to

decrease uptake of GLUT1 substrates *in vitro*, and for which there is a proposed

inhibitory mechanism of interaction with GLUT1.

Identification of antipsychotics that interact with GLUT

157   We also performed a Pubmed search to identify antipsychotics which interacted with

158   GLUT transporters, using the parameters (((GLUT) AND (antipsychotic)) AND

159 (("1985"[Date - Publication]: "2021"[Date - Publication])) (accessed 04/03/2022). In

160 the antipsychotics group we included both typical and atypical antipsychotics that

161 decreased the uptake of GLUT1 substrates *in vitro*.

Physicochemical characterisation of GLUT1 substrates and inhibitors, and GLUT-interacting antipsychotics

The physicochemical characteristics of each member of the three groups (i.e. GLUT1

substrates, GLUT1 inhibitors and GLUT-interacting antipsychotics) was obtained from

the chemical property databases, DrugBank (30) or MarvinSketch (version 22.9.0,

2022, ChemAxon) (31) and tabulated.

Specifically, information about the structure and molecular weight (MW) was obtained

from the DrugBank (30). The mean MW ± SEM, and the median MW of the three

groups was then calculated and the results compared.

The gross charge distribution at pH 7.4 of each molecule was then obtained from

MarvinSketch. The mean gross charge distribution at physiological pH (± SEM) of the

three groups was then calculated and the results compared. The number of

microspecies of each molecule at pH7.4 was examined and the percentage prevalence

and charge of the top two microspecies was then reported (MarvinSketch). The

physicochemical characteristics of the three groups were compared to those obtained

for amisulpride.

In silico molecular docking

In silico molecular docking was used to examine the molecular level interactions of amisulpride, alpha-D-glucose, beta-D-glucose, and sucrose with GLUT1 using the molecular docking tool GOLD. The main endogenous substrate of GLUT1 is D-glucose, which is a monosaccharide present in the body as anomers: 36% alpha-D-glucose and 64% beta-D-glucose (32). In this molecular docking study alpha-D-glucose and beta-D-glucose were used as positive controls. Sucrose is a plant disaccharide which is thought not to interact with GLUT1 and was used as the negative control.

The GLUT1 Protein Database (PDB) code used was 5EQI – the crystal structure of human GLUT1. The 5EQI crystal structure is for the inward open conformation of the transporter which has been reported to be the most favourable for ligand binding (33).

The results from the molecular docking simulations were expressed as a free energy binding and chem score. A molecule is considered a substrate for a given transporter if their interaction has a free energy binding of ≤5kcal/mol and has a high chem score. The chem score is one of the scoring functions of the molecular docking program. It provides information on the strength of the interaction between ligand and binding sites.

In vitro studies in model of the human BBB (hCMEC/D3 cells)

The human cerebral microvessel endothelial cells/D3 (hCMEC/D3) are an immortalised human adult brain endothelial cell line. This is a well-established and characterised model, regularly used to study the human BBB (16,34). The hCMEC/D3 cell line originated from human brain tissue obtained following surgical excision of an area from the temporal lobe of an adult female with epilepsy (34).

hCMEC/D3 cells were used to study the interaction between GLUT1 and amisulpride. To do this, the expression of a functional GLUT1 was first confirmed using Western blot (WB) and accumulation assays with [^{14}C]D-glucose. This was followed by further accumulation assays with [^{14}C]D-glucose and non-labelled amisulpride. [^{3}H]mannitol was used as a marker of cellular integrity and non-specific binding to the membranes and plastic wear. The hCMEC/D3 cells (passages 30-35) were grown as described previously by Sekhar et al., 2019 (16). All experiments were carried out at King's College London, in accordance with the guidelines of the Local Ethics Committee and research governance guidelines.

Western blot studies – expression of GLUT1 in hCMEC/D3 cells

Cells were grown for four to five days in T-75 flasks until they formed a fully confluent monolayer, and then for another two to three days before harvesting and preparing lysates for WB. Lysates were prepared by placing the flask on ice, aspirating the medium and washing the cells in ice-cold phosphate buffered saline (#70011-36 Gibco, Fischer Scientific Ltd, UK). The cells were then incubated on a shaker for 10 minutes in radio-immunoprecipitation assay (RIPA) buffer (#R0278, Sigma, UK) and 1% (v/v)

protease inhibitors (#78441, Thermo Fisher Scientific, UK) for cell lysis and protein

solubilisation. The cells were then scraped off from the bottom of the flask and

transferred to an Eppendorf tube. The tube was then agitated for 20 to 30 minutes at

4°C. Next, the cell suspension was centrifuged (Biofuge Fresco, Heraeus Instruments,

UK) for 20 minutes at 11,753 x G at 4°C. The resulting supernatant (protein lysate) was

snap frozen in liquid nitrogen. Before the start of the WB procedure, the protein lysates

were thawed and the total protein concentration in each lysate was estimated by

bicinchoninic acid (BCA) assay using bovine serum albumin standards (Thermo

Scientific, UK) as described by Sekhar et al., 2019 (16).

The protein lysates for WB were mixed with sample loading buffer (1:4) (#NP0007

NuPAGE LDS Sample Buffer (4x), Invitrogen, Carlsbad, USA), reducing agent (1:10)

(#B0009, Novex by Life Technologies, USA), and RIPA buffer. Once prepared, they

were heated at 95°C for 5 minutes. The rest of the WB procedure was performed as

described in (16) except, 30 µg of protein was loaded in each well, antibody solutions

were made in 5% milk in TBS-T, and washes were performed in TBS-T (for antibodies

dilutions see Supplementary Table 1, S1 File). Quantification of protein expression was

determined by calculating the intensity ratio of the band of interest and the band of the

loading control (GAPDH). Band intensity ratio analysis was conducted using ImageJ

software (35).

Accumulation assays – function of GLUT1 in hCMEC/D3 cells

The cell culture method and experimental design of the accumulation studies are

described in the following sections. The cells were grown as already described in the

"*In vitro* studies" section. They were split at 80-90% confluence and were seeded at

25,000 cells/cm^2 in the central 60 wells of 96 well plates for accumulation assay

experiments (#10212811, Fisher Scientific, UK). They were grown for four to five days

until they formed a fully confluent monolayer, and then for another two to three days

before the start of the accumulation assay.

For the control accumulation assay experiments, the hCMEC/D3 cells were incubated

in an accumulation buffer (135 mM NaCl, 10 mM HEPES, 5.4 mM KCl, 1.5 mM

$CaCl_2H_2O$, 1.2 mM $MgCl_2.H_2O$, 1.1 mM D-glucose - the standard non-labelled D-

glucose concentration for the accumulation buffer, and water, pH = 7.4) for 1 h. The

buffer also contained [^{14}C]D-glucose (3 µM) and [^{3}H]mannitol (0.05 µM).

[^{3}H]Mannitol was used as a marker for non-specific binding, and barrier integrity

(36,37). All treatment conditions contained 0.05% DMSO. Incubation was performed

in a shaker at 37°C and 120 rotations per minute. Next, the cells were washed with ice-

cold phosphate buffered saline (#BR0014G, Oxoid Limited, England) to stop the

accumulation process and to remove any radiolabelled molecules and buffer that had

not entered the cells. Then the cells were incubated in 1% Triton X (200 µl per well)

for 1 h at 37°C, to solubilise the cell membranes, and to free any accumulated

radiolabelled compounds via lysis. Half of the cell Triton X lysate from each well was

pipetted into a scintillation vial and 4 ml of scintillation fluid (#6013329, Ultima

GoldTM, Perkin Elmer, USA) was added. The amount of ^{14}C and ^{3}H radioactivity in

the sample was measured in disintegrations per minute (dpm) on a Tricarb 2900TR

liquid scintillation counter. The dpm of each sample were corrected for background

dpm. The background dpm was determined from a vial containing 100 µl 1% Triton X

and 4 ml liquid scintillation fluid only. The remaining cell lysate in the plate was used

for total protein concentration control in each well, determined by BCA analysis (17).

In our self-inhibition studies, investigating the presence of a functional GLUT1

transporter in the hCMEC/D3 cells, the cells were incubated with accumulation buffer,

containing [^{14}C]D-glucose (3 µM), [^{3}H]mannitol (0.05 µM) and 4 mM non-labelled D-

glucose. The 4 mM total concentration of non-labelled D-glucose included the standard

1.1mM non-labelled D-glucose normally present in accumulation buffer. Radioactivity

uptake was compared to the control condition in which the cells were treated with

accumulation buffer of the same composition apart from the non-labelled D-glucose

content which was 1.1 mM (the standard non-labelled D-glucose concentration in the

accumulation buffer). All treatments contained 0.05% DMSO. After 1 h incubation, the

cells were lysed and used for liquid scintillation counting and BCA assay as already

described. The non-labelled D-glucose concentration for the treatment condition (4

mM) was selected based on previous reports that normal plasma glucose concentrations

are kept in a narrow range between 4 and 8 mM in healthy subjects (38) and on

preliminary experiments studying the effect of 4, 6, and 9 mM non-labelled D-glucose

on the membrane integrity of hCMEC/D3 cells (see Supplementary Fig 2, S1 File).

In order to assess the effect of amisulpride on the transport of D-glucose through the

hCMEC/D3 membrane, non-labelled amisulpride (20, 50, or 100 µM) was added to the

accumulation buffer, containing [^{14}C]D-glucose (3 µM), [^{3}H]mannitol (0.05 µM), and

non-labelled D-glucose (1.1 mM - total standard concentration of non-labelled D-

glucose in the accumulation buffer). All treatment conditions contained 0.05% of

DMSO. The cells were incubated with this mixture for 1 h, the BCA assay and liquid

scintillation counting was performed as described previously.

The total cellular accumulation of [^{14}C]D-glucose was expressed as a volume of

distribution (V_d; µl/mg). V_d was calculated as the ratio of disintegrations per minute

(dpm)/mg protein in the Triton X lysate to dpm/ µl of the accumulation buffer. The V_d

values for [^{3}H]mannitol were subtracted from the V_d values for [^{14}C]D-glucose to

correct for cell membrane integrity. This value was then divided by the protein content

in each well to correct for protein content. Values were expressed as a mean ± SEM.

Statistical analysis of the results from the self-inhibition studies was performed using

unpaired one-tailed Student's t-test. The results from the studies of the interaction

between amisulpride and GLUT1 were analysed using One-way ANOVA. All

statistical analysis was performed with GraphPad Prism 9.0.

309   **Animal model studies in WT and 5xFamilial AD (FAD) mice**

The 5xFAD mice are a model of AD which carry human amyloid-β precursor

protein (APP) with the Swedish (K670N, M671L), Florida (I716V), and London

(V717I) FAD mutations along with human PSEN-1 with two FAD mutations: M146L

and L286V under the murine Thy-1 promotor (39). This model has been described as

one of the few models that shows several AD hallmarks, including neural loss,

neurodegeneration, gliosis, and spatial memory deficits (40).

We confirmed the model phenotype in the following ways: comparing the weight of

females and males from the two genotypes; using transmission electron microscopy

(TEM) to confirm the presence of amyloid plaques in the brain of 5xFAD mice;

comparing the expression of APP in brain capillary enriched pellets isolated from WT

and 5xFAD mice. Enrichment of the pellets with brain endothelial cells was confirmed

using transferrin receptor 1 (TfR1) as an endothelial cell marker.

Animal husbandry

All *in vivo* experiments were performed in accordance with the Animal Scientific

Procedures Act (1986) and Amendment Regulations 2012 and with consideration to the

Animal Research: Reporting of *In Vivo* Experiments (ARRIVE) guidelines. The study

was approved by the King's College London Animal Welfare and Ethical Review

Body. The UK government home office project license number was 70/7755.

The mice were housed at King's College London in groups of three or four under

standard conditions (20-22°C, 12 h light/dark cycle) with food and water *ad libitum*.

They were housed in guideline compliant cages. Animal welfare was assessed daily by

animal care technicians. Animals were identified by earmarks. The experimenter was

not blinded to the mouse genotype.

The WT (C57/BL6) mice and the 5xFAD mice (on C57/BL6 background) were

between 4.5 and 15 months old. The average lifespan of the C57/BL6 mice has been

reported to be 30 months (41). Whereas the 5xFAD mice have been reported to have

lifespan of approximately 15 months (42), with some authors reporting a median

lifespan of up to 24.6 months (41).

TEM ultrastructural study

To assess Aβ plaque presence in the brain in WT and 5xFAD mice we used TEM. All

studied animals were female. The animals of each genotype were grouped into two age

ranges. The groups were WT 4.5-6 months, 5xFAD 6 months, WT 12 months, 5xFAD

12 months with n=2 mice per group. The weight of the animals was between 21.9 g and

26.7 g. For brain dissection, all animals were terminally anaesthetised with

intraperitoneal injection of pentobarbital (100 µl/animal) (Fort Dodge, Southampton,

UK).

Once each mouse was anaesthetised, it was perfuse-fixed. The left ventricle of the heart

was infused with ice-cold phosphate buffered saline and the right atrium sectioned to

provide an open circuit. Once the vasculature was free of blood, 4% paraformaldehyde

(PFA, #F017/2, TAAB, UK) was infused via the heart. Fixation with 4% PFA was

performed for 10 minutes. The brain was then removed, and the frontal cortex was

dissected, and cut into 1 mm^3 samples.

The brain samples were processed, and sectioned for TEM, and then imaged. The

samples were incubated overnight at 4°C in TEM fixative, containing 2.5% (v/v)

glutaraldehyde in 0.1 M cacodylate buffer (pH 7.3). The tissues were post fixed in 1%

(w/v) osmium tetroxide (#O021, TAAB, UK) for 1.5 h at 4°C. Then they were washed

and dehydrated through serial graded incubations in ethanol: 10%, followed by 70%,

and 100%. The tissue was infiltrated in embedding resin medium (#T028, TAAB, UK)

for 4 h at room temperature. The samples were embedded on flat moulds and

polymerised at 70°C for 24 h. Ultrathin sections (100-120 nm) were cut on a Leica

ultramicrotome (Leica microsystems, Germany) and mounted on mesh copper grids.

They were contrasted with uranyl acetate for 2 minutes and with lead citrate for 1

minute. The sections were imaged on an EM-1400 (Plus) transmission microscope

operated at 120 kV (JEOL USA, Inc.).

For each frontal cortex sample, a minimum of three images were collected, and

amyloid plaques were identified and labelled with reference to electron microscopic

atlas of cells, tissues, and organs (43). The dimensions of each image were 3296x2472

pixels.

Western blot studies

We isolated the brain capillaries of each mouse used in the *in situ* brain perfusion

experiments. The mouse brain was *in situ* perfused via the heart with artificial plasma,

infused with rabiolabelled amisulpride and radiolabelled sucrose for 10 minutes. Then

it was homogenised in capillary depletion buffer (10.9 mM HEPES, 141 mM NaCl, 4

mM KCl, 2.8 mM $CaCl_2$ (aqueous solution - 1M), 1 mM $MgSO_4.7H_2O$, 1 mM

NaH_2PO_4, 10 mM glucose) (brain weight x 3) and 26% dextran (MW 504.4 g/mol)

(#J14495.A1, VWR, UK) (brain weight x 4). The homogenate was centrifuged at 5,400

G for 15 min at 4°C. This resulted in separation of an endothelial cell-enriched pellet

and a supernatant containing the brain parenchyma and interstitial fluid. Half of the

capillary pellets were snap frozen in liquid nitrogen and used for WB analysis to test

for TfR1, and BBB transporter expression, including GLUT1, and P-glycoprotein (P-

gp).

For WB, the mouse endothelial cell enriched capillary pellets were thawed,

homogenised in 250 µl RIPA buffer with added protease inhibitor (1% v/v). The tissue

was incubated in the buffer at 4°C for 30 minutes and then centrifuged (Biofuge

Fresco, Heraeus Instruments, UK) at 7,999 G for 15 minutes at 4°C. The resulting

supernatant was used for WB analysis. The rest of the WB procedure was performed as

previously described in this paper. For antibodies used, see Supplementary Table 1, S1

File.

Quantification of protein expression was determined by calculating the intensity ratio

of the band of interest and the band of the loading control (GAPDH or tubulin). Band

intensity ratio analysis was conducted using ImageJ software (35).

Unpaired two-tailed Student's t-test was used for statistical analysis of the difference in

expression of each transporter studied between the WT and the 5xFAD mice.

***In situ* brain perfusions**

The *in situ* brain perfusion technique allows examination of the movement of slowly

moving molecule across the BBB in the absence of systemic metabolism. This method

was used to compare [^{3}H]amisulpride and [^{14}C]sucrose uptake into the brain in WT and

in 5xFAD mice.

The mice were terminally anaesthetised with medetomidine hydrochloride (2 mg/kg,

Vetoquinol UK Limited) and ketamine (150 mg/kg, Pfizer, UK, and Chanelle, UK),

injected intraperitoneally. Heparin was injected intraperitoneally before the perfusion

(100 units heparin dissolved in 0.9 % NaCl (aqueous solution) (heparin – batch

number: PS40057; NaCl - Sigma Aldrich, Denmark). Two experimental groups were

perfused – WT (12-15 months old, n=7, n=4 females, n=3 males) and 5xFAD (12-15

months old, n=7, n=4 males, n=3 females). Weights of the perfused mice were between

16.8 g and 41.5 g. Animals with a weight lower than 25 g were excluded from the

perfusion analysis as perfusion at a flow rate of 5.5 ml/min could cause loss of BBB

integrity (44) but their capillary lysates were used in WB experiments.

Artificial plasma was used in the perfusion experiments. It contained 117 mM NaCl,

4.7 mM KCl, 2.46 mM MgSO$_4$.7H$_2$O, 24.8 mM NaHCO$_3$, 1.2 mM KH$_2$PO$_4$, 2.5 mM

CaCl$_2$ (aqueous solution - 1M), 39 g/L Dextran, 10 mM glucose, 1g/L bovine serum

albumin, and it was mixed with Evan's blue dye (0.0551 g per 1 L of artificial plasma,

#E2129-10G, Sigma Life Science, India). It was warmed to 37°C and oxygenated by

95% O$_2$/5%CO$_2$ gas bubbled through the solution. The artificial plasma also contained

[^{3}H]amisulpride (6.5 nM) and [^{14}C]sucrose (9.4 µM). Perfusion time was 10 minutes.

After the perfusion, the brain was dissected out and weighed. The frontal cortex,

striatum, thalamus, and hypothalamus were dissected under a microscope (Leica,

Wetzlar, Germany), weighed, and solubilised in Solvable (#6NE9100, PerkinElmer,

Inc., USA) for 2-3 days, then they were taken for liquid scintillation counting. These

regions were selected to compare with data from other *in situ* brain perfusion

experiments and observations from human brain data sets (16) focused on similar areas,

including the caudate nucleus and the putamen which are part of the striatum (45).

The rest of the brain was used for capillary depletion as already described. The whole

brain homogenate, supernatant (containing brain parenchyma and interstitial fluid), and

half of the capillary pellets were also solubilised and used for liquid scintillation

counting.

The concentration of [^{3}H] or [^{14}C] radioactivity present in the brain (disintegrations per

minute per gram of tissue – dpm/g) was expressed as a percentage of the concentration

of radioactivity detected in the artificial plasma (disintegration per minute per

millilitre). The value obtained was named %Uptake showing the radioactivity in ml/g

of tissue x 100. The %Uptake values for [^{3}H]amisulpride were corrected for vascular

space by subtracting the corresponding [^{14}C]sucrose Uptake values from the

[^{3}H]amisulpride Uptake values.

Statistical analysis of the difference between the uptake of [^{14}C]sucrose corrected

[^{3}H]amisulpride into the brain of WT and 5xFAD was performed using unpaired two-

tailed Student's t-test for the capillary pellet, supernatant, and homogenate . The

difference between WT and 5xFAD mice in the [^{3}H]amisulpride uptake in the brain

areas studied was analysed with Mixed-effects analysis with Holm-Sidak post hoc test.

The same tests were used to analyse the difference in the $[^{14}C]$sucrose brain uptake

between WT and 5xFAD mice.

Human control and AD tissue studies

452   We used TEM to examine the brain and brain capillary ultrastructure of an AD case.

453   Brain capillary depletion samples from age-matched human control and AD cases were

454   used to evaluate and compare the total protein levels and the expression levels of TfR1,

455   GLUT1 and P-gp in the two groups.

Ethics statement

Human tissue brain samples were provided via Brains for Dementia Research (BDR)

and were anonymised. Written consent was provided by BDR and the specific BDR

reference numbers were: TRID_170, TRID_170 amendment, TRID_265 and

TRID_287.

BDR has ethical approval granted by the National Health Service (NHS) health

research authority (NRES Committee London-City & East, UK: REC

reference:08/H0704/128+5. IRAS project ID:120436). Tissue samples were supplied

by The Manchester Brain Bank and the London Neurodegenerative Diseases Brain

Bank, both of which are part of the BDR programme, jointly funded by Alzheimer's

Research UK and Alzheimer's Society. Tissue was received on the basis that it will be

handled, stored, used, and disposed of within the terms of the Human Tissue Act 2004.

The human samples were collected during February 2012 to February 2019. The

human tissue studies were conducted during 1st October 2018 to 29th January 2021.

The authors conducting the human tissue analysis studies did not have access to

information that could reveal the identity of the human tissue donors.

TEM ultrastructural study

The tissue used was from one case which was received from the London

Neurodegenerative Diseases Brain Bank, Denmark Hill, King's College London. Case

details: BBN002.32856; sex: female; age: 74; post-mortem delay: 19 h; pathological

diagnosis: Alzheimer's disease, modified Braak staging (BrainNet Europe – BNE

staging) stage VI.

Samples with size of up to 1 x 5 x 1 mm were dissected from the frontal cortex, caudate

and putamen. Each sample was immersed in TEM fixative (2.5% glutaraldehyde in

0.1M cacodylate buffer) and incubated overnight at 4°C. The samples were then post-

fixed in 1% (w/v) osmium tetroxide (#O021, TAAB, UK) for 1.5 h at 4°C. Then they

were washed and dehydrated through serial graded incubations in ethanol - 10%

ethanol, followed by 70%, followed by 100%. The tissue was infiltrated in embedding

resin medium (#T028, TAAB, UK) for 4 h at room temperature. Next, the samples

were embedded on flat moulds and polymerised at 70°C for 24 h. Ultrathin sections

(100-120 nm) were cut on a Leica ultramicrotome (Leica microsystems, Germany) and

mounted on mesh copper grids. They were contrasted with uranyl acetate for 2 minutes

and with lead citrate for 1 minute. Finally, the sections were imaged on an EM-1400

(Plus) transmission microscope operated at 120 kV (JEOL USA, Inc.). For each section

of each brain region (frontal cortex, caudate, putamen), 5 to 10 pictures were examined

for pathological changes in the capillary and neurovascular unit. The original

dimensions of the images are 3296□x2472 pixels.

The presence of a single layer of endothelial cell surrounded by a layer of basement

membrane was considered to be an intact capillary. The observation of oedematous

space and vacuoles around the vessels, vacuolated pericytes, large vacuoles in the

endothelial cells or endoplasmic reticulum swelling, as well as oedema around the

capillary and multiple layers of basement membrane were considered as signs of

pathological changes (46). Neurodegeneration was identified by the presence of

lipofuscin granules, loss of myelin compactness, neurite degeneration and fibrillary

deposits (47,48).

Western blot studies

Post-mortem brain capillaries from neurologically healthy individuals and AD cases

were used to investigate the expression of transporters. The Braak stage of the control

cases was between I, and II (age-related pathology only). For the AD cases, the Braak

stage was between IV and VI. See Supplementary Information about the sex, age,

post-mortem delay (PMD), clinical diagnosis, and Braak stage of the individual cases

(Supplementary Table 2, S1 File).

Brain capillaries isolated from healthy individuals and AD patients were used to

investigate the expression levels of BBB transporters of interest by WB analysis. Brain

capillaries were isolated after homogenising brain tissue from the frontal cortex or the

caudate and carrying out a dextran-based density-gradient centrifugation to produce a

capillary enriched pellet. The capillary pellet was then homogenised in capillary

depletion buffer (brain weight x 3) and 26% dextran (brain weight x 4). The

homogenate was subjected to density gradient centrifugation (5,400 G for 15 minutes at

4°C) to give an endothelial cell-enriched pellet, the resulting supernatant was discarded

(49). The pellet was further lysed in 150-200 µl ice-cold RIPA buffer with added

protease inhibitors at 4°C, and then centrifuged at 8,000 G for 15 minutes at 4°C.

The protein concentration in each lysate was determined using a BCA assay and the

WB procedure was carried out as already described. For antibodies used, see

Supplementary Table 1, S1 File.

We used anti-TfR1 antibodies to detect TfR1 - an endothelial cell marker. This way we

aimed to confirm that the capillary pellets were enriched in endothelial cells. We also

used antibodies to detect GLUT1, and P-gp.

Quantification of protein expression was determined by calculating the intensity ratio

of the band of interest and the band of the loading control (GAPDH). Band intensity

ratio analysis was conducted using ImageJ software (35).

For statistical analysis, we used two-tailed unpaired Student's t-test to compare the

difference of transporter expression in the frontal cortex and the caudate between

control and AD cases.

Results

Physicochemical characteristics of amisulpride and glucose

We determined the physicochemical characteristics of amisulpride and D-glucose using

DrugBank and MarvinSketch. Amisulpride (chemical abstracts service (CAS) number

71675-85-9) has a MW of 369.48 g/mol, a pKa of 9.37 and exists as two microspecies

at physiological pH. The predominant (96.77%) microspecies is positively charged and

has a single positive charge at pH 7.4. The other microspecies (3.23%) has no charge

(Fig 1). The gross charge distribution at pH 7.4 of amisulpride is +0.968.

Fig 1. The percentage distribution and chemical structures of the two amisulpride

microspecies found at physiological pH. Microspecies A has a single positive charge

and Microspecies B has no charge, according to MarvinSketch 22.9.0.

D-glucose has a MW of 180.16 g/mol, and a pKa of 11.8. In solution, at equilibrium,

D-glucose exists as two anomers that interconvert spontaneously: ~36% alpha-D-

glucose, and ~64% beta-D-glucose, with less than 0.01% being present as an open-

chain form (linear glucose) (32,50). The major microspecies (99.99%) of both alpha-

D-glucose and beta-D-glucose has no charge (Supplementary Table 2, S1 File). The

other microspecies (0.01%) of each anomer has a single negative charge at

physiological pH. The linear form of D-glucose was found to be 100% neutral. Only

alpha- and beta-D-glucose have been included in the GLUT1 substrates group

(Supplementary Table 2, S1 File).

Identification of GLUT1 substrates and inhibitors

Our two PubMed searches identified 99 reviews, and 16 primary research articles, which discussed GLUT1 substrates, GLUT1 inhibitors or antipsychotics which interacted with GLUT transporters and matched our inclusion criteria. In these reviews and primary research articles, 9 GLUT1 substrates, 33 GLUT1 inhibitors, and 11 GLUT-interacting antipsychotics (including 6 typical antipsychotics and 5 atypical antipsychotics) were described and are listed in Supplementary Table 3-5, S1 File. They include the typical antipsychotics: chlorpromazine, fluphenazine, loxapine, pimozide and spiperone. Haloperidol is also included although some studies suggest no interaction with GLUT (23)(51). Atypical antipsychotics include: clozapine, desmethylclozapine, olanzapine, quetiapine and risperidone. Supplementary Table 6 (S1 File) provides specific details about the experimental evidence that indicates that these 11 antipsychotics interact with GLUT. In all cases this interaction was considered to be inhibitory.

Physicochemical characteristics of published substrates and inhibitors of GLUT1

Molecular Weight

The physiochemical characteristics of the identified GLUT1 substrates, GLUT1 inhibitors and the antipsychotics that interact with GLUT are summarised and compared to amisulpride (Fig 2, Table 1, and Supplementary Table 3-5, S1 File). The MW range for the GLUT1 substrates was 164.16 g/mol to 232.28 g/mol, with a

mean±SEM of 184.95±6.45 g/mol, the MW range of the GLUT1 inhibitors was 180.16

g/mol to 518.55 g/mol with a mean±SEM of 325.50±14.4 g/mol and the MW range for

GLUT interacting antipsychotics was 312.44 to 461.55 g/ mol with a mean±SEM of

369.38±16.04 g/mol (Table 1). One-way ANOVA showed differences in the MW F (2,

50) = 18.72, Tukey's multiple comparisons test showed significant difference between

the MW of GLUT1 substrates and inhibitors (p<0.0001), and between GLUT1

substrates and antipsychotics (p<0.0001) (Fig 2A). No significant difference was

observed between the MW of the GLUT1 inhibitors and the group of antipsychotics

(Fig 2A).

Fig 2. Physicochemical characteristics of GLUT1 substrates and inhibitors.

Evaluation of the PubMed database identified 9 substrates and 33 inhibitors of GLUT1,

and 11 antipsychotics inhibiting cell entry of GLUT1 substrates. A) Comparison of the

molecular weight (g/mol) of substrates, inhibitors of GLUT1, and antipsychotics. Each

square, dot, and triangle represents a compound. Comparison of the predicted gross

charge of substrates, inhibitors of GLUT1, and antipsychotics at pH=7.4, according to

MarvinSketch 22.9.0. Data was analysed using One-way ANOVA, GraphPad Prism 9.

B) The pie charts show the molecular weight of the substrates and inhibitors of

GLUT1, and of antipsychotics inhibiting cell entry of GLUT1 substrates. C) The pie

charts show the charge of the most prevalent microspecies of the GLUT1 substrates,

inhibitors, and antipsychotics interacting with GLUT at physiological pH according to

MarvinSketch 22.9.0.

The median for the GLUT1 substrates was 180.16 g/mol, for the GLUT1 inhibitors it

was 308.34 g/mol, for the antipsychotics it was 375.86 g/mol (Table 1). We found that

88.89% of the GLUT1 substrates were small molecules with a MW between 101 and

200 g/mol (Fig 2B). The GLUT1 inhibitors showed a wider range of molecular weight

with the majority falling (69.7 %) between 250 and 400 g/mol (Figure 2B).   The

majority (72.73%%) of the antipsychotics affecting GLUT1 substrates uptake were in

the range of 301-400 g/mol (Fig 2B).   There was no significant difference in the

molecular weight between the typical (386.18±23.42) and the atypical antipsychotics

(349.21±20.14) (t=1.169, df=9, p= 0.27, two-tailed unpaired t-test).

Table 1: Summary of the MW range, Mean±SEM, Median (g/mol), and gross charge at

	GLUT1 substrates n=9	GLUT1 inhibitors n=33	Antipsychotics n=11		
			All	Typical n=6	Atypical n=5
MW Range (g/mol)	164.16-232.28	180.16-518.55	312.44-461.55	318.86-461.55	312.44-410.49
Mean ± SEM (g/mol)	184.95±6.45	325.50±14.4	369.38±16.04	386.18±23.42	349.21±20.14
Median (g/mol)	180.16	308.34	375.86	385.76	326.82
Gross charge at physiological pH	+0.089 ± 0.09	-0.30 ±0.06	+0.89 ±0.02	+0.90 ±0.02	+0.87 ±0.05

physiological pH of GLUT1 substrates, inhibitors, and antipsychotics interacting with GLUT.

Charge

We investigated the predicted gross charge distribution at physiological pH of the

identified substrates and inhibitors of GLUT1 as well as the GLUT interacting

antipsychotics (Fig 2A), and also reported the predicted charge and the percentage

distribution of the top two microspecies (Supplementary Table3-5, S1 File).

Overall, we found that GLUT1 substrates, GLUT1 inhibitors and GLUT-interacting

antipsychotics had an average gross charge distribution at pH 7.4 of +0.089±0.09,

-0.30±0.06, and +0.89±0.02, respectively (Fig 2A, and Table 1). There was a

significant difference between all three groups (One-way ANOVA $F_{(2,50)}$ = 58.96,

p<0.0001, Tukey's multiple comparisons test GLUT1 substrates vs GLUT1 inhibitors

(p=0.0057); GLUT1 substrates vs antipsychotics p<0.0001; GLUT1 inhibitors vs

antipsychotics p<0.0001). There was no significant difference in the gross charge

between the typical (0.90±0.02) and the atypical (0.87±0.05) antipsychotics (t=0.678,

df=9, p= 0.5147, two-tailed unpaired t-test).

Importantly, there were two molecules which could be outliers within their groups.

This included the GLUT1 substrate, glucosamine, which had a much greater charge of

+0.827, and the atypical antipsychotic, quetiapine, which had a lower charge of +0.696

(Fig 2A).

GLUT1 substrates existed as either one, two or three microspecies at physiological pH.

GLUT1 inhibitors could exist as non-ionizable molecules (e.g. mercuric chloride) or as

up to 25 microspecies (e.g. morin) at physiological pH. The antipsychotics suggested

to interact with GLUT existed at physiological pH as either two, three or four

microspecies. This information plus the predicted charge of the top two microspecies

(if present) at physiological pH is tabulated in Supplementary Tables 3-5, S1 File. The

charge of the major microspecies of each group is presented in the form of pie charts

(Fig 2C). The major microspecies of GLUT1 substrates were neutral or had +1 charge,

GLUT inhibitors were neutral or had -1 charge and the anti-psychotics which interacted

with GLUT all had +1 charge at physiological pH.

In silico molecular docking

In silico molecular docking studies revealed that amisulpride could interact with

GLUT1 based on the low free energy binding of the interaction: -29.04 kcal/mol, and

the high chem score: 26.79. The molecular docking predicted conventional hydrogen

bonds between oxygen atoms from amisulpride and the amino acids: threonine (THR)

A: 137, and tryptophan (TRP) A: 412, and between hydrogen atom in the NH_2 group of

amisulpride and asparagine (ASN) A: 411. Alkyl interaction between amisulpride and

the amino isoleucine (ILE) A: 164, and pi-pi stacking interaction between amisulpride

and TRP A: 412 were also observed (Fig 3A).

Fig 3. Molecular level interactions of amisulpride, alpha-D-glucose, beta-D-

glucose, and sucrose with the binding sites of GLUT1. 2D (left) and 3D (right)

representations. In the 2D representations, green dotted lines are used to show

hydrogen bonds, pink dotted lines are used to depict hydrophobic interactions. A) Stick

representation is used for amisulpride, and line representation for the amino acid

residues from GLUT1. B) Stick representation is used for alpha-D-glucose, and line

representation for the amino acid residues from GLUT1. C) Stick representation is used

for beta-D-glucose, and line representation for the amino acid residues from GLUT1.

D) Stick representation is used for sucrose, and line representation for the amino acid

residues from GLUT1.

We also studied the interaction of GLUT1 with alpha-D-glucose and beta-D-glucose.

This interaction was used as a positive control. Molecular docking study found that

both alpha-D-glucose, and beta-D-glucose are substrates for GLUT1 with free binding

energy of -15.39 kcal/mol and chem score of 15.29. Conventional hydrogen bonds

between alpha-D-glucose and beta-D-glucose, and GLUT1 were observed at TRP A:

388, glutamine (GLN) A: 282, ASN A: 411, and THR A: 137 (Fig 3B and 3C). The

negative control sucrose, a disaccharide, showed higher free binding energy and lower

chem score than amisulpride, and alpha- and beta-D-glucose (monosaccharide). The

free energy binding for the interaction of sucrose with GLUT1 was -8.58 kcal/mol, and

the chem score was 6.55 (Fig 3D).

688   Expression and Function of GLUT1 in hCMEC/D3 cells

689   We examined the expression and function of GLUT1 in an established cell model of

690   the human BBB - the hCMEC/D3 immortalised cell line. Verification studies of the

691   experimental design were also performed.

692   Expression of GLUT1 in hCMEC/D3 cells

The hCMEC/D3 cells were found to express the GLUT1 transporter (40-60 kDa).

GAPDH was used as a loading control. Lysates from human colon adenocarcinoma

(Caco-2) cells were used as a positive control, and lysate from the human lung cancer

cell line (Calu-3) or human embryonic kidney cell line (HEK-293) were used as a

negative control (Fig 4A).

Fig 4. Function of GLUT1 in hCMEC/D3 cells – self inhibition. A) GLUT1

expression in hCMEC/D3 cells. Three passages of hCMEC/D3 cells (P30, 31, 33) (30

μg of protein per well) were tested for GLUT1 (40-60 kDa) expression. The figure is an

example membrane of three technical repeats. Caco-2 cell lysate was used as a positive

control; HEK-293 cell lysate was used as a negative control. GAPDH (37 kDa) was

used as a loading control. Antibodies used: anti-GLUT1 antibody – 1:100 000,

#ab115730; anti-GAPDH antibody – 1:2500, #ab9485, Abcam; secondary anti-rabbit

IgG, HRP-linked antibody – 1:2000, #7074, Cell Signalling Technology. B) V_d of

[^{3}H]mannitol was not significantly different between the control and the experimental

conditions. C) Non-labelled glucose decreased significantly the accumulation of

[^{14}C]D-glucose ([^{3}H]mannitol corrected) in hCMEC/D3 cells after 1 h of incubation.

Results are expressed as mean ± SEM, n=3-4 plates, passages (P33, 34, and 35) with

five well replicates per treatment in each plate. Data were analysed with an unpaired

one-tailed Student's t-test, using GraphPad Prism 9, each point represents a plate.

Accumulation assays – function of GLUT1 in hCMEC/D3 cells

To check that excess concentrations of D-glucose did not affect the membrane integrity

of the hCMEC/D3 cells, verification studies were performed. Cells were incubated

with [^{14}C]D-glucose and [^{3}H]mannitol as a control (containing the standard amount of

non-labelled D-glucose – 1.1 mM), and with [^{14}C]D-glucose, [^{3}H]mannitol, and non-

labelled D-glucose (4 mM) as a test condition. There was no significant change in the

permeability of [³H]mannitol (cell permeability marker) when the cells were treated

with 4 mM non-labelled D-glucose for 1 h (Fig 4B). Further analysis and studies

could therefore be performed using similar D-glucose concentrations.

To investigate the function of GLUT1 in hCMEC/D3 cells, we performed a self-

inhibition experiment. Incubation of the cells with 4 mM non-labelled D-glucose for 1

h led to a significant decrease in the V_d of [¹⁴C]D-glucose ([³H]mannitol and protein

corrected) (from 97.5 ± 22.3 µl/mg to 37.3 ±10.8 µl/mg; t=2.161, df=5, p= 0.0415,

unpaired one-tailed Student's t-test, data is presented as mean ± SEM) (Fig 4C).

When we tested the interaction of amisulpride with GLUT1 in hCMEC/D3 cells, the

entry of [³H]mannitol (0.05 µM) into the cells was used as a cell integrity marker.

There was no significant change in the [³H]mannitol permeability when the cells were

treated with amisulpride (20, 50, or 100 µM) (Fig 5A). There was also no significant

effect on the V_d of [¹⁴C]D-glucose when the cells were treated with amisulpride (20, 50,

or 100 µM) (Fig 5B).

Fig 5. Interaction of amisulpride with GLUT1 in hCMEC/D3 cells. A) Amisulpride

at three different concentrations (20, 50, or 100 µM) did not have a significant effect on

the accumulation of [³H]mannitol in hCMEC/D3 cells after 1 h of incubation. B)

Amisulpride at three different concentrations (20, 50, or 100 µM) did not have a

significant effect on the accumulation of [¹⁴C]glucose ([³H]mannitol corrected) in

hCMEC/D3 cells after 1 h of incubation. Results are expressed as mean ± SEM, n=3-4

plates, passages (P33, 34, and 35) with five well replicates per treatment in each plate.

Data were analysed with a One-way ANOVA, using GraphPad Prism 9, each point

represents a plate.

Animal model of AD - validation

Mouse weight comparison

We compared the weight of the WT and 5xFAD mice and of the two sexes (age

between 12 and 15 months). All data are presented as mean ± SEM. The WT mice had

higher average weight than the 5xFAD mice (32.31 ± 2.93 g vs 26.15 ± 2.5 g), and all

the 5xFAD mice with weight lower than 25 g were female. We observed that the

females had significantly lower weight (g) than the males in the 5xFAD group (19.7 ±

1.46 vs 31 ± 1.61; t=4.983, df=5, p=0.0042; unpaired two-tailed Student's t-test). In the

WT group, weight difference between the sexes was smaller and did not reach

significance (Supplementary Fig 3, S1 File).

TEM ultrastructural study

We examined frontal cortex samples of WT and 5xFAD mice for the presence of Aβ

plaques using TEM. Structures were identified as Aβ plaques by comparison to Aβ

plaques which had previously been identified in TEM images (40). Amyloid plaques

were not observed in the frontal cortex of the young WT mice (4.5-6 months; weight

25.4 g, 26.1 g). Importantly, structures identical to Aβ plaques previously reported in

the literature (40) were present in the 5xFAD mice in both the young (6 months; weight

23.8 g, 23.5g) and the old (12 months 21.9 g, 22 g, data not shown) group (Fig 6).

Fig 6. TEM ultrastructural study in WT and 5xFAD mice. TEM image of the

frontal cortex of A) WT mouse, which is free of Aβ plaques, and B) 5xFAD mouse,

with Aβ plaques. Mouse age – 4.5 – 6 months. Magnification – 1200x, scale bar – 10

μm, n=2 mice per group.

Western blot studies

We performed WB for human APP to examine the genotype of each mouse used in this

study. APP is the precursor protein from which amyloid-β (Aβ) is cleaved by β-

secretase and γ-secretase (52). Our studies confirmed human APP expression was not

detectable in the WT mice, but was detectable in the 5xFAD mice cerebral capillaries

(See Supplementary Fig 4, S1 File)

We observed TfR1 and transporter expression in the mouse brain endothelial cell

lysates from WT and 5xFAD mice used in the *in situ* brain perfusions (See

Supplementary Fig 5, S1 File). Thus, these experiments confirmed we have

successfully isolated the BBB compartment from the whole brain.

In particular, brain capillaries from WT and 5xFAD mice were found to express

GLUT1, PMAT, multi-drug and toxin extrusion protein 1 (MATE1), OCT1, and P-gp.

No significant effect of the genotype was found on the expression of these proteins

(See Supplementary Figs 5-9, S1 File).

In situ brain perfusions

[^{3}H]amisulpride uptake (uncorrected for [^{14}C]sucrose) was significantly greater than

[^{14}C]sucrose uptake in all brain areas studied apart from the hypothalamus, thalamus,

and capillary pellet in the WT mice (paired two-tailed t-test, data not shown). In the

5xFAD mice the [^{3}H]amisulpride uptake (uncorrected for [^{14}C]sucrose) was

significantly greater than [^{14}C]sucrose uptake in all brain areas studied (paired two-

tailed t-test, data not shown).

When comparing WT and 5xFAD mice, there was a trend for an increase in

[^{3}H]amisulpride uptake into specific brain regions, whole brain homogenate, and

capillary pellet in the 5xFAD mice but this failed to reach statistical significance (Fig

7A and 7B). However, there was a statistically significant increase in the uptake of

radiolabelled amisulpride in the supernatant (made of brain parenchyma and interstitial

fluid) of 5xFAD mice compared to WT mice (t=2.550, df=8, p=0.0342, unpaired two-

tailed t-test, by 104.86%). These results were [^{14}C]sucrose corrected. None of the

regions showed a significant difference in the [^{14}C]sucrose uptake between the WT and

the 5xFAD mice (Fig 7A and 7B).

Fig 7. [^{3}H]Amisulpride (sucrose corrected) and [^{14}C]sucrose uptake into the brain

of WT and 5xFAD mice. WT and 5xFAD mice were perfused with [^{3}H]amisulpride

and [^{14}C]sucrose. Perfusion time - 10 minutes, fluid flow rate – 5.5 ml/min. Mouse age:

12-15 months. WT n=6, 5xFAD n=4, apart for the hypothalamus where WT n=5,

5xFAD n=4; the capillary pellet where WT n=3, 5xFAD n=3, and the homogenate

where WT n=6, 5xFAD n=3. No significant differences in paracellular permeability

and membrane integrity were observed in any region. Each dot represents data from

one mouse. All data are expressed as mean ± SEM, Mixed effects analysis and

unpaired two-tailed Student's t-test, GraphPad Prism 9.

Human control and AD tissue studies

TEM ultrastructural study

For each brain region (frontal cortex, caudate, putamen), 10 to 20 pictures were examined, and the endothelial cells, basement membranes, capillary lumens, axons, myelin sheets and markers of degeneration were identified and labelled with reference to Electron Microscopic Atlas of cells, tissues and organs (43). We observed thickened basement membrane, vacuolisation of the endothelial cell, fibrillary deposits, and oedema around brain capillaries in the frontal cortex, caudate and the putamen (Fig 8). Various other features of degeneration were observed in our AD case (BNE stage VI), including: lipofuscin granules in the frontal cortex and the putamen, and myelin degeneration in the caudate (Fig 9).

Fig 8. TEM image of capillaries in the frontal cortex (A) putamen (B) and caudate (C) of a human AD case. Frontal cortex and caudate magnification – 3000x scale bar – 2 µm. Putamen magnification – 2000x, scale bar – 5 µm. n=1 case, two sections per brain area were stained and imaged, 5 to 10 images were examined per section. Case number: BBN002.32856; Sex: F; Age: 74; PM delay: 19 h; Alzheimer's disease, BNE stage VI.

Fig 9. TEM image of the A) frontal cortex (A and C), putamen (B), and caudate (D) of human AD case.

A) TEM image of the frontal cortex of human AD case.

B) TEM image of the putamen of human AD case.

A and B showing electron dense and lucid portions of lipofuscin granules.

Magnification – 1500x, scale bar – 5 µm.

C) TEM image of degenerating neurites and degenerating myelinated axon in the

frontal cortex of a human AD case. Magnification – 6000x, scale bar – 2 µm.

D) TEM image of myelin and axon degeneration in the caudate of a human AD case.

Magnification – 2500x, scale bar – 5 µm.

Case, n=1, two sections per brain area were stained and imaged, 5 to 10 images were

examined per section. Case number: BBN002.32856; Sex: F; Age: 74; PM delay: 19 h;

Alzheimer's disease, BNE stage VI.

Total protein concentration

A BCA assay was used to compare the total protein concentration in the frontal cortex

and caudate capillary lysates of control vs AD cases. For the frontal cortex control

cases n=9; AD cases n=9) and caudate (control cases n=9; AD cases n=5), an unpaired

two-tailed t-tests showed no significant difference between AD and control

(Supplementary Fig 10, S1 File).

Western blot studies

TfR1 was detected in the capillary protein lysates from the frontal cortex and caudate

of human brains both in the control and in AD cases. There was no significant

difference between the two groups in the frontal cortex or the caudate (unpaired two-

tailed t-test). The expression of TfR1 in the samples confirms they are endothelial cell

enriched (Fig 10).

Fig 10: TfR1, GLUT1 and P-gp expression in human frontal cortex and caudate

brain capillary lysates. Comparing control and AD expression. Intensity ratio was

calculated using the intensity of the band of TfR1, GLUT1 or P-gp, and controlling it

for the intensity of the loading control GAPDH. Results are presented as mean±SEM,

each dot represents a case, data was analysed using unpaired two-tailed Student's t-test.

All analysis was performed using Image J, Excel, and GraphPad Prism 9.

We studied the expression of GLUT1 in control and AD age-matched cases. We did

not see a significant change in the expression of GLUT1 between control and AD in the

frontal cortex or the caudate (Fig 10).

We also studied the expression of P-gp in control and AD age-matched cases. The

brain areas we examined were the frontal cortex and the caudate. We observed a

decrease in P-gp expression in the AD group comparted to control, both in the frontal

cortex and the caudate. In the caudate this decrease was significant (t=2.841, df=16,

p=0.0118), unpaired two-tailed Student's t-test (Fig 10; Supplementary Figs 11-13, S1

File).

Discussion

This study investigated the role of GLUT1 in the transport of the antipsychotic,

amisulpride at the BBB, to understand better its role in the hypersensitivity of AD

patients to the side effects of antipsychotics compared to healthy aged patients (4). We

used an integrative approach to test the hypotheses that amisulpride interacts with

GLUT1 at the BBB, and that expression of BBB transporters and the transport of

amisulpride and glucose into the brain is affected by AD. Analysis of the published

literature allowed us to identify three groups of molecules: GLUT1 substrates, GLUT1

inhibitors and antipsychotics that interacted with GLUT.   We then utilized specialist

chemical property databases to obtain a clearer picture of the physicochemical

characteristics of these molecules and their groups as well as amisulpride. *In silico*

docking studies allowed us to explore the specific molecular interactions of amisulpride

with GLUT1. WB and *in vitro* accumulation assays in hCMEC/D3 cells were used to

confirm the expression of GLUT1 protein in these BBB cells, the presence of

functional GLUT1 transporter, and to determine possible interactions between GLUT1

and amisulpride. We also examined the neurovascular unit architecture in WT and

5xFAD mice at an ultrastructural level using TEM and assessed the suitability of the

5xFAD mouse model to study the BBB permeability of amisulpride in AD. WB was

used to study the expression of SLC and ABC transporters, including GLUT1 at the

BBB in WT and 5xFAD mice. The *in situ* brain perfusion technique allowed us to

compare amisulpride uptake into compartments of the brain in WT and 5xFAD mice.

We also compared transporter expression in capillaries from human cases with and

without Alzheimer's dementia. TEM was utilised to directly visualise cellular

structures in an individual with AD.

Physicochemical characteristics of GLUT1 substrates and

inhibitors

Amisulpride is an atypical antipsychotic and has a MW of 369.48 g/mol. The main

microspecies (96.77%) at physiological pH has one positive charge. The other

microspecies (3.23%) of amisulpride at physiological pH has no charge. The gross

charge distribution at pH 7.4 of amisulpride is +0.968.

Our literature review and analysis of the chemical property databases revealed that

GLUT1 substrates are typically neutral, however, we have identified one substrate, D-

glucosamine, which has a positive gross charge distribution physiological pH of

+0.827. GLUT1 inhibitors are predominately neutral, but we identified inhibitors which

had a negative gross charge distribution at pH 7.4 (for example, -1.043 - Lavendustin

B).

Importantly, both typical and atypical antipsychotics were identified which impede

uptake of GLUT1 substrates *in vitro*. They all have a gross charge distribution which is

positive at physiological pH and their main microspecies at physiological pH has a

single positive charge, similar to amisulpride. An example is olanzapine for which

there is also *in silico* molecular docking data for an inhibitory interaction with a

bacterial glucose/H^+ symporter from *Staphylococcus epidermidis*. It can impede the

alternating opening and closing of the substrate cavity necessary for glucose transport

(53). Another example is risperidone (24), which has been reported to interact with

GLUT, and to inhibit glucose uptake *in vitro* (in rat PC12 cells). Thus, it is plausible

that other atypical antipsychotics such as amisulpride could interact with GLUT1.

Although it is important to consider that not all antipsychotics interact with GLUT. For

example, sulpiride (CAS 23756-79-8; MW 341.43, gross charge at physiological pH

+0.972) and clozapine N-oxide (a clozapine metabolite: CAS 34233-69-7; MW 342.83,

gross charge at physiological pH +1.616) were reported to not have a significant effect

on glucose uptake *in vitro* (23,51).

When it comes to MW, all the established substrates of GLUT1 are smaller than

amisulpride – 164.16 to 232.28 g/mol (MW of the main substrate - glucose is 180.16

g/mol). Amisulpride, has a MW similar to the established GLUT1 inhibitors and to

the antipsychotics reported to inhibit uptake of GLUT1 substrates. Amisulpride is

therefore more likely to represent a GLUT1 inhibitor than substrate. However, this

interpretation is limited by the small number of established GLUT1 substrates, even

though a large number of reviews (99) were examined.

First generation antipsychotics are known to inhibit dopaminergic neurotransmission

most effectively by blocking about 72% of the D2 dopamine receptors in the brain.

They also block noradrenergic, cholinergic, and histaminergic receptors. Whereas,

second generation antipsychotics block D2 dopamine receptors and serotonin receptors

(5-HT), mainly the 5-HT2A subtype (54).

Interestingly, amisulpride (second generation antipsychotic) shows high and similar

affinities for the D2 and D3 dopamine receptor subtypes but it does not have significant

affinity to the other receptor subtypes (55).

Although atypical antipsychotics are usually considered to have a broader mechanism

of action compared to typical antipsychotics, there were no significant differences in

physicochemical characteristics of the two groups or in their reported type of

interaction with GLUT1.

955   *In silico* studies

In silico, GLUT1 was found to interact with amisulpride with a free energy binding of -

29.04 kcal/mol (the positive control, beta-D-glucose, showed free energy binding of -

15.39 kcal/mol, and the negative control, sucrose, showed much higher free energy

binding of -8.58 kcal/mol). Previous studies have shown that interactions with TRP412,

TRP388, phenylalanine (PHE) 291, PHE379 and glutamate (GLU) 380 may play

critical roles in ligand binding to GLUT1 in the inward open conformation (33). Also,

another study reported interactions between cytochalasin B, a competitive inhibitor of

glucose exit via GLUT1 (56), and THR137, ASN411, GLN282, ASN288, glycine

(GLY) 384 and TRP388 in GLUT1 (57). In addition, ASN411 and TRP412 were found

to be important for the binding of inhibitors in the central binding site of GLUT1 (57).

Importantly, we see interactions between amisulpride and GLUT1 at some of the

reported amino acid residues – TRP412, THR137 and ASN411. The interaction of

amisulpride with these specific amino acid residues in GLUT1, taken together with the

size of amisulpride compared to GLUT1 substrates and inhibitors, and its positive

charge, suggests amisulpride might competitively inhibit glucose delivery to the CNS

via GLUT1.

Dysfunction of GLUT1 is likely to cause further dysregulation of the neurovascular

unit (NVU), resulting in the loss of BBB integrity as well as the observed altered

transporter expression and increased Aβ toxicity in AD. Interestingly, increased

neuronal uptake of glucose has been shown to protect against Aβ toxicity, and altering

glucose delivery to the brain could influence progression of AD pathology (58). A

situation that may be exacerbated by the additional medications AD patients usually

receive (59). Antipsychotics are commonly prescribed with antidepressant or sedative

drugs (60), i.e., citalopram (2) (MW: 324.4; +1 charge at pH 7.4) (30). It can be

assumed that if two of these psychotropics are substrates (or inhibitors) for the same

transporter and are prescribed together then this is likely to change individual drug

delivery and potentially nutrient delivery to the CNS and contribute to the drug

hypersensitivity. In fact, polypharmacy is an important predictor of adverse drug

reactions for people with and without dementia (59).

In vitro **studies in model of the human BBB (hCMEC/D3**

cells)

We confirmed the expression of GLUT1 in the hCMEC/D3 cells with WBs. These

results are in line with previous studies. GLUT1 protein expression has been detected

in hCMEC/D3 cells and in brain microvascular endothelial cells derived from healthy

patient-derived induced pluripotent stem cells (61,62). Other glucose transporters'

mRNA has also been detected in the hCMEC/D3 cells, namely GLUT3 and sodium-

dependent glucose transporter 1 (SGLT1) (62,63). Protein expression of GLUT3 and

GLUT4, but not SGLT1, has also been reported in hCMEC/D3 cells (61). Nevertheless,

there is consensus that GLUT1 is the most highly expressed transporter in the

hCMEC/D3 cells and the BBB (64).

In vitro, our self-inhibition study using 4 mM non-labelled glucose, showed that 4 mM

glucose significantly decreased the uptake of [^{14}C]D-glucose into the hCMEC/D3 cells

but did not affect membrane integrity as measured with [^{3}H]mannitol. This suggests the

presence of functional glucose transporter on the hCMEC/D3 cells.

1003

Our *in vitro* experiments looking at the interaction between antipsychotics and GLUT1

showed no effect of micromolar concentrations of amisulpride on the accumulation of

[^{14}C]D-glucose into the hCMEC/D3 cells. This suggests that clinically relevant

concentrations of amisulpride do not inhibit GLUT1 (amisulpride has a plasma Cmax

of 0.17-1.19 μM) (4,65). However, these results are difficult to interpret conclusively

as the GLUT1 is partially saturated due to the presence of non-labelled glucose in the

control accumulation assay buffer. The non-labelled glucose being essential for

endothelial cell survival.

1012

Overall, the high affinity (Km = ~2mM (66)) of the GLUT1 transporter, high

expression of GLUT1 at the BBB, excess glucose concentrations in buffer/plasma and

the low concentrations of amisulpride could make the interaction between amisulpride

and GLUT1 difficult to detect *in vitro*. Amisulpride has been associated with

hyperglycemia as a side effect in 1 in 100 people (67). The hyperglycaemia possibly

being caused by inhibition of GLUT1 transport. Interestingly, recent studies did not

associate its use with a higher prevalence of diabetes (68)

1020

1021 Cell culture studies have shown that another second-generation antipsychotic -

1022 clozapine, inhibits glucose uptake (at 20 μM) in PC12 rat cells after a 30-minute

1023 incubation. Exposure to clozapine beyond 24 h (at concentrations up to 20 μM), there

1024 was a significant increase in the cellular expression of GLUT1 and GLUT3 (23).

1025

Later studies have confirmed that both clozapine and risperidone interact with glucose

transporters and inhibit glucose transport. It has been suggested that this could be

mediated by directly binding to glucose transporters and allosterically modulating

either a glucose binding side or the conformational change in the protein conformation

required for transport (24). Importantly, risperidone which is the only antipsychotics

licensed for use in AD, shows treatment response and emergent side effects at very low

doses and low plasma concentrations (7), similar to amisulpride (10). Also, clozapine is

used at low doses and at correspondingly low plasma concentrations in the treatment of

psychosis in Parkinson's disease patients (69).

1035

Animal model studies in WT and 5xFamilial AD (FAD) mice

In order to study the effect of AD on the BBB and how this contributes to the increased

sensitivity of AD patients to antipsychotics, we need good preclinical models. In this

study we utilized the 5xFAD mice. The significantly lower weight of the female mice,

compared to the male mice of the 5xFAD genotype which we observed, matches with

previous reports that female mice develop the pathology earlier than male mice (70).

This could be related to increased expression of the Thy1 promoter which drives the

transgenes in the 5xFAD mice and has an oestrogen response element (71) resulting in

generation of higher levels of Aβ (72).

1045

1046   **TEM ultrastructural study**

1047   Using TEM, we detected Aβ plaques in the frontal cortex of 5xFAD mice from 6

1048   months of age. This is in agreement with previous data, reporting amyloid deposition in

1049   these mice from the age of 2 months (40) and with our Western blot studies which

1050   confirmed APP expression.

1051

In situ **brain perfusions**

When we studied the permeability of [^{3}H]amisulpride across the BBB in a whole

animal *in vivo*, we observed low permeability of the drug across the BBB in WT and in

5xFAD mice (both groups 12-15 months of age).

Interestingly, as measured by [^{14}C]sucrose (MW 359.48 g/mol), there was no

significant change in vascular integrity between age-matched WT and 5xFAD mice in

all areas that we investigated even though there was a trend for increased permeability

in the 5xFAD mice in the frontal cortex, thalamus, supernatant, and homogenate.

Importantly, there was significantly higher uptake of [^{3}H]amisulpride (corrected for

[^{14}C]sucrose) in the supernatant of 5xFAD mice compared to WT mice. An earlier

animal study has also reported difference in the brain uptake of amisulpride in an AD

mouse model compared to WT, without significant change in the [^{14}C]sucrose uptake,

i.e. without significant change in BBB integrity (16). The 3xTg mouse model of AD

had increased [^{3}H]amisulpride uptake in the frontal cortex, but not the occipital cortex,

compared to WT mice at 24 months of age (age matched) (16). Thus, suggesting that

the increased uptake of amisulpride in the brain of the 5xFAD mice could be related to

regional BBB changes in transporter expression associated with AD.

Western blot studies

A decrease in GLUT1 expression in brain samples has been observed previously using

WB in 5xFAD mice compared to WT at 9 months of age (73). This study used whole

brain samples and pooled the samples from all of their WT mice together and from all

of their 5xFAD mice together (73). In our study we confirmed the expression of

GLUT1 in 5xFAD and WT mouse capillaries at 12-15 months of age. We did not

observe significant difference between the genotypes, this is likely related to the fact

that we did not pool our samples but analysed each mouse separately which would have

increased variability.

1078

Other studies using the 5xFAD mouse model have reported a decrease in the

expression of P-gp and GLUT1 in the cortex capillaries of 6-month-old 5xFAD mice,

compared to age-matched WT mice (74). We did not detect any effect of the genotype;

this could be because we used the whole mouse brain to prepare the capillary samples

and to perform WB, and the expression changes could be region-specific.

Finally, expression of PMAT, MATE1, OCT1, and P-gp has been reported in the BBB

of 3xTg mice. No difference in expression levels was observed when they were

compared to age matched WT mice (16). These results are in line with our observations

in the 5xFAD model. This could be because both studies used the whole brain to do

capillary isolation, whereas the AD-associated transporter expression changes might be

regional, as seen in human control and AD cases (16).

1090

Human control and AD tissue studies

TEM ultrastructural study

In our AD case we were able to identify brain capillaries, myelinated axons and

neurites using TEM. Our images showed signs of NVU, and brain degeneration such as

swollen basement membrane and oedema around the capillaries, lipofuscin granules,

degenerating neurites, degenerating myelin, and fibrillary depositions.

1097

Previous EM studies have also reported abnormalities in the brain capillaries,

associated with AD, these include, splitting and duplication of the basement membrane,

reduction of the length of the tight junctions, morphological alterations of the

mitochondria of the endothelial cells, the pericytes and the perivascular astrocytic

processes. The number of the pinocytic vesicles was substantially increased in the

endothelium of the brain capillaries in AD in comparison with age-matched controls.

Thus, it has been suggested that abnormalities in the brain capillaries may result in the

release of neurotoxic factors and abnormal Aβ homeostasis in the brain and contribute

to AD pathology (75).

1107

1108   **Western blot studies**

1109   Expression of the endothelial marker TfR1 in our human brain capillary lysates

1110   confirms that we have isolated brain capillary pellets enriched in endothelial cells.

1111   There was no significant change in TfR1 expression between control and AD samples

1112   observed.

1113

Diminished glucose uptake has been reported in the hippocampus, parietotemporal

cortex and/or posterior cingulate cortex in individuals at genetic risk for AD (76),

positive family history (77) and/or mild or no cognitive impairment who develop AD

(78,79). Reduced levels of GLUT1 in cerebral microvessels have also been reported in

AD in the caudate nucleus (80), frontal cortex (protein was decreased but not mRNA)

(19), and the hippocampus (18). It has been suggested that GLUT1 deficiency can

contribute to the disease process, acting in tandem with Aβ to initiate or amplify

vascular damage and Aβ accumulation (81). We did not observe a significant change in

GLUT1 expression in the frontal cortex and caudate between controls and AD patients.

This could be related to the insufficient selectivity of the antibody, the need of more

cases, or the heterogeneity in the group of cases.

1125

P-gp is ubiquitously and abundantly expressed in the brain capillaries (82).

Importantly, protein expression has been reported to decrease significantly in the

prefrontal cortex of AD patients, compared to healthy ageing controls (61 to 100 years

old) (83). In line with other studies, we observed a decrease in the P-gp expression in

the caudate capillaries of AD patients, compared to healthy controls. Importantly, P-gp

protects the brain from potentially toxic substances and has been reported to extrude

Aβ from the brain (84). It has been suggested that downregulation of P-gp could allow

pharmaceuticals into the central nervous system and may increase the accumulation of

Aβ (83).

1135

Conclusion

1136

In conclusion, a literature review identified GLUT1 substrates, GLUT1 inhibitors and

antipsychotics that interacted with GLUT. Physicochemical characterization of these

groups using chemical property databases established that amisulpride had similar

properties to the GLUT-interacting antipsychotics group.   Our *in silico* molecular

docking studies revealed that amisulpride interacts with GLUT1 and could potentially

affect glucose delivery to the CNS. We also have *in vitro* evidence for the presence of

functional glucose transporter in the hCMEC/D3 cells line.   However, we could not

detect any interaction of amisulpride with GLUT1 in this assay.   This is possibly

because glucose being essential for endothelial cell survival limits the sensitivity of this

assay for exploring antipsychotic interaction with GLUT1 *in vitro*.   TEM and WB

analysis validated the 5xFAD mouse model for our study. The *in situ* brain perfusion

studies showed limited entry of amisulpride across the BBB in both WT and 5xFAD

mice, and an increased uptake into the brain of the 5xFAD mice compared to WT mice,

although the cerebrovascular space was similar in both genotypes. Our WB work with

P-gp further confirms that transporter expression at the human BBB is altered in AD,

although a significant difference was not observed for GLUT1 expression in our cases.

It is possible that amisulpride competitively inhibits glucose entry via GLUT1, which

may further compromise the neurovascular unit and increase BBB permeability, and

therefore increase central drug access, and contribute to the amisulpride sensitivity

observed in AD. This research further confirms the national guidance that antipsychotic

drugs should only be prescribed at the lowest dose possible for the shortest durations in

AD. It is also plausible that, in the longer term, the impact on energy delivery to the

brain may lead to further cellular degeneration. The implications of our findings extend

to other antipsychotic drugs.

1161

List of abbreviations

3xTgAD: triple transgenic, Aβ: amyloid beta, AD model; 5xFAD: five times familial

Alzheimer's disease mouse model; ABC: Adenosine triphosphate binding cassette;

AD: Alzheimer's disease; APP: amyloid precursor protein; ASN: asparagine; BBB:

blood–brain barrier; BCA: bicinchoninic acid; BDR: Brain for Dementia Research;

dpm: disintegrations per minute; CAS: chemical abstracts service number; FBS: foetal

bovine serum; GLN: glutamine; GLU: glutamate; GLUT1: glucose transporter 1; GLY:

glycine; hCMEC/D3: immortalized human cerebral microvessel endothelial cell line;

HEK cells: human embryonic kidney cells; MATEs: multi-drug and toxin extrusion

proteins; MW: molecular weight; OCT: organic cation transporters; NVU:

neurovascular unit; PFA: paraformaldehyde; P-gp: P-glycoprotein; PHE:

phenylalanine; PMAT: plasma membrane monoamine transporter; PMD: post-mortem

delay; RIPA: radio-immunoprecipitation assay; SGLT1: sodium-dependent glucose

transporter 1; SLC: solute carrier; TEM: transmission electron microscopy; TfR1:

transferrin receptor 1; THR: threonine; TRP: tryptophan; V_d: volume of distribution;

WB: Western blot; WT: wild type.

1178

Declarations

Ethics approval and consent to practice

All *in vivo* animal experiments were performed in accordance with the Animal

Scientific Procedures Act (1986) and Amendment Regulations 2012 and with

consideration to the Animal Research: Reporting of *In Vivo* Experiments (ARRIVE)

guidelines. The study was approved by the King's College London Animal Welfare

and Ethical Review Body. The UK government home office project license number

was 70/7755.

1187

Human tissue brain samples were provided via Brains for Dementia Research (BDR)

and were anonymised. Written consent was provided by BDR and the specific BDR

reference numbers were: TRID_170, TRID_170 amendment, TRID_265 and

TRID_287.

BDR has ethical approval granted by the National Health Service (NHS) health

research authority (NRES Committee London-City & East, UK: REC

reference:08/H0704/128+5. IRAS project ID:120436). Tissue samples were supplied

by The Manchester Brain Bank and the London Neurodegenerative Diseases Brain

Bank, which are both part of the BDR programme, jointly funded by Alzheimer's

Research UK and Alzheimer's Society. Tissue was received on the basis that it will be

handled, stored, used, and disposed of within the terms of the Human Tissue Act 2004.

1199

Consent for publication

1204

Availability of data and materials

1206   The datasets supporting the conclusions of this article are included within the article

1207   and supplementary S1 File.

Competing interests

1209   The authors declare that they have no competing interests.

1210

1211

1212

Acknowledgements

We acknowledge the support of Professor D.M. Mann from the Manchester Brain Bank

who provided some of the human brain tissue samples used in this study. We would

also like to thank Ms S. Selvackadunco from the London Neurodegenerative Diseases

Brain Bank for her assistance with acquiring the human brain samples used in our WB

and TEM studies. We also acknowledge the support of Professors I. Romero, B.

Weksler, and P. Couraud who provided the hCMEC/D3 cell line under MTA.

1220

References

1. Jeste D V, Jin H, Golshan S, Mudaliar S, Glorioso D, Fellows I, et al. Discontinuation of quetiapine from an NIMH-funded trial due to serious adverse events. Am J Psychiatry. 2009 Aug;166(8):937–8.

2. Jeste D V, Blazer D, Casey D, Meeks T, Salzman C, Schneider L, et al. ACNP White Paper: update on use of antipsychotic drugs in elderly persons with dementia. Neuropsychopharmacology [Internet]. 2008 Apr;33(5):957–70. Available from: http://dx.doi.org/10.1038/sj.npp.1301492

3. Murray PS, Kumar S, Demichele-Sweet MAA, Sweet RA. Psychosis in Alzheimer's disease. Biol Psychiatry [Internet]. 2014 Apr;75(7):542–52. Available from: http://dx.doi.org/10.1016/j.biopsych.2013.08.020

4. Reeves S, Bertrand J, D'Antonio F, McLachlan E, Nair A, Brownings S, et al. A population approach to characterise amisulpride pharmacokinetics in older people and Alzheimer's disease. Psychopharmacology (Berl) [Internet]. 2016 Sep;233(18):3371–81. Available from: http://dx.doi.org/10.1007/s00213-016-4379-6

5. Ballard C, Howard R. Neuroleptic drugs in dementia: benefits and harm. Nat Rev Neurosci [Internet]. 2006 Jun;7(6):492–500. Available from: http://dx.doi.org/10.1038/nrn1926

6. Schneider LS, Tariot PN, Dagerman KS, Davis SM, Hsiao JK, Ismail MS, et al. Effectiveness of atypical antipsychotic drugs in patients with Alzheimer's disease. N Engl J Med [Internet]. 2006 Oct;355(15):1525–38. Available from: http://dx.doi.org/10.1056/NEJMoa061240

7. Reeves S, Bertrand J, Uchida H, Yoshida K, Otani Y, Ozer M, et al. Towards safer risperidone prescribing in Alzheimer's disease. British Journal of Psychiatry. 2021 May 1;218(5):268–75.

8. Schlösser R, Gründer G, Anghelescu I, Hillert A, Ewald-Gründer S, Hiemke C, et al. Long-Term Effects of the Substituted Benzamide Derivative Amisulpride on Baseline and Stimulated Prolactin Levels. Neuropsychobiology. 2002;46(1):33–40.

9. Reeves S, Eggleston K, Cort E, McLachlan E, Brownings S, Nair A, et al. Therapeutic D2/3 receptor occupancies and response with low amisulpride blood concentrations in very late-onset schizophrenia-like psychosis (VLOSLP). Int J Geriatr Psychiatry. 2018;33(2):396–404.

10. Reeves S, Bertrand J, D'Antonio F, McLachlan E, Nair A, Brownings S, et al. A population approach to characterise amisulpride pharmacokinetics in older people and Alzheimer's disease. Psychopharmacology (Berl) [Internet]. 2016 Sep;233(18):3371–81. Available from: http://dx.doi.org/10.1007/s00213-016-4379-6

11. Reeves S, McLachlan E, Bertrand J, D'Antonio F, Brownings S, Nair A, et al. Therapeutic window of dopamine D2/3 receptor occupancy to treat psychosis in Alzheimer's disease. Brain. 2017;140(4):1117–27.

12. Clark-Papasavas C, Dunn JT, Greaves S, Mogg A, Gomes R, Brownings S, et al. Towards a therapeutic window of D2/3 occupancy for treatment of psychosis in Alzheimer's disease, with [18F]fallypride positron emission tomography. Int J Geriatr Psychiatry [Internet]. 2014 Oct;29(10):1001–9. Available from: http://dx.doi.org/10.1002/gps.4090

13. Hiemke C, Baumann P, Bergemann N, Conca A, Dietmaier O, Egberts K, et al. AGNP Consensus Guidelines for Therapeutic Drug Monitoring in Psychiatry: Update 2011. Pharmacopsychiatry. 2011 Sep;44(6):195–235.

14.   Lako IM, van den Heuvel ER, Knegtering H, Bruggeman R, Taxis K. Estimating dopamine
D☐ receptor occupancy for doses of 8 antipsychotics: a meta-analysis. J Clin
Psychopharmacol. 2013 Oct;33(5):675–81.
15.   Sparshatt A, Taylor D, Patel MX, Kapur S. Amisulpride - dose, plasma concentration,
occupancy and response: implications for therapeutic drug monitoring. Acta Psychiatr Scand.
2009 Dec;120(6):416–28.
16.   Sekhar GN, Fleckney AL, Boyanova ST, Rupawala H, Lo R, Wang H, et al. Region-specific
blood–brain barrier transporter changes leads to increased sensitivity to amisulpride in
Alzheimer's disease. Fluids Barriers CNS. 2019 Dec 17;16(1):38.
17.   dos Santos Pereira JN, Tadjerpisheh S, Abu Abed M, Saadatmand AR, Weksler B, Romero
IA, et al. The poorly membrane permeable antipsychotic drugs amisulpride and sulpiride are
substrates of the organic cation transporters from the SLC22 family. AAPS J. 2014
Nov;16(6):1247–58.
18.   Horwood N, Davies DC. Immunolabelling of hippocampal microvessel glucose transporter
protein is reduced in Alzheimer's disease. Virchows Arch. 1994;425(1):69–72.
19.   Mooradian AD, Chung HC, Shah GN. GLUT-1 expression in the cerebra of patients with
Alzheimer's disease. Neurobiol Aging. 18(5):469–74.
20.   Kalaria RN, Harik SI. Reduced glucose transporter at the blood-brain barrier and in cerebral
cortex in Alzheimer disease. J Neurochem. 1989 Oct;53(4):1083–8.
21.   Hooijmans CR, Graven C, Dederen PJ, Tanila H, van Groen T, Kiliaan AJ. Amyloid beta
deposition is related to decreased glucose transporter-1 levels and hippocampal atrophy in
brains of aged APP/PS1 mice. Brain Res. 2007 Nov 21;1181:93–103.
22.   Merlini M, Meyer EP, Ulmann-Schuler A, Nitsch RM. Vascular β-amyloid and early
astrocyte alterations impair cerebrovascular function and cerebral metabolism in transgenic
arcAβ mice. Acta Neuropathol. 2011 Sep;122(3):293–311.
23.   Dwyer DS, Pinkofsky HB, Liu Y, Bradley RJ. Antipsychotic drugs affect glucose uptake and
the expression of glucose transporters in PC12 cells. Prog Neuropsychopharmacol Biol
Psychiatry. 1999 Jan;23(1):69–80.
24.   Ardizzone TD, Bradley RJ, Freeman AM, Dwyer DS. Inhibition of glucose transport in PC12
cells by the atypical antipsychotic drugs risperidone and clozapine, and structural analogs of
clozapine. Brain Res [Internet]. 2001 Dec 27;923(1–2):82–90. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11743975
25.   Lean MEJ, Pajonk FG. Patients on Atypical Antipsychotic Drugs. Diabetes Care. 2003 May
1;26(5):1597–605.
26.   Dwyer DS, Donohoe D. Induction of hyperglycemia in mice with atypical antipsychotic
drugs that inhibit glucose uptake. Pharmacol Biochem Behav. 2003 May;75(2):255–60.
27.   Dwyer DS, Pinkofsky HB, Liu Y, Bradley RJ. Antipsychotic drugs affect glucose uptake and
the expression of glucose transporters in PC12 cells. Prog Neuropsychopharmacol Biol
Psychiatry. 1999 Jan;23(1):69–80.
28.   Boyanova S, Wang H, Fleckney AL, Gatt A, Farag DB, Rahman KM, et al. Heightened
sensitivity of people with Alzheimer's disease to the side effects of antipsychotic drug
amisulpride may be mediated through an interaction with glucose transporter 1 at the
blood☐brain barrier. In: Alzheimer's Dement 2020;16(Suppl3):e047395. 2020.
29.   Bethesda (MD): National Library of Medicine (US) NC for BI. PubMed.
https://pubmed.ncbi.nlm.nih.gov/ Accessed 16 March 2021.
30.   Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a major
update to the DrugBank database for 2018. Nucleic Acids Res. 2018;46(D1):D1074–82.

31.   MarvinSketch version 22.9.0. ChemAxon http://chemaxon.com Accessed April 2022 and
1313   April 2023.
1314   32.   Oliva L, Fernandez-Lopez JA, Remesar X, Alemany M. The Anomeric Nature of Glucose
1315   and Its Implications on Its Analyses and the Influence of Diet: Are Routine Glycaemia
1316   Measurements Reliable Enough? J Endocrinol Metab. 2019;9(3):63–70.
1317   33.   Almahmoud S, Wang X, Vennerstrom JL, Zhong HA. Conformational Studies of Glucose
1318   Transporter 1 (GLUT1) as an Anticancer Drug Target. Molecules. 2019 Jun 7;24(11).
1319   34.   Weksler BB, Subileau EA, Perrière N, Charneau P, Holloway K, Leveque M, et al. Blood-
1320   brain barrier-specific properties of a human adult brain endothelial cell line. FASEB J. 2005
1321   Nov;19(13):1872–4.
1322   35.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
1323   Vol. 9, Nature Methods. 2012. p. 671–5.
1324   36.   Wolff A, Antfolk M, Brodin B, Tenje M. In Vitro Blood-Brain Barrier Models-An Overview
1325   of Established Models and New Microfluidic Approaches. J Pharm Sci. 2015
1326   Sep;104(9):2727–46.
1327   37.   Cloyd JC, Snyder BD, Cleeremans B, Bundlie SR, Blomquist CH, Lakatua DJ. Mannitol
1328   pharmacokinetics and serum osmolality in dogs and humans. J Pharmacol Exp Ther. 1986
1329   Feb;236(2):301–6.
1330   38.   Sprague JE, Arbeláez AM. Glucose counterregulatory responses to hypoglycemia. Pediatr
1331   Endocrinol Rev. 2011 Sep;9(1):463–73; quiz 474–5.
1332   39.   Richard BC, Kurdakova A, Baches S, Bayer TA, Weggen S, Wirths O. Gene Dosage
1333   Dependent Aggravation of the Neurological Phenotype in the 5XFAD Mouse Model of
1334   Alzheimer's Disease. J Alzheimers Dis. 2015;45(4):1223–36.
1335   40.   Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, et al. Intraneuronal beta-amyloid
1336   aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial
1337   Alzheimer's disease mutations: potential factors in amyloid plaque formation. J Neurosci.
1338   2006 Oct 4;26(40):10129–40.
1339   41.   Rae EA, Brown RE. The problem of genotype and sex differences in life expectancy in
1340   transgenic AD mice. Neurosci Biobehav Rev. 2015 Oct;57:238–51.
1341   42.   Szu J, Jullienne A, Obenaus A, Territo PR, Consortium M. Lifespan neuroimaging of the
1342   5xFAD mouse model of Alzheimer's disease: Evolution of metabolic and vascular
1343   perturbations. Alzheimer's & Dementia. 2020 Dec 7;16(S4).
1344   43.   Jastrow H. Dr. Jastrows electron microscopic atlas.
1345   http://www.drjastrow.de/WAI/EM/EMAtlas.html Accessed 10 October 2019.
1346   44.   Mason BL, Pariante CM, Jamel S, Thomas SA. Central Nervous System (CNS) Delivery of
1347   Glucocorticoids Is Fine-Tuned by Saturable Transporters at the Blood-CNS Barriers and
1348   Nonbarrier Regions. Endocrinology. 2010 Nov 1;151(11):5294–305.
1349   45.   Báez-Mendoza R, Schultz W. The role of the striatum in social behavior. Front Neurosci.
1350   2013;7.
1351   46.   Garbuzova-Davis S, Rodrigues MCO, Hernandez-Ontiveros DG, Louis MK, Willing AE,
1352   Borlongan C v, et al. Amyotrophic lateral sclerosis: a neurovascular disease. Brain Res. 2011
1353   Jun 29;1398:113–25.
1354   47.   Merlo S, Nakayama ABS, Brusco J, Rossi MA, Carlotti CG, Moreira JE. Lipofuscin
1355   Granules in the Epileptic Human Temporal Neocortex with Age. Ultrastruct Pathol.
1356   2015;39(6):378–84.

48. Pickett EK, Rose J, McCrory C, McKenzie CA, King D, Smith C, et al. Region-specific
depletion of synaptic mitochondria in the brains of patients with Alzheimer's disease. Acta
Neuropathol. 2018;136(5):747–57.
49. Sanderson L, Dogruel M, Rodgers J, de Koning HP, Thomas SA. Pentamidine movement
across the murine blood-brain and blood-cerebrospinal fluid barriers: effect of trypanosome
infection, combination therapy, P-glycoprotein, and multidrug resistance-associated protein. J
Pharmacol Exp Ther. 2009 Jun;329(3):967–77.
50. Ouellette RJ, Rawn JD. Carbohydrates. In: Organic Chemistry. Elsevier; 2018. p. 889–928.
51. Dwyer DS, Liu Y, Bradley RJ. Dopamine receptor antagonists modulate glucose uptake in
rat pheochromocytoma (PC12) cells. Neurosci Lett. 1999 Oct;274(3):151–4.
52. Chow VW, Mattson MP, Wong PC, Gleichmann M. An Overview of APP Processing
Enzymes and Products. Neuromolecular Med. 2010 Mar 24;12(1):1–12.
53. Babkin P, George Thompson AM, Iancu C v, Walters DE, Choe JY. Antipsychotics inhibit
glucose transport: Determination of olanzapine binding site in Staphylococcus epidermidis
glucose/H(+) symporter. FEBS Open Bio. 2015;5:335–40.
54. Chokhawala K, Stevens L. Antipsychotic Medications [Internet]. In: StatPearls [Internet].
Treasure Island (FL): StatPearls Publishing; 2023 [cited 2023 Apr 6]. Available from:
https://www.ncbi.nlm.nih.gov/books/NBK519503/
55. Natesan S, Reckless GE, Barlow KBL, Nobrega JN, Kapur S. Amisulpride the 'atypical'
atypical antipsychotic — Comparison to haloperidol, risperidone and clozapine. Schizophr
Res. 2008 Oct;105(1–3):224–35.
56. Devés R, Krupka RM. Cytochalasin B and the kinetics of inhibition of biological transport. A
case of asymmetric binding to the glucose carrier. Biochimica et Biophysica Acta (BBA) -
Biomembranes. 1978 Jul;510(2):339–48.
57. Kapoor K, Finer-Moore JS, Pedersen BP, Caboni L, Waight A, Hillig RC, et al. Mechanism
of inhibition of human glucose transporter GLUT1 is conserved between cytochalasin B and
phenylalanine amides. Proc Natl Acad Sci U S A. 2016 Apr 26;113(17):4711–6.
58. Niccoli T, Cabecinha M, Tillmann A, Kerr F, Wong CT, Cardenes D, et al. Increased
Glucose Transport into Neurons Rescues Aβ Toxicity in Drosophila. Current Biology. 2016
Sep;26(17):2291–300.
59. Kable A, Fullerton A, Fraser S, Palazzi K, Hullick C, Oldmeadow C, et al. Comparison of
Potentially Inappropriate Medications for People with Dementia at Admission and Discharge
during An Unplanned Admission to Hospital: Results from the SMS Dementia Study.
Healthcare. 2019 Jan 9;7(1):8.
60. Orsel K, Taipale H, Tolppanen AM, Koponen M, Tanskanen A, Tiihonen J, et al.
Psychotropic drugs use and psychotropic polypharmacy among persons with Alzheimer's
disease. European Neuropsychopharmacology. 2018 Nov;28(11):1260–9.
61. Al-Ahmad AJ. Comparative study of expression and activity of glucose transporters between
stem cell-derived brain microvascular endothelial cells and hCMEC/D3 cells. Am J Physiol
Cell Physiol. 2017 Oct 1;313(4):C421–9.
62. Ohtsuki S, Ikeda C, Uchida Y, Sakamoto Y, Miller F, Glacial F, et al. Quantitative targeted
absolute proteomic analysis of transporters, receptors and junction proteins for validation of
human cerebral microvascular endothelial cell line hCMEC/D3 as a human blood-brain
barrier model. Mol Pharm. 2013 Jan 7;10(1):289–96.
63. Weksler B, Romero IA, Couraud PO. The hCMEC/D3 cell line as a model of the human
blood brain barrier. Fluids Barriers CNS. 2013 Mar 26;10(1):16.

64.   Meireles M, Martel F, Araújo J, Santos-Buelga C, Gonzalez-Manzano S, Dueñas M, et al.
Characterization and modulation of glucose uptake in a human blood-brain barrier model. J
Membr Biol. 2013 Sep;246(9):669–77.
65.   Coukell AJ, Spencer CM, Benfield P, Carpenter Tr WT. DRUG EVALUATION
Amisulpride eNS A Review of its Pharmacodynamic and Pharmacokinetic Properties and
Therapeutic Efficacy in the Management of Schizophrenia. Vol. 6, Drugs. 1996.
66.   Burant CF, Sivitz WI, Fukumoto H, Toshiaki K, Nagamatsu S, Susumo S, et al. Mammalian
Glucose Transporters: Structure and Molecular Regulation. Vol. 47, RECENT PROGRESS
IN HORMONE RESEARCH: Proceedings of the 1990 Laurentian Hormone Conference.
1991.
67.   Zentiva. Patient Information Leaflet: Amisulpride.
https://www.medicines.org.uk/emc/files/pil.3965.pdf Accessed 13 January 2022.
68.   Holt RIG. Association Between Antipsychotic Medication Use and Diabetes. Curr Diab Rep.
2019 Oct 2;19(10):96.
69.   Ruggieri S, de Pandis MF, Bonamartini A, Vacca L, Stocchi F. Low dose of clozapine in the
treatment of dopaminergic psychosis in Parkinson's disease. Clin Neuropharmacol. 1997
Jun;20(3):204–9.
70.   Forner S, Kawauchi S, Balderrama-Gutierrez G, Kramár EA, Matheos DP, Phan J, et al.
Systematic phenotyping and characterization of the 5xFAD mouse model of Alzheimer's
disease. Sci Data. 2021;8(1):270.
71.   Sadleir KR, Eimer WA, Cole SL, Vassar R. Aβ reduction in BACE1 heterozygous null
5XFAD mice is associated with transgenic APP level. Mol Neurodegen. 2015 Jan 7;10:1.
72.   Maarouf CL, Kokjohn TA, Whiteside CM, Macias MP, Kalback WM, Sabbagh MN, et al.
Molecular Differences and Similarities Between Alzheimer's Disease and the 5XFAD
Transgenic Mouse Model of Amyloidosis. Biochem Insights. 2013;6:1–10.
73.   Ahn KC, Learman CR, Dunbar GL, Maiti P, Jang WC, Cha HC, et al. Characterization of
Impaired Cerebrovascular Structure in APP/PS1 Mouse Brains. Neuroscience.
2018;385:246–54.
74.   Park R, Kook SY, Park JC, Mook-Jung I. Aβ1-42 reduces P-glycoprotein in the blood-brain
barrier through RAGE-NF-κB signaling. Cell Death Dis. 2014 Jun 26;5:e1299.
75.   Baloyannis SJ. Brain capillaries in Alzheimer's disease. Hell J Nucl Med. 18 Suppl 1:152.
76.   Ossenkoppele R, Prins ND, van Berckel BN. Amyloid imaging in clinical trials. Alzheimers
Res Ther. 2013;5(4):36.
77.   Mosconi L, Rinne JO, Tsui WH, Murray J, Li Y, Glodzik L, et al. Amyloid and metabolic
positron emission tomography imaging of cognitively normal adults with Alzheimer's
parents. Neurobiol Aging. 2013 Jan;34(1):22–34.
78.   Mosconi L, Mistur R, Switalski R, Tsui WH, Glodzik L, Li Y, et al. FDG-PET changes in
brain glucose metabolism from normal cognition to pathologically verified Alzheimer's
disease. Eur J Nucl Med Mol Imaging. 2009 May;36(5):811–22.
79.   Landau SM, Harvey D, Madison CM, Koeppe RA, Reiman EM, Foster NL, et al.
Associations between cognitive, functional, and FDG-PET measures of decline in AD and
MCI. Neurobiol Aging. 2011 Jul;32(7):1207–18.
80.   Simpson IA, Chundu KR, Davies-Hill T, Honer WG, Davies P. Decreased concentrations of
GLUT1 and GLUT3 glucose transporters in the brains of patients with Alzheimer's disease.
Ann Neurol. 1994 May;35(5):546–51.

81.   Winkler EA, Nishida Y, Sagare AP, Rege S v, Bell RD, Perlmutter D, et al. GLUT1
reductions exacerbate Alzheimer's disease vasculo-neuronal dysfunction and degeneration.
Nat Neurosci. 2015 Apr;18(4):521–30.
82.   Löscher W, Langer O. Imaging of P-glycoprotein function and expression to elucidate
mechanisms of pharmacoresistance in epilepsy. Curr Top Med Chem. 2010;10(17):1785–91.
83.   Chiu C, Miller MC, Monahan R, Osgood DP, Stopa EG, Silverberg GD. P-glycoprotein
expression and amyloid accumulation in human aging and Alzheimer's disease: preliminary
observations. Neurobiol Aging. 2015 Sep;36(9):2475–82.
84.   Chai AB, Leung GKF, Callaghan R, Gelissen IC. P-glycoprotein: a role in the export of
amyloid-β in Alzheimer's disease? FEBS J. 2020;287(4):612–25.
1458

1459

Supporting Information

1461

1462   • S1 File.docx

1463

1464   • 6 Supplementary Tables

1465

1466   • 13 Supplementary Figures

Amisulpride microspecies A (96.77%)

Amisulpride microspecies B (3.23%)

A

GLUT1

GAPDH

P30 P31 P33 Caco2   HEK

40-60 kDa

37 kDa

B

[³H]Mannitol

C

[¹⁴C]D-glucose

A:

B:

* – electron dense portions of lipofuscin granules, ⬈ – lucid portions of lipofuscin granules

* – vacuolisation,   ▲ – myelinated whorled masses,   * – condensed axonal cytoplasm

Frontal Cortex