Title:

Spectral consistency in sound sequence affects perceptual accuracy

in discriminating subdivided rhythmic patterns

Running title:

Spectral consistency affects perceptual accuracy of subdivided

rhythm

Author:

Jun Nitta[1], Sotaro Kondoh [1], Kazuo Okanoya[1,2**], Ryosuke O. Tachibana[1*]

Affiliation

[1] Graduate School of Arts and Sciences, the University of Tokyo, Tokyo, Japan

[2] Advanced Comprehensive Research Organization, Teikyo University, Tokyo, Japan

[**] co-corresponding author

E-mail: kazuookanoya@gmail.com

[*] corresponding author

E-mail: rtachi@gmail.com

Abstract:

Musical compositions are distinguished by their unique rhythmic patterns, determined by subtle differences in how regular beats are subdivided. Precise perception of these subdivisions is essential for discerning nuances in rhythmic patterns. While musical rhythm typically comprises sound elements with a variety of timbres or spectral cues, the impact of such spectral variations on the perception of rhythmic patterns remains unclear. Here, we show that consistency in spectral cues affects perceptual accuracy in discriminating subdivided rhythmic patterns. We conducted online experiments using rhythmic sound sequences consisting of band-passed noise bursts to measure discrimination accuracy. Participants were asked to discriminate between a swing-like rhythm sequence, characterized by a 2:1 interval ratio, and its more or less exaggerated version. This task was also performed under two additional rhythm conditions: inversed-swing rhythm (1:2 ratio) and regular subdivision (1:1 ratio). The center frequency of the band noises was either held constant or alternated between two values. Our results revealed a significant decrease in discrimination accuracy when the center frequency was alternated, irrespective of the rhythm ratio condition. This suggests that rhythm perception is not only shaped by temporal structure but also affected by spectral properties.

Introduction

We detect temporal regularity and capture complex structures of sound events in almost all styles of music. Each perceived regular pulse is called a beat. Subdividing each beat interval provides various characteristics in the rhythmic pattern of the music. In other words, we perceptually organize various sound events into a rhythm structure according to their temporal properties by detecting and subdividing regular patterns in sound sequences [1]. For example, subdividing beats with simple integer ratios such as 1:1 and 2:1 is far more frequently observed in Western music [2]. In contrast, complex ratios such as 5:3 are very common in drum ensemble music in West Africa [3]. One of the famous subdivisions is swing. Swing is a long-short duration alternating pattern considered an essential factor in jazz music. The integer ratio of swing is generally considered to be 2:1, but it approaches 1:1 as the tempo increases. [4]. This study focuses on the perceptual accuracy of such subdivision rhythms.

Various temporal factors influence our rhythm perception. We can easily perceive regular beats even though the actual timings of the performed beats usually fluctuate and deviate from an exact regular pattern [5]. Despite such variations in beat durations, audiences perceive temporal regularity in the performed sound pattern and interpret the variation as intentional aesthetic expressions [6,7]. When beats are subdivided into finer and/or uneven intervals, rhythm perception becomes more variable. Equal subdivisions cause the tempo to be perceived as slower, even while the actual tempo remains constant [8]. Conversely, sound sequences with an identical subdivision pattern but different tempi tend to be perceived as different rhythmic patterns [9]. Precise subdivision perception relies on detecting subtle relative differences among time onsets in a regular beat sequence. We can notice approximately 2.5 % displacement of one note

onset in an isochronous sequence with a tempo of 240-ms inter-onset interval [4],

although such detection threshold increases when the tempo becomes faster [11].

Moreover, asymmetries have been reported in which subdivisions are perceived as

equally spaced, even though they are short-long patterns, and this does not happen in

long-short patterns [15].

It is known that not only the temporal property but also the sound timbre, or

spectral property, influences how we organize the sound sequence structure. This issue

has been intensively investigated in the research field of auditory stream segregation

[16,17]. For example, a study using band noise suggested that differences in the center

frequency of the band noise best predicted the degree to which stream segregation

occurred [18]. Another study used a sound sequence alternating between pure tone and

narrow-band noise regularly and delaying only even-numbered sounds gradually [19].

The result showed that the greater the difference in spectral aspect between the two sound

elements, the less likely the subjects were to notice the delay of even-numbered sounds.

The authors claimed that differences in timbre enhanced the stream segregation, thereby

making it more difficult to perceive the temporal relationship of sound elements across

different streams. This raises the possibility that spectral variations might influence the

accuracy of rhythm perception. In real music pieces, rhythmic patterns typically consist

of sounds with different timbres, such as those produced by a drum set. Consequently, an

experimental assessment of how varying spectral cues affects the accuracy of rhythmic

pattern perception offers novel insights for understanding rhythm perception in practical

musical contexts.

In the present study, we examined the effect of different timbres on rhythm

perception by psychophysically measuring the discrimination accuracy of multiple

rhythmic patterns, including swing rhythms. We used a sound sequence consisting of

band noise bursts, and manipulated the spectral cue by changing their center frequency.

Experiments were performed online via a web browser to recruit many participants. We

conducted three experiments corresponding to three rhythmic patterns: long-short rhythm

(Exp. 1), short-long rhythm (Exp. 2), and straight rhythm (Exp. 3). To confirm the

reliability of online experiments, particularly in the aspect of data reproducibility across

arbitrary sound listening environments, we additionally conducted an experiment with

participants who performed their tasks in the local laboratory so that we could confirm

the listening environment (Exp. 1').

Methods

Participants

We initially recruited a hundred participants for each of three online experiments via a

crowd-sourcing service (CrowdWorks, Inc., Japan). None of them reported a history of

hearing problems. A subset of participants was excluded from the analysis (5, 6, and 14

participants in Exps. 1, 2, and 3, respectively) due to incomplete data retrieval caused by

malfunctions in the online experiment system. Then, we screened the dataset according

to correct rates in screening trials and goodness of fit to the psychometric function (see

Analysis section). After the screening, data from 40 (20 males, 20 females; age: $M \pm SD$

$= 39.2 \pm 8.10$ yo), 43 (21 males, 21 females, 1 other; age: 36.3 ± 8.66), and 30 (18 males,

12 females, age: 38.9 ± 8.96) participants for Exps. 1, 2, and 3 were included in further

analysis (**Table 1**). We allowed participants to join multiple experiments and additionally

assessed intra-individual factors across different experiments. Several people participated

in two experiments: after screening, 12, 10, and 12 participants overlapped in Exps. 1-2,

2-3, and 3-1, respectively. Three participated in all experiments. We additionally

conducted the Exp. 1' on 16 people (8 males, 8 females; age: 27.5 ± 6.74) in our local

laboratory without using the crowd-sourcing service so that we were able to confirm the

acoustic environment of the participants.

We confirmed music expertise with a questionnaire. None of the participants was

an expert musician, but almost half had experienced intensive music instrument practices

as amateurs: 46, 47, and 46 participants in Exps. 1, 2, and 3, respectively. Note that the

intensive music experience refers to training at least two hours per day and may include

experience in music classes and club activities but excludes experience in compulsory

music classes. All participants were given informed consent prior to the experiment. All

experimental procedures were approved by Ethics Review Committee on Experimental

Research with Human Subjects of the Graduate School of Arts and Sciences, The

University of Tokyo (No. 718-3).

Table 1. The number of participant information.

	Exp. 1	Exp. 2	Exp. 3	Exp. 1'
screened (recruited)	40 (95)	43 (94)	30 (86)	16 (16)
male	20 (53)	21 (51)	18 (50)	8 (8)
female	20 (42)	21 (42)	12 (36)	8 (8)
other	0 (0)	1 (1)	0 (0)	0 (0)

Sound stimuli

We employed sound sequences of brief band-passed noises as stimuli (**Figure 1**). The

center frequency of the band noise was manipulated to form two conditions: the same-

frequency condition, in which the center frequency was fixed at 1500 Hz, and the

different-frequency condition, where the center frequency was alternated between 1000

and 2250 Hz (14 semitones apart). The bandwidth was fixed at 1000 Hz. Each band noise

was created from white noise by band-pass filtering (FIR, for 1000-Hz noise: 582 taps,

1500-Hz: 1020 taps, 2250-Hz: 2038 taps), which was designed by the 'band-pass'

function in MATLAB. The duration of each noise burst was 30 ms with a rise and fall

time of 10 ms.

Each stimulus sequence consisted of eight noise bursts. Inter-onset intervals of

odd number-th noises were fixed at 300 ms, which formed musical beats (hence, we call

this inter-beat interval: IBI). Onsets of even number-th noises were varied according to

an interval ratio index *r*, defined as follows:

$$r = 1 - (T/2 - d)/(T/2 + d) \qquad (1)$$

here, *T* represents IBI, and *d* shows onset displacement from the middle timepoint

between beats (*T*/2). The *r* value becomes 0, 0.5, or −0.5 for the interval ratio of 1:1

(straight rhythm), 2:1 (swing), or 1:2 (inverse-swing), respectively. In the different-

frequency condition, either the odd-number-th (on-the-beat) noises or the even-number-

th (off-the-beat) ones had the higher frequency band (2250-Hz centered), and the others

were placed at the lower band (1000-Hz centered). All sound stimuli were digitally

generated at 44.1 kHz, 16 bits sampling. Their amplitudes were adjusted to have the same

root-mean-square level.

The stimuli in the different-frequency condition had two different types: high-

low (on-beat noises had higher center frequency) and low-high (on-beat noises had lower),

though this factor was out of our interest in this study. We randomly assigned participants

to the one of two conditions (each participant experienced only either high-low or low-

high stimuli). To examine the effect of high-low order on discrimination accuracy, we

used a linear mixed model. The model fitting showed that neither high-low order or

interaction had a significant effect on σ (see Table S1). This effect was consistent in Exps.

1, 2, and 3. This suggests that not high-low order but center-frequency condition has an

effect on discrimination accuracy. Thus, we ignored this factor by pooling data from these

two groups.

Figure 1. Schematic drawing of stimuli design for interval ratio r = 0.5. Upper indicates a

stimulus used in the same-frequency condition. Lower shows a stimulus used in the different-

frequency condition. The thick solid line (blue) indicates beats.

Task and condition

In each experimental trial, two stimulus sequences were successively presented with a

300-ms interval between them. The former sequence was the standard stimulus, and the

latter was the comparison. We conducted three experiments (Exps. 1, 2, and 3) in which

the standard stimuli had the interval ratio of r = 0.5, −0.5, and 0 (corresponding to 2:1,

1:2, and 1:1), respectively. The comparison stimuli were prepared with varying r values

in five small steps: 0.2, 0.4, 0.5, 0.6, 0.8 (Exp. 1); −0.2, −0.4, −0.5, −0.6, −0.8 (Exp. 2); −

0.3, −0.15, 0, 0.15, 0.3 (Exp. 3). Participants were asked to choose "bouncier one"

(corresponding to larger absolute r value) from the two stimulus sequences in Exps. 1 and

2, while asked to select "equally spaced one" in Exp. 3. We randomized the order of the

standard and comparison stimuli in Exp. 3 to avoid selection bias since the standard was

always "equally spaced" in this experiment.

Participants were required to respond by mouse click within 2.5 s after the offset

of the second stimulus sequence. After participants responded, the correctness of the

answer was feedbacked as symbols ○ (correct) or × (incorrect) on the computer screen.

To confirm whether participants appropriately focused on the task, we inserted screening

trials in experimental sessions. In the screening trial, the interval ratio of comparison

stimuli was $r = 0.2$, −0.2, or 0.3 in Exps. 1, 2, or 3, respectively, so that participants could

easily choose the correct response if they kept paying attention to the presented stimuli.

Procedure

Each of Exps. 1-3 consisted of 240 trials (2 frequency conditions × 5 interval ratio

conditions × 24 trial repetitions) that were equally separated into six blocks with inserting

brief rests. In Exp. 1', the number of repetitions was increased to 40 trials (total 400 trials)

to obtain stable data. The frequency conditions and onset displacement conditions were

randomized within the block. Screening trials were added at the beginning and end of the

first and second blocks, respectively. Before starting the experiment, participants

underwent a practice session that consisted of 8 trials, one for each of the eight conditions,

except for the one with an onset displacement of 0 ms. Correct/incorrect feedback was

provided after each response during the practice session. When the response time

exceeded the required time (2.5 s), the participant was immediately informed that the time

was up.

Before the experiment session, we conducted a headphone-screening task to

confirm whether the participants wore headphones. Participants were asked to select the

sound with the lowest loudness among the three different sound stimuli, according to a

previous report [21]. The three sound stimuli were (i) a diotic presentation of 200-Hz pure

tone, (ii) a 6-dB lower 200-Hz pure tone, and (iii) a dichotic presentation of 200-Hz pure

tone to one ear side and its antiphase waveform to the other side. After the experiment,

we asked participants about their music experience using a questionnaire. This

questionnaire included questions about musical instrument experience, daily music

listening, favorite genres and songs, and dance or rhythm game experience.

All experiments were programmed using a JavaScript-based experiment builder

(lab.js [20]) and conducted online via Open Lab (Open Lab Online UG, Konstanz,

Germany). The entire experiment took around 30 minutes.

Analysis

To assess whether discrimination accuracy varies with spectral cue, we classified data

into the same- and different-frequency conditions, and approximated by the curves

described below, respectively. For each participant, we calculated the mean response rate

for each interval ratio. In Exps. 1 and 2, the obtained response rate was approximated by

a sigmoid function (cumulative normal distribution). The shape of sigmoid curve is

determined by two parameters: μ gives the point of maximal gradient, which corresponds

to the point of subjective equality (PSE); and σ shows the steepness of the gradient, which

is equivalent to the discrimination accuracy. We calculated μ and σ for each subject and

compared them with frequency conditions. For Exp. 3, we fitted a simple Gaussian

function with fixing μ at 0 and the maximum value at 0.5 instead of using the sigmoid

function because the response rate in Exp. 3 were supposed to be the highest (0.5) at the

standard stimulus interval ratio ($r = 0$). Additionally, we performed partial fittings for data

either in positive or negative interval ratio values ($0 \leq r < 1$ or $-1 < r \leq 0$) to assess the

positive-negative asymmetry of discrimination accuracy.

In Exps. 1, 2 and 3, data with one or more incorrect answers in the screening

trials were excluded as participants did not respond appropriately. We set further

screening criteria to maintain the data reliability as follows: (1) the adjusted coefficient

of determination (R^2) in the sigmoid fitting must be 0.5 or larger; (2) the absolute value

of PSE ($|\mu|$) must range between 0 and 1; (3) the discrimination accuracy (σ) must be 0 or

larger. We excluded participants from further analyses if the data did not satisfy these

criteria either in the same-frequency or different-frequency condition.

In Exps. 1, 2 and 3, we performed a supplementary analysis of the effects of

high-low order and center-frequency conditions on PSE and discrimination accuracy

using the linear mixed model as follows: *Y ~ frequency + order + frequency × order +*

(1|participant). *Y* indicates independent parameter, μ (PSE) or log σ (discrimination

accuracy), *frequency* indicates dummy variable representing center-frequency conditions

(0: same-frequency; 1: different-frequency), *order* indicates dummy variable representing

high-low order (0: high-low; 1: low-high), and *(1|participant)* indicates random effect for

each participant. We logarithmically transformed σ to treat it as a parametric variable. We

performed the analysis in R version 4.3.1. To fit the model, we used 'lmer' function of

the lme4 and lmerTest package. We also adopted 'r.squaredGLMM' function of the

MuMIn package to calculate the goodness of fit with R-squared value. To examine the

effect of high-low order on discrimination accuracy, we used a linear mixed model. After

screening, participants were divided into two groups roughly in half (Exp. 1: $n_{\text{high-low}}$ =

19, $n_{\text{low-high}}$ = 21; Exp. 2: $n_{\text{high-low}}$ = 22, $n_{\text{low-high}}$ = 21; Exp. 3: $n_{\text{high-low}}$ = 12, $n_{\text{low-high}}$ = 18).

The model fitting showed that σ was significantly higher in the different-frequency

condition, and neither high-low order or interaction had a significant effect on σ (Table

2). This effect was consistent in Exps. 1, 2, and 3. This suggests that only center-frequency

condition has an effect on discrimination accuracy. The model fitting performance was

moderate (Exp.1: R^2 = 0.52; Exp.2: R^2 = 0.51; Exp. 3: R^2 = 0.56). These suggest that

neither high-low order nor interaction effects contribute to discrimination accuracy. The

linear-mixed model fitted to μ showed that no consistent effects were found in

Experiments 1 and 2 (Table 3). In Exp. 1, the interaction was significant, but the goodness

of fit was low (R^2 = 0.23). In Exp. 2, μ was significantly higher in the different-frequency

condition (R^2 = 0.63). This suggests that the effect of high-low order on PSE differs

depending on the long-short and short-long rhythms.

260   Homogeneities of variance in μ and σ data were examined by the two-sample F

test, which failed to guarantee the homogeneity in μ in Exp. 1 ($F_{(39, 39)}$ = 2.32, p =

0.010), σ in Exp. 1 ($F_{(39, 39)}$ = 7.88, $p < 0.001$), σ in Exp. 2 ($F_{(42, 42)}$ = 3.02, $p <$

0.001), and σ in Exp. 3 (negative: $F_{(17, 17)}$ = 7.47, $p < 0.001$; positive: $F_{(17, 17)}$ = 3.22,

p = 0.021). Therefore, we employed Wilcoxon signed-rank tests for this analysis. The

significance level α was set at 0.05. Post hoc analysis confirmed that the statistical power

$(1-\beta)$ was more than 0.8 in Exps. 1, 2, and 3 (Exp. 1: $1-\beta$ = 0.99; Exp. 2: $1-\beta$ = 0.98; Exp.

3: $1-\beta$ = 0.99). Note that there were some data (n = 3) in which σ was very close to zero

in Exp. 3, and we conducted the test in each case when they were excluded and not

excluded.

Result

Reliability of the online experiment

First, we confirmed the reliability of the online experiments regarding hearing environments by comparing the results of Exp. 1 with its offline version Exp. 1'. The critical but uncontrollable factor in online experiments is the potential difference in acoustic conditions, such as the frequency characteristics of headphones and background noise from the environment. In our case, we used band-passed noise to reduce the influence of variability in the frequency characteristics of the participants' headphones. However, it is almost impossible to control the background noise factor. Therefore, we here performed the same experiment as in Exp. 1 in the local laboratory for a subset of participants, that is, under identical background conditions. Sigmoidal approximations for Exps. 1 and 1' showed almost exactly overlapping curves for the same-frequency condition (**Figure 2a**), demonstrating that the potential variability in listening environments did not affect the main results of our online experiments.

Exp. 1&2: long-short rhythm (2:1), short-long rhythm (1:2)

We then assessed the effect of spectral differences in sound sequences on the discrimination accuracy of the long-short rhythm and short-long rhythm in Exp. 1 and 2, respectively. The accuracy was measured as the steepness (σ) of the fitted sigmoid curve. We screened the data according to the exclusion criteria and analyzed the remaining data obtained from 40 participants in Exp. 1 (**Figure 2a-c**). We calculated the response rate in each interval ratio and found that the steepness of fitted curve in the different-frequency condition was less than that in the same-frequency condition (**Figure 2a**). The estimated

PSE (μ) (**Figure 2b**) was not significantly different between the two conditions (Wilcoxon

signed-rank test, $Z = 1.83$, $p = 0.068$). In contrast, the discrimination accuracy (σ) (**Figure**

2c) was significantly larger in the different-frequency condition than in the same-

frequency condition ($Z = 4.73$, $p < 0.001$).

We also analyzed the screened data obtained from 43 participants in Exp. 2 (**Figure**

2d-f), and found that the steepness in the different-frequency condition was less than that

in the same-frequency condition (**Figure 2d**), similar to the long-short rhythm in Exp. 2.

Both the estimated μ (**Figure 2e**) and σ (**Figure 2f**) were significantly greater in the

different-frequency condition (μ: $Z = 3.90$, $p < 0.001$; σ: $Z = 3.75$, $p < 0.001$).

These results demonstrate that the discrimination accuracy decreases when the center

frequency of narrow-band noises is alternated; thus, spectral inconsistency in the sound

sequence deteriorates the discrimination of subdivided rhythms.

Exp. 3: Straight Rhythm (1:1)

In Exp. 3, data from 30 participants survived after the screening. Different from Exp. 1

and 2, the response rate was assumed to be 0.5 at the regular interval ($r = 0$) and 0 at both

ends ($r = 1$ and -1) because of the task design. We found again that the fitted curve in

the different-frequency condition was less steep than that in the same-frequency condition

(**Figure 3a**). The symmetric fitting showed that the estimated σ was significantly larger

in the different-frequency condition ($Z = 4.78$, $p < 0.001$). The result indicates that the

discrimination accuracy decreases with the spectral inconsistency even for the straight

rhythm, similar to the long-short and short-long rhythms.

We also performed the partial fitting for each of the positive and negative interval

ratios to assess the asymmetry of the response rate data (shown as dotted lines in

Figure 3a). The estimated σ of the asymmetric fitting (**Figure 3c**) was significantly larger

for the negative interval ratio than for the positive ratio in both frequency: conditions

(same-frequency: $Z = 4.22$, $p < 0.001$ different-frequency: $Z = 3.18$, $p = 0.001$).

Effect of music experience

To examine the effect of musical instrument experience, participants in Exp. 1 were

divided into two groups (with experiences: $n = 23$; without experiences: $n = 20$) by

reporting whether they had instrumental experience, and σ was calculated for each

frequency condition. The results showed a significant difference in σ between the

frequency conditions in both groups (with experiences: $Z = -3.47$, $p < 0.001$; without

experiences: $Z = -3.54$, $p < 0.001$). This suggests that discrimination accuracy decreased

under the frequency conditions, regardless of musical experience.

Figure 2. PSE and discrimination accuracy in long-short (Exp. 1 and 1', a-c) and short-long rhythm (Exp. 2, d-f). (a) Response rates that comparison stimuli were "more bouncing" for each interval ratio. Red line indicates the different-frequency condition, and blue line indicates the same-frequency condition in Exp. 1. Light-blue dotted line indicates the same-frequency condition in Exp. 1'. Error bar shows standard error. Curves are sigmoidal approximations. **(b)** Distribution of PSE (μ) for same- and different-frequency conditions. Black dots and lines indicate participants in the group of high-low order, and gray indicates those in low-high order. **(c)** Distribution of discrimination accuracy (σ) for same- and different-frequency conditions. **(d)** Response rates that the comparison stimuli were "more bouncing" for each interval ratio. Red line indicates the different-frequency condition, and blue line indicates the same-frequency condition. Error bars are standard errors. Curves are sigmoidal approximations. **(e)** Distribution of PSE (μ) for same- and different-frequency conditions. **(f)** Distribution of discrimination accuracy (σ) for same- and different-frequency conditions. * $p <$ 0.05.

Figure 3. Discrimination accuracy in straight rhythm (Exp. 3). (a) Response rates that the comparison stimuli were "more bouncing" for each interval ratio. Red line indicates the different-frequency condition, and blue line indicates the same-frequency condition. Error bars are standard errors. Curves drawn by a solid line indicate symmetric fitting. Curves drawn by dotted line indicates asymmetric fitting. (b) Distribution of PSE ($|\mu|$) for same- and different-frequency conditions. Black dots and lines indicate participants in the group of high-low order, and gray indicates those in low-high order. (c) Distribution of accuracy (σ) for same- and different-frequency conditions. * $p < 0.05$.

Discussion

The present study aimed to clarify how spectral consistency in sound sequences affects the perception of subdivision rhythm patterns. We assessed participants' rhythm discrimination of band-pass noise sequences with manipulating sound timbre by differentiating their center frequencies. We found that discrimination accuracies (measured as σ of the fitted sigmoid function) decreased when the center frequencies alternated, irrespective of the rhythm pattern difference. This result indicates that the spectral consistency of sound sequences influences the accuracy of subdivided rhythm perception.

Effect of spectral consistency on perceptual accuracy

Our analysis revealed that the discrimination accuracy decreased when the center frequency of the sequential band noises alternated (**Figure 2, 3**). This may be because a sound sequence that includes two spectral cues can be perceived as two segregated streams. Similarly, previous studies have reported that spectral differences promote stream segregation, which makes change detection more difficult for regular-interval (straight) rhythms [18,19], even though our results showed the same effect on non-straight rhythms. This decrease in perceptual accuracy can be quantified as the discrimination threshold. A study reported 6.2 ms displacement in the isochronous tone sequence with 130 ms intervals (corresponding to 260 ms IBI in our case) as a just-noticeable difference [4]. Another study estimated the threshold for discriminating swing from straight rhythms to be approximately 18 ms for 300-ms IBI [22]. We roughly estimated the thresholds in our data as interval ratios crossing 25% and 75% of the fitted curves. The mean estimated thresholds for the same-frequency condition in Exps. 1, 2, and 3 were 8.7, 7.1, and 9.6

ms, respectively. That for the different-frequency condition were 15.5, 10.7, and 19.5 ms, showing an apparent increase in the thresholds (corresponding to the decrease in perceptual accuracy) even though the estimated values were within the range of thresholds that were previously reported.

These findings suggest the following mechanism. Perceived timbre difference among sound elements produced by the center frequency manipulation enhances the segregation of auditory streams, and then the segregation elevates the threshold of discriminating subtle temporal shifts among sound elements across different streams, making the perceptual accuracy decrease. Note that we did not ask the participants to self-report whether they perceived sound streams as integrated or segregated. The precise relationship between our results and stream segregation should be addressed.

Response asymmetry

The point of subjective equality (PSE) of the interval ratio in both Exps. 1 and 2 shifted towards 0 in the same-frequency condition (**Figure 2b, 2e**), indicating that participants perceived two stimuli equivalent when the comparison stimulus was slightly closer to the straight rhythm. The shift was not observed in the different-frequency condition. This effect may be due to the fixed presentation order of the standard and comparison stimuli in these experiments, e.g., short-term memory of the standard deformed before hearing the comparison; though such discussions were out of the scope of this study.

In Exp. 3, the estimated σ was significantly different between positive and negative interval ratios in both the same-frequency and different-frequency conditions (**Figure 3c**). This asymmetry can be explained by the time-shrinking effect [23]. When three consecutive tones with two onset intervals (T1 and T2), we tend to perceive the

second time interval as shorter than the first (T2 < T1). This effect becomes weaker when

the second interval is physically shorter than the first. A previous study showed that

participants detected a negative shift of interval ratio more accurately than a positive shift

in 1:2 rhythm [14], which is consistent with our result.

Individual variability

Our analysis did not find a significant effect of musical expertise. Contrary to our result,

several studies have suggested that musical experience increases the accuracy of rhythm

perception. For instance, people who have high musical experiences or music experts

detect smaller changes in regular interval patterns than people with low musical

experience [24] [25]. This discrepancy may be because no experts were recruited in our

study.

One might wonder if participants exhibit consistent tendencies across different

experiments. We started three online experiments at almost the same time, and some

participants have joined two or three of them (see Participants subsection in Methods).

Thus, we additionally performed *ad hoc* analysis for this issue by comparing individual

discrimination abilities between Exp. 1 and 2 for people who participated in both

experiments. We used less strict criteria for screening to increase the sample size ($n = 23$),

in which we allowed to include participants who made a incorrect response once in the

screening trials either in Exp. 1 or 2. As a result, almost all participants showed consistent

tendency: both $|\mu|$ and σ were larger in the different-frequency condition than in the same-

frequency condition (**Figure 4**). These indicate consistent tendencies in PSE and

discrimination accuracy across different experiments.

Figure 4. Comparison of μ and σ for overlap participants in Exp. 1 (a) and 2 (b).

Scatterplots of the absolute value of PSE ($|\mu|$) in the different-frequency condition for each $|\mu|$

in the same-frequency condition. Open circles indicate the values in Exp. 1, and filled circles

indicate the values in Exp. 2. Red indicates all correct answers to the screening trials, blue

indicates one incorrect answer to the screening trials in Exp. 1, and green indicates one incorrect

answer to the screening trials in Exp. 2. Solid lines connect the same participants.

Reliability of online experiment

The acoustic environment was variable among participants since they potentially heard

stimuli from different audio devices with various background noises. We tried controlling

such acoustic factors by introducing the headphone check and instructing them to join the

experiment in a silent place. Further, we screened data according to the screening criteria,

i.e., performance of screening trials and R^2 of the sigmoid approximation. These criteria

were determined arbitrarily, although we confirmed their validity as follows. We designed

screening trials with clearly different interval ratios between standard and comparative

stimuli to detect reliable participants who performed all these trials correctly. The R^2

criteria were useful to remove data that exhibited extraordinary response profiles, e.g., a

linear or U-shape pattern, and were far from the sigmoidal shape. We successfully

removed such data at $R^2 = 0.5$ with keeping a sufficient sample size. These criteria were

rather conservative (we omitted 58.9% of data in total) but necessary to ensure the

reliability of the data. The close match in the sigmoid approximation curves between Exps.

1 and 1' (in-laboratory experiment) suggested further reliability of results obtained from

our online experiments.

Implications for cultural aspects of music

Our study also provides implications for the cultural aspects of music. The findings

suggest that a decrease in perceptual accuracy may also occur in the musical performance

since rhythmic patterns usually consist of multiple musical instruments. For example,

drummers often use hi-hats and ride cymbals with the same center frequency in jazz music,

where the swing ratio is considered an essential factor. The audience may feel more swing

by composing rhythmic patterns with such instruments. However, many other parameters

are involved in the actual rhythmic pattern, such as intensity, accents, and duration. For

more accurate implications, it would be essential to investigate the effects of these factors

on the accuracy of rhythm perception.

A previous study found a relationship between the perceptual accuracy of

interval ratio and nationality [26]. For example, the accuracy of American participants

was high only in the 2:1 ratio, while that of Turkish was high in 2:1 and 3:2, which were

familiar swing ratios for them. This suggests perceptual accuracy may vary depending on

culture-specific listening experiences and acquired musical knowledge. Therefore, it will

be essential to investigate in future studies how the properties of rhythm perception,

observed in the present study, vary with musical experience and belonging to culture to

understand how we acquire rhythm perception.

Our paradigm can also be used to examine the effect of cultural differences on

perceptual accuracy. For example, prior probability distributions over integer ratio on

musical rhythm differ by cultural experience [27]. This may be due to the varying

accuracy of discrimination depending on the cultural experience. A similar online

experiment recruiting participants from different cultural backgrounds may provide

mechanistic insights into the cultural evolution of music.

Conclusion

We found that differences in center frequency in band noise lower the perceptual accuracy

of subdivided rhythm regardless of the rhythmic pattern. This indicates that rhythm

perception is affected not only by temporal structure but also by spectral parameters. Our

findings bridge the gap between rhythm perception and auditory stream segmentation

studies. Our paradigm also has the potential to examine the mechanism of cultural

differences in music.

Acknowledgment

This work was supported by JSPS KAKENHI Grant Number JP20K21804. We thank Dr.

Kazutoshi Kudo for his helpful comments on early manuscripts. Furthermore, we want to

express our gratitude to the colleagues from Okanoya Laboratory who provided insight

and expertise that greatly assisted the research.

Reference

1. Large EW, Palmer C. Perceiving temporal regularity in music. Cognitive Science. 2002;26(1):1–37.

2. Lerdahl F, Jackendoff RS. A Generative Theory of Tonal Music, reissue, with a new preface. MIT Press; 1983. 390 p.

3. Polak R. Rhythmic Feel as Meter: Non-Isochronous Beat Subdivision in Jembe Music from Mali. Music Theory Online [Internet]. 2010 Dec 1 [cited 2022 Aug 9];16(4). Available from: https://mtosmt.org/issues/mto.10.16.4/mto.10.16.4.polak.html

4. Friberg A, Sundströöm A. Swing Ratios and Ensemble Timing in Jazz Performance: Evidence for a Common Rhythmic Pattern. Music Perception. 2002 Mar 1;19(3):333–49.

5. Repp BH. Patterns of expressive timing in performances of a Beethoven minuet by nineteen famous pianists. The Journal of the Acoustical Society of America. 1990 Aug;88(2):622–41.

6. Palmer C. On the Assignment of Structure in Music Performance. Music Perception: An Interdisciplinary Journal. 1996;14(1):23–56.

7. Drake C, Penel A, Bigand E. Tapping in Time with Mechanically and Expressively Performed Music. Music Perception: An Interdisciplinary Journal. 2000;18(1):1–23.

8. Repp BH. Metrical Subdivision Results in Subjective Slowing of the Beat. Music Perception. 2008 Sep 1;26(1):19–39.

9. Handel S. The effect of tempo and tone duration on rhythm discrimination. Perception & Psychophysics. 1993 May 1;54(3):370–82.

10. Fitch WT, Rosenfeld AJ. Perception and Production of Syncopated Rhythms. Music Perception. 2007 Sep 1;25(1):43–58.

11. Ehrlé N, Samson S. Auditory discrimination of anisochrony: Influence of the tempo and musical backgrounds of listeners. Brain and Cognition. 2005 Jun;58(1):133–47.

12. Povel DJ. Internal representation of simple temporal patterns. Journal of Experimental Psychology: Human Perception and Performance. 1981 Feb 1;7(1):3–18.

13. Repp BH. Sensorimotor synchronization: a review of the tapping literature. Psychonomic Bulletin & Review. 2005 Dec 1;12(6):969–92.

14. Repp BH, London J, Keller PE. Perception–production relationships and phase correction in
synchronization with two-interval rhythms. Psychological Research-psychologische Forschung.
2011 May 1;75(3):227–42.

15. Hasuo E, Arao H. Certain non-isochronous sound trains are perceived as more isochronous
when they start on beat. Atten Percept Psychophys. 2020 May;82(4):1548–57.

16. Van Noorden LPAS. Temporal coherence in the perception of tone sequences. PhD thesis,
Eindhoven University of Technology [Internet]. 1975 [cited 2022 Jan 6]; Available from:
https://ci.nii.ac.jp/naid/10012929296/

17. Bregman AS. Auditory Scene Analysis: The Perceptual Organization of Sound. MIT Press;
1990. 800 p.

18. Bregman AS, Ahad PA, Van Loon C. Stream segregation of narrow-band noise bursts.
Perception & Psychophysics. 2001 Jul 1;63(5):790–7.

19. Cusack R, Roberts B. Effects of differences in timbre on sequential grouping. Perception &
Psychophysics. 2000 Jul 1;62(5):1112–20.

20. Henninger F, Shevchenko Y, Mertens UK, Kieslich PJ, Hilbig BE. lab.js: A free, open, online
study builder. Behav Res. 2022 Apr;54(2):556–73.

21. Woods KJP, Siegel MH, Traer J, McDermott JH. Headphone screening to facilitate web-based
auditory experiments. Atten Percept Psychophys. 2017 Oct 1;79(7):2064–72.

22. Frane AV, Shams L. Effects of tempo, swing density, and listener's drumming experience, on
swing detection thresholds for drum rhythms. The Journal of the Acoustical Society of America.
2017 Jun;141(6):4200–8.

23. Nakajima Y, ten Hoopen G, Sasaki T, Yamamoto K, Kadota M, Simons M, et al. Time-
Shrinking: The Process of Unilateral Temporal Assimilation. Perception. 2004 Sep;33(9):1061–
79.

24. Ellis MC. Research Note. Thresholds for Detecting Tempo Change. Psychology of Music. 1991
Oct 1;19(2):164–9.

25. Repp BH. Sensorimotor synchronization and perception of timing: effects of music training and
task experience. Human Movement Science. 2010 Apr 1;29(2):200–13.

26. Hannon EE, Soley G, Ullal S. Familiarity overrides complexity in rhythm perception: A cross-

cultural comparison of American and Turkish listeners. Journal of Experimental Psychology:

Human Perception and Performance. 2012;38(3):543–8.

27. Jacoby N, McDermott JH. Integer Ratio Priors on Musical Rhythm Revealed Cross-culturally

by Iterated Reproduction. Current Biology. 2017 Feb 6;27(3):359–70.