Direct detection of deformation modes on varying length scales in active biopolymer networks

Samantha Stam[1,2,3], Margaret L. Gardel[2,4,5], Kimberly L. Weirich[6]

[1]Biophysical Sciences Graduate Program, University of Chicago, Chicago, IL 60637
[2]Institute for Biophysical Dynamics, University of Chicago, Chicago, IL 60637
[3]Department of Oncological Sciences, University of Utah, Salt Lake City, UT 84112
[4]James Franck Institute, University of Chicago, Chicago, IL 60637
[5]Department of Physics, University of Chicago, Chicago, IL 60637
[6]Department of Materials Science & Engineering, Clemson University, Clemson, SC 29634

Correspondence: samantha.stam@hci.utah.edu, gardel@uchicago.edu, weirich@clemson.edu

ABSTRACT

Correlated flows and forces that emerge from active matter orchestrate complex processes such as shape regulation and deformations in biological cells and tissues. The active materials central to cellular mechanics are cytoskeletal networks, where molecular motor activity drives deformations and remodeling. Here, we investigate deformation modes in contractile actin networks driven by the molecular motor myosin II through quantitative fluorescence microscopy. We examine the deformation anisotropy at different length scales in networks of sparsely cross-linked and bundled actin. In sparsely cross-linked networks, we find myosin-dependent biaxial buckling modes across length scales. Interestingly, both long and short-wavelength buckling may contribute to network contractility. In cross-linked bundled networks, uniaxial contraction predominates on long length scales, while the uniaxial or biaxial nature of the deformation depends on bundle microstructure at shorter length scales. The anisotropy of deformations may provide insight to the mechanical origins of contractility in actin networks and regulation of collective behavior in a variety of active materials.

INTRODUCTION

Active materials are ubiquitous in biology and of increasing interest in engineered materials due to their emergent collective behavior and responsive properties, such as correlated motion, large density fluctuations, and directed force transmission (*1*). These intriguing materials are made of mechanically active building blocks that exert internal forces and produce local motion, resulting in emergent collective phenomena. In biology, active materials emerge from collections of active particles at various length scales, including flocks of birds, suspensions of swarming bacteria, layers of motile cells, and networks of polymers, while engineered active matter includes collections of chemically reactive colloids and mechanically driven particles (*2*). Developing a fundamental understanding of the mechanical properties that emerge from active material microstructure is necessary to characterize functions such as roles within biological tissues and potential for engineered applications such as microrobotics (*3*).

In biological cells, active materials comprise the cytoskeleton, networks and bundles of filament-forming proteins such as actin and microtubules. A primary way that the cytoskeleton is made active is through molecular motors, such as myosin II, that translate the chemical energy of nucleotide hydrolysis into forces that drive filament motion and deformation in actin networks. Myosin II motor heads take force-generating steps, while interacting with multiple actin filaments, resulting in actin filament sliding and buckling that leads to overall network deformation. The activity of motors can also drive relative sliding of filaments and gather filament ends into central locations in a process known as polarity sorting (*4-7*).

These cytoskeletal materials exhibit classic active matter characteristics such as correlated motion (*8, 9*) and large density fluctuations (*8*). However, in contrast to many other active materials, the constituent particles in the cytoskeleton are filaments that are highly deformable and polar, resulting in deformations that depend on orientation and mechanical properties of the filament. For example, actin filaments are semi-flexible polymers, giving rise to an asymmetry in a filament's response to compression relative to extension. This asymmetric mechanical response leads to actin filaments preferentially buckling under load and is hypothesized to be critical to producing the net contractile behavior characteristic of active cytoskeletal networks (*10-15*). Previous experimental and theoretical studies have suggested that the types of network deformation modes, buckling and sliding, are regulated through the network microstructure and activity such as filament rigidity (*16*), cross-linking between filaments (*13, 16-19*), or motor concentration and properties (*11, 13, 20*). These deformation modes underlie differences in emergent network properties, including whether the material is predominately contractile or extensile. To understand the microscopic origin of the active material behavior, most previous experimental studies have relied on indirect assessment of deformation modes via comparison of contraction rate or other metrics to theoretical models (*17, 18*).

Here, we investigate how active network deformation modes depend on length scale and internal stress through quantitative image analysis of deformation anisotropy (*16*) in cytoskeletal networks with different microstructures. We find that uniaxial deformations, which indicate actin filament sliding (*16*), are the predominate deformation mode in networks of semi-flexible actin at sufficiently low internal stress. At higher myosin densities, the internal stress increases, leading to length-scale dependent biaxial buckling

deformations. At intermediate myosin densities, long-wavelength buckling deformations continually rearrange the network persist for tens of minutes without causing network clustering typically observed at higher densities (*12, 21-23*). In networks of cross-linked rigid bundles, by contrast, we observe different patterns of deformation as a function of internal stress and length scale than in networks of filaments. Networks of bundles deform uniaxially on sufficiently large length scales. However, bundles can have different microstructures, where filaments are aligned with the same polarity or with randomly oriented filaments. We find that these different bundle microstructures exhibit either stress-independent (unipolar bundles) or stress and length-scale dependent deformation modes (random polarity bundles). These results reveal how active materials tune deformations at different length scales with microstructure and internal force, emphasizing the importance of considering length scale in building physical models.

METHODS

Protein purification, fluorescent labeling, and storage

Actin, skeletal muscle myosin II, filamin, fascin, and fimbrin were purified as described previously and stored as frozen stocks at -80°C (*16, 24, 25*). Actin was purified from rabbit acetone powder (Pel-Freez Biologicals) and stored 2 mM Tris·HCl, pH 8.0, 0.2 mM ATP, 0.2 mM $CaCl_2$, 0.2 mM DTT, and 0.005% NaN_3 (*26*). Myosin II was prepared from fresh chicken skeletal muscle 5 mM PIPES, pH 7.0, 450 mM KCl (*27*). Filamin was purified from chicken gizzard and stored in 10 mM Tris-HCl, pH 7.4, 120 mM NaCl, 0.5 mM EGTA, 1 mM DTT (*28*). Human fascin (GST tag fusion) and pombe fimbrin (HisTag fusion) were purified from E. coli using plasmids from D. Kovar Lab (University of Chicago) (*29, 30*). Actin and myosin II were fluorescently labeled with tetramethylrhodamine-6-maleimide (TMR, Life Technologies) and Alexa-647 maleimide (Life Technologies), respectively (*31*).

Sample preparation

To prepare microscopy samples, a chamber was constructed from a coverslip passivated with an oil-surfactant layer or lipid bilayer. To prepare an oil-surfacant passivated sample chamber, PFPE-PEG-PFPE surfactant (008; RAN Biotechnologies) was dissolved in a fluorinated oil (Novec-7500 Engineered Fluid; 3M) at a concentration of 2 wt% (*9*). Coverslips (no. 1.5, Fisherbrand) were cleaned by sonication and rinsed in pure water and ethanol. Coverslips were then treated to make hydrophobic by incubating in a 2% vol/vol solution of triethoxy(octyl)silane (440213; Sigma-Aldrich) in isopropanol. The hydrophobic coating was removed from entire coverslip surface except for a 2 mm x 2 mm square region by exposing to UV-ozone for 10 minutes with a square Teflon mask covering the region. This constrained the oil to the square region and prevented flow generated by seeping at the coverslip edges. A glass cylinder was epoxied to the coverslip to produce a microscopy chamber. A minimal volume of oil (~ 3.5 μL) was added to the chamber to coat the surface, and excess solution removed immediately prior to adding the actin polymerization mixture.

To prepare a lipid bilayer passivated sample chamber (*32*), coverslips (no. 1.5; Fisherbrand) were first rinsed with water and ethanol then cleaned through exposure to UV-ozone for 20 min. A glass cylinder was epoxied to the clean coverslip, and vesicle

buffer (10 mM sodium phosphate, pH 7.5, and 140 mM NaCl) added as soon as possible. DOPC vesicles (1,2-dioleoyl-sn-glycero-3-phosphocholine; Avanti Polar Lipids) were then added to a concentration of 100 µM. The coverslip was incubated with the vesicles for 15 min to allow bilayer formation, before the bilayer was rinsed with a buffer comprising of 1x F-buffer (below, in actin polymerization) prior to adding the final sample.

Sparsely cross-linked actin networks

To prepare cross-linked actin networks crowded to a coverslip surface (*12*), a mixture of 0.3 wt% 15 cp methylcellulose, 1 µM actin (1:10 TMR-maleimide labeled:unlabeled), 0.002 µM filamin crosslinker, and an oxygen scavenging system to prevent photobleaching [4.5 mg/mL glucose, 2.7 mg/mL glucose oxidase (345486; Calbiochem), 1,700 units/mL catalase (02071; Sigma), and 0.5 vol/vol % β-mercaptoethanol], was made in 1x F-buffer (10 mM imidazole, pH 7.5, 1 mM $MgCl_2$, 50 mM KCl, 0.2 mM EGTA, and 4 mM ATP). Actin was allowed to polymerize and crowd to the coverslip surface for 30 minutes before addition of dimeric Alexa-647 labeled myosin.

Bundled actin networks

Bundled actin networks were constructed similarly to sparsely cross-linked networks, but have no cross-linker (filamin) during the initial 30 min polymerization period. To prepare fascin cross-linked bundles, fascin was added at a ratio of 1:10 fascin:actin to the polymerized actin sample and incubated for 20 minutes to allow bundle formation. For additional filamin cross-linking, filamin was subsequently added at a ratio of 1:500 filamin:actin and incubated for an additional 20 minutes. For fimbrin bundles, the fimbrin was added 20 minutes after actin polymerization at a ratio of 1:10 and incubated for 20 minutes. Myosin filaments were polymerized separately for 10 minutes in the same buffer conditions described above for actin polymerization before addition to the bundled actin networks.

Image acquisition and analysis

An inverted microscope (Eclipse Ti-E; Nikon) with a spinning-disk confocal head (CSU-X; Yokagawa Electric) was used for all imaging. Imaging was done under magnification of a 40× 1.15 N.A. water-immersion objective (Apo LWD; Nikon) or a 60×/1.49 NA oil immersion objective (Zeiss). Images were acquired with a CMOS camera (Zyla-4.2-USB3; Andor) or a CCD camera (Coolsnap HQ2, Photometrics). Images were acquired every 1 s for Figs. 1-5 and every 5 s for Fig. 6. Calculation of flow vectors and deformation anisotropy was conducted as we described previously (*16*) using PIV software (*33*) and custom Matlab scripts. Values of ρ_{myo} and $P_{biaxial}$ were smoothed such that data points report the average value over overlapping windows of 50 s in fimbrin-bundled sample and 20 s in all other samples.

RESULTS

To create a cytoskeletal material inspired by the cellular cortex, we crowd actin filaments to a thin, dense layer with a depletion agent (see Methods) (*12*). Filamin, an actin cross-linking protein, connects entangled filaments into networks without causing bundling when incorporated at a low concentration (0.01 µM; 1:100 molar ratio with

respect to actin monomer). A thin oil-surfactant layer passivates the coverslip surface against protein adhesion (*9*). To introduce activity, we add myosin II dimers. We define the time the motors are first added to the sample as time t = 0 s. Upon dilution in the sample buffer, which contains lower salt concentration than the storage buffer, the myosin II dimers polymerize into puncta (referred to as myosin, Fig. 1, white puncta), which are filaments that each contain several hundred myosin II motors (*12*).

Myosin incorporates into the sparsely cross-linked network, initially increasing from 0 to 0.05 puncta/μm^2 over the first ~100 s, as myosin polymerizes into elongated puncta that bind to actin filaments (ρ_{myo}, Fig. 1D, open circles). During this initial period of myosin accumulation, there is an increase in the fluctuations along the contour of actin filaments, indicating myosin-induced stresses bending actin filaments. This filament bending induces local filament rearrangement without substantially changing network density (Fig. 1A, 0:00-1:40). As myosin continues to accumulate, the actin and myosin filaments rearrange into clusters with a central myosin core and intervening voids (Fig. 1A, 5:00). This is consistent with previous reports of actomyosin network coarsening *in vitro*, where dense actomyosin clusters form given a sufficiently high myosin concentration (*12, 21-23*). As a result of clustering, the number density of myosin puncta begins to decrease near t = 100 s (Fig. 1D).

We previously found that cross-linked networks of unipolar rigid bundles contract via uniaxial deformations arising from relative sliding of the bundles (*16*). These networks are constructed with two different actin cross-linkers. One type of cross-linker, fascin, rigidly links actin filaments together into bundles with a uniform polarity (*34*). We observe that these bundles appear as straight rods with diffraction limited width that may overlap with other bundles (Fig. 1B, (*16, 24*)). We then add a flexible cross-linker, filamin, after the fascin-mediated bundles form to ensure that the bundles form an inter-connected network. Effectively, the unipolar bundles act as polar filaments with a higher stiffness than the unbundled, semi-flexible actin (*24*). The unipolar bundle networks form asters upon addition of myosin (Fig. 1B, 1:40, (*16*)). As with the sparsely cross-linked network in Fig. 1A, the density of myosin puncta initially increases as myosin accumulates in the network, and then decreases as clusters form (Fig 1D, open squares).

To more broadly understand the influence of actin bundle structure on network deformation modes, we use a different rigid actin cross-linker, fimbrin, to make actin bundles. These bundles have similar packing of the actin filaments as fascin bundles (*35*), and therefore might also contract uniaxially. In contrast to fascin bundles, within fimbrin bundles are aligned with random polarity (*36*). Myosin filaments can simultaneously bind to actin filaments with opposite polarity and build internal stress within a bundle (*24*).

Myosin activity induces the random polarity bundle network to coarsen into dense actomyosin clusters (Fig. 1C), similar to the unipolar bundle network. However, the intermediate stages of network reorganization are distinct from the unipolar bundle network. While the unipolar bundles appear rigid, there is visual evidence of buckling or bending of the random polarity bundles (Fig. 1C, 1:40). Similar to the previous networks, the myosin puncta density initially rises and then falls during network clustering (Fig 1D, filled squares (*16*)).

The average divergence of actin displacement vectors calculated by particle imaging velocimetry (PIV, *Materials and Methods*) has been used previously to quantify contractility in actomyosin networks (*20*). The typical behavior of the average divergence

over time is a decrease to negative (contractile) values as myosin accumulates and contracts the network followed by a return to near zero values as the network forms clusters (*16, 20*). The bundled networks from Fig. 1B and 1C both display this expected behavior with minimum values of the divergence (Fig. 1E, orange open squares and blue closed squares) reached near the respective myosin puncta density peaks in Fig. 1D. However, the average divergence of the sparsely cross-linked sample oscillates near zero throughout most of the network rearrangement and only decreases to consistently negative values long after network clustering begins (Fig. 1E, black open circles). This is surprising given that the buckling/compression of filaments observed in Fig. 1A is expected to produce a negative divergence (*16, 20*).

To obtain a better understanding of differences in the contraction of these networks, we develop a metric for contraction on different length scales. We split the actin images into boxes of size s which contain local PIV vectors, $\vec{v}_{act}(\vec{r})$ (Fig. 2A). The contraction of each box may be characterized by the contractile moment:

$$Tr(M_{ij})/s = \sum_{\vec{r}} \vec{v}_{act}(\vec{r}) \cdot \frac{\vec{r}}{s}$$

where $\vec{v}_{act}(\vec{r})$ is the PIV vector located at position $\vec{r}$ from the center of the box and M_{ij} is the flow dipole moment tensor (*16*). This has been rescaled from our previous quantification of contraction (*16*) to facilitate comparison of contractile moments with different s (length scale) values.

At a sufficiently large length scale, a decrease of $<Tr(M_{ij})/s>$ to negative (contractile) values during the initial stages of contraction occurs in the sparsely cross-linked network (Fig. S1A). To explain the trends revealed by the average contractile moment, we visualize contraction on mesoscopic length scales by calculating $Tr(M_{ij})/s$ with $s = 50$ µm for overlapping boxes and plotting the value using the color scales in Fig. 2B and Fig. S2. In all networks, large areas with contractile (red) $Tr(M_{ij})/s$ values are separated by regions of extensile (blue) $Tr(M_{ij})/s$ values. Interestingly, these contractile regions begin developing at early time points and low myosin densities and increase in intensity as the contraction proceeds (Fig. S2).

Given that the deformation modes of these networks are expected to vary between filament buckling and sliding (*16*), we anticipate that the contractility on different length scales will depend on increasing internal stress in divergent ways. To examine this possibility, we plot the average contractile moment in non-overlapping boxes against ρ_{myo} during the myosin accumulation stage (as defined Fig. 1D). In the sparsely cross-linked network, $<Tr(M_{ij})/s>$ at $s=50$ µm begins decreasing to negative values near $\rho_{myo} =0.01$ µm^{-2} before oscillating slightly between $\rho_{myo}=0.02$ µm^{-2} and 0.04 µm^{-2} (Fig. 2C, green open circles). A steeper decrease then occurs near $\rho_{myo}=0.04$ µm^{-2} followed by a return to somewhat less negative values at $\rho_{myo}=0.05$ µm^{-2}. The magnitude of $<Tr(M_{ij})/s>$ at $s=10$ µm is smaller and increases to slightly positive values at a similar ρ_{myo} as the decrease in $<Tr(M_{ij})/s>$ at $s=50$ µm (Fig. 2C, open black squares).

The changes of $<Tr(M_{ij})/s>$ with increasing ρ_{myo} differ in the bundled networks. In the unipolar bundle network, the magnitude of $<Tr(M_{ij})/s>$ is also smaller for the shorter length scale, $s=10$ µm, than $s=50$ µm (Fig. 2D). However, the trends of the curves at both length scales are similar, with both starting at initially positive values before decreasing to negative ones. This decrease continues after the peak in myosin density (Fig. S1B).

Similarly, the curves for $<Tr(M_{ij})/s>$ versus ρ_{myo} at longer and shorter length scales in the random polarity bundled network have different magnitudes but otherwise resemble one another closely (Fig. 2E, $s=6.5$ μm, open black squares and $s=45$ μm, open green circles). In this network, both remain near zero until they decrease to negative values when ρ_{myo} exceeds approximately 0.03 μm^{-2}. Overall, the trends of contractility with internal stress and length scale are very different between these networks.

To characterize the origin of contractility in terms of filament sliding versus buckling deformation modes, we diagonalize the flow dipole moment tensor, M_{ij}, of boxes with varying s (16). The resulting eigenvalues, M_{min} and M_{max}, characterize the local deformation anisotropy (Fig. 3A). A value of M_{min}/M_{max} of 0 indicates that the deformation of actin network within the defined box is along a single axis. At the other extreme, a value of M_{min}/M_{max} of 1 indicates that the deformation is completely biaxial. We superimpose ellipses on the actin images with minor and major axes M_{min} and M_{max} respectively to visualize the anisotropy of deformations. For $s = 20$ μm in the sparsely cross-linked network, the size of the ellipses (magnitudes of M_{min} and M_{max}) increases between time 0:00 to 1:40 (Fig. 3B), indicating an increase in extent of deformation as myosin accumulates in the network. At a smaller length scale, $s = 5$ μm, we also find that M_{min} and M_{max} increase as myosin accumulates (Fig 3C).

The deformation anisotropy also changes as the sparsely cross-linked network reorganizes. Deformations are predominantly uniaxial when myosin density in the network is initially low (left panels in Fig. 3B and 3C, quantified in Fig. 3D and 3E, blue curves). This differs from our previously reported data of networks with greater initial myosin density, which exhibited biaxial deformations arising from filament buckling (16). As network deformation proceeds, corresponding to increased myosin accumulation, we observe a large shift in the probability distribution of M_{min}/M_{max} at $s = 20$ μm. The probability of biaxial deformations ($M_{min}/M_{max}>0.5$) increases such that the distribution becomes peaked at $M_{min}/M_{max} \sim 0.8$ (Fig. 3D). At a smaller length scale ($s = 5$ μm, Fig. 3E), the distribution also shifts to more biaxial deformations after the addition of myosin, but distribution of M_{min}/M_{max} always remains more uniaxial and the changes over time are smaller. At long times on both length scales, the distribution eventually shifts back to a distribution with fewer biaxial deformations.

To understand how these patterns of deformation give rise to the differences in contractility seen in Fig. 2, we define $P_{biaxial}$ as the proportion of deformations in the network at a given time and s value with M_{min}/M_{max} values greater than 0.5. We then plot $P_{biaxial}$ as a function of ρ_{myo} during the myosin accumulation stage similarly to the plots of $<Tr(M_{ij})/s>$ in Figs. 2C-2E. We find that $P_{biaxial}$ with $s = 5$-10 μm (Fig. 4A, open black squares) is consistently lower than $P_{biaxial}$ with $s=45$-50 μm (Fig. 4A, open green circles) in the sparsely cross-linked network at all values of ρ_{myo}. At both length scales, $P_{biaxial}$ is initially constant over a range of low ρ_{myo}, but sharply increases above a threshold ρ_{myo}. The increase in $P_{biaxial}(s)$ at $s = 5$-10 μm occurs at a higher value of ρ_{myo}, which is consistent with a higher internal stress requirement for small wavelength buckling modes. Also consistent with this stress requirement, the value of ρ_{myo} required to raise $P_{biaxial}$ by

20% of its baseline value ($\rho_{activate}$) decreases with s in an average of four samples (Fig. S3).

Intriguingly, the sharp increase of the long-wavelength biaxial buckling modes in Fig. 4A occurs at approximately the same ρ_{myo} value as the initial decrease of <Tr(M_{ij})/s> with s=50 μm in Fig. 2C ($\rho_{myo} \approx 0.02$ μm^{-2}). This suggests that long-wavelength buckling starts contributing to contractility before the short-wavelength buckling begins. The additional decrease of <Tr(M_{ij})/s> with s=50 μm at $\rho_{myo} \approx 0.04$ μm^{-2} occurs while $P_{biaxial}$ (s=5-10 μm) is increasing and $P_{biaxial}$ (s=45-50 μm) is reaching a plateau. However, the value of <Tr(M_{ij})/s> at s=10 μm in Fig. 2C remains slightly extensile even during the increase of $P_{biaxial}$ at the shorter length scale.

The unipolar bundle network (Fig. 4B) similarly has a baseline value of $P_{biaxial}$ that is lower for s = 5-10 μm (open black squares) than s = 45-50 μm (open green circles). Unlike the sparsely cross-linked network in Fig. 4A however, there is no sharp increase in $P_{biaxial}$ with increasing ρ_{myo}. The difference of $P_{biaxial}$ from its initial value to its extremum at or before the peak in ρ_{myo}, $\Delta P_{biaxial}$, remains similarly near zero over a range of s (Fig. S4). Therefore, the uniaxial sliding deformations underlying contractility in this network (16) dominate the deformation anisotropy at both length scales.

For the random polarity bundle network, a different pattern of length-scale dependent changes in $P_{biaxial}$ with increasing myosin density is observed (Fig. 4C). At a smaller value of s=3-7 μm (open black squares), $P_{biaxial}$ begins increasing at $\rho_{myo} \approx 0.04$ μm^{-2} while at s=42-46 (open green circles), $P_{biaxial}$ decreases. This suggests that buckling of bundles occurs locally, but the bundles remain sufficiently intact to transmit anisotropic deformations on longer length scales, which is consistent with the images in Fig. 1C. The value of $\Delta P_{biaxial}$ crosses over from positive to negative values at s > 25 μm (Fig. S4).

Previous studies have suggested that contraction may originate at the microscale in proximity to myosin filaments and propagate outward into the network (15, 19). However, the sparsely cross-linked network appears to contrast with this expectation given that the onset of buckling with increasing ρ_{myo} first occurs at higher s. To specifically inhibit the buckling modes at small s and examine the influence on contractility, we investigate networks with lower myosin densities. In a sparsely cross-linked networks with ~ 5 times lower ρ_{myo} than the network in Fig. 1A (Fig. 5A, filled squares), we observe no coarsening of the actin network even after 15 minutes of myosin activity (Fig. 5B). When we investigate network with a slightly increased ρ_{myo}, (Fig. 5A, open circles), we observe a restoration of some actin coarsening at t=15:00. (Fig. 5C). Clustering of the myosin occurs in both networks as indicated by the decreases in ρ_{myo} following their peaks in Fig. 5A. Large regions with contractile Tr(M_{ij})/s values are present in both samples (Fig 5D, red areas) but are lower magnitude than the data in Fig. 2B.

We repeat our analysis of <Tr(M_{ij})/s> with s=50 μm in these samples to assess how contractility varies with deformation mode differences. In the sample with the lowest ρ_{myo}, <Tr(M_{ij})/s> (s=50 μm) remains mostly extensile and oscillates with increasing ρ_{myo} (Fig. 5E, left panel). This contrasts with the decrease to negative values seen in Fig. 2C. In the intermediate ρ_{myo} sample, the decrease to negative <Tr(M_{ij})/s> with increasing ρ_{myo} is qualitatively restored (Fig. 5E, right panel), though the minimum value is somewhat less negative than Fig. 2C. Interestingly, the small decrease between ρ_{myo}=0.01 μm^{-2} and ρ_{myo}=0.02 μm^{-2} and the subsequent steeper decrease at ρ_{myo}>0.02 μm^{-2} resembles the trend in Fig. 2C.

Examining $P_{biaxial}$ during these changes in $<Tr(M_{ij})/s>$ reveals potential contributions of individual deformation modes to contractility. In the lowest ρ_{myo} network, $P_{biaxial}$ (s=45-50 μm) sharply increases near ρ_{myo} = 0.01 μm^{-2} while only a small increase in $P_{biaxial}$ (s=5-10 μm) occurs (Fig. 5F, left panel). In the intermediate ρ_{myo} network, the trends of both $P_{biaxial}$ curves are initially similar, but both reach slightly higher peak values (Fig. 5F, right panel). Similar to the high ρ_{myo} network in Fig. 2C and 4A, the initial decrease of $<Tr(M_{ij})/s>$ in the intermediate ρ_{myo} network occurs during the increase in $P_{biaxial}$ (s=45-50 μm) between ρ_{myo} = 0.01 μm^{-2} and ρ_{myo} =0.02 μm^{-2} (Fig. 5E and Fig. 5F, right panels). This suggests that long wavelength buckles are again contributing to contractility. However, $<Tr(M_{ij})/s>$ does not decrease during the rise of $P_{biaxial}$ (s=45-50 μm) in the low ρ_{myo} network (Fig. 5E and Fig. 5F, left panels). Similar maximum values of $P_{biaxial}$ (s=45-50 μm) happen in the low ρ_{myo} network compared to the more contractile networks, but they require less myosin and are reached at ~700 s versus ~100 s after myosin addition (Fig. S5). Thus, buckling at low ρ_{myo} might be mechanically different and less contractile and/or contractile buckling is counterbalanced by extensile deformations.

DISCUSSION

Deformations in active, soft materials are critical during processes requiring changes of material shape, size, or transport of components. The mode of deformation at the microscale can include both translocation of components with respect to one another and deformation of components, regulating material scale mechanical responses such as contraction or expansion (*13*), flow-induced transport (*9, 37*), or force transmission to external physical attachments such as cellular adhesions. To understand how microscopic material structure influences deformations and mechanical response, substantial efforts have been made connecting theoretical and computational models of material deformation to available experimentally observable quantities (*38*). Here, we build on our previous work in which deformation anisotropy was analyzed at a single length scale to identify actin buckling and sliding (*16*) to bridge the gap between experiments in which deformation modes typically must be inferred from indirect measurements and modeling in which the deformations on all length scales are directly accessible. For example, our analysis may be useful for testing models for how energy is injected and dissipated on varying length scales in active matter systems (*39*). While models describing contraction as propagating out from a local buckling deformation and being amplified by the network (*14, 40*) may be appropriate for some contracting materials, to our knowledge there is currently no model predicting whether buckling can cascade to smaller length scales. Such a possibility may be consistent with the sparsely cross-linked network deformations we observe here and is an interesting topic for future studies.

Analysis of deformation modes may also clarify contraction mechanisms in the more complex, composite cytoskeletal materials inside of living cells. Intracellular actin networks include disordered meshworks, branched networks that may inhibit contractility (*41, 42*), and bundles that may regulate myosin-II force magnitudes (*24*) and transmit contractile forces to other actin structures (*43*). As a result of these actin architectures, multiple deformations modes could exist within the same cell. Strategies to label actin

and detect flow in cells (*44*) are likely adaptable to linking specific modes with contractility similarly to the analysis performed here. Downstream analysis could examine co-regulatory roles of actin binding proteins and specific contractile deformations. For example, small versus large-wavelength buckling could lead to filament severing (*12, 15*) and affect the activity of actin polymerizing proteins that act on the generated filament barbed ends. Force transmission also occurs between different filaments of the cytoskeleton, with microtubules potentially becoming bent or broken in response to contractile forces generated by actin networks (*45, 46*). Additionally, given the presence of feedback between localized signaling and contractility (*47-49*), future studies could also establish actin deformations as indicators of proximal signaling dynamics.

Force transmission may also occur in organelles outside of the cytoskeleton, and analysis of the resulting deformations could be used as a step toward understanding when, how, and why forces are transmitted in these active materials. For example, recent research has indicated that organelle-organelle contacts are tethers and potentially capable of exerting forces to divide mitochondria (*50, 51*). This suggests that force transmission occurring at these sites may be important for maintaining normal organelle structure and distribution. Altering the physical structure of mitochondria via deformation and translocation may influence their metabolism (*52*) and, consequently, their ability to produce energy or induce detrimental effects that lead to cell aging or death. As another example, mechanosensitive processes such as the opening of ion channels may be related to deformations in both cells and the fibrous networks of the extracellular matrix (*53*). More generally, analysis of deformation modes in cells could function as an accessory to omics-type approaches that help link individual or sets of molecules to physiological processes even when the specific roles of the deformations are unknown. Development of additional tools to track organelle motions with comparable resolution as is possible with cytoskeletal filaments may be necessary for this analysis.

ACKNOWLEDGEMENTS
We thank members of David Kovar's laboratory, particularly J. Christensen, J. Winkelman, C. Suarez, and A. Harker for the purified protein cross-linker fimbrin, as well as the construct for fascin and advice on purifying proteins. This research was partially supported by the University of Chicago MRSEC, funded through the National Science Foundation award DMR-2011854. This research was also supported in part by the National Science Foundation EPSCoR Program under NSF Award # OIA-1655740 and the National Institutes of Health National Institute of Biomedical Imaging and Bioengineering Training grant number T32EB009412.

REFERENCES

1.   G. Gompper *et al.*, The 2020 motile active matter roadmap. *Journal of Physics: Condensed Matter* **32**, 193001 (2020).

2.	M. C. Marchetti *et al.*, Hydrodynamics of soft active matter. *Reviews of Modern Physics* **85**, 1143.

3.	H. Jia *et al.*, 3D printed protein-based robotic structures actuated by molecular motor assemblies. *Nature Materials* **21**, 703-709 (2022).

4.	F. J. Nedelec, T. Surrey, A. C. Maggs, S. Leibler, Self-organization of microtubules and motors. *Nature* **389**, 305-308 (1997).

5.	R. Urrutia, M. A. McNiven, J. P. Albanesi, D. B. Murphy, B. Kachar, Purified Kinesin Promotes Vesicle Motility and Induces Active Sliding between Microtubules Invitro. *Proceedings of the National Academy of Sciences of the United States of America* **88**, 6701-6705 (1991).

6.	V. Wollrab *et al.*, Polarity sorting drives remodeling of actin-myosin networks. *Journal of Cell Science* **132**, jcs219717 (2018).

7.	A. Sciortino, A. R. Bausch, Pattern formation and polarity sorting of driven actin filaments on lipid membranes. *Proceedings of the National Academy of Sciences* **118**, e2017047118 (2021).

8.	V. Schaller, C. Weber, C. Semmrich, E. Frey, A. R. Bausch, Polar patterns of driven filaments. *Nature* **467**, 73-77 (2010).

9.	T. Sanchez, D. T. Chen, S. J. DeCamp, M. Heymann, Z. Dogic, Spontaneous motion in hierarchically assembled active matter. *Nature* **491**, 431-434 (2012).

10.	T. B. Liverpool, M. C. Marchetti, J. F. Joanny, J. Prost, Mechanical response of active gels. *EPL (Europhysics Letters)* **85**, 18007 (2009).

11.	M. Lenz, T. Thoresen, M. L. Gardel, A. R. Dinner, Contractile Units in Disordered Actomyosin Bundles Arise from F-Actin Buckling. *Physical Review Letters* **108**, (2012).

12.	M. P. Murrell, M. L. Gardel, F-actin buckling coordinates contractility and severing in a biomimetic actomyosin cortex. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 20820-20825 (2012).

13.	J. M. Belmonte, M. Leptin, F. Nédélec, A theory that predicts behaviors of disordered cytoskeletal networks. *Mol Syst Biol* **13**, 941 (2017).

14.	P. Ronceray, C. P. Broedersz, M. Lenz, Fiber networks amplify active stress. *Proceedings of the National Academy of Sciences* **113**, 2827-2832 (2016).

15.	S. K. Vogel, Z. Petrasek, F. Heinemann, P. Schwille, Myosin motors fragment and compact membrane-bound actin filaments. *Elife* **2**, (2013).

16.	S. Stam *et al.*, Filament rigidity and connectivity tune the deformation modes of active biopolymer networks. *Proceedings of the National Academy of Sciences* **114**, E10037 (2017).

17.	H. Ennomani *et al.*, Architecture and Connectivity Govern Actin Network Contractility. *Curr Biol* **26**, 616-626 (2016).

18.	K. T. Stanhope, V. Yadav, C. D. Santangelo, J. L. Ross, Contractility in an extensile system. *Soft Matter* **13**, 4268-4277 (2017).

19.	R. Blackwell *et al.*, Microscopic origins of anisotropic active stress in motor-driven nematic liquid crystals. *Soft Matter* **12**, 2676-2687 (2016).

20.	M. Murrell, M. L. Gardel, Actomyosin sliding is attenuated in contractile biomimetic cortices. *Molecular Biology of the Cell* **25**, 1845-1853 (2014).

21.	M. Soares e Silva *et al.*, Active multistage coarsening of actin networks driven by myosin motors. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 9408-9413 (2011).

22.	F. Backouche, L. Haviv, D. Groswasser, A. Bernheim-Groswasser, Active gels: dynamics of patterning and self-organization. *Phys Biol* **3**, 264-273 (2006).

23.	S. Koehler, V. Schaller, A. R. Bausch, Structure formation in active networks. *Nature Materials* **10**, 462-468 (2011).

24.	K. L. Weirich, S. Stam, E. Munro, M. L. Gardel, Actin bundle architecture and mechanics regulate myosin II force generation. *Biophys J* **120**, 1957-1970 (2021).

25.	K. L. Weirich *et al.*, Liquid behavior of cross-linked actin bundles. *Proceedings of the National Academy of Sciences* **114**, 2131-2136 (2017).

26.	J. Spudich, S. Watt, Regulation of Rabbit Skeletal Muscle Contraction. 1. Biochemical Studies of Interactions of Tropomyosin-Troponin Complex with Actin and Proteolytic Fragments of Myosin. *Journal of Biological Chemistry* **246**, 4866-& (1971).

27.	S. S. Margossian, S. Lowey, Preparation of Myosin and Its Subfragments from Rabbit Skeletal-Muscle. *Methods in Enzymology* **85**, 55-71 (1982).

28.	S. Craig, C. Lancashire, J. Cooper, Preparation of Smooth-Muscle Alpha-Actinin. *Methods in Enzymology* **85**, 316-321 (1982).

29.	D. Vignjevic, J. Peloquin, G. G. Borisy, in *Regulators and Effectors of Small GTPases: Rho Family*. (Academic Press, 2006), vol. Volume 406, pp. 727-739.

30.	C. T. Skau, D. R. Kovar, Fimbrin and tropomyosin competition regulates endocytosis and cytokinesis kinetics in fission yeast. *Curr Biol* **20**, 1415-1422 (2010).

31.	A. B. Verkhovsky, G. G. Borisy, Non-Sarcomeric Mode of Myosin-Ii Organization in the Fibroblast Lamellum. *Journal of Cell Biology* **123**, 637-652 (1993).

32.	K. Weirich, J. Israelachvili, D. Fygenson, Bilayer Edges Catalyze Supported Lipid Bilayer Formation. *Biophysical Journal* **98**, 85-92 (2010).

33.	K.-A. Chang, N. Mori. (2008).

34.	J. R. Bartles, Parallel actin bundles and their multiple actin-bundling proteins. *Current opinion in cell biology* **12**, 72-78 (2000).

35.	A. Bretscher, Fimbrin Is a Cytoskeletal Protein That Crosslinks F-Actin In vitro. *Proceedings of the National Academy of Sciences of the United States of America* **78**, 6849-6853 (1981).

36.	C. T. Skau *et al.*, Actin Filament Bundling by Fimbrin Is Important for Endocytosis, Cytokinesis, and Polarization in Fission Yeast. *Journal of Biological Chemistry* **286**, 26964-26977 (2011).

37.	N. Kumar, R. Zhang, J. J. de Pablo, M. L. Gardel, Tunable structure and dynamics of active liquid crystals. *Science Advances* **4**, eaat7779.

38.	S. Banerjee, M. L. Gardel, U. S. Schwarz, The Actin Cytoskeleton as an Active Adaptive Material. *Annual Review of Condensed Matter Physics* **11**, 421-439 (2020).

39.	R. Alert, J. Casademunt, J.-F. Joanny, Active Turbulence. *Annual Review of Condensed Matter Physics* **13**, 143-170 (2022).

40.	J. Notbohm, A. Lesman, P. Rosakis, D. A. Tirrell, G. Ravichandran, Microbuckling of fibrin provides a mechanism for cell mechanosensing. *Journal of The Royal Society Interface* **12**, 20150320 (2015).

41.	Q. Yang, X.-F. Zhang, T. D. Pollard, P. Forscher, Arp2/3 complex–dependent actin networks constrain myosin II function in driving retrograde actin flow. *Journal of Cell Biology* **197**, 939-956 (2012).

42.	C. G. Muresan *et al.*, F-actin architecture determines constraints on myosin thick filament motion. *Nature Communications* **13**, 7008 (2022).

43.	T. Vignaud *et al.*, Stress fibres are embedded in a contractile cortical network. *Nature Materials* **20**, 410-420 (2021).

44.	G. Danuser, C. M. Waterman-Storer, Quantitative fluorescent speckle microscopy of cytoskeleton dynamics. *Annu Rev Biophys Biomol Struct* **35**, 361-387 (2006).

45.	N. Wang *et al.*, Mechanical behavior in living cells consistent with the tensegrity model. *Proc Natl Acad Sci U S A* **98**, 7765-7770 (2001).

46.   C. M. Waterman-Storer, E. D. Salmon, Actomyosin-based retrograde flow of microtubules in the lamella of migrating epithelial cells influences microtubule dynamic instability and turnover and is associated with microtubule breakage and treadmilling. *J Cell Biol* **139**, 417-434 (1997).

47.   M. F. Staddon, E. M. Munro, S. Banerjee, Pulsatile contractions and pattern formation in excitable actomyosin cortex. *PLOS Computational Biology* **18**, e1009981 (2022).

48.   M. Nishikawa, S. R. Naganathan, F. Jülicher, S. W. Grill, Controlling contractile instabilities in the actomyosin cortex. *eLife* **6**, e19595 (2017).

49.   J. B. Michaux, F. B. Robin, W. M. McFadden, E. M. Munro, Excitable RhoA dynamics drive pulsed contractions in the early C. elegans embryo. *Journal of Cell Biology* **217**, 4230-4252 (2018).

50.   M. Bohnert, Tether Me, Tether Me Not—Dynamic Organelle Contact Sites in Metabolic Rewiring. *Developmental Cell* **54**, 212-225 (2020).

51.   J. R. Friedman *et al.*, ER tubules mark sites of mitochondrial division. *Science* **334**, 358-362 (2011).

52.   M. Roy, P. H. Reddy, M. Iijima, H. Sesaki, Mitochondrial division and fusion in metabolism. *Current Opinion in Cell Biology* **33**, 111-118 (2015).

53.   Y.-C. Chuang, C.-C. Chen, Force From Filaments: The Role of the Cytoskeleton and Extracellular Matrix in the Gating of Mechanosensitive Channels. *Frontiers in Cell and Developmental Biology* **10**, (2022).

FIGURE CAPTIONS

Figure 1: Actomyosin network contractility is dependent on filament cross-linking and bundling. (A) Fluorescence microscopy images of a network of unbundled actin filaments (red) sparsely cross-linked with the flexible cross-linker filamin as myosin (white) accumulates over time. (B) Images of unipolar actin bundles (red) formed with a rigid actin cross-linker (fascin) and then linked together with the flexible cross-linker filamin as myosin (white) accumulates. (C) Images of actin bundles (red) with random actin polarity formed by the cross-linker fimbrin as myosin (white) accumulates. In each image sequence, the myosin is added to the sample at time 0:00 and subsequently polymerizes into punctate filaments. Scale bars are 10 µm. (D) Myosin filament puncta density, ρ_{myo}, plotted against time after myosin addition, for the sparsely cross-linked (black open circles), unipolar bundles (orange open squares) and random polarity bundle (filled blue squares) samples. (E) Divergence as a function of time for representative samples of sparsely cross-linked (open black circles), unipolar bundle (open orange squares), and random polarity bundle (filled blue squares) actin networks.

Figure 2: Contractility is dependent on actin network architecture and length scale. (A): Particle imaging velocimetry generates vectors describing the local material flow in an actin network. The network images are separated into boxes of side length s, and the contractility of each box is quantified using the contractile moment. (B): Contractile moment values of overlapping boxes in the sparsely cross-linked network during contraction. The color represents $\text{Tr}(M_{ij})/s$ of a box with s = 50 µm centered at each position. (C): Spatial average of $\text{Tr}(M_{ij})/s$ in non-overlapping boxes with s = 50 µm (open green circles) and s = 10 µm (open black squares) in sparsely cross-linked network. (D): Spatial average of $\text{Tr}(M_{ij})/s$ in non-overlapping boxes with s = 50 µm (open green circles) and s = 10 µm (open black squares) in unipolar bundle network. (E): Spatial average of $\text{Tr}(M_{ij})/s$ in non-overlapping boxes with s = 45 µm (open green circles) and s = 6.5 µm (open black squares) in random polarity bundle network.

Figure 3: Anisotropy analysis reveals varying deformation mechanisms in contractile networks. (A): The deformation anisotropy of each box of PIV vectors is quantified using the flow dipole tensor and represented by an ellipse. (B): Deformation anisotropy ellipse images in sparsely cross-linked network at length scale s = 20 µm. (C): Deformation anisotropy ellipse images in same network at s = 5 µm. (D): Distributions of the deformation anisotropy, M_{min}/M_{max}, at varying time points with s = 20 µm. (E): Distributions of M_{min}/M_{max} with s = 5 µm. Scale bars in (B) and (D) are 10 µm.

Figure 4: Length-scale dependent activation of deformation modes varies for different actin networks. $P_{biaxial}(s)$, defined as the fraction of M_{min}/M_{max} values greater than 0.5 as a function of myosin puncta density during the myosin accumulation stage for (A) sparsely cross-linked networks, with s = 5-10 µm (open black squares) and s = 45-50 µm (open green circles); (B) unipolar bundle networks, with s = 5-10 µm (open black squares) and s = 45-50 µm (open green circles); and (C) random polarity bundle networks, with s = 3-7 µm (open black squares) and s = 42-46 µm (open green circles).

Figure 5: Reducing the myosin density in sparsely cross-linked actin networks lowers the frequency and net contractility of buckling. (A) Myosin filament puncta density, ρ_{myo}, plotted against time after myosin addition in two sparsely cross-linked networks with low ρ_{myo} (open circles) and intermediate ρ_{myo} (filled squares). (B) Image of actin in low ρ_{myo} network after 15:00 of myosin activity. (C) Image of actin in intermediate ρ_{myo} network after 15:00 of myosin activity. (D) Heat maps of $Tr(M_{ij})/s$ in overlapping, 50 μm boxes in low ρ_{myo} (left) and intermediate ρ_{myo} (right) networks. (E) Spatial average of $Tr(M_{ij})/s$ with non-overlapping boxes, s = 50 μm, in low ρ_{myo} (left) and intermediate ρ_{myo} (right) networks. (F) $P_{biaxial}$ with s = 45-50 μm (green circles) and s = 5-10 μm (black squares) in low ρ_{myo} (left) and intermediate ρ_{myo} (right) networks.

Figure 1

Figure 2

A *length scale s*

PIV vectors, $\vec{v}_{act}$

Contractile moment, $Tr(M_{ij})/s$

B $Tr(M_{ij})/s$ (μ/s)

C $\langle Tr(M_{ij})/s\rangle$ (10^{-3} μ/s) $\langle Tr(M_{ij})/s\rangle$ (μ/s)

D $\langle Tr(M_{ij})/s\rangle$ (10^{-2} μ/s) $\langle Tr(M_{ij})/s\rangle$ (μ/s)

E $\langle Tr(M_{ij})/s\rangle$ (10^{-3} μ/s) $\langle Tr(M_{ij})/s\rangle$ (μ/s)

ρ_{myo} (μm^{-2})

Figure 3

Figure 4

Figure 5