Tree Species Identity Drives Soil Carbon and Nitrogen Stocks in Nutrient-Poor Sites

Estela Covre Foltran[1] , Norbert Lamersdorf [1]

[1] Soil Science of Temperate Ecosystems - University of Göttingen

Author contact: estelacfoltran@gmail.com

Abstract

The establishment of mixed forest stands can be seen as an option to improve soil organic

carbon and N storages and to protect forest ecosystems from various impacts of climate

change. We analyzed groups of pure mature European beech (*Fagus sylvatica*), Douglas fir

(*Pseudotsuga menziesii*) and Norway spruce (*Picea abies*) stands as well as mixtures of

beech with either Douglas fir or spruce along Northern Germany. To understand the potential

effects of species mixtures on carbon and nitrogen storage, we examined the content and

stocks of organic carbon and nitrogen in the O-horizon (organic layer) and mineral soil. The

C stock of the organic layer was largely depending on tree species, whereas the C stock of the

mineral soil varied among soil types. Total soil organic C (SOCt) stocks (Organic horizon +

0-30 cm soil depth) showed significant species identities and mixing effects were most

obvious due to the high contribution of the SOC stocks in the organic layer (SOCo) to total

SOCt. These contribution was site-dependent. Overall, under sandy soil conditions, conifers

and mixed forests allocated 10% more SOC and N at the organic layer compared to loamy

soils, whereas the SOC and N stocks under beech maintained the same proportion,

independent of the site condition.

The interaction between species and sites was significant only for Douglas Fir and mixed

Douglas Fir/beech stands, indicating that the effect of species on carbon and nitrogen varied

across sites, being significantly high at sandy soils.

In summary, the higher potential for carbon storage in mixed-species forests compared to

pure stands, along with the shift of carbon into more stable pools (such as from the Ol layer

to the Oh layer), emphasizes the capacity of mixed forest to provide valuable ecosystem

services while reducing the risk of unintended carbon losses.

Keywords: mixed forests; broadleaves; conifers; *Fagus sylvatica; Pseudotsuga menziesii;*

Picea abies; SOC.

Introduction

Forest soils are complex ecosystems and their management, especially through the selection

of trees with distinct species identities, may have a major impact on soil organic carbon

(SOC) and nitrogen (N) dynamics and sequestration potentials (Jandl et al., 2007; Vesterdal

et al., 2008).

Typical driving forces of such tree species identities include the type of foliage, the quality of

leaf litter, indicated e.g. by the C/N ratio, or the type of rooting characteristic in forest soils.

Thus tree species identity may have an important impact on the ecosystem level such as soil

C stock and C/N ratios, particularly in the O-horizon and top mineral soil layers (Dawud et

al., 2017; Vesterdal et al., 2013). Furthermore, the species diversity drives the fluxes of

carbon and nitrogen in soils (Cepáková et al., 2016; Mueller et al., 2012; Vesterdal et al.,

2002) and their role on SOC sequestration are been highlighted to improve the ecosystem

functions and to cope with climate change scenarios.

Different patterns of C storage in soil profile and organic horizons has been reported for

conifers and deciduous trees on temperate ecosystems (Bolte and Villanueva, 2006) .

However, it is highly uncertain which processes are responsible for differences in soil C

stocks in organic horizon and mineral soil. Differences in litter input (Díaz-Pinés et al.,

2011), root distribution (Oulehle et al., 2007) as well as decomposition rates (Cools et al.,

2014) have been suggested as the most likely explanation for tree species influence on soil C

stocks. Often, Douglas fir (*Pseudotsuga menziesii menziesii*) and European beech (*Fagus*

sylvatica) shows high fine root density in deeper soil layers (Bolte and Villanueva, 2006;

Oulehle et al., 2007), increasing C allocation in mineral soil layers. In contrast, the native

conifer Norway spruce (*Picea abies*), known as shallow-rooted tree species (Bolte and

Villanueva, 2006) accumulate C in the upper mineral soil layers.

Compositionally and structurally diverse forests represent an important element of

approaches to deliver a wide range of ecosystem goods and services (Ammer, et al., 2020).

Moreover, the establishment and management of mixed stands is also discussed as an

effective measure to adapt forests stands to climate change (Berg and McClaugherty, 2020;

Frouz et al., 2009) and C sequestration in soil. Admixtures of beech in conifer forests (either

with Douglas fir or Norway spruce) has been reported to modify C and N input through the

fine root distribution in the soil profile (Vesterdal and Raulund-Rasmussen, 1998).

Additionally, the enhanced nutrient concentrations of beech litter increase litter

decomposition of conifers needles (mixed stands)(Krishna and Mohan, 2017; Neumann et al.,

2018; Vesterdal et al., 2008), followed by less C accumulation in the organic soil layers and

shift to stable C pools in mineral soil. However, still inconsistent information on how mixed

stands of beech with Douglas would affect forest soil C and N vertical distribution. Due the

differences in litter input (Díaz-Pinés et al., 2011) as well as decomposition rates (Cools et

al., 2014) its most likely that this mixture enhance the organic C stocks at organic layers as

well. Thus, the deep root system reported to Douglas fir and beech and the high SOC on

organic layer might suggest that the admixture of beech into Douglas fir stands potentially

results in higher total C in the whole soil profile.

Therefore, the main objective of our study was to analyze the accumulation, vertical

distribution as well as litter quality indices (C/N ratios) of the upper mineral C and N stocks

of different pure and mixed stand types (pure European beech, pure Norway spruce, pure

Douglas-fir, mixed European beech/Norway spruce and mixed European beech/Douglas-fir)

along Northern Germany. We hypothesized that i) the admixture of beech to Douglas fir

forests will increase C stocks at the mineral soil compared to respectively beech

monocultures, ii) the OC stocks at litter layer will decrease under the admixture of beech to

Douglas fir and, shift to stables C pools (H-layer and mineral soil), compared to respectively

monocultures, and iii) on rather nutrient poor sites (sandy soil), species-identity effects on

organic layer will be stronger pronunced than on rich soils (loamy soil).

Material and Methods

Study sites

We investigated seven sites in Lower Saxony, northern Germany. Each site contained quintet

neighboring forest stands. Three of these stands were monospecific stands of European beech

(*Fagus sylvatica*), Norway spruce (*Picea abies*), or Douglas fir (*Pseudotsuga menziesii*). The

other two were mixed stands, one composed of spruce/beech and one composed of Douglas

fir/beech.

Essential criteria for the selection of the sites were the presence of the respective quintets in a
similar advanced silvicultural development status, altogether, we employed a multi-criteria
approach, considering factors such as climate, soil type (based on German soil inventory),
stand age, and existing research on land-use, as well as a relevant single tree admixture in the
mixed stands.

However, on the basis of the available forest site mapping and also confirmed by Foltran et
al. (2023), two groups with relatively homogeneous site conditions were distinguished in our
data set, in particular via the texture data (Table 1), but also addressing the geological
substrate classes, the correspondently developed soil types as well as the given range of water
and nutrient levels (Foltran et al., 2023).

In the first region in southern Lower Saxony, we chose three sites (Dassel, Winnefeld, and
Nienover). These three sites are located in the hills of the Solling plateau and lie between 300
and 450 m.a.s.l. They share similar climatic conditions, i.e. a mean annual air temperature of
7.2°C and mean annual precipitation of about 1040 mm/year. The age ranges between 45 and
90 years and the texture is relatively rich (23% of clay; Table 1). The soils at these locations
are Dystric Cambisols (FAO, 2015) developed from Triassic sandstone material.

In the second region, located in the northern Lower Saxony, we selected four sites (Unterlüß,
Nienburg, and Göhrde I and II). These sites are in lowland Lower Saxony and lie between 80
to 150 m.a.s.l. and all have similar climatic conditions, i.e., a mean annual air temperature of
8.4°C, mean annual precipitation of 720 mm/year. The age ranges between 53 and 130 years.
Most frequent soil types are therefore Podzols (Table 1) and their transitional forms towards
Cambisols.

With respect to nutrients and according to the official Lower Saxony forest site mapping and
Foltran et al. (2023), the southern sites (loamy soils) are nutrient-rich, and the northern sites
(sandy soils) are nutrient-poor.

Table 1. Soil classification from each plot is given following (FAO, 2014) and the German
classification (BGR) (Düwel et al., 2007). Soil parent material was identified matching the
plot coordinates with the German National inventory database (LBEG). Soil texture (10-30
cm*) was measured by the integral suspension pressure method (ISP) and determined by
PARIO (Foltran et al., 2023).

Region	Sites	Soil (FAO-WRB, 2014)	German Classification (BGR)	Parent material (LBEG)	Clay %	Silt %	Sand %
Solling Region (SR)	Dassel	Dystric Cambisols from quartzitic sandstones and conglomerates with low base status	Podsolige Braunerde aus basenarmen quarzitischen Sandsteinen und Konglomeraten	Medium colored sandstone / sandstone, siltstone, claystone	21	53	26
	Winnefeld				23	57	20
	Nienover				23	57	20
Northern Region (NR)	Nienburg	Haplic Podzols / Dystric Regosols from dry dystrophic sand deposits	Eisenhumus-Podsol / Podsol-Regosol aus trockenen, nährstoffarmen Sanden	Drenthe stage of the Saale glaciation / sand, gravel /melt water deposits	7	13	80
	Unterlüß				6	15	79
	Göhrde II				6	15	79
	Göhrde I	Spodic Arenosols from dry dystrophic sand deposits	Podsol-Braunerde aus trockenen, nährstoffarmen Sanden	Warthe stage of the Saale glaciation / silt / clayey, sandy, gritty / basic moraine	3	24	73

WRB: Working Group World Reference Base
BGR:Bundesanstalt für Geowissenschaften und Rohstoffe
122 LBEG: Landesamt für Bergbau, Energie und Geologie
123
124

Soil sampling

In each forest stand type (50 m x 50 m) at all sites, 4 randomly selected points were chosen as

representative sampling points. The selected points were oriented at stand-level, e.g., we

standardized two meters minimal distance from the trees to avoid coarse roots. At each

sampling plot, the forest floor was collected using a steel frame and sorted by identifiable

foliar (O_L – Litter), non-foliar (O_F – decay layer), and non-identifiable and humified (O_H –

Humus) layers of the organic layer. Mineral soil was sampled using a core auger (d=8 cm)

and, separated at 0-5, 5-10, and 10-30 cm soil depth. Bulk soil density from each depth was

calculated using soil metal rings (250 cm³) to further stock analysis.

Partly missing bulk density data due to frozen soil conditions, interfering tree roots or stones

during sampling were estimated by Adams equation (ADAMS, 1973) adapted by CHEN et

al. (2017). The approach uses SOM and pH as bulk density predictors.

Sample preparation and analysis

All mineral soil samples were oven-dried at 40°C until constant weight and sieved through 2

mm mesh, subsamples from the fine soil fractions (<2 mm diameter) were ground with a

Retsch mortar grinder RM 200 (Retsch, Germany) for 10 min. The organic layer samples

were dried at 60°C until constant weight, weighted and ball milled (MM2, Fa Retsch) for

further analysis.

All the samples were analyzed for total C and N by dry combustion in a Leco CSN 2000

analyzer. There was no inorganic C ($CaCO_3$) within 30 cm depth in soils and all measured C

were consequently considered to be organic.

The soil pH was measured in all mineral soil samples and are described in detail in the

Foltran et al. (2023).

150   Calculations of Stocks

We estimated the soil bulk density from the oven-dried and moisture corrected (105 ∘C) fine

soil mass and its volume. The fine soil volume was estimated from the difference between the

volume of the soil corer and the volume of stones and roots. Carbon and N stocks in each

layer were estimated from the organic horizon mass (O-layers) and soil bulk density (mineral

soil), concentrations of C and N and depth of the individual soil depths for each sampling

point.

158   Statistical Analyses

To address the non-independent nature of multiple horizons within one soil profile for this

subset of data, we chose linear mixed effect models (Rasmussen et al., 2018). To estimate the

effect of forest stand and site condition (loamy vs sandy), we fitted linear mixed models

(LMMs) to log-transformed response variables (C and N stocks) and then applied planned

contrasts (Piovia-Scott et al., 2019). All LMMs included forest stands (European beech,

Douglas fir, Douglas fir/beech, Norway spruce, spruce/beech), site conditions (loamy and

sandy sites), and soil depths (O_L, O_F, O_H [organic layers] and 0–5, 5–10 and 10-30 cm

[mineral soil]) as fixed effects. Models were stepwise selected by likelihood ratio test, and

minimal models included all main effects and the interaction of forest type and region.

The mean values for each site are given in the supplementary material (Table S3 and S4).

There, the effect of the forest stand (pure or mixed forest) on the soil chemical parameters for

individual depth was assessed by least significant difference test. As the assumption of

normal distribution was not met (tested for with Shapiro Wilk test), we used the Kruskal

Wallis test, followed by pairwise Mann-Whitney tests with a correction factor for multiple

pairwise testing, to identify statistically significant differences among forest stand and site

conditions by individual depth.

All analyses were done in R 4.0.3 (R Core Team, 2020). We used the 'nlme' package to fit

LMMs and the 'emmeans' package for planned contrasts. All mixed models met the

assumptions of normality of residuals and homogeneity of variance.

Results

Organic horizon mass and soil density

No effects of forest stand and region on organic horizon masses (OHM) was observed.

Instead, depth and the interactions Species:depth and Species:Region were consistency

significant affecting the organic layer masses. The model was found to be statistically

significant ($p < 0.001$; r^2=0.397), indicating that approximately 40% of the variation in the

response variable (OHM) could be explained by the model (Figure S2).

The effect of depth was significant, with O_H having a positive effect ($p<0.001$) on OHM, its

interaction term depth:Region shows that O_H layer shows negative effects at southern sites

than northern sites ($p<0.001$). The interaction terms show that both mixed stands (Douglas

fir/beech and spruce/beech) and O_H layer had significant and positive effects on OHM.

The mineral soil bulk density ranges from 0.8 to 1.4 g cm^{-3}. The southern sites showed

significant lower soil bulk density ($p<0.001$) than northern sites, and Douglas fir and Douglas

fir/beech showed significant lower bulk soil density ($p<0.001$) compared to beech. A

significant interaction among forest stands and site conditions was observed (Figure 1),

where Douglas fir and its mixture with beech (Douglas fir/beech) showed higher soil

densities at loamy soils (Southern sites) than at sandy soils (Northern sites) (Table S1).

Figure 1. Linear mixed-effects model on soil bulk density. The figure (a) shows the estimate coefficient (β), t-test and significance *p* for each term included in the model. Species (D: Douglas fir, S: spruce; Be: beech; Douglas fir/beech: DB; and, spruce/beech: SB), Region (Southern sites [loamy soils] and Northern sites [sandy soils]) and depth (0-5, 5-10 and 10-30 cm]).

Organic-horizon and mineral soil C and N concentrations

We used a linear mixed model to investigate the relationship between carbon (C) and nitrogen (N) concentration and species, depth, and their interaction across the sites (Northern and Southern sites). The results showed that all five stands had a significant effect on carbon concentration. The highest carbon concentration was found in the O_H layer and decreased with depth. Furthermore, the effect of depth varied across species, with higher carbon concentrations found in the top layer of the organic horizon for all species except Douglas fir. In terms of the species effect, the carbon concentration was significantly higher under beech and mixed spruce/beech stands compared to conifers, Douglas fir and spruce. Except at the O_H layer where both conifers and mixed stands had higher concentrations of carbon than

beech stands. Moreover, the interaction between species and sites was significant only for
Douglas Fir and mixed Douglas Fir/beech stands, indicating that the effect of species on
carbon concentration varied across sites, being significantly high at sandy soils (Northern
sites).
The forest stands also showed a significant effect on the N concentration. Specifically, the N
concentration was higher for both mixture forests (Douglas fir/beech, spruce/beech), and
spruce than for the pure beech stand. The depth also significantly affected the N
concentration. Compared to the reference depth (10-30 cm), all other depths were
significantly associated with higher N concentration. Additionally, the interaction between
Species and Sites was also found to be significant, indicating that the effect of species on the
N concentration varies depending on the site.
236

238 Figure 2. Linear mixed-effects model on Carbon (a), Nitrogen (b) concentration and CN ratio
239 (c). The figure (a) shows the estimate coefficient (β), t-test and significance *p* for each term
240 included in the model. Species (D: Douglas fir, S: spruce; Be: beech; Douglas fir/beech: DB;
241 and, spruce/beech: SB), Region (Southern sites [loamy soils] and Northern sites [sandy soils])
242 and depth [O_L, O_F, O_H, 0-5, 5-10 and 10-30 cm]).
243
244
245

C and N stocks

The results indicated that C stock was higher in the beech stands at the Southern sites (loamy

soils) (23.33 Mg ha^{-1}) than in Douglas-fir stands (19.60 Mg ha^{-1}). On the contrary, at

Northern sites (sandy soils), Douglas-fir stands had higher C stock (23.46 Mg ha^{-1}) than

beech stands (16.29 Mg ha^{-1}).

The multiple stepwise regression model explained 49% of the variance in C stock and was

found to be statistically significant ($p < 0.001$). Species, soil conditions, and depth were

significant predictors of C stock ($p < 0.05$) (Figure 2). The coefficient estimate for sandy soils

was positive, indicating that C stock were higher in sandy soils compared to loamy soils. The

interaction term between species and sandy soil was statistically significant ($p < 0.001$) and

negative, indicating that the effect of species on C stock differed between sandy and loamy

soils, and it was attenuated in sandy soils. There, Douglas fir and Douglas fir/beech showed

the highest C stocks compared to beech.

The effect of species on C stock was stronger at litter layer (O_L). Additionally, the

interactions between Species:depth and Species:sandy soils were found to be statistically

significant ($p < 0.05$) with negative coefficients, suggesting that the effect of species on C

stock was more pronounced in the litter layer at sandy soils.

Figure 3. Linear mixed-effects model on Carbon stock (a) and Carbon stock (b; Mg ha^{-1}). The figure (a) shows the estimate coefficient (β), t-test and significance *p* for each term included in the model. Species (D: Douglas fir, S: spruce; Be: beech; Douglas fir/beech: DB; and, spruce/beech: SB), Region (Southern sites [loamy soils] and Northern sites [sandy soils]) and depth [L,F,H,0-5, 5-10 and 10-30 cm]). The figure (b) shows the C stock in the organic layer (brown bars) and mineral soil (green bars), and the standard error (n=16).

Figure 4. Linear mixed-effects model on Nitrogen stock (a) and Nitrogen stock (b; Mg ha⁻¹). The figure (a) shows the estimate coefficient (β), t-test and significance *p* for each term included in the model. Species (D: Douglas fir, S: spruce; Be: beech; Douglas fir/beech: DB; and, spruce/beech: SB), Region (Southern sites [loamy soils] and Northern sites [sandy soils]) and depth [L,F,H,0-5, 5-10 and 10-30 cm]). The figure (b) shows the N stock in the organic layer (brown bars) and mineral soil (green bars), and the standard error (n=16).

The model was statistically significant ($p < 0.001$) and explained 63.53% of the variance in N
stock. The results showed that Species had a significant effect on N stock ($p = 0.02$).
Aditionally, the interaction effect between Species and soil condition was also significant (p
>0.001), indicating that the effect of Species on N stock was attenueted in sandy soils
compared to loamy soils. At the loamy soils (Southern sites), total N stocks were higher
under spruce stands than under Douglas fir stands. In contrast, at the sandy soils (Northern
sites), higher N stocks under Douglas fir stand than under spruce stand were observed (Figure
4).
At the southern sites, significantly larger N stocks were observed under mixed spruce/beech
stand than Douglas/beech, whereas at nouthern sites, the opposite were observed
(Douglas/beech > Spruce/beech).
At all sites, N stocks at organic layer under mixed beech–conifer stands (either
Douglas/beech or spruce/beech), significantly exceeded beech stands. The effect of the
mixtures showed the same site-dependent patterns as reported for SOC stocks, where the
stocks under both beech–conifer mixtures were similar to those under the respective pure
conifer stands.
The soil depth was also a significant predictor of N stock, with 10-30 cm showing the highest
N stock ($p >0.001$). However, none of the interaction terms between Species and depth were
significant, indicating that the effect of Species on N stock did not vary across different soil
depths.
297
SOC Distribution

The contribution of the organic layer stocks (SOC_{Org}) to the total stocks (SOCt) was site-
dependent. At southern sites, the SOC_{Org} contributed between 15% and 22% for Douglas fir
and spruce stands, respectively, whereas for beech it was only 8%. The mixture stands
showed intermediate results, ranging from 10% (Douglas fir/beech) to 17% (spruce/beech).
At northern sites, the contribution of SOC_{Org} on SOCt stock was strong pronounced. Under
the spruce stand, 33% of the SOC stock is allocated to the organic horizon, whereas 20% was
observed in the Douglas fir stand. The contribution of SOC_{org} to SOCt for mixed forests
ranged between 23% (Douglas/beech) and 30% (spruce/beech) and only 10% for beech
stands.

The N allocation corresponded to the SOC vertical distribution. At the northern sites, the relative contribution ranged from 20–26% for conifers and mixed forest to 11% at the beech forest, whereas at the southern sites, it ranged from 13–17% for conifers to only 6% under beech forest. At northern sites, mixture stands showed a clear contrast: at the spruce/beech stand, the contribution of organic layer N stock to the total N stock was 30%, while at the Douglas/beech stand, it was only 7%.

Overall, under sandy soil conditions, conifers and mixed forests allocated 10% more SOC and N at the organic layer compared to loamy soils, whereas the SOC and N stocks under beech maintained the same proportion (>90% at SOC_{ms}), independent of the site condition.

Effect of abiotic factors on C and N stocks

The effect of abiotic factors like soil pH on OC stock was investigated using linear mixed models (Figure 5). The increase of pH caused significant decreased (P<0.001) on SOC at mineral soil. The increase of 1 unit of pH (3.75 to 4.75) reduced in about 25 % the total SOC. Whereas, the opposite was observed for N stocks, where the increased of pH showed increases in N stocks.

Figure 5. Relationship between the soil organic carbon (SOC) (a) and Nitrogen (b) on mineral soil. The grey line denotes a linear mixed-effect model (LMM; P < 0.05) with the predictor variables as fixed effect and species as random intercept terms. The dashed line represents a

95% confidence interval. All response variables (C stock, and N stock) were transformed

with natural logarithm, whereas predictor variable (pH) were standardized by subtracting the

mean and dividing by the standard deviation.

Discussion

Tree species effects

We observed significant effect of tree species on organic C and N stocks as well as its

vertical distribution across all investigated sites. Overall, the organic-horizon (O) stocks of C

and N were significantly higher under conifer forests (Douglas fir and spruce) than under

beech (Figure 1). At the mineral soil (MS) the effects of tree species were site-dependent.

The southern sites (loamy soil), beech and spruce stands presented higher C and N stocks

than Douglas fir forest, confirming earlier results of Antisari et al. (2015) where smallest

SOC stocks of the organic horizon under beech stand are accompanied by enhanced SOC

stock in the mineral topsoil. In contrast, at northern sites (sandy soil), the beech stands

showed the smallest OC stock at both layers (O and MS).

According to Neumann et al. (2018), broadleaves litter tends to decompose faster compared

to conifer litter. This can be attributed to lower lignin and phenol concentrations, which

promote rapid decomposition rates and efficient accumulation of mineral-associated organic

matter, as reported by Rasse et al. (2005). In our study, we observed a higher carbon (C)

input from the organic horizon to the mineral soil in the beech forest under loamy soil

conditions, supporting the findings of previous research. The higher C allocation on mineral

soil than organic horizon under beech than conifers has also been reported in common garden

studies (Vesterdal et al., 2013). The differences in vertical distribution has been attributed to

the associated community of macrofauna species in beech dominated forests (Achilles et al.,

2021).

Therefore, the quality of the soil organic matter (SOM) might be significantly different under

coniferous and deciduous trees (Jaffrain et al., 2007). The O_H layer under conifers results in a

more recalcitrant and hydrophobic composition than that of O_H layer at beech stand (Thomas

et al., 2014) . The low mineralization quotient consequently leads to larger C accumulation

due the slow C turnover, as observed in our study under both conifers stands.

Besides the litter quality, rooting patterns and microbial activity among beech, Douglas fir

and Norway spruce might change the C inputs between tree species. Simultaneous studies on

the turnover of fine root biomasses carried out by Lwila et al. (2021) have shown that beech

forests showed higher fine roots biomass than Douglas fir at the sandy soils, whereas the fine

root necromass was higher for Douglas fir compared to beech. High microbial activity at the

upper mineral soil is reported in our study area (Lu and Scheu, 2021), suggesting high

rhizosphere carbon inputs at the beech forest stand compared to Douglas fir.

In general, conifer roots contain lower lignin concentrations than roots of beech (Thomas et

al., 2014) (Newman and Hart 2006), which might lead to a lower longevity and a faster

decomposition (Berger et al., 2002; Cremer et al., 2016). Additionally, fine roots necromass

deliver a considerable amount of organic material (Cremer et al., 2016; Dawud et al., 2017)

enhancing the root-derived SOM. Due the higher productivity of Douglas fir than beech (in

particular in dry conditions as at northern sites) (Schmid et al., 2014; Schmidt et al., 2015)

higher turnover rate of Douglas fir fine roots is expected and consequently fast fine root

turnover might cause favorable changes on SOC stocks, as observed in our study and similar

studies (Lorenz and Lal, 2014).

Mixed forest effect

Independent of the site conditions, the pure stands of beech showed relatively small total

SOC stocks, whereas the pure conifer sites enhanced total SOC and N stocks, accumulated to

a large extent in the organic layer. Furthermore, the admixture of conifers to beech enlarged

the total SOC and N storage only in the northern sites, i.e. under sandy soil conditions,

whereas in the southern sites, i.e., under loamy soil conditions, both mixtures of conifers to

beech presented similar storages compared to pure beech. Overall, the carbon (C) stocks in

the O_H layer, as well as the total soil organic carbon (SOC) and nitrogen (N) stocks, were

generally similar between mixed beech-conifer stands and their respective pure stands. The C

stocks at both mixed stands resembled those of beech stands at southern sites and approached

the levels observed in conifer stands at northern sites.

The admixture of beech into conifers stands in nutrient-poor soil conditions, particularly

observed at the northern sites, had significant effects on soil organic carbon (SOC) and

nitrogen (N) stocks, which aligns with the results reported by Dawud et al. (2017). In these

soil conditions, both the mixture of Douglas fir and beech and pure Douglas fir stands

exhibited higher SOC content in the humus layer (O_H) and the 5-30 cm soil depth. This can

be attributed to the competitive advantage of beech fine roots, which have the ability to

access deeper soil layers and exploit areas less occupied by competing species. In contrast,

Douglas fir fine roots tend to be restricted to the topsoil (15-30 cm), resulting in a shift in the

distribution of beech fine roots towards the subsoil. This phenomenon has been reported in

previous studies (Hendriks and Bianchi, 1995; Schmid and Kazda, 2002). These observed

shifts of root vertical distribution are known as below-ground complementarity effects, e.g.,

through vertical segregation of roots of different species which exploit soil at different depths

allowing for reduced competition (Loreau M, 2001), has been suggested as a mechanism by

which species mixtures store more C in deeper soil layers than monocultures (Bauhus et al.,

2009; Forrester et al., 2006). Additionally, the admixture of beech leaf litter to the more

recalcitrant needle litter induced a faster litter mass loss and consequently shift of SOC stock

from O_L to O_H layer, i.e., similar decomposition rate (k) for mixed stands and pure beech and

higher than Douglas fir and spruce forests, suggesting similar decomposability of litter in the

mixed forest and in the beech forest (not published).

Thus, allocation of C in soils is an important ecosystem service provided by forests (Pretzsch

et al., 2017). Owing to the potential of higher SOC storages in mixed-species forests

compared to pure stands and to shift the C into stable pools i.e., the shift from O_L layers to O_H

layer, enhance the capacity of mixed forest to provide ecosystem services e.g., improve soil

functioning like soil biodiversity and nutrient availability and, furthermore the resilience of

the forest ecosystem to cope with future climate change scenarios.

Abiotic effects

The persistence of SOC is largely due to complex interactions between SOC and its

environment, such as reactive mineral surfaces, climate, water availability and soil acidity

(Leifeld et al., 2013; Mikutta et al., 2009; Zhou et al., 2019). The most important factor in

SOC stabilization at sites with high clay contents, as partly observed in our study, is probably

the association with soil minerals, irrespective of forest type (Fierer and Jackson, 2006; Meier

et al., 2020). Indeed, enhanced clay and silt content increased the SOC in the mineral soil at

our investigated sites and, in contrast decreased the SOC storage at organic horizon. Loamy

soils are known to buffer influences by tree species more strongly than sandy soils (MEIER

and LEUSCHNER, 2010) and this would explain why we see more effects of species on the

northern sites dominated by sandy soils than at southern sites.

Therefore, differences in soil pH often go along with differences in soil mineralogy as well

and the latter exerts control on the stabilization of mineral associated organic matter (Meier et

al., 2020) and microbiota community (MEIER and LEUSCHNER, 2010). The accumulation

of organic matter in soils is influenced by relatively low pH levels, which can reduce the

decomposition of soil organic matter (SOM), as highlighted in the study by Meier et al.

(2020). This, in turn, leads to noticeable effects of tree species on organic matter

accumulation, as observed in our study. Interestingly, Meier et al. (2020) also suggested that

soil acidity is associated with increased rates of root exudation in beech forests, particularly

in the topsoil of glacial sandy soils found in the northern sites. This increased root exudation

can be seen as an adaptation of the trees to low nutrient availability, a finding consistent with

our study (Foltran et al., 2023). In these conditions, the majority of nutrient uptake occurs in

the AE horizon, which is enriched with organic material, further explaining the adaptability

of beech forests in response to varying nutrient availability.

Conclusion

Site dependent effects of tree species (European beech, Douglas fir and Norway spruce) on

SOC and N stocks as well as on SOC and N concentration were observed. The SOC and N

stocks generally were smallest in pure beech stands compared with Douglas fir and Norway

spruce. The SOC and N stocks in mixed stands of beech with Douglas fir or Norway spruce

are generally between those of the respective pure stands. However, the results indicate that

the effects of admixture of beech into conifers stands (Douglas fir and spruce) are site-

dependent At the Northern sites, the adaptability of Douglas fir under dry and poor-nutrient

site conditions combined with high plasticity of beech promoted a favorable effect on total

SOC and N, enhancing its storages as observed at pure Douglas fir and the mixture Douglas

fir/beech. In contrast, at the Southern sites, spruce/beech showed higher SOC and N stocks

than Douglas/beech.

The Douglas fir/beech stand mixture showed significant increases of total SOC under sandy

soils (Northern sites). Overall, the higher potential for carbon storage in mixed-species

forests compared to pure stands, along with the shift of carbon into more stable pools (such as

from the O_L layer to the O_H layer), emphasizes the capacity of mixed forest to provide

valuable ecosystem services while reducing the risk of unintended carbon losses.

Author contributions

All authors contributed to the study conception and design. Material preparation, data

collection and analysis were performed by EF. The first draft of the manuscript was written

by EF and all authors commented on previous versions of the manuscript. All authors read

and approved the final manuscript.

Acknowledgements

The study was conducted as part of the Research Training Group 2300 funded by the German

research funding organization (Deutsche Forschungsgemeinschaft – DFG). We gratefully

acknowledge the administrative support by Serena Müller and the indispensable help of

Julian Meyer and Dirk Böttger during soil sampling. Furthermore, we thank Sylvia Bondzio,

Karin Schmidt for their valuable advice during laboratory work.

References

Ammer, Christian; Annighöfer, Peter; Balkenhol, Niko; Hertel, Dietrich; Leuschner,

Christoph; Polle, Andrea; Lamersdorf, Norbert; Scheu, Stefan; Glatthorn, J. (2020),

2020. RTG 2300 - study design, location, topography and climatic conditions of

research plots in 2020. Pangaea.

Antisari, L.V., Falsone, G., Carbone, S., Marinari, S., Vianello, G., 2015. Douglas-fir

reforestation in North Apennine (Italy): Performance on soil carbon sequestration,

nutrients stock and microbial activity. Applied Soil Ecology 86, 82–90.

https://doi.org/10.1016/j.apsoil.2014.09.009

Berg, B., McClaugherty, C., 2020. Plant Litter - Decomposition, Humus Formation, Carbon
Sequestration, 4th ed. Springer International Publishing. https://doi.org/10.1007/978-3-
030-59631-6

Berger, T.W., Neubauer, C., Glatzel, G., 2002. Factors controlling soil carbon and nitrogen
stores in pure stands of Norway spruce (Picea abies) and mixed species stands in
Austria. Forest Ecology and Management 159, 3–14. https://doi.org/10.1016/S0378-
1127(01)00705-8

Bolte, A., Villanueva, I., 2006. Interspecific competition impacts on the morphology and
distribution of fine roots in European beech (fagus sylvatica L.) and Norway spruce
(picea abies (L.) karst.). European Journal of Forest Research 125, 15–26.
https://doi.org/10.1007/s10342-005-0075-5

Cepáková, Š., Tošner, Z., Frouz, J., 2016. The effect of tree species on seasonal fluctuations
in water-soluble and hot water-extractable organic matter at post-mining sites.
Geoderma 275, 19–27. https://doi.org/10.1016/j.geoderma.2016.04.006

Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K., Paul, E., 2013. The Microbial
Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter
decomposition with soil organic matter stabilization: Do labile plant inputs form stable
soil organic matter? Global Change Biology 19, 988–995.
https://doi.org/10.1111/gcb.12113

Cremer, M., Kern, N.V., Prietzel, J., 2016. Soil organic carbon and nitrogen stocks under
pure and mixed stands of European beech, Douglas fir and Norway spruce. Forest
Ecology and Management 367, 30–40. https://doi.org/10.1016/j.foreco.2016.02.020

Cremer, M., Prietzel, J., 2017. Soil acidity and exchangeable base cation stocks under pure
and mixed stands of European beech, Douglas fir and Norway spruce. Plant and Soil
415, 393–405. https://doi.org/10.1007/s11104-017-3177-1

Dawud, S.M., Raulund-Rasmussen, K., Domisch, T., Finér, L., Jaroszewicz, B., Vesterdal,
L., 2016. Is Tree Species Diversity or Species Identity the More Important Driver of Soil
Carbon Stocks, C/N Ratio, and pH? Ecosystems 19, 645–660.
https://doi.org/10.1007/s10021-016-9958-1

Dawud, S.M., Vesterdal, L., Raulund-Rasmussen, K., 2017. Mixed-species effects on soil C
and N stocks, C/N ratio and pH using a transboundary approach in adjacent common
garden douglas-fir and beech stands. Forests 8. https://doi.org/10.3390/f8040095

FAO, 2014. World reference base for soil resources 2014. International soil classification
system for naming soils and creating legends for soil maps, World Soil Resources
Reports No. 106.

Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial
communities. Proceedings of the National Academy of Sciences of the United States of
America 103, 626–631. https://doi.org/10.1073/pnas.0507535103

Foltran, E.C., Lamersdorf, N., Ammer, C., 2023. Do admixed conifers change soil nutrient
conditions of European beech stands? *Soil Research* 63 https://doi.org/10.1071/
SR22218

bioRxiv 2020.09.25.313213. https://doi.org/10.1101/2020.09.25.313213

Frouz, J., Pižl, V., Cienciala, E., Kalčík, J., 2009. Carbon Storage in Post-Mining Forest Soil,
the Role of Tree Biomass and Soil Bioturbation. Biogeochemistry 94, 111–121.

Krishna, M.P., Mohan, M., 2017. Litter decomposition in forest ecosystems: a review.
Energy, Ecology and Environment 2, 236–249. https://doi.org/10.1007/s40974-017-
0064-9

Leifeld, J., Bassin, S., Conen, F., Hajdas, I., Egli, M., Fuhrer, J., 2013. Control of soil pH on
turnover of belowground organic matter in subalpine grassland. Biogeochemistry 112,
59–69. https://doi.org/10.1007/s10533-011-9689-5

Lorenz, K., Lal, R., 2014. Soil organic carbon sequestration in agroforestry systems. A
review. Agronomy for Sustainable Development 34, 443–454.
https://doi.org/10.1007/s13593-014-0212-y

Lu, J.-Z., Scheu, S., 2020. Mixing coniferous and deciduous trees (Fagus sylvatica): Site-
specific response of soil microorganisms 1–29. https://doi.org/:
https://doi.org/10.1101/2020.07.21.213900

Meier, I.C., Tückmantel, T., Heitkötter, J., Müller, K., Preusser, S., Wrobel, T.J., Kandeler,
E., Marschner, B., Leuschner, C., 2020. Root exudation of mature beech forests across a
nutrient availability gradient: the role of root morphology and fungal activity. New
Phytologist 226, 583–594. https://doi.org/10.1111/nph.16389

MEIER, I.N.A.C., LEUSCHNER, C., 2010. Variation of soil and biomass carbon pools in
beech forests across a precipitation gradient. Global Change Biology 16, 1035–1045.
https://doi.org/https://doi.org/10.1111/j.1365-2486.2009.02074.x

Mikutta, R., Schaumann, G.E., Gildemeister, D., Bonneville, S., Kramer, M.G., Chorover, J.,
Chadwick, O.A., Guggenberger, G., 2009. Biogeochemistry of mineral-organic
associations across a long-term mineralogical soil gradient (0.3-4100 kyr), Hawaiian
Islands. Geochimica et Cosmochimica Acta 73, 2034–2060.
https://doi.org/10.1016/j.gca.2008.12.028

Mueller, K.E., Eissenstat, D.M., Hobbie, S.E., Oleksyn, J., Jagodzinski, A.M., Reich, P.B.,
Chadwick, O.A., Chorover, J., 2012. Tree species effects on coupled cycles of carbon,
nitrogen, and acidity in mineral soils at a common garden experiment. Biogeochemistry
111, 601–614. https://doi.org/10.1007/s10533-011-9695-7

Oulehle, F., Hofmeister, J., Hruška, J., 2007. Modeling of the long-term effect of tree species
(Norway spruce and European beech) on soil acidification in the Ore Mountains.
Ecological Modelling 204, 359–371. https://doi.org/10.1016/j.ecolmodel.2007.01.012

Rasse, D.P., Rumpel, C., Dignac, M.F., 2005. Is soil carbon mostly root carbon? Mechanisms
for a specific stabilisation. Plant and Soil 269, 341–356. https://doi.org/10.1007/s11104-
004-0907-y

Renger, M.;, Bohne, K.;, Facklam, M.;, Harrach, T.;, Riek, W.;, Schäfer, W.;, Wessolek, G.;,
Zacharias, S., 2008. Ergebnisse und Vorschläge der DBG-Arbeitsgruppe „Kennwerte
des Bodengefüges" zur Schätzung bodenphysikalischer Kennwerte. Bodenökologie und
Bodengenese 51.

Robertson, A.D., Paustian, K., Ogle, S., Wallenstein, M.D., Lugato, E., Francesca Cotrufo,
M., 2019. Unifying soil organic matter formation and persistence frameworks: The

MEMS model. Biogeosciences 16, 1225–1248. https://doi.org/10.5194/bg-16-1225-
2019

Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A.,
Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P., Kleber, M., Ko, I., 2011.
Persistence of soil organic matter as an ecosystem property. Nature 478.
https://doi.org/10.1038/nature10386

Thomas, F.M., Molitor, F., Werner, W., 2014. Lignin and cellulose concentrations in roots of
Douglas fir and European beech of different diameter classes and soil depths. Trees -
Structure and Function 28, 309–315. https://doi.org/10.1007/s00468-013-0937-2

Vesterdal, L., Clarke, N., Sigurdsson, B.D., Gundersen, P., 2013. Do tree species influence
soil carbon stocks in temperate and boreal forests? Forest Ecology and Management
309, 4–18. https://doi.org/10.1016/j.foreco.2013.01.017

Vesterdal, L., Raulund-Rasmussen, K., 1998. Forest floor chemistry under seven tree species
along a soil fertility gradient. Canadian Journal of Forest Research 28, 1636–1647.
https://doi.org/10.1139/cjfr-28-11-1636

Vesterdal, L., Schmidt, I.K., Callesen, I., Nilsson, L.O., Gundersen, P., 2008. Carbon and
nitrogen in forest floor and mineral soil under six common European tree species. Forest
Ecology and Management 255, 35–48.
https://doi.org/https://doi.org/10.1016/j.foreco.2007.08.015

Zhou, W., Han, G., Liu, M., Li, X., 2019. Effects of soil pH and texture on soil carbon and
nitrogen in soil profiles under different land uses in Mun River Basin, Northeast
Thailand. PeerJ 2019. https://doi.org/10.7717/peerj.7880