A retinotopic code structures the interaction between perception and memory systems

Adam Steel[1*], Edward H. Silson[2*], Brenda D. Garcia[1], & Caroline E. Robertson[1]

[1]Department of Psychology and Brain Sciences, Dartmouth College, Hanover, NH, 03755
[2]Psychology, School of Philosophy, Psychology, and Language Sciences, University of Edinburgh, Edinburgh, UK EH8 9JZ
Corresponding author: Adam Steel, Department of Psychology and Brain Sciences, Dartmouth College, 3 Maynard Street, Hanover, NH, 03755; email: adam.steel@dartmouth.edu; tel: +1 (202) 640 9340

* equal author contribution

Author contributions: AS, EHS, CER conceived and designed the experiment. EHS and BDG contributed stimulus code. AS and BDG collected the data. AS processed the data. AS and EHS analyzed the data. All authors wrote and edited the manuscript.

Conflict of interest statement

The authors declare no conflict of interest.

Data availability statement

Data will be made available via datalad upon publication.

Code availability statement

Code used for data analysis will be made available via GitHub upon publication. Analysis code will be made available to editors and reviewers during review via Dropbox upon request.

Acknowledgements

We thank Iris Groen for assistance with specific analysis code. This work was supported by the National Institute of Mental under award number 1R01MH130529-01 (C.E.R.) and by the Neukom Institute for Computational Science (A.S.).

1 Introduction

We encode the visual world retinotopically, imposing a spatial code on visual information processing[1–4].
However, models of brain organization generally assume that retinotopic coding is replaced by abstract,
amodal coding as information propagates through the visual hierarchy[5,6] towards memory structures[7–12].
This view raises a puzzle for constructive accounts of visual memory: how can mnemonic and visual
information interact effectively in the brain if they are represented using fundamentally different neural
codes[13]? Recent work has suggested that even high-level cortical areas, including the default mode
network, exhibit retinotopic coding: they contain visually evoked population receptive fields (pRFs) with
inverted response amplitudes[14,15]. But, the functional relevance of this retinotopic coding at the cortical
apex remains unclear. Here, we report that retinotopic coding at the cortical apex structures interactions
between mnemonic and perceptual areas in the brain. Using fine-grained, individual participant functional
magnetic resonance imaging (fMRI), we show that just beyond the anterior edge of category-selective
visual cortex, category-selective memory areas exhibit a robust, inverted retinotopic code. The negative
and positive pRF populations in mnemonic and perceptual areas, respectively, have closely matched visual
field representations, reflecting their tight functional coupling. Moreover, the +/- pRFs in perceptual and
mnemonic cortex exhibit spatially-specific opponent responses during both bottom-up visual processing
and top-down memory recall, suggesting that these areas are interlocked in a mutually-inhibitory
dynamic. This spatially-specific opponency further generalizes to familiar scene perception, a task that
requires mnemonic-perceptual interplay. Together, these results show that retinotopic coding structures
interactions between perceptual and mnemonic systems in the brain, thereby scaffolding their dynamic
interaction.

23 Main text

Recent reports have described the presence of retinotopic coding in high-level cortical areas, including
the default mode network [14,15]. Yet, this finding is at odds with the conventional view of cortical
representation, which assumes that retinotopic coding is replaced by amodal, abstract coding as
information propagates through the visual hierarchy towards the cortical apex [7–12]. What is the functional
role of retinotopic coding in areas atop the cortical hierarchy? Here, we tested the hypothesis that the
retinotopic code spanning levels of the cortical hierarchy structures interactions between functionally-
coupled mnemonic and perceptual areas of the brain.

We began by determining whether retinotopic coding was present in mnemonic and perceptual regions
spanning levels of the cortical hierarchy. In Exp. 1, participants (N=17) underwent visual population
receptive field (pRF) mapping[16,17] **(Fig. 1A)** with advanced multi-echo fMRI [18,19] to maximize signal-to-noise
and pRF model fitting in anterior regions of the temporal and parietal lobes [19]. As expected, an initial
group-level whole-brain analysis revealed robust, positive retinotopic responses extending from early to
high-level visual areas (positive pRFs (+pRFs)) **(Fig. 1B)**. We also observed robust and reliable pRFs beyond
the anterior extent of known retinotopic maps[20] in anterior ventral temporal and lateral parietal cortex,

regions of the cortical apex historically considered amodal [9,10] **(Fig. 1B).** In striking contrast with classically-defined visual areas, a large portion of these anterior voxels had pRFs with an inverted visual response (i.e., negative visually-evoked pRFs, -pRF[14,15]). For brevity, we refer to voxels with positive or negative pRF response amplitudes as +/- pRFs, respectively.

At the group-level, -pRFs only appeared beyond the anterior extent of known retinotopic maps[20], in areas associated with the cortical apex, consistent with prior reports[14,15] **(Fig. 1B)**. This topographic profile of pRF amplitude was evident at the single participant level, with robust pRF responses evident throughout posterior cerebral cortex and -pRFs emerging at the anterior edge of high-level visual cortex **(Fig. S1)**. In general, these anterior pRFs were not arranged topographically on the cortical surface in a manner that recapitulates the layout of the retina (i.e., did not constitute a 'retinotopic map'). However, these -pRFs were robust and reliable within individuals: a half-split analysis confirmed the signed amplitude (i.e., positive or negative) was highly reproducible **(see Supplemental Material)**. Importantly, these anterior -pRF populations on the ventral and lateral surfaces are distinct from the population of -pRFs observed on the medial wall in peripheral early visual areas, which arise from stimulation outside the pRF's receptive field (i.e., surround suppression) [15,21].

Figure 1. *Population receptive field (pRF) mapping reveals retinotopic coding throughout posterior cortex, comprised of both positive and negative amplitude pRFs.* ***A.*** *PRF paradigm and modeling. In Exp. 1, participants underwent pRF mapping. Participants viewed visual scenes through a bar aperture that gradually traversed the visual field. Each visual field traversal lasted 36 seconds (18, 2s positions), and the bar made eight traversals per run. The direction of motion varied between traversals. To estimate the pRF for each voxel, a synthetic timeseries is generated for 400 visual field locations (200 x and y positions), and 100 sizes (sigma). This results in 4 million possible timeseries that are fit to each voxel's activity. The fit results in four parameters describing each receptive field: x, y, sigma, and amplitude.* ***B.*** *Negative amplitude pRFs fall anterior to cortex typically considered visual. Group-average (n=17) pRF amplitude map (thresholded at $R^2>0.08$) is shown on partially inflated representations of the left hemisphere, alongside regions-of-interest the scene perception areas (OPA, PPA), place memory areas (LPMA, VPMA) (localized in an independent group of participants[22]), and the default mode network [23]. The known retinotopic maps in posterior cortex (black*

If the inverted retinotopic code at the cortical apex serves a functional role of structuring interactions with perceptual areas, we hypothesized that -pRFs would concentrate in mnemonic, as compared with perceptual functional areas. Consistent with this prediction, we observed that -pRFs tended to fall within swaths of cortex that selectively responded during top-down recall of familiar places (place memory areas, PMAs[22]) on the lateral and ventral surfaces (lPMA and vPMA, respectively) at the group level [22]**(Fig. 1B)**. These two mnemonic areas each lie immediately anterior to one of the scene perception areas[24] (SPAs) of the human brain (occipital place area, OPA[25,26], and parahippocampal place area, PPA [27]). In contrast to the PMAs, the SPAs tended to contain +pRFs. We confirmed this observation in individual participants by calculating the percentage of +/- pRFs within individually localized SPAs and PMAs (lateral: LSPA, LPMA; ventral: VSPA, VPMA) (**Fig. 2A**, see Methods) to better understand their nuanced functional topography[22,28–30] **(Fig. 2A, S1-S2).** This analysis confirmed that, unlike the perceptual SPAs, which almost exclusively contained +pRFs, the mnemonic PMAs contained a significant percentage of -pRFs **(Fig. 2B)** (two-way rmANOVA, main effect of ROI ($F(1, 16)=85.83, p<0.0001$; ROI:Hemisphere interaction ($F(1, 16)=11.24$, $p=0.004$); post-hoc tests: lh: OPA v LPMA $t(16)=10.74$, $p_{corr}=0.000002$, D=3.43; rh: OPA v LPMA $t(16)=5.97$, $p_{corr}=0.00002$, D=1.89). On the ventral surface, this effect was generally stronger in the left compared to the right hemisphere (two-way rmANOVA, main effect of ROI ($F(1, 16)=10.01, p=0.006$; no ROI:Hemisphere interaction ($F(1, 16)=3.12$, $p=0.09$; lh: PPA v VPMA $t(16)=2.62$, $p_{corr}=0.02$, D=0.94; rh: PPA v VPMA $t(16)=2.57$, $p_{corr}=0.04$, D=0.90). Notably, in many subjects, the PMAs contained -pRFs and +pRFs; both -/+ pRFs populations in the PMAs were equally driven by recall (ts <0.5, ps > 0.6), suggesting their primary functional difference is their response to visual stimulation. Together, these results show that –pRFs are disproportionately represented in category-selective mnemonic areas (PMAs), as compared with their perceptual counterparts (SPAs).

Figure 2. *Transition to mnemonic cortex is marked by the appearance of negative population receptive fields (pRFs). **A**. PRF modeling reveals posterior-anterior inversion of pRF amplitude in individual participants. Left. PRF amplitude for a representative participant overlaid onto a lateral view of the left hemisphere*

Interestingly, both +pRFs and -pRFs in the PMAs tended to be smaller than +pRFs in the SPAs on both surfaces (Lateral surface: Main effect of ROI (F(2, 32)=10.27, p=0.0003; OPA vs +pRFs in LPMA: t(16)=5.61, Pcorr=0.0001, D=1.83; OPA vs -pRFs in LPMA: t(16)=2.59, Pcorr=0.01, D=1.31; Ventral surface: F(2, 22)=15.46 p=0.000006; PPA v +VPMA: t(11)=2.83, Pcorr=0.03, D=0.99; PPA v -VPMA(t(11)=5.43, Pcorr=0.00006, D=1.85). PRF sizes were not significantly different between +/- pRFs in LPMA (t(16)=1.52, Pcorr=0.14, D=0.48), but on the ventral surface +pRFs in VPMA were significantly larger than -pRFs in VPMA (t(11)=2.87, Pcorr=0.03, D=1.06). This result particularly notable, as it runs counter to the typical pattern of pRFs becoming larger moving anteriorly from early visual areas towards higher-level brain regions[6,16,31,32]. Although how the additional spatial precision manifests in these anterior areas is not clear, this result suggests that the PMAs could potentially represent highly specific information even compared to their perceptual counterparts **(Fig. 3)**.

Figure 3. *Memory areas contain smaller pRFs compared to their paired perceptual areas. **Left.** Group average pRF size with memory areas and perception areas overlaid. Nodes are thresholded at R^2 > 0.08. **Right.** Bars represent the mean pRF size for +pRFs in SPAs (OPA, PPA) and +/- pRFs in PMAs (LPMA, VPMA). Individual data points are shown for each participant. Across both surfaces, pRFs were significantly smaller on average in the PMAs than their perceptual counterparts.*

The enriched concentration of -pRFs in mnemonic (PMA) as compared with perceptual (SPA) areas led to a specific hypothesis regarding their functional role. Specifically, the PMAs are thought to act as a bridge between the cortical SPAs and the spatio-mnemonic system in the medial temporal lobe[22], but the format

for sharing information among these areas is not clear. We hypothesized that shared retinotopic coding
could serve as a shared substrate to scaffold the interaction between perceptual and mnemonic systems.
If this proposal is correct, we should observe evidence for a functional interaction between these areas
that depends on retinotopic position. We explored three tests of this hypothesis: do paired +/- pRFs
represent similar portions of the visual field (Exp. 1)?; do the +/- pRFs identified during visual stimulation
maintain their opponent activation profiles during top-down memory recall (Exp., 2)?; is the functional
link between +/- pRFs recapitulated when viewing familiar scene images (Exp. 3)?

First, we reasoned that if the retinotopic code scaffolds communication between perceptual and
mnemonic systems, -pRFs within mnemonic areas should exhibit the same differential visual field biases
as their perceptual counterparts[17]. In particular, visual field representations in OPA and PPA are biased
towards the lower and upper visual fields, respectively. Do -/+ pRFs in their respective PMAs share these
biases? Using pRF data (Experiment 1), we calculated visual field coverage estimates for each pRF
population (+pRFs in SPAs, +pRFs and - pRFs in PMAs). Consistent with prior reports, we found that OPA
and PPA showed lower and upper field biases, respectively (Elevation bias OPA vs PPA: t(16)=2.42, p=0.02,
D=0.98) **(Fig. 4A)**. Critically, we found that the visual field representations of the -pRFs in memory areas
closely matched their paired perceptual counterparts **(Fig. 4B)**. Specifically, -pRFs in LPMA were biased
towards the lower visual field (matching OPA), whereas -pRFs in VPMA were biased towards the upper
visual field (matching PPA) (Elevation bias -LPMA vs -VPMA: t(11)=4.53, p=0.0008, D=0.94). In contrast,
the +pRF populations in LPMA/VPMA did not show biases that corresponded to their perceptual
counterparts: +pRFs in LPMA were biased towards the upper visual field, the opposite of OPA, and +pRFs
in VPMA showed no clear elevation bias (Elevation bias +LPMA vs +VPMA: t(16)=2.78, p=0.01, D=0.65).
These data show that the -pRFs in the PMAs represent similar visual field locations as their paired SPAs,
and based on the known functional interaction between the paired PMAs and SPAs during scene
processing, we suggest that the visual field representation may be, in part, inherited from feedforward
connections originating in the SPAs.

*Figure 4. Shared visual field representations between paired perception and memory areas. **A.** Ventral and*
lateral SPAs/PMAs differentially represent the upper (ventral areas) and lower (lateral areas) visual fields.
Group visual field coverage plots for SPAs and PMAs. X-axis represents ipsilateral to contralateral visual field

Second, if, as hypothesized, +/- pRF populations reflect bottom-up sensory and top-down internal mnemonic processes, we predicted that the sign of the spatially-specific opponent interactions observed during visual mapping (Exp. 1) would reverse during a top-down memory paradigm. Specifically, Exp. 1 showed that when activity was high in +pRFs within the SPAs, activity was low in -pRFs within the PMAs. In contrast, we hypothesized that during a recall task, when -pRF activity would be high, +pRF activity would be low. To test for this competitive interaction during recall, in Experiment 2, we examined the activation profile of +/- pRFs during a place memory task, wherein participants recalled personally familiar visual environments (e.g., their kitchen, **Fig. 5A**). For this analysis, we first partitioned LPMA and VPMA into their +pRF and -pRF populations, respectively. Then, for each subject, we calculated the average activity (activation values versus baseline) from each of the 36 trials of the memory task for each population and ROI (e.g., -pRFs in the place memory areas, +pRFs scene perception area) and z-scored the activation values in each ROI. We compared the z-scored values using a partial correlation between the -pRF/+pRFs in the PMAs and the SPAs (e.g., correlation between -pRFs in LPMA with +pRFs in LSPA, while controlling for +pRFs in LPMA, and vice versa) for each subject. Partial correlation allowed us to compare the unique impact of these distinct neural populations, while simultaneously controlling for non-specific effects (like motion and attention) that impact beta estimates on each trial. We compared the partial correlation values from the different pRF populations (i.e., -LPMA x OPA vs +LPMA x OPA) for all subjects using paired t-tests separately for the lateral and ventral surfaces.

As predicted, we observed an opponent relationship between activation of the -pRFs in the PMAs and the +pRFs in the SPAs during the place memory recall task (-LPMA x OPA vs +LPMA x OPA: t(16)=3.10, p=0.0006, D=1.52; -VPMA x PPA vs +VPMA x PPA: t(11)=5.27, p<0.0001, D=2.98). Specifically, on trials where the activity of -pRFs in PMAs was high, SPA activation was reduced **(Fig. 5B)** (t-test vs. zero – lateral t(16)=-2.19, p=0.04, D=0.53; ventral: t(11)=-3.63, p=0.003, D=1.04). In contrast, the +pRFs in the PMAs did not show a push-pull dynamic: on trials where activity of +pRFs in the PMA was high, activation of the SPA also increased (t-test vs. zero: lateral: t(16)=3.76, p=0.001, D=1.69); ventral: t(11)=5.88, p=0.0001, D=1.69) **(Fig. 5B)**. We found a similar pattern when we considered all trials from all subjects pooled together (**Fig. 5B insets, Fig. S4**).

Is the interaction between perceptual and mnemonic spatially specific? Importantly, if the interaction between the PMAs and SPAs is structured by a retinotopic code, we expect that the opponent dynamic between -pRFs in the PMAs with +pRFs in the SPAs would be stronger between pRFs representing shared regions of visual space. To test this, we again compared the partial correlation in trial x trial activation between +/- pRFs of the perception and memory areas, this time focusing on subpopulations of +/- pRFs with matched (versus unmatched) visual field representations (in x, y, and sigma) (see Methods; **Fig. 5C**). Because of our stringent pRF matching criteria (see Methods), the overall number of subjects included in this analysis was reduced (Lateral: N=14; Ventral: N=7).

This analysis revealed that the inhibitory interaction between -pRFs in PMAs and +pRFs in SPAs is, in fact, structured by a retinotopic code (i.e., is spatially specific) **(Fig 5D, 5E)**. On both the lateral and ventral surfaces, the anti-correlation in trial x trial activity during recall is stronger for matched vs unmatched pRFs (Lateral: Amplitude x Matching: $F(1, 13)=4.64$, $p=0.05$; $t(13)=2.15$, $p=0.05$, $D=0.77$; Ventral: Amplitude x Matching: $F(1, 6)=5.96$, $p=0.05$; $t(6)=2.44$, $p=0.05$, $D=0.60$). Importantly, this interaction was driven by the sign of the association between -pRFs and + pRFs in the PMAs with the SPA +pRFs: the correlation between trial x trial activation was significantly stronger (i.e., more negative) for matched vs unmatched -pRFs on the lateral and ventral surfaces: (Lateral: $t(13)=2.48$, $p_{one-tailed}=0.01$; Ventral: $t(6)=2.06$, $p_{one-tailed}=0.04$), and we found a similar trend for +pRFs (i.e., stronger positive correlation (Lateral: $t(13)=1.62$, $p_{one-tailed}=0.06$; Ventral: $t(6)=2.45$ $p_{one-tailed}=0.02$). These data support the hypothesis that the spatially-specific opponent relationship between -pRFs in the PMAs and +pRFs in the SPAs evidenced during bottom up visual stimulation (i.e., pRF mapping in Exp. 1) is reversed during top-down memory recall. Specifically, -pRFs in the PMAs (defined during pRF mapping) exert top-down inhibition of pRFs in the SPAs representing similar regions of visual space.

Figure 5. *Positive pRFs in scene perception areas (SPAs) and negative pRFs in place memory areas (PMAs) exhibit a spatially-specific push-pull dynamic during memory recall.* ***A.*** *Recall task paradigm. In Exp. 2,*

*participants visually recalled personally familiar places for 10 s (cued with stimulus name; 36 trials per participant), and we calculated activation versus baseline at each trial. **B.** Negative and positive pRF (-pRF, +pRF) populations in SPAs and PMAs exhibit an opponent interaction during recall (Lateral: t(13)=3.30, p=0.005; Ventral: t(11)=5.66, p=0.00001). On both the lateral and ventral surfaces, when activity of -pRFs in the PMA is high, activity in the corresponding SPA is reduced (t-test versus zero – Lateral: t(14)=-2.12, p=0.05; Ventral t(14)=-3.08, p=0.01). In contrast, when activity in +pRFs in a PMA is high, activity in the adjacent SPA is also high (t-test versus zero – Lateral: t(14)=4.22, p=0.0004; Ventral: t(14)=7.37, p=0.000006). Inset scatter plots show normalized trial-wise activation of each region for all trials in all participants. **C.** Schematic of pRF matching between OPA with -pRFs in LPMA. Schematic pRFs are depicted as circles. The OPA pRF is shown in orange, along with its best matched -pRF in LPMA (shown in dark blue) and randomly matched -pRF in LPMA (light blue). Inset. Histogram indicates the dice coefficients for iterative matching between OPA pRFs and randomly paired pRFs from LPMA (light blue) compared to matched pRFs from LPMA (dark blue) for one example participant, showing better correspondence between matched vs random, "unmatched" pRFs. This pattern was consistent in all participants (ts>6, ps<0.0001). **D,E.** Matched pRFs had a significantly stronger association in trial x trial activation than unmatched pRFs on the lateral **(D)** and ventral **(E)** surfaces (Lateral: Amplitude x Matching: F(1, 13)=4.64, p=0.05; Ventral: Amplitude x Matching: F(1, 6)=5.96, p=0.05) which suggests that the interaction between the perceptual and memory areas is spatially specific (Lateral: t(13)=2.15, p=0.05, D=0.77; Ventral: t(6)=2.44, p=0.05, D=0.60). In other words, on trials where the activity of a PMA -pRF was relatively high, the activity of its spatially-matched SPA +pRF was relatively low. Importantly, this interaction was driven by the sign of the association between -pRFs and + pRFs the PMAs with the SPAs: the correlation between trial x trial activation was significantly stronger (i.e., more negative) for matched vs unmatched -pRFs on the lateral and ventral surfaces: (Lateral: t(13)=2.48, $p_{one-tailed}$=0.01; Ventral: t(6)=2.06, $p_{one-tailed}$=0.04), and we found a similar trend for +pRFs (i.e., stronger positive correlation (Lateral: t(13)=1.62, $p_{one-tailed}$=0.06; Ventral: t(6)=2.45 $p_{one-tailed}$=0.02). *: p < 0.05, **: p<0.01, ***: p<0.001.*

Although the spatially-specific push-pull dynamic observed between +/- pRFs during both perception (Exp. 1) and recall (Exp. 2) is compelling, these two tasks represent situations wherein activation of the visual and memory systems should be maximally opposed (i.e., focusing attention externally on a visual stimulus vs. focusing attention internally during visual recall). Do -pRFs in the PMAs and +pRFs in the SPAs interact in a mutually inhibitory fashion in contexts where perceptual and memory systems are expected to interact, such as during familiar scene perception? As a final test of the interaction between the SPA and PMA pRFs, we asked whether the mutually inhibitory interaction between -pRFs in the PMAs and +pRFs in the SPAs is evident when participants view images of locations that were familiar to them in real life. We tested two possible hypotheses. On one hand, processing familiar scenes could be mutually excitatory for both the memory and perception areas, resulting in a positive trial x trial relationship between the SPAs and both +pRFs and -pRFs in PMAs. On the other hand, the opponent interaction might persist if the SPAs and -pRFs in the PMAs were interlocked in a mutually inhibitory relationship, like predictive coding [33–35].

For this experiment (Exp. 3), a subset of participants (N=8) passively viewed portions of images (lower left and right quadrants) of recognizable Dartmouth College landmarks **(Fig. 6A).** We focused on the lateral surface (OPA and LPMA), because -pRFs could be localized most robustly in LPMA in our main sample of participants. Familiar scene stimuli were presented in the lower quadrants to maximally stimulate these areas, and we specifically considered +/- pRFs with centers in the contralateral lower visual field. For this analysis, we tested whether the -pRFs in LPMA and +pRFs in OPA are correlated or anti-correlated during familiar scene perception. We hypothesized that activation of +pRFs in OPA would be inversely related to activation of -pRFs in LPMA, consistent with a mutually inhibitory interaction. Further, we predicted that this interaction would be spatially specific: the inhibitory dynamic should be stonger when visual

information was presented in the contralateral lower visual field (i.e., preferred quadrant), relative to the
ipsilateral lower visual field (i.e., non-preferred quadrant).

We found that the spatially-specific, inhibitory interaction between +pRFs in OPA with -pRFs in LPMA
persists during familiar scene viewing (Interaction between visual field and pRF association: $F_{(1,7)}=8.27$,
p=0.02; **Fig. 6**). The trial x trial opponent dynamic between OPA pRFs, and -pRFs and +pRFs in the PMAs
differed by visual field (t(7)=2.79, p=0.026, D=0.98). As we observed in recall, this dynamic was driven by
the opposing sign of the relationship between -/+ pRFs. We found that the relationship between pRFs in
OPA and -pRFs in LPMA was negative in 6/8 participants in the contralateral visual field, and the strength
of the association between OPA and -pRF activity for stimuli in the contralateral visual field was
significantly stronger than the ipsilateral visual field (t(7)=2.71, p=0.03, D=0.95). In contrast, the
relationship between OPA and +pRFs in LPMA was positive in 8/8 participants, and was numerically
stronger for the contralateral hemifield (t(7)=1.75, p=0.12, D=0.62). Together, the opposing responses
during visual stimulation (Exp. 1 & 3) as well as during recall (Exp. 2), strongly suggest that the spatially-
specific, mutually inhibitory interaction between -pRFs in mnemonic regions and +pRFs in perceptual
regions is a generalizable description of their interaction that extends to naturalistic contexts, like familiar
scene perception.

Figure 6. *Negative and positive pRFs in the place memory areas exhibit a push-pull dynamic during perception*
of familiar scene images. ***Right.*** *In Exp. 3, participants passively viewed portions of images depicting two*
familiar Dartmouth College landmarks in the lower visual field. We investigated the trial x trial correlation in
activation between +pRFs OPA and -/+ pRFs in LPMA. ***Left.*** *-pRFs in LPMA exhibit an opponent interaction*
with +pRFs in OPA. 6/8 participants exhibited a negative correlation between these pRF populations, and this
relationship was significantly stronger when stimuli were presented in the contralateral compared to
ipsilateral visual field. In contrast, activation of the +pRFs in LPMA were significantly positive for both
*contralateral and ipsilateral visual field presentations. *: p < 0.05, ***: p<0.001.*

To summarize, in contrast to classic models of brain organization, which generally assume that retinotopic
coding is replaced by amodal, abstract codes as information propagates through the visual hierarchy
towards memory structures [7,8,11], we observe that retinotopic coding persists anterior to regions classically

considered visual in the form of -pRFs. The shared retinotopic code between perceptual and mnemonic areas structures their mutual activity, such that pRFs representing similar areas in visual space have highly anti-correlated activation during both bottom-up scene perception and also top-down memory recall. Together with recent reports showing retinotopic coding persisting as far as the 'cortical apex'[14,15,36], including the default mode network[14,15], as well as the hippocampus[36,37], our work suggests that retinotopic coding may play an integral role in structuring information processing across the brain.

Understanding how mnemonic and sensory representations functionally interface in the brain while avoiding interference is a central puzzle in neuroscience[38–41]. Theoretical work has shown that representational interference can be reduced through operations like orthogonalization[14,38,42], pattern separation[42,43], and areal segregation[42], but the principles that preserve interaction across mnemonic and perceptual populations are less clear. Our findings speak to both sides of this puzzle. On one side, within the functionally-defined memory areas, perceptual and mnemonic information may be maintained in distinct, spatially separated +/- pRF populations, avoiding interference. On the other side, the shared retinotopic code between the SPAs and PMAs enables functional interaction between their representations. In particular, the observed mutually inhibitory dynamic between the -pRFs in memory areas and +pRFs in perceptual areas suggests a specific class of functional interaction, where top-down predictions cancel out bottom-up sensory information consistent with those predictions[33–35]. These top-down retinotopic signals from memory might help bias perception towards probable scene views to enable efficient visually guided actions[44,45].

This conclusion has significant implications for our systems-level understanding of information processing at the cortical apex. Our work demonstrates that high-level brain areas approaching the cortical apex[9,10] explicitly represent visual information in the environment: these regions have an inverted retinotopic code[14,15] that organizes their functional interaction with visual areas. This view contrasts strikingly with the view of the cortical apex as an amodal and "internally-oriented" neural system. Large-scale deactivation of mnemonic areas at the cortical apex (e.g., the DMN) during visual tasks and activation during internally-focused tasks is among the most striking and widely-replicated network-level patterns in functional neuroimaging[46–52]. Critically, the deactivation within the DMN was not thought to represent stimulus information, but rather the negative response resulted from a trade-off between internal and externally oriented neural processes[48–50,53,54]. In contrast, our results suggest that the opponent activation profiles of DMN/non-DMN brain areas may reflect a functional interaction between mnemonic and perceptual areas that are involved in processing the same stimulus, rather than a trade-off between dissociable processes (like internal and external cognition). Thus, it may be necessary to reconsider the default mode network's role in tasks that require externally-directed attention, as well as what representational content might be conveyed in negative BOLD responses.

How general is retinotopic coding for structuring perceptual-mnemonic interactions? On the one hand, because visually guided navigation has unique mnemonic demands (e.g., representing visual information out of view) scene areas might have privileged access to memory information. Indeed, the scene memory and perception areas may uniquely span the boundary of DMN/visual cortex (**Fig. 1B, S3**)[22]. Likewise, these unique mnemonic processes could lead to preferential coding of retinotopic information in scene memory areas, making the opponent interaction observed here "scene specific". On the other hand, the present

work and others[14,36,37] show that pRFs, including -pRFs, are widely distributed throughout the brain. As such, retinotopic representations could be well-poised to structure interactions between many functionally coupled brain areas broadly in cortex beyond the domain of scenes, including other functional subnetworks situated within the default mode network[22,29,30,55–59]. Relatedly, parietal, occipital, and temporal areas utilize multiple visuospatial coding motifs[2–4,6,60–63], including retinotopic[1], spatiotopic[2–4], and head/body-centered formats[60–63], suggesting that the format of visual spatial coding within a region may depend on the computational roles a given region plays. Thus, while our data clearly shows the mutual retinotopic code structures the shared activity between paired perceptual and mnemonic areas, future studies will be necessary to further elaborate the nature of the visuospatial coding across cognitive domains and sensory modalities.

Surprisingly, the PMAs contained a large proportion of +pRFs that, unlike the -pRFs, did not share the biased visual field representations of their SPA counterparts. This raises two questions. First, what regions contribute visual information to the memory areas' +pRFs? While the present data offers no definitive answer, it is possible that they represent information arriving locally from other visual areas like the visual maps in the dorsal intraparietal sulcus[64] or via longer range connections like the vertical occipital fasciculus[65]. Second, what do the +pRFs contribute to the PMAs' functions? The robust activation of both +pRFs and -pRFs in the PMAs during memory recall suggests that these populations have related activity and differ primarily in their response to visual stimulation. Notably, the PMAs are defined based on a single response property (response during recall of places versus people), and functional regions may contain sub-regions with different functional, structural, and cytoarchitectonic characteristics (similar to the presence of multiple retinotopic maps within larger, functionally-defined visual regions like OPA[66]). Future work is needed to elucidate what differential roles the +/- pRF in the PMAs might play.

To conclude, our results suggest that retinotopic coding is used in regions spanning the cortical hierarchy, from visual areas into the default mode network. Maintenance of a retinotopic code in high-level cortex, alongside other visuospatial, multisensory, semantic, and abstract codes, may be crucial for scaffolding communication between perceptual and mnemonic cortical areas.

Methods

Participants

17 adults (13 female; age=22.8±3.5 STD years old) completed the fMRI experiments. Participants had normal or correct-to-normal vision, were not colorblind, and were free from neurological or psychiatric conditions. Written consent was obtained from all participants in accordance with the Declaration of Helsinki and with a protocol approved by the Dartmouth College Institutional Review Board (Protocol #31288).

Visual Stimuli and Tasks

Population Receptive Field Mapping (Experiment 1)

During pRF mapping sessions, a bar aperture traversed gradually through the visual field, whilst revealing randomly selected scene fragments from 90 possible scenes. During each 36 s sweep, the aperture took 18 evenly spaced steps every 2 s (1 TR) to traverse the entire screen. Across the 18 aperture positions all 90 possible scene images were displayed once. A total of eight sweeps were made during each run (four orientations, two directions). Specifically, the bar aperture progressed in the following order for all six runs: Left to Right, Bottom Right to Top Left, Top to Bottom, Bottom Left to Top Right, Right to Left, Top Left to Bottom Right, Bottom to Top, and Top Right to Bottom Left). The bar stimuli covered a circular aperture (diameter = 11.4° of visual angle). Participants performed a color detection task at fixation, indicating via button press when the white fixation dot changed to red. Color fixation changes occurred semi-randomly, with approximately two-color changes per sweep. Stimuli were presented using PsychoPy (version 3.2.3) [67].

Scene Perception Area Localizer

The scene perception areas (SPAs, i.e. occipital place area, OPA; parahippocampal place area, PPA) are defined as regions that selectively activate when an individual perceives places (i.e., a kitchen) compared with other categories of visual stimuli (i.e., faces, objects, bodies) [22,27,55,68]. To identify these areas in each individual, participants performed an independent functional localizer scan. On each run of the localizer (2 runs), participants passively viewed blocks of scene, face, and object images presented in rapid succession (500 ms stimulus, 500 ms ISI). Blocks were 24 s long, and each run comprised 12 blocks (4 blocks/condition). There was no interval between blocks.

Place Memory Area Localizer (Recall task; Experiment 2)

The place memory areas (PMAs) are defined as regions that selectively activate when an individual recalls personally familiar places (i.e., their kitchen) compared with personally familiar people (i.e., their mother)[22]. To identify these areas in each individual, participants performed an independent functional localizer. Prior to fMRI scanning, participants generated a list of 36 personally familiar people and places to establish individualized stimuli (72 stimuli total). These stimuli were generated based on the following instructions.

> "For your scan, you will be asked to visualize people and places that are personally familiar to you. So, we need you to provide these lists for us. For personally familiar people, please choose people that you know in real life (no celebrities) that you can visualize in great detail. You do not need to be contact with these people now, as long as you knew them personally and remember what they look like. So, you could choose a childhood friend even if you are no longer in touch with this person. Likewise, for personally familiar places, please list places that you have been to and can richly visualize. You should choose places that are personally relevant to you, so you should avoid choosing places that you have only been to one time. You should not choose famous places where you have never been.

You can choose places that span your whole life, so you could do your current kitchen, as
well as the kitchen from your childhood home."

During fMRI scanning, participants recalled these people and places. On each trial, participants
saw the name of a person or place and recalled them in as much detail as possible for the duration
that the name appeared on the screen (10 s). Trials were separated by a variable ISI (4-8 s). Place
memory areas were localized by contrasting activity when participants recalled personally
familiar places compared with people (see ROI definitions section).

Perception of Familiar Scenes (Experiment 3)

A subset of participants in the main experiment (n=9) took part in this experiment, which involved
viewing an image of two prominent Dartmouth College buildings (Baker Library and Rollins
Chapel; one image per landmark). One participant was excluded for lack of familiarity with one
building (Rollins Chapel). All remaining participants were familiar with the landmarks and had
lived in the Hanover area for at least one year.

During scanning, participants passively viewed the familiar scene images. On each trial,
participants maintained fixation while passively viewing the lower left or lower right quadrant of
each image (display time: 1s). Images subtended as much of the whole lower visual field quadrant
possible (0°- 8° visual angle). We focused on the lower quadrants to maximally stimulate OPA
and LPMA. Prior to scanning we showed participants the full image of each location to familiarize
them with the specific image they would be seeing. Each image was presented eight times per
location per run. Trials were separated by a variable ISI (4-8 s). We collected two imaging runs,
resulting in 32 trials for each visual field location.

fMRI Data Processing

MRI Acquisition

All data was collected at Dartmouth College on a Siemens Prisma 3T scanner (Siemens, Erlangen,
Germany) equipped with a 32-Channel head coil. Images were transformed from dicom to nifti
format using dcm2niix (v1.0. 20190902) [69].

T1 image

For registration purposes, a high-resolution T1-weighted magnetization-prepared rapid
acquisition gradient echo (MPRAGE) imaging sequence was acquired (TR = 2300 ms, TE = 2.32 ms,
inversion time = 933 ms, Flip angle = 8°, FOV = 256 × 256 mm, slices = 255, voxel size = 1 ×1 ×
1 mm). T1 images segmented and surfaces were generated using Freesurfer [70–72] (version 6.0)
and aligned to the fMRI data using align_epi_anat.py and @SUMA_AlignToExperiment[73].

Functional MRI Acquisition

FMRI data were acquired using a multi-echo T2*-weighted sequence. The sequence parameters
were: TR=2000 ms, TEs=[14.6, 32.84, 51.08], GRAPPA factor=2, Flip angle=70°, FOV=240 x 192

mm, Matrix size=90 x 72, slices=52, Multi-band factor=2, voxel size=2.7 mm isotropic. The initial
two frames of data acquisition were discarded by the scanner to allow the signal to reach steady
state.

Preprocessing

Multi-echo data processing was implemented based on the multi-echo preprocessing pipeline
from afni_proc.py in AFNI[74]. Signal outliers in the data were attenuated (3dDespike[75]). Motion
correction was calculated based on the second echo, and these alignment parameters were
applied to all runs. The optimal combination of the three echoes was calculated, and the echoes
were combined to form a single, optimally weighted time series (T2smap.py). Multi-echo ICA
denoising[18,76–78] was then performed (see *Multi-echo ICA*, below). Following denoising, signals
were normalized to percent signal change.

Multi-echo ICA

The data were denoised using multi-echo ICA denoising (tedana.py[18,77,78]). In brief, PCA was
applied, and thermal noise was removed using the Kundu decision tree method. Subsequently,
data were decomposed using ICA, and the resulting components were classified as signal and
noise based on the known properties of the T2* signal decay of the BOLD signal versus noise.
Components classified as noise were discarded, and the remaining components were
recombined to construct the optimally combined, denoised timeseries.

Population Receptive Field Modeling

Detailed description of the pRF model implemented in AFNI is provided elsewhere[17]. Briefly,
given the position of the stimulus in the visual field at every time point, the model estimates the
pRF parameters that yield the best fit to the data: pRF amplitude (positive, negative), pRF center
location (x, y), and size (diameter of the pRF). Both Simplex and Powell optimization algorithms
are used simultaneously to find the best time-series/parameter sets (amplitude, x, y, size) by
minimizing the least-squares error of the predicted time-series with the acquired time-series for
each voxel. Relevant to the present work, the amplitude measure refers to the signed (positive
or negative) degree of linear scaling applied to the pRF model, which reflects the size of the neural
response to visual stimulation of its receptive field.

Sampling of fMRI Data to the Cortical Surface

In each participant, the analyzed functional data were projected onto surface reconstructions of
each individual participant's hemispheres in the SUMA standard mesh (std.141[79]) derived from
the Freesurfer autorecon script using the Surface Mapping with AFNI (SUMA[73]) software and the
3dvol2surf commands.

Regions of Interest (ROI) Definition

Scene perception regions (OPA, PPA) were established using the same criterion used in our prior work[22]. Scene perception areas were drawn based on a general linear test comparing the coefficients of the GLM during scene versus face blocks. Comparable results were observed when identifying the SPAs by comparing scene versus object blocks. A vertex-wise significance of p < 0.001 along with expected anatomical locations was used to define the regions of interest[68,80].

To define category-selective memory areas, the familiar people/places memory data was modeled by fitting a gamma function of the trial duration for trials of each condition (people and places) using 3dDeconvolve. Estimated motion parameters were included as additional regressors of no-interest. Place memory areas were drawn based on a general linear test comparing coefficients of the GLM for people and place memory. A vertex-wise significance threshold of p < 0.001 was used to draw ROIs.

To control for differing ROI sizes across regions and people, we restricted our analysis to 300 vertices centered on the center of mass of the thresholded regions. Consistent with prior work, we chose 300 vertices to ensure that no region or participant disproportionately contributed to any effects[22]. Results were qualitatively similar when 600 vertices were considered, suggesting our findings did not depend on ROI size.

ROI Analysis of pRF Amplitude

To calculate the percentage of -pRFs in each ROI we applied the following procedures. First, pRFs were thresholded on variance explained by the pRF model ($R_2 > 0.08$), which is consistent with prior work using R^2 thresholds ranging between 0.05 to 0.1[15,37,81,82]. Importantly, our results were consistent at R^2 thresholds between 0.05 to 0.2. Next, to avoid analyzing only very few pRFs within an ROI, only ROIs consisting of >25 suprathreshold pRFs (>8.3 % of total ROI) were included. The percentage of suprathreshold pRFs with a negative amplitude within each ROI was then calculated and submitted for statistical analysis.

Visual Field Coverage

Visual field coverage plots represent the group average sensitivity of each ROI to different positions in the visual field. To compute these, individual participant visual field coverage plots were first derived. These plots combine the best Gaussian receptive field model for each suprathreshold voxel within each ROI. Here, a max operator, at each point in the visual field, the maximum value from all pRFs within the ROI. The resulting coverage plot thus represents the maximum envelope of sensitivity across the visual field. Individual participant visual field coverage plots were averaged across participants to create group-level coverage plots.

To compute the elevation biases we calculated the mean pRF value (defined as the mean value in a specific portion of the visual field coverage plot) in the contralateral upper visual (UVF) and contralateral lower visual field (LVF) and computed the difference (UVF-LVF) for each participant, ROI and amplitude (+/-) separately. A positive value thus represents an upper visual field bias,

whereas a negative value represents a lower visual field bias. Analysis of the visual field biases
considers pRF center location (like the center of mass calculation does), as well as pRF size and
R2. This makes mean pRF value a preferable summary metric than analyzing pRF center position
alone[17,37,55,66].

Recall Trial x Trial Analysis

To assess the interaction between +pRFs in SPAs and -pRFs in SPAs during memory recall, we
adopted the following procedures. For each participant and ROI, we sampled the pattern of
activity (t-value versus baseline) elicited during the recall of each personally familiar place (36
places per participant) from the place memory localizer experiment. Suprathreshold pRFs were
separated according to amplitude (+pRFs, -pRFs) before averaging the recall responses across
pRFs. This produced 36 responses (one for each recalled place) per pRF amplitude in each
participant/ROI. We then z-scored these values for each participant and ROI separately.

After normalization, in each participant, we computed the partial correlation (Pearson's R)
between responses during memory recall of +pRFs in SPAs with -pRFs in PMAs, while controlling
for the responses of +pRFs in PMAs and vice-versa. To determine whether the +pRF and -pRF
from the PMAs had differential influence on activity in the SPAs, we compared the correlation
coefficients from each population against each other using paired t-tests. To determine whether
the influence of the -pRFs and +pRFs was significant, we compared the Fisher's transformed
correlation coefficients from each population (+pRFs and -pRFs in the PMAs) against zero (no
correlation).

Spatial Specificity Analysis

First, we matched +pRFs in the scene perception areas with both +/- pRFs in the place memory
areas separately (e.g., OPA -> -LPMA & OPA -> +LPMA) using the following procedure. On each
iteration (1000 iterations total; randomized PMA pRF order): 1) For every pRF in a PMA, we
computed the pairwise Euclidean distance (in x, y, and sigma) to all +pRFs in the paired SPA and
found the SPA pRF with the smallest distance that was smaller than the median distance of all
possible pRF pairs, 2) we required that all pRF matches were uniquely matched, so if an SPA pRF
was the best match for two PMA pRFs, then the second PMA pRF was excluded. To prevent under
sampling, only subjects with greater than 10 matched -/+ pRFs on average in each region were
considered (Lateral surface: N=14, Ventral surface: N=7).

Second, we compared the correlation in trial x trial activation matched (versus non-matched)
pairs of pRFs with "unmatched" pRFs. To create the "unmatched", random pRF pairings, we
randomly sampled pRFs in the memory area (repeated 1000 times). We then computed the
unique correlation in trial x trial activation during recall between SPA pRFs and PMA pRFs, using
the same procedure as in our main analysis (e.g., the partial correlation between SPA pRFs with
PMA -pRFs, controlling for PMA +pRFs) for each iteration of the pRF matching. We compared the
mean of the Fisher transformed partial correlation values across the iterations for the matched
pRFs with the mean of random (i.e., unmatched) pRFs. To ensure that matched pRFs had better
corresponding visual field representations than unmatched pRFs, we calculated the visual field

overlap between pRF pairs in the matched samples, compared with the random samples (average
dice coefficient of the visual field coverage for all matched versus unmatched iterations.

Perception of Familiar Scenes Trial x Trial Analysis

To assess the interaction between pRFs in OPA and +/- pRFs in LPMA during perception, we
adopted the following procedure. We task-evoked activity using a GLM with each image
presentation fit as a separate regressor and calculated the average trial-wise activation of pRFs
in OPA and +/- pRFs in LPMA. Importantly, because of our interest in spatially-specific
interactions, we only considered pRFs with centers in the contralateral lower visual field (i.e., left
hemisphere OPA pRFs had to have centers in the lower right quadrant). We then Fisher-
transformed these values for each participant and ROI separately.

After normalization, in each participant we calculated the partial correlation between the trial x
trial activation of OPA with the negative and positive pRFs in LPMA separately for each hemifield
(i.e., OPA with -pRFs in LPMA, controlling for +pRFs in LPMA when images were presented in the
left hemifield). We compared the contralateral preference in the association (Fisher transformed
partial correlation values) between the +/- pRFs using a repeated measures ANOVA with visual
field (ipsilateral, contralateral) and pRF association (OPA x +LPMA, OPA x -LPMA) as factors.

Statistical Analysis

Statistics were calculated using the R Studio package (version 1.3)[83]. For each scene-perception
and scene-recall pair (e.g., OPA, LPMA), we predicted a greater percentage of negative pRFs in
the scene-recall (e.g., LPMA) compared with scene-perception regions (e.g., OPA). We conducted
repeated measures analysis of variance (rmANOVA) using the ezANOVA function from the 'ez'
package[84]. Alpha level of $p < 0.05$ was used to assess significance. All tests were two-tailed, except
for the spatial specificity analysis in which our strong *a priori* prediction warranted the use of
one-tailed tests. We applied Bonferroni correction for multiple comparisons where appropriate.

Supplemental Material

Determining reliability of pRF signed pRF amplitude (Odd/Even split-half analysis)

To quantify the reliability of pRF estimates, pRF models were computed on the average time-series of all odd runs (1, 3, 5) and even runs (2, 4, 6) separately, in each participant. For each ROI we pooled the pRF amplitude (positive, negative) of suprathreshold pRFs ($R^2 > 0.08$) across participants. First, we computed the percentage of pRFs in each ROI whose amplitude sign (i.e., positive or negative) remained consistent across splits. Importantly, we found that the signed amplitudes were highly stable across splits: for all ROIs greater than 78% of pRFs had consistent signed amplitude. Second, we computed the Pearson's correlation coefficient in amplitude across independent splits (Odd, Even). We observed a significant positive correlations all ROIs in both hemispheres (LH; OPA: $r(4404)=0.71$, $p<0.0001$; PPA: $r(3689)=0.55$, $p<0.0001$; LPMA: $r(2502)=0.67$, $p<0.0001$; VPMA: $r(1955)=0.32$, $p<0.0001$; RH; OPA: $r(4285)=0.83$, $p<0.0001$; PPA: $(4100)=0.79$, $p<0.0001$; LPMA: $r(2239)=0.48$, $p<0.0001$; VPMA: $r(2834)=0.69$, $p<0.0001$).

Individual participant amplitude maps

5 ***Figure S1.*** *Transition from positive to negative amplitude population receptive fields (+pRF, -pRF) moving*
6 *anteriorly from posterior cerebral cortex is evident in individual participants. Figure depicts amplitude maps*
7 *from all participants' left hemispheres. Only vertices surviving the threshold applied in the main text ($R^2 >$*
8 *0.08) are shown. Individual participant SPAs and PMAs used for analysis are drawn in white (PMAs) and*
9 *black (SPAs).*

Prominence of retinotopic coding within SPAs and PMAs

To quantify the extent to which retinotopic coding is expressed within each ROI we first calculated the percentage of suprathreshold pRFs ($R^2 > 0.08$) within our ROIs for each subject separately before testing each against a non-retinotopic prediction using t-tests (i.e., t-test versus zero, with Bonferroni correction). Retinotopic coding was significantly present within each ROI (LH; OPA: $t(12.51)=$, $p_{corr}=3.35^{-9}$, $D=2.99$; PPA: $t(16)=8.68$, $p_{corr}=5.67^{-7}$, $D=2.17$; LPMA: $t(16)=6.23$, $p_{corr}=3.58^{-5}$, $D=1.59$; VPMA: $t(16)=6.65$, $p_{corr}=1.64^{-5}$, $D=1.66$; RH; OPA: $t(16)=12.15$, $p_{corr}=5.10^{-9}$, $D=3.03$; PPA: $t(16)=11.97$, $p_{corr}=6.32^{-9}$, $D=3.12$; LPMA: $t(16)=8.75$, $p_{corr}=5.05^{-7}$, $D=2.18$; VPMA: $t(16)=6.39$, $p_{corr}=2.68^{-5}$, $D=1.55$).

Figure S2. *Retinotopic coding in SPAs and PMAs. Bars represent the mean percentage of suprathreshold pRFs ($R^2>0.08$) in each ROI/hemisphere for the lateral (left) and ventral (right) surfaces, respectively. Individual data points are overlaid. Each ROI exhibited a significant percentage of suprathreshold pRFs.*

1   Comparison of individual participant PMAs and SPAs to the default mode network

3   **Figure S3.** *Comparison between the location of the SPAs, PMAs, and default mode network in one*
4   *participant (example participant from Main text **Fig. 2**). This pattern was consistent in all individuals and at*
5   *the group level (**Main text Fig. 1B**). Default mode network defined using the Yeo et al., 2011 parcellation.*

Differential correlation in trial x trial activation during recall across all trials

Figure S4. *Differential interaction between pRFs in SPAs with -/+ pRFs in memory areas is evident across all*
trials. Each scatter plot and corresponding correlation values depict the unique correlation between pRFs in
the SPAs with -pRFs (blue) and +pRFs (red) in the PMAs (e.g., correlation between +pRFs in OPA with -pRFs
in LPMA, controlling for +pRFs in LPMA). Each data point represents the z-scored activation on a given trial
for all pRFS in the population (i.e., all -LPMA pRFS) for a given subject on a trial.

References

1. Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual Field Maps in Human Cortex. *Neuron* **56**, 366–383 (2007).

2. Pertzov, Y., Avidan, G. & Zohary, E. Multiple Reference Frames for Saccadic Planning in the Human Parietal Cortex. *The Journal of Neuroscience* **31**, 1059 (2011).

3. Golomb, J. D. & Kanwisher, N. Higher Level Visual Cortex Represents Retinotopic, Not Spatiotopic, Object Location. *Cerebral Cortex* **22**, 2794–2810 (2012).

4. Gardner, J. L., Merriam, E. P., Movshon, J. A. & Heeger, D. J. Maps of Visual Space in Human Occipital Cortex Are Retinotopic, Not Spatiotopic. *The Journal of Neuroscience* **28**, 3988 (2008).

5. Holmes, G. DISTURBANCES OF VISION BY CEREBRAL LESIONS. *Br J Ophthalmol* **2**, 353–384 (1918).

6. Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual field maps in human cortex. *Neuron* **56**, 366–383 (2007).

7. Popham, S. F. *et al.* Visual and linguistic semantic representations are aligned at the border of human visual cortex. *Nature Neuroscience 2021 24:11* **24**, 1628–1636 (2021).

8. Güçlü, U. & van Gerven, M. A. J. Deep Neural Networks Reveal a Gradient in the Complexity of Neural Representations across the Ventral Stream. *Journal of Neuroscience* **35**, 10005–10014 (2015).

9. Huntenburg, J. M., Bazin, P. L. & Margulies, D. S. Large-Scale Gradients in Human Cortical Organization. *Trends Cogn Sci* **22**, 21–31 (2018).

10. Margulies, D. S. *et al.* Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proc Natl Acad Sci U S A* **113**, 12574–12579 (2016).

11. Huth, A. G., De Heer, W. A., Griffiths, T. L., Theunissen, F. E. & Gallant, J. L. Natural speech reveals the semantic maps that tile human cerebral cortex. *Nature* **532**, 453–458 (2016).

12. Bellmund, J. L. S., Gärdenfors, P., Moser, E. I. & Doeller, C. F. Navigating cognition: Spatial codes for human thinking. *Science (1979)* **362**, eaat6766 (2018).

13. Groen, I. I. A., Dekker, T. M., Knapen, T. & Silson, E. H. Visuospatial coding as ubiquitous scaffolding for human cognition. *Trends Cogn Sci* **26**, 81–96 (2022).

14.	Szinte, M. & Knapen, T. Visual Organization of the Default Network. *Cerebral Cortex* **30**, 3518–3527 (2020).

15.	Christiaan Klink, P., Chen, X., Vanduffel, W. & Roelfsema, P. R. Population receptive fields in non-human primates from whole-brain fmri and large-scale neurophysiology in visual cortex. *Elife* **10**, (2021).

16.	Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual cortex. *Neuroimage* **39**, 647 (2008).

17.	Silson, E. H., Chan, A. W. Y., Reynolds, R. C., Kravitz, D. J. & Baker, C. I. A retinotopic basis for the division of high-level scene processing between lateral and ventral human occipitotemporal cortex. *Journal of Neuroscience* **35**, 11921–11935 (2015).

18.	Kundu, P., Inati, S. J., Evans, J. W., Luh, W. M. & Bandettini, P. A. Differentiating BOLD and non-BOLD signals in fMRI time series using multi-echo EPI. *Neuroimage* **60**, 1759–1770 (2012).

19.	Steel, A., Garcia, B. D., Silson, E. H. & Robertson, C. E. Evaluating the efficacy of multi-echo ICA denoising on model-based fMRI. *Neuroimage* **264**, 119723 (2022).

20.	Wang, L., Mruczek, R. E. B., Arcaro, M. J. & Kastner, S. Probabilistic maps of visual topography in human cortex. *Cerebral Cortex* **25**, 3911–3931 (2015).

21.	Shmuel, A. *et al.* Sustained Negative BOLD, Blood Flow and Oxygen Consumption Response and Its Coupling to the Positive Response in the Human Brain. *Neuron* **36**, 1195–1210 (2002).

22.	Steel, A., Billings, M. M., Silson, E. H. & Robertson, C. E. A network linking scene perception and spatial memory systems in posterior cerebral cortex. *Nature Communications 2021 12:1* **12**, 1–13 (2021).

23.	Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 1125–1165 (2011).

24.	Epstein, R. A. & Baker, C. I. Scene Perception in the Human Brain. *Annu Rev Vis Sci* **5**, annurev-vision-091718-014809 (2019).

25.	Dilks, D. D., Julian, J. B., Paunov, A. M. & Kanwisher, N. The occipital place area is causally and selectively involved in scene perception. *Journal of Neuroscience* **33**, 1331–1336 (2013).

26.	Hasson, U., Levy, I., Behrmann, M., Hendler, T. & Malach, R. Eccentricity bias as an organizing principle for human high-order object areas. *Neuron* **34**, 479–490 (2002).

27. Epstein, R. & Kanwisher, N. A cortical representation the local visual environment. *Nature* **392**, 598–601 (1998).

28. Gordon, E. M. *et al.* Precision Functional Mapping of Individual Human Brains. *Neuron* **95**, 791-807.e7 (2017).

29. Braga, R. M. & Buckner, R. L. Parallel Interdigitated Distributed Networks within the Individual Estimated by Intrinsic Functional Connectivity. *Neuron* **95**, 457-471.e5 (2017).

30. DiNicola, L. M., Braga, R. M. & Buckner, R. L. Parallel distributed networks dissociate episodic and social functions within the individual. *J Neurophysiol* **123**, 1144–1179 (2020).

31. Breedlove, J. L., St-Yves, G., Olman, C. A. & Naselaris, T. Generative Feedback Explains Distinct Brain Activity Codes for Seen and Mental Images. *Current Biology* 1–14 (2020) doi:10.1016/j.cub.2020.04.014.

32. Favila, S. E., Kuhl, B. A. & Winawer, J. Perception and memory have distinct spatial tuning properties in human visual cortex. *Nature Communications 2022 13:1* **13**, 1–21 (2022).

33. Bastos, A. M. *et al.* Canonical Microcircuits for Predictive Coding. *Neuron* **76**, 695–711 (2012).

34. Hindy, N. C., Ng, F. Y. & Turk-Browne, N. B. Linking pattern completion in the hippocampus to predictive coding in visual cortex. *Nature Neuroscience 2016 19:5* **19**, 665–667 (2016).

35. Mumford, D. & Lee, T. S. Hierarchical Bayesian inference in the visual cortex. *JOSA A, Vol. 20, Issue 7, pp. 1434-1448* **20**, 1434–1448 (2003).

36. Knapen, T. Topographic connectivity reveals task-dependent retinotopic processing throughout the human brain. *Proc Natl Acad Sci U S A* **118**, (2021).

37. Silson, E. H., Zeidman, P., Knapen, T. & Baker, C. I. Representation of Contralateral Visual Space in the Human Hippocampus. *Journal of Neuroscience* **41**, 2382–2392 (2021).

38. Libby, A. & Buschman, T. J. Rotational dynamics reduce interference between sensory and memory representations. *Nature Neuroscience 2021 24:5* **24**, 715–726 (2021).

39. Kiyonaga, A., Scimeca, J. M., Bliss, D. P. & Whitney, D. Serial dependence across perception, attention, and memory. *Trends Cogn Sci* **21**, 493 (2017).

40. Summerfield, C. & de Lange, F. P. Expectation in perceptual decision making: neural and computational mechanisms. *Nat Rev Neurosci* **15**, 745–756 (2014).

41. Rademaker, R. L., Chunharas, C. & Serences, J. T. Coexisting representations of sensory and mnemonic information in human visual cortex. *Nature Neuroscience 2019 22:8* **22**, 1336–1344 (2019).

42. Favila, S. E., Lee, H. & Kuhl, B. A. Transforming the Concept of Memory Reactivation. *Trends Neurosci* **43**, 939–950 (2020).

43. Yassa, M. A. & Stark, C. E. L. Pattern separation in the hippocampus. *Trends Neurosci* **34**, 515–525 (2011).

44. Albright, T. D. On the perception of probable things: neural substrates of associative memory, imagery, and perception. *Neuron* **74**, 227–245 (2012).

45. Ryan, J. D. & Shen, K. The eyes are a window into memory. *Curr Opin Behav Sci* **32**, 1–6 (2020).

46. Zabelina, D. L. & Andrews-Hanna, J. R. Dynamic network interactions supporting internally-oriented cognition. *Curr Opin Neurobiol* **40**, 86–93 (2016).

47. Shulman, G. L. *et al.* Common Blood Flow Changes across Visual Tasks: II. Decreases in Cerebral Cortex. *J Cogn Neurosci* **9**, 648–663 (1997).

48. Fox, M. D. *et al.* The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci U S A* **102**, 9673–9678 (2005).

49. Raichle, M. E. The Brain's Default Mode Network. *Annu Rev Neurosci* **38**, 433–447 (2015).

50. Christoff, K., Gordon, A. M., Smallwood, J., Smith, R. & Schooler, J. W. Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proc Natl Acad Sci U S A* **106**, 8719–8724 (2009).

51. Fox, K. C. R., Spreng, R. N., Ellamil, M., Andrews-Hanna, J. R. & Christoff, K. The wandering brain: Meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought processes. *NeuroImage* Preprint at https://doi.org/10.1016/j.neuroimage.2015.02.039 (2015).

52. Chun, M. M., Golomb, J. D. & Turk-Browne, N. B. A Taxonomy of External and Internal Attention. *https://doi.org/10.1146/annurev.psych.093008.100427* **62**, 73–101 (2010).

53. Smallwood, J. & Schooler, J. W. The science of mind wandering: Empirically navigating the stream of consciousness. *Annu Rev Psychol* **66**, 487–518 (2015).

54. Smallwood, J. *et al.* The neural correlates of ongoing conscious thought. *iScience* **24**, 102132 (2021).

55. Silson, E. H., Steel, A. D. & Baker, C. I. Scene-Selectivity and Retinotopy in Medial Parietal Cortex. *Front Hum Neurosci* **10**, 412 (2016).

56. Silson, E. H., Steel, A., Kidder, A., Gilmore, A. W. & Baker, C. I. Distinct subdivisions of human medial parietal cortex support recollection of people and places. *Elife* **8**, (2019).

57. Deen, B. & Freiwald, W. A. Parallel systems for social and spatial reasoning within the cortical apex. *bioRxiv* 2021.09.23.461550 (2021) doi:10.1101/2021.09.23.461550.

58. Ranganath, C. & Ritchey, M. Two cortical systems for memory-guided behaviour. *Nature Reviews Neuroscience* vol. 13 713–726 Preprint at https://doi.org/10.1038/nrn3338 (2012).

59. Ritchey, M. & Cooper, R. A. Deconstructing the Posterior Medial Episodic Network. *Trends Cogn Sci* **xx**, (2020).

60. Mullette-Gillman, O. A., Cohen, Y. E. & Groh, J. M. Eye-centered, head-centered, and complex coding of visual and auditory targets in the intraparietal sulcus. *J Neurophysiol* **94**, 2331–2352 (2005).

61. Fetsch, C. R., Wang, S., Gu, Y., DeAngelis, G. C. & Angelaki, D. E. Spatial Reference Frames of Visual, Vestibular, and Multimodal Heading Signals in the Dorsal Subdivision of the Medial Superior Temporal Area. *Journal of Neuroscience* **27**, 700–712 (2007).

62. Chen, X., DeAngelis, G. C. & Angelaki, D. E. Flexible egocentric and allocentric representations of heading signals in parietal cortex. *Proc Natl Acad Sci U S A* **115**, E3305–E3312 (2018).

63. Duhamel, J. R., Bremmer, F., BenHamed, S. & Graf, W. Spatial invariance of visual receptive fields in parietal cortex neurons. *Nature 1997 389:6653* **389**, 845–848 (1997).

64. Swisher, J. D., Halko, M. A., Merabet, L. B., McMains, S. A. & Somers, D. C. Visual Topography of Human Intraparietal Sulcus. *Journal of Neuroscience* **27**, 5326–5337 (2007).

65. Yeatman, J. D. *et al.* The vertical occipital fasciculus: A century of controversy resolved by in vivo measurements. *Proc Natl Acad Sci U S A* **111**, E5214–E5223 (2014).

66. Silson, E. H., Groen, I. I. A., Kravitz, D. J. & Baker, C. I. Evaluating the correspondence between face-, scene-, and object-selectivity and retinotopic organization within lateral occipitotemporal cortex. *J Vis* **16**, 14–14 (2016).

67. Peirce, J. W. PsychoPy—Psychophysics software in Python. *J Neurosci Methods* **162**, 8–13 (2007).

68. Weiner, K. S. *et al.* Defining the most probable location of the parahippocampal place area using cortex-based alignment and cross-validation. *Neuroimage* **170**, 373–384 (2018).

69. Li, X., Morgan, P. S., Ashburner, J., Smith, J. & Rorden, C. The first step for neuroimaging data analysis: DICOM to NIfTI conversion. *J Neurosci Methods* **264**, 47–56 (2016).

70. Fischl, B. FreeSurfer. *NeuroImage* vol. 62 774–781 Preprint at https://doi.org/10.1016/j.neuroimage.2012.01.021 (2012).

71. Fischl, B. *et al.* Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron* **33**, 341–355 (2002).

72. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical surface-based analysis: I. Segmentation and surface reconstruction. *Neuroimage* **9**, 179–194 (1999).

73. Saad, Z. S. & Reynolds, R. C. SUMA. *Neuroimage* **62**, 768–773 (2012).

74. Cox, R. W. AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research* **29**, 162–173 (1996).

75. Jo, H. J. *et al.* Effective Preprocessing Procedures Virtually Eliminate Distance-Dependent Motion Artifacts in Resting State FMRI. *J Appl Math* **2013**, (2013).

76. DuPre, E. *et al.* TE-dependent analysis of multi-echo fMRI with *tedana*. *J Open Source Softw* **6**, 3669 (2021).

77. DuPre, E. *et al.* ME-ICA/tedana: 0.0.6. Preprint at https://doi.org/10.5281/ZENODO.2558498 (2019).

78. Evans, J. W., Kundu, P., Horovitz, S. G. & Bandettini, P. A. Separating slow BOLD from non-BOLD baseline drifts using multi-echo fMRI. *Neuroimage* **105**, 189–197 (2015).

79. Argall, B. D., Saad, Z. S. & Beauchamp, M. S. Simplified intersubject averaging on the cortical surface using SUMA. *Hum Brain Mapp* **27**, 14–27 (2006).

80. Julian, J. B., Fedorenko, E., Webster, J. & Kanwisher, N. An algorithmic method for functionally defining regions of interest in the ventral visual pathway. (2012) doi:10.1016/j.neuroimage.2012.02.055.

81. Gomez, J., Barnett, M. & Grill-Spector, K. Extensive childhood experience with Pokémon suggests eccentricity drives organization of visual cortex. *Nature Human Behaviour 2019* 1 (2019) doi:10.1038/s41562-019-0592-8.

82. Gomez, J. *et al.* Development of population receptive fields in the lateral visual stream improves spatial coding amid stable structural-functional coupling. *Neuroimage* **188**, 59–69 (2019).

83. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computingitle. Preprint at http://www.r-project.org/ (2013).

84. Lawrence, M. A. ez: Easy analysis and visualization of factorial experiments. Preprint at (2016).