Gap junction-dependent amino acid transfer promotes germ cell growth

Caroline Vachias[1], Camille Tourlonias[1], Louis Grelée[1], Nathalie Gueguen[1], Yoan Renaud[1], Parvathy Venugopal[1], Graziella Richard[1], Pierre Pouchin[1], Émilie Brasset[1], Vincent Mirouse[1]*

1) Institute of Genetics, Reproduction and Development (iGReD), Université Clermont Auvergne - UMR CNRS 6293 - INSERM U1103, Faculté de Médecine, 28 Place Henri-Dunant, 63000 Clermont-Ferrand, France
*contact: vincent.mirouse@uca.fr

Abstract 180 words

Gap junctions allow the exchange of small molecules between cells. How this function could be used to promote cell growth has never been explored. During *Drosophila* ovarian follicle development, germ cells, which are surrounded by epithelial somatic cells, undergo massive growth. We found that this growth depends on gap junctions between these cell populations, with a requirement for Innexin4 and Innexin2, in the germ cells and the somatic cells, respectively. Somatic cells express enzymes and transporters involved in amino acid metabolism that are absent in germ cells. Among them, we identified a proline transporter required for germline growth. Its ectopic expression in the germline can compensate for its absence in somatic cells, confirming the metabolic exchange between these cell types. Moreover, in somatic cells, innexin2 expression and gap junction assembly are regulated by the insulin receptor/PI3K kinase pathway. Altogether these data illustrate how metabolic exchange through gap junctions can promote cell growth and how such mechanism can be integrated into a developmental programme, coupling growth control by extrinsic systemic signals with the intrinsic coordination between cell populations.

Key words

Drosophila, innexins, germ cells, metabolism, SLC36, P-bodies

 Running title

Cell growth control by gap junctions

Summary statement between 15 and 30 words,

We demonstrate that gap junction-dependent metabolic flow from a cell population to another promotes cell growth and can be regulated to coordinate cell growth.

Introduction

Gap junctions are channels between adjacent cells. Each cells harbour a hemichannel made of six subunits, connexins in vertebrates and innexins in other animal phyla where gap junctions are found. These two protein families are very different in terms of primary sequence, but similar in terms of conformation (Bauer et al 2005; Evans 2015; Oshima et al 2016; Skerrett, & Williams 2017). Hemichannels can be homomeric or heteromeric in composition and channels can be formed from two identical or different hemichannels, leading to a large range of channel combinations with different regulations or solute specificities. Gap junctions generally assemble in plaques that contain many channels and each channel allows the direct communication, and thus molecular flows, between the cytoplasm of two cells. However, due to the channel size and conformation, only passive transfer of small molecules and ions, up to approximatively 1KDa, including second messengers (e.g. inositol triphosphate and calcium) is allowed. Gap junction proteins also participate in other cellular mechanisms, such as cell migration (Elias et al 2007; Miao et al 2020). Moreover, many molecules that can diffuse through gap junctions are linked to energetic metabolism (e.g. glucose and pyruvate) and to anabolic metabolism (e.g. amino acids). However, despite many studies on these metabolite flows, their functional relevance has been elusive.

Such metabolite exchanges could promote cell growth. Available genetic data suggest that the growth of mammalian oocytes could be an illustration of such a mechanism. Indeed, mutation of connexin 37 (*Cx37*) in germ cells or of *Cx43* in somatic granulosa cells, which surround and are in contact with the oocyte, strongly impairs oocyte and follicle growth (Gittens, & Kidder 2005; Simon et al 1997). Moreover, amino acid import and pyruvate production are more efficient in follicle cells, suggesting a potential metabolic flow towards the oocyte (Eppig et al 2005; Sugiura et al 2005). Nonetheless, it is not known: i) to which extent this putative transfer of metabolites contributes to explain gap junction impact on oocyte growth; and ii) whether such mechanism is coupled with follicle cell growth and more generally, is integrated in the genetic programme controlling follicle development.

A major feature of oocytes is their large size throughout animal evolution, despite important variation among species (Doherty et al 2022; Church et al 2019). Their size prefigures the early embryo size and their content will determine the early embryo development success rate and quality (Petkova et al 2014; Burn et al 2015). Hence, oocyte growth is an essential step that requires robust underlying mechanisms (Doherty et al 2021; Vachias et al 2014b). Oocyte development and growth usually occur in ovarian follicles where germ cells are surrounded by follicle cells (i.e. somatic epithelial cells). *Drosophila* oogenesis takes place in a structure called ovariole, in which follicles continuously arise and develop from the anterior to the posterior end. Each ovary is subdivided in about 16 ovarioles (Fig. 1A). Follicles bud from a structure called the germarium that contains germ cells and somatic stem cells. Follicle development is divided in 14

morphological stages during which the oocyte volume increases 1000 times in about 3 days before being fertilized and laid (Spradling et al 1993). At stage 1, a follicle contains a germline cyst of 16 interconnected cells: 15 nurse cells that will grow by endoreplication and 1 oocyte at the posterior end, blocked in early meiosis. Each germline cyst is encapsulated by the follicular epithelium that is initially composed of about 30 cells. These cells proliferate giving 800 cells at stage 6 and then they undergo endoreplication (Chen et al 2019). Both germ cell growth and follicle cell growth are cell autonomously sensitive to the usual growth systemic signals, such as the Insulin Receptor/Phosphatidylinositol 3 Kinase (InR/PI3K) and Target of Rapamycin (TOR) pathways (Drummond-Barbosa, & Spradling 2001; LaFever et al 2010; Sun et al 2010; Vachias et al 2014b). These pathways affect also non-cell autonomously the growth of the adjacent tissue in both direction: the soma influences germline growth and *vice versa* (LaFever et al 2010; Maines et al 2004; Sun et al 2010; Vachias et al 2014b; Wang, & Riechmann 2007). The most striking illustration of this reciprocal coordination is observed when *Pten*, a InR/PI3K repressor, is mutated in follicle somatic cells. The faster growth of these somatic cells induces the faster growth of the surrounded wild-type germ cells. Regulation of somatic cell growth by germline growth involves the Hippo pathway and its modulation by tension induced on the epithelium due to the germline cyst volume increase (Borreguero-Muñoz et al 2019). However, it is not known how the follicle somatic cells control germ cell growth.

Here, we tested the hypothesis of an evolutionary conservation of gap junction function between germ cells and follicle cells to regulate *Drosophila* female germ cell growth and explored the underlying mechanism. Several reports indicate that gap junctions are present during fly oogenesis. In gem cells, innexin 4 (Inx4) forms plaques at the contact with follicle cells (Tazuke et al 2002; Bohrmann, & Zimmermann 2008; Sahu et al 2021). Several innexins, including Inx2, are expressed in follicle cells and partially colocalize with or are juxtaposed to Inx4 (Bohrmann, & Zimmermann 2008; Sahu et al 2021). Inx4 is required for germ cell survival before follicle formation, and Inx2 is involved in the germline cyst encapsulation by follicle cells during follicle budding (Gilboa et al 2003; Mukai et al 2011; Tazuke et al 2002). Inx2 and Inx4 are also required for morphogenetic events during oogenesis. such as the stretching of anterior follicle cells and the migration of border cells during stage 9 (Sahu et al 2021; Miao et al 2020). Moreover, Inx4 and Inx2 form gap junctions in the testis where they are required at different steps of sperm development (Tazuke et al 2002; Pesch et al 2022; Smendziuk et al 2015).

We found that Inx4/Inx2 gap junctions are required for female germline growth once follicles are formed. Moreover, follicle cells specifically express genes involved in amino acid biology, and proline import in follicle cells is necessary for germline growth. This germline growth control by somatic follicle cells is linked to the formation of processing bodies (P-bodies) and can be partially bypassed by the direct expression of a proline transporter in the germline. Moreover, gap junction

110   assembly is controlled by the InR/PI3K pathway, thus connecting growth coordination between

111   these two cell types with systemic growth control.

Results

Inx2 and Inx4 form gap junctions at the soma-germline interface that are required for germline growth

As gap junctions are required for oocyte growth in mammals, we asked whether they could play a similar role in *Drosophila* oogenesis (Gittens, & Kidder 2005; Simon et al 1997). Therefore, first, we characterized the gap junction composition and profile in somatic and germ cells, focusing on stages 1 to 8 when follicle development is globally limited to growth without major morphogenetic changes. As previously described (Tazuke et al 2002; Bohrmann, & Zimmermann 2008; Sahu et al 2021), we detected Inx2 and Inx4 forming plaques at the interface between germ and somatic cells. Inx2 also formed plaques on the lateral domain of follicle cells (Fig. 1B-E). Inx2 expression and plaque size tended to increase as follicles developed (Fig. S1). The cytoplasmic signal localization suggested that Inx2 and Inx4 were mainly expressed in the soma and germline, respectively. Accordingly, *Inx2* knock-down (*Inx2* RNAi) induction in somatic clones led to the cell-autonomous loss of Inx2 staining, whereas induction of *Inx4* knock-down (*Inx4* RNAi) in the germline led to the loss of Inx4 staining (Fig. 1F,G). Notably, we did not detect Inx4 at the contact of *Inx2* RNAi follicle cells, confirming published results (Fig. 1F''') (Sahu et al 2021). Similarly, when *Inx4* was knocked down in germline cysts, we did not detect Inx2 plaques at the germline contact, although Inx2 protein expression was increased in follicle cells (Fig. 1G'''). This observation could suggest a feedback mechanism between gap junction formation and Inx2 expression. Nonetheless, these data clearly established the presence of gap junction plaques composed of Inx4 in the germline and of Inx2 in follicle cells, and the Inx4-Inx2 interdependence for plaque assembly.

We then tested whether these gap junctions influenced germline growth by generating large *Inx2* mutant clones. In such conditions, we observed follicles that were smaller than the younger one at their anterior, a phenomenon never observed in the wild-type ovarioles. The quantification of the ratio between the volumes of the defective growth follicle and the follicle at its anterior confirmed these observations (Fig. 1H,I). We observed this phenotype with three different alleles, and only when all (or almost) epithelial cells of a follicle harboured the mutated *Inx2* while the anterior one was wild-type or contained a smaller percentage of mutant cells. A similar effect on the growth of the whole follicle was previously described for mutations affecting follicle cell growth (LaFever et al 2010; Maines et al 2004; Sun et al 2010; Vachias et al 2014b). However, follicles containing a mosaic of wild-type and *Inx2* mutant follicle cells grew normally and we did not observe any difference in cell size and in the proportion of cells in S phase (EDU-positive) between these populations, indicating that Inx2 did not influence somatic growth in a cell autonomous manner (Fig. 1J-K). This suggests that gap junctions are directly required for germline growth. We could not confirm this hypothesis using germline null *Inx4* mutant clones because in these mutants germline cyst development is blocked very early in the germanium (Tazuke et al 2002). Therefore,

150   we generated *Inx4* RNAi clones in the germline, directly detected by the absence of Inx4

151   expression (Fig. 1F). *Inx4* RNAi cysts were not larger than the younger wild-type follicle, indicating

152   defective germline growth (Fig. 1G). Altogether, these data indicate that Inx2 and Inx4 form

153   homomeric and heterotypic gap junctions between somatic and germ cells that are required for

154   germ cell growth.

Genes implicated in amino acid metabolism are enriched in somatic follicular cells

Gap junction requirement for germ cell growth supports a model in which metabolites diffuse from

follicle cells to germ cells. However, the direct identification of such metabolites is technically

challenging. We hypothesized that if follicle cells produce or import metabolites not present in the

germline, we might identify enzymes or transporters that are involved in this process and are

expressed only in follicle cells. To this aim, we performed translating ribosome affinity purification

(TRAP), an approach that allows identifying the tissue-specific translatome. We used

nanos:Gal4VP16 and *trafficjam:Gal4* drivers to express a GFP-tagged ribosomal protein

(UASp:RPL10a-GFP) specifically in the germline and in somatic cells, respectively (Fig. 2A). Then,

we immunoprecipitated the GFP-tagged polysomes and isolated and sequenced the associated

mRNAs under translation. Our biological replicates were highly reproducible (Table S1). To extract

genes with a specific or strongly enriched somatic expression we used a fold-change of 5 between

soma and germline as cut-off that gave a list of 811 genes (Fig. 2B). We then selected genes

involved in small molecule metabolic processes, reducing the list to 52 genes. The whole

ecdysone synthesis pathway (six genes), which is exclusively active in follicle cells, was enriched

in follicle cells, thus validating our TRAP approach (Domanitskaya et al 2014). Moreover, six genes

encoding enzymes involved in amino acid biosynthesis were specifically expressed in follicle cells

(Fig. 2B). By analysing this GO class, we found that 29 enzymes were expressed in both cell types,

but none was germline-specific (Fig. 2C). Similarly, among the 60 amino acid transporters

encoded by the fly genome, 21 were expressed in both tissues, 12 were specifically expressed in

the soma, and none in the germline alone (Fig. 2B and D). These data indicated that among the

genes specifically expressed in follicle cells there is a strong bias for genes involved in amino acid

synthesis or import. Given the importance of these metabolites for growth, they are good

candidates to be transferred through gap junctions.

Proline import in follicle cells is required for germline growth

Then, we performed a reverse genetic screen by inducing the silencing (RNAi) of the genes

involved in amino acid synthesis or transport identified by TRAP in follicle cells and then looking

for ovary growth defects. For each amino acid many redundancies may occur between anabolic

pathways or transporters that may preclude the observation of clear phenotypes. However, the

knock-down of one of the 18 tested genes reproductively induced an ovary growth defect in
independent RNAi lines (Fig. 3B-C). This gene, *CG43693*, was uncharacterized in *Drosophila* but
showed a clear homology with SLC36 family transporters, and was especially close to mammalian
SLC36A1 and SLC36A that are mainly involved in proline transport (Boll et al 2002; Metzner et al
2006; Pillai, & Meredith 2011).

We confirmed by RNA-FISH that *CG43693* was specifically expressed in follicle cells (Fig 3D-E).
Moreover, we detected *CG43693* mRNA in all follicle cells and at all stages of oogenesis, and its
expression progressively increased with the stage (Fig. 3D). A role in importing amino acids from
the haemolymph would require a cell membrane localization, while some amino acid transporters
are specific to some intracellular vesicular compartments. Therefore, to determine CG43693
protein localization, we generated a GFP knock-in with the MiMIC system to insert in-frame an
EGFP-encoding exon (Venken et al 2011). This insertion was in the non-conserved N-terminal part
common to five of the seven described *CG43693* isoforms. Importantly, it tags isoforms RA and
RB that were expressed in follicle cells according to our TRAP mRNA sequence data. The line
CG43693-GFP was homozygous viable and fertile without visible phenotype, suggesting that the
insertion did not affect protein function. We observed a strong GFP signal in the whole cortex of
follicle cells that tended to increase throughout oogenesis, but no signal in the germline (Fig 3F-
G). Thus, both CG43693 tissue distribution and subcellular localization were in agreement with
the hypothesis that it is implicated in the import of metabolites, such as proline, to promote follicle
growth.
We then tested whether CG43693 was required for follicle growth. The Minos element insertion
(*CG43693^{MI101960}*) used for the GFP knock-in initially contains an exon with STOP codons in
different frames. As it is located in the protein N-terminus, it should induce null mutation of the
affected isoforms, which include the ones expressed in follicle cells. This line was homozygous
lethal. Replacing the STOP codon-containing exon with the GFP exon suppressed lethality,
indicating that it was due to the gene mutation. We generated mutant clones in the follicular
epithelium. Analysis of small clones that did not cover the whole epithelium did not show any
difference in cell size and proliferation, suggesting that follicle cell growth was not affected (Fig.
3J,K). Conversely, we observed follicles smaller (or of the same size) than the younger wild-type
one when the whole epithelium of a follicle was mutated (Fig. 3H,I). As this growth defect led to
tiny and round follicles, we called this gene *cochonnet (coch)* after the nickname of the small wood
ball used with larger metal boules for pétanque, a traditional game in the South of France.
Importantly, these experiments indicated that *coch* expression in somatic follicle cells was
required for germline growth.
220
221

Genetic evidence of a metabolic transfer between soma and germline cells

Our results support a model in which proline imported in follicle cells diffuses through gap junctions to sustain germline growth. In this case, germline expression of *coch* should rescue genetic conditions in which it is absent in somatic follicle cells. To test this hypothesis we combined the UAS/Gal4 system with the QUAS/QF system (Potter et al 2010). We generated a matTub:QF driver and a QUASp promoter, inspired by the UASp promoter, for proper expression in the germline (Rørth 1998). The Q*UASp:GFP* transgene driven by matTub:QF was expressed in the germline at all stages, starting slightly earlier than what described for matTub:Gal4Vp16, indicating that both driver and QUASp promoter were functional (Fig. S2). Expression of the *QUASp:coch* transgene in the germline with the same driver (*MatTub>coch*) had a slight negative effect on ovary size (Fig. 4A-C,F). However, when it was combined with somatic knock-down of *coch* (*tj>cochRNAi),* it rescued the effect of the latter on ovary size (Fig. 4A,D-F)*.* These genetic data demonstrated that the requirement of *coch* expression in follicular cells for germline growth can be compensated by its germline expression, strongly arguing for a direct metabolic exchange between these cell types.

Blocking gap junctions or proline import induces processing bodies.

Previous studies indicated that somatic growth impairment affects germ cell development (LaFever et al 2010; Maines et al 2004; Sun et al 2010; Vachias et al 2014b). Moreover, the mRNA binding protein Me31B, which is typically concentrated where the anterior-posterior axis determinants localize in the oocyte, becomes enriched in large cytoplasmic structures in the germline when somatic growth is impaired (Burn et al 2015; Shimada et al 2011). These condensates are reminiscent of P-bodies and stress granules observed upon various stresses, including amino acid deprivation (Riggs et al 2020). In follicles, their formation can be induced by reducing protein availability in fly food and are more easily seen at stage 9 in Me31B-GFP-expressing follicles (Shimada et al 2011) (Fig. 5A,B). Therefore, we set up a semi-automated method for their quantification by measuring the fluorescence fraction found in condensates in stage 9 nurse cells (Fig. 5C). P-body formation is associated with translation arrest usually induced by inhibitory phosphorylation of the eukaryotic translation initiation factor 2 subunit alpha (eIF2α) on serine 51 (S51) **(Riggs et al 2020)**. Overexpression in the germline of a transgene mimicking this phosphorylated form (eIF2α-S51D), but not wild-type eIF2α, strongly induced P-body formation, although we did not quantify them because these follicles never reached stage 9 (Fig. 5D,E). Importantly, eIF2α-S51D overexpression was sufficient to completely block germline growth, suggesting a direct link between P-body formation and growth control (Fig. 5E). As P-bodies potentially repress germline growth, we tested whether the absence of gap junctions between germ cells and follicle cells or defective proline import in the follicle cells could induce their formation. We knocked down *Inx2* and *coch* in follicle cells using *tj:Gal4* in the presence of

Me31B-GFP. Constitutive *Inx2* RNAi in the somatic lineage completely blocked oogenesis and
precluded the observation of stages 8-9. Therefore, we added a constitutive *Gal80^ts* transgene to
allow expression of the RNAi construct only during a short period of time at the permissive
temperature (14 hours, at 30°C). *Coch* and *Inx2* knock-down strongly induced the formation of P-
bodies, suggesting that the observed growth limitation was linked to their formation and the
subsequent arrest of translation (Fig 5F-I).
Formation of P-bodies or stress granules following amino acid deprivation can occur
independently or dependently of eIF2α S51 phosphorylation. In the first case, which is not the
most frequently described, it is associated with the repression of the TOR pathway (Candena
Sandoval et al 2021). However, induction of *Tor* RNAi in the germline does not induce P-body
formation, excluding this possibility (Burn et al 2015). In the second case, eIF2α is phosphorylated
by GCN2 that acts as an indirect sensor of amino acid availability. We generated null mutant alleles
for *gcn2* that were homozygous viable, as similar alleles described during the course of this project
(Kosakamoto et al 2022; Srivastava et al 2022; Kang et al 2017). Protein deprivation in *gcn2*
transheterozygous females still led to P-body formation in the germ cells (Fig. S3A-E). Moreover,
an indirect reporter of GCN2 activity and eIF2α−S51 phosphorylation showed no signal in the
germline of wild-type flies, even after protein deprivation (Fig. S3F-G). Lastly, we tested more
directly *gcn2* role in the germline when proline import in follicle cells is blocked (*tj>cochRNAi*).
However, *gcn2* mutant germline clones did not increase the growth rate of wild-type and *coch*
RNAi follicles (Fig. S3H,I). These results strongly argued against the implication of GCN2 as a germ
cell growth repressor when proline availability is reduced.
Altogether these data indicate that although the two well-established pathways involved in amino
acid sensing do not seem implicated, P-body formation is linked to germ cell growth control by
gap junctions and proline import in follicle cells.
283
Gap junction assembly links intrinsic growth coordination and systemic control
Published data indicated that InR/PI3K pathway inhibition in follicle cells induces the formation of
P-bodies in the germline (Burn et al 2015). This result was reproduced by silencing (RNAi) *akt*, an
essential actor of this pathway, in follicle cells (Fig. 6A-C). Since somatic activity of the InR/PI3K
pathway also strongly influences germline growth, we asked whether there was a link between
the InR/PI3K pathway activity in follicle cells and their ability to transfer metabolites to the
germline. Coch-GFP expression level and localization were similar in wild-type and in follicle cells
with a PI3K gain of function or mutated for *akt,* suggesting no impact of this pathway on proline
import in follicle cells (Fig. S4A,B). Conversely, in *akt* mutant cells*,* Inx2 was almost undetectable
(Fig. 6D), whereas Inx2 expression was strongly increased in *Pten* mutant cells and plaques size
was increased (Fig. 6E). These data indicated that Inx2 expression is sensitive to InR/PI3K
pathway gain and loss of function. We also observed that *Inx2* mRNA level was strongly increased

in *Pten* mutant follicle cells, suggesting that the observed effects on protein level and plaque

assembly were due *Inx2* upregulation (Fig. 6F). These observations supported a model in which

the non-cell autonomous effect of the InR/PI3K pathway from somatic cells to germ cells is

mediated by gap junctions. To test this hypothesis, we performed an epistasis experiment. As

previously described (Vachias et al 2014a), large *Pten* mutant clones in the follicular epithelium

accelerated germline growth in a non-cell autonomous manner (Fig. 6G). This effect was

abrogated upon *Inx2* RNAi induction in *Pten* mutant cells (Fig. 6H). Thus, gap junctions are

required for germline growth control via InR/PI3K activity in the follicular epithelium. Altogether,

our data link the systemic control of somatic cells to the growth coordination between somatic

and germline cells via the modulation of gap junction assembly.

Discussion

In this article we show how gap junctions participate in cell growth control by allowing a metabolic flow between cells. This is illustrated by the identification of a proline transporter specifically expressed in follicle cells and required for germline cell growth. Importantly, its expression in the germline can partially bypass its silencing in the soma, providing evidence for a functional metabolic exchange between these cell types.

Inx2 and Inx4 have various functions during *Drosophila* oogenesis (Mukai et al 2011; Sahu et al 2021; Tazuke et al 2002; Miao et al 2020). However, our data showing their involvement in germ cell growth can be more easily compared with gap junction role in mammal follicles where Cx37 is expressed in the oocyte and Cx43 in follicle cells (Gittens, & Kidder 2005; Simon et al 1997). This indicates that gap junction requirement for oocyte growth is a conserved feature throughout evolution. Gap junctions are present in most animal tissues and can directly connect different cell types, for instance, neurons and glial cells. Therefore, their involvement in growth control might be more general. We found that Inx2 expression and the subsequent formation of plaques were strongly regulated by the InR/PI3K pathway in a cell autonomous manner. This indicates that gap junctions are a permissive prerequisite for germline growth, and it also provides an effective mechanism to link metabolic flow with growth systemic signals and coordinate the growth of the two connected cell populations.

Our results also established that Coch (CG43693), a putative proline transporter of the SLC36A family, must be expressed in somatic cells for germline growth. Proline import is important for cell growth in different contexts, including in cancer (Geng et al 2021; Newton et al 2020). It is usually proposed that TOR signalling links proline availability and cell growth control (Shang et al 2017; Yoshida et al 2019; Zhao et al 2019). However, this pathway is unlikely implicated in the mechanism described here. Indeed, first, when germline growth is blocked due to alteration of somatic cell growth upon loss of *akt*, which also regulates gap junction assembly, the TOR pathway is still active in the germline (Vachias et al 2014b). Moreover, overactivation of the TOR pathway in the germline in such conditions does not suppress growth inhibition. Finally, TOR inhibition in the *Drosophila* female germline does not induce P-body formation (Burn et al 2015). Therefore, we tested the involvement of the other well-described amino acid sensing pathway that relies on GCN2. Our results also excluded this mechanism to explain P-body formation and growth inhibition in the absence of proline import in somatic cells or of gap junctions. Therefore, more studies are needed to determine the precise mechanism underlying germline growth control. Nevertheless, several arguments suggest that proline import might just be the tip of the iceberg of the metabolic cooperativity between follicle cells and germ cells. *Inx2* mutant clones have a stronger impact on germline growth than *coch* mutant clones, suggesting that other metabolites from follicle cells also may promote germline growth. Accordingly, our translatome analysis showed that different genes involved in amino acid synthesis or import were specifically

expressed in follicle cells, but none in germ cells. The absence of phenotype when these genes
were knocked down, except *coch*, could be explained by multiple potential levels of redundancy
among synthesis pathways, transporters, and synthesis and import. Nonetheless, detailed
analysis of the genes overexpressed in follicle cells compared with germ cells suggest that other
amino acids (e.g. tryptophane, valine, tyrosine) could flow from the soma to the germline.
Moreover, in the mouse, the alanine transporter SLC38A3 is strongly enriched in granulosa cells
that are required for efficient alanine import in the oocyte (Eppig et al 2005). Thus, altogether these
data suggest that gap junction-dependent amino acid flow is not restricted to proline. Moreover,
glucose intake and pyruvate production are more efficient in granulosa cells, suggesting that
follicle cells could provide energetic molecules to the oocyte (Sugiura et al 2005). In line with this
observation in mammals, TRAP data indicated that transporter for trehalose (tret1-1), the
circulating sugar in insects, was expressed in somatic cells and not in the germline, whereas
germline development is highly dependent on sugar and the pentose phosphate pathway
(Carvalho-Santos et al 2020). Importantly, it has been recently shown that glucose availability and
ATP production regulate stress granules in yeast, suggesting that a link between growth control
and energy molecule disposal could also exist during germline development (Cereghetti et al
2021). Sperm development also relies on a specific metabolic programme suggesting that gap
junction-dependent metabolic cooperativity could participate in spermatogenesis (Hudry et al
2019). Thus, a contribution of energetic metabolism in the soma-germline cooperativity via gap
junctions will be an interesting avenue for future investigations.
One might ask what is the evolutive advantage of this metabolic cooperativity compared with
direct import or synthesis in the germline. Two main hypotheses could be proposed. First, female
germ cells undergo massive growth and there is a non-linear relation between volume and cell
surface increases. Consequently, surface exchange with the extracellular medium might not allow
sufficient metabolic import and may require the support of follicle cells. This hypothesis was
proposed to explain gap junction requirement for mammal oocyte growth (Su et al 2009).
However, in this case, both tissues should be able to import the required metabolites and our data
does not support such a model because follicle cells expressed a whole set of anabolic enzymes
and transporters that were not expressed in the germline. Alternatively, such a mechanism may
provide a protective effect for the germline. It is well established that the cell metabolic activity
can induce stress, with for instance, the production of reactive oxygen species (ROS). However,
the germline must be protected, especially its DNA content that is transmitted to the embryo. A
recent study demonstrated that mammalian oocytes block ROS production by suppressing
mitochondrial complex I (Rodríguez-Nuevo et al 2022). Thus, a gap junction-mediated metabolic
exchange might allow externalizing the stress-generating metabolic activity to follicle cells that
will anyway die few days later. Notably, *coch* expression in germ cells had a slight but significant
negative impact on ovary size, suggesting that its presence in germ cells is detrimental for follicle

development. Although, the exact reason for this deleterious effect is unknown, this observation
fits with a model in which externalization of metabolic import and activity could facilitate proper
oocyte development. However, Coch is a transporter, and not an enzyme, and thus, the protective
effect might not be directly due to proline exclusion, but to one of the many possible downstream
metabolic activities (Patriarca et al 2021).
Altogether, our data establish that metabolic flow through gap junctions is essential for cell growth
and that gap junction assembly can be modulated to adjust cell growth rate. Moreover, it shows
that amino acids, and potentially other metabolites, are important actors in this mechanism ending
with the formation of P-bodies, opening a large field for investigation to obtain a comprehensive
view of this metabolic cooperativity.
393

Methods

Fly genetics and handling

All fly stocks used are detailed in Table S2. The final genotypes, temperature and heat-shock conditions are in Table S3. *gcn2* null alleles were generated by inducing indel mutations using an available gRNA line. Alleles are described in Table S2. Unless specified, flies were kept on a corn meal-based medium with 80 g/L fresh yeast. Protein starvation was performed on grape juice agar plate. Coch-EGFP was obtained by inserting the EGFP-FlAsH-StrepII-TEV-3xFlag cassette in the Minos element insertion MI01960, as previously described (Venken et al 2011).

TRAP experiments

Ovaries of 100 females of each genotype (tj>RPL10-GFP or Nos>RPL10-GFP) were dissected on ice. Then, TRAP was performed as described in Bertin et al, 2015. Briefly, after homogenization, ovary extracts were preabsorbed with magnetic beads. Immunoprecipitation was performed with anti-GFP antibodies already coupled with magnetic beads before mRNA extraction. Tissue specificity and enrichment of the extracted mRNAs were checked by RT-qPCR with *GFP* (enrichment), *traffic-jam* (soma specific), *Ago3* and *Aubergine* (germline specific) primers. mRNA libraries were made with the Nugen Ovation 1-16 droso Universal RNA-seq kit according to the manufacturer's instructions. Sequencing was performed by Fasteris. Data were deposited on GEO (GSE230452) and described in Table S1.

Molecular cloning and transgenesis

The *QF* sequence was amplified from the pAttB-QF-sv40 vector and cloned in the vector that contains the alpha4-tubulin promoter (mat-tub). The pQUASp vector was constructed from pUAST in which the promoter was replaced by QUAS sites and the minimal P-element promoter was amplified from the pUASp vector. Then, this vector was used to clone *EGFP* or *coch* coding sequence and P-element insertions were generated. *eIF2α* and *eIF2α-S51D* coding sequences were cloned in the pUASz vector, and transgenes were inserted at the AttP40 landing site.

Immunostaining, FISH, EDU incorporation, imaging and quantitative analyses

Resources and reagents are listed in Table S1. Immunostaining was performed as described in Vachias et al, 2013. Stellaris SM-FISH oligo-probes against *Inx2* and *CG43693/coch* mRNAs were produced by Biosearch Technologies. FISH was performed according to the manufacturer's instruction. Images were acquired on a Zeiss LSM800 confocal microscope or a Zeiss Cell observer spinning disc microscope. P-bodies were quantified using a homemade macro initially designed to quantify basement membrane fibrils (Cerqueira Campos et al 2020). Quantifications were done on 5 μm z-stack projections of spinning-disc images acquired with a x20 lens. After manual selection of a ROI corresponding to nurse cells, P-bodies were detected by keeping objects with a minimal size of 5 adjacent pixels and a minimal fluorescence intensity of x1.75 of the mean intensity. Then, the total fluorescence contained in these P-bodies was quantified and

reported as a fraction of the total fluorescence of the ROI. For EDU incorporation, ovaries were

incubated with 10μM EDU in complemented Schneider medium for 15 minutes. After fixation,

staining was performed according to the manufacturer's instructions (Kit EDU C10638, Thermo

Fisher). Mutant-control follicle volume ratios were calculated after estimating the follicle volume

by multiplying their length by their width to the square. Mutant and control follicles in position 2

or 3 of the ovariole starting from the anterior were analysed. Cell size was automatically

determined after cell segmentation using Tissue Analyzer (Aigouy et al 2016). Statistical analyses

were performed with Prism. For all experiments, the minimum sample size is indicated in the figure

legends. For each experiment, multiple females were dissected. Randomization or blinding was

not performed. The sample normality was calculated using the D'Agostino and Pearson normality

test. The unpaired t-test was used to compare samples with normal distribution (or one-way

ANOVA for multiple comparisons), and the unpaired Mann–Whitney test for samples without

normal distribution. Figures were prepared using ScientiFig (Aigouy, & Mirouse 2013).

Acknowledgements

We are grateful to G. Junion, P Phelan, G Tanentzapf, H.D. Ryoo for sharing antibodies or fly stocks. This research was financed by the French government IDEX-ISITE initiative 16-IDEX-0001 (CAP 20-25). We also thank the CLIC facility (Clermont Imagerie Confocale).

References

Aigouy, B. & Mirouse, V., 2013, ScientiFig: a tool to build publication-ready scientific figures, *Nature methods*, 10(11), p. 1048.

Aigouy, B., Umetsu, D. & Eaton, S., 2016, Segmentation and Quantitative Analysis of Epithelial Tissues, *Methods in molecular biology (Clifton, N.J.)*, 1478, pp. 227-39.

Bauer, R., Löer, B., Ostrowski, K., Martini, J., Weimbs, A., Lechner, H. & Hoch, M., 2005, Intercellular communication: the Drosophila innexin multiprotein family of gap junction proteins, *Chemistry & biology*, 12(5).

Bohrmann, J. & Zimmermann, J., 2008, Gap junctions in the ovary of Drosophila melanogaster: localization of innexins 1, 2, 3 and 4 and evidence for intercellular communication via innexin-2 containing channels, *BMC developmental biology*, 8, p. 111.

Boll, M., Foltz, M., Rubio-Aliaga, I., Kottra, G. & Daniel, H., 2002, Functional characterization of two novel mammalian electrogenic proton-dependent amino acid cotransporters, *The Journal of biological chemistry*, 277(25).

Borreguero-Muñoz, N., Fletcher, G.C., Aguilar-Aragon, M., Elbediwy, A., Vincent-Mistiaen, Z.I. & Thompson, B.J., 2019, The Hippo pathway integrates PI3K-Akt signals with mechanical and polarity cues to control tissue growth, *PLoS biology*, 17(10), p. e3000509.

Burn, K.M., Shimada, Y., Ayers, K., Vemuganti, S., Lu, F., Hudson, A.M. & Cooley, L., 2015, Somatic insulin signaling regulates a germline starvation response in Drosophila egg chambers, *Developmental biology*, 398(2).

Cadena Sandoval, M., Heberle, A.M., Rehbein, U., Barile, C., JM, R.P. & Thedieck, K., 2021, mTORC1 Crosstalk With Stress Granules in Aging and Age-Related Diseases, *Frontiers in aging*, 2.(M

Carvalho-Santos, Z., Cardoso-Figueiredo, R., Elias, A.P., Tastekin, I., Baltazar, C. & Ribeiro, C., 2020, Cellular metabolic reprogramming controls sugar appetite in Drosophila, *Nature metabolism*, 2(9).

Cereghetti, G., Wilson-Zbinden, C., Kissling, V.M., Diether, M., Arm, A., Yoo, H., Piazza, I., Saad, S., Picotti, P., Drummond, D.A., Sauer, U., Dechant, R. & Peter, M., 2021, Reversible amyloids of pyruvate kinase couple cell metabolism and stress granule disassembly, *Nature cell biology*, 23(10).

Cerqueira Campos, F., Dennis, C., Alégot, H., Fritsch, C., Isabella, A., Pouchin, P., Bardot, O., Horne-Badovinac, S. & Mirouse, V., 2020, Oriented basement membrane fibrils provide a memory for F-actin planar polarization via the Dystrophin-Dystroglycan complex during tissue elongation, *Development*, 147(7).

Chen, D.Y., Crest, J., Streichan, S.J. & Bilder, D., 2019, Extracellular matrix stiffness cues junctional remodeling for 3D tissue elongation, *Nature communications*, 10(1).

Church, S.H., Donoughe, S., BAS, D.M. & Extavour, C.G., 2019, Insect egg size and shape evolve with ecology but not developmental rate, *Nature*, 571(7763).

Doherty, C.A., Amargant, F., Shvartsman, S.Y., Duncan, F.E. & Gavis, E.R., 2022, Bidirectional communication in oogenesis: a dynamic conversation in mice and Drosophila, *Trends in cell biology*, 32(4).

Doherty, C.A., Diegmiller, R., Kapasiawala, M., Gavis, E.R. & Shvartsman, S.Y., 2021, Coupled oscillators coordinate collective germline growth, *Developmental cell*, 56(6).

Domanitskaya, E., Anllo, L. & Schüpbach, T., 2014, Phantom, a cytochrome P450 enzyme essential for ecdysone biosynthesis, plays a critical role in the control of border cell migration in Drosophila, *Developmental biology*, 386(2).

Drummond-Barbosa, D. & Spradling, A.C., 2001, Stem cells and their progeny respond to nutritional changes during Drosophila oogenesis, *Developmental biology*, 231(1), pp. 265-78.

Elias, L.A., Wang, D.D. & Kriegstein, A.R., 2007, Gap junction adhesion is necessary for radial migration in the neocortex, *Nature*, 448(7156).

Eppig, J.J., Pendola, F.L., Wigglesworth, K. & Pendola, J.K., 2005, Mouse oocytes regulate metabolic cooperativity between granulosa cells and oocytes: amino acid transport, *Biology of reproduction*, 73(2).

Evans, W.H., 2015, Cell communication across gap junctions: a historical perspective and current developments, *Biochemical Society transactions*, 43(3).

Geng, P., Qin, W. & Xu, G., 2021, Proline metabolism in cancer, *Amino acids*, 53(12).

Gilboa, L., Forbes, A., Tazuke, S.I., Fuller, M.T. & Lehmann, R., 2003, Germ line stem cell
differentiation in Drosophila requires gap junctions and proceeds via an intermediate state,
Development, 130(26).

Gittens, J.E. & Kidder, G.M., 2005, Differential contributions of connexin37 and connexin43 to
oogenesis revealed in chimeric reaggregated mouse ovaries, *Journal of cell science*, 118(Pt 21).

Hudry, B, E, D.G., Mineo, A., Gaspar, P., Hadjieconomou, D., Studd, C., Mokochinski, J.B.,
Kramer, H.B., Plaçais, P.Y., Preat, T. & Miguel-Aliaga, I., 2019, Sex Differences in Intestinal
Carbohydrate Metabolism Promote Food Intake and Sperm Maturation, *Cell*, 178(4).

Kang, M.J., Vasudevan, D., Kang, K., Kim, K., Park, J.E., Zhang, N., Zeng, X., Neubert, T.A.,
Marr, M.T. & Ryoo, H.D., 2017, 4E-BP is a target of the GCN2-ATF4 pathway during Drosophila
development and aging, *The Journal of cell biology*, 216(1).

Kosakamoto, H., Okamoto, N., Aikawa, H., Sugiura, Y., Suematsu, M., Niwa, R., Miura, M. &
Obata, F., 2022, Sensing of the non-essential amino acid tyrosine governs the response to
protein restriction in Drosophila, *Nature metabolism*, 4(7).

LaFever, L., Feoktistov, A., Hsu, H.J. & Drummond-Barbosa, D., 2010, Specific roles of Target of
rapamycin in the control of stem cells and their progeny in the Drosophila ovary, *Development*,
137(13), pp. 2117-26.

Maines, J.Z., Stevens, L.M., Tong, X. & Stein, D., 2004, Drosophila dMyc is required for ovary
cell growth and endoreplication, *Development*, 131(4), pp. 775-86.

Metzner, L., Neubert, K. & Brandsch, M., 2006, Substrate specificity of the amino acid
transporter PAT1, *Amino acids*, 31(2).

Miao, G., Godt, D. & Montell, D.J., 2020, Integration of Migratory Cells into a New Site In Vivo
Requires Channel-Independent Functions of Innexins on Microtubules, *Developmental cell*,
54(4).

Michel, D., Arsanto, J.P., Massey-Harroche, D., Béclin, C., Wijnholds, J. & Le Bivic, A., 2005,
PATJ connects and stabilizes apical and lateral components of tight junctions in human
intestinal cells, *Journal of cell science*, 118(Pt 17), pp. 4049-57.

Mukai, M., Kato, H., Hira, S., Nakamura, K., Kita, H. & Kobayashi, S., 2011, Innexin2 gap
junctions in somatic support cells are required for cyst formation and for egg chamber formation
in Drosophila, *Mechanisms of development*, 128(7-10).

Newton, H., Wang, Y.F., Camplese, L., Mokochinski, J.B., Kramer, H.B., AEX, B., Fets, L. &
Hirabayashi, S., 2020, Systemic muscle wasting and coordinated tumour response drive
tumourigenesis, *Nature communications*, 11(1).

Oshima, A., Tani, K. & Fujiyoshi, Y., 2016, Atomic structure of the innexin-6 gap junction channel
determined by cryo-EM, *Nature communications*, 7.

Patriarca, E.J., Cermola, F., D'Aniello, C., Fico, A., Guardiola, O., D, D.C. & Minchiotti, G., 2021,
The Multifaceted Roles of Proline in Cell Behavior, *Frontiers in cell and developmental biology*, 9.

Pesch, Y.Y., Dang, V., Fairchild, M.J., Islam, F., Camp, D., Kaur, P., Smendziuk, C.M.,
Messenberg, A., Carr, R., McFarlane, C.R., Musso, P.Y., F, V.P. & Tanentzapf, G., 2022, Gap
junctions mediate discrete regulatory steps during fly spermatogenesis, *PLoS genetics*, 18(9).

Petkova, M.D., Little, S.C., Liu, F. & Gregor, T., 2014, Maternal origins of developmental
reproducibility, *Current biology : CB*, 24(11), pp. 1283-8.

Pillai, S.M. & Meredith, D., 2011, SLC36A4 (hPAT4) is a high affinity amino acid transporter when
expressed in Xenopus laevis oocytes, *The Journal of biological chemistry*, 286(4).

Potter, C.J., Tasic, B., Russler, E.V., Liang, L. & Luo, L., 2010, The Q system: a repressible
binary system for transgene expression, lineage tracing, and mosaic analysis, *Cell*, 141(3).

Riggs, C.L., Kedersha, N., Ivanov, P. & Anderson, P., 2020, Mammalian stress granules and P
bodies at a glance, *Journal of cell science*, 133(16).

Rodríguez-Nuevo, A., Torres-Sanchez, A., Duran, J.M., De Guirior, C., Martínez-Zamora, M.A. &
Böke, E., 2022, Oocytes maintain ROS-free mitochondrial metabolism by suppressing complex
I, *Nature*, 607(7920), pp. 756-61.

Rørth, P., 1998, Gal4 in the Drosophila female germline, *Mechanisms of development*, 78(1-2).

Sahu, A., Karmakar, S., Halder, S., Ghosh, G., Acharjee, S., Dasgupta, P., Ghosh, R.,
Deshpande, G. & Prasad, M., 2021, Germline soma communication mediated by gap junction
proteins regulates epithelial morphogenesis, *PLoS genetics*, 17(8).
Shang, P., Valapala, M., Grebe, R., Hose, S., Ghosh, S., Bhutto, I.A., Handa, J.T., Lutty, G.A., Lu,
L., Wan, J., Qian, J., Sergeev, Y., Puertollano, R., Zigler, J.S., Xu, G.T. & Sinha, D., 2017, The
amino acid transporter SLC36A4 regulates the amino acid pool in retinal pigmented epithelial
cells and mediates the mechanistic target of rapamycin, complex 1 signaling, *Aging cell*, 16(2).
Shimada, Y., Burn, K.M., Niwa, R. & Cooley, L., 2011, Reversible response of protein localization
and microtubule organization to nutrient stress during Drosophila early oogenesis,
Developmental biology, 355(2).
Simon, A.M., Goodenough, D.A., Li, E. & Paul, D.L., 1997, Female infertility in mice lacking
connexin 37, *Nature*, 385(6616).
Skerrett, I.M. & Williams, J.B., 2017, A structural and functional comparison of gap junction
channels composed of connexins and innexins, *Developmental neurobiology*, 77(5).
Smendziuk, C.M., Messenberg, A., Vogl, A.W. & Tanentzapf, G., 2015, Bi-directional gap
junction-mediated soma-germline communication is essential for spermatogenesis,
Development, 142(15).
Spradling, A.C., Bate, M. & Arias, A.M., 1993, The development of Drosophila melanogaster, *The
Development of Drosophila Melanogaster*, 1, pp. 1-70.
Srivastava, A., Lu, J., Gadalla, D.S., Hendrich, O., Grönke, S. & Partridge, L., 2022, The Role of
GCN2 Kinase in Mediating the Effects of Amino Acids on Longevity and Feeding Behaviour in
Drosophila, *Frontiers in aging*, 3.
Su, Y.Q., Sugiura, K. & Eppig, J.J., 2009, Mouse oocyte control of granulosa cell development
and function: paracrine regulation of cumulus cell metabolism, *Seminars in reproductive
medicine*, 27(1), pp. 32-42.
Sugiura, K., Pendola, F.L. & Eppig, J.J., 2005, Oocyte control of metabolic cooperativity
between oocytes and companion granulosa cells: energy metabolism, *Developmental biology*,
279(1).
Sun, P., Quan, Z., Zhang, B., Wu, T. & Xi, R., 2010, TSC1/2 tumour suppressor complex
maintains Drosophila germline stem cells by preventing differentiation, *Developmen)*, 137(15),
pp. 2461-9.
Tazuke, S.I., Schulz, C., Gilboa, L., Fogarty, M., Mahowald, A.P., Guichet, A., Ephrussi, A.,
Wood, C.G., Lehmann, R. & Fuller, M.T., 2002, A germline-specific gap junction protein required
for survival of differentiating early germ cells, *Development*, 129(10).
Vachias, C., Fritsch, C., Pouchin, P., Bardot, O. & Mirouse, V., 2014a, Tight coordination of
growth and differentiation between germline and soma provides robustness for drosophila egg
development, *Cell reports*, 9(2), pp. 531-41.
Vachias, C., Fritsch, C., Pouchin, P., Bardot, O. & Mirouse, V., 2014b, Tight coordination of
growth and differentiation between germline and soma provides robustness for drosophila egg
development, *Cell reports*, 9(2), pp. 531-41.
Venken, K.J., Schulze, K.L., Haelterman, N.A., Pan, H., He, Y., Evans-Holm, M., Carlson, J.W.,
Levis, R.W., Spradling, A.C., Hoskins, R.A. & Bellen, H.J., 2011, MiMIC: a highly versatile
transposon insertion resource for engineering Drosophila melanogaster genes, *Nature methods*,
8(9), pp. 737-43.
Wang, Y. & Riechmann, V., 2007, The role of the actomyosin cytoskeleton in coordination of
tissue growth during Drosophila oogenesis, *Current biology : CB*, 17(15), pp. 1349-55.
Yoshida, A., Bu, Y., Qie, S., Wrangle, J., Camp, E.R., Hazard, E.S., Hardiman, G., R, D.L.,
Knudsen, K.E. & Diehl, J.A., 2019, SLC36A1-mTORC1 signaling drives acquired resistance to
CDK4/6 inhibitors, *Science advances*, 5(9).
Zhao, L., Zhang, X., Ji, X., Jin, Y. & Liu, W., 2019, The amino acid transporter PAT1 regulates
mTORC1 in a nutrient-sensitive manner that requires its transport activity, *Cellular signalling*, 53.
611
612

Figure legends

Figure 1: Soma-germline gap junctions are required for germline growth

A) Scheme of an ovariole, oriented from the anterior (A) to the posterior (P) end, as all the subsequent images. The ovariole starts with the germarium from which young follicles bud before undergoing massive growth until the formation of an egg at the posterior end. In each follicle, somatic follicle cells (in purple) surround a germline cyst with the nurse cells (pink) and the oocyte (in blue). B) Sagittal view and C) Maximum intensity projection images of a stage 7 follicle after immunostaining for Inx2 (green in B and C, white in B' and C') and Inx4 (magenta in B and C, white in B'' and C''). D) and E) Higher magnification of the insets in B and C to illustrate the formation of plaques at the germline-soma interface. F) Immunostaining for Inx2 (green in F and F', white in F''') and Inx4 (magenta in F', white in F'') in a follicle containing a *Inx2* RNAi-expressing clone in follicle cells marked by GFP expression (white in F). G) Immunostaining for Inx2 (green in G, white in G'' and G''') and Inx4 (magenta in G, white in G') in follicles containing *Inx4* RNAi-expressing germline clones visualized by Inx4 absence. Note the absence of Inx2 plaques at germline contacts at higher magnification (G''') and the growth defect of *Inx4* RNAi follicles compared with the wild-type younger follicle (G). H) Somatic *Inx2^A* mutant clones, marked by the absence of RFP expression, induce a germline growth defect when they cover the whole epithelium. I) Quantification of the ratio between the volume of the follicle showing a growth defect (fully covered by *Inx2* mutant cells) and the volume of the n-1 follicle of the same ovariole (n= 10 control and 7 *Inx2* mutant follicles). J-K) quantification of J) cell surface incorporation and K) EDU-positive cells in small mutant *Inx2* clones compared with the surrounding wild-type cells (n= 20, 20 clones or groups of wild-type cells for J and K). For all graphs, data are the mean ± SD. ***p <0.01. Scale bars: 20 μm.

Figure 2: TRAP analysis identifies soma-specific expression of amino acid metabolism and transport genes

A) Representative images of follicles expressing RPL10a-GFP in the soma (driven by tj:Gal4) or the germline (driven by nos:Gal4p16). After polysome immunoprecipitation (IP), mRNAs were sequenced. B) The expression of 811 genes was significantly enriched in somatic cells compared with germline cells (fold change >5, p-value <0.05). After filtering for genes involved in small molecule metabolism or transport, genes related to amino acid biology were identified. C) Distribution (soma and/or germline) of genes encoding proteins implicated in amino acid biosynthesis or transport according to their expression profile. No gene was enriched exclusively in the germline.

Figure 3: Proline import in follicle cells promotes germline growth

A-B) Representative images of ovaries in A) control and B) after *CG43693* knock-down by RNAi in somatic cells. C) Ovary size (surface) quantification in control and after *CG43693* somatic knock-down using two RNAi lines. D-E) In situ hybridization analysis of *CG43693* expression in D) a whole ovariole and in E) stage 4 and 7 follicles. F) Expression of endogenous CG43693 protein tagged with GFP from germarium to stage 8. G) Higher magnification of follicle cells showing CG43693-GFP localization in the cell cortex. H) *CG43693/cochonnet (coch)* mutant clones marked by the absence of RFP. When all follicle cells are mutated, follicle growth is affected. I) Quantification of the ratio between the volume of follicle with a growth defect and fully covered by mutant cells and the n-1 follicle in the same ovariole (n= 10 control and 6 *coch* mutant follicles). J-K) quantification of J) EDU-positive cells and K) cell surface in small *coch* mutant clones compared with the surrounding wild-type cells (n=12 control and 12 mutant clones or groups of wild-type cells). For all graphs, data are the mean ± SD. **p <0.01, ***p <0.001. Scale bars: 50 μm in D,E,F,H and 20 μm in G.

Figure 4: Ectopic proline import in the germline compensates its absence in the soma

A) scheme representing *coch* expression in green in the different genotypes used on this figure. B-E) Representative images of a full ovary from B) a control female, C) a *MatTub> coch* female D) a *tj > coch RNAi* female, and E) a *tj > coch RNAi, MatTub> coch* female. C) Quantification of ovary size (surface) in the indicated genotypes (n = 14, 10, 19, and 27 ovaries). For all graphs, data are the mean ± SD., ***p <0.001, ****p <0.0001. Scale bars: 50 μm.

Figure 5: Defective gap junctions or proline import induces P-body formation

A-B) Endogenous GFP-tagged Me31B protein expression in stage 9 follicles in A) well fed conditions or B) after 15 hours of protein starvation. C) P-body quantification in normal and starved conditions. D-E) Me31B-GFP expression in ovarioles germline that overexpresses D) wild-type eIF2α or E) or eIF2α-S51D (inhibitory phosphorylation). F and H) Representative images of Me31B-GFP expression in stage 9 follicles after RNAi-based knock-down in somatic cells of F) *Inx2* or H) *coch*. G and I) P-body quantification in follicles after RNAi-based knock-down in somatic cells of G) *Inx2* and I) *coch.* For all graphs, data are the mean ± SD, ****p <0.0001. Scale bars: 50 μm.

Figure 6: Gap junction are controlled by the InR/PI3K pathway

A,B) Representative images of Me31B-BFP expression in stage 9 follicles from A) a control female and B) a *tj:Gal4, Tub:Gal80^{ts} > akt RNAi* female. C) P-body quantification (fluorescence intensity) in the indicated genotypes (n = 8 and 9 follicles). Data are the mean ± SD. ****p <0.0001. D) Inx2 expression in an *akt^q* mutant clone (marked by the absence of GFP expression). E-F) Inx2 protein (E) and *Inx2* mRNA expression by FISH (F) in a *Pten^{dj189}* mutant clone marked by the absence of RFP expression. G) Large MARCM *Pten^{dj189}* mutant clone marked by the absence of GFP

687   expression showing faster follicle growth. H) Large MARCM $Pten^{dj189}$ mutant clone in which faster

688   follicle growth was abolished after expression of an RNAi against *Inx2*. Scale bars: 50 μm in A, B,

689   G, H and 20 μm in D,E,F.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplemental data
693

694 Supplemental data include :
695   - 4 supplemental figures
696   - Table S1 : Excel file of soma vs germline TRAP analysis
697   - Table S2 : Reagents and resources
698   - Table S3 : Genotypes and specific conditions
699

700 **Figure S1:**
701 Maximum intensity projection images of an ovariole after immunostaining for Inx2 (green) and
702 Inx4 (red). Note the progressive increase of Inx2 expression.
703

704 **Figure S2:**
705 Ovariole expressing a QUASP:GFP transgene under the control of the matTub:QF driver.
706

Figure S3:
A,B) Representative images of Me31B-BFP expression in stage 9 follicles from a: A) control
female, B) starved control female, C) *gcn2* transheterozygous mutant female, and D protein
starved *gcn2* transheterozygous mutant female. E) P-body quantification (fluorescence intensity)
in follicles of the indicated genotypes and conditions. Data are the mean ± SD. F,G) Absence of
Atf4-GFP protein expression used a read-out of GCN2 activity in F) normal and G) protein-starved
conditions. H-I) Ovarioles with germline *gcn2* mutant clones marked by the absence of RFP
expression (magenta) and stained for F-actin (green) in H) wild-type background and I) in flies
harbouring *coch* RNAi in follicle cells.
716

717 **Figure S4:**
718 A) Coch-GFP expressing follicle with *akt* mutant clones (RFP-negative cells). B) Coch-GFP
719 expressing follicle with flip-out clones that express a constitutively active form of PI3K (RFP-
720 positive cells).
721

Figure S1

matTub:QF>QUASP:GFP

Figure S2

Figure S3

Figure S4

Table S2: Resources and reagents

RESOURCE OR REAGENT	SOURCE	IDENTIFIER
Experimental models: Organisms/Strains		
tj:Gal4 : P{w+mW.hs=GawB}NP1624	Kyoto Stock Center (DGRC)	FlyBase ID: FBti0034540; DGRC: 104055
FRT19A, Inx2^A	Bloomington Drosophila Stock Center	FlyBase ID: FBal0291304; BDSC: 54481
FRT19A, Inx2^B	Bloomington Drosophila Stock Center	FlyBase ID: FBal0291305; BDSC: 52363
FRT19A, Inx2^{G0173a}	Bloomington Drosophila Stock Center	FlyBase ID: FBal0151828; DGRC: 111858
UAS :Inx2 RNAi (JF02446)	Bloomington Drosophila Stock Center	FlyBase ID: FBti0128670; BDSC: 29306
UAS : Inx4 RNAi (GL00447)	Bloomington Drosophila Stock Center	FlyBase ID: FBtp0068601; BDSC: 35607
nos:Gal4Vp16	Bloomington Drosophila Stock Center	FlyBase ID: FBti0012410; BDSC: 4937
Mat-tub :Gal4Vp16 (V2H)	Bloomington Drosophila Stock Center	FlyBase ID: FBti0016915; BDSC: 7062
UASp-RPL10a-GFP	Junion G.	Nd
UAS : CG43693 RNAi (GLV21061)	Bloomington Drosophila Stock Center	FlyBase ID: FBti0144657; BDSC: 35696
UAS : CG43693 RNAi (GL01304)	Bloomington Drosophila Stock Center	FlyBase ID: FBti0149385; BDSC: 41873
CG43693/Coch ^{MI01960} : Mi (Michel et al 2005)CG43693^{MI01960}	Bloomington Drosophila Stock Center	FlyBase ID: FBti0132690; BDSC: 34264

Coch/CG43693-EGFP : CG43693 [MI01960-GFSTF.0]	This study	
Me31B-GFP : P{PTT-GB}me31B[CB05282]	Bloomington Drosophila Stock Center	FlyBase ID: FBal0211797 BDSC: 51530
UASz : eIF2a	This study	
UASz : eIF2a-S51A	This study	
UASz : eIF2a-S51D	This study	
Mat-tub :QF	This study	
QUASp :EGFP	This study	
QUASp :Coch	This study	
UAS :gcn2 sgRNA	Vienna Drosophila Resource Center	VDRC: 341108
nos:Gal4Vp16, ; UAS:Cas9	Bloomington Drosophila Stock Center	BDSC: 54593
FRT gnc2[CB20]	This study, contains a 2n postion 878 (gg) on gcn2-RA transcript	
FRT gcn2[CB14]	This study, contains a 1n deletion at position 877 (a) and a 9n insertion (tttccgacc) on gcn2-RA transcript	
4E-BPintron-dsRed	Kang et al, J Cell Biol, 2017	FlyBase ID: FBtp0116205
ATF4-GFP : Mi{PT-GFSTF.1}crcMI02300-GFSTF.1	Bloomington Drosophila Stock Center	FlyBase ID: FBal0314614; BDSC: 59608
FRT40A, Pten[dj189]	Pan D.	FlyBase ID: FBal0117670
FRT82B, akt[q]	S. Cohen	Nd
UAS : akt RNAi (HM04007)	Bloomington Drosophila Stock Center	FlyBase ID: FBti0130466 BDSC 31701
UAS : PI3K-CA (Chromosome II)	This study, obtained by P-element mobilization of UAS:PI3K92E-CAAX (FBal0156077)	
Antibodies		

anti-Inx4, rabbit	G. Tanentzapf	1/2000
anti-Inx2, Guinea Pig	G. Tanentzapf	1/400
anti-Inx2 rabbit	P. Phelan	1/5000
anti-coracle mouse	DSHB	C615.16 1/100
anti-mouse –Alexa 488	Life technologies	A21202 1/1000
anti-rabbit -Cy5, Donkey	Jackson Immunoresearch	711-175-152 dilution 1/1000
Anti-rabbit pig-Cy3, Donkey	Jackson Immunoresearch	711-165-152 dilution 1/1000
anti-guinea pig-Cy5, Donkey	Jackson Immunoresearch	703-175-155; dilution 1/1000
Anti-guinea pig-Cy3, Donkey	Jackson Immunoresearch	706-165-148 dilution 1/1000
Chemicals, Peptides, kits		
Phalloidin Atto-488, -550, -633	Sigma	#49409, #19083, #68825
DAPI	Sigma	100 µg/ml D8417-1MG
EdU Click-iT™ Plus Alexa 555	Thermo Fisher	C10638
Ovation 1-16 droso Universal RNA-seq kit	Nugen	No longer distributed
Software and Algorithms		
Fiji	(Schindelin et al, 2012)	
Prism		

727 **Table S3: Genotypes and specific conditions**

728 (h: hours; HS: heat-shock, GP : grape juice agar plate)

Figure		Genotype	Condition
Figure 1	BE	*w¹¹¹⁸*	25°C
	F	*y,w,hs:flp122; Act5C:FRT-Stop-FRT-Gal4, UAS:GFP , UAS:Inx2 RNAi*	1HS, 4 days at 30°C
	G H,I	*y,w,hs:flp122; Act5C:FRT-Stop-FRT-Gal4, UAS:GFP , UAS:Inx4 RNAi*	1HS, 4 days at 30°C
		FRT19A, Inx2 / FRT19A, Ubi:nlsRFP, hs:flp122	1HS, 10-12 days at 25°C
	J,K	*FRT19A, Inx2 / FRT19A, Ubi:nlsRFP, hs:flp122*	1HS, 2 days at 25°C
Figure 2	A	*tj:Gal4 / UASp :Rpl10a-GFP* *and* *UASp :Rpl10a-GFP; nos:Gal4VP16*	25°C
Figure 3	A	*tj:Gal4*	4 days at 30°C
	B	*tj:gal4 / UAS: CG43693 RNAi ᴳᴸ⁰¹³⁰⁴*	4 days at 30°C
	C	*tj:Gal4*	4 days at 30°C
		tj:gal4 / UAS: CG43693 RNAiᴳᴸⱽ²¹⁰¹⁶¹	
		tj:gal4 / UAS: CG43693 RNAiᴳᴸ⁰¹³⁰⁴	
	D,E	*w¹¹¹⁸*	25°C
	F,G	*CG43693-GFP (or coch-GFP)*	25°C
	H,I	*y,w,hs:flp122; FRT2A, cochᴹᴵ⁰¹⁹⁶⁰/ FRT2A, Ubi:nlsRFP*	2HS, 8 days at 25°C
	J,K	*y,w,hs:flp122; FRT2A, cochᴹᴵ⁰¹⁹⁶⁰/ FRT2A, Ubi:nlsRFP*	1HS, 2 days at 25°C
Figure 4	A,B,F	*tj:gal4, matTub:QF, / +* *tj:gal4, matTub:QF,; QUASP:coch*	4 days at 30°C
	A,C,F	*tj:gal4, matTub:QF,; UAS: coch RNAiᴳᴸⱽ²¹⁰⁶¹*	
	A,D,F A,E,F	*tj:gal4, matTub:QF, QUASP:coch; UAS: coch RNAiᴳᴸⱽ²¹⁰⁶¹*	
Figure 5		*Me31B-GFP*	25°C
	A,C	*Me31B-GFP*	25°C, 14h on GP
	C		25°C
	D	*matTub :Gal4Vp16, Me31B-GFP , UASz: eIF2α*	25°C
	E,F	*matTub :Gal4Vp16, Me31B-GFP, UASz: eIF2α-S51D*	18°C, 14h at 30°C
	G,H	*tj:gal4,Tub:Gal80ᵗˢ,Me31B-GFP / UAS:Inx2RNAi*	2 days at 25 °C
		tj:gal4, Me31B-GFP / UAS: coch RNAiᴳᴸⱽ²¹⁰⁶¹	
Figure 6	A,C	*tj:Gal4, Tub:Gal80ᵗˢ Me31B-GFP*	2 days at 30°C
	B,C	*tj:Gal4, Tub:Gal80ᵗˢ, Me31B-GFP / UAS: akt RNAi*	2 days at 30°C
	D	*y,w,hs:flp122; FRT82B, aktᑫ / FRT82B, Ubi:GFP*	1HS, 4 days at 25°C

E,F G H	*y,w,hs:flp122 ; FRT40A,Pten^{dj189} / FRT40A, Ubi:nlsRFP* *y,w,hs:flp122, Tub:Gal4, UAS :GFP ; FRT40A,Pten^{dj189} /* *FRT40A, Tub :Gal80* *y,w,hs:flp122, Tub:Gal4, UAS :GFP ; FRT40A,Pten^{dj189} /* *FRT40A, Tub :Gal80; UAS :Inx2 RNAi*	1HS, 4 days at 25°C 1HS, 5 days at 25°C 1HS, 6 days at 25°C 1HS, 6 days at 25°C
Figure S1	*w¹¹¹⁸*	25°C
Figure S2	*MatTub:QF / QUASP:GFP*	25°C
Figure S3 A,C B,C D,E H I	*Me31B-GFP* *Me31B-GFP, gcn2^{CB20}/ gcn2^{CB14}* *ATF4-GFP* *y,w,hs:flp122 ; FRT82B, gcn2^{CB14} / FRT82B, Ubi:nlsRFP* *y,w,hs:flp122 ; tj :Gal4 ; UAS: coch RNAi^{GL01304}, FRT82B,* *gcn2^{CB14} / FRT82B RFP*	25°C 25°C 25°C 1HS, 5 days at 30°C 1HS, 5 days at 30°C
Figure S4 A B	*y,w,hs:flp122 ; coch-GFP, FRT82B, akt^q / FRT82B,* *Ubi:nlsRFP* *y,w, hsflp122; Act5C:FRT-STOP-FRTGal4; UAS:RFP/* *UAS:PI3K-CAAX ; coch-GFP*	1HS, 5 days at 25°C 1HS, 5 days at 25°C