The contribution of an X chromosome QTL to non-Mendelian inheritance and

unequal chromosomal segregation in *A. freiburgense*

Authors: Talal Al-Yazeedi[1,3], Sally Adams[1], Sophie Tandonnet[1], Anisa Turner[1], Jun

Kim[2,4], Junho Lee[2], and Andre Pires-daSilva[1*]

Affiliations:

[1] School of Life Sciences, University of Warwick, Coventry, CV4 7AL, UK

[2] Institute of Molecular Biology and Genetics, Seoul National University, Seoul,

08826, South Korea

Current address:

[3] Liverpool School of Tropical Medicine, Pembroke Place Liverpool, L3 5QA, UK

[4] Department of Convergent Bioscience and Informatics, Chungnam National

University, Daejeon, 34134, South Korea

*Corresponding author (andre.pires@warwick.ac.uk)

Abstract

Auanema freiburgense is a trioecious nematode with co-existing males, females, and selfing hermaphrodites. Crosses of XO males with XX females result in a low percentage of XO sons due to the elimination of the nullo-X spermatids by the fathers. This process yields most viable sperm containing an X chromosome, leading to a higher transmission probability of the X chromosome compared to expected transmission via random segregation. The mechanism underlying this process involves the asymmetric distribution of essential cellular organelles during sperm formation, which likely depends on the X chromosome. Specifically, sperm components segregate with the X chromosome daughter cell, while other components are discarded in the nullo-X daughter cell. Here we found that intercrossing two strains of *A. freiburgense* results in lines in which males produce viable nullo-X sperm. Thus, crosses of those hybrid males with females result in a high percentage of sons. To uncover the genetic basis of nullo-spermatid elimination and X-chromosome drive, we generated a genome assembly for *A. freiburgense* and genotyped the intercrossed lines. We identified a QTL encompassing several genes on the X chromosome that are associated with its non-Mendelian inheritance observed in *A. freiburgense*. This finding provides valuable clues to the underlying factors involved in asymmetric organelle partitioning during male meiotic division and thus non-Mendelian transmission of the X chromosome and sex ratios.

Introduction

Animal reproduction involves the production of gametes, with distinct characteristics
and patterns of inheritance. Male and female gametes differ not only in size but also
in the transmission of cytoplasmic elements. Typically, cytoplasmic components
such as mitochondria and bacterial endosymbionts are maternally inherited (but see
[1-4]). In contrast, the nuclear genome is usually inherited equally from both parents,
ensuring a balanced contribution of genetic material. However, exceptions to this
symmetrical nuclear inheritance exist across various species, ranging from maternal-
only to paternal-only inheritance of the nuclear genome (for review, see [5]). These
variations in inheritance patterns for cytoplasmic elements and the nuclear genome
highlight the dynamic nature of reproductive strategies in animals. Understanding
these exceptions contributes to our broader understanding of reproductive biology
and the diverse mechanisms employed by organisms to propagate their genetic
material.

One such variation is the asymmetric transmission of sex chromosomes. In XX:XY
sex determination systems, for instance, half of the gametes produced by the male
have one X chromosome and the other half have one Y chromosome. The
transmission is asymmetric because only the male can transmit the Y chromosome
to the next generation and females transmit twice the number of X chromosomes.
Although the expected XX:XY ratio in the offspring for this type of sex determination
is 1:1, selfish genetic elements in one of the sex chromosomes may drive a non-
Mendelian transmission, resulting in a bias towards female or male offspring [6].

Similarly, XX:XO sex-determining systems may also result in non-Mendelian

transmission of the X chromosome, resulting in sex ratio bias after crosses. In this

type of sex-determining system, the male is heterogametic and therefore is expected

to produce an equal number of X-bearing and nullo-X gametes. In some organisms,

however, the nullo-X sperm is not produced by males, resulting in gametes that only

carry an X chromosome [7-9]. Thus, the following generation is biased towards XX

animals.

Nematodes are known to have diverse sex determination systems [10], which makes

them well-suited for studying sex ratios due to their short life cycles, prolific

reproduction, and ease of husbandry. In some nematode clades, crosses between

XX and XO individuals result in highly biased XX offspring (for review, see [11]). In only

a few cases the mechanisms underlying this sex ratio bias are known. In nematodes

that include the genus *Auanema* and *Strongyloides*, for instance, the male-producing

nullo-X spermatid is obligatorily eliminated during spermatogenesis [7,12-14].

Nematodes of the genus *Auanema* are trioecious, consisting of XX hermaphrodites,

XX females, and XO males. In hermaphrodites and males, an asymmetric division

occurs during spermatogenesis, resulting in organelles partitioning to different sides

of the dividing cell (Figure 1A) [12,13,15]. During spermatogenesis in *A. freiburgense*,

components essential for sperm function, such as mitochondria and organelles

containing cytoskeletal proteins necessary for sperm motility, co-segregate with the

X chromosome [16]. Meanwhile, the sister cell lacking an X chromosome and

consisting only of autosomes undergoes differentiation into a residual body that is

subsequently eliminated together with organelles such as Golgi complex and

endoplasmic reticulum [12,13,15]. Consequently, crosses between males and XX

females result in the production of mostly XX offspring [7,12,17].

Based on the timing of the X chromosome co-segregation relative to the organelles

and on the symmetric segregation of organelles in masculinized XX mutants, it has

been suggested that the X chromosome may serve as a polarizing signal for this

asymmetric cell division [13,15]. In this study, we report the use of Recombinant Inbred

Advanced Intercross Lines (RIAILs), whole genome sequencing, and Quantitative

Trait Locus (QTL) mapping to identify the genetic components controlling

asymmetric segregation of organelles in male meiosis in *Auanema freiburgense* [13,18].

Some of the RIALs displayed a high rate of male production after outcrossing males.

We show that males in these lines produce viable nullo-X sperm, and that this new

phenotype maps to a region in the X chromosome. These findings are consistent

with the hypothesis that loci in the X chromosome are acting as a polarizing signal

during sperm formation, although loci in other chromosomes may also be involved.

Figure 1. Models of meiosis in *Auanema*. (**A**) In wild-type male spermatogenesis,

the chromatids of the X chromosome separate in meiosis I. In anaphase II, the cell

that receives the X chromosome also inherits mitochondria and the fibroid bodies

containing the Major Sperm Protein (MSP). The cell without the X chromosome

(nullo-X cell) inherits non-sperm materials and is discarded as a polar body (not to

scale). (**B**) *Auanema* females have canonical meiosis, where the X chromosomes

undergo recombination and there is the production of an X-bearing oocyte and three

polar bodies. Therefore, most offspring from crosses between females and males

are XX. However, if non-disjunction occurs during female oogenesis, it can result in a

nullo-X oocyte and male offspring (not pictured in the diagram). (**C**) During

hermaphrodite spermatogenesis, diplo-X sperm is produced: the two homologous X

chromosomes are segregated into one daughter cell during anaphase II. (**D**) In

hermaphrodite oogenesis, the resulting oocyte does not harbor an X chromosome.

Results

Auanema freiburgense RIAILs exhibit a transgressive phenotype

A. freiburgense is a species with three sexual morphs (XX females, XX

hermaphrodites, and XO males) [18]. Previous research has shown that when an *A.*

freiburgense female crosses with a male, the resulting offspring has a skewed sex

ratio against males [13,18]. An XX sex ratio bias occurs because the X-bearing sperm

of the male is viable, whereas the nullo-X spermatid is not [13].

Here we tested two *A. freiburgense* inbred strains, APS7 and APS14 [19], which

produced 18% or fewer sons after outcrossing (Table 1) [20].The high ratio of XX

offspring confirms previous studies with other strains of the same species [13,18]. To

investigate sex determination in *A. freiburgense*, we created a genetic linkage map

using 100 Recombinant Inbred Advanced Intercross Lines (RIAILs) (Figure 2).

Approximately 60% of these recombinant lines exhibited a transgressive phenotype,

indicating the presence of traits not observed in either parental strain. When males

from these hybrid lines crossed with APS7 females, they produced a higher

percentage of sons (High-male RIAILs, or HM-RIAILs) than the parental strains

(Figure 3) [20].

143
Figure 2. Construction of *A. freiburgense* Recombinant Inbred Advanced

Intercross Lines (RIAILs). The breeding scheme involved seven generations of

crosses and ten generations of bottlenecking by the propagation of individuals

derived from the self-fertilization of a single hermaphrodite. Cells illustrate somatic

cells at different stages of RIAILs construction, including mitochondrial and

chromosomal background. RIAILs were created by crossing an APS14 female with

an APS7 male. F1s resulting from the cross were left to mate in a large mating pool.

F1 hermaphrodites and females mated with male siblings were isolated to 100

individual plates to produce F2s, each with unique recombination patterns. From the

F2 to the F7 generation, lines were intercrossed in an inbreeding avoidance scheme.

In this scheme, each line was crossed with another line only once to maximize

haplotype breakpoints. Then, each line was selfed by single worm descent for ten

generations to increase genome homozygosity. By the F17 generation, each line had

a unique mixed genome from both parental strains and was homozygous at most

loci.

Figure 3. Percentage of male offspring from RIAILs crosses with wildtype

APS7 female. In an outcross with a wild-type female, males from most RIAILs

produce more sons than the APS7 and APS14 strains. * Indicates that crossing data

is not available.

HM lines produce more viable nullo-X sperm than the parental and LM lines

To monitor the inheritance of the X chromosome after a cross, we used a

polymorphic marker on the X chromosome. We found that females and males from

APS7 and APS14 produce rare viable nullo-X oocytes and rare viable nullo-X sperm,

respectively (Table 1). For instance, in reciprocal crosses between APS7 and

APS14, sometimes the sons inherited the X from the mother (indicating fertilization

between an X-bearing oocyte and nullo-X sperm), and sometimes from the father

(fertilization between a nullo-X oocyte and X-bearing sperm) (Table 1). Overall, the

percentage of viable nullo-X sperm produced by APS7 and APS14 males is relatively

low (<20%).

Given that the high-male lines described above produce a large percentage of sons,

we hypothesized that males in those strains were producing more nullo-X sperm

than the parental lines. To explore this further, we focused on two high-male (HM)

lines and two low-male (LM) lines. Males from these lines carried the APS14 X

genotyping marker and were crossed with an APS7 mother to enable tracking of the

X chromosome. Males from the HM lines (lines 12 and 19) produced around 50% of

sons, whereas males from the LM lines (lines 2 and 38) generated 5% or less of

sons after outcrossing (Table 1). Sons from the LM males inherited the paternal X

chromosome in all samples tested, indicating that these lines produce viable X-

bearing sperm and few or no nullo-X sperm. In contrast, almost all the sons from the

HM males inherited the maternal X chromosome, indicating that ~50% of the sperm

in these lines is composed of viable nullo-X sperm.

Table 1. The percentage of male XO progeny and nullo-X sperms from RIAILs,

intra- and inter-strain crosses in *A. freiburgense.* The father's genotype was

determined using the X-linked marker X_{634} (see Material and Methods) [20].

Mother	Father	Father's genotype	% of sons	Maternal X in son (N)	Paternal X in son (N)	% nullo-X sperm
APS7	APS7	N/A	18	N/A	N/A	N/A
APS14	APS14	N/A	8	N/A	N/A	N/A
APS14	APS7	APS7	19	31	4	17
APS7	APS14	APS14	13	16	15	7
APS7	Line 12	APS14	54	29	0	54
APS7	Line 19	APS14	50	25	4	43
APS7	Line 2	APS14	2	0	19	0
APS7	Line 38	APS14	5	0	25	0

X-chromosome independent high-density genetic linkage map improved X

chromosome assembly

To map the locus involved in the production of sons after crossing, the genome of *A.*

freiburgense (strain APS7) was sequenced using long reads of Pacific Biosciences

(PacBio) and short Illumina pair-end and mate-pair data. The initial genome

assembly with this sequencing data resulted in 75 scaffolds (Afr-genome-v1). To

improve the genome assembly, the scaffolds were anchored and ordered onto a

genetic linkage map. To create this map, we used sequencing data from 100 RIAILs,

the parental lines APS7 and APS14, and using Afr-genome-v1 as a reference.

In the related species *A. rhodense,* the X chromosomes do not recombine in

hermaphrodites [7]. Therefore, we expected that the bottlenecking of the RIAILs by

selfing single hermaphrodites would still result in an X chromosome containing

significant heterozygosity. To accommodate this unique biology, we employed three

different approaches to building the genetic linkage map, with parameter settings as

follows (see Materials and Methods): (1) using only homozygous markers for all

chromosomes (Figure S1A and S2). The resulting map consisted of seven linkage

groups, with the X linkage group containing fewer markers than the autosomes

(Table S1, Table 2); (2) using both homozygous and heterozygous markers for all

chromosomes. However, the markers clustered in several groups (more than seven)

(data not shown); (3) a map was constructed independently for the autosomes and

another one for the X chromosome (Table 2, Figure S1B and Figure S2). For this

latest approach, we used only homozygous markers to group the autosomal markers

and both homozygous and heterozygous markers to group the X chromosome

markers. Using this approach, the final length of the X chromosome doubled

compared to the first approach (Table 2 and Table S1).

When the coding genes from this integrated map-scaffold genome were compared to

those of *C. elegans*, a high degree of synteny was observed between L.7 and the *C.*

elegans X chromosome (Figure S5). The final genetic linkage map consists of seven

linkage groups with 16,792 markers from 29 scaffolds, accounting for 97% of the

sequenced genome (Figure 5, Figure S2) [20]. The overall percentage of missing

markers in the GLM is 6% (Figure 5 and Figure S3). Around 3% of sequenced DNA

(1,689,119 bp in 47 scaffolds) were left unplaced, as they did not cluster with any

linkage group (Table S2). 15 kb of those unplaced sequences correspond to the

mitochondrial genome.

230
Figure 5. Parental background of the RIAILs, genotype distribution, and

recombination across the genetic linkage map. Genotypes clustered into seven

linkage groups, with each group appearing in a separate panel on the x-axis. (**A**)

Each row in the y-axis represents the genotype of a RIAIL, color-coded according to

its genetic background. The x-axis shows the genetic distance in Mb along each

linkage group. (**B**) Genome-wide parental genotype distribution, with the color

indicating the proportion of parental genotype occurrence (on the y-axis) per genetic

distance (on the x-axis). Linkage groups are in separate panels, and the X

chromosome (L.7) was plotted separately to show the distribution of heterozygous

genotypes compared to homozygous genotypes. Parental genotypes vary

consistently per linkage group due to differences in recombination frequencies

between linkage groups and within a linkage group. For details about the genotype

composition and genotype versus the distance per phenotype of individual RIAILs,

see Supplementary Figures S3 and S4.

Characteristics of the A. freiburgense genome

The resulting chromosomal-scale assembly, which was the result of the integration

of the genetic and physical maps, spanned 53.5 Mb (Table 2 and Table S2). The

genome contained 89.2% of complete nematode BUSCO orthogroups (Table S3).

The annotation of repeat sequences (as described in the Methods section) revealed

that 10 Mb (18.2%) of the genome was repetitive (Figure 6 and listed in Table S4).

Using a combination of *ab initio* predictors and evidence-based methods (including

curated proteins and transcriptomic data), we predicted 8,759 protein-coding genes

(Figure 6, Table 2). To assess the completeness and quality of the predicted protein-

coding genes, we used BUSCO (nematoda_odb10 database) on the annotated

transcript dataset. The annotated transcripts contained 84.8% complete BUSCO

orthogroups with 1.5% duplicated BUSCOs. The total percentage of repeat DNA in

the table excludes the percentage of repeat DNA in unplaced scaffolds (1.5%).

Table 2. Characteristics of the *A. freiburgense* genome.

LG	N markers	Length (Mb)	N genes	% repeat DNA
L.1	1079	7.2	1112	19.7
L.2	4404	13.0	1964	20.4
L.3	3337	11.4	1833	13.6
L.4	1638	5.3	896	14.9
L.5	2290	8.0	1442	10.9
L.6	1412	5.0	979	8.8
L.7	2632	3.3	496	35.5
Total	16792	53.5	8759	16.7

Figure 6. The density of genes, repeats, and GC content across the seven chromosomal-scale scaffolds of *A. freiburgense*. Gene density (blue), repeats (orange) and GC content (red) were plotted along the *A. freiburgense* chromosomal-scale assembly using a window size of 200 kb.

Next, we examined the chromosomal-scale genome of *A. freiburgense* to examine the evolution of chromosomal gene content in nematodes. Previous research has identified seven possible ancestral chromosomal units of Rhabditida nematodes, called Nigon elements [21,22]. They represent the reconstructed genic content of the hypothesized seven chromosomes of the ancestor of all Rhabditida nematodes. The

Nigon elements in *A. freiburgense* are mostly fragmented into different

chromosomes (Figure 7). Notably, Nigon B is fragmented into three different

chromosomes in *A. freiburgense* (although it is intact in *A. rhodense*) (Fig 7A). In

contrast, Nigon N, which is present in two chromosomes in *A. rhodense*, is mainly in

one chromosome in *A. freiburgense* (Figure 7B). When analyzed in context with

other nematodes, *A. freiburgense* chromosomes underwent recent fission and fusion

events of Nigon elements (Figure 7B, Figure S5, Table S5).

Figure 7. Gene painting according to their Nigon Element association. (**A**)
Nigon elements in *A. freiburgense*. (**B**) Model of chromosome evolution in
Rhabditida.

Identifying a candidate region on the X chromosome for the rates of
production of males

Next-generation sequencing combined with bulk segregant analysis (NGS-BSA) is a

high throughput strategy to identify potential QTL regions associated with traits of

interest [23]. We used NGS-BSA as a first approach, by preparing four pools of DNA,

two for HM lines and two for LM lines, each containing DNA from 10 different RIAILs

(see Materials and Methods). The delta SNP index refers to the ratio of the

frequency of a specific single nucleotide polymorphism (SNP) in a pooled sample of

individuals with a particular phenotype (e.g., high-male or low-male) to the frequency

of the same SNP in a pooled sample of individuals with the opposite phenotype [24]. In

this case, the SNP index represents the ratio of the frequency of the alternative allele

(APS14) to the frequency of the reference allele (APS7). It is used as a measure of

the degree of association between a particular SNP and a trait of interest in bulk

segregant analysis. After sequencing the four pools, we found two peaks of high

delta SNP-index associated with high male rates (Figure 8 and S6). The first region

is on L.5, about 6.3 Mb long, and the second is on L.7 (ChrX), approximately 1.0 Mb

long (Figure 8, Table S6).

To narrow the candidate locus further, we performed QTL analysis using genome-

wide SNPs from individual RIAILs. With this approach, we confirmed the QTL on the

X chromosome associated with male rates (Figure 9). The QTL region above the

genome-wide threshold at p-value 0.2 spans 0.85 Mb and is located between the

positions 1,949,677 bp and 2,806,499 bp on the X chromosome (Figure 10). This

region overlaps with the region in the X chromosome identified in the NGS-BSA

analysis (Table S6). The peak of the QTL region was determined using the genome-

wide threshold at p-value 0.05, which extends from 1,949,688 bp to 2,372,281 bp,

with the highest LOD score of 3.37 at position 2,356,542 (Figure 9 [20].

Figure 8. Next-generation sequencing coupled with bulk segregant analysis

(NGS-BSA). (**A**) RIAILs were classified as either HM or LM based on the ratio of

male progeny resulting from crosses with APS7 females. Variants were identified in

two pools of 20 individuals each from each category. The SNP-index was calculated

for each SNP from LM-pool (**B**), HM-pool (**C**), and the difference in SNP-index (delta

SNP-index) between the two pools (**D**, green dots). The black line represents a

tricube-smoothed SNP-index in a sliding window of 1 Mb. The confidence intervals

for delta-SNP-index are indicated in red (95%) and blue (99%).

Figure 9. Genome-wide QTL association with the male rate phenotype. The

threshold was obtained based on a significance level (p-value) of <0.05 and <0.2,

corresponding to LOD (logarithm of odds) scores of 2.98 (blue line) and 2.35 (red

line), respectively.

The QTL region contains significant variation between APS7 and APS14

The variant calling analysis [25] detected 354,705 variants between APS7 and APS14,

resulting in an average density of one variant per 155 bp. Most of these variants

(84.1%) were single nucleotide polymorphisms (SNPs) (Table S7). Of the identified

SNPs, 0.1% were nonsense, and 5% were missense SNPs. In addition to SNPs,

8.6% of all variants are insertions, and 7.2% are deletions (Table S7).

The QTL region contains 122 predicted protein-coding genes, 367 repeat motifs, and

307 tRNA sequences [20]. SnpEff is a variant annotation and effect prediction tool to

annotate and predict the effects of genetic variants, such as single nucleotide

polymorphisms (SNPs), insertions, deletions, and structural variants, on genes and

proteins [26]. Using this tool, 362 missense variants were predicted within the QTL

region on the X chromosome [20]. 73/122 (59.8%) of the predicted protein-coding

genes have at least one missense variation, causing an amino acid change [20]. In

addition, we found 390 insertions, 353 small deletions, and 12 large structural

deletions [20]. The tRNA sequences, specific to APS7, clustered in a region free of

protein-coding genes (2,566,140 bp - 2,703,460 bp) (Figure 10) [20]. Due to the high

number of variants within the QTL region, we were unable to pinpoint a particular

functional component that regulates organelle distribution during spermatogenesis in

A. freiburgense. Further testing with functional studies of candidate genes will be

required to identify the locus associated with the rates of male production.

361 **Figure 10. Variants between APS7 and APS14 in the QTL region.** The top half of

362 the plot shows the APS14 variants (relative to the reference genome of APS7) with

363 missense variants, large structural variant deletions, and small indels. The bottom

364 half of the plot shows the APS7 annotation of repeats, genes, and tRNAs.

Discussion

Auanema populations consist of selfing hermaphrodites, females, and males [17,18].

Female versus hermaphrodite development is determined intergenerationally

according to maternal age or exposure to social cues [19,22,27-31], while males are

determined chromosomally [12].

In *Auanema*, the process of spermatogenesis in XO males and XX hermaphrodites

involves the elimination of nullo-X spermatids [7]. Males produce predominantly X-

bearing sperm [12,13], resulting in non-Mendelian transmission of the X chromosome,

resulting in mostly XX offspring when mating with females [17]. When mating with

hermaphrodites, males predominantly sire sons [7]. This occurs due to meiotic non-

disjunction of the X chromosome in hermaphrodite oocytes [7], which leads to the

production of nullo-X oocytes. As a result, the resulting embryos lack an X

chromosome, leading to the development of male individuals when fertilized by the

X-bearing sperm of the male [7].

Unlike in *C. elegans* spermatogenesis, where non-sperm components are deposited

into a residual body without DNA [16], *Auanema* eliminates those cytoplasmic

components together with autosomes. Similar phenomena are observed in other

organisms, such as sciarid flies, which undergo asymmetric segregation of

chromosomes during meiosis and eliminate an entire set of chromosomes into a

residual body (for review, see [32,33]). Scale insects also remove sets of

heterochromatic chromosomes during spermatogenesis [34].

The removal of DNA observed in *Auanema* and other organisms represents a type of

Programmed DNA Elimination (PDE), which has independently evolved in different

taxa [35-38]. PDE processes may influence sex ratios in various species. In *Nasonia*

wasps, for instance, PDE removes sets of chromosomes to convert female embryos

into males [39,40]. In the nematode *Strongyloides*, nullo-X sperm, responsible for male

development, is eliminated during spermatogenesis [14], while in XX embryos, specific

portions of the X chromosome are targeted for degradation to generate males [41].

Hermaphrodites of the nematode *Rhabdias* also expel one of their X chromosomes

into a residual body during spermatogenesis [42]. The mechanisms of how specific

chromosomes are targeted for PDE are still not well understood.

The study of meiotic drive and PDE mechanisms has often been hindered by the

lack of genetic tools available for many organisms that exhibit these processes.

However, *Auanema* has emerged as a promising model system for investigating

these mechanisms due to its advantageous features, including easy husbandry,

short life cycle, large brood sizes, and the recent development of genetic tools and

resources [22,31,43,44].

In this study, we aimed to improve our understanding mechanisms of X-chromosome

drive in *A. freiburgense,* generating genetic resources in the form of a chromosomal-

scale assembly of its genome. To enhance the contiguity of the assembly, we

constructed independent genetic linkage maps for the autosomes and the X

chromosome. By accounting for differences in recombination events between these

chromosomes during the inbreeding phase of Recombinant Inbred Advanced

Intercross Lines (RIAILs), we were able to improve the contiguity of the X

chromosome assembly. Our efforts resulted in a 47.3% increase in the size of the X

chromosome compared to the initial genetic linkage map (Table S1). This improved

assembly will provide valuable insights into the unique biology and mechanisms

underlying PDE in *A. freiburgense.*

We serendipitously discovered that by crossing two inbred strains of *A. freiburgense*,

males of some hybrid lines can produce more sons than the parental strains. By

using genetic markers specific to the X chromosome, we found that males in these

transgressive lines retain the nullo-X sperm, which is absent in the parental lines. As

a result, when these males mate with females, there is an increased proportion of

sons. This 'new phenotype' is typical of transgression, which is defined as the

appearance of a phenotype in F2s that is more extreme than the parental

phenotypes [45-48].

Novel allele combinations can result in phenotypes that exceed the range observed

in either parent, often with extreme or unexpected traits [47]. However, this mechanism

is unlikely to be the case for high male production in *A. freiburgense*, given that all

lines with transgressive phenotypes are inbred and therefore likely to be

homozygous in most loci. Furthermore, the parental inbred strains APS7 and APS14,

which were originally derived from the outbred strains SB372 [18,49] and JU1782 [29,50]

respectively, did not show transgressive phenotypes during the process of

inbreeding.

Lines with high-frequency males had an overrepresentation of homozygous APS14

alleles in a region of the X chromosome, indicating that this region is associated with

the phenotype. Since the same allele combination was present in the parental line

APS14, we can discard a model based on dominance or exposure of recessive

alleles [47]. Instead, the 'high male' phenotype is likely to be the result of epistatic

interactions between a combination of homozygous alleles of more than one locus.

Although the BSA indicated that a second locus associated with high production of

males is located in chromosome 5, we could not confirm this by QTL mapping using

whole genome sequencing of all RIAILs. The success and precision of NGS-BSA

rely on the high number of individuals used per bulk and the coverage of

sequencing. Even though we sequenced bulks at high coverage, there were only 10

lines per bulk. We reason that the genetic variation in the small number of lines

selected per bulk did not represent the genetic variation in all of the RIAILs sharing a

similar phenotype. As a result, another candidate region that is probably not

associated with the phenotype was detected. To improve confidence in NGS-BSA,

future experiments should increase the number of lines per bulk by pooling all the

HM-RIAILs in one bulk and the LM-RIAILs in another bulk to capture all the genetic

variations within the RIAILs.

The nature of nullo-X elimination by a polarizing cue from the X chromosome is still

unclear. In the ~800 kb mapped QTL region there are 123 predicted protein-coding

genes, and 75 of the candidate protein-coding genes have at least one missense

mutation. In addition, there are structural variants in non-coding regions that may

mediate the polarization of cytoplasmic components during meiosis. Further

investigation of those variants could shed more light on the biological mechanism

underlying DNA elimination during sex determination by polarizing the spermatocyte

cytoplasm to generate viable sperm and non-viable spermatids.

Figure 11. Proposed models of spermatogenesis in HM and LM RIAILs. (A)

Male cross-progeny resulting from a cross between APS7 females and LM lines

inherited the paternal X chromosome, indicating that nullo-X sperm are discarded

during the spermatogenesis of males from those lines. **(B)** Nullo-X sperm could be

produced by symmetrical distribution of sperm components between X-bearing and

nullo-X sperm during anaphase II. The symmetric segregation could be total, where

subcellular compartments, including sperm components, segregate equally, or a *C.*

elegans-like anaphase II, where a central residual body is formed for discarded

materials.

Author contributions

T.A. and A.P.S. designed the study. T.A., S.A., A.T., and J.L. conducted the

experimental work. Genome assembly, annotation, and analysis were conducted by

T.A., S.T., S.A., and J.K. The article was written by T.A. and A.P.S. with input from

all other authors.

Acknowledgments

A.P.-d.S and S.A. were supported by grants from BBSRC (BB/L019884/1) and

Leverhulme Trust (RPG-2019-329). S.T. was funded by a full Ph.D. scholarship from

the program Ciência sem Fronteiras (CNPq agency, process number 201116/2014-

6). A.T. was funded by the Doctoral Training Program from Natural Environment

Research Council (NERC CENTA). J. L. was supported by Samsung Science and

Technology Foundation SSTF-BA1501-52.

Materials and methods

Strains and their maintenance

A. freiburgense (previously known as *Rhabditis* sp. SB372, or *A. freiburgensis*) was

first isolated in Freiburg, Germany, by Prof. Walter Sudhaus from a horse dung pile

[18,49]. The *A. freiburgense* SB372 strain underwent eleven generations of

bottlenecking (expansion from a single-selfing hermaphrodite) to produce the inbred

strain APS7 [19]. *A. freiburgense* JU1782 strain was isolated from a rotting *Petasites*

stem sampled in Ivry (Val-de-Marne, France) [30] by Marie-Anne Felix. JU1782

underwent bottlenecking for ten generations and was renamed inbred strain APS14.

Nematodes were maintained on NGM plates seeded with *Escherichia coli* OP50-1 at

20 °C.

Sexual morph identification and female-male cross

In uncrowded conditions, *A. freiburgense* hermaphrodites produce mostly female

and male progeny [29,50]. *A. freiburgense* dauer larvae invariantly develop into self-

fertilizing hermaphrodite adults [18,29]. Therefore, to isolate female and male sexual

morphs for crosses, dauer larvae were incubated on NGM OP50-1 plates at a low

density (3 dauers per 2 cm diameter OP50-1 bacterial lawn) until they reached

adulthood and began laying eggs (approximately 48 h after collection) [19].

Approximately 36 h after the start of egg-laying, L2 stage female larvae were

differentiated from males by tail morphology and moved to fresh female-only plates,

to prevent fertilization. After 24 h, virgin females reached adulthood and were used in

crosses. Young adult males were isolated from the original low-density plates.

The crosses were conducted with a ratio of one female to one male. The male was

subsequently removed after 24 hours, and the mother was transferred to a new

plate. The male and non-male progeny were counted 48 to 72 h after egg laying and

the counts were combined for all 3 plates for each cross.

Generation of A. freiburgense *Recombinant Inbred Advanced Intercross Lines*

(RIAILs)

One hundred *A. freiburgense* recombinant inbred advanced intercross lines (RIAILs)

were generated by crossing APS7 males with APS14 females. Hybrid F1 progeny

resulting from the cross was left to mate (or self-reproduce) in a large mating pool.

F1 self-reproducing hermaphrodites and females, each mated with one or more

males, were isolated to establish the lines. Three to five F2 females were picked

from each line and crossed with two males from a different line, in an inbreeding

avoidance scheme. Inter-crosses between lines from F2 to F7 were established in a

way that every two lines were only crossed once to maximize haplotype breakpoints.

After seven generations of inter-line crosses, lines were inbred by single worm

descent for ten generations to bring alleles into homozygosity at most loci. Once the

17th generation was reached, the 100 lines were maintained by sampling animals

from a crowded plate to a fresh new plate once a month.

Crossing males from A. freiburgense *Recombinant Inbred Advanced Intercross Lines*

(RIAILs) with wild-type APS7 females

In each cross, a male from each RIAIL was crossed with a wild-type APS7 female for

~24h. After the cross, males were removed. The fertilized female was moved to a

new plate every day until it stopped laying eggs, to synchronize the growth of the F1

progeny. Ratios of male-to-female progeny were scored for each cross. Males from

all lines were crossed except lines number 17, 48, 79, and 107.

DNA Extraction from A. freiburgense *Recombinant Inbred Advanced Inter-crosses*

Lines (RIAILs) and sequencing

For each line, the nematodes were harvested with water from five NGM plates (10

cm in diameter). They were collected in a conical tube and washed 2-3 times with

water. After each wash, nematodes were let to settle naturally to the bottom of the

tube rather than by centrifugation. 500 µl of lysis buffer (100 mM Tris (pH 8.5), 100

mM NaCl, 50 mM EDTA, 1% SDS, and 1% beta-Mercaptoethanol) was added and

tubes were frozen at -80 °C overnight.

Three cycles of thawing and freezing were performed before adding 2.5 µl of

proteinase K (20 mg/ml) to each tube, followed by incubation at 65 °C for 3-4 hours.

DNA extraction was performed with the Gentra Puregen Core kit (Qiagen) following

the manufacturer's instructions.

Sequencing libraries were generated using TruSeq DNA nano gel free at the

GenePool facility at the University of Edinburgh. Sequencing libraries were

sequenced using the Illumina HiSeq platform to generate 150-bp paired-end reads,

with an insert size estimation of 350 bp.

DNA extraction from A. freiburgense *for PacBio, Illumina mate-pair, and pair-end*

sequencing

For PacBio long-read sequencing, DNA was extracted from plates containing *A.*

freiburgense APS7 of various stages. To lyse worms, we used 6 mL of 55 °C Cell

Lysis Solution from The Gentra Puregene® Cell and Tissue Kit (Qiagen) with 0.1

mg/mL proteinase K and 1% β-mercaptoethanol. The solution was directly poured

into a 200-µL worm pellet, and worms were lysed in the solution at 55 °C for 8 h with

occasional inverting. Polynucleotides were purified from the mixture by using phenol-

chloroform-isoamyl alcohol (25:24:1 v/v) DNA extraction and ethanol precipitation

methods coupled with phase-lock gel to minimize pipetting and DNA shearing.

Purified polynucleotides were re-dissolved in the TE buffer and treated with 10

µg/mL RNase for 2 h. DNA was extracted by the same DNA purification procedure,

and dissolved in 10 mM Tris-HCl (pH 8.0). Macrogen (South Korea,

https://www.macrogen.com/en/main) performed PacBio library preparation and

sequencing in the Sequel platform with the continuous long-read sequencing mode.

DNA extraction for Illumina mate pair and pair-end sequencing of APS7 was

performed as described for the RIAIL sequencing, using DNA from dauer larvae

collected from 200 plates, as previously described [51].

Genome assembly

The assembly of the PacBio data was performed with the Canu software (version

1.6) [52], using a minimum read length (minReadLength) of 3 kb and setting the

corrected error rate (correctedErrorRate) to 0.030, as this resulted in the most

contiguous assembly. The estimated genome size (genomeSize) was set to 55 Mb.

Bacterial contamination was identified and removed through BLASTn (version 2.7.1)

[53] alignments to a contaminant database that included 3,000 bacterial genomes

downloaded from the European Nucleotide Archive (ENA) on March 30, 2018. The

following BLAST parameters were used "-task megablast -evalue 1e-06 -outfmt 6 -

perc_identity 50".

We first polished our preliminary draft genome using Quiver with PacBio raw reads

aligned to the assembly using pbalign (version 0.3.1). We then further polished the

assembly using Pilon with all available Illumina short reads. Illumina short reads

were first preprocessed using skewer (parameters " -n -Q 20 -l 51"). Trimmed reads

were then aligned against the assembly using BWA. These alignments were then

used to fix base-level inconsistencies between the Illumina read and the genome

assembly using Pilon ("--changes --fix bases --chunksize 8000000 --diploid [54].

A list of all the programs used in all bioinformatics analyses, their versions, and

parameters are included in Table S10.

RIAILs genotyping

The assembled genome polished with Illumina reads was used as a reference

genome to genotype the 100 RIAILs and the APS14 parent samples. The quality of

individual samples' raw reads was checked using FastQC to ensure sequencing

adaptors were removed. Each line/strain paired-end DNA sample was aligned

against the long-read genome assembly using BWA aligner [55]. Sequence Alignment

Map (SAM) files were converted into Binary Alignment Map (BAM) files, then sorted

according to the position in the reference genome [56]. Aligned reads in every BAM file

were assigned a new read-group tag using the Picard AddOrReplaceReadGroups

tool to make it compatible with the genome analysis toolkit GATK pipeline [57].

Variants from every alignment bam file with a new read-group tag were identified

against the APS7 reference genome using genome analysis toolkit GATK tools,

producing a single variant calling file containing the variants of each sample against

the reference APS7 genome. Low-quality variants were filtered with VariantFiltration

provided by GATK tools using default parameters [25,57-59]. Then variants were further

filtered using vcftools keeping only variants with a minimum depth of 10, and a

minimum genotype quality of 30, removing variants that are missing in more than

25% of RIAILs, removing all the indels, and only keeping bi-allelic SNP variants [60].

The final 274,394 SNPs were used as markers to construct a genetic linkage map.

Construction of a high-density genetic linkage map

The genetic linkage map was constructed using R/qtl and ASMap R packages [61,62].

R/qtl was used to process markers pre-construction of the genetic linkage map and

ASMap for the genetic linkage map. Markers were imported into R using rqtl as RILs

data, crosstype = "riself", expecting no heterozygous markers in the dataset,

heterozygous markers were assigned a missing value. 10,326 markers either

missing the genotype of one of the parents or found heterozygous were omitted from

the data set, leaving 264,068 markers from 100 samples on 40 scaffolds. Using the

rqtl function 'drop.markers' the marker set was further refined by (i) removing

markers that were absent in 90% of the data allowing only 10% of missing values, (ii)

removing markers that shared similar genotypes to reduce redundancy in the data

set and (iii) removing markers with abnormal segregation distortion (P. value < 0.05).

After that, ASMap R package was used to construct a genetic linkage map with the

remaining 14,955 markers from 100 samples on 25 scaffolds. The genetic map was

constructed using the 'mstmap.cross' function provided by the ASMap package

using a p value of 1e-11. Genetic distance between markers was computed and

markers were linked and organized into 7 main linkage groups. An initial genetic

linkage map of seven linkage groups was constructed using 14,884 markers from

100 samples on 22 scaffolds representing 93.7% of the genome.

To improve X chromosome assembly, another genetic linkage map was constructed

specifically for the X chromosome by including all the heterozygous and

homozygous markers and removing all autosomal scaffolds. Linkage group 7 from

the initial genetic linkage map was identified using synteny mapping as the X

chromosome in *A. freiburgense*. To construct an independent genetic linkage map

for the X chromosome, markers were imported into rqtl as "f2 intercross" to include

all the heterozygous and homozygous markers. In total, 264,068 markers from 40

scaffolds were imported. Using the rqtl function 'drop.markers', we removed markers

that were not present in 75% of the data allowing only 25% of missing values, and

redundancy in the data set was reduced by removing markers that share similar

genotypes. Then, markers belonging to autosomal linkage groups (L.1 to L.6) were

removed leaving only 21 scaffolds with 3,128 markers from X chromosome scaffolds

and unplaced scaffolds. Before linkage map construction, using ASMap "pullCross"

function, markers with more than 10% missing values and those with a high

segregation distortion (segregation ratio less than 1:98:1) were removed to be

pushed back into the map after construction. The X chromosome genetic map was

constructed using the 'mstmap.cross' function provided by the ASMap package

using a p value of 1e-12. Using the "pushCross" function most markers with high

segregation distortion were pushed back to the constructed map using a segregation

ratio threshold of (0.5:99:0.5). Small linkage groups were subsetted leaving only the

main linkage group representing the X chromosome with 2,632 markers. The new X

chromosome genetic linkage map was added to the previously constructed genetic

linkage map replacing Linkage Group 7 (L.7). The final genetic linkage map of 7

linkage groups contained 16,792 markers from 29 scaffolds and represents 97% of

the genome [20].

Synteny analysis and identification of ancestral chromosomal elements (Nigon

Elements)

Scaffolds were anchored to the genetic map using ALLMAPS software, producing a

chromosomal-scale assembly, where scaffolds were ordered and oriented into their

respective position within each linkage group [63]. The chromosomal assembly was

aligned to the *C. elegans* genome using MUMmer with default parameters and

macro-synteny patterns were visualized using Circos [64,65].

To assess the completeness of the genome assembly, a BUSCO analysis was

conducted in the genome mode using Augustus gene discovery against the latest

nematode database (nematoda_odb10) (Table S3 and S8). To examine the patterns

of Nigon Elements in each chromosomal-scale scaffold, we used the program vis-

ALG (https://github.com/pgonzale60/vis_ALG) [21]. Briefly, we used the location of the

BUSCO genes identified in *A. freiburgense* and the association between BUSCO

genes and Nigon Elements previously determined to paint the chromosomal-scale

scaffolds according to the Nigon Elements.

X-chromosome genotyping and calculation of nullo-X ratio

To follow the X chromosome inheritance, we used an X-linked polymorphic marker

(X_{634}) where a HindIII restriction site (AAGCTT) is present in APS7 but not in the

APS14 strain (AAACTT). X chromosome genotyping was conducted by PCR

amplification of a region spanning the SNP followed by digestion of the product using

HindIII. Genomic DNA was extracted from individual males using a modified single-

worm PCR method. A single male was frozen in 20 µl of 1x PCR buffer at -80 °C for

a minimum of 24 h. After thawing, the tissue was lysed and genomic DNA was

released by the addition of 0.5 µl of proteinase K (20 mg/ml) and incubation at 65 °C

for 60 mins, followed by 95 °C for 15 mins, to inactivate the enzyme. Samples were

frozen at -20 °C for at least 24 h before use in PCR. Each PCR reaction was

conducted with 2 µl of DNA, 10 µl of GoTaq Green Master Mix (Promega), 10 µM

forward primer (UW634_F 5'-AGGGACACGATTGCCTTCTG-3'), and 10 µM of the

reverse primer (UW635_R 5'-AATGCCGCGGAGGTCTTTAA-3') in a final volume of

20 µl. The following cycling conditions were applied: 94 °C for 5 min, followed by 30

cycles of 94 °C for 15 sec, 55 °C for 30 sec, and 72 °C for 1 min. The PCR products

were digested by direct addition of 0.5 µl of HindIII (Promega) and incubation for 1 h

at 37 °C. The genotype of each sample was determined by agarose gel

electrophoresis of the digested sample. The APS7 X allele gave 2 fragments (329 bp

and 234 bp) and the APS14 allele remained undigested (563 bp). Note the X_{634}

marker region is outside the QTL region (coordinates 438,785 to 439,347).

To determine the percentage of nullo-X sperm that contributed to the generation of

sons, the paternal X chromosome was genotyped using the X-linked marker X_{634}

(Table 1). To determine the percentage of viable nullo-X sperm for each strain, we

multiplied the percentage of sons resulting from crosses by the proportion of sons

that inherited the maternal X chromosomes from the total number of genotyped

males [20], as only sons that have a maternal X are derived from the fertilization of a

nullo-X sperm.

Structural annotation of the genome

A comprehensive repeat library was produced for the genome assembly using

multiple repeat-finding programs. The pipeline was based on the TransposableELMT

wrapper script (https://github.com/PlantDr430/TransposableELMT). Repeats were

identified using RepeatModeler v2.0.1 [66], TransposonPSI v08222010 [67], LTRfinder

v1.0.7 [68] and LTRharvest [69](implemented in GenomeTools v1.6.1 [70]. To limit the

identification of false positives, the LTRharvest output was post-processed with

LTRdigest [71] (implemented in GenomeTools v1.6.1 [70]). The resulting libraries were

combined, classified using RepeatClassifier v2.0.1 (part of the RepeatModeler v2.0.1

package [68] and redundancy removed using USEARCH [72] based on 90% similarity.

The non-redundant custom library was used to soft mask repeat regions in the

assembly with RepeatMasker v4.1.0 [73].

Gene predictions were made using the *ab initio* and evidence-driven gene predictors

GeneMark-ES [74], SNAP (implemented in Maker2) [75], Maker2 [76] and Augustus

[77](trained with BUSCO v5.1.3 [78]).

An *A. freiburgense* transcriptome assembled using Trinity [79], the *C. elegan*s protein

database (UP000001940_6239), and the UniProt/Swiss-prot database

(uniprot_sprot.fasta) were used as evidence-based inputs for the first round of

Maker2. The resulting output was then used to train Augustus and SNAP. The

unmasked genome was used as input into GeneMark-ES. Finally, the outputs from

the first round of Maker2, Augustus, SNAP, and GeneMark were used as inputs in

Maker2. The gene predictions from the second round of Maker2 were used for our

analyses. The annotation was later lifted over to the final chromosomal-scale

assembly using 'flo' using chain files [80].

Functional annotation of the genome

The functional annotation of the genome was conducted using the Blast2Go [81]. A

local Blast search of the *A. freiburgense* predicted proteins was conducted using a

searchable database of the Swissprot proteins (swissprot.gz downloaded from NCBI

July 2021) prepared within the Blast2Go software. The same predicted proteins were

also analyzed with InterProScan (within the Blast2Go software) and the annotations

merged to give the final functional annotation.

The completeness and quality of the predicted protein-coding genes were assessed

using BUSCO on the transcript dataset in genome mode with gene discovery via

Augustus against the nematode database (nematoda_odb10) (Table S9).

Mitochondrial genome

The mitochondrial scaffold was identified by BLASTn (Version 2.9.0+)[53], using the *A.*

rhodense mitochondrial genome as a reference database. Only one scaffold was

identified as the mitochondrion. This scaffold was analyzed and annotated using

MITOS [82], which revealed a large duplication, probably due to a misassembly due to

the mitochondrial genome being circular. We used the pair-wise local aligner Water

[83] to identify precisely the junctions of the duplicated region and removed it using the

subseq function of seqkit (version 0.16.1) [84]. The curated mitochondrial genome was

re-integrated into the assembly and replaced with the un-curated one. The

annotations associated with the un-curated mitochondrial genome were removed

and replaced by the MITOS annotations.

Pooling DNA from lines with similar phenotypes into discrete pools and sequencing

Equal amounts of DNA from 10 RIAIL lines with the same phenotype were mixed to

create pools of DNA from High Male lines (HM-Pools) and Low Male lines (LM-

Pools). Two pools for each category were prepared, each containing 1.5 µg of DNA.

The RIAIL lines used for the first and second HM-Pools were 23, 24, 26, 28, 29, 30,

33, 45, 57, 61 and 35, 42, 46, 55, 56, 59, 63, 65, 72 and 97, respectively. The RIAIL

lines used for the first and second LM-Pools were 2, 5, 9, 13, 14, 17, 20, 54, 74, and

77, and 6, 8, 15, 18, 25, 27, 37, 40, 69, and 95, respectively. The quality of DNA was

examined by running 1 µl of DNA on 1.8 % agarose gel, and concentration was

measured using a Qubit fluorometer. Five DNA samples; 2 HM-pools, 2 LM-pools,

and DNA from APS14 maternal strain, were sequenced. Sequencing libraries were

generated using TruSeq DNA nano gel free at the GenePool facility at the University

of Edinburgh on an Illumina HiSeq platform to generate 150-bp paired-end reads,

with an insert size estimation of 350 bp.

Variants calling for APS14 and next-generation sequencing Bulk segregant analysis

(NGS-BSA)

The quality of the reads was assessed using FastQC software to have an overview

of the reads' quality [85]. Paired-end reads were cleaned using Skewer using the

following parameters"-n -Q 20 -l 51 -t 32 -m pe" [86]. The quality of reads was

reexamined after cleaning using FastQC software. Then, BWA program was used to

align short reads from each pool and APS14 strain separately to the APS7 reference

genome [55]. Alignment files in SAM format were converted to BAM files using the

samtools "view" command [56]. BAM files from the HM-pools and the LM-pools were

merged to produce a single BAM for each phenotype and were subsequently sorted

using samtools [56]. Variants from sorted BAM files of both pool samples and APS14

strain were called Individually using "Haplotypecaller" from GATK (Genomic Analysis

Toolkit) [25]. Low-quality variants were filtered out with 'VariantFiltration' provided by

GATK tools using default parameters [25,57-59]. Variants were filtered further using

bcftools based on Genotype Quality >= 30 and depth >= 10 while keeping

homozygous sites only [60]. SnpEff (v5.0.1) [26] and SnpSift (v4.3.1) [87] were used to

annotate and predict the effects of the genetic variants between APS14 and APS7

reference genome. Large structural variants between APS14 and APS7 genomes

were identified using the Parliament2 pipeline [88]. Structural variants were called

using the following software: Breakdancer [89], CNVnator, [90], Manta [91], Lumpy [92] and

DELLY [93]. To avoid false positive calls we only considered structural variants that

were called with more than one software.

Pools' variant calling files were merged into one file using GATK "GenotypeGVCFs"

command [25]. Joint genotyping using "GenotypeGVCFs" combined all SNPs and

indels records from both pools to produce correct genotype likelihood outputting a

single combined variant calling file (VCF) [94]. Variants were filtered using vcftools [60]

removing all sites with missing values, removing all Indels, and keeping only biallelic

sites. In total 334,230 SNPs were kept. The joint variant calling file was converted to

a table using "VariantToTable" command provided by the GATK [25]. Regions

displaying differences between the HM-pools and LM-pools were identified using the

R-package "QTLseqr" [95]. QTLseq analysis for NGS-BSA was used to calculate the

allele frequency difference (SNP-index) from the allele depth at individual SNP [24].

Candidate regions were identified by setting a sliding window size to 1 Mb, and the

number of SNPs was counted in that window. Within each sliding window, a tricube-

smoothed delta-SNP-index was calculated by constant local regression. A simulation

was performed where the delta-SNP-index per bulk was calculated and simulated

over 1000 replications based on RILs F2 population with a bulk size of 20.

Confidence intervals at 95% and 99% were estimated using the quantile from the

simulation. An alternative approach, G-statistics, was used to identify significant

QTLs from BSA [96]. G-statistics was calculated genome-wide, and a tricube-

smoothed G-statistics (G') was predicted in a sliding window of 1M bp. P-values

were estimated and adjusted (Benjamini-Hochberg method), and negative log10-

were calculated from (G'). Significant regions were identified using a genome-wide

false discovery rate of 0.1 [97].

QTL analysis

The QTL analysis was conducted using rqtl on all the markers from all the scaffolds

before they were organized into linkage groups to include markers with high

segregation distortion that were omitted from the genetic linkage map and to include

all the scaffolds, including those that were excluded from the final chromosomal

assembly [61,98]. The 264,068 markers dataset imported into rqtl as RILs (see section

"Construction of high-density genetic linkage map") were filtered using the rqtl

function 'drop.markers' by removing markers that are not present in 90% of the data

allowing only 10% of missing values and redundancy in the data set was reduced by

removing markers that shared similar genotypes. All the markers with high

segregation distortion were retained in the final data set of 18,513 markers from the

100 samples on 28 scaffolds. Markers segregation distortion is a natural

phenomenon and the inclusion of distorted markers in QTL mapping increases the

power of detecting a QTL [98,99]. The interval mapping genotype probability was

calculated using the kosambi map function with a 1 CM step size and an error

probability of 0.001. A whole-genome scan on all RIAILs was performed with a single

QTL model using Haley–Knott regression, with the percentage of male progeny after

an outcross for each RIAIL as a phenotype [100]. The result from the first scan was

permuted 1000 times to obtain a genome-wide LOD score (logarithm of odds)

significance threshold [101]. Genome-wide thresholds were obtained at p values < 0.05

and < 0.2 corresponding to a LOD score of 2.98 and 2.35 respectively. Interval

estimates of the location of the QTL were obtained via 1.5-LOD support intervals,

which could be calculated via the function (lodint). Three QTLs on three different

scaffolds belonging to the X chromosome were detected from the single scan.

Scanning for the likelihood of interacting QTLs using a two-dimensional genome

scan, for all pairwise combinations of intervals, was impossible due to a large

number of markers [102]. Hence, a multiple QTL analysis using QTLs identified from

the single scan was performed. Briefly, a qtl object for multiple qtl analysis was

created containing all the identified QTLs from the single scan using the rqtl function

(makeqtl). QTLs were fitted into an additive model using the 'addinit' function, which

adds one interaction at a time in the context of a multiple interval mapping (MIM)

model, model formula: y ~ QTL1 + QTL2 + QTL3. An interaction was detected

between the three QTLs identified from the single scan at p-value (Chi2) < 0.05. The

interaction was expected as the three QTLs were on scaffolds assembled adjacent

to each other on the X Chromosome. The estimate of the qtl peak location was

refined using the 'refinqtl' function. Finally, the function 'addqtl' was used to scan for

an additional QTL to be added to the model. Markers outlining the start and the end

of QTL regions identified on scaffolds were overlapped with the genetic linkage map

to determine the physical location of the QTL region on the final genome assembly.

The three QTLs were adjacent to each other on the X linkage group, therefore they

were merged into a single QTL. In a separate analysis, the 18,513 markers dataset

from 22 scaffolds (nonredundant and includes all markers with segregation

distortion) used to scan for a QTL was merged with the genetic linkage map to obtain

the genetic position of each marker [20]. The analysis was repeated using all the steps

above. A single QTL was detected on the X chromosome confirming the three

identified QTLs in the earlier analysis are part of one continuous X chromosome

QTL.

Availability of the data

We deposited all sequencing data at the European Nucleotide Archive (ENA) under

BioProject numbers PRJEB55706 (Illumina genomic Data), PRJEB60474 and

PRJEB50372 (Illumina transcriptomic data), and PRJNA640723 (PacBio data).

Furthermore, we uploaded the genome assembly, annotation, and predicted protein

dataset to NCBI with accession PRJNA947217.

References

1   Hoeh, W. R., Blakley, K. H. & Brown, W. M. Heteroplasmy suggests limited
biparental inheritance of Mytilus mitochondrial DNA. *Science* **251**, 1488-1490
(1991). https://doi.org:10.1126/science.1672472

2   Zouros, E., Freeman, K. R., Ball, A. O. & Pogson, G. H. Direct evidence for
extensive paternal mitochondrial DNA inheritance in the marine mussel
Mytilus. *Nature* **359**, 412-414 (1992). https://doi.org:10.1038/359412a0

3   Zouros, E., Oberhauser Ball, A., Saavedra, C. & Freeman, K. R. An unusual
type of mitochondrial DNA inheritance in the blue mussel Mytilus. *Proc Natl*
Acad Sci U S A **91**, 7463-7467 (1994).
https://doi.org:10.1073/pnas.91.16.7463

4   Hurst, L. D. The incidences. mechanisms and evolution of cytoplasmic sex
ratio distorters in animals. *Biological Reviews* **68**, 121-194 (1993).
https://doi.org:https://doi.org/10.1111/j.1469-185X.1993.tb00733.x

5   Ross, L., Mongue, A. J., Hodson, C. N. & Schwander, T. Asymmetric
Inheritance: The Diversity and Evolution of Non-Mendelian Reproductive
Strategies. *Annual Review of Ecology, Evolution, and Systematics* **53**, 1-23
(2022). https://doi.org:10.1146/annurev-ecolsys-021822-010659

6   Burt, A. & Trivers, R. *Genes in conflict : the biology of selfish genetic*
elements. (Belknap Press of Harvard University Press, 2006).

7   Tandonnet, S. *et al.* Sex- and gamete-specific patterns of X chromosome
segregation in a trioecious nematode. *Curr Biol* **28**, 93-99 e93 (2018).
https://doi.org:10.1016/j.cub.2017.11.037

8   Blackman, R. L. Spermatogenesis in the aphid Amphorophora tuberculata
(Homoptera, Aphididae). *Chromosoma* **92**, 357-362 (1985).
https://doi.org:10.1007/BF00327467

9   Wilson, A. C. C., Sunnucks, P. & Hales, D. F. Random loss of X chromosome
at male determination in an aphid, Sitobion near fragariae, detected using an
X-linked polymorphic microsatellite marker. *Genetical Research* **69**, 233-236
(1997).

10   Pires-daSilva, A. Evolution of the control of sexual identity in nematodes.
Semin Cell Dev Biol **18**, 362-370 (2007).
https://doi.org:10.1016/j.semcdb.2006.11.014

11   Van Goor, J., Shakes, D. C. & Haag, E. S. Fisher vs. the Worms:
Extraordinary Sex Ratios in Nematodes and the Mechanisms that Produce
Them. *Cells* **10**, 1793 (2021).

12   Shakes, D. C., Neva, B. J., Huynh, H., Chaudhuri, J. & Pires-daSilva, A.
Asymmetric spermatocyte division as a mechanism for controlling sex ratios.
Nat Commun **2**, 157 (2011). https://doi.org:10.1038/ncomms1160

13   Winter, E. S. *et al.* Cytoskeletal variations in an asymmetric cell division
support diversity in nematode sperm size and sex ratios. *Development* **144**,
3253-3263 (2017). https://doi.org:10.1242/dev.153841

14   Dulovic, A., Koch, I., Hipp, K. & Streit, A. Strongyloides spp. eliminate male-
determining sperm post-meiotically. *Mol Biochem Parasitol* **251**, 111509
(2022). https://doi.org:10.1016/j.molbiopara.2022.111509

15   Al-Yazeedi, T., Xu, E. L., Kaur, J., Shakes, D. C. & Pires-daSilva, A. Lagging
X chromatids specify the orientation of asymmetric organelle partitioning in XX
spermatocytes of Auanema rhodensis. *Genetics* **222** (2022).
https://doi.org:10.1093/genetics/iyac159

16 Ward, S., Argon, Y. & Nelson, G. A. Sperm morphogenesis in wild-type and
fertilization-defective mutants of *Caenorhabditis elegans*. *J Cell Biol* **91**, 26-44
(1981).

17 Félix, M. A. Alternative morphs and plasticity of vulval development in a
rhabditid nematode species. *Dev Genes Evol* **214**, 55-63 (2004).

18 Kanzaki, N. *et al.* Description of two three-gendered nematode species in the
new genus *Auanema* (Rhabditina) that are models for reproductive mode
evolution. *Sci Rep* **7**, 11135 (2017). https://doi.org:10.1038/s41598-017-
09871-1

19 Adams, S. *et al.* Sexual morph specialisation in a trioecious nematode
balances opposing selective forces. *Sci Rep* **12**, 6402 (2022).
https://doi.org:10.1038/s41598-022-09900-8

20 Al-Yazeedi, T. *et al.* A QTL influences sex ratios by controlling asymmetric
organelle positioning during male spermatogenesis in *Auanema freiburgense*.
figshare, 10.6084/m6089.figshare.22820627 (2023).
https://doi.org:10.6084/m9.figshare.22820627

21 Gonzalez de la Rosa, P. M. *et al.* A telomere-to-telomere assembly of
Oscheius tipulae and the evolution of rhabditid nematode chromosomes. *G3*
(Bethesda) **11** (2021). https://doi.org:10.1093/g3journal/jkaa020

22 Tandonnet, S. *et al.* Chromosome-wide evolution and sex determination in the
three-sexed nematode *Auanema rhodensis*. *G3: Genes|Genomes|Genetics* **9**,
1211-1230 (2019). https://doi.org:10.1534/g3.119.0011

23 Li, Z. & Xu, Y. Bulk segregation analysis in the NGS era: a review of its
teenage years. *Plant J* **109**, 1355-1374 (2022).
https://doi.org:10.1111/tpj.15646

24 Takagi, H. *et al.* QTL-seq: rapid mapping of quantitative trait loci in rice by
whole genome resequencing of DNA from two bulked populations. *Plant J* **74**,
174-183 (2013). https://doi.org:10.1111/tpj.12105

25 Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls:
the Genome Analysis Toolkit best practices pipeline. *Curr Protoc*
Bioinformatics **43**, 11 10 11-11 10 33 (2013).
https://doi.org:10.1002/0471250953.bi1110s43

26 Cingolani, P. *et al.* A program for annotating and predicting the effects of
single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)* **6**, 80-92 (2012).
https://doi.org:10.4161/fly.19695

27 Chaudhuri, J., Kache, V. & Pires-daSilva, A. Regulation of sexual plasticity in
a nematode that produces males, females, and hermaphrodites. *Curr Biol* **21**,
1548-1551 (2011). https://doi.org:10.1016/j.cub.2011.08.009

960 28 Chaudhuri, J. *et al.* Mating dynamics in a nematode with three sexes and its
961   evolutionary implications. *Sci Rep* **5**, 17676 (2015).
962   https://doi.org:10.1038/srep17676

963 29 Zuco, G. *et al.* Sensory neurons control heritable adaptation to stress through
964   germline reprogramming. *bioRxiv*, 406033 (2018).
965   https://doi.org:https://doi.org/10.1101/406033

967 30 Robles, P. *et al.* Parental energy-sensing pathways control intergenerational
968   offspring sex determination in the nematode Auanema freiburgensis. *BMC*
969   *Biol* **19**, 102 (2021). https://doi.org:10.1186/s12915-021-01032-1

970 31 Tandonnet, S. *et al.* De Novo Genome Assembly of *Auanema melissensis*, a
971   Trioecious Free-Living Nematode. *J Nematol* **54**, 20220059 (2022).
972   https://doi.org:10.2478/jofnem-2022-0059

973 32 Goday, C. & Esteban, M. R. Chromosome elimination in sciarid flies.
974   *Bioessays* **23**, 242-250 (2001).

975 33 Gerbi, S. A. in *Germ Line — Soma Differentiation* (ed Wolfgang Hennig) 71-
976   104 (Springer Berlin Heidelberg, 1986).

977 34 Bongiorni, S., Fiorenzo, P., Pippoletti, D. & Prantera, G. Inverted meiosis and
978   meiotic drive in mealybugs. *Chromosoma* **112**, 331-341 (2004).
979   https://doi.org:10.1007/s00412-004-0278-4

980 35 Drotos, K. H. I., Zagoskin, M. V., Kess, T., Gregory, T. R. & Wyngaard, G. A.
981   Throwing away DNA: programmed downsizing in somatic nuclei. *Trends in*
982   *Genetics* **38**, 483-500 (2022).
983   https://doi.org:https://doi.org/10.1016/j.tig.2022.02.003

984 36 Dedukh, D. & Krasikova, A. Delete and survive: strategies of programmed
985   genetic material elimination in eukaryotes. *Biol Rev Camb Philos Soc* **97**, 195-
986   216 (2022). https://doi.org:10.1111/brv.12796

987 37 Kloc, M., Kubiak, J. Z. & Ghobrial, R. M. Natural genetic engineering: A
988   programmed chromosome/DNA elimination. *Developmental Biology* **486**, 15-
989   25 (2022). https://doi.org:https://doi.org/10.1016/j.ydbio.2022.03.008

990 38 Wang, J. & Davis, R. E. Programmed DNA elimination in multicellular
991   organisms. *Curr Opin Genet Dev* **27**, 26-34 (2014).
992   https://doi.org:10.1016/j.gde.2014.03.012

993 39 Nur, U., Werren, J. H., Eickbush, D. G., Burke, W. D. & Eickbush, T. H. A
994   "selfish" B chromosome that enhances its transmission by eliminating the
995   paternal genome. *Science* **240**, 512-514 (1988).
996   https://doi.org:10.1126/science.3358129

40　Werren, J. H. & Stouthamer, R. PSR (paternal sex ratio) chromosomes: the
ultimate selfish genetic elements. *Genetica* **117**, 85-101 (2003).
https://doi.org:10.1023/a:1022368700752

41　Nemetschke, L., Eberhardt, A. G., Hertzberg, H. & Streit, A. Genetics,
chromatin diminution, and sex chromosome evolution in the parasitic
nematode genus *Strongyloides*. *Curr Biol* **20**, 1687-1696 (2010).

42　Runey, W. M., Runey, G. L. & Lauter, F. H. Gametogenesis and fertilization in
Rhabdias ranae Walton 1929: I. The parasitic hermaphrodite. *J Parasitol* **64**,
1008-1014 (1978).

43　Kranse, O. *et al.* Toward genetic modification of plant-parasitic nematodes:
delivery of macromolecules to adults and expression of exogenous mRNA in
second stage juveniles. *G3 Genes|Genomes|Genetics* **11**, jkaa058 (2021).
https://doi.org:10.1093/g3journal/jkaa058

44　Adams, S., Pathak, P., Shao, H., Lok, J. B. & Pires-daSilva, A. Liposome-
based transfection enhances RNAi and CRISPR-mediated mutagenesis in
non-model nematode systems. *Sci Rep* **9**, 483 (2019).
https://doi.org:10.1038/s41598-018-37036-1

45　deVicente, M. C. & Tanksley, S. D. QTL analysis of transgressive segregation
in an interspecific tomato cross. *Genetics* **134**, 585-596 (1993).

46　Grant, V. *Genetics of flowering plants*. (Columbia University Press, 1975).

47　Rieseberg, L. H., Archer, M. A. & Wayne, R. K. Transgressive segregation,
adaptation and speciation. *Heredity (Edinb)* **83 (Pt 4)**, 363-372 (1999).
https://doi.org:10.1038/sj.hdy.6886170

48　Rieseberg, L. H., Widmer, A., Arntz, A. M. & Burke, J. M. The genetic
architecture necessary for transgressive segregation is common in both
natural and domesticated populations. *Philos Trans R Soc Lond B Biol Sci*
358, 1141-1147 (2003). https://doi.org:10.1098/rstb.2003.1283

49　Sudhaus, W. An update of the catalogue of paraphyletic 'Rhabditidae'
(Nematoda) after eleven years. *SOIL ORGANISMS* **95**, 95–116 (2023).
https://doi.org:10.25674/so95iss1id312

50　Robles, P. *et al.* Chromatin-associated effectors of energy-sensing pathways
mediate intergenerational effects. *bioRxiv*, 2020.2008.2031.275727 (2020).
https://doi.org:10.1101/2020.08.31.275727

51　Pires-daSilva, A. *Pristionchus pacificus* protocols. *WormBook*, 1-20 (2013).
https://doi.org:10.1895/wormbook.1.114.2

52　Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive
k-mer weighting and repeat separation. *Genome Res* **27**, 722-736 (2017).
https://doi.org:10.1101/gr.215087.116

53   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC*
Bioinformatics **10**, 421 (2009). https://doi.org:10.1186/1471-2105-10-421

54   Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial
variant detection and genome assembly improvement. *PLoS One* **9**, e112963
(2014). https://doi.org:10.1371/journal.pone.0112963

55   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-
Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009).
https://doi.org:10.1093/bioinformatics/btp324

56   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**, 2078-2079 (2009).
https://doi.org:10.1093/bioinformatics/btp352

57   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework
for analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297-
1303 (2010). https://doi.org:10.1101/gr.107524.110

58   DePristo, M. A. *et al.* A framework for variation discovery and genotyping
using next-generation DNA sequencing data. *Nat Genet* **43**, 491-498 (2011).
https://doi.org:10.1038/ng.806

59   Poplin, R. *et al.* Scaling accurate genetic variant discovery to tens of
thousands of samples. *bioRxiv*, 201178 (2018).
https://doi.org:10.1101/201178

60   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**,
2156-2158 (2011). https://doi.org:10.1093/bioinformatics/btr330

61   Broman, K. W., Wu, H., Sen, S. & Churchill, G. A. R/qtl: QTL mapping in
experimental crosses. *Bioinformatics* **19**, 889-890 (2003).

62   Taylor, J. & Butler, D. R Package ASMap: Efficient Genetic Linkage Map
Construction and Diagnosis. *Journal of Statistical Software* **79**, 29 (2017).
https://doi.org:10.18637/jss.v079.i06

63   Tang, H. *et al.* ALLMAPS: robust scaffold ordering based on multiple maps.
Genome Biol **16**, 3 (2015). https://doi.org:10.1186/s13059-014-0573-1

64   Krzywinski, M. *et al.* Circos: an information aesthetic for comparative
genomics. *Genome Res* **19**, 1639-1645 (2009).
https://doi.org:10.1101/gr.092759.109

65   Kurtz, S. *et al.* Versatile and open software for comparing large genomes.
Genome Biol **5**, R12 (2004). https://doi.org:10.1186/gb-2004-5-2-r12

66   Smit, A. F. A. & Hubley, R. *RepeatModeler Open-1.0*,
<http://www.repeatmasker.org> (2008-2015).

67 Haas, B. *TransposonPSI: An Application of PSI-Blast to Mine (Retro-*
)Transposon ORF Homologies, <https://transposonpsi.sourceforge.net/>
(2010).

68 Xu, Z. & Wang, H. LTR_FINDER: an efficient tool for the prediction of full-
length LTR retrotransposons. *Nucleic Acids Res* **35**, W265-268 (2007).
https://doi.org:10.1093/nar/gkm286

69 Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible
software for de novo detection of LTR retrotransposons. *BMC Bioinformatics*
9, 18 (2008). https://doi.org:10.1186/1471-2105-9-18

70 Gremme, G., Steinbiss, S. & Kurtz, S. GenomeTools: a comprehensive
software library for efficient processing of structured genome annotations.
IEEE/ACM Trans Comput Biol Bioinform **10**, 645-656 (2013).
https://doi.org:10.1109/TCBB.2013.68

71 Steinbiss, S., Willhoeft, U., Gremme, G. & Kurtz, S. Fine-grained annotation
and classification of de novo predicted LTR retrotransposons. *Nucleic Acids*
Res **37**, 7002-7013 (2009). https://doi.org:10.1093/nar/gkp759

72 Edgar, R. C. Search and clustering orders of magnitude faster than BLAST.
Bioinformatics **26**, 2460-2461 (2010).
https://doi.org:10.1093/bioinformatics/btq461

73 Smit, A. F. A. & Hubley, R. *RepeatMasker-4.0*, <www.repeatmasker.org>
(2013-2015).

74 Lomsadze, A., Ter-Hovhannisyan, V., Chernoff, Y. O. & Borodovsky, M. Gene
identification in novel eukaryotic genomes by self-training algorithm. *Nucleic*
Acids Res **33**, 6494-6506 (2005). https://doi.org:10.1093/nar/gki937

75 Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59 (2004).
https://doi.org:10.1186/1471-2105-5-59

76 Holt, C. & Yandell, M. MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. *BMC*
Bioinformatics **12**, 491 (2011). https://doi.org:10.1186/1471-2105-12-491

77 Stanke, M. & Waack, S. Gene prediction with a hidden Markov model and a
new intron submodel. *Bioinformatics* **19 Suppl 2**, ii215-225 (2003).

78 Seppey, M., Manni, M. & Zdobnov, E. M. BUSCO: Assessing Genome
Assembly and Annotation Completeness. *Methods Mol Biol* **1962**, 227-245
(2019). https://doi.org:10.1007/978-1-4939-9173-0_14

79 Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data
without a reference genome. *Nat Biotechnol* **29**, 644-652 (2011).
https://doi.org:10.1038/nbt.1883

80 Pracana, R., Priyam, A., Levantis, I., Nichols, R. A. & Wurm, Y. The fire ant
social chromosome supergene variant Sb shows low diversity but high

divergence from SB. *Mol Ecol* **26**, 2864-2879 (2017).
https://doi.org:10.1111/mec.14054

81   Götz, S. *et al.* High-throughput functional annotation and data mining with the
Blast2GO suite. *Nucleic Acids Res* **36**, 3420-3435 (2008).
https://doi.org:10.1093/nar/gkn176

82   Donath, A. *et al.* Improved annotation of protein-coding genes boundaries in
metazoan mitochondrial genomes. *Nucleic Acids Res* **47**, 10543-10552
(2019). https://doi.org:10.1093/nar/gkz833

83   Smith, T. F. & Waterman, M. S. Identification of common molecular
subsequences. *J Mol Biol* **147**, 195-197 (1981). https://doi.org:10.1016/0022-
2836(81)90087-5

84   Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast
Toolkit for FASTA/Q File Manipulation. *PLoS One* **11**, e0163962 (2016).
https://doi.org:10.1371/journal.pone.0163962

85   Patel, R. K. & Jain, M. NGS QC Toolkit: a toolkit for quality control of next
generation sequencing data. *PLoS One* **7**, e30619 (2012).
https://doi.org:10.1371/journal.pone.0030619

86   Jiang, H., Lei, R., Ding, S. W. & Zhu, S. Skewer: a fast and accurate adapter
trimmer for next-generation sequencing paired-end reads. *BMC*
Bioinformatics **15**, 182 (2014). https://doi.org:10.1186/1471-2105-15-182

87   Cingolani, P. *et al.* Using Drosophila melanogaster as a Model for Genotoxic
Chemical Mutational Studies with a New Program, SnpSift. *Front Genet* **3**, 35
(2012). https://doi.org:10.3389/fgene.2012.00035

88   Zarate, S. *et al.* Parliament2: Accurate structural variant calling at scale.
Gigascience **9** (2020). https://doi.org:10.1093/gigascience/giaa145

89   Chen, K. *et al.* BreakDancer: an algorithm for high-resolution mapping of
genomic structural variation. *Nat Methods* **6**, 677-681 (2009).
https://doi.org:10.1038/nmeth.1363

90   Abyzov, A., Urban, A. E., Snyder, M. & Gerstein, M. CNVnator: an approach
to discover, genotype, and characterize typical and atypical CNVs from family
and population genome sequencing. *Genome Res* **21**, 974-984 (2011).
https://doi.org:10.1101/gr.114876.110

91   Chen, X. *et al.* Manta: rapid detection of structural variants and indels for
germline and cancer sequencing applications. *Bioinformatics* **32**, 1220-1222
(2016). https://doi.org:10.1093/bioinformatics/btv710

92   Layer, R. M., Chiang, C., Quinlan, A. R. & Hall, I. M. LUMPY: a probabilistic
framework for structural variant discovery. *Genome Biol* **15**, R84 (2014).
https://doi.org:10.1186/gb-2014-15-6-r84

93   Rausch, T. *et al.* DELLY: structural variant discovery by integrated paired-end
and split-read analysis. *Bioinformatics* **28**, i333-i339 (2012).
https://doi.org:10.1093/bioinformatics/bts378

94   Brouard, J. S., Schenkel, F., Marete, A. & Bissonnette, N. The GATK joint
genotyping workflow is appropriate for calling variants in RNA-seq
experiments. *J Anim Sci Biotechnol* **10**, 44 (2019).
https://doi.org:10.1186/s40104-019-0359-0

95   Mansfeld, B. N. & Grumet, R. QTLseqr: An R Package for Bulk Segregant
Analysis with Next-Generation Sequencing. *Plant Genome* **11** (2018).
https://doi.org:10.3835/plantgenome2018.01.0006

96   Magwene, P. M., Willis, J. H. & Kelly, J. K. The Statistics of Bulk Segregant
Analysis Using Next Generation Sequencing. *PLOS Computational Biology* **7**,
e1002255 (2011). https://doi.org:10.1371/journal.pcbi.1002255

97   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A
Practical and Powerful Approach to Multiple Testing. *Journal of the Royal*
Statistical Society: Series B (Methodological) **57**, 289-300 (1995).
https://doi.org:https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

98   Zuo, J. F. *et al.* Effect of marker segregation distortion on high density linkage
map construction and QTL mapping in Soybean (Glycine max L.). *Heredity*
(Edinb) **123**, 579-592 (2019). https://doi.org:10.1038/s41437-019-0238-7

99   Lyttle, T. W. Segregation distorters. *Ann Rev Ecol Evol Syst.* **25**, 511-557
(1991).

100   Haley, C. S. & Knott, S. A. A simple regression method for mapping
quantitative trait loci in line crosses using flanking markers. *Heredity (Edinb)*
69, 315-324 (1992). https://doi.org:10.1038/hdy.1992.131

101   Churchill, G. A. & Doerge, R. W. Empirical threshold values for quantitative
trait mapping. *Genetics* **138**, 963-971 (1994).
https://doi.org:10.1093/genetics/138.3.963

102   Dupuis, J., Brown, P. O. & Siegmund, D. Statistical methods for linkage
analysis of complex traits from high-resolution maps of identity by descent.
Genetics **140**, 843-856 (1995). https://doi.org:10.1093/genetics/140.2.843

1179