A chromosome-level reference genome for the common octopus, *Octopus vulgaris* (Cuvier, 1797)

Dalila Destanović[1,*], Darrin T. Schultz[1,*,+], Ruth Styfhals[2,5], Fernando Cruz[6], Jèssica Gómez-Garrido[6], Marta Gut[6], Ivo Gut[6], Graziano Fiorito[5], Oleg Simakov[1,+], Tyler S. Alioto[6+], Giovanna Ponte[5,+], Eve Seuntjens[2,3,4,+]

Affiliations:

[1]Department of Neurosciences and Developmental Biology, University of Vienna, Djerassiplatz 1, 1030, Vienna, Austria

[2]Lab of Developmental Neurobiology, Animal Physiology and Neurobiology Division, Department of Biology, KU Leuven, Leuven, Belgium

[3]KU Leuven Institute for Single Cell Omics (LISCO), KU Leuven, Leuven, Belgium

[4]Leuven Brain Institute, KU Leuven, Leuven, Belgium

[5]Department of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121, Naples, Italy

[6]Centro Nacional de Análisis Genómico (CNAG), C/Baldiri Reixac 4, 08028 Barcelona, Spain

* - equal contribution
+ - corresponding authors

Short title: Chromosome-scale genome of *Octopus vulgaris*
Keywords: coleoid cephalopods, chromosome-scale, Hi-C

Abstract

Cephalopods are emerging animal models and include iconic species for studying the link between genomic innovations and physiological and behavioral complexities. Coleoid cephalopods possess the largest nervous system among invertebrates, both for cell counts and brain-to-body ratio. *Octopus vulgaris* has been at the center of a long-standing tradition of research into diverse aspects of cephalopod biology, including behavioral and neural plasticity, learning and memory recall, regeneration, and sophisticated cognition. However, no chromosome-scale genome assembly is available for *O. vulgaris* to aid in functional studies. To fill this gap, we sequenced and assembled a chromosome-scale genome of the common octopus, *O. vulgaris*. The final assembly spans 2.8 billion basepairs, 99.34% of which are in 30 chromosome-scale scaffolds. Hi-C heatmaps support a karyotype of 1n=30 chromosomes. Comparisons with other octopus species' genomes show a conserved octopus karyotype, and a pattern of local genome rearrangements between species. This new chromosome-scale genome of *O. vulgaris* will further facilitate research in all aspects of cephalopod biology, including various forms of plasticity and the neural machinery underlying sophisticated cognition, as well as an understanding of cephalopod evolution.

Introduction

Coleoid cephalopods (cuttlefish, squid, and octopus) comprise about 800 extant species characterized by highly diversified lifestyles, body plans, and adaptations. Cephalopod-specific traits, such as complex nervous systems (Young 1964; Hochner et al. 2006; Hochner 2012; Fiorito et al. 2014; Wang and Ragsdale 2019; Ponte et al. 2021), advanced learning abilities (reviewed in Marini et al. 2017), and the richness in body patterning considered to be involved in camouflaging and communication (Borrelli et al. 2006; Chiao and Hanlon 2019) have made this taxon ideal for studying evolutionary novelties. The neural plasticity of cephalopod brains and the existence of evidence for functionally analogous structures shared with mammalian brains have made cephalopods into a model comparative clade for neurophysiology research (Shigeno et al. 2018; Styfhals et al. 2022).

Despite the technical difficulties of sequencing their typically large and repetitive genomes, the available cephalopod genomes have given insights into the genomic basis for the evolution of novelty (Albertin et al. 2015, 2022; Jiang et al. 2022; Kim et al. 2018; Li et al. 2020; Marino et al. 2022; Schmidbaur et al. 2022). The first-published cephalopod genome, that of *Octopus bimaculoides* (Albertin et al. 2015), made it clear that cephalopod genomic novelties were not attributable to whole-genome duplication, as occurred in the vertebrate ancestor (Meyer and Schartl 1999; Dehal and Boore 2005). Comparisons of recently available chromosome-scale genome assemblies, including those of the Boston market squid *Doryteuthis pealeii* (Albertin et al. 2022) and the Hawaiian bobtail squid *Euprymna scolopes* (Schmidbauer et al. 2022), have shown the impact of genome reorganization on novel regulatory units in coleoid cephalopods. Still, it is not yet known how these units are made in terms of their gene content or their evolution in separate squid and octopus lineages. In this respect, it is crucial that the growing cephalopod genomics resources and approaches help obtain high-quality genomes for the established experimental species.

The common octopus, *Octopus vulgaris*, has long been used as a model for the study of learning and cognitive capabilities in invertebrates (reviewed in: Young 1964; Marini et al. 2017), and is also used as a comparative system in the study of neural organization and evolution (Shigeno et al. 2018; Ponte et al. 2022). Furthermore, recent advances in the culture of this species' early life stages have increased its suitability for molecular approaches and have provided important developmental staging information (Deryckere et al. 2020).

One bottleneck to studying *O. vulgaris* is the lack of a chromosome-scale genome assembly. While the reported karyotype of *O. vulgaris* is 1n=28 (Inaba 1959; Vitturi et al. 1982) or 1n=30 (Gao & Natsukari, 1990), to date there is no definitive answer. Existing genomic resources for *O. vulgaris* include a short read-based genome assembly (Zarrella et al. 2019), and a genome annotation based on the closely

related *O. sinensis* genome that is supported with PacBio Iso-Seq reads and FLAM-seq curation (Styfhals
et al. 2022; Zolotarov et al. 2022). These resources have been valuable in characterizing the molecular
and cellular diversity of the developing brain (Styfhals et al. 2022), the evolution of cephalopod brains
(Zolotarov et al. 2022), and the non-coding RNA repertoire unique to cephalopods (Petrosino et al. 2022).
Further improvements to the *O. vulgaris* genome assembly and genome annotation will provide a
valuable resource to the cephalopod and neuroscience communities.
Here we describe a chromosome-scale genome assembly and annotation of the common octopus,
O. vulgaris. We have validated our assembly using available chromosome-scale genomes of octopus
species (Li et al. 2020, Albertin et al. 2022; Jiang et al. 2022). Our analyses reveal large-scale
chromosomal homologies, yet a pattern of local rearrangement within chromosomes between species.
84
85

Materials and methods

Sample Collection

One adult male *Octopus vulgaris* (780 g body weight, specimen tube3-27.05.21-GP, BioSamples ERS14895525 and ERS14895526) was collected in the Gulf of Naples, Italy (40°48'04.1"N 14°12'32.7"E) by fishermen in May 2021. The animal was immediately sacrificed humanely following EU guidelines and protocols for collection of tissues from wild animals (Andrews et al. 2013; Fiorito et al. 2015) (see Data Availability for animal welfare information). The central brain masses (optic lobes, OL; supra-, SEM; sub-esophageal, SUB) were dissected out (ERS14895525), and the spermatophores (ERS14895526) were collected as described in Zarrella et al. (2019). All dissections were carried out on a bed of ice in seawater, and the excised tissues were then weighed and flash-frozen in liquid nitrogen.

High Molecular Weight Genomic DNA Extraction

High molecular weight genomic DNA (HMW gDNA) was extracted from frozen spermatophore sample (160 mg) (ERS14895526) using a salt-extraction protocol at the Stazione Zoologica Anton Dohrn (Italy) following Albertin et al. 2022. Briefly, two cryopreserved sample aliquots were each lysed for 3 hours at 55°C in separate tubes of 3 mL lysis buffer containing proteinase K. Then 1 mL of NaCl (5M) was added to each tube. The tubes were mixed by inversion then spun down for 15 minutes at 10,000 rcf. The supernatants were then transferred to a new tube and 2 volumes of cold ethanol (100%) was added. The DNA precipitate was then spooled, washed, resuspended in elution buffer (10 mM Tris, 0.1 mM EDTA, pH 8.5), and stored at 4°C. The DNA concentration was quantified using a Qubit DNA BR Assay kit (Thermo Fisher Scientific), and the purity was evaluated using Nanodrop 2000 (Thermo Fisher Scientific) UV/Vis measurements.

10X Genomics Library Preparation and Sequencing

A 10 ng aliquot of the spermatophore HMW DNA was used to prepare a 10X Genomics Chromium library (Weisenfeld et al. 2017) at the National Center for Genomic Analysis (Centre Nacional d'Anàlisi Genòmica - CNAG, Spain) using the Chromium Controller instrument (10X Genomics) and Genome Reagent Kits v2 (10X Genomics) following the manufacturer's protocol. The library was indexed with both P5 and P7 indexing adaptors. The resulting sequencing library was checked that the insert size matched the protocol specifications on an Agilent 2100 BioAnalyzer with the DNA 7500 assay (Agilent).

The library was sequenced at CNAG with an Illumina NovaSeq 6000 with a read length of 2x151bp, and was demultiplexed with dual indices (Supplementary Data 1).

 Long-read Whole Genome Library Preparation and Sequencing

The spermatophore HMW DNA was also used to prepare one Oxford Nanopore Technologies (ONT)
1D sequencing library (kit SQK-LSK110) at CNAG. Briefly, 2.0 μg of the HMW DNA was treated with
the NEBNext FFPE DNA Repair Mix (NEB) and the NEBNext UltraII End Repair/dA-Tailing Module
(NEB). ONT sequencing adaptors were then ligated to the DNA, then the DNA was purified with 0.4X
AMPure XP Beads and eluted in Elution Buffer.

Two sequencing runs were performed at CNAG on an ONT PromethIon 24 using ONT R9.4.1 FLO-
PRO 002 flow cells. The libraries were sequenced for 110 hours. The quality parameters of the
sequencing runs were monitored by the MinKNOW platform version 21.05.8 (Oxford Nanopore
Technologies)   and   base   called   with   Guppy,   version   5.0.11   (available   through
https://community.nanoporetech.com) (Supplementary Data 1).

Omni-C Library Preparation and Sequencing

A DoveTail Genomics Omni-C library was prepared at SciLifeLab (Solna, Sweden) using the flash-
frozen brain tissue from the same individual used to generate the ONT long reads and 10X Genomics
chromium reads (ERS14895525). One hundred milligrams of brain tissue were pulverized to a fine
powder using a mortar and pestle under liquid nitrogen. Two 20 mg aliquots of the pulverized tissue were
fixed in PBS with formaldehyde and disuccinimidyl glutarate (DSG), and were prepared according to the
manufacturer's protocol as two separate libraries. To increase the final complexity, the two libraries
bound to streptavidin beads were pooled together into a single tube prior to P7 indexing PCR. The
amplified library was sequenced at SciLifeLab on an Illumina NovaSeq 6000 with a read length of 2x150
bp, and was demultiplexed with one index (Supplementary Data 1).

Nuclear Genome Assembly

Sequencing produced 77Gb of ONT WGS reads (27.5x coverage) and 230.25 Gb of 10X Genomics
linked reads (77.7x coverage). These data were assembled with the CNAG Snakemake assembly pipeline
v1.0 (https://github.com/cnag-aat/assembly_pipeline) to obtain an optimal base assembly for further Hi-C
scaffolding. In brief, this pipeline first preprocessed the 10X reads with *LongRanger basic* v2.2.2
(https://github.com/10XGenomics/longranger) and filtered the ONT reads with *FiltLong* v0.2.0
(https://github.com/rrwick/Filtlong), and then the ONT reads were assembled with both *Flye* v2.9
(Kolmogorov et al. 2019) and *NextDenovo* v2.4.0 (Hu et al. 2023). The following evaluations were run on
both assemblies and after each subsequent step of the pipeline: BUSCO v5.2.2 (Manni et al. 2021) with
metazoan_odb10 and *Merqury* v1.1 (Rhie et al. 2020) to estimate the consensus accuracy (QV) and k-mer
statistics, and *fasta-stats.py* for contiguity statistics. The best contig assembly was obtained with
NextDenovo (see assembly metrics Supplementary Data 2), so the remaining steps of the pipeline were
run on this assembly (Supplementary Figure 1 and Supplementary Data 2).

The assembly was polished with 10X Illumina and ONT reads using *Hypo* v1.0.3 (Kundu et al.
2019); collapsed with *purge_dups* v1.2.5 (Guan et al. 2020); then scaffolded with the 10X chromium
reads using *Tigmint* v1.2.4 (Jackman et al. 2018), *ARKS* v1.2.2 (Coombe et al. 2018) and *LINKS* v1.8.6
(Warren   et   al.   2015)   following   the   Faircloth's   Lab   protocol   (http://protocols.faircloth-
lab.org/en/latest/protocols-computer/assembly/assembly-scaffolding-with-arks-and-links.html).   The
specific parameters and versions used to assemble the *O. vulgaris* specimen are listed in Supplementary

Data 3. Finally, 310 scaffolds shorter than 1 Kb were removed from the assembly. This assembly was
used for scaffolding with Omni-C data.
164
Omni-C Scaffolding
165
The Omni-C reads (863.85 million read pairs) were then mapped to the assembly (Supplementary
Data 4) using the recommended procedure from Dovetail Genomics (https://omni-
c.readthedocs.io/en/latest/fastq_to_bam.html). In short, the reads were mapped to the reference using *bwa*
mem v0.7.17-r1188 (Li 2013) with flags *-5SP -T0*, converted to a sorted *.bam* file, and filtered to reads
with a minimum mapping quality of 30 with *samtools* v1.9 (Li et al. 2009) with *htslib* v1.9, and filtered to
keep uniquely mapping pairs with *pairtools* v0.3.0 (Open2C et al. 2023). The minimum mapping quality
threshold of 30 was used to accommodate for the organism's heterozygosity and repetitiveness (1.22%
and 68.68%, respectively. see supplementary table Supplementary Data 5). After excluding PCR
duplicates and improperly mated reads with *pairtools*, 231.59 million Hi-C read pairs were used to
scaffold the assembly with *YaHS* v1.1 (Zhou et al. 2023) in the default mode, thus initially detects and
corrects errors in contigs, introducing breaks at misjoins.
177
Generation of the Hi-C Heatmaps and Manual Curation
178
179
We then manually curated the scaffolded assembly using an editable Hi-C heatmap to improve the
assembly's quality and to correct misassemblies. The process described below was repeated for five
rounds until there were no obvious improvements to make based on the Hi-C heatmap signal.
Chromap v0.2.3 (Zhang et al. 2021) was used to align the Omni-C reads to the genome with a read
alignment quality cutoff of Q0. The resulting *.pairs* file (quality cutoffs: 2,10) was converted using *awk* v
4.2.1(Aho et al. 1988) to a *.longp* file, a format used by *Juicebox Assembly Tools* (Dudchenko et al.
2018). We ran the script *run-assembly-visualizer.sh* from the *3D-DNA* pipeline (Dudchenko et al. 2017)
on the *.longp* file to generate a *.hic* file. The *generate-assembly-file-from-fasta.awk* script from the *3D-*
DNA pipeline (Dudchenko et al. 2017), and the *assembly-from-fasta.py* from the *Artisanal* pipeline
(Bredeson et al. 2022) were used to generate the *.assembly* files necessary to curate the *.hic* heatmap file
in *Juicebox Assembly Tools* (Dudchenko et al. 2018).
The resulting *.hic* heatmap file was visualized using the visualization tool *Juicebox* v1.11.08 (Durand
et al. 2016). Using the signal in the Hi-C heatmap we corrected the order and orientation of contigs within
the chromosome-scale scaffolds, and placed small contigs and scaffolds onto the chromosome-scale
scaffolds. A new *.fasta* assembly was generated from the corrected *.assembly* file by using the *assembly-*
to-fasta.py script from the *Artisanal* pipeline.
The corrected assembly was aligned to the chromosome-scale *O. sinensis* (GCA_006345805.1) (Li et
al. 2020), *O. bimaculoides* (GCA_001194135.2) (Albertin et al. 2022), and *A. fangsiao* (Jiang et al. 2022)
genomes using *minimap2* v2.24 (Li, 2018), *snakemake* v7.19.1-3.11.1 (Köster and Rahmann 2012) and
the *snakemake* script *GAP_dgenies_prep* (https://doi.org/10.5281/zenodo.7826771). The resulting *.paf*
file was visualized with *D-GENIES* v1.4.0 (Cabanettes and Klopp 2018). Regions of the *O. vulgaris*
chromosome-scale scaffolds that had ambiguous Hi-C heatmap signal, or regions that had no obvious
homology to other *Octopus* spp. chromosome-scale scaffolds were removed from the chromosome-scale
scaffolds and retained as smaller scaffolds at the end of the genome assembly *.fasta* file. Scaffolds were
renamed based on homology with *O. bimaculoides* chromosomes.
205

 Decontamination

After curation, we ran the *BlobToolKit* INSDC pipeline (Challis et al. 2020), using the NCBI *nt*
database (updated on December 2022) and the following BUSCO *odb10* databases: eukaryota, fungi,
bacteria, metazoa and mollusca. This analysis identified 226 scaffolds either matching the phylum
Mollusca or having no-hit in the database (Supplementary Figure 2). A total of 47 small scaffolds
matching other phyla (Supplementary Data 6 and Supplementary Figure 3) were considered contaminants
and removed from the assembly. This scaffolded and decontaminated assembly was then carried forward
for annotation and comparative analyses, and is available at https://denovo.cnag.cat/octopus and the
INSDC (ENA, NCBI, and DDBJ) accession number GCA_951406725.1.

Nuclear Genome Annotation

The gene annotation of the octopus genome assembly was obtained by combining transcript
alignments, protein alignments, and *ab initio* gene predictions as described below. A flowchart of the
annotation process is shown in Supplementary Figure 4.
Repeats present in the genome assembly were annotated with *RepeatMasker* v4-1-2 (Smit et al. 2013-
2015) using the custom repeat library available for Mollusca. Moreover, a new repeat library specific to
the assembly was made with *RepeatModeler* v1.0.11. After excluding repeats from the resulting library
that were part of repetitive protein families by performing a *BLAST* (Altschul et al. 1990) search against
Uniprot, *RepeatMasker* was rerun with this new library to annotate species-specific repeats.
PacBio Iso-Seq reads from several developmental stages were downloaded from NCBI
(PRJNA718058, PRJNA791920, PRJNA547720) (García-Fernández et al. 2019; Deryckere et al. 2021;
Zolotarov et al. 2022). Bulk RNA-seq from an adult octopus (Petrosino et al. 2022) was downloaded from
the *ArrayExpress* database under accession number E-MTAB-3957. The short and long reads were
aligned to the genome using *STAR* v-2.7.2a (Dobin et al. 2013) and *minimap2* v2.14 (Li, 2018) with the
option *-x splice:hq*. Transcript models were subsequently generated using *Stringtie* v2.1.4 (Pertea et al.
2015) on each *.bam* file, and then all the transcript models were combined using *TACO* v0.6.3 (Niknafs
et al. 2017). High-quality junctions to be used during the annotation process were obtained by running
Portcullis v1.2.0 (Mapleson et al. 2018) after mapping with *STAR* and *minimap2*. Finally, *PASA*
assemblies were produced with *PASA* v2.4.1 (Haas et al. 2008). The *TransDecoder* program, part of
the *PASA* package, was run on the *PASA* assemblies to detect coding regions in the transcripts.
The complete proteomes of *O. vulgaris*, *O. bimaculoides,* and *Sepia pharaonis* were downloaded
from *UniProt* in October 2022 and aligned to the genome using *Spaln* v2.4.03 (Iwata and Gotoh 2012).
Ab initio gene predictions were performed on the repeat-masked assembly with two different
programs: *Augustus* v3.3.4 (Stanke et al. 2006) and *Genemark-ES* v2.3e (Lomsadze et al. 2014) with and
without incorporating evidence from the RNA-seq data. Before gene prediction, *Augustus* was trained
with octopus-specific evidence. The gene candidates used as evidence for training *Augustus* were
obtained after selecting *Transdecoder* annotations that were considered complete and did not overlap
repeats, clustering them into genes, and selecting only one isoform per gene. These candidates were
aligned to the *Swissprot* NCBI database with *blastp* v2.7.1 (Altschul et al. 1990) to select only those with
homology to proteins. The final list of candidate genes was made of 1764 genes with *BLAST* hits to
known proteins with e-values smaller than 10^{-9} and greater than 55% identity.
Finally, all the data were combined into consensus CDS models using *EvidenceModeler* v1.1.1
(EVM) (Haas et al. 2008). Additionally, UTRs and alternative splicing forms were annotated via two
rounds of *PASA* annotation updates. Functional annotation was performed on the annotated proteins with

Blast2go v1.3.3 (Conesa et al. 2005). First, a *DIAMOND* v2.0.9 *blastp* (Buchfink et al. 2021) search was
made against the *nr* database. Furthermore, *Interproscan* v5.21-60.0 (Jones et al. 2014) was run to detect
protein domains on the annotated proteins. All these data were combined by *Blast2go* v1.3.3, which
produced the final functional annotation results.
Identification of long non-coding RNAs (lncRNAs) was done by first filtering the set of *PASA-*
assemblies that had not been included in the annotation of protein-coding genes to retain those longer than
200bp and not covered more than 80% by repeats. The resulting transcripts were clustered into genes
using shared splice sites or significant sequence overlap as criteria for designation as the same gene.
257
Nuclear Genome and Annotation Completeness Assessment
The final *O. vulgaris* genome assembly, the annotated transcripts, the proteins from the annotated
transcripts, and the other available octopus genomes were assessed for completeness using BUSCO
databases as described above (Materials and Methods - Genome Assembly). To compare the qualities of
each assembly, we used *fasta_stats* (Chapman et al. 2011) shown in (Table 1). We calculated the
percentage of bases in the chromosome-scale scaffolds (Table 1) with *bioawk* v1.0
(https://github.com/lh3/bioawk).

 Mitogenome Assembly and Annotation

To assemble the mitochondrial genome we employed a strategy that uses a reference bait to select the
mitochondrial nanopore reads, assembles those reads into a single circular contig, and then performs two
rounds of polishing. To obtain the mitochondrial sequences, all ONT reads with a mean quality of ≥10
were mapped with *minimap2* v2.24 (Li 2018) against the circular complete, 15,744 bp mitochondrial
genome of another specimen of *O. vulgaris* (NC_006353.1) (Yokobori 2004) with the minimap2
parameter *-ax map-ont*. We retained all reads with a mapping quality ≥ 13. Approximately 5,000 ONT
reads passed these filters including 15 reads accounting for 181,644 total basepairs (12x coverage) with a
mean length of 12,112 bp.

All the retained ONT reads were assembled with *Flye* v2.9 (Kolmogorov et al. 2019) using the
options *flye --scaffold -i 2 -g 15744 --nano-raw --min-overlap 7000*. This produced one circular contig.
The *-i 2* option specified for flye caused this contig to be polished twice with the input ONT reads. After
polishing the length of the circular contig was 15,651 bp, and a web blastn search revealed that it spanned
the length of the NC_006353.1 mitochondrial genome. The circular mitogenome contig was rotated and
oriented as follows. First, we annotated the contig using *MITOS* v2.1.3 (Bernt et al. 2013) with
parameters *-c 5 --linear --best -r refseq81m*. Second, we use the coordinates in the *results.bed* file to
orient the mitogenome, so it starts with the conventional tRNA Phenyl-Alanine (trnF) (Formenti et al.
2021).

To evaluate the assembly accuracy, we first aligned the selected ONT reads back to the assembly with
minimap2 and visually inspected the alignment with *IGV* v2.14.1 (Robinson et al. 2023). Finally, the
xcOctVulg1 mitogenome was aligned against the mitogenome of other species using *DNAdiff* v1.3 from
mummer package v3.23 (Kurtz et al. 2004). These species included the mitogenomes of another specimen
of *O. vulgaris* (NC_006353.1), *O. sinensis* (NC_052881.1), *O. bimaculoides* (NC_029723.1), and *A.*
fangsiao (AB240156.1). From these pairwise alignments, we calculated the percent identity.

289 Results and Discussion

DNA Sequencing

Sequencing the ONT WGS library yielded 8.3 million ONT PromethIon reads containing 82.57
billion base pairs (Gbp) with 29.47x coverage per library. Sequencing of the 10X Genomics Chromium
library yielded 762 million read pairs containing 228.69 Gbp with 81.64x coverage per library. The
Omni-C library sequencing yielded 863.85 million read pairs, containing 259.16 Gbp of data with 33.02X
coverage. Details about sequence data can be found in Supplementary Data 1.

Manual Curation and Decontamination of the Assembly

Manually curating the genome assembly improved the quality of the final assembly, as 495 scaffolds
were placed in the chromosome-scale scaffolds, and 47 additional scaffolds were removed through the
contamination analysis (Table 1). The final 2.80 Gb assembly, xcOctVulg1.1, has a scaffold N50 of 118.9
Mb, an N90 of 18.2 Mb, QV39 and gene completeness estimated using BUSCO v5.3.2 with
mollusca_odb10 of C:86.5% [S:85.8%, D:0.7%], F:3.4%, M:10.1%, n:5295 (Fig. 1C). The BUSCO score
with *metazoa_odb10* for the final assembly is C:92.3% [S:91.8%, D:0.5%], F:2.7%, M:5.0% (Table 2).
The statistics for all intermediate assemblies are shown in Supplementary Data 2. Also, in Supplementary
Figure 3 we show that the final assembly has been properly decontaminated.

The Octopus *Karyotype*

The genome assembly from this study contains 30 large scaffolds with Hi-C heatmap signal (Fig. 1D) that is consistent with each scaffold representing a single chromosome and resembles the Hi-C heatmaps of other chromosome-scale octopus genome assemblies (Li et al. 2020; Albertin et al. 2022; Jiang et al. 2022). The first reported *O. vulgaris* karyotypes from Japan and Italy were 1n=28 chromosomes (Inaba 1959; Vitturi et al. 1982), but later studies also using *O. vulgaris* individuals sampled in Japan reported at 1n=30 (Gao and Natsukari 1990). The karyotype 1n=30 has been reported in four other octopus species: *Callistoctopus minor, Amphioctopus fangsiao, Cistopus sinensis*, and *Amphioctopus areolatus* (Gao and Natsukari 1990; Adachi et al. 2014; Wang and Zheng 2017). The only exception is *Hapalochlaena maculosa* which does not have a confirmed karyotype, but 47 linkage groups were suggested for this species (Whitelaw et al. 2022).

In light of the recent taxonomic designation of a new species *O. sinensis* (East Asian Common Octopus) from the previously synonymous *O. vulgaris* (Gleadall 2016; Amor et al. 2017, 2019; Amor 2023), this suggests that the reported *O. vulgaris* karyotypes probably belong to *O. sinensis*. Dot plot analyses, described below, show that *O. vulgaris* and *O. sinensis* share 30 homologous, largely collinear, chromosomes (Fig. 2).

The final version of the *O. vulgaris* genome was aligned to the genomes of three octopus species, *O. sinensis*, *O. bimaculoides*, and *A. fangsiao* (Fig. 2). *O. vulgaris* and *O. sinensis* have a less diverged genome sequence and few inversions between homologous, collinear chromosomes. General chromosomal collinearity was also observed in comparison to *O. bimaculoides* (Fig. 2). We have found large-scale inversions (megabase-scaled, larger than 1Mb) throughout the genomes of two species. The overall sequence similarity is lower compared to the previous pair, and a greater number of chromosomal rearrangements are present, confirming that they are more diverged. This is expected considering that *O. bimaculoides* and the *O. vulgaris-O. sinensis clade* diverged around 34 million years ago (mya) (Jiang et al. 2022), while *O. sinensis* and *O. vulgaris* diverged just 2.5 mya (Amor et al. 2019). In Figure 2, the collinearity between *O. vulgaris* and *A. fangsiao* chromosomes is visible. As expected, as *A. fangsiao* is the most distant to *O. vulgaris* of the compared species, the genomes are even more rearranged.

Our whole-genome alignment analyses support the hypothesis that *O. vulgaris, O. sinensis*, *O. bimaculoides,* and *A. fangsiao* share 30 homologous chromosomes (Fig. 2). Given the divergence time of these species, these results suggest that the karyotype of the common ancestor of this clade, and perhaps the common ancestor of octopuses, also had 30 chromosomes that still exist in extant species.

Karyotype stability was described in the squid lineage (Decapodiformes) on loliginid and sepiolid squids (Albertin et al. 2022). This study has suggested that the smaller karyotype found in octopuses (1n=30) compared to squids (1n=46) results from secondary fusions of a more ancestral squid chromosomal complement. Recently, it has been suggested that chromosomal fusions impact recombination, as well as chromosomal nuclear occupancy, in mice (Vara et al. 2021). Therefore, chromosomal fusions in the common ancestor of the octopus lineage might be one of the drivers of diversification, as this changes the chromosomal interactions and is hypothesized to lead to the formation of novel regulatory units (Vara et al. 2021). Such events are important in light of understanding the emergence of octopus-specific traits. We infer from the genome-genome comparisons that a similar pattern of intrachromosomal rearrangements with the conservation of individual chromosomes is seen in octopus species, as described in squids (Albertin et al. 2022). However, the loliginids and sepiolids are estimated to have diverged 100 mya (Albertin et al. 2022), while the genera *Octopus* and *Amphioctopus* are estimated to have diverged 44 mya (Jiang et al. 2022). Therefore, a more-distant species chromosome-

scale genome is needed to claim karyotype stasis in Octopodiformes. Nevertheless, future comparative
studies of the genomes of these closely-related species will shed light on the evolutionary history of
octopuses as a separate lineage of coleoid cephalopods. In addition to this, *O. vulgaris* is a model animal
in neurobiological studies, and having a high-quality genome will facilitate further studies of the
cephalopod brain.

Nuclear Genome Annotation

In total, we annotated 23,423 protein-coding genes that produce 31,799 transcripts (1.36 transcripts
per gene) and encode 30,121 unique protein products. We were able to assign functional labels to 53.5%
of the annotated proteins. The annotated transcripts contain 8.42 exons on average, with 87% of them
being multi-exonic (Table 3). In addition, 1,849 long non-coding transcripts have been annotated. The
number of protein-coding genes annotated here is slightly lower than those reported for other octopus
genome assemblies, like *O. sinensis* (Li et al. 2020). After checking the general statistics of both
annotations (Table 3), we can observe that the genes annotated here tend to be longer (both in the number
of exons and global length). After comparing both methods, the main difference that we believe is
responsible for this difference in length is the source of the transcriptomic data, the inclusion of long-read
Iso-seq data in the annotation process is known to result in less fragmented and longer annotations.

Nuclear Genome and Annotation Completeness Assessment

The BUSCO score was calculated for the *O. vulgaris*, *O. bimaculoides*, *O. sinensis*, and *A. fangsiao*
genomes. For the chromosome-scale *O. vulgaris* genome, the BUSCO score for a whole-genome
nucleotide sequence using the metazoan reference dataset was 92.3% for complete genes (954 core
genes). The full score can be seen in Table 2. This is an improvement considering the BUSCO score of
the previous *O. vulgaris* genome assembly (GCA_003957725.1) for complete genes was 63.1% (Zarrella
et al. 2019). Additionally, we assessed the completeness of the annotated proteome and transcriptome by
calculating the BUSCO score against the *metazoa_odb10* and *mollusca_odb10* databases (Supplementary
Data 2).

Mitogenome Assembly and Annotation

The mitogenome assembly of the *O. vulgaris* specimen (xcOctVulg1) has a length of 15,651 bp and
contains 13 protein-coding, 23 ncRNA, 2 rRNA, and 21 tRNA genes. The ONT read alignment to the
mitogenome shows a high consensus support for each nucleotide except for 16 positions (Supplementary
Figure 5). These 16 positions are single nucleotide polymorphisms, not indels, and the base at each
position is the base with the highest coverage in the reads at that position (Supplementary Figure 6).
Therefore, the mitochondrial genome has a high per-base accuracy.

The percentages of identity (See Supplementary Data 7) between the *O. vulgaris* and other octopus
mitochondrial genome sequences are consistent with the phylogeny topology (Fig. 2, Supplementary Data
7), and previous research on octopus taxonomy. The mitochondrial genome of the specimen collected in
Japan and identified as *O. vulgaris* (NC_006353.1) shows a higher identity to *O. sinensis* (99.85%) than
to our *O. vulgaris* specimen (96.79%). The 3.21% difference between the mitogenomes of the specimen
from this study and NC_006353.1 is close to the estimated divergence rate (~2% divergence/million years
(Arbogast and Slowinski 1998)) for *O. vulgaris* and *O. sinensis* (estimated time of divergence: 2.5mya
(Amor et al. 2019). These results suggest that the specimen collected in Japan and identified as *O.*
vulgaris (NC_006353.1) is more likely to be *O. sinensis*. This possibility is consistent with recent

morphological, molecular, and geographic delimitations made between the *O. sinensis* and *O. vulgaris*
species complex (Gleadall 2016; Amor et al. 2017, 2019; Amor 2023).
397

Conclusion

Octopus vulgaris is an important emerging model in comparative neuroscience, cognition research,
and evolutionary studies of cephalopods. The chromosome-scale genome assembly and annotation
reported here provide an improved reference for single-cell multiomics and the study of non-coding
regions and gene regulatory networks, that require the context of chromosome-scale sequences. This
assembly and annotation will also facilitate many avenues of cephalopod research, in particular analyses
of genome evolutionary trends in octopus and cephalopods within invertebrates. Furthermore, the
chromosome-scale *O. vulgaris* genome assembly will allow the estimation of chromosome rearrangement
rates, the emergence of novel coding and non-coding genes among octopuses, and the turnover rates of
putative regulatory regions. The scientific interest in *O. vulgaris* as a model animal in many fields
including (evolutionary) developmental biology and neuroscience will be facilitated by the availability of
a high-quality genome.

These efforts may help bridge the traditional *O. vulgaris* research on neurobiology, behavior, and
development to the molecular determinants involved in these fields.

413
414
415
416

Main Manuscript Figures

419 Dataset: xcOctVulg1.1

Figure 1 | ***Octopus vulgaris*** **assembly statistics and quality control.** (A) A specimen of *O. vulgaris*. (B) A cladogram showing the phylogenetic relationship between the compared species and the family Argonautidae as an outgroup (Taite et al. 2023). Chromosome-scale genome assemblies are available for the starred species (*). (C) The snail plot shows that the final version of the chromosome-scale *O. vulgaris* assembly has N50 of 119Mb, the longest scaffold is 225Mb long, and a BUSCO score for complete genes of 86.6% against the *mollusca_odb10* database. (D) The Hi-C heatmap of the final genome assembly shows 30 chromosome-scale scaffolds with very few sequences in unplaced scaffolds. Photography credit © Antonio, Valerio Cirillo (BEOM SZN), 2023 (A).

428
429
Figure 2 | Comparative analyses of available chromosome-scale Octopodidae genomes. The figure shows the inferred
phylogenetic relationship (Amor et al. 2017; Jiang et al. 2022; Taite et al. 2023) and the inferred divergence times (Amor et al.
2019; Jiang et al. 2022) of four octopus species. The diagrams show genome-genome alignments for each species compared to
O. vulgaris.

Main Manuscript Tables

Table 1 | Octopus genome assembly statistics.

Assembly	Number of scaffolds	Number of contigs	Scaffold sequence total	Scaffold N50/L50	Number of scaffolds > 50 KB	% of the bases in chromosome-scaled scaffolds
Final chromosome-scale *O. vulgaris* genome	226	2758	2800.4 MB	118.9 MB/9	57	99.34
Pre-curation scaffolded assembly *O. vulgaris*	768	2776	2801.6 MB	118.3 MB/9	296	95.82
Chromosome-scale *O. bimaculoides* (Albertin et al. 2022)	145326	713915	2342.5 MB	96.9 MB/9	85	95.46
Chromosome-scale *O. sinensis* (Li et al., 2020)	13516	20491	2719.2 MB	105.9 MB/10	1800	86.09
Chromosome-scale *A. fangsiao* (Jiang et al. 2022)	6409	9099	4341.1 MB	169.7 MB/10	1769	93.05

Table 2 | Octopus *metazoa_odb10* BUSCO scores.

Genome	Complete BUSCO	Single BUSCO	Duplicated BUSCO	Fragmented BUSCO	Missing BUSCO
Chromosome-scale *O. vulgaris*	92.3% [881]	91.8% [876]	0.5% [5]	2.7% [25]	5.0% [48]
Contig-level *O. vulgaris* (Zarrella et al. 2019)	63.1% [602]	62.6% [597]	0.5% [5]	24.8% [237]	12.1% [115]
Chromosome-scale *O. bimaculoides* (Albertin et al. 2022)	94.6% [903]	94.2% [899]	0.4% [4]	3.2% [31]	2.2% [20]
Chromosome-scale *O. sinensis* (Li et al., 2020)	95.7% [913]	90.5% [863]	5.2% [50]	2.6% [25]	1.7% [16]
Chromosome-scale *A. fangsiao* (Jiang et al. 2022)	93.5% [892]	91.6% [874]	1.9% [18]	3.5% [33]	3.0% [29]

439
440

441 **Table 3 | Genome annotation statistics.**

	OctVul6B annotation	Osinensis ASM634580v1 (Li et al. 2020)
Number of protein-coding genes	23,423	31,676
Median gene length (bp)	20,288	4,403
Number of transcripts	31,799	31,676
Number of exons	168,570	184,658
Number of coding exons	161,430	180,943
Median UTR length (bp)	1,255	441
Median intron length (bp)	2,467	1,520
Exons/transcript	8.42	5.83
Transcripts/gene	1.36	1
Multi-exonic transcripts	87%	81%
Gene density (gene/Mb)	8.36	11.7

Data Availability Statement

The data are available at https://denovo.cnag.cat/octopus. On the INDSC databases (ENA, NCBI, DDBJ) the genome is available at accession GCA_951406725.1, and the data in BioProject PRJEB61268. Euthanizing cephalopods solely for tissue removal does not require authorization from the National Competent Authority under Directive 2010/63/EU and its transposition into National Legislation. Samples were taken from local fishermen, and humane killing followed principles detailed in Annex IV of Directive 2010/63/EU as described in the Guidelines on the Care and Welfare of Cephalopods (Fiorito et al. 2015). The sampling of octopuses from the wild included in this study was authorized by the Animal Welfare Body of Stazione Zoologica Anton Dohrn (Ethical Clearance: case 06/2020/ec AWB-SZN). Genomes of *O. sinensis* (GCA_006345805.1) (Li et al., 2020) and *O. bimaculoides* (GCA_001194135.2) (Albertin et al. 2022) were downloaded from NCBI, while the *A. fangsiao* genome (Jiang et al. 2022) was downloaded from Figshare (https://figshare.com/s/fa09f5dadcd966f020f3).

Acknowledgments

The authors acknowledge that some of the computational results of this work have been achieved using the Life Science Compute Cluster (LiSC) of the University of Vienna. We also acknowledge support from the National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council, and NAISS/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure.

Conflict of Interest

D.T.S. is a shareholder of Pacific Biosciences of California, Inc. All other authors declare no competing interests.

Author Contributions

Dalila Destanovic = D.D.
Darrin T. Schultz = D.T.S.
Oleg Simakov = O.S.
Eve Seuntjens = E.S.
Ruth Styfhals = R.S.
Giovanna Ponte = G.P.
Tyler S. Alioto = T.S.A.
Fernando Cruz = F.C.
Jessica Gomez-Garrido = J.G.G.
Ivo Gut = I.G.
Marta Gut = M.G.

E.S., O.S. and G.P. conceived of the study design. D.D., D.T.S., O.S., T.S.A., M.G., and F.C. wrote the first draft of the manuscript. G.P. collected and dissected the octopus individual sequenced for this study. T.S.A., F.C., D.D., and D.T.S. assembled the genome. J.G.G. annotated the genome. D.D., J.G.G.,

482 and F.C. performed genomic analyses. D.D., D.T.S., F.C., J.G.G. and created the figures. All authors
483 contributed to the interpretation, presentation, and writing of the manuscript.

Funding

485   EASI-Genomics - This project has received funding from the European Union's Horizon 2020:
486 European Union Research and Innovation Programme (ERC-H2020-EURIP) under grant agreement No
487 824110. D.T.S., D.D., and O.S. were supported by ERC-H2020-EURIP grant to O.S., grant number
488 945026. E.S. and R.S. were supported by grants IDN/20/007 and C14/21/065 from KU Leuven.

Literature Cited:

Adachi K, Ohnishi K, Kuramochi T, Yoshinaga T, Okumura S-I. 2014. Molecular cytogenetic study in
Octopus (*Amphioctopus*) *areolatus* from Japan. Fish Sci. 80(3):445–450. doi:10.1007/s12562-
014-0703-4.

Aho AV, Kernighan BW, Weinberger PJ. 1988. The AWK programming language. Reading, Mass:
Addison-Wesley Pub. Co.

[GCF_001194135.2] Albertin CB, Medina-Ruiz S, Mitros T, Schmidbaur H, Sanchez G, Wang ZY,
Grimwood J, Rosenthal JJC, Ragsdale CW, Simakov O, et al. 2022. Genome and transcriptome
mechanisms driving cephalopod evolution. Nat Commun. 13(1):2427. doi:10.1038/s41467-022-
29748-w.

Albertin CB, Simakov O, Mitros T, Wang ZY, Pungor JR, Edsinger-Gonzales E, Brenner S, Ragsdale
CW, Rokhsar DS. 2015. The octopus genome and the evolution of cephalopod neural and
morphological novelties. Nature. 524(7564):220–224. doi:10.1038/nature14668.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J Mol
Biol. 215(3):403–410. doi:10.1016/S0022-2836(05)80360-2.

Amor MD, Doyle SR, Norman MD, Roura A, Hall NE, Robinson AJ, Leite TS, Strugnell JM. 2019.
Genome-wide sequencing uncovers cryptic diversity and mito-nuclear discordance in the *Octopus*
vulgaris species complex. bioRxiv 573493. [accessed 2023 Feb 5]. doi:10.1101/573493.

Amor MD, Norman MD, Roura A, Leite TS, Gleadall IG, Reid A, Perales-Raya C, Lu C, Silvey CJ,
Vidal EAG, et al. 2017. Morphological assessment of the *Octopus vulgaris* species complex
evaluated in light of molecular-based phylogenetic inferences. Zool Scr. 46(3):275–288.
doi:10.1111/zsc.12207.

Amor, MD. 2023. Untangling the *Octopus vulgaris* species complex using a combined genomic and
morphological approach [dissertation]. Melbourne: La Trobe University.
doi:10.26181/21854307.V1.

Andrews PLR, Darmaillacq A-S, Dennison N, Gleadall IG, Hawkins P, Messenger JB, Osorio D, Smith
VJ, Smith JA. 2013. The identification and management of pain, suffering and distress in
cephalopods, including anaesthesia, analgesia and humane killing. J Exp Mar Biol Ecol. 447:46–
64. doi:10.1016/j.jembe.2013.02.010.

Arbogast BS, Slowinski JB. 1998. Pleistocene speciation and the mitochondrial DNA clock. Science.
282(5396):1955-1955. doi:10.1126/science.282.5396.1955a

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, Stadler PF.
2013. MITOS: Improved de novo metazoan mitochondrial genome annotation. Mol Phylogenet
Evol. 69(2):313–319. doi:10.1016/j.ympev.2012.08.023.

Borrelli L, Gherardi F, Fiorito G. 2006. A Catalogue of Body Patterning in Cephalopoda. 1st ed. Firenze:
Firenze University Press (Cataloghi e collezioni). doi:10.36253/88-8453-376-7.

Bredeson JV, Lyons JB, Oniyinde IO, Okereke NR, Kolade O, Nnabue I, Nwadili CO, Hřibová E, Parker
M, Nwogha J, et al. 2022. Chromosome evolution and the genetic basis of agronomically
important traits in greater yam. Nat Commun. 13(1):2001. doi:10.1038/s41467-022-29114-w.

Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using
DIAMOND. Nat Methods. 18(4):366–368. doi:10.1038/s41592-021-01101-x.

Cabanettes F, Klopp C. 2018. D-GENIES: dot plot large genomes in an interactive, efficient and simple
way. PeerJ. 6:e4958. doi:10.7717/peerj.4958.

Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. 2020. BlobToolKit – Interactive Quality
Assessment of Genome Assemblies. G3 GenesGenomesGenetics. 10(4):1361–1374.
doi:10.1534/g3.119.400908.

Chapman JA, Ho I, Sunkara S, Luo S, Schroth GP, Rokhsar DS. 2011. Meraculous: De Novo Genome
Assembly with Short Paired-End Reads. Salzberg SL, editor. PLoS ONE. 6(8):e23501.
doi:10.1371/journal.pone.0023501.

Chiao C-C, Hanlon RT. 2019. Rapid Adaptive Camouflage in Cephalopods. In: Oxford Research
Encyclopedia of Neuroscience. Oxford University Press. doi:10.1017/CBO9780511852053.009.

Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. 2005. Blast2GO: a universal tool for
annotation, visualization and analysis in functional genomics research. Bioinformatics.
21(18):3674–3676. doi:10.1093/bioinformatics/bti610.

Coombe L, Zhang J, Vandervalk BP, Chu J, Jackman SD, Birol I, Warren RL. 2018. ARKS:
chromosome-scale scaffolding of human genome drafts with linked read kmers. BMC
Bioinformatics. 19(1):234. doi:10.1186/s12859-018-2243-x.

Dehal P, Boore JL. 2005. Two rounds of whole genome duplication in the ancestral vertebrate. PLoS
Biol. 3(10):e314. doi:10.1371/journal.pbio.0030314

Deryckere A, Styfhals R, Elagoz AM, Maes GE, Seuntjens E. 2021. Identification of neural progenitor
cells and their progeny reveals long distance migration in the developing octopus brain. eLife.
10:e69161. doi:10.7554/eLife.69161.

Deryckere A, Styfhals R, Vidal EAG, Almansa E, Seuntjens E. 2020. A practical staging atlas to study
embryonic development of *Octopus vulgaris* under controlled laboratory conditions. BMC Dev
Biol. 20(1):7. doi:10.1186/s12861-020-00212-6.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29(1):15–21.
doi:10.1093/bioinformatics/bts635.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol I, Lander
ES, Aiden AP, et al. 2017. De novo assembly of the *Aedes aegypti* genome using Hi-C yields
chromosome-length scaffolds. Science. 356(6333):92–95. doi:10.1126/science.aal3327.

Dudchenko O, Shamim MS, Batra SS, Durand NC, Musial NT, Mostofa R, Pham M, Glenn St Hilaire B,
Yao W, Stamenova E, et al. 2018. The Juicebox Assembly Tools module facilitates *de novo*
assembly of mammalian genomes with chromosome-length scaffolds for under $1000. bioRxiv
254797. [accessed 2023 Apr 4]. doi:10.1101/254797.

Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL. 2016. Juicebox
Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. Cell Syst.
3(1):99–101. doi:10.1016/j.cels.2015.07.012.

Fiorito G, Affuso A, Anderson DB, Basil J, Bonnaud L, Botta G, Cole A, D'Angelo L, De Girolamo P,
Dennison N, et al. 2014. Cephalopods in neuroscience: regulations, research and the 3Rs. Invert
Neurosci. 14(1):13–36. doi:10.1007/s10158-013-0165-x.

Fiorito G, Affuso A, Basil J, Cole A, de Girolamo P, D'Angelo L, Dickel L, Gestal C, Grasso F, Kuba M,
et al. 2015. Guidelines for the Care and Welfare of Cephalopods in Research –A consensus based
on an initiative by CephRes, FELASA and the Boyd Group. Lab Anim. 49(2_suppl):1–90.
doi:10.1177/0023677215580006.

Formenti G, Rhie A, Balacco J, Haase B, Mountcastle J, Fedrigo O, Brown S, Capodiferro MR, Al-Ajli
FO, Ambrosini R, et al. 2021. Complete vertebrate mitogenomes reveal widespread repeats and
gene duplications. Genome Biol. 22(1):120. doi:10.1186/s13059-021-02336-9.

Gleadall IG. 2016. *Octopus sinensis* d'Orbigny, 1841 (Cephalopoda: Octopodidae): Valid Species Name
for the Commercially Valuable East Asian Common Octopus. Species Divers. 21(1):31–42.
doi:10.12782/sd.21.1.031.

Gao YM, Natsukari Y. 1990. Karyological Studies on Seven Cephalopods. Venus Jpn J Malacol. 49(2):
126-145. doi:10.18941/venusjjm.49.2_126.

García-Fernández P, Prado-Alvarez M, Nande M, Garcia de la serrana D, Perales-Raya C, Almansa E,
Varó I, Gestal C. 2019. Global impact of diet and temperature over aquaculture of *Octopus*
vulgaris paralarvae from a transcriptomic approach. Sci Rep. 9(1):10312. doi:10.1038/s41598-
019-46492-2.

Guan D, McCarthy SA, Wood J, Howe K, Wang Y, Durbin R. 2020. Identifying and removing haplotypic
duplication in primary genome assemblies. Valencia A, editor. Bioinformatics. 36(9):2896–2898.
doi:10.1093/bioinformatics/btaa025.

Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR. 2008.
Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to
Assemble Spliced Alignments. Genome Biol. 9(1):R7. doi:10.1186/gb-2008-9-1-r7.

Hochner B. 2012. An embodied view of octopus neurobiology. Curr Biol. 22(20):R887-R892.
doi:10.1016/j.cub.2012.09.001

Hochner B, Shomrat T, Fiorito G. 2006. The octopus: a model for a comparative analysis of the evolution
of learning and memory mechanisms. Biol Bull. 210(3):308-317. doi:10.2307/4134567

Hu J, Wang Z, Sun Z, Hu B, Ayoola AO, Liang F, Li J, Sandoval JR, Cooper DN, Ye K, et al. 2023. An
efficient error correction and accurate assembly tool for noisy long reads. bioRxiv
2023.03.09.531669. [accessed 2023 Mar 24]. doi:10.1101/2023.03.09.531669.

Inaba A. 1959. Notes on the Chromosomes of Two Species of Octopods (Cephalopoda, Mollusca). Jpn J

Genet. 34(5):137–139. doi:10.1266/jjg.34.137.

Iwata H, Gotoh O. 2012. Benchmarking spliced alignment programs including Spaln2, an extended
version of Spaln that incorporates additional species-specific features. Nucleic Acids Res.
40(20):e161–e161. doi:10.1093/nar/gks708.

Jackman SD, Coombe L, Chu J, Warren RL, Vandervalk BP, Yeo S, Xue Z, Mohamadi H, Bohlmann J,
Jones SJM, et al. 2018. Tigmint: correcting assembly errors using linked reads from large
molecules. BMC Bioinformatics. 19(1):393. doi:10.1186/s12859-018-2425-6.

Jiang D, Liu Q, Sun J, Liu S, Fan G, Wang L, Zhang Y, Seim I, An S, Liu X, et al. 2022. The gold-ringed
octopus (Amphioctopus fangsiao) genome and cerebral single-nucleus transcriptomes provide
insights into the evolution of karyotype and neural novelties. BMC Biol. 20(1):289.
doi:10.1186/s12915-022-01500-2.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka
G, et al. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics.
30(9):1236–1240. doi:10.1093/bioinformatics/btu031.

Kim B-M, Kang S, Ahn D-H, Jung S-H, Rhee H, Yoo JS, Lee J-E, Lee S, Han Y-H, Ryu K-B, et al. 2018
Sep 25. The genome of common long-arm octopus *Octopus minor*. GigaScience.
doi:10.1093/gigascience/giy119.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat
graphs. Nat Biotechnol. 37(5):540–546. doi:10.1038/s41587-019-0072-8.

Köster J, Rahmann S. 2012. Snakemake—a scalable bioinformatics workflow engine. Bioinformatics.
28(19):2520–2522. doi:10.1093/bioinformatics/bts480.

Kundu R, Casey J, Sung W-K. 2019. HyPo: Super Fast & Accurate Polisher for Long Read Genome
Assemblies. bioRxiv 2019.12.19.882506. doi:10.1101/2019.12.19.882506.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile
and open software for comparing large genomes. Genome Biol. 5(2):R12. doi:10.1186/gb-2004-
5-2-r12.

[GCF_006345805.1] Li F, Bian L, Ge J, Han F, Liu Z, Li X, Liu Y, Lin Z, Shi H, Liu C, et al. 2020.
Chromosome-level genome assembly of the East Asian common octopus (*Octopus sinensis*)
using PacBio sequencing and Hi-C technology. Mol Ecol Resour. 20(6):1572–1582.
doi:10.1111/1755-0998.13216.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
doi:10.48550/ARXIV.1303.3997.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Birol I, editor. Bioinformatics.
34(18):3094–3100. doi:10.1093/bioinformatics/bty191.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000
Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and
SAMtools. Bioinformatics. 25(16):2078–2079. doi:10.1093/bioinformatics/btp352.

Lomsadze A, Burns PD, Borodovsky M. 2014. Integration of mapped RNA-Seq reads into automatic
training of eukaryotic gene finding algorithm. Nucleic Acids Res. 42(15):e119–e119.
doi:10.1093/nar/gku557.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and
Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
Eukaryotic, Prokaryotic, and Viral Genomes. Kelley J, editor. Mol Biol Evol. 38(10):4647–4654.
doi:10.1093/molbev/msab199.

Mapleson D, Venturini L, Kaithakottil G, Swarbreck D. 2018. Efficient and accurate detection of splice
junctions from RNA-seq with Portcullis. GigaScience. 7(12). doi:10.1093/gigascience/giy131.

Marini G, De Sio F, Ponte G, Fiorito G. 2017. Behavioral Analysis of Learning and Memory in
Cephalopods ☆. In: Learning and Memory: A Comprehensive Reference. Elsevier. p.
441–462. doi:10.1016/B978-0-12-809324-5.21024-9.

Marino A, Kizenko A, Wong WY, Ghiselli F, Simakov O. 2022. Repeat Age Decomposition Informs an
Ancient Set of Repeats Associated With Coleoid Cephalopod Divergence. Front Genet.
13:793734. doi:10.3389/fgene.2022.793734.

Meyer A, Schartl M. 1999. Gene and genome duplications in vertebrates: the one-to-four (-to-eight in
fish) rule and the evolution of novel gene functions. Curr Opin Cell Biol. 11(6):699-704.
doi:10.1016/s0955-0674(99)00039-3

Niknafs YS, Pandian B, Iyer HK, Chinnaiyan AM, Iyer MK. 2017. TACO produces robust multisample
transcriptome assemblies from RNA-seq. Nat Methods. 14(1):68–70. doi:10.1038/nmeth.4078.

Open2C, Abdennur N, Fudenberg G, Flyamer IM, Galitsyna AA, Goloborodko A, Imakaev M, Venev
SV. 2023. Pairtools: from sequencing data to chromosome contacts. bioRxiv 2023.02.13.528389.
[accessed 2023 Apr 4]. doi:10.1101/2023.02.13.528389

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie enables
improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol. 33(3):290–295.
doi:10.1038/nbt.3122.

Petrosino G, Ponte G, Volpe M, Zarrella I, Ansaloni F, Langella C, Di Cristina G, Finaurini S, Russo MT,
Basu S, et al. 2022. Identification of LINE retrotransposons and long non-coding RNAs
expressed in the octopus brain. BMC Biol. 20(1):116. doi:10.1186/s12915-022-01303-5.

Ponte G, Chiandetti C, Edelman DB, Imperadore P, Pieroni EM, Fiorito G. 2022. Cephalopod Behavior:
From Neural Plasticity to Consciousness. Front Syst Neurosci. 15:787139.
doi:10.3389/fnsys.2021.787139.

Ponte G, Taite M, Borrelli L, Tarallo A, Allcock AL, Fiorito G. 2021. Cerebrotypes in Cephalopods:
Brain Diversity and Its Correlation With Species Habits, Life History, and Physiological
Adaptations. Front Neuroanat. 14:565109. doi:10.3389/fnana.2020.565109.

Rhie A, Walenz BP, Koren S, Phillippy AM. 2020. Merqury: reference-free quality, completeness, and
phasing assessment for genome assemblies. Genome Biol. 21(1):245. doi:10.1186/s13059-020-
02134-9.

Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP. 2023. igv.js: an embeddable JavaScript
implementation of the Integrative Genomics Viewer (IGV). Alkan C, editor. Bioinformatics.
39(1):btac830. doi:10.1093/bioinformatics/btac830.

Schmidbaur H, Kawaguchi A, Clarence T, Fu X, Hoang OP, Zimmermann B, Ritschard EA,

Weissenbacher A, Foster JS, Nyholm SV, et al. 2022. Emergence of novel cephalopod gene
regulation and expression through large-scale genome reorganization. Nat Commun. 13(1):2172.
doi:10.1038/s41467-022-29694-7.

Shigeno S, Andrews PLR, Ponte G, Fiorito G. 2018. Cephalopod Brains: An Overview of Current
Knowledge to Facilitate Comparison With Vertebrates. Front Physiol. 9:952.
doi:10.3389/fphys.2018.00952.

Smit AFA, Hubley R, Green P. 2013-2015. RepeatMasker Open-4.0. http://www.repeatmasker.org.

Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a
generalized hidden Markov model that uses hints from external sources. BMC Bioinformatics.
7(1):62. doi:10.1186/1471-2105-7-62.

Styfhals R, Zolotarov G, Hulselmans G, Spanier KI, Poovathingal S, Elagoz AM, De Winter S,
Deryckere A, Rajewsky N, Ponte G, et al. 2022. Cell type diversity in a developing octopus brain.
Nat Commun. 13(1):7392. doi:10.1038/s41467-022-35198-1.

Taite M, Fernández-Álvarez FÁ, Braid HE, Bush SL, Bolstad K, Drewery J, Mills S, Strugnell JM,
Vecchione M, Villanueva R, et al. 2023. Genome skimming elucidates the evolutionary history of
Octopoda. Mol Phylogenet Evol. 182:107729. doi:10.1016/j.ympev.2023.107729.

Vara C, Paytuví-Gallart A, Cuartero Y, Álvarez-González L, Marín-Gual L, Garcia F, Florit-Sabater B,
Capilla L, Sanchéz-Guillén RA, Sarrate Z, et al. 2021. The impact of chromosomal fusions on 3D
genome folding and recombination in the germ line. Nat Commun. 12(1):2981.
doi:10.1038/s41467-021-23270-1.

Vitturi R, Rasotto MB, Farinella-Ferruzza N. 1982. The chromosomes of 16 molluscan species. Bolletino
Zool. 49(1–2):61–71. doi:10.1080/11250008209439373.

Wang J, Zheng X. 2017. Comparison of the genetic relationship between nine Cephalopod species based
on cluster analysis of karyotype evolutionary distance. Comp Cytogenet. 11(3):477–494.
doi:10.3897/compcytogen.v11i3.12752.

Wang ZY, Ragsdale CW. 2019. Cephalopod Nervous System Organization. In: Oxford Research
Encyclopedia of Neuroscience. Oxford University Press.
doi:10.1093/acrefore/9780190264086.013.181

Warren RL, Yang C, Vandervalk BP, Behsaz B, Lagman A, Jones SJM, Birol I. 2015. LINKS: Scalable,
alignment-free scaffolding of draft genomes with long reads. GigaScience. 4(1):35.
doi:10.1186/s13742-015-0076-3.

Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. 2017. Direct determination of diploid genome
sequences. Genome Res. 27(5):757–767. doi:10.1101/gr.214874.116.

Whitelaw BL, Jones DB, Guppy J, Morse P, Strugnell JM, Cooke IR, Zenger K. 2022. High-Density
Genetic Linkage Map of the Southern Blue-ringed Octopus (Octopodidae: Hapalochlaena
maculosa). Diversity. 14(12):1068. doi:10.3390/d14121068.

Yokobori S -i. 2004. Long-Term Conservation of Six Duplicated Structural Genes in Cephalopod
Mitochondrial Genomes. Mol Biol Evol. 21(11):2034–2046. doi:10.1093/molbev/msh227.

Young JZ. 1964. A Model of the Brain. Oxford University Press, USA.

Zarrella I, Herten K, Maes GE, Tai S, Yang M, Seuntjens E, Ritschard EA, Zach M, Styfhals R, Sanges
R, et al. 2019. The survey and reference assisted assembly of the *Octopus vulgaris* genome. Sci
Data. 6(1):13. doi:10.1038/s41597-019-0017-6.

Zhang H, Song L, Wang X, Cheng H, Wang C, Meyer CA, Liu T, Tang M, Aluru S, Yue F, et al. 2021.
Fast alignment and preprocessing of chromatin profiles with Chromap. Nat Commun. 12(1):6566.
doi:10.1038/s41467-021-26865-w.

Zhou C, McCarthy SA, Durbin R. 2023. YaHS: yet another Hi-C scaffolding tool. Alkan C, editor.
Bioinformatics. 39(1):btac808. doi:10.1093/bioinformatics/btac808.

Zolotarov G, Fromm B, Legnini I, Ayoub S, Polese G, Maselli V, Chabot PJ, Vinther J, Styfhals R,
Seuntjens E, et al. 2022. MicroRNAs are deeply linked to the emergence of the complex octopus
brain. Sci Adv. 8(47):eadd9938. doi:10.1126/sciadv.add9938.

Supplementary Information for:

A chromosome-level reference genome   for the

common octopus, *Octopus vulgaris* (Cuvier, 1797)

Dalila Destanovic[1,*], Darrin T. Schultz[1,*,+], Ruth Styfhals[2,5], Fernando Cruz[6], Jèssica Gómez-Garrido[6],
Marta Gut[6], Ivo Gut[6,7], Graziano Fiorito[5], Oleg Simakov[1,+], Tyler S. Alioto[6,7,+], Giovanna Ponte[5,+], Eve
Seuntjens[2,3,4,+]

736
737   **This PDF file includes:**

738
Supplementary Figure 1 [Genome assembly pipeline]
Supplementary Figure 2 [Contamination analysis of the manually curated genome]
Supplementary Figure 3 [Contamination analysis of the final version of *O. vulgaris* genome]
Supplementary Figure 4 [Annotation workflow]
Supplementary Figure 5 [Alignment of ONT reads to the *O. vulgaris* mitochondrial genome.]
Supplementary Figure 6 [Coverage of ambiguous positions in the mitochondrial genome]

745
746   **Other Supplementary Data for this manuscript include the following:**

747
Supplementary Data 1 [Supplementary_Data_1.xlsx]
Supplementary Data 2 [Supplementary_Data_2.xlsx]
Supplementary Data 3 [Supplementary_Data_3.xlsx]
Supplementary Data 4 [Supplementary_Data_4.xlsx]
Supplementary Data 5 [Supplementary_Data_5.txt]
Supplementary Data 6 [Supplementary_Data_6.xlsx]
Supplementary Data 7 [Supplementary_Data_7.xlsx]
755

     Supplementary Figures

758 Supplementary Figure 1 | Genome assembly pipeline. Snakemake workflow is used to generate the scaffolded *Octopus*
759 *genome* assembly.

Supplementary Figure 2 | Contamination analysis of the manually curated genome. The original manually curated assembly
had some sequences belonging to other phyla. These fragments were found in the unplaced scaffolds in the assembly.

761
762

Supplementary Figure 3 | Contamination analysis of the final version of *O. vulgaris* genome. Decontamination of the assembly was successful, as the molluscan and no-hit sequences were kept. This is the final version of the chromosome-scale genome that was generated.

Supplementary Figure 4 | Annotation workflow. Combined data of protein sequences from different cephalopod species, and RNA sequences generated from adult and embryonic tissue were used to annotate the genome. This resulted in the annotation of CDS, UTRs, and alternative splice variants for the genome.

768
769
770
771

Supplementary Figure 5 | Alignment of ONT reads to the *O. vulgaris* mitochondrial genome. The ONT reads aligned to the
mitochondrial genome support the consensus of each position except for 16 nucleotides (see vertical coloured bars along the
coverage track).

Supplementary Figure 6 | Coverage of ambiguous positions in the mitochondrial genome. The 16 positions that were
polymorphic were single-nucleotide differences. The consensus sequence is that with the highest frequency in the reads. One
example is shown here, with the G in the mitogenome appearing in most of the reads except one with an A.