Quantifying network behavior in the rat prefrontal cortex:

a reproducibility crisis

Congzhou M. Sha[1,2], Jian Wang[2], Richard B. Mailman[2,3,*], Yang Yang[2,*], Nikolay V. Dokholyan[1,2,4,5,6,*]

[1] *Department of Engineering Science and Mechanics, Penn State University, University Park, PA USA*

[2] *Department of Pharmacology, Penn State College of Medicine, Hershey, PA USA*

[3] *Department of Neurology, Penn State College of Medicine, Hershey, PA USA*

[4] *Department of Biochemistry and Molecular Biology, Penn State College of Medicine, Hershey, PA USA*

[5] *Department of Chemistry, Penn State University, University Park, PA USA*

[6] *Department of Biomedical Engineering, Penn State University, University Park, PA USA*

Corresponding authors

R.B.M. rbm17@psu.edu

Y.Y.: yyang1@pennstatehealth.psu.edu

N.V.D.: dokh@psu.edu

Abbreviations

Neuroelectrophysiology (NEP), dynamic time warping (DTW), medial prefrontal cortex

(mPFC), Kolmogorov-Smirnov test (KS-test), receiver operating characteristic (ROC),

area under the curve (AUC)

Abstract

The question of how consciousness and behavior arise from neural activity is

fundamental to understanding the brain, and to improving the diagnosis and treatment

of neurological and psychiatric disorders. There is significant murine and primate

literature on how behavior is related to the electrophysiological activity of the medial

prefrontal cortex and its role in working memory processes such as planning and

decision-making. Existing experimental designs, however, have insufficient statistical

power to unravel the complex processes of the prefrontal cortex. We therefore

examined the theoretical limitations of such experiments, providing concrete guidelines

for robust and reproducible science. We piloted the use of dynamic time warping and

associated statistical tests to data from neuron spike trains and local field potentials, to

quantify neural network synchronicity and correlate neuroelectrophysiology with rat

behavior. Our results indicate the statistical limitations of existing data, making

meaningful comparison between dynamic time warping with traditional Fourier and

wavelet analysis currently impossible until larger and cleaner datasets are available.

Significance Statement

The prefrontal cortex is important in decision-making, yet no robust method currently

exists to correlate neuron firing in the PFC to behavior. We argue that existing

experimental designs are ill-suited to addressing these scientific questions, and we

propose a potential method using dynamic time warping to analyze PFC neural

electrical activity. We conclude that careful curation of experimental controls is needed

to separate true neural signals from noise accurately.

Introduction

Hodgkin and Huxley's model of neuron action potentials derived from experiments with the squid giant axon[1] was the seminal event in neurophysiology[2–4]. The central dogma of modern neuroscience is that neuron electrochemical activity and connectivity at the microscopic level can provide a clear understanding of complex behaviors. Thus, measuring electrical signals can be sufficient to integrate the chemical signaling and activity of neural networks.

The visual cortex and its relationship to visual processing is one of the better-understood neural networks because there are significant spatial correlations in signals that result from local connections between neurons. These spatial autocorrelations result in a high signal-to-noise ratio due to the strong intensity of the electric field produced by local synchronized neuron firing. Additionally, the neuroelectrophysiology of individual cells in the visual pathway has allowed mathematical modeling of the performance of specific neurons[5,6]. The spatial autocorrelations of visual neural activity are sufficiently high that low-resolution, indirect measurements of neuron activity, such as local blood flow[7–9] and extracranial electrodes[10], are sufficient to reconstruct perceived images. The organization of the visual cortex may be explained by the fact that images projected on the retina are spatially correlated, and therefore biological neural networks have adapted to take advantage of these correlations.

From an evolutionary perspective, the visual cortex is an ancient structure, and therefore the neural networks involved in visual processing have optimized over time. In contrast, the prefrontal cortex as part of the neocortex is one of the least developed brain regions, especially through evolutionary analysis of its size in mammals[11,12].

Unlike the visual cortex, the prefrontal cortex does not seem to possess highly local and

regular spatial organization[13]. Because of the lack of regular spatial organization of the

prefrontal cortex and the consequent low intensity of electrical signals,

electrophysiology is one of the only methods with adequate sensitivity to probe the fine

structure of the prefrontal neural network. Unfortunately, such measurements are highly

invasive, and experiments are only feasible on laboratory animals such as rats[14–16]. To

find the relationship between neuron firing and behavior, methods are required to

analyze these complex electrical measurements.

The characteristics of this neuron prefrontal network constrain the firing patterns

of neurons. Therefore, methods are required to capture key network characteristics from

recorded spike trains and local field potentials (LFPs). Our hypothesis is that these

network characteristics are embodied by the level of synchronicity of neuron firing:

neurons that work in concert will have spike trains and/or LFPs that are highly

correlated or anti-correlated, whereas neurons that process independent data will not be

correlated.

When analyzing complex data such as neuroelectrophysiology tracings, the

major statistical concern is bias and overfitting[17]. Although large amounts of data are

collected in these experiments, the degrees of freedom are severely restricted by the

number of animals and recordings collected[18]. When researchers choose a statistical

model for their data, certain assumptions are made about the structure of the data.

Commonly, a class of statistical models is chosen with hyperparameters tuned to best fit

the data. The more hyperparameters that must be tuned, the fewer degrees of freedom

remain for the subsequent goodness-of-fit tests. In this work, we sought to maximize the

degrees of freedom remaining after modeling to address our scientific questions.

Schematically, we must allocate the degrees of freedom in the data to the modelling we

perform and the scientific questions we seek to answer:

$$\mathrm{dof}_{data} = \mathrm{dof}_{model} + \mathrm{dof}_{science}$$

95   (1)

The more degrees of freedom we can devote to science, the more confident we can be

in our statistical tests.

Therefore, we chose to investigate neuron synchronicity using the dynamic time

warping (DTW) method[19]. DTW is an efficient, non-parametric approach to determine

the best alignment of two time series, such that the overall shapes of the time series are

matched. Compared to classical Fourier and wavelet analysis[14,15], DTW gives a global

alignment of the data that is robust to local variations in timing. We predicted that DTW

could discern if two neurons fire synchronously or asynchronously. In contrast, local

timing variations in low frequency signals such as firing rate and voltage lead to

destructive interference when performing the Fourier or wavelet transform, decreasing

the signal-to-noise ratio. In practice, these techniques require the tuning of various

hyperparameters to smooth and denoise the data. Examples include a window size for

the spike-triggered LFP average[14], cutoff frequencies for band-pass filters[14], various

bootstrapping techniques[14], and choice of kernel for support vector machines[15]. In

contrast, traditional DTW uses no additional parameters to yield a dissimilarity index for

each pair of spike trains or LFP tracings, therefore retaining more degrees of freedom

for answering statistical questions.

Thus, we analyzed electrophysiology data collected from rats performing the T-

maze task, a task that evaluates working memory. We use previously published spike

train and LFP recordings taken from the rat medial prefrontal cortex (mPFC)[14–16], along

with previously unpublished data from three additional rats (rats A, B, and C) recorded

using published methods[16]. We investigated if DTW would be useful in elucidating the

connection between neural activity and behavior (Results), and we examined the

underlying assumptions of neuroelectrophysiology experiments (Discussion).

Results

Our theoretical arguments for the low statistical power of existing experimental

designs are presented in the Discussion and Methods (*Reproducibility of electrode*

recordings). The data from the three published studies (Table 1) had rat neuron spike

trains and/or LFPs recorded as the rats performed in T-maze (Figure 1-A). The exact

experimental details for the three studies differed slightly[14–16], but their combined

scientific goal was to analyze the link between neuron recordings and working memory

performance in the T-maze task. We focused on the four-second window of time

centered on the moment at which the rat leaves the T-intersection (decision box in

Figure 1-A), presumably making its choice. This window contained two seconds of pre-

decision neural activity (e.g., cognition and decision-making process) and two seconds

of post-decision neural activity (e.g., evaluation of reward or lack of reward). Since the

experimental designs differed among the studies (e.g., the rat either traversed the T-

maze continuously[14,15] or was picked up by the researcher in between traversals[16]), we

used this limited window of time for our analysis to limit the effects of the design,

constituting one of the only hyperparameters chosen in our analysis. Our hypothesis

was that neural activity was affected significantly by the independent variables of (1)

timing of the recording relative to the choice, (2) correctness of the choice, and (3)

sampled neurons (Table 2). Null and alternative hypotheses for each statistical test we

performed are included in the Supplemental Tables.

We used the DTW method on the neuron spike trains from all three studies and

from the local field potentials for Ito et al.[14] and Yang et al.[16] (Figure 1-B & 1-C). Since

DTW provides a numerical measurement of dissimilarity between firing of neurons, we

converted the resulting DTW matrices into undirected, unweighted graphs

(Supplemental Methods). For each set of neuron spike trains and/or LFPs, we

quantified the connectivity of the resulting DTW graph by a single number, d_{crit}. To

evaluate for the presence of differences among sets of experiments, we used non-

parametric statistical tests (Kruskal-Wallis, Kolmogorov-Smirnov, Mantel, and Boschloo

exact tests) to mitigate both our lack of knowledge of the true underlying probability

distributions and the small sample sizes. These statistical tests and the calculation of

d_{crit} for a given DTW matrix are described in detail in the Supplemental Methods. We

visualized d_{crit} as a function of our independent variables in Figure 2 (spike trains) and

Supplemental Figure 1 (LFPs).

We defined a ***trial*** as a single traversal of the T-maze by a rat. We defined a

session as all trials recorded on a single day for a single rat. In the Ito[14] and Stout[15]

studies, electrodes were adjusted between sessions, and, therefore, a different set of

neurons were recorded between sessions. We set the level of statistical significance

(false positive rate) to α = 0.05.

Neural network firing is consistent across a single session

Our first question was whether in analyzing d_{crit}, could we pool trials from a single

session? The null hypothesis was that during a single session, we may assume that the

rat basal activity is unchanged between recordings in the same session, as measured

by d_{crit}, The alternative hypothesis is that specific factors affect d_{crit} between trials in the

same session, such as the categorical variables in Table 2.

First, we analyzed the normality of d_{crit} with the Shapiro-Wilk test (Supplemental

Tables 1 and 2). For neuron spike trains, we found that in 96.7% (60 out of 62) of

sessions, the samples were not normally-distributed ($p < 0.05$), and for LFPs, we found

that 80.0% (20 out of 25) were not normally-distributed. Comparing spike trains to LFP

with Boschloo's exact test, these fractions were not significantly different (p=0.615). We

performed subgroup analysis on the categorical variables (1) correctness and (2) timing

and found that this proportion was unaffected by further stratification by these variables.

Therefore, we concluded that the Kruskal-Wallis non-parametric test was more

appropriate than classical one-way analysis of variance (ANOVA), due to violation of the

assumption of normally-distributed data and due to the limited number of trials per

session.

Only 4.8% (3 out of 62) of the spike train sessions displayed significant (Kruskal-

Wallis $p < 0.05$) trial-to-trial variance of d_{crit}, regardless of stratification by correctness

and/or timing; the corresponding proportion for LFPs was 12% (3 out of 25). For

completeness, we also performed ANOVA and found that 11.2% (7 out of 62) of the

spike train sessions displayed significant trial-to-trial variance, regardless of stratification;

the corresponding proportion for LFPs was 20% (5 out of 25). We tested if Kruskal-

Wallis gave different results as compared to ANOVA using Boschloo's exact test, and

found that neither the spike train data (p = 0.238) nor the LFPs (p = 0.378) were

significantly different.

Since few of the sessions (within our 5% margin of error for false positives from α =

0.05) had a significant trial-to-trial variance in the spike trains, we concluded that we

may perform pooled analysis of d_{crit} for the spike trains of trials across a single session.

Boschloo's exact test on the LFPs also demonstrated a non-significant deviation from

the 5% margin of error (p = 0.413 for Kruskal-Wallis, p = 0.140 for ANOVA).

Neural network structure is revealed by the DTW matrix

Instead of characterizing network connectivity with a single number (d_{crit}), we

used all the information available in the DTW matrix to determine if firing patterns were

consistent, using non-parametric Mantel tests. We performed all pairwise comparisons

for trials in the same rat to ask the question: given a single session, is there a difference

in Pearson correlation coefficient of before vs. after depending on if we compare (1)

before vs. after of the same trial to (2) before vs. after of different trials? We performed

pairwise Mantel test comparisons among trials in the same session. We performed

Kruskal-Wallis and Kolmogorov-Smirnov tests to determine if these distributions of the

correlation coefficient and *p*-value were different (Supplemental Figures 2 and 3,

Supplemental Table 3). Overall, we found that there was no significant difference trial-

to-trial for the before vs. after DTW matrix correlation.

When we stratified by correctness (Supplemental Table 3), we did observe an effect

for Kolmogorov-Smirnov tests on the Pearson correlation coefficient in 35.7% (5 out of

14) sessions. When we used d_{crit} alone, this proportion was 7.1% (1 out of 14). To

evaluate if the KS-test/Mantel test combination was more powerful than the Kruskal-

Wallis test on d_{crit}, we performed a Boschloo exact test on the contingency table but this

failed to reach statistical significance (p = 0.159). We concluded that there was mixed

evidence for the efficacy of Mantel tests for before vs. after DTW matrices in

determining if a rat made the correct or incorrect choice at the T-intersection.

Neural network differences exist across multiple sessions for a single rat

Our second question was: when analyzing d_{crit}, can we pool all the trials for a

single rat, regardless of the day the recordings were taken? Using the Kruskal-Wallis

test, we found that 70% (7 of 10) rats displayed significant ($p < 0.05$) day-to-day

variance of d_{crit} in their spike trains, regardless of additional stratification by (1)

correctness and (2) timing (Supplemental Table 1). The caveat was that we excluded

six rats from the Yang study[16] in this analysis, because we were unable to apply the

Kruskal-Wallis test since recordings for those rats were only performed on a single day

(dof = 0). Additionally, the three rats which did not display significant differences across

sessions were 17914 and 18471 from the Ito study[14] and rat A from Yang[16]. The same

analysis was performed for LFPs, in which 80% (4 out of 5) rats (all rats except rat A

from Yang[16]) displayed significant variance (Supplemental Table 2). For the LFPs, the

five rats from Stout[15] were excluded since the LFPs were not recorded for those rats.

Boschloo's exact test showed no difference between spike trains and LFPs ($p = 0.922$).

We conclude that the resting cognitive state for each rat may be subject to change

depending on the day that the recording is performed.

Significant neural network heterogeneity exists between rats and studies

Our third question was: does d_{crit} depend on the specific rat? Using the Kruskal-

Wallis test, we found that the Stout[15] and Yang[16] studies displayed significant ($p < 0.001$)

rat-to-rat differences in their spike trains, whereas Ito[14] was non-significant ($p = 0.206$).

Ito[14], however also had the fewest rats ($n = 3$) potentially leading to the insufficient

statistical power of the Kruskal-Wallis test. The LFPs showed significant ($p < 0.001$) rat-

to-rat variation in both Ito[14] and Yang[16]. Pooling all rats, we confirmed ($p < 0.001$) that

the specific study under consideration affects the value of d_{crit} for both spike trains and

LFPs. We conclude that significant heterogeneities exist among studies, animals, and

sessions. The statistical tests supporting this conclusion are summarized in

Supplemental Tables 1 (spike trains) and 2 (LFPs).

DTW distance is inconsistently correlated with physical electrode distance

Do correlations in firing among neurons reflect the underlying spatial distance

between the neurons? We used both Welch's t-test for unequal variances and the non-

parametric Kolmogorov-Smirnov test (KS-test) to test for differences in the distribution

of the DTW distance when grouped by the spatial distance of the associated electrodes.

In all three studies, the d_{crit} between neurons recorded from the same electrode

versus from different electrodes were statistically significantly different ($p < 0.001$) by

both the t-test and KS-test (Supplemental Figures 4, 6-8); however, this classification

was useless as a predictor or regressor[20]. For example, using d_{cutoff} as a classifier of d_{crit}

for determining if two neurons were recorded from the same electrode vs. from different

electrodes, the receiver operator characteristic (ROC) curve is essentially no different

from random guessing (Supplemental Figure 5), with area under the curve (AUC) of

0.51. If we stratify data by session (Supplemental Figures 6-8), however, the ROC curve

and AUC vary for specific rats/sessions (Supplemental Figure 9), and in some cases

demonstrate high predictive value (AUC).

For the Yang study[16], we were able to obtain the exact geometry of the electrode.

Therefore, we were able to stratify DTW distances by the physical distance of the

neurons measured (Figure 3). For this stratification, we were unable to determine a

consistent trend. Rats B and C demonstrate decreased DTW distance (increased

synchronicity) in both neural spike trains and LFPs, whereas the remaining rats

demonstrated slightly increasing DTW distance (decreased synchronicity) in both,

except for rat S. For rat S, however, the large decrease in DTW distance seen in the

spike trains may not be captured in the LFPs, because there was no zero-distance

comparison for LFPs, whereas multiple spike trains recorded from the same electrode

could be thought of as having zero distance.

Discussion

We demonstrated that the DTW distance and the computed parameter d_{crit}

captured some of the mPFC neural network firing dynamics for both the spike trains and

LFPs, which were associated with T-maze task performance. We found weak evidence

that the correctness of rat choice influences the firing dynamics (Supplemental Table 3).

More importantly, we also found that significant heterogeneities exist among studies,

animals, and sessions, as measured by DTW distance and d_{crit}. To demonstrate

meaningful associations between behavior (e.g., T-maze task performance) and neural

network activities (in the mPFC), the data and computed results must be consistent. We

first address the question of statistical consistency through a theoretical discussion of

the experiments and the assumptions made in neuroelectrophysiology.

There are two major issues with attempting to measure a specific population of

neurons. First, precise surgical implantation of electrodes is required; second, enough

neurons must be sampled to capture overall neural network dynamics that can be

consistently associated with certain behaviors. These two issues synergize at our

current level of intracranial electrode technology, complicating fine measurements of

neurons. Wheras the rat brain is far smaller than a human brain (ca. 1 cm^3), it still

contains an estimated 21 million neurons[21]. Reproducible surgical implantation in a

specific area of the rat brain such as the medial prefrontal cortex is therefore highly

dependent on the fine-motor skill of the researcher, and there is no guarantee that the

same neurons or circuits will be sampled.

In fact, in Ito[14] and Stout[15] the electrodes were purposefully adjusted after each

session so that a different set of neurons would be sampled. Due to the lack of spatial

organization of the prefrontal cortex, we postulate that the surgical sampling of neurons

and circuits in the prefrontal cortex is essentially random, and that electrode adjustment

similarly resamples the neurons and circuits being recorded. The only way to counteract

the statistical effects of random sampling is through the measurement of a sufficiently

large fraction of the neurons in the brain area under consideration. In the Methods, we

illustrate these issues with a simple mathematical illustration (*Reproducibility of*

electrode recordings, Figure 1-D).

To provide a guide for researchers investigating the connection between

neuroelectrophysiology and behavior, we asked what did each of the three studies do

right and what could be improved? There are three experimental parameters we

considered: $n_{neurons}$ (the number of neurons recorded); $n_{network\ repeat}$ (the number of times

the same neural network is recorded); and n_{rats} (the number of biological replicates). We

contend that all three of these parameters must be relatively high to make meaningful

conclusions, especially if one wishes to use more powerful parametric statistical tests

over the non-parametric tests we used in this study.

To have any hope of identifying the specific neural network firing patterns which

lead to decision-making in the T-maze, we must consistently identify the neurons

responsible ($n_{neurons}$), the circuits responsible ($n_{network\ repeat}$), and demonstrate that the

conclusions generalize across organisms (n_{rats}). By this simple reasoning, we contend

that the three recent studies we have examined here do not allow us to draw

conclusions, regardless of data analysis method (DTW or otherwise). Stout[15] recorded

too few neurons ($n_{neurons}$ ~1), and changed the neurons recorded across sessions

($n_{network\ repeat} = 1$) for $n_{rats} = 5$. Ito[14] recorded a moderate number of neurons ($n_{neurons}$

~20), but still changed the neurons recorded across sessions ($n_{network\ repeat} = 1$), with n_{rats}

= 3. Yang[16] recorded a moderate number of neurons ($n_{neurons}$ ~20), and kept the same

neurons recorded across sessions ($n_{network\ repeat} = 5$), but only performed these

replicates for $n_{rats} = 2$; the other $n_{rats} = 7$ had ($n_{network\ repeat} = 1$). In short, larger electrodes

($n_{neurons}$), more sessions per experimental condition ($n_{network\ repeat}$), and more animals

(n_{rats}) are needed to achieve a robust and reproducible conclusion. Our

recommendation for electrode size ($n_{neurons}$) depends on the number of neural clusters

or circuits one wishes to investigate (Figure 1-D), whereas $n_{network\ repeat}$ and n_{rats} should

be chosen according to traditional recommendations such that parametric statistical

tests can be performed[22,23].

We now turn to the efficacy of DTW in potentially uncovering connections

between neuron electrical activity and behavior. In our analysis, we demonstrated that

the DTW distance and the computed parameter d_{crit} captured some of the mPFC neural

network firing dynamics for both the spike trains and LFPs. We found weak evidence

that the correctness of rat choice influences the firing dynamics (Supplemental Table 3).

We expect, however, that many more experiments are needed to confirm the

classification of the correctness of a decision based on mPFC activity because our

results indicated that whereas firing trials could be pooled across the same session,

they could not be pooled across different sessions or studies (Supplemental Tables 1

and 2). In terms of spatial dependence of firing synchronicity, we found that for specific

rats and sessions, the classification of DTW distances for recordings taken from the

same vs. different electrodes was both statistically significant (Supplemental Figures 4,

6-8) and useful as a regressor (Supplemental Figure 9). For the Yang study[16], the

spatial dependence of DTW distances was highly dependent on the specific rat sampled

(Figure 3), providing evidence of the lack of spatial organization of the prefrontal cortex.

The difficulties we encountered in analyzing spike trains and LFPs can be

addressed through changes to the experimental design. The number of neurons

sampled is an important characteristic to consider, especially if only a few neurons were

recorded and some recordings represented only a single neuron (e.g., Stout[15]). If the

mPFC is composed of separate neural networks that perform specific, modular tasks, it

is unlikely that recording a low number of neurons will provide meaningful, interpretable

results. Since firing dynamics were consistent across a single session, we posit that

multiple recordings of a single neuron population are the best approach to

characterizing network firing behavior. The DTW spatial dependence remained

consistent across multiple sessions for rats A and C (Figure 3), therefore we are

reasonably confident that we recorded the same population of neurons across the

sessions. In contrast, because the electrodes were adjusted after each session, Ito[14]

and Stout[15] only recorded a single session per neuron population that complicates the

modelling we must perform. Is the additional variation across sessions in Yang[16] due to

changes in the resting cognitive state of the rats? For the data analyzed in this work, it

was not possible for us to consistently distinguish between variance due to noisy

measurements and variance due to rat behavioral changes. It may be necessary to

perform many control sessions to characterize in adequate detail all of the resting

cognitive states of a specific rat. Crucially, recording a single control session as Yang[16]

did for the seven other rats (B, K, Q, R, S, T, X) likely fails to detect variations in the

resting state for a single rat. To complement the recording of many control sessions,

recording additional rat behaviors may yield insight as to how the rat resting cognitive

state influences the network firing dynamics.

Recent advances in imaging and brain modelling provide hope that we may one

day understand the relationship between neuron firing and consciousness. Whole brain

connectomes have been mapped for simple organisms (*C. elegans*[24], *Ciona intestinalis*

larvae[25], and *Platynereis dumerilii*[26]), and recently all 3,000 neurons and 548,000

synapses of a fruit fly larva (*Drosophila melanogaster*) were mapped[27]. Although

mammalian brains are orders of magnitude larger than insect brains, whole-brain

connectomes for rats may not be out of reach in the coming decades. These

connectomes are essential for understanding cognition, but they provide only a static,

anatomic reference upon which neural physiology and pathophysiology must be built.

Our study highlights the data requirements for the next steps in investigating the

function and dynamics of biological neural networks.

Methods and Materials

Acquisition of additional rat recordings

Besides the data we used from the three published studies, new data were

collected from three additional male Sprague-Dawley rats (Charles River Laboratories).

They weighed 300–350 g when received and were named as rats A, B, and C. The

experimental procedure of T-maze alternation and neural recording followed published

protocol[16]. All animal care and surgical procedures were in accordance with the National

Institutes of Health Guide for the Care and Use of Laboratory Animals and Penn State

Hershey Animal Resources Program, and were reviewed and approved by the local

IACUC.

Dynamic time warping and statistical tests

Mathematical details on dynamic time warping, the Mantel test, and the

Kolmogorov-Smirnov test are included in the Supplemental Methods.

Estimated number of neurons in the rat medial prefrontal cortex

Our goal here is to provide a reasonable estimate for the number of neurons that

one must record to reconstruct the dynamics of the entire mPFC. We used existing

estimates of neuron and synaptic density in the rat prefrontal cortex as starting points

for our modeling. We set the total number of neurons in the rat brain[21] to N_{brain} = 2.1 x

10^7 with volume[28] V_{brain} = 2,500 mm^3. We set the volume of the mPFC[28] to V_{mPFC} = 20

mm^3. Assuming a uniform distribution of neuron count throughout the brain, the

estimated number of neurons in the mPFC was $N_{mPFC} = N_{brain} \times \frac{V_{mPFC}}{V_{brain}} = 1.7 \times 10^5$.

Reproducibility of electrode recordings

Here, we discuss the problem of reproducibility in neuroelectrophysiology.

Because of the apparent lack of spatial organization in the mPFC, when electrodes are

implanted, they uniformly sample the estimated N_{mPFC} = 10^5 neurons in the mPFC.

Depending on how cooperative and synchronized neuron firing is, we assume that there

is a certain number of neural clusters $n_{cluster}$ for which the measurement of a single

neuron sufficiently captures the behavior of the entire cluster, and that measurement of

these clusters correlates with rat behavior. We also assumed that sampling is done with

replacement, i.e., $n_{cluster}$ << $N_{mPFC.}$ For reproducibility, one must (1) record the same

neural clusters across biological replicates and (2) identify or classify the neural clusters

to show that findings in one animal generalize to other animals.

(1) The first issue of capturing a certain percentage of the neural clusters in the

rat brain corresponds to a classical problem in combinatorial probability known as the

coupon collection problem. The mathematical question is: given $c = n_{cluster}$ categories,

what is the expected number of samples S which needs to be drawn from those

categories so that all categories are represented at least once? In the case of equal

probabilities for drawing each category,

$$S = c \cdot \sum_{i=1}^{c} \frac{1}{i} = c H_c \; ,$$

404   (2)

405   where H_c is the c^{th} harmonic number. Alternatively, we may require that a certain

406   number $1 < k < c$ out of all the categories is represented[29], so that Eq. (2) is a special

407   case of

$$S(k) = c\,(H_c - H_{c-k}) = c \sum_{i=0}^{k-1} \frac{1}{c - i} \; .$$

408   (3)

In both cases, $S \sim c \log c$ asymptotically. We have plotted the expected number

of clusters we may identify depending on the maximum number of neurons recorded

from each of the studies (Figure 1-D). In practice, it is necessary to first determine the

number of neural clusters that exist and to choose neuron sample size that can

adequately sample all the clusters with high probability.

(2) The second issue of identifying neural clusters across different animals relies

on the researcher's ability to determine which of the $n_{cluster}!$ possible matchings is the

correct matching between any two rats. To reduce the number of possibilities, one must

rely either on prior information (such as typical firing rates or patterns of certain clusters)

or on other measurements of network behavior that are invariant under permutations of

nodes in the graph.

Computational analysis and plotting

All computational analysis was carried out in Python 3.10 and R 4.1, on an M1 Max MacBook Pro with 64 GB RAM. For statistical tests: Shapiro-Wilk, Kruskal-Wallis, and ANOVA were carried out using pingouin 0.5.3[30], Mantel tests[31,32] were carried out using scikit-bio 0.5.7[31], and Kolmogorov-Smirnov and Boschloo exact tests were carried out using SciPy 1.8.1[33]. We implemented dynamic time warping[19] using Cython 0.29.32[34]. We used the DataFrame structure from Pandas 1.5.1[35] to organize our data, and we used seaborn 0.12.1[36] and Matplotlib 3.5.2[37] for plotting. We used NumPy 1.23.3[38,39] for numerical array operations. We used ggpubr 0.6.0[40] for the balloon plots.

Acknowledgements

We thank Hiroshi T. Ito, May-Britt Moser, John J. Stout, and Amy L. Griffin for supplying rat neuroelectrophysiology recordings and precise experimental details for our data analysis.

We acknowledge support from the National Institutes of Health (NIH) 1R35 GM134864 (N.V.D.), 1RF1 AG071675 (Y.Y., N.V.D., R.B.M.), 1R01 AT012053 (N.V.D.), the National Science Foundation 2210963 (N.V.D.), and the Passan Foundation (N.V.D.).

Author contributions (CRediT statement)

C.M.S. Conceptualization, methodology, software, validation, formal analysis, investigation, data curation, writing – original draft, writing – review & editing, visualization.

J.W.　Conceptualization, methodology, software, formal analysis, writing – review &

editing.

R.B.M.　Conceptualization, methodology, validation, resources, writing – review &

editing, supervision, project administration, funding acquisition.

Y.Y.　Conceptualization, methodology, validation, investigation, resources, data

curation, writing – review & editing, supervision, project administration, funding

acquisition.

N.V.D.　Conceptualization, methodology, software, validation, formal analysis,

resources, writing – review & editing, supervision, project administration,

funding acquisition.

Competing interests

We confirm that we have no financial or other interests related to this work which

could affect or have the perception of affecting our objectivity or the content of this

article.

Code and Data Availability

All code and data which are necessary to reproduce the results in this work are

posted on Zenodo (https://doi.org/10.5281/zenodo.7933346).

Tables

Table 1: Characteristics of the three studies analyzed in this work.

	Ito[14]	**Stout**[15]	**Yang**[16]
Year of publication	2018	2020	2022
Number of rats	3	5	9
Total number of sessions (including sessions in which only one neuron was recorded)	13	45	13
Sessions excluded (sessions in which only a single neuron was recorded)	0	9	0
Number of trials (excluding single-neuron recordings)	790	1508	472
Mean neurons recorded (s.d.)	17.56 (4.24)	3.31 (2.21)	31.8 (28.55)
Range of neurons recorded	[10, 26]	[1, 10]	[5, 76]
Fixed electrode	No	No	Yes
Spatial metadata	Yes	Yes	Yes

461   *Table 2: Listing of categorical variables used for stratification/grouping in our*
462   *analyses.*

Categorical variable	**Values**	**Interpretation**
timing	before, after	Data from before vs. after the rat left the

		T-intersection
correctness	true, false	True if rat chose the correct (i.e. alternate) branch of the T-maze during the task
study	Ito, Stout, Yang	The study to which the rat belongs
rat	17 total animals	The rat identity
session	62 total sessions (with more than 1 neuron recorded)	The rat identity and the date of recording

Figures

Figure 1: Outline of experiments, our working model, and data analysis. (a) Rats

complete the T-maze task, in which the reward switches locations after each completion

of the task. This task tests the working memory and decision-making capacity of the rat.

(b) Neurons in the medial prefrontal cortex are recorded using intracranial electrodes,

resulting in the measurement of specific clusters of neurons. (c) Dynamic time warping

computes an optimal alignment of neuron spike trains and local field potentials,

producing a single number characterizing the synchronicity of the neurons. (d)

Depending on the number of distinct neuron clusters in the region of interest, a certain

number of neurons must be sampled to produce a representative sample of all the

clusters; in combinatorics, this is known as the coupon collector problem. We plot the

number of clusters that the maximum number of neurons recorded in each of the three

studies can theoretically discern.

Figure 2: Balloon plot of neuron spike train d_{crit}, grouped by rat, correctness of T-arm choice, and timing with respect to T-arm choice. Each box represents a single rat. The x-axis represents timing (before vs. after the rat visits the T-maze intersection) and the y-axis represents correctness of T-arm taking (True vs. False). The size and color of each marker represents $\log_{10} d_{\text{crit}}$. For rats with multiple trials, we took the mean of $\log_{10} d_{\text{crit}}$. Rats 17913, 17914, and 18471 are from Ito[14] (2018); single letter rats are from Yang[16] (2022); the remaining five rats are from Stout[15] (2020). The most

489   visually striking difference is in the Ito[14] rats, in which there is a stark contrast for spike

490   trains recorded before vs. after making a choice.

Figure 3: DTW distance for neuron spike trains (left) and LFPs (right) as a function of electrode distance (from Yang[16]). The median DTW distance is plotted for each electrode distance, across all sessions for the indicated rat (rat A had 4 sessions, rat C had 3 sessions, and the remaining rats had a single session). 95% confidence intervals were estimated using 1,000 bootstrap samples. There is no consistent correlation between the DTW and electrode distance for either dataset, supporting the idea that there is no discernible spatial organization of neurons in the mPFC. Note that the DTW metric is different for the two datasets, with the distance for neuron spike trains reported as a time (ms) and the distance for LFPs reported as a voltage (V). Physical distance is reported in multiples of 0.25 mm.

References

1. Hodgkin, A. L. & Huxley, A. F. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol* **117**, 500–544 (1952).

2. Coull, J. T. Neural correlates of attention and arousal: insights from electrophysiology, functional neuroimaging and psychopharmacology. *Prog Neurobiol* **55**, 343–361 (1998).

3. Citron, F. M. M. Neural correlates of written emotion word processing: A review of recent electrophysiological and hemodynamic neuroimaging studies. *Brain Lang* **122**, 211–226 (2012).

4. Kucyi, A. & Davis, K. D. The Neural Code for Pain: From Single-Cell Electrophysiology to the Dynamic Pain Connectome. *The Neuroscientist* **23**, 397–414 (2017).

5. Hartline, H. K. Visual Receptors and Retinal Interaction. *Science (1979)* **164**, 270–278 (1969).

6. Gilbert, C. D. & Wiesel, T. N. Receptive field dynamics in adult primary visual cortex. *Nature* **356**, 150–152 (1992).

7. Miyawaki, Y. *et al.* Visual Image Reconstruction from Human Brain Activity using a Combination of Multiscale Local Image Decoders. *Neuron* **60**, 915–929 (2008).

8. Shen, G., Dwivedi, K., Majima, K., Horikawa, T. & Kamitani, Y. End-to-End Deep Image Reconstruction From Human Brain Activity. *Front Comput Neurosci* **13**, (2019).

9. Ren, Z. *et al.* Reconstructing seen image from brain activity by visually-guided cognitive representation and adversarial learning. *Neuroimage* **228**, 117602 (2021).

10. Wakita, S., Orima, T. & Motoyoshi, I. Photorealistic Reconstruction of Visual Texture From EEG Signals. *Front Comput Neurosci* **15**, (2021).

11. Teffer, K. & Semendeferi, K. Human prefrontal cortex. in 191–218 (2012). doi:10.1016/B978-0-444-53860-4.00009-X.

12. Preuss, T. M. & Wise, S. P. Evolution of prefrontal cortex. *Neuropsychopharmacology* **47**, 3–19 (2022).

13. O'Reilly, R. C. The What and How of prefrontal cortical organization. *Trends Neurosci* **33**, 355–361 (2010).

14. Ito, H. T., Moser, E. I. & Moser, M.-B. Supramammillary Nucleus Modulates Spike-Time Coordination in the Prefrontal-Thalamo-Hippocampal Circuit during Navigation. *Neuron* **99**, 576-587.e5 (2018).

15. Stout, J. J. & Griffin, A. L. Representations of On-Going Behavior and Future Actions During a Spatial Working Memory Task by a High Firing-Rate Population of Medial Prefrontal Cortex Neurons. *Front Behav Neurosci* **14**, (2020).

16.   Yang, Y., Kocher, S. D., Lewis, M. M. & Mailman, R. B. Dose-Dependent
Regulation on Prefrontal Neuronal Working Memory by Dopamine D1 Agonists:
Evidence of Receptor Functional Selectivity-Related Mechanisms. *Front Neurosci*
16, (2022).

17.   Cawley, G. C. & Talbot, N. L. C. On over-fitting in model selection and
subsequent selection bias in performance evaluation. *Journal of Machine*
Learning Research **11**, (2010).

18.   Marek, S. *et al.* Reproducible brain-wide association studies require thousands of
individuals. *Nature* **603**, (2022).

19.   Vintsyuk, T. K. Speech discrimination by dynamic programming. *Cybernetics* **4**,
52–57 (1972).

20.   Lo, A., Chernoff, H., Zheng, T. & Lo, S.-H. Why significant variables aren't
automatically good predictors. *Proceedings of the National Academy of Sciences*
112, 13892–13897 (2015).

21.   Korbo, L. *et al.* An efficient method for estimating the total number of neurons in
rat brain cortex. *J Neurosci Methods* **31**, 93–100 (1990).

22.   de Winter, J. C. F. Using the student's t-test with extremely small sample sizes.
Practical Assessment, Research and Evaluation **18**, (2013).

23.   Curtis, M. J. *et al.* Experimental design and analysis and their reporting: New
guidance for publication in BJP. *British Journal of Pharmacology* vol. 172 Preprint
at https://doi.org/10.1111/bph.12856 (2015).

24.   Cook, S. J. *et al.* Whole-animal connectomes of both Caenorhabditis elegans
sexes. *Nature* **571**, 63–71 (2019).

25.   Ryan, K., Lu, Z. & Meinertzhagen, I. A. The CNS connectome of a tadpole larva of
Ciona intestinalis (L.) highlights sidedness in the brain of a chordate sibling. *Elife*
5, (2016).

26.   Verasztó, C. *et al.* Whole-animal connectome and cell-type complement of the
three-segmented Platynereis dumerilii larva. *bioRxiv* 2020.08.21.260984 (2020)
doi:10.1101/2020.08.21.260984.

27.   Winding, M. *et al.* The connectome of an insect brain. *Science (1979)* **379**, (2023).

28.   Hamezah, H. S. *et al.* Volumetric changes in the aging rat brain and its impact on
cognitive and locomotor functions. *Exp Gerontol* **99**, 69–79 (2017).

29.   Ferrante, M. & Frigo, N. On the expected number of different records in a random
sample. *ArXiv* (2012).

30.   Vallat, R. Pingouin: statistics in Python. *J Open Source Softw* **3**, 1026 (2018).

31.   The scikit-bio development team. scikit-bio: A Bioinformatics Library for Data
Scientists, Students, and Developers. Preprint at (2023).

32.   Mantel, N. The Detection of Disease Clustering and a Generalized Regression
Approach. *Cancer Res* **27**, 209–220 (1967).

33.   Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in
Python. *Nat Methods* **17**, 261–272 (2020).

34.   Behnel, S. *et al.* Cython: The Best of Both Worlds. *Comput Sci Eng* **13**, 31–39
(2011).

35.   The pandas development team. pandas-dev/pandas: Pandas. Preprint at (2022).

36.   Waskom, M. seaborn: statistical data visualization. *J Open Source Softw* **6**,
(2021).

37.   Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput Sci Eng* **9**, 90–95
(2007).

38.   Van Der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy array: A structure
for efficient numerical computation. *Comput Sci Eng* **13**, 22–30 (2011).

39.   Harris, C. R. *et al.* Array programming with NumPy. *Nature* vol. 585 357–362
Preprint at https://doi.org/10.1038/s41586-020-2649-2 (2020).

40.   Kassambara, A. ggpubr: 'ggplot2' Based Publication Ready Plots.

596