Breakthrough report

Gametophytic epigenetic regulators MEA and DME synergistically suppress
ectopic shoot formation in *Arabidopsis*

Mohit Rajabhoj[1], Sudev Sankar[1,3], Ramesh Bondada[1,4], Anju P Shanmukhan[2],
Kalika Prasad[2]*, Ravi Maruthachalam[1]*

[1]*School of Biology, IISER-Thiruvananthapuram, Trivandrum, Kerala 695551, India*
[2]*Department of Biology, IISER-Pune, Pune, Maharashtra 411008, India*
[3]*Present address: Department of Plant and Microbial Biology & Zurich-Basel Plant Science*
Center, University of Zurich, Zollikerstrasse 107, Zurich 8008, Switzerland
[4]*Present address: Department of Biology and Gus R. Douglass Institute, West Virginia State*
University, WV 25112-1000, USA
*Corresponding authors: ravi@iisertvm.ac.in, kalika.prasad@iiserpune.ac.in

Keywords: MEDEA, DEMETER, shoot development, epigenetics, histone
methylation, DNA methylation

Running title: MEA and DME control shoot development

Abstract

The gametophyte to sporophyte transition is an important event facilitating the alternation of generations in a plant life cycle. The paternally imprinted epigenetic regulators DEMETER(DME) and MEDEA(MEA) synergistically control central cell proliferation and differentiation, ensuring proper gametophyte to sporophyte transition in *Arabidopsis.* Mutant alleles of *DME* and *MEA* are maternal lethal, eluding the recovery of recessive homozygotes to examine their role in the sporophyte. Here, we exploited the paternal transmission of these mutant alleles coupled with CENH3-haploid inducer to generate *mea*-1;*dme*-2 sporophytes. Strikingly, the simultaneous loss of function of MEA and DME leads to the emergence of multiple shoot meristems at the apical pole of the plant body axis. *DME* and *MEA* are expressed in the developing shoot apex and regulate the expression of various shoot-promoting factors. Chromatin immunoprecipitation(ChIP) and DNA methylation analysis revealed several shoot regulators as potential targets of MEA and DME, with *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings showcasing their misexpression. Knockdown of upregulated shoot-promoting factors STM, CUC2, and PLT5 rescued the ectopic shoot phenotypes. Our findings reveal a previously unrecognized synergistic role of MEA and DME in restricting the meristematic activity at the shoot apex during sporophytic development to orchestrate the plant architecture.

Introduction

Sporophyte development in higher plants results from a fine balance between cell growth and differentiation at the meristems. Primary meristems at the shoot and root apices are responsible for all post-embryonic organ development, coordinated by the polar gradients of plant hormones (Stahl and Simon, 2010; Ten Hove *et al.*, 2015). Throughout this process, epigenetic regulatory complexes provide an additional fine-tuning of gene expression, further contributing to plant developmental plasticity. Regulators of Histone methylation like Polycomb repressive complex 1 & 2 (PRC1 & PRC2) transcriptionally repress embryonic regulators such as *BABYBOOM* (*BBM*),

LEAFY COTYLEDON1 & 2 (*LEC1 & LEC2*) (Chen *et al*., 2010). In contrast, the EM-
BRYONIC FLOWER (EMF)- and VERNALIZATION (VRN)-PRC2s regulate sporo-
phyte development during the transition from the vegetative to the reproductive life
phase (Yoshida *et al*., 2001; De Lucia *et al*., 2008). Genes expressed during shoot
development like *WUSCHEL* (*WUS*), *SHOOT MERISTEMLESS* (*STM*), *AGAMOUS*
(*AG*), *BREVIPEDICELLUS* (*BP*), *KNOTTED-LIKE FROM ARABIDOPSIS THALIANA*
2 (*KNAT2*), and *KNUCKLES* (*KNU*) are known targets of PRC2 components, and
their expression domain is instructed by trimethylation of histone H3 at Lys27
(H3K27me3) repression mark (Kwon *et al*., 2006; Lodha *et al*., 2013; Sun *et al*.,
2019; Xiao and Wagner, 2015; Xiao *et al*., 2017). *PLETHORA* (*PLT*) transcription
factors are known for regulating lateral organ initiation at both the root and shoot api-
ces (Prasad *et al*., 2011; Hofhuis *et al*., 2013). Although auxin gradients mainly regu-
late the PLT expression domains, the root meristem-specific PLTs are also epigenet-
ically regulated (Aichinger *et al*., 2011; Lee and Seo, 2018). *PLTs*, in turn, control the
shoot-promoting gene family *CUP-SHAPED COTYLEDON* (*CUC*) (Kareem *et al*.,
2015). CUCs and STM positively regulate each other to maintain stem cell popula-
tions and promote organ formation at the shoot apical meristem (SAM) boundary
(Spinelli *et al*., 2011). CUCs are also positively regulated by chromatin remodeling
factors like BRAHMA (BRM), which is required to establish leaf primordia (Shen and
Xu, 2009). These reports indicate the maintenance of a delicate balance of CUCs
expression during shoot development *via* direct or indirect epigenetic regulators.
Apart from controlling shoot development, epigenetic marks such as H3K27me3 and
DNA methylation are often reset during major developmental phase transitions, such
as gamete and embryo formation (Slotkin *et al*., 2009; Iwasaki and Paszkowski,
2014; Borg *et al*., 2020). Such a reset helps demarcate embryonic and meristematic
programs in growing shoots while maintaining a pool of undifferentiated cells at the
meristem (Kuwabara and Gruissem, 2014; Pikaard and Scheid, 2014).

An epigenetic reset essential for normal embryo development in *Arabidopsis thaliana*
is mediated by the components of FERTILIZATION INDEPENDENT SEED-
Polycomb Repressive Complex 2 (FIS-PRC2) and DNA glycosylase DEMETER
(DME) (Xiao *et al*., 2003; Gehring *et al*., 2006). DME and members of FIS-PRC2
complex like MEDEA (MEA) are primarily expressed in central cells of the embryo
sac (Choi *et al*., 2002; Luo *et al*., 2000). The loss-of-function mutations in these

genes result in female gametophyte lethality reflected as parent-of-origin defects dur-
ing seed development. DME and MEA are known to regulate seed development to-
gether, where DME activates the expression of MEA in the central cell by removing
DNA methylation marks from *MEA* promoter and intergenic regions (Xiao *et al.*,
2003; Gehring *et al.*, 2006). Apart from gametogenesis and seed development, his-
tone methylation mediated by FIS-PRC2 also shapes seedling transitions by modu-
lating seed dormancy and repression of embryonic traits in developing seedlings
(Bouyer *et al.*, 2011).

Similarly, DNA methylation is reported to be high in developing seedlings to sup-
press transposon element expression and maintain an undifferentiated stem cell
niche (Bouyer *et al.*, 2017; Higo *et al.*, 2020). These reports indicate the vital role of
DNA and histone methylation in shaping plant architecture, with PRC2 members like
MEA and DNA glycosylases like DME likely to participate in transcriptional regula-
tion. Thus, a complete loss-of-function of DME and MEA provides an ideal system
for examining their function during sporophytic development.

Mutant alleles of *DME* and *MEA* are transmitted through male gamete as micro-
sporogenesis and microgametogenesis proceeds unhampered in most such mu-
tants, but not through female lineage (Grossniklaus *et al.*, 1998; Choi *et al.*, 2002;
Köhler *et al.*, 2003). The generation of homozygous gametophytic lethal mutants
thus is challenged by conventional genetic approaches, which rely on self-pollination
and inbreeding of heterozygous single/double mutants to generate respective homo-
zygotes barring those alleles with the leaky transmission. Hence, it is difficult to deci-
pher the role of such gametophyte-lethal genes, if any, in sporophytic develop-
ment. Studies involving maternal lethals of *DME, MEA,* and their epigenetic regula-
tion in sporophytes thus relied either on examining heterozygotes (Roszak and Köh-
ler, 2011; Schoft *et al.*, 2011; Figueiredo *et al.*, 2015; Park *et al.*, 2017; Zhang *et al.*,
2018; Khouider *et al.*, 2021) or homozygous plants carrying a weak allele (Park *et al.*, 2017; Kim *et al.*, 2021; Simonini *et al.*, 2021; Williams *et al.*, 2022), thereby re-
stricting complete understanding of their function. Alternate methods to study DME
and MEA in sporophyte development have been used previously, including paternal
haploids from single mutants, conditional homozygotes, and homozygotic progeny of
transheterozygote mutants (Ravi *et al.*, 2014; Roy *et al.*, 2018; Simonini *et al.*, 2021;

Kim *et al.*, 2021). The extended roles of MEA and DME in sporophytes deciphered

using *mea* and *dme* single mutants prompted us to generate and investigate the spo-

rophytic phenotypes of *mea*; *dme* double homozygotes.

Here, we employed haploid genetics to generate viable *mea* $^{-/-}$;*dme* $^{-/-}$ doubled hap-

loid (DH) sporophytes. Using these DH mutant homozygotes, we show that MEA and

DME are essential for the later sporophyte development of *Arabidopsis thaliana*

plants. Loss of function of MEA and DME leads to the emergence of more than one

functional shoot meristem at the apical pole of the plant body axis. We further show

that these epigenetic regulators control the shoot meristem development by regulat-

ing key shoot-promoting factors such as *STM*, *PLT5*, and *CUC2*. Our studies reveal

a new role of MEA and DME mediated epigenetic regulation in controlling shoot ar-

chitecture during post-embryonic development.

Results

***mea*-1 $^{-/-}$;*dme*-2 $^{-/-}$ double mutants are viable and initiate functional ectopic SAM**

Previous studies have established that MEA and DME play a key role in seed devel-

opment by regulating polar nuclei and endosperm division after double fertilization

(Grossniklaus *et al.* 1998; Choi *et al.* 2002). Several *dme* and *mea* alleles have been

isolated by induced mutagenesis from some select natural *Arabidopsis thaliana* ac-

cessions (Choi *et al.* 2002; Guitton *et al.* 2004; Schoft *et al.* 2011). Irrespective of the

accession, both *dme* and *mea* mutant alleles can be transmitted through pollen

(male) with rare or no maternal female transmission resulting in >25% seed lethality

upon self-fertilization of the corresponding heterozygotes. We have earlier shown

that generating *mea* and *dme* paternal haploid plants is possible using CenH3-

mediated haploid induction (Ravi *et al.*, 2014; Roy *et al.*, 2018). Here, we advanced

further to generate doubled haploid(DH) plants of *dme*-2 $^{-/-}$ (L*er*), *dme*-6 $^{-/-}$, *dme*-7 $^{-/-}$

(Col-0), and *mea*-1 $^{-/-}$ (L*er*), *mea*-6 $^{-/-}$, *mea*-7 $^{-/-}$ (C24) mutants to examine the recessive

vegetative phenotypes associated with *dme* and *mea* mutants (Figure 1A, Supple-

mentary Table 1). Here, we used *dme*-2 and *mea*-1 alleles for generating double

homozygotes for the following reasons; firstly, among all the mutant alleles charac-

terized, *mea*-1 and *dme*-2 are reported to show strong loss-of-function phenotypes

(Grossniklaus *et al.*, 1998; Choi *et al.*, 2002; Simonini *et al.*, 2021; Kim *et al.*, 2021),

which is ideal for double mutant studies. Secondly, both alleles are of the Landsberg
erecta (L*er*) ecotype, which yields a high number of spontaneous DH seeds than Co-
lumbia (Col-0) (Ravi and Chan, 2010). This feature is essential to maximize seed
yield to obtain viable spontaneous doubled haploid seeds in the strong gametophyte
lethal alleles with high seed abortion like *mea*-1 and *dme*-2. Thirdly, identical L*er*
backgrounds of single mutants of *mea* and *dme* ensure genetic homogeneity in the
resultant double mutants, thereby avoiding confounding effects due to different ge-
netic backgrounds.

mea-1 haploids, when germinated in Murashige and Skoog agar (MSA) plates,
showed no visible phenotypic defects and grew normal as that of their wild-type sib-
lings. *mea*-1 haploids produced a good number of spontaneous DH seeds, of which
2% (n= 21/1178, pooled from 14 haploid plants) were viable, giving rise to *mea*-1$^{-/-}$
DH offspring. Most seeds produced by DH *mea*-1$^{-/-}$ plants were atrophied (81%, n=
1407/1754, supplementary figure 1) in confirmation with previous reports (Simonini
et al., 2021). A small fraction of germinated seedlings (4%, n= 14/338) failed to grow
beyond the cotyledonary leaf stage and perished two to four days post-germination
(dpg). Unlike *mea*-1 haploids, a fraction of *mea*-1$^{-/-}$ (33%, n= 107/324) displayed
stunted and aberrant vegetative growth (Supplementary Figure 2). However, when
transferred to the soil medium, these seedlings recovered and showed normal vege-
tative growth with no discernible phenotypes.

Like *mea*-1 haploids, *dme*-2 haploids also showed normal vegetative growth but se-
vere seed abortion during reproductive development (98%, n= 1286/1308), owing to
the strong loss-of-function of *dme*-2. Spontaneous *dme*-2$^{-/-}$ DH seeds harvested from
dme-2 haploid plants showed less than 2% (n= 22/1308) germination on the MSA
medium. About 10% (n= 43/478) of *dme*-2$^{-/-}$ seedlings ceased to grow beyond the
cotyledon leaf stage (Supplementary Figure 1D). Unlike *mea*-1$^{-/-}$ seedlings, a vast
majority of *dme*-2$^{-/-}$ seedlings (78%; n= 341/435) exhibited developmental abnormali-
ties such as fused cotyledons, delayed emergence of true leaves, stunted hypocotyl,
and retarded root growth (Supplementary Figure 2). Despite these abnormalities,
most of the *dme*-2$^{-/-}$ seedlings (n= 153/157) recovered from the early vegetative
growth defects when they transitioned to the adult stage, similar to that observed for
mea-1$^{-/-}$ homozygotes. However, unlike *mea*-1$^{-/-}$ plants, *dme*-2$^{-/-}$ plants displayed flo-

ral defects like improper petal number and positioning, and abnormalities in gynoeci-
um, as reported in earlier studies (Supplementary Figure 3, Ravi *et al.*, 2014; Kim *et*
al., 2021).

Next, we attempted to generate *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ double homozygotes to examine
the loss of histone methylation (H3K27me3) and DNA demethylation mediated by
MEA and DME, respectively, during early vegetative development of seedlings. To
this end, we crossed *mea*-1$^{+/-}$;*dme*-2$^{+/-}$ as a pollen parent, with haploid inducer GFP-
tailswap as female, and selected *mea*-1;*dme*-2 haploids(n=9/32) by PCR-based
genotyping (Figure 1A, Supplementary Figure 1).

The spontaneous DH seeds obtained from *mea*-1;*dme*-2 haploid plants showed less
than 1% (n= 13/1950) viability up to germination (Supplementary Figure 1). The
mea-1;*dme*-2 haploids showed normal vegetative development as that of *mea*-1 and
dme-2 haploids. The *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ offspring were grown further for bulk seed
production. The selfed *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ plants continued to produce atrophied seeds
(n= 1678/1681), as observed in the parent generation. Around 38% (n= 87/226) of
germinated *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ were arrested in development and displayed pale and
fused cotyledons (Supplementary Figure 2). The remaining viable *mea*-1$^{-/-}$;*dme*-2$^{-/-}$
seedlings displayed severely stunted growth and shorter hypocotyl compared to wild-
type (Figure 1B). Such seedlings also showed underdeveloped and fused cotyledons
without petioles compared to WT (Figure 1C-E). We also observed smaller, asym-
metric true leaves, indicating defects during shoot development (n= 107/139). Also,
nearly half the number of *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings (n= 71/139) displayed ectopic
shoot emerging from the existing shoot apex (Figure 1F, G). Further microscopic
analysis of such *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ plants shoot apices revealed the presence of two
SAMs consistent with the presence of ectopic shoots (n= 25/48). Figure 1L, M). Like
mea-1$^{-/-}$ and *dme*-2$^{-/-}$ single mutant plants, *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ plants also recovered
from their early developmental defects and displayed normal vegetative growth in the
later stages (n= 129/132, Figure 1I, Supplementary Figure 3). In *mea*-1$^{-/-}$;*dme*-2$^{-/-}$
plants with ectopic shoot, both SAMs developed independently, which upon vegeta-
tive to reproductive transition, gave rise to two inflorescence shoots from a single
vegetative body (rosette) (Figure 1I, K, Supplementary Figure 3). This "twin-plant"
phenotype is reminiscent of conjoined twins having two independent upper trunks

with a common lower portion of the body. The "twin plant" phenotype arising from

persistent ectopic shoots suggests a role for DME and MEA in regulating shoot meri-

stematic activity, as reported previously (Kim *et al.* 2021; Simonini *et al.* 2021).

Interestingly, we note that a fraction of single mutant *mea*-1, *dme*-2, and *mea*-1;*dme*-

2 double mutants show no visible developmental defects and display WT phenotype.

Even single and double mutant seedlings that show early developmental defects

post-germination show resilience at later stages of vegetative development. Howev-

er, the ectopic shoots initiated during early developmental stages persist throughout

the plant life cycle, displaying twin plant phenotype. This observation suggests that

the function of both DME and MEA is predominantly required only during juvenile

growth but dispensable for later vegetative growth.

MEA and DME are expressed in developing shoot meristems and leaf primor-

dia

To decipher if the aberrant vegetative phenotypes in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings re-

sult from the loss-of-function of MEA and DME, we sought to examine their expres-

sion patterns of *MEA* and *DME* in germinated seedlings. Recent reports have shown

that in addition to its gametophyte expression in the central cell of the female game-

tophyte and sperm cells of the pollen, *DME* is also expressed in sporophytic tissues

such as the developing SAMs, root apical meristems (RAMs) and inflorescence me-

ristems (IMs) (Kim *et al.*, 2008; Park *et al.*, 2017). Likewise, *MEA* is expressed in the

developing embryo till the torpedo stage (Vielle-Calzada *et al.*, 1999; Spillane *et al.*,

2007) and induced in sporophytic tissues on demand when challenged with patho-

gens (Roy *et al.*, 2018). Further, transcriptome data also indicates the expression of

MEA and DME mRNA in the seedlings and other sporophytic tissues (Schmid *et al.*,

2005). Besides these reports, a systematic expression analysis

of *DME* and *MEA* during early seedling development needs to be improved. The ec-

topic shoot phenotype in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings prompted us to examine the de-

tailed expression pattern of MEA and DME in early seedlings. Hence, we first at-

tempted real-time PCR to examine the transcript levels of *MEA* and *DME* mRNA. For

this, we extracted total RNA from 1, 3, 5, 7, and 10-day-old seedlings, respectively,

from *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ and wild-type (WT) seedlings as control. Here, we found high

expression of *DME* mRNA until 2 dpg, with a decline in expression after 3

dpg. *DME* mRNA expression again increases after 5 dpg with plant growth (Figure

2A). On the other hand, we observed *MEA* mRNA expression to be low in the early

stage of vegetative development but steadily increased over time till the 10 dpg

seedling stage.

Next, we examined the spatiotemporal expression pattern of DME and MEA during

early sporophytic growth by confocal microscopy of the WT embryos and seedlings

expressing *pMEA*::*gMEA*:*vYFP* and *pDME*::*gDME*:*vYFP* translational   fusions.

Both, *pMEA*::*gMEA*:*vYFP* and *pDME*::*gDME*:*vYFP* transgenes were functional and

rescued the *mea*-1 $^{-/-}$ and *dme*-2 $^{-/-}$ mutant phenotypes (Supplementary Figure 4,

Supplementary Table T1). The rescue in seed lethality and expression pattern sug-

gests that both the transgenes harbor necessary cis-regulatory sequences capable

of recapitulating the endogenous expression pattern of *MEA* and *DME*. We further

monitored   the   expression   patterns   of *MEA* and *DME* using *pMEA*::3X-

VENUS and *pDME*::3X-*VENUS* in developing embryos and shoots (Supplementary

Figure   4).   Consistent   with   the   previous   studies,   we   observed   ac-

tive *MEA* transcription in developing embryos (Vielle-Calzada *et al.*,1999; Spillane *et*

al., 2007). *pMEA-driven* VENUS reporter is visible until the globular stage embryo,

with scattered expression observed in the later embryonic stages (n= 33/45) (Figure

2B-F). On the other hand, we observed *DME* was observed to be expressed in all

developmental stages of the embryo, i.e., from the four-celled to the walking-stick

stage embryos (n= 66/71) (Figure 2G-K).

We further examined the expression pattern of *MEA* and *DME* in the shoot apex.

Previous studies have documented the expression of *DME* in developing leaves of

seedlings (Kim *et al.*, 2008; Park *et al.*, 2017; Jullien *et al.*, 2008). However, it is un-

clear from these studies where MEA and *DME* are expressed in the shoot meristem.

We, therefore, examined the expression patterns in shoot meristem during post-

embryonic development. *MEA* was predominantly expressed in the peripheral do-

main of the shoot meristem (Figure 2L). Its expression was also detected in develop-

ing leaf primordia (Figure 2M). As reported earlier, *DME* was expressed uniformly

throughout the shoot meristem (Figure 2N) (Kim *et al.*, 2008; Park *et al.*, 2017).

Like *MEA*, its expression was also detected in developing leaf primordia (Figure 2O).

Taken together, the expression patterns of *MEA* and *DME* corroborate their role in
maintaining shoot meristems.

MEA and DME repress the expression of shoot-promoting factors

We next examined if the ectopic shoot meristems in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ make function-
al stem cell niches. Toward this, we examined the expression *pWUS*::3X-
VENUS:*tWUS.* WUS is a marker for SAM stem cell niche (Mayer *et al.*, 1998;
Schoof *et al.*, 2000; Yadav *et al.*, 2013). Strikingly, *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings
showed ectopic *WUS* promoter activity at the shoot apex (n= 21/48, Figure 3G, G(i)
to G (iii)) as compared to WT seedlings where WUS-YFP is confined to one organiz-
ing center (Figure 3F). Ectopic WUS-YFP corroborates with shoot phenotype seen
in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$. We also observed ectopic WUS-YFP activity in developing coty-
ledons of *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ embryos and SAM (n= 26/78) (Figure 3D-E). This indi-
cates de novo organizing centers specified in the non-canonical regions in the em-
bryos, as reported earlier (Simonini *et al.*, 2021).

We next examined the expression of genes essential for lateral organ initiation. PIN1
expression is an indicator of auxin transport whose expression is instrumental in or-
gan initiation in the actively dividing SAM (Heisler *et al.*, 2005). PIN1-GFP was de-
tected in shoot meristem and in developing leaf primordia of ectopic shoots of *mea*-1^{-}
$^{/-}$;*dme*-2$^{-/-}$ mutant (n=29/55, Figure 3U). Ectopic transport of auxin suggests ectopic
initiation of aerial organs in SAM. As expected, PIN1-GFP was expressed in wild-
type and mutant embryos (Figure 3H-S). Similar to WUS, we found ectopic PIN1 ex-
pression in the mutant embryos (Figure 3L-S).

We further analyzed the expression of genes that regulate the boundary between the
meristem and developing lateral organ primordia. The transcription factor, CUC2, is
an early marker of shoot organ boundary specification at SAM (Heisler *et al.*, 2005).
CUC2 expression in developing embryo displays characteristic stripe formation in the
central domain away from protoderm cells till the heart shape stage but later sur-
rounds the SAM between its boundary with cotyledons (Aida *et al.*, 1999; Aida *et al.*,
2002) (Figure 3V, W). For *pCUC2*::*CUC2*:*vYFP*; *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ embryos, the typi-
cal stripe pattern of CUC2-vYFP in the central domain is lost, and we observed more
dispersed expression in SAM initial and protoderm cells (n= 31/77). (Figure 3Z, A').

In a few embryos, we observed CUC2-vYFP expression in the cotyledon, defining a

new region for organ initiation compared to wild-type (n=15/77) (Figure 3B', C').

Similarly, CUC2-vYFP expression in *mea*-1$^{-/-}$*;dme*-2$^{-/-}$ shoots is dispersed along the

peripheral and central zone instead of being restricted at the organ primordia (Figure

3E'). The ectopic expression of CUC2-vYFP indicates the de-repression of *CUC2* in

SAM, which is essential for proper organ initiation and patterning (Burian *et al.*,

2015). Corroborating with the mutant phenotype, we also observed an increase in

the   transcript   levels   of   a   variety   of   shoot   regulators   such   as

CUC2, CUC3, YUC4, PLT5, STM, KNAT1, and *KNAT6* in *mea*-1$^{-/-}$*;dme*-2$^{-/-}$ seedlings

(Figure 3A). These data demonstrate the de-repression of most canonical shoot reg-

ulators in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings.

Shoot regulators are targets of MEA and DME during early shoot development

The ectopic shoot phenotype in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ plants, coupled with the expression

of *DME* and *MEA* in the early sporophyte, prompted us to identify critical targets of

MEA and DME that regulate the shoot meristem specification. The interplay between

DNA methylation and histone modifications in specifying cell fate and differentiation

during development is well-known in animals and plants (Du *et al.*, 2015; Williams

and Gehring, 2020). For instance, a positive feedback loop between DNA methyla-

tion and H3K9 methylation maintains the heterochromatin in a repressed state in

plants (Du *et al.*, 2015). H3K27 methylation, similarly, has been observed predomi-

nantly in the heterochromatin state associated with non-CG methylation. The prefer-

ential binding of the histone demethylase RELATIVE OF EARLY FLOWERING 6

(REF6) to hypo-methylated DNA leads to the loss of the H3K27me3 mark from the

chromatin (Cui *et al.*, 2016; Qiu *et al.*, 2019). Hence, we hypothesized that a similar

interaction between DNA methylation and histone methylation could be jointly medi-

ated by the action of DME and MEA, leading to the regulation of the shoot regula-

tors.

As a histone methyltransferase, MEA catalyzes the trimethylation of H3K27. Hence,

to identify targets of MEA, we performed whole genome profiling of H3K27me3 lev-

els by Chromatin Immunoprecipitation (ChIP)-seq experiment. Here, we observed

4994 significant peaks in WT compared to *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings. Out of anno-

tated 3885 H3K27me3 binding loci, we selected 431 enriched gene loci for further

analysis based on their role in shoot development as predicted by the Gene Ontolo-

gy Resource (http://geneontology.org/)(Figure 4A). These 431 genes encompass

known shoot regulators like *STM*, *KNAT1*, *CUC2*, *PLT5*, *ESR1* & *2*, *CKX3*, *AHP6*,

KNU, and *YUC4*. Upon closer examination, we found H3K27me3 binding more per-

sistent in the promoter region (90.4%) of these genes, as reported by earlier studies

(Zhang *et al.*, 2007; Xiao *et al.*, 2016). We also observed differential H3K27me3

binding in the gene body region of *CUC2*, *PLT5*, *STM*, and *WUS* (Figure 4B-E). This

loss of repressive H3K27me3 mark in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ for such genes predicts ac-

tive and ectopic transcription. We next examined the expression levels of potential

shoot regulators identified genome-wide in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ mutant seedlings by RT-

qPCR. Regulators of organ initiation, like *CUC3*, *YUC4*, and *ESR2,* were upregulat-

ed in the mutant seedlings compared to WT plants (Figure 4A). Similarly, genes in-

volved in stem cell maintenance, like *STM*, *KNAT1*, and *KNAT6*, were upregulated

compared to WT (Figure 4A). Overall, we see a de-repression of various shoot regu-

lators in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings. This observation is consistent with the loss of

repressive H3K27me3 mark in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ plants (Figure 3A).

To identify the likely targets of DME, we carried out Chop PCR for particular genes

that affect shoot development (Zhang *et al.*, 2014). Chop-PCR employs methylation-

sensitive restriction enzymes (MSRE) to differentially digest methylated and un-

methylated DNA, followed by PCR of the digested product. Here, we used McrBC,

which recognizes at least two methylated cytosines, preceded by purines: A or G, in

the range of 40-3000 bp to cleave 30bp aside of methylated cytosine (Sutherland *et*

al., 1992). Therefore, a heavily methylated DNA region will be less amplified upon

McrBC-digested chop-PCR (Dasgupta and Chaudhari, 2019). We chose upstream

regions of genes like *WUS* and *STM* for Chop-PCR. Based on available methylome

data (1001 Genome Project; http://neomorph.salk.edu/1001.aj.php) and their known

role in shoot development, we examined their DNA methylation pattern in *mea*-1 $^{-/-}$

;*dme*-2 $^{-/-}$ seedlings. We identified DNA methylation patches in the upstream region

of both genes and designed gene-specific $^{\text{oligonucleotides}}$ to amplify those regions after

McrBC   digestion   (Figure   4F,   G). We   observed   weak   amplification

for *WUS* and *STM* promoter regions in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings, indicating methyl-

ated DNA region (Figure 4H). Change in DNA methylation pattern may be associat-

ed with the differential expression pattern of these genes in the *mea*-1[-/-];*dme*-2[-/-] de-

veloping seedlings.

377 Downregulation of shoot regulators in *mea*-1[-/-]*;dme*-2[-/-] leads to a partial res-

378 cue of the ectopic shoot phenotype

Since the majority of shoot regulators are over-expressed in the SAM of *mea*-1[-/-]

;dme-2[-/-] seedlings, we sought if down-regulation of select shoot regulators viz.,

STM, PLT5, and CUC2 would suppress the ectopic shoot formation. Our selection of

these genes for further analysis is based on previously implicated role in promoting

the shoot meristem formation. While CUC2 and STM were earlier shown to be suffi-

cient for ectopic shoot meristem initiation (Daimon *et al.*, 2003; Scofield *et al.*, 2008),

we found inducible PLT5 over-expression sufficient for triggering ectopic cell prolifer-

ation in the shoot (Supplementary Figure 6).To examine the consequence of down-

regulating STM, PLT5, and CUC2 in the *mea*-1[-/-]*;dme*-2[-/-], we performed estradiol-

inducible RNAi-mediated gene silencing. Upon estradiol-induced RNAi, *STM*

mRNA levels reduced from 5.42 fold to 0.71 fold in *STM-dsRNAi; mea*-1[-/-]*;dme*-2[-/-]

seedlings vs. WT seedlings (Figure 5A). Similarly, *CUC2-dsRNAi* and *PLT5-*

dsRNAi constructs decreased *CUC2* and *PLT5 mRNA* levels from 2.52 and 3.54 fold

to 0.93 and 0.6 fold, respectively (Figure 5A). The reduction in mRNA levels

of *STM*, *CUC2*, and *PLT5* correlated with the reduction of ectopic shoots in *mea*-1[-/-]

;dme-2[-/-] plants (Figure 5B). Here, estradiol-induced downregulation of *STM* in *mea*-

1[-/-]*;dme*-2[-/-] germinating seedlings reduced the frequency of ectopic shoot from 49%

(N=4, n= 68/139, uninduced mock-treated seedlings) to 9% (N=4, n= 9/98, estradiol-

induced seedlings), suggesting a partial rescue to WT phenotype (N=4, n= 89/98,

Figure 5B, C). Similarly, RNAi-mediated knock-down of *CUC2* and *PLT5*

mRNA in *mea*-1[-/-]*;dme*-2[-/-] germinating seedlings resulted in the reduction of ectopic

shoots to 23% (N=3, n= 32/134) and 12% (N=4, n= 28/231), respectively (Figure 5B,

Supplementary Figure 5). We interpret the partial rescue of ectopic shoot phenotype

to WT in these RNAi lines as an indicator of multiple shoot regulators regulated by

MEA and DME, as observed in their expression pattern (Figure 3A). Together, our

results suggest that MEA and DME act synergistically to regulate early shoot devel-

opment by repressing ectopic induction of shoot-promoting factors.

406 Discussion

407   The histone methyltransferase, MEDEA, and the DNA glycosylase, DEME-
408 TER, were initially identified as paternally imprinted genes expressed in the central
409 cell of the female gametophyte. Both act synergistically to restrict the over-
410 proliferation of the central cell in the absence of double fertilization (Moreno-Romero
411 *et al.*, 2016). Ovules inheriting the strong mutant alleles of both *mea* and *dme* abort
412 seed development, making it impossible to recover homozygotes of either mutant.
413 The maternal lethality of the *mea* and *dme* alleles makes it challenging to examine
414 their roles beyond the gametophyte stage. However, recent studies have successful-
415 ly implemented various genetic strategies to obtain viable, single mu-
416 tant, *mea,* and *dme* homozygotes (Ravi *et al.*, 2014; Pires *et al.*, 2016; Kim *et al.*,
417 2021; Williams *et al.*, 2022). These single mutants have helped demonstrate the role
418 of DME and MEA in various phases of sporophyte development, in addition to its pri-
419 or established role in the gametophytes. For instance, MEA is required for proper
420 embryonic patterning by regulating cell division via D-type cyclin 1.1 (CYCD1;1) (Si-
421 monini *et al.*, 2021). MEA also facilitates *P. syringae* infection in *Arabidopsis* by
422 downregulating RESISTANCE TO P. SYRINGAE2 (RPS2) (Roy *et al.*, 2018). Like
423 MEA, DME also has a sporophytic role in controlling cell division rate in sporophytic
424 tissues like root hair and stomata, the maintenance of inflorescence meristem, and
425 *de novo* shoot regeneration (Kim *et al.*, 2021). DME, along with its somatic DNA de-
426 methylating counterparts, is also shown to regulate flowering time and inflorescence
427 meristem activity (Williams *et al.*, 2022; Kim *et al.*, 2021). Like MEA, DME also re-
428 stricts pathogen infection by repressing transposable elements controlling disease-
429 related proteins in *Arabidopsis* (Schumann *et al.*, 2019). In contrast to the docu-
430 mented single mutant phenotypes, *mea*-1$^{-/-}$*;dme*-2$^{-/-}$ plants show a unique presence
431 of twin shoots arising from seed with a single embryo suggesting the interplay of
432 both MEA and DME during seed to the seedling transition phase of sporophyte de-
433 velopment.

434   The rescue of *mea*-1$^{-/-}$*;dme*-2$^{-/-}$ haploid sporophytes using CenH3-
435 mediated genome elimination as a tool facilitated us to generate a population
436 of *mea*-1$^{-/-}$*;dme*-2$^{-/-}$ DH plants which in turn shed light on the synergistic role of MEA
437 and DME in suppressing the activation of ectopic shoot meristems during early spo-
438 rophyte development. Shoot developmental processes like regeneration and floral
439 transitions are known to be restricted by the repressive histone and DNA methylation

marks (Li *et al.*, 2011; Sun *et al.*, 2014). However, only a few studies have reported
the involvement of such epigenetic players during early sporophyte development
(Bouyer *et al.*, 2011). Seed to seedling transition is controlled by FIE, a subunit of the
PRC2 complex (Bouyer *et al.*, 2011). Here we propose that, during sporophytic de-
velopment, MEA, another subunit of the PRC2 complex, and DME restrict the for-
mation of a single meristem at the apical pole by repressing shoot regulators. Con-
sistent with this hypothesis, we show that an array of shoot regulators gets upregu-
lated than WT levels in the *mea*-1$^{-/-}$*;dme*-2$^{-/-}$ mutants. A closer inspection of the epi-
genetic signatures (DNA methylation) near selected shoot regulators revealed per-
turbance from the normal levels (Figure 4). The loss of H3K27me3 and gain of DNA
methylation is likely sufficient to trigger a change in the chromatin state near these
genes, causing a de-repression of the genes controlling shoot development early in
the plant life cycle. Restoration of wild-type shoot apex by down-regulation of a se-
lect few shoot-promoting factors in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ mutant strengthen this mecha-
nistic framework. Likely, DME- and MEA-mediated epigenetic suppression of shoot-
promoting factors such STM, PLT5, and CUC2 restrict the coordinated interactions
between several fundamental cellular processes such as cell division and differentia-
tion at the apical pole of the plant's body axis during post-embryonic development.

In conclusion, we demonstrate an extended synergistic role for the gameto-
phytic epigenetic regulators DME and MEA during the seed-to-seedling transition of
sporophyte development. Our study provides the basic framework for understanding
the role of gametophytic lethals in sporophytic tissue development. Further studies
utilizing haploid induction for generating homozygous mutants may offer more in-
sights into plant growth development.

Figure 1. Generation and phenotypic examination of *mea*-1 $^{-/-}$;*dme*-2 $^{-/-}$ plants

(A) Schematic diagram illustrating rescue of *mea*-1$^{-/-}$;*dme*-2$^{-/-}$plants using CENH3-

mediated genome elimination. **(B,C)** Phenotype of *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings as

compared to WT at 2- leaf (5 dpg) and 6-leaf (10 dpg) stage respectively. Repre-

sentative images of L*er* WT plants at **(D)** 2-leaf (5dpg), **(F)** 6-leaf (10 dpg), **(H)** inflo-

rescence emergence (21 dpg) and **(J)** flowering (28 dpg) stage as compared with

mea-1$^{-/-}$;*dme*-2$^{-/-}$ plants with ectopic shoot at **(E)** 2-leaf , **(G)** 6-leaf, **(I)** inflorescence

emergence and **(K)** flowering stages.

Longitudinal section of shoot apex of the 3-leaf stage (7 dpg) **(L)** L*er* WT and **(M)**

mea-1$^{-/-}$;*dme*-2$^{-/-}$ seedling. Yellow arrowhead is pointing at the shoot apex. Scale bar:

(B-G)= 1 mm; **(H-K)**= 5 mm; **(L-M)**= 20 µm. dpg: days post germination.

Figure 2. Expression analysis of MEA and DME in sporophytic tissues (A) Relative abundance of MEA and DME mRNA in developing seedlings. *DME* mRNA levels are abundant in 1- and 2-dpg seedlings with sharp decrease in its level on 3- and 5-dpg (1-dpg: 7.62±2.37, **p= 0.0016; 2-dpg: 7.69±0.30, ****p= 0.000003; 3-dpg:

481 1.15±0.13, ^{ns}p= 0.11; 5-dpg: 2.06±0.20, **p= 0.003; 7-dpg: 3.28±0.29, ***p=0.00084;

482 10-dpg: 5.79±0.99, **p= 0.0033). *MEA* mRNA levels gradually increased as the

483 seedling transitioned from 1- to 10-dpg (1-dpg: 2.76±0.69, *p= 0.011; 2-dpg:

484 6.12±0.89, **p= 0.0023; 3-dpg: 4.27±0.76, **p= 0.0034; 5-dpg: 3.97±0.62, **p=

485 0.0033; 7-dpg: 15.66±1.08, ****p=0.0001; 10-dpg: 51.96±5.12, ***p= 0.00033). Sta-

486 tistical test: Multiple unpaired *t*-tests (Holm-Sidak method), with alpha = 0.05. Error

487 bar: SD.

488 **(B-O)** Representative images of embryos and shoot apices showing expression pat-

489 terns of **(B-F, L,N)** *pMEA*::3X- *VENUS* and **(G-K, M,O)** *pDME::*3X-*VENUS.*

490 *pMEA*::3X- *VENUS* expression (yellow) in developing embryo **(B)** octant, **(C)** globu-

491 lar, **(D)** early heart-shape **(E,F)** heart-shape **(L)** SAM and in **(N)** shoot apex.

492 *pDME*::3X- *VENUS* expression (yellow) in developing embryo **(G)** octant, **(H)**

493 dermatogen **(I)** globular, **(J)** heart-shape **(K)** late heart-shape **(M)** SAM and **(O)** shoot

494 apex. SAM- shoot apical meristem, LP-leaf primordium, TL- true leaf. Scale bar: **(B-**

495 **K)**= 10 µm; **(L-O)**= 20 µm.

Figure 3. Deregulated expression pattern of shoot regulators in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ embryos and seedling (A) Relative abundance of shoot regulators in 2-3 dpg old *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings compared with WT (fold change). *AHP6*: 3.78±0.24, ***p=0.0011; *CLE2*: 4.43±0.78, *p=0.019; *CLV1*: 1.08±0.22, ^{ns}p=0.571; *CUC2*: 2.52±0.87, **p=0.014; *CUC3*: 9.18±1.54, ***p=0.01; *ESR1*: 3.15±0.19, ***p=0.0006; *ESR2*: 9.02±0.58, ***p=0.0003; *KNAT1*: 2.30±0.19, ****p=0.00002; *KNAT6*: 2.35±0.59, *p=0.016; *PIN1*: 3.31±0.51, *p=0.019; *PIN2*: 2.47±0.53, *p=0.049; *PLT1*: 1.26±0.19, ^{ns}p=0.13; *PLT3*: 5.71±1.14, *p=0.021; *PLT5*: 2.73±0.82, *p=0.021; *PLT7*: 5.71±1.19, *p=0.021; *STM*: 3.74±0.65, *p=0.021; *WUS*: 0.26±0.22, *p=0.03; *YUC1*: 2.49±0.49, *p=0.042; *YUC4*: 7.32±1.03, **p=0.0067. N=3, statistical test: Multiple unpaired *t*-tests (Holm-Sidak method), with alpha = 0.05. Error bar: SD.

(B-E') Representative images of embryos and shoot apices showing expression pattern of **(B-H3)** *pWUS*::3X-*VENUS,* **(H-U)** PIN1-GFP and **(V-E')** CUC2-vYFP shoot regulators.

pWUS::3X-*VENUS* expression (yellow) in organizing center of embryos of **(B-C)** wild-type and **(D-E)** *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ *(*n= 26/78) showing ectopic organizing center in cotyledon (marked with yellow arrowhead). *pWUS*::3X-VENUS is marking center of shoot meristem in **(F)** wild-type, inset is showing organizing center in SAM and **(G to G(iii))** *mea*-1$^{-/-}$;*dme*-2$^{-/-}$,along with **(G(i))** primary shoot meristem **(G(ii))** axillary meristem and **(G(iii))** ectopic shoot meristem, marked with blue arrowhead (n=21/48). The meristem dome (SAM) and leaf primordia (LP) are labeled and indicated with a white dashed line. The scale bar represents 20 μm in **(B-F)** and **(G(i) to G(iii))**, and 100 μm in **(G)**.

(H-U) PIN1-GFP (green) expression in embryos of **(H-K)** wild-type and **(L-S)** *mea*-1 $^-$

$^{/-}$;*dme*-2 $^{-/-}$ showing ectopic PIN1 expression. Dashed lines outline the embryo shape.

PIN1-GFP expression in shoot meristem of **(T)** wild-type **(U)** *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seed-

ling showed PIN1 expression in ectopic shoot along with primary shoot (n=29/55)

along with red as autofluorescence. The scale bar represents 20 µm in **(H-S)** and100

µm in **(T-U)**.

(V-E') CUC2-vYFP (yellow) expression in embryos of **(V-Y)** wild-type and **(Z- C')**

mea-1$^{-/-}$;*dme*-2$^{-/-}$ showing ectopic expression at SAM and developing cotyledons (n=

31/77). CUC2-vYFP expression in shoot apex of **(D')** wild-type and **(E')** *mea*-1$^{-/-}$

;*dme*-2$^{-/-}$ showing ectopic expression in SAM apart from leaf primordia (n= 15/77).

The scale bar represents 20 µm. SAM- shoot apical meristem, AM- axillary meri-

stem, LP-leaf primordium.

Figure 4. Validation of targets of MEA and DME (A) Gene ontology (GO) of en-
riched genes in ChIP using αH3K27me3. **(B-E)** Graphical representation of ChIP-seq
using αH3K27me3 for **(B)** *STM*, **(C)** *CUC2* **(D)** *PLT5* and **(E)** *WUS* gene and their
upstream region. Pink boxes denote prominent enrichment in WT ChIP DNA as
compared to *mea*-1[-/-];*dme*-2[-/-] ChIP DNA. Input DNA is WT DNA without αH3K27me3
treatment. **(F-G)** Cartoon representation of *WUS* and *STM* upstream regions show-
ing putative methylated regions (yellow box) and corresponding forward (red) and
reverse (green) oligonucleotides used for Chop PCR. **(H)** Lower amplification in
McrBC-chop PCR of hypermethylated regions of *WUS*, and *STM* upstream regions

544   in *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ as compared to hypomethylated WT. Black line: upstream region,

545   gray line: UTR, blue box: exon, blue dotted line: intron.

Figure 5. RNAi mediated downregulation of select target genes. (A) Relative abundance of *CUC2*, *PLT5*, and *STM* transcripts in respective RNAi lines. *CUC2*: WT- 1±0, *mea*-1$^{-/-}$;*dme*-2$^{-/-}$- 2.52±0.87, *CUC2*-dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$- 0.94±0.24, *p=0.038; *PLT5*: WT- 1±0, *mea*-1$^{-/-}$;*dme*-2$^{-/-}$- 3.54±0.36, *PLT5*-dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$- 0.60±0.08, ***p=0.00048; *STM*: WT- 1±0, *mea*-1$^{-/-}$;*dme*-2 $^{-/-}$- 5.42±0.75, *STM*-

dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$- 0.71±0.15, ***p=0.0008. Statistical test: Multiple unpaired

t-tests (Holm-Sidak method), with alpha = 0.05. Error bar: SD.

(B) Quantification of shoot number in *CUC2*-, *PLT5*- and *STM*-RNAi lines as com-

pared to WT and *mea*-1$^{-/-}$;*dme*-2$^{-/-}$. WT: n= 161 (Normal shoot= 161, ectopic shoot=

0); *mea*-1$^{-/-}$;*dme*-2$^{-/-}$: n= 139 (Normal shoot= 71, ectopic shoot= 68); *CUC2*-

dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$: n= 134 (Normal shoot= 102, ectopic shoot= 32, ****p<

0.0001); *PLT5*-dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$: n= 231 (Normal shoot= 203, ectopic shoot=

32, ****p< 0.0001); *STM*-dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$: n= 98 (Normal shoot= 89, ectopic

shoot= 9, ****p< 0.0001). Statistical analysis: two-way ANOVA (Sidak method), with

alpha = 0.05. Error bar: SEM.

(C) Rescue of ectopic shoot in estradiol-induced *STM*-dsRNAi;*mea*-1$^{-/-}$;*dme*-2$^{-/-}$

plants. Asterix(*) marks plants with ectopic shoots. White dotted box corresponds to

the zoomed plants shown in the below panel. Yellow arrowhead points to the shoot

apex. Scale bar: 1cm.

Materials and Methods

Plant materials and growth conditions

Arabidopsis thaliana seeds were surface sterilized by fumigation with Chlorine (Cl_2)
gas (Lindsey III *et al.*, 2017). Fumigated seeds were then plated onto Murashige and
Skoog (MS) growth medium with 0.8% agar and cold stratified at 4℃ in dark for four
days. After cold stratification, the plates were transferred to the growth chamber for
germination. Seedlings and plants in pots were grown under white fluorescent lights
(7000 lux at 20 cm) at 22 °C with a 16 h light/8 h dark cycle at 75% RH in a controlled
growth cabinet (Percival Inc.). Heterozygous *MEA*/*mea*-1 (Grossniklaus *et al.*, 1998)
and *DME*/*dme*-2 (Choi *et al.*, 2002) mutants were previously described. Following
heterozygous seeds were obtained from Arabidopsis Biological Resource Center:
MEA/*mea*-6 (CS6996), *MEA*/*mea*-7 (CS6997), *DME*/*dme*-6 (GK-252E03), and
DME/*dme*-7 (SALK107538).

Generation of maternal gametophytic lethal homozygotes

To generate *dme*-2 (AT5G04560) and *mea*-1 (AT1G02580) homozygous mutants,
heterozygous *DME*/*dme*-2 and *MEA*/*mea*-1 plants were crossed as the male to hap-
loid inducer GFP-*tailswap* plants as female (Ravi *et al.*, 2014). The resultant F1
seeds were collected and germinated on Murashige and Skoog agar plates and later
transferred to soil for further growth. Haploids were identified phenotypically and cy-
tologically as described earlier (Ravi and Bondada, 2016). All the haploids were PCR
genotyped to identify the *DME* WT allele using MR146 and MR147 oligo combina-
tions amplifying a 791 bp fragment. Similarly, the *dme*-2 mutant haploids were identi-
fied using MR147 and a left border-specific oligo MR148 producing 381 bp fragment.
For *MEDEA*, the WT allele was identified using oligonucleotides combinations MR8
and MR319 (722 bp), and the *mea*-1 allele using MR8 and a left border-specific oligo
MR10 (312 bp). The DNA sequences for all oligos used in this study is given in Sup-
plementary Table 4. The mutant haploids were screened for spontaneous mitot-
ic/meiotic doubling events for obtaining doubled haploid seeds which gives rise to
homozygous mutant diploid sporophytes. For generating homozygous *mea*-1;*dme*-2
double mutant in L*er* ecotype, *mea*-1 homozygous plants serving as female parent
were crossed with *dme*-2 heterozygous plants as male parent. The resulting progeny
PCR genotyped for both the *mea*-1 and *dme*-2 alleles, and the plants heterozygous

plant both *mea*-1 and *dme*-2 alleles were identified and used as male parent in a
cross with GFP-*tailswap* as female. *mea*-1;*dme*-2 haploid plants were identified us-
ing PCR-based genotyping. Spontaneous double haploid seeds from *mea*-1;*dme*-2
plants were collected to generate *mea*-1;*dme*-2 doubled haploid (diploid) sporo-
phytes used in this study.

Quantitative Real Time-PCR

For expression analysis of *DME* and *MEA* mRNA, we harvested shoot apices from 1,
3, 5, 7, and 10 dpg old WT seedlings and stored at -80℃ for further use. Shoot api-
ces from *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings were used as control. Similarly, for expression
analysis of selective shoot regulators and somatic epigenetic factors, we harvested
shoot apices from 2 to 3-day-old WT and *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings. Total RNA was
isolated using TRIzol^{TM} reagent as per the manufacturer's instructions. The extracted
RNA was treated with NEB(New England Biolabs) DNase I (RNase free) (M0303)
according to NEB guidelines to remove the genomic DNA contamination. One mi-
crogram of total RNA was used for complementary DNA (cDNA) synthesis by
PrimeScript first strand cDNA synthesis kit (TAKARA-Bio, Cat# 6110) using an oli-
go(dT) and random hexamer as oligonucleotides. qPCRs were performed on Mas-
tercycler BIORAD CFX96^{TM} using the oligonucleotides mentioned in Supplementary
Table T4. *ACTIN2* (*ACT2*) served as a reference gene control. The reactions were
carried out using TB Green® Premix Ex Taq™ II (TAKARA-Bio, Cat no. RR82WR)
and incubated at 95 °C for 2 minutes, followed by 40 cycles of 95 °C for 15 s and 60
°C for 30 s. PCR specificity was checked by melting curve analysis. *ACT2* was used
for the normalization for all the reactions. At least three independent biological sam-
ples were analyzed for each time point mentioned above, and qPCR reactions were
set up with three technical replicates for each biological replica. mRNA abundance of
target genes was analyzed using the normalized $2^{-\Delta\Delta Ct}$ (cycle threshold) method
(Livak and Schmittgen, 2001).

ChIP followed by Next-Generation Sequencing

To identify the targets of MEA protein and differentiate histone methylation, we per-
formed chromatin immunoprecipitation (ChIP) followed by next-generation sequenc-
ing as described previously (Salvi *et al.*, 2020; Yamaguchi *et al.*, 2014) on two inde-
pendent biological replicates. 500 mg of micro dissected shoot apices, devoid of cot-

yledons and hypocotyl, from 2 to 3-day-old WT and *mea*-1[-/-];*dme*-2[-/-] seedlings were

used for each sample. The samples were fixed in 1.8% formaldehyde in 1X PBS and

vacuum infiltrated for 15 mins at room temperature to cross-link the tissue. After fixa-

tion, formaldehyde solution was removed and replaced by ice-cold 0.125 M glycine

solution (in 1X PBS) followed by vacuum infiltration for 5 mins to quench the reac-

tion. The glycine solution was filtered away and samples were washed twice with ice-

cold 1X PBS. Samples were blotted on Kim wipe paper to remove excess 1X PBS.

Samples were frozen and powdered in liquid nitrogen. The powdered samples were

resuspended in nuclei extraction buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1%

Triton X-100, 0.1% sodium deoxycholate, 2.5 mM EDTA, 10% glycerol, freshly sup-

plemented with cOmplete™, EDTA-free protease inhibitor cocktail (Cat#

04693132001) and 1 mM PMSF (Sigma). After brief incubation of 15 minutes, sam-

ple solutions were filtered using Falcon® 40 µm cell strainer (Cat# 352340). Filtrate

was centrifuged at 4°C to pellet the nuclei. The pellet was resuspended in an ice-

cold nuclei lysis buffer (50 mM Tris-HCl, pH 8; 5 mM EDTA, pH 8; 0.5% SDS). The

solution was sonicated for 8 cycles of 10 seconds with 20 seconds gap at 39% am-

plitude (Branson Sonifier, Marshall Scientific). 1.1% triton was added to the sonicat-

ed sample and centrifuged to isolate the supernatant. Preclearing of the sample was

done by adding 20 µl of Dynabeads™ Protein G (Invitrogen, Cat# 10003D) and in-

cubated at 4 ℃ for 2h. An aliquot of the sample was stored as Input for the experi-

ment. The sample was incubated with mouse monoclonal antibody to H3K27me3

(ab6002; Abcam) overnight at 4℃, and dynabeads protein was added to immuno-

precipitate the DNA fragment for 5 hr at 4℃. Sequential washes were performed in

the beads using cold low salt wash buffer (150 mM NaCl, 0.2% SDS, 0.5% Triton X-

100, 2 mM EDTA pH 8, 20 mM Tris-HCl pH 8), cold high salt wash buffer (500 mM

NaCl, 0.2% SDS, 0.5% Triton X-100, 2 mM EDTA pH 8, 20 mM Tris-HCl pH 8), cold

250mM LiCl wash buffer (0.25 M LiCl, 0.5% NP-40, 0.5% sodium deoxycholate, 1

mM EDTA pH 8, 10 mM Tris-HCl pH 8) and 0.5X TE. All washes were done twice.

Magnetic beads were collected and resuspended in TES (25 mM Tris-HCl pH 8, 5

mM EDTA pH 8, 0.5% SDS). 5 M NaCl was added to the sample and input control to

reverse the cross-links overnight at 65℃. DNA fragments were purified using a PCR

purification kit (Cat #28106, Qiagen). Library constructions were performed, and val-

idated using Qubit and Agilent TapeStation, followed by 150bp paired-end sequenc-
ing on Illumina NovaSeq 6000 by miBiome Therapeutics LLP, Mumbai, India.
The trimmed reads were aligned to reference genome TAIR10, using BWA 0.7.17-
r1188 (Li, 2013). The quality check of the alignments was done using Qualimap
2.2.2-dev (Okonechnikov *et al.*, 2015). Duplicated reads, reads with low mapping
quality, and mitochondrial and plastid reads were identified and removed with
Samtools version 1.2 (Li *et al.*, 2009). Enriched intervals between the IP sample and
its relative Input were identified by MACS2 (version 2.0.10) program (Zhang *et al.*,
2008) with the default setting. The big-Wig files were generated from the BAM files
using Deeptools 3.5.1 (Ramirez *et al.*, 2016). The coverage is calculated as the
number of reads per bin and normalized to 1 x sequencing depth (reads per genome
coverage, RPGC). The resulting bigWig files were visualized in Integrated Genome
Viewer (https://igv.org/app/)(Robinson *et al.*, 2011) to observe the protein binding on
the chromatin. Peak annotation and GO and KEGG analysis for the target regions
were done using ChIPseeker 1.32.1 (Yu *et al.*, 2015) and ClusterProfiler 4.4.4 (Yu *et*
al., 2012).
684

Chop-PCR assay

Genomic DNA from 2 to 3 dpg old WT and *mea*-1$^{-/-}$;*dme*-2$^{-/-}$ seedlings was isolated
using the standard CTAB method. One microgram of DNA was digested with meth-
ylation-sensitive restriction enzyme McrBC (NEB, Cat# M0272) at a concentration of
10 units/µg of DNA template in a 30 µl reaction mix, as per manufacturer's instruc-
tion, for at least 6 h. After heat inactivation, 4 µl of the digested DNA was used as a
template in a 20 µl reaction volume for PCR for 25 cycles.
692

Microscopy and imaging

Bright-field images of seedlings were captured using Leica M205FA stereo micro-
scope. Confocal laser scanning microscopic images were acquired using Zeiss LSM
880 and Leica TCS SP5 II confocal laser-scanning microscope. The confocal imag-
ing was carried out on dissected embryos and 3 to 5 dpg old seedlings using Argon
laser source at 40% power. The excitation wavelengths were as follows: 488 nm- for
GFP and 514 nm- for YFP. Auto-fluorescence was captured at the wavelength of
633 nm. Images were captured with 40x (oil) or 63x (oil) objectives with 550-750
master gain, 100-200 range pinhole, and frame size of 1,024 by 1,024 pixels. Master

gain and the pinhole adjustment varied with samples. The line averaging was adjust-

ed to four and the digital zoom to 0.6. Z-stack images were taken in 5 μm or 1 μm

intervals. The whole plant images are captured using a Canon 70D SLR camera.

The images are edited in Adobe Photoshop CS4.

Plasmid construction, molecular cloning, and plant transformation

For cloning the gene and its regulatory sequences, the sequence of interest was

PCR amplified from total genomic DNA isolated by CTAB method. For cDNA cloning,

total RNA was isolated using TRIzol^{TM} reagent as per the manufacturer's instruc-

tions. One microgram of DNase-free RNA was used for cDNA synthesis by Prime-

Script 1st strand cDNA synthesis kit using an oligo(dT) and random hexamer as oli-

gonucleotides . Polymerase chain reaction (PCR) was carried out using PrimeSTAR$^{®}$

GXL DNA Polymerase (TAKARA-Bio, Cat# R050). The PCR conditions were specific

to each sequence and amplified using 10 μM sequence-specific paired oligonucleo-

tides (see Supplementary Table T4) using Applied Biosystems™ ProFlex™ 3 x 32-

well PCR System. All entry clones were propagated in DH5α strain of *E.coli* using a

single Gateway cloning system. To knock down the expression of *STM*, 272 bp exon

sequence was amplified from *cDNA* using region-specific oligonucleotides and

cloned in the p2R3E-As-OScT vector (sense) and p221Z-S-Intr vector (antisense)

using *Bam*HI and *Xho*I restriction digestion and ligation. All entry clones were propa-

gated in the DH5α strain of E.coli using a single gateway cloning system. The sense

and antisense for STM were cloned under *pG1090* inducible promoter (Zuo *et al.*,

2000) and subsequently in *pCAMBIA-based R4R3* destination vector. Similarly, the

PLT5 (299 bp) and *CUC2* (Varapparambath *et al.*, 2022) knockdown constructs were

cloned under *pG1090* inducible promoter to silence the transcript level of *PLT5* and

CUC2, respectively. To generate *pDME*::*3X-VENUS*:*3AT*, 5000bp upstream regula-

tory sequence of *DME* is tagged with *3X-VENUS* and *3AT* terminator. Similarly,

5142bp of *MEA* promoter was tagged with *3X-VENUS* to generate *pMEA*::*3X-*

VENUS:*3AT*. Translation fusion lines of DME (*pDME*::*gDME*:*vYFP*) and MEA

(*pMEA*::*gMEA*:*vYFP*) were generated using three fragments multisite recombination

gateway cloning system where the above-mentioned upstream regulatory sequences

of *DME* and *MEA* were cloned to drive *DME* (8303 bp) and *MEA* (4199 bp) genomic

sequences fused with *vYFP*.

Estradiol treatment for inducible gene knock-down

Inducible RNAi for *STM*, *PLT5*, and *CUC2* genes was carried out, as mentioned earlier (Varapparambath *et al.*, 2022). MS agar medium supplemented with 5μM β-estradiol was used to induce conditional gene silencing using *pG1090:XVE* artificial promoter throughout seed germination. Estradiol-treated and mock (DMSO) treated seedlings were scored for ectopic shoot emergence at 15 dpg.

Statistical analysis

The details of the statistical analysis followed in this study are described in the respective figure legends (number of samples (n), mean± S.E.M). The statistical significance levels were defined as $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. For mRNA quantitative analysis, statistical significance was determined using the Multiple unpaired *t*-tests (Holm-Sidak method), with alpha = 0.05. For RNAi phenotype quantitative analysis, statistical significance was determined using two-way ANOVA (Sidak method), with alpha = 0.05. All the statistical tests and analyses were performed using GraphPad PRISM (v9.5.1) statistical software.

Data Availability

The ChIP-sequencing data generated and analyzed during the current study are available from NCBI under the bio project accession PRJNA972183 consisting of six SRA accession identifiers: SRR24547737, SRR24547736, SRR24547733, SRR24547734, SRR24547735, and SRR24547732. All other source data are included in the article as Supplementary Table T3.

Acknowledgments

M.P.R. thanks to the IISER Thiruvananthapuram for providing a Ph.D. research fellowship. S.S. is a recipient of the Department of Science and Technology (DST) - INSPIRE scholarship for higher education (SHE). R.B. is the recipient of a Junior and Senior Research Fellowship (J.R.F. and S.R.F.) for a Ph.D. funded by the University Grants Commission (U.G.C.), Govt. of India. A.P.S. thanks to the Council of Scientific and Industrial Research (CSIR) for the Ph.D. fellowship. K.P. acknowledges grants from the Department of Biotechnology (D.B.T.), Government of India (BT/PR41931/BRB/10/1964/2021), the Science and Engineering Research Board-Scientific and Useful Profound Research Advancement (SERB-SUPRA), Govern-

770 ment of India (S.P.R./2021/000109). R.M. acknowledges financial support from Ra-
771 malingaswami's re-entry fellowship awarded by the Department of Biotechnology
772 (D.B.T., Govt. of India) and the Dupont Young Professor grant, Dupont U.S.A. We
773 thank the Indian Institute of Science Education and Research (IISER)- Thiruvanan-
774 thapuram for intramural financial and infrastructural support.
775

Author contributions

R.M. conceptualized, designed, and supervised the study. M.P.R. contributed to cer-
tain aspects of the study design and carried out most of the experiments. S.S. gen-
erated and performed the phenotypic assessment of RNAi lines along M.P.R. R.B.
helped generate double mutant lines. A.P.S. performed cloning experiments for pro-
ducing reporter lines. K.P. designed, supervised, and critically analyzed the study.
M.P.R., K.P., and R.M. wrote the manuscript. R.M. and K.P. acquired funds for the
study. All the authors read and approved the draft.

Declaration of interests

The authors declare no competing interests.

References

1. Aichinger, E., Villar, C.B.R., Di Mambro, R., Sabatini, S., and Köhler, C.
(2011). The CHD3 Chromatin Remodeler PICKLE and Polycomb Group Pro-
teins Antagonistically Regulate Meristem Activity in the *Arabidopsis* Root.
Plant   Cell   23:   1047–1060.
793
794 2. Aida, M., Ishida, T., and Tasaka, M. (1999). Shoot apical meristem and coty-
795   ledon formation during *Arabidopsis* embryogenesis: Interaction among the
796   *CUP-SHAPED COTYLEDON* and *SHOOT MERISTEMLESS* genes. Devel-
797   opment   126:   1563–1570.
798
3. Aida, M., Vernoux, T., Furutani, M., Traas, J., and Tasaka, M. (2002). Roles of
PIN-FORMED1 and MONOPTEROS in pattern formation of the apical region
of the *Arabidopsis* embryo. Development 129: 3965–3974.
4. Borg, M., Jacob, Y., Susaki, D. *et al.* (2020).Targeted reprogramming of
H3K27me3 resets epigenetic memory in plant paternal chromatin. Nat Cell
Biol   22:   621–629.
805

806   5. Bouyer, D., Kramdi, A., Kassam, M., Heese, M., Schnittger, A., Roudier, F.,
807   and Colot, V. (2017). DNA methylation dynamics during early plant life. Ge-
808   nome   Biol.   18:   1–12.
809

810   6. Bouyer, D., Roudier, F., Heese, M., Andersen, E.D., Gey, D., Nowack, M.K.,
811   Goodrich, J., Renou, J.P., Grini, P.E., Colot, V., and Schnittger, A. (2011).
812   Polycomb repressive complex 2 controls the embryo-to-seedling phase transi-
813   tion.   PLoS   Genet.   7.
814

815   7. Burian, A., Raczyńska-Szajgin, M., Borowska-Wykręt, D., Piatek, A., Aida, M.,
816   and Kwiatkowska, D. (2015). The *CUP-SHAPED COTYLEDON2* and *3* genes
817   have a post-meristematic effect on *Arabidopsis thaliana* phyllotaxis. Ann. Bot.
818   115:   807–820.
819

820   8. Chaudhury, A.M., Ming, L., Miller, C., Craig, S., Dennis, E.S., and Peacock,
821   W.J. (1997). Fertilization-independent seed development in *Arabidopsis thali-*
822   *ana*.   Proc.   Natl.   Acad.   Sci.   U.   S.   A.   94:   4223–4228.
823

824   9. Chen, D., Molitor, A., Liu, C., and Shen, W.H. (2010). The *Arabidopsis* PRC1-
825   like ring-finger proteins are necessary for repression of embryonic traits during
826   vegetative   growth.   Cell   Res.   20:   1332–1344.
827

828   10. Choi, Y., Gehring, M., Johnson, L., Hannon, M., Harada, J.J., Goldberg, R.B.,
829   Jacobsen, S.E., and Fischer, R.L. (2002). DEMETER, a DNA glycosylase
830   domain protein, is required for endosperm gene imprinting and seed viability
831   in   *Arabidopsis*.   Cell   110:   33–42.
832

833   11. Cui, X. *et al.* (2016). REF6 recognizes a specific DNA sequence to demethyl-
834   ate H3K27me3 and regulate organ boundary formation in *Arabidopsis*. Nat.
835   Genet. 48: 694–699.
836

837   12. Daimon, Y., Takabe, K., and Tasaka, M. (2003). *The CUP-SHAPED COTY-*
838   *LEDON* Genes Promote Adventitious Shoot Formation on Calli. Plant Cell
839   Physiol.   44:   113–121.
840

841   13. Dasgupta, P. and Chaudhuri, S. (2019). Analysis of DNA Methylation Profile
842   in Plants by Chop-PCR. In Plant Innate Immunity, W. Gassmann, ed, Methods
843   in Molecular Biology. (Springer New York: New York, NY), pp. 79–90.
844

845   14. De Lucia, F., Crevillen, P., Jones, A.M.E., Greb, T., and Dean, C. (2008). A
846   PHD-polycomb repressive complex 2 triggers the epigenetic silencing of FLC
847   during vernalization. Proc. Natl. Acad. Sci. U. S. A. 105: 16831–16836.
848

849 15. Du, J., Johnson, L.M., Jacobsen, S.E., and Patel, D.J. (2015). DNA methyla-
850 tion pathways and their crosstalk with histone methylation. Nat. Rev. Mol. Cell
851 Biol. 16: 519–532.
852

853 16. Feng, J., Liu, T., Qin, B., Zhang, Y., and Liu, X.S. (2012). Identifying ChIP-seq
854 enrichment using MACS. Nat. Protoc. 7: 1728–1740.
855

856 17. Figueiredo, D.D., Batista, R.A., Roszak, P.J., and Köhler, C. (2015). Auxin
857 production couples endosperm development to fertilization. Nat. Plants 1.
858

859 18. Gehring, M., Bubb, K.L., and Henikoff, S. (2009). Extensive Demethylation of
860 Repetitive Elements During Seed Development Underlies Gene Imprinting.
861 Science 324: 1447–1451.
862

863 19. Gehring, M., Huh, J.H., Hsieh, T.F., Penterman, J., Choi, Y., Harada, J.J.,
864 Goldberg, R.B., and Fischer, R.L. (2006). DEMETER DNA glycosylase estab-
865 lishes MEDEA polycomb gene self-imprinting by allele-specific demethylation.
866 Cell 124: 495–506.
867

868 20. Gong, Z., Morales-Ruiz, T., Ariza, R.R., Roldán-Arjona, T., David, L., and Zhu,
869 J.K. (2002). ROS1, a repressor of transcriptional gene silencing in *Arabidop-
870 sis*, encodes a DNA glycosylase/lyase. Cell 111: 803–814.
871

872 21. Grossniklaus, U., Vielle-Calzada, J.P., Hoeppner, M.A., and Gagliano, W.B.
873 (1998). Maternal control of embryogenesis by *MEDEA*, a Polycomb group
874 gene in *Arabidopsis*. Science 280: 446–450.
875

876 22. Guitton, A.-E., Page, D.R., Chambrier, P., Lionnet, C., Faure, J.-E.,
877 Grossniklaus, U., and Berger, F. (2004). Identification of new members of Fer-
878 tilization Independent Seed Polycomb Group pathway involved in the control
879 of seed development in *Arabidopsis thaliana*. Development 131: 2971–81.
880

881 23. Gutzat, R. *et al.* (2020). *Arabidopsis* shoot stem cells display dynamic tran-
882 scription and DNA methylation patterns. EMBO J. 39.
883

884 24. Heisler, M.G., Ohno, C., Das, P., Sieber, P., Reddy, G.V., Long, J.A., and
885 Meyerowitz, E.M. (2005). Patterns of auxin transport and gene expression
886 during primordium development revealed by live imaging of the *Arabidopsis*
887 inflorescence meristem. Curr. Biol. 15: 1899–1911.
888

889 25. Higo, A. *et al.* (2020). DNA methylation is reconfigured at the onset of repro-
890 duction in rice shoot apical meristem. Nat. Commun. 11: 1–12.
891

892 26. Hofhuis, H., Laskowski, M., Du, Y., Prasad, K., Grigg, S., Pinon, V., and
893   Scheres, B. (2013). Phyllotaxis and rhizotaxis in *Arabidopsis* are modified by
894   three plethora transcription factors. Curr. Biol. 23: 956–962.
895

896 27. Ibarra, C.A. *et al.* (2012). Active DNA Demethylation in Plant Companion Cells
897   Reinforces Transposon Methylation in Gametes. Science 337: 1360–1364.

898 28. Iwasaki M and Paszkowski J (2014) Epigenetic memory in plants. EMBO J.
899   33: 1987-1998.
900   a.
901

902 29. Kareem, A., Durgaprasad, K., Sugimoto, K., Du, Y., Pulianmackal, A.J.,
903   Trivedi, Z.B., Abhayadev, P.V., Pinon, V., Meyerowitz, E.M., Scheres, B., and
904   Prasad, K. (2015). *PLETHORA* genes control regeneration by a two-step
905   mechanism. Curr. Biol. 25: 1017–1030.
906

907 30. Khouider, S., Borges, F., LeBlanc, C., Ungru, A., Schnittger, A., Martienssen,
908   R., Colot, V., and Bouyer, D. (2021). Male fertility in *Arabidopsis* requires ac-
909   tive DNA demethylation of genes that control pollen tube function. Nat. Com-
910   mun. 12: 1–10.
911

912 31. Kim, M., Ohr, H., Lee, J.W., Hyun, Y., Fischer, R.L., and Choi, Y. (2008).
913   Temporal and spatial downregulation of *Arabidopsis* MET1 activity results in
914   global DNA hypomethylation and developmental defects. Mol. Cells 26: 611–
915   615.
916

917 32. Kim, S., Park, J.S., Lee, J., Lee, K.K., Park, O.S., Choi, H.S., Seo, P.J., Cho,
918   H.T., Frost, J.M., Fischer, R.L., and Choi, Y. (2021). The DME demethylase
919   regulates sporophyte gene expression, cell proliferation, differentiation, and
920   meristem resurrection. Proc. Natl. Acad. Sci. U. S. A. 118.
921

922 33. Köhler, C., Hennig, L., Spillane, C., Pien, S., and Gruissem, W. (2003). The
923   Polycomb -group protein MEDEA regulates seed development by controlling
924   expression of the MADS-box gene *PHERES1*. Genes Dev. 17: 1540–1553.
925

926 34. Kuwabara, A. and Gruissem, W. (2014). *Arabidopsis* Retinoblastoma-related
927   and Polycomb group proteins: Cooperation during plant cell differentiation and
928   development. J. Exp. Bot. 65: 2667–2676.
929

930 35. Kwon, C.S., Hibara, K.I., Pfluger, J., Bezhani, S., Metha, H., Aida, M., Tasaka,
931   M., and Wagner, D. (2006). A role for chromatin remodeling in regulation of
932   CUC gene expression in the *Arabidopsis* cotyledon boundary. Development
933   133: 3223–3230.
934

36. Leroy, O., Hennig, L., Breuninger, H., Laux, T., and Köhler, C. (2007). Polycomb group proteins function in the female gametophyte to determine seed development in plants. Development 134: 3639–3648.

37. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

38. Li, W., Liu, H., Cheng, Z.J., Su, Y.H., Han, H.N., Zhang, Y., and Zhang, X.S. (2011). DNA methylation and histone modifications regulate de novo shoot regeneration in *Arabidopsis* by modulating WUSCHEL expression and auxin signaling. PLoS Genet. 7.

39. Lindsey, B.E., Rivero, L., Calhoun, C.S., Grotewold, E., and Brkljacic, J. (2017). Standardized method for high-throughput sterilization of *Arabidopsis* seeds. J. Vis. Exp. 2017: 1–7.

40. Livak, K.J. and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. Methods 25: 402–408.

41. Lodha, M., Marco, C.F., and Timmermans, M.C.P. (2013). The ASYMMETRIC LEAVES complex maintains repression of *KNOX* homeobox genes via direct recruitment of Polycomb-repressive complex2. Genes Dev. 27: 596–601.

42. Luo, M., Bilodeau, P., Dennis, E.S., Peacock, W.J., and Chaudhury, a (2000). Expression and parent-of-origin effects for FIS2, MEA, and FIE in the endosperm and embryo of developing *Arabidopsis* seeds. Proc. Natl. Acad. Sci. U. S. A. 97: 10637–10642.

43. Makarevich, G., Villar, C.B.R., Erilova, A., and Köhler, C. (2008). Mechanism of *PHERES1* imprinting in *Arabidopsis*. J. Cell Sci. 121: 906–912.

44. Mayer, K.F.X., Schoof, H., Haecker, A., Lenhard, M., Jürgens, G., and Laux, T. (1998). Role of WUSCHEL in Regulating Stem Cell Fate in the *Arabidopsis* Shoot Meristem. Cell 95: 805–815.

45. Okonechnikov, K., Conesa, A., and García-Alcalde, F. (2016). Qualimap 2: Advanced multi-sample quality control for high-throughput sequencing data. Bioinformatics 32: 292–294.

46. Park, J., Frost, J.M., Park, K., Ohr, H., Tae, G., Kim, S., and Eom, H. (2017). Control of DEMETER DNA demethylase gene transcription in male and fe-

979   male gamete companion cells in *Arabidopsis thaliana*. Proc. Natl. Acad. Sci.
980   U.   S.   A.   114:   2078–2083.

47. Pikaard, C.S. and Scheid, O.M. (2014). Epigenetic regulation in plants. Cold Spring Harb. Perspect. Biol. 6: 1–31.

48. Prasad, K. *et al.* (2011). *Arabidopsis* PLETHORA transcription factors control phyllotaxis. Curr. Biol. 21: 1123–1128.

49. Qiu, Q., Mei, H., Deng, X., He, K., Wu, B., Yao, Q., Zhang, J., Lu, F., Ma, J., and Cao, X. (2019). DNA methylation repels targeting of *Arabidopsis* REF6. Nat. Commun. 10: 1–9.

50. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Res. 44: W160–W165.

51. Ravi, M. *et al.* (2014). A haploid genetics toolbox for *Arabidopsis thaliana*. Nat. Commun. 5: 5334.

52. Ravi, M. and Bondada, R. (2016). Genome Elimination by Tailswap CenH3: In Vivo Haploid Production in *Arabidopsis thaliana*. Methods Mol Biol. 1469:77-99.

53. Ravi, M. and Chan, S.W.L. (2010). Haploid plants produced by centromere-mediated genome elimination. Nature 464: 615–618.

54. Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Bio-technol. 29: 24–26.

55. Roszak, P.J. and Köhler, C. (2011). Polycomb group proteins are required to couple seed coat initiation to fertilization. Proc. Natl. Acad. Sci. U. S. A. 108: 20826–20831.

56. Roy, S., Gupta, P., Rajabhoj, M.P., Maruthachalam, R., and Nandi, A.K. (2018). The polycomb-group repressor MEDEA attenuates pathogen defense. Plant Physiol.

57. Saleh, A., Al-Abdallat, A., Ndamukong, I., Alvarez-Venegas, R., and Avramova, Z. (2007). The *Arabidopsis* homologs of trithorax (ATX1) and en-hancer of zeste (CLF) establish "bivalent chromatin marks" at the silent *AGA-*

MOUS   locus.   Nucleic   Acids   Res.   35:   6290–6296.

58. Salvi, E., Rutten, J.P., Di Mambro, R., Polverari, L., Licursi, V., Negri, R., Dello Ioio, R., Sabatini, S., and Ten Tusscher, K. (2020). A Self-Organized PLT/Auxin/ARR-B Network Controls the Dynamics of Root Zonation Development in *Arabidopsis thaliana*. Dev. Cell 53: 431-443.e23.

59. Schmid, M., Davison, T.S., Henz, S.R., Pape, U.J., Demar, M., Vingron, M., Schölkopf, B., Weigel, D., and Lohmann, J.U. (2005). A gene expression map of *Arabidopsis thaliana* development. Nat. Genet. 37: 501–506.

60. Schoft, V.K., Chumak, N., Choi, Y., Hannon, M., Garcia-Aguilar, M., Machlicova, A., Slusarz, L., Mosiolek, M., Park, J.-S., Park, G.T., Fischer, R.L., and Tamaru, H. (2011). Function of the DEMETER DNA glycosylase in the *Arabidopsis thaliana* male gametophyte. Proc Natl Acad Sci USA 108: 8042–7.

61. Schoof, H., Lenhard, M., Haecker, A., Mayer, K.F.X., Jürgens, G., and Laux, T. (2000). The stem cell population of *Arabidopsis* shoot meristems is maintained by a regulatory loop between the *CLAVATA* and *WUSCHEL* genes. Cell   100:   635–644.

62. Schumann, U., Lee, J.M., Smith, N.A., Zhong, C., Zhu, J.K., Dennis, E.S., Millar, A.A., and Wang, M.B. (2019). DEMETER plays a role in DNA demethylation and disease response in somatic tissues of *Arabidopsis*. Epigenetics 14: 1074–1087.

63. Scofield, S., Dewitte, W., and Murray, J.A. (2008). A model for *Arabidopsis* class-1 KNOX gene function. Plant Signal. Behav. 3: 257–259.

64. Shen, W.-H. and Xu, L. (2009). Chromatin Remodeling in Stem Cell Maintenance in *Arabidopsis thaliana*. Mol. Plant 2: 600–609.

65. Simonini, S., Bemer, M., Bencivenga, S., Gagliardini, V., Pires, N.D., Desvoyes, B., van der Graaff, E., Gutierrez, C., and Grossniklaus, U. (2021). The Polycomb group protein MEDEA controls cell proliferation and embryonic patterning in *Arabidopsis*. Dev. Cell 56: 1945-1960.e7.

66. Spillane, C., Schmid, K.J., Laoueillé-Duprat, S., Pien, S., Escobar-Restrepo, J.-M., Baroux, C., Gagliardini, V., Page, D.R., Wolfe, K.H., and Grossniklaus, U. (2007). Positive darwinian selection at the imprinted *MEDEA* locus in plants.   Nature   448:   349–352.

1065   67. Spinelli, S.V., Martin, A.P., Viola, I.L., Gonzalez, D.H., and Palatnik, J.F.
1066   (2011). A Mechanistic Link between *STM* and *CUC1* during *Arabidopsis* De-
1067   velopment.   Plant   Physiol.   156:   1894–1904.
1068

1069   68. Stahl, Y. and Simon, R. (2010). Plant primary meristems: shared functions
1070   and   regulatory   mechanisms.   Curr.   Opin.   Plant   Biol.   13:   53–58.
1071

1072   69. Su, Y.H., Zhou, C., Li, Y.J., Yu, Y., Tang, L.P., Zhang, W.J., Yao, W.J.,
1073   Huang, R., Laux, T., and Zhang, X.S. (2020). Integration of pluripotency
1074   pathways regulates stem cell maintenance in the *Arabidopsis* shoot meristem.
1075   Proc.   Natl.   Acad.   Sci.   U.   S.   A.   117:   22561–22571.
1076

1077   70. Sun, B., Looi, L.S., Guo, S., He, Z., Gan, E.S., Huang, J., Xu, Y., Wee, W.Y.,
1078   and Ito, T. (2014). Timing mechanism dependent on cell division is invoked by
1079   Polycomb   eviction   in   plant   stem   cells.   Science   343.
1080

1081   71. Sutherland, E., Coe, L., and Raleigh, E.A. (1992). McrBC: a multisubunit
1082   GTP-dependent   restriction   endonuclease.   J.   Mol.   Biol.   225:   327–348.
1083

1084   72. Ten Hove, C.A., Lu, K.J., and Weijers, D. (2015). Building a plant: Cell fate
1085   specification in the early *Arabidopsis* embryo. Development 142: 420–430.
1086

1087   73. Thomas, J., Spillane, C., Coluccio, A., Hoeppner, M.A., and Grossniklaus, U.
1088   (1999). Maintenance of genomic imprinting at the *Arabidopsis MEDEA* locus
1089   requires   zygotic   DDM1   activity.:   2971–2982.
1090

1091   74. Varapparambath, V. *et al.* (2022). Mechanical conflict caused by a cell-wall-
1092   loosening enzyme activates de novo shoot regeneration. Dev. Cell 57: 2063–
1093   2080.
1094

1095   75. Williams, B.P., Bechen, L.L., Pohlmann, D.A., and Gehring, M. (2022). Somat-
1096   ic DNA demethylation generates tissue-specific methylation states and im-
1097   pacts   flowering   time.   Plant   Cell   34:   1189–1206.
1098

1099   76. Williams, B.P. and Gehring, M. (2020). Principles of Epigenetic Homeostasis
1100   Shared Between Flowering Plants and Mammals. Trends Genet. 36: 751–
1101   763.
1102

1103   77. Xiao, J. *et al.* (2017). Cis and trans determinants of epigenetic silencing by
1104   Polycomb repressive complex 2 in *Arabidopsis*. Nat. Genet. 49: 1546–1552.
1105

1106   78. Xiao, J., Lee, U.-S., and Wagner, D. (2016). Tug of war: adding and removing
1107   histone lysine methylation in *Arabidopsis*. Curr. Opin. Plant Biol. 34: 41–53.
1108

1109 79. Xiao, J. and Wagner, D. (2015). Polycomb repression in the regulation of
1110 growth and development in *Arabidopsis*. Curr. Opin. Plant Biol. 23: 15–24.
1111

1112 80. Xiao, W., Gehring, M., Choi, Y., Margossian, L., Pu, H., Harada, J.J.,
1113 Goldberg, R.B., Pennell, R.I., and Fischer, R.L. (2003). Imprinting of the *MEA*
1114 polycomb gene is controlled by antagonism between MET1 methyltransferase
1115 and DME glycosylase. Dev. Cell 5: 891–901.
1116

1117 81. Yadav, R.K., Perales, M., Gruel, J., Ohno, C., Heisler, M., Girke, T., Jönsson,
1118 H., and Reddy, G.V. (2013). Plant stem cell maintenance involves direct tran-
1119 scriptional repression of differentiation program. Mol. Syst. Biol. 9: 654.
1120

1121 82. Yamaguchi, N., Winter, C.M., Wu, M.-F., Kwon, C.S., William, D.A., and Wag-
1122 ner, D. (2014). PROTOCOL: Chromatin Immunoprecipitation from *Arabidopsis*
1123 Tissues. Arab. Book 12: e0170.
1124

1125 83. Yoshida, N., Yanai, Y., Chen, L., Kato, Y., Hiratsuka, J., Miwa, T., Sung, Z.R.,
1126 and Takahashi, S. (2001). EMBRYONIC FLOWER2, a novel Polycomb group
1127 protein homolog, mediates shoot development and flowering in *Arabidopsis*.
1128 Plant Cell 13: 2471–2481.
1129

1130 84. Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). ClusterProfiler: An R pack-
1131 age for comparing biological themes among gene clusters. OMICS J. Integr.
1132 Biol. 16: 284–287.
1133

1134 85. Yu, G., Wang, L.G., and He, Q.Y. (2015). ChIP seeker: An R/Bioconductor
1135 package for ChIP peak annotation, comparison and visualization. Bioinformat-
1136 ics 31: 2382–2383.
1137

1138 86. Zhang, H., Tang, K., Wang, B., Duan, C.G., Lang, Z., and Zhu, J.K. (2014).
1139 Protocol: A beginner's guide to the analysis of RNA-directed DNA methylation
1140 in plants. Plant Methods 10: 1–9.
1141

1142 87. Zhang, S., Wang, D., Zhang, H., Skaggs, M.I., Lloyd, A., Ran, D., An, L.,
1143 Schumaker, K.S., Drews, G.N., and Yadegari, R. (2018). FERTILIZATION-
1144 INDEPENDENT SEED-Polycomb Repressive Complex 2 Plays a Dual Role in
1145 Regulating Type I MADS-Box Genes in Early Endosperm Development. Plant
1146 Physiol. 177: 285 LP – 299.
1147

1148 88. Zhang, X., Clarenz, O., Cokus, S., Bernatavichute, Y.V., Pellegrini, M.,
1149 Goodrich, J., and Jacobsen, S.E. (2007). Whole-Genome Analysis of Histone
1150 H3 Lysine 27 Trimethylation in *Arabidopsis*. PLOS Biol. 5: e129.
1151

1152   89. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
1153   Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-
1154   based Analysis of ChIP-Seq (MACS). Genome Biol. 9: R137.
1155

1156   90. Zuo, J., Niu, Q.W., and Chua, N.H. (2000). An estrogen receptor-based trans-
1157   activator XVE mediates highly inducible gene expression in transgenic plants.
1158   Plant J. 24: 265–273.

Parsed Citations

Aichinger, E., Villar, C.B.R., Di Mambro, R., Sabatini, S., and Köhler, C. (2011). The CHD3 Chromatin Remodeler PICKLE and Polycomb Group Proteins Antagonistically Regulate Meristem Activity in the Arabidopsis Root. Plant Cell 23: 1047–1060.
  Google Scholar: Author Only Title Only Author and Title

Aida, M., Ishida, T., and Tasaka, M. (1999). Shoot apical meristem and cotyledon formation during Arabidopsis embryogenesis: Interaction among the CUP-SHAPED COTYLEDON and SHOOT MERISTEMLESS genes. Development 126: 1563–1570.
  Google Scholar: Author Only Title Only Author and Title

Aida, M., Vernoux, T., Furutani, M., Traas, J., and Tasaka, M. (2002). Roles of PIN-FORMED1 and MONOPTEROS in pattern formation of the apical region of the Arabidopsis embryo. Development 129: 3965–3974.
  Google Scholar: Author Only Title Only Author and Title

Borg, M., Jacob, Y., Susaki, D. et al. (2020).Targeted reprogramming of H3K27me3 resets epigenetic memory in plant paternal chromatin. Nat Cell Biol 22: 621–629.
  Google Scholar: Author Only Title Only Author and Title

Bouyer, D., Kramdi, A, Kassam, M., Heese, M., Schnittger, A, Roudier, F., and Colot, V. (2017). DNA methylation dynamics during early plant life. Genome Biol. 18: 1–12.
  Google Scholar: Author Only Title Only Author and Title

Bouyer, D., Roudier, F., Heese, M., Andersen, E.D., Gey, D., Nowack, M.K., Goodrich, J., Renou, J.P., Grini, P.E., Colot, V., and Schnittger, A. (2011). Polycomb repressive complex 2 controls the embryo-to-seedling phase transition. PLoS Genet. 7.
  Google Scholar: Author Only Title Only Author and Title

Burian, A, Raczyńska-Szajgin, M., Borowska-Wykręt, D., Piatek, A, Aida, M., and Kwiatkowska, D. (2015). The CUP-SHAPED COTYLEDON2 and 3 genes have a post-meristematic effect on Arabidopsis thaliana phyllotaxis. Ann. Bot. 115: 807–820.
  Google Scholar: Author Only Title Only Author and Title

Chaudhury, A.M., Ming, L., Miller, C., Craig, S., Dennis, E.S., and Peacock, W.J. (1997). Fertilization-independent seed development in Arabidopsis thaliana. Proc. Natl. Acad. Sci. U. S. A. 94: 4223–4228.
  Google Scholar: Author Only Title Only Author and Title

Chen, D., Molitor, A, Liu, C., and Shen, W.H. (2010). The Arabidopsis PRC1-like ring-finger proteins are necessary for repression of embryonic traits during vegetative growth. Cell Res. 20: 1332–1344.
  Google Scholar: Author Only Title Only Author and Title

Choi, Y., Gehring, M., Johnson, L., Hannon, M., Harada, J.J., Goldberg, R.B., Jacobsen, S.E., and Fischer, R.L. (2002). DEMETER, a DNA glycosylase domain protein, is required for endosperm gene imprinting and seed viability in Arabidopsis. Cell 110: 33–42.
  Google Scholar: Author Only Title Only Author and Title

Cui, X. et al. (2016). REF6 recognizes a specific DNA sequence to demethylate H3K27me3 and regulate organ boundary formation in Arabidopsis. Nat. Genet. 48: 694–699.
  Google Scholar: Author Only Title Only Author and Title

Daimon, Y., Takabe, K., and Tasaka, M. (2003). The CUP-SHAPED COTYLEDON Genes Promote Adventitious Shoot Formation on Calli. Plant Cell Physiol. 44: 113–121.
  Google Scholar: Author Only Title Only Author and Title

Dasgupta, P. and Chaudhuri, S. (2019). Analysis of DNA Methylation Profile in Plants by Chop-PCR. In Plant Innate Immunity, W. Gassmann, ed, Methods in Molecular Biology. (Springer New York: New York, NY), pp. 79–90.
  Google Scholar: Author Only Title Only Author and Title

De Lucia, F., Crevillen, P., Jones, A.M.E., Greb, T., and Dean, C. (2008). A PHD-polycomb repressive complex 2 triggers the epigenetic silencing of FLC during vernalization. Proc. Natl. Acad. Sci. U. S. A. 105: 16831–16836.
  Google Scholar: Author Only Title Only Author and Title

Du, J., Johnson, L.M., Jacobsen, S.E., and Patel, D.J. (2015). DNA methylation pathways and their crosstalk with histone methylation. Nat. Rev. Mol. Cell Biol. 16: 519–532.
  Google Scholar: Author Only Title Only Author and Title

Feng, J., Liu, T., Qin, B., Zhang, Y., and Liu, X.S. (2012). Identifying ChIP-seq enrichment using MACS. Nat. Protoc. 7: 1728–1740.
  Google Scholar: Author Only Title Only Author and Title

Figueiredo, D.D., Batista, R.A, Roszak, P.J., and Köhler, C. (2015). Auxin production couples endosperm development to fertilization. Nat. Plants 1.
  Google Scholar: Author Only Title Only Author and Title

Gehring, M., Bubb, K.L., and Henikoff, S. (2009). Extensive Demethylation of Repetitive Elements During Seed Development

Underlies Gene Imprinting. Science 324: 1447–1451.
Google Scholar: Author Only Title Only Author and Title

Gehring, M., Huh, J.H., Hsieh, T.F., Penterman, J., Choi, Y., Harada, J.J., Goldberg, R.B., and Fischer, R.L. (2006). DEMETER DNA glycosylase establishes MEDEA polycomb gene self-imprinting by allele-specific demethylation. Cell 124: 495–506.
Google Scholar: Author Only Title Only Author and Title

Gong, Z, Morales-Ruiz, T., Ariza, R.R., Roldán-Arjona, T., David, L., and Zhu, J.K. (2002). ROS1, a repressor of transcriptional gene silencing in Arabidopsis, encodes a DNA glycosylase/lyase. Cell 111: 803–814.
Google Scholar: Author Only Title Only Author and Title

Grossniklaus, U., Vielle-Calzada, J.P., Hoeppner, M.A., and Gagliano, W.B. (1998). Maternal control of embryogenesis by MEDEA, a Polycomb group gene in Arabidopsis. Science 280: 446–450.
Google Scholar: Author Only Title Only Author and Title

Guitton, A.-E., Page, D.R., Chambrier, P., Lionnet, C., Faure, J.-E., Grossniklaus, U., and Berger, F. (2004). Identification of new members of Fertilization Independent Seed Polycomb Group pathway involved in the control of seed development in Arabidopsis thaliana. Development 131: 2971–81.
Google Scholar: Author Only Title Only Author and Title

Gutzat, R. et al. (2020). Arabidopsis shoot stem cells display dynamic transcription and DNA methylation patterns. EMBO J. 39.
Google Scholar: Author Only Title Only Author and Title

Heisler, M.G., Ohno, C., Das, P., Sieber, P., Reddy, G.V., Long, J.A., and Meyerowitz, E.M. (2005). Patterns of auxin transport and gene expression during primordium development revealed by live imaging of the Arabidopsis inflorescence meristem. Curr. Biol. 15: 1899–1911.
Google Scholar: Author Only Title Only Author and Title

Higo, A. et al. (2020). DNA methylation is reconfigured at the onset of reproduction in rice shoot apical meristem. Nat. Commun. 11: 1–12.
Google Scholar: Author Only Title Only Author and Title

Hofhuis, H., Laskowski, M., Du, Y., Prasad, K., Grigg, S., Pinon, V., and Scheres, B. (2013). Phyllotaxis and rhizotaxis in Arabidopsis are modified by three plethora transcription factors. Curr. Biol. 23: 956–962.
Google Scholar: Author Only Title Only Author and Title

Ibarra, C.A. et al. (2012). Active DNA Demethylation in Plant Companion Cells Reinforces Transposon Methylation in Gametes. Science 337: 1360–1364.
Google Scholar: Author Only Title Only Author and Title

Iwasaki M and Paszkowski J (2014) Epigenetic memory in plants. EMBO J. 33: 1987-1998.
Google Scholar: Author Only Title Only Author and Title

Kareem, A, Durgaprasad, K., Sugimoto, K., Du, Y., Pulianmackal, A.J., Trivedi, Z.B., Abhayadev, P.V., Pinon, V., Meyerowitz, E.M., Scheres, B., and Prasad, K. (2015). PLETHORA genes control regeneration by a two-step mechanism. Curr. Biol. 25: 1017–1030.
Google Scholar: Author Only Title Only Author and Title

Khouider, S., Borges, F., LeBlanc, C., Ungru, A, Schnittger, A, Martienssen, R., Colot, V., and Bouyer, D. (2021). Male fertility in Arabidopsis requires active DNA demethylation of genes that control pollen tube function. Nat. Commun. 12: 1–10.
Google Scholar: Author Only Title Only Author and Title

Kim, M., Ohr, H., Lee, J.W., Hyun, Y., Fischer, R.L., and Choi, Y. (2008). Temporal and spatial downregulation of Arabidopsis MET1 activity results in global DNA hypomethylation and developmental defects. Mol. Cells 26: 611–615.
Google Scholar: Author Only Title Only Author and Title

Kim, S., Park, J.S., Lee, J., Lee, K.K., Park, O.S., Choi, H.S., Seo, P.J., Cho, H.T., Frost, J.M., Fischer, R.L., and Choi, Y. (2021). The DME demethylase regulates sporophyte gene expression, cell proliferation, differentiation, and meristem resurrection. Proc. Natl. Acad. Sci. U. S. A. 118.
Google Scholar: Author Only Title Only Author and Title

Köhler, C., Hennig, L., Spillane, C., Pien, S., and Gruissem, W. (2003). The Polycomb -group protein MEDEA regulates seed development by controlling expression of the MADS-box gene PHERES1. Genes Dev. 17: 1540–1553.
Google Scholar: Author Only Title Only Author and Title

Kuwabara, A. and Gruissem, W. (2014). Arabidopsis Retinoblastoma-related and Polycomb group proteins: Cooperation during plant cell differentiation and development. J. Exp. Bot. 65: 2667–2676.
Google Scholar: Author Only Title Only Author and Title

Kwon, C.S., Hibara, K.I., Pfluger, J., Bezhani, S., Metha, H., Aida, M., Tasaka, M., and Wagner, D. (2006). A role for chromatin remodeling in regulation of CUC gene expression in the Arabidopsis cotyledon boundary. Development 133: 3223–3230.

Google Scholar: Author Only Title Only Author and Title

Leroy, O., Hennig, L., Breuninger, H., Laux, T., and Köhler, C. (2007). Polycomb group proteins function in the female gametophyte to determine seed development in plants. Development 134: 3639–3648.
  Google Scholar: Author Only Title Only Author and Title

Li, H., Handsaker, B., Wysoker, A, Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.
  Google Scholar: Author Only Title Only Author and Title

Li, W., Liu, H., Cheng, Z.J., Su, Y.H., Han, H.N., Zhang, Y., and Zhang, X.S. (2011). DNA methylation and histone modifications regulate de novo shoot regeneration in Arabidopsis by modulating WUSCHEL expression and auxin signaling. PLoS Genet. 7.
  Google Scholar: Author Only Title Only Author and Title

Lindsey, B.E., Rivero, L., Calhoun, C.S., Grotewold, E., and Brkljacic, J. (2017). Standardized method for high-throughput sterilization of Arabidopsis seeds. J. Vis. Exp. 2017: 1–7.
  Google Scholar: Author Only Title Only Author and Title

Livak, K.J. and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. Methods 25: 402–408.
  Google Scholar: Author Only Title Only Author and Title

Lodha, M., Marco, C.F., and Timmermans, M.C.P. (2013). The ASYMMETRIC LEAVES complex maintains repression of KNOX homeobox genes via direct recruitment of Polycomb-repressive complex2. Genes Dev. 27: 596–601.
  Google Scholar: Author Only Title Only Author and Title

Luo, M., Bilodeau, P., Dennis, E.S., Peacock, W.J., and Chaudhury, a (2000). Expression and parent-of-origin effects for FIS2, MEA, and FIE in the endosperm and embryo of developing Arabidopsis seeds. Proc. Natl. Acad. Sci. U. S. A. 97: 10637–10642.
  Google Scholar: Author Only Title Only Author and Title

Makarevich, G., Villar, C.B.R., Erilova, A, and Köhler, C. (2008). Mechanism of PHERES1 imprinting in Arabidopsis. J. Cell Sci. 121: 906–912.
  Google Scholar: Author Only Title Only Author and Title

Mayer, K.F.X., Schoof, H., Haecker, A, Lenhard, M., Jürgens, G., and Laux, T. (1998). Role of WUSCHEL in Regulating Stem Cell Fate in the Arabidopsis Shoot Meristem. Cell 95: 805–815.
  Google Scholar: Author Only Title Only Author and Title

Okonechnikov, K., Conesa, A., and García-Alcalde, F. (2016). Qualimap 2: Advanced multi-sample quality control for high-throughput sequencing data. Bioinformatics 32: 292–294.
  Google Scholar: Author Only Title Only Author and Title

Park, J., Frost, J.M., Park, K., Ohr, H., Tae, G., Kim, S., and Eom, H. (2017). Control of DEMETER DNA demethylase gene transcription in male and female gamete companion cells in Arabidopsis thaliana. Proc. Natl. Acad. Sci. U. S. A. 114: 2078–2083.
  Google Scholar: Author Only Title Only Author and Title

Pikaard, C.S. and Scheid, O.M. (2014). Epigenetic regulation in plants. Cold Spring Harb. Perspect. Biol. 6: 1–31.
  Google Scholar: Author Only Title Only Author and Title

Prasad, K. et al. (2011). Arabidopsis PLETHORA transcription factors control phyllotaxis. Curr. Biol. 21: 1123–1128.
  Google Scholar: Author Only Title Only Author and Title

Qiu, Q., Mei, H., Deng, X., He, K., Wu, B., Yao, Q., Zhang, J., Lu, F., Ma, J., and Cao, X. (2019). DNA methylation repels targeting of Arabidopsis REF6. Nat. Commun. 10: 1–9.
  Google Scholar: Author Only Title Only Author and Title

Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Res. 44: W160–W165.
  Google Scholar: Author Only Title Only Author and Title

Ravi, M. et al. (2014). A haploid genetics toolbox for Arabidopsis thaliana. Nat. Commun. 5: 5334.
  Google Scholar: Author Only Title Only Author and Title

Ravi, M. and Bondada, R. (2016). Genome Elimination by Tailswap CenH3: In Vivo Haploid Production in Arabidopsis thaliana. Methods Mol Biol. 1469:77-99.
  Google Scholar: Author Only Title Only Author and Title

Ravi, M. and Chan, S.W.L. (2010). Haploid plants produced by centromere-mediated genome elimination. Nature 464: 615–618.
  Google Scholar: Author Only Title Only Author and Title

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative

genomics viewer. Nat. Biotechnol. 29: 24–26.
Google Scholar: Author Only Title Only Author and Title

Roszak, P.J. and Köhler, C. (2011). Polycomb group proteins are required to couple seed coat initiation to fertilization. Proc. Natl. Acad. Sci. U. S. A. 108: 20826–20831.
Google Scholar: Author Only Title Only Author and Title

Roy, S., Gupta, P., Rajabhoj, M.P., Maruthachalam, R., and Nandi, A.K. (2018). The polycomb-group repressor MEDEA attenuates pathogen defense. Plant Physiol.
Google Scholar: Author Only Title Only Author and Title

Saleh, A., Al-Abdallat, A., Ndamukong, I., Alvarez-Venegas, R., and Avramova, Z. (2007). The Arabidopsis homologs of trithorax (ATX1) and enhancer of zeste (CLF) establish "bivalent chromatin marks" at the silent AGAMOUS locus. Nucleic Acids Res. 35: 6290–6296.
Google Scholar: Author Only Title Only Author and Title

Salvi, E., Rutten, J.P., Di Mambro, R., Polverari, L., Licursi, V., Negri, R., Dello Ioio, R., Sabatini, S., and Ten Tusscher, K. (2020). A Self-Organized PLT/Auxin/ARR-B Network Controls the Dynamics of Root Zonation Development in Arabidopsis thaliana. Dev. Cell 53: 431-443.e23.
Google Scholar: Author Only Title Only Author and Title

Schmid, M., Davison, T.S., Henz, S.R., Pape, U.J., Demar, M., Vingron, M., Schölkopf, B., Weigel, D., and Lohmann, J.U. (2005). A gene expression map of Arabidopsis thaliana development. Nat. Genet. 37: 501–506.
Google Scholar: Author Only Title Only Author and Title

Schoft, V.K., Chumak, N., Choi, Y., Hannon, M., Garcia-Aguilar, M., Machlicova, A., Slusarz, L., Mosiolek, M., Park, J.-S., Park, G.T., Fischer, R.L., and Tamaru, H. (2011). Function of the DEMETER DNA glycosylase in the Arabidopsis thaliana male gametophyte. Proc Natl Acad Sci USA 108: 8042–7.
Google Scholar: Author Only Title Only Author and Title

Schoof, H., Lenhard, M., Haecker, A., Mayer, K.F.X., Jürgens, G., and Laux, T. (2000). The stem cell population of Arabidopsis shoot meristems is maintained by a regulatory loop between the CLAVATA and WUSCHEL genes. Cell 100: 635–644.
Google Scholar: Author Only Title Only Author and Title

Schumann, U., Lee, J.M., Smith, N.A., Zhong, C., Zhu, J.K., Dennis, E.S., Millar, A.A., and Wang, M.B. (2019). DEMETER plays a role in DNA demethylation and disease response in somatic tissues of Arabidopsis. Epigenetics 14: 1074–1087.
Google Scholar: Author Only Title Only Author and Title

Scofield, S., Dewitte, W., and Murray, J.A. (2008). A model for Arabidopsis class-1 KNOX gene function. Plant Signal. Behav. 3: 257–259.
Google Scholar: Author Only Title Only Author and Title

Shen, W.-H. and Xu, L. (2009). Chromatin Remodeling in Stem Cell Maintenance in Arabidopsis thaliana. Mol. Plant 2: 600–609.
Google Scholar: Author Only Title Only Author and Title

Simonini, S., Bemer, M., Bencivenga, S., Gagliardini, V., Pires, N.D., Desvoyes, B., van der Graaff, E., Gutierrez, C., and Grossniklaus, U. (2021). The Polycomb group protein MEDEA controls cell proliferation and embryonic patterning in Arabidopsis. Dev. Cell 56: 1945-1960.e7.
Google Scholar: Author Only Title Only Author and Title

Spillane, C., Schmid, K.J., Laoueillé-Duprat, S., Pien, S., Escobar-Restrepo, J.-M., Baroux, C., Gagliardini, V., Page, D.R., Wolfe, K.H., and Grossniklaus, U. (2007). Positive darwinian selection at the imprinted MEDEA locus in plants. Nature 448: 349–352.
Google Scholar: Author Only Title Only Author and Title

Spinelli, S.V., Martin, A.P., Viola, I.L., Gonzalez, D.H., and Palatnik, J.F. (2011). A Mechanistic Link between STM and CUC1 during Arabidopsis Development. Plant Physiol. 156: 1894–1904.
Google Scholar: Author Only Title Only Author and Title

Stahl, Y. and Simon, R. (2010). Plant primary meristems: shared functions and regulatory mechanisms. Curr. Opin. Plant Biol. 13: 53–58.
Google Scholar: Author Only Title Only Author and Title

Su, Y.H., Zhou, C., Li, Y.J., Yu, Y., Tang, L.P., Zhang, W.J., Yao, W.J., Huang, R., Laux, T., and Zhang, X.S. (2020). Integration of pluripotency pathways regulates stem cell maintenance in the Arabidopsis shoot meristem. Proc. Natl. Acad. Sci. U. S. A. 117: 22561–22571.
Google Scholar: Author Only Title Only Author and Title

Sun, B., Looi, L.S., Guo, S., He, Z., Gan, E.S., Huang, J., Xu, Y., Wee, W.Y., and Ito, T. (2014). Timing mechanism dependent on cell division is invoked by Polycomb eviction in plant stem cells. Science 343.
Google Scholar: Author Only Title Only Author and Title

Sutherland, E., Coe, L., and Raleigh, E.A. (1992). McrBC: a multisubunit GTP-dependent restriction endonuclease. J. Mol. Biol. 225: 327–348.
  Google Scholar: Author Only Title Only Author and Title

Ten Hove, C.A, Lu, K.J., and Weijers, D. (2015). Building a plant: Cell fate specification in the early Arabidopsis embryo. Development 142: 420–430.
  Google Scholar: Author Only Title Only Author and Title

Thomas, J., Spillane, C., Coluccio, A, Hoeppner, M.A, and Grossniklaus, U. (1999). Maintenance of genomic imprinting at the Arabidopsis MEDEA locus requires zygotic DDM1 activity.: 2971–2982.
  Google Scholar: Author Only Title Only Author and Title

Varapparambath, V. et al. (2022). Mechanical conflict caused by a cell-wall-loosening enzyme activates de novo shoot regeneration. Dev. Cell 57: 2063–2080.
  Google Scholar: Author Only Title Only Author and Title

Williams, B.P., Bechen, L.L., Pohlmann, D.A, and Gehring, M. (2022). Somatic DNA demethylation generates tissue-specific methylation states and impacts flowering time. Plant Cell 34: 1189–1206.
  Google Scholar: Author Only Title Only Author and Title

Williams, B.P. and Gehring, M. (2020). Principles of Epigenetic Homeostasis Shared Between Flowering Plants and Mammals. Trends Genet. 36: 751–763.
  Google Scholar: Author Only Title Only Author and Title

Xiao, J. et al. (2017). Cis and trans determinants of epigenetic silencing by Polycomb repressive complex 2 in Arabidopsis. Nat. Genet. 49: 1546–1552.
  Google Scholar: Author Only Title Only Author and Title

Xiao, J., Lee, U.-S., and Wagner, D. (2016). Tug of war: adding and removing histone lysine methylation in Arabidopsis. Curr. Opin. Plant Biol. 34: 41–53.
  Google Scholar: Author Only Title Only Author and Title

Xiao, J. and Wagner, D. (2015). Polycomb repression in the regulation of growth and development in Arabidopsis. Curr. Opin. Plant Biol. 23: 15–24.
  Google Scholar: Author Only Title Only Author and Title

Xiao, W., Gehring, M., Choi, Y., Margossian, L., Pu, H., Harada, J.J., Goldberg, R.B., Pennell, R.I., and Fischer, R.L. (2003). Imprinting of the MEA polycomb gene is controlled by antagonism between MET1 methyltransferase and DME glycosylase. Dev. Cell 5: 891–901.
  Google Scholar: Author Only Title Only Author and Title

Yadav, R.K., Perales, M., Gruel, J., Ohno, C., Heisler, M., Girke, T., Jönsson, H., and Reddy, G.V. (2013). Plant stem cell maintenance involves direct transcriptional repression of differentiation program. Mol. Syst. Biol. 9: 654.
  Google Scholar: Author Only Title Only Author and Title

Yamaguchi, N., Winter, C.M., Wu, M.-F., Kwon, C.S., William, D.A., and Wagner, D. (2014). PROTOCOL: Chromatin Immunoprecipitation from Arabidopsis Tissues. Arab. Book 12: e0170.
  Google Scholar: Author Only Title Only Author and Title

Yoshida, N., Yanai, Y., Chen, L., Kato, Y., Hiratsuka, J., Miwa, T., Sung, Z.R., and Takahashi, S. (2001). EMBRYONIC FLOWER2, a novel Polycomb group protein homolog, mediates shoot development and flowering in Arabidopsis. Plant Cell 13: 2471–2481.
  Google Scholar: Author Only Title Only Author and Title

Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). ClusterProfiler: An R package for comparing biological themes among gene clusters. OMICS J. Integr. Biol. 16: 284–287.
  Google Scholar: Author Only Title Only Author and Title

Yu, G., Wang, L.G., and He, Q.Y. (2015). ChIP seeker: An R/Bioconductor package for ChIP peak annotation, comparison and visualization. Bioinformatics 31: 2382–2383.
  Google Scholar: Author Only Title Only Author and Title

Zhang, H., Tang, K., Wang, B., Duan, C.G., Lang, Z, and Zhu, J.K. (2014). Protocol: A beginner's guide to the analysis of RNA-directed DNA methylation in plants. Plant Methods 10: 1–9.
  Google Scholar: Author Only Title Only Author and Title

Zhang, S., Wang, D., Zhang, H., Skaggs, M.I., Lloyd, A, Ran, D., An, L., Schumaker, K.S., Drews, G.N., and Yadegari, R. (2018). FERTILIZATION-INDEPENDENT SEED-Polycomb Repressive Complex 2 Plays a Dual Role in Regulating Type I MADS-Box Genes in Early Endosperm Development. Plant Physiol. 177: 285 LP – 299.
  Google Scholar: Author Only Title Only Author and Title

Zhang, X., Clarenz, O., Cokus, S., Bernatavichute, Y.V., Pellegrini, M., Goodrich, J., and Jacobsen, S.E. (2007). Whole-Genome Analysis of Histone H3 Lysine 27 Trimethylation in Arabidopsis. PLOS Biol. 5: e129.
Google Scholar: Author Only Title Only Author and Title

Zhang, Y., Liu, T., Meyer, C.A, Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based Analysis of ChIP-Seq (MACS). Genome Biol. 9: R137.
Google Scholar: Author Only Title Only Author and Title

Zuo, J., Niu, Q.W., and Chua, N.H. (2000). An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. Plant J. 24: 265–273.
Google Scholar: Author Only Title Only Author and Title