The AMP deaminase of the mollusk *Helix pomatia* is an unexpected member of the adenosine deaminase-related growth factor (ADGF) family

George Tzertzinis[1], Mehul B. Ganatra[1], Cristian Ruse[1#], Christopher H. Taron[1], Bryce Causey[1], Liang Wang[1], Ira Schildkraut[1]*

[1] New England Biolabs, Ipswich, MA, 01938, USA

#Current Address: Moderna Therapeutics, Cambridge, MA, 02139, USA.

*Corresponding author

E-mail: schildkraut@neb.com

Abstract

We report here the first occurrence of an adenosine deaminase-related growth factor (ADGF) that deaminates adenosine 5' monophosphate (AMP) in preference to adenosine. The ADGFs are a group of secreted deaminases found throughout the animal kingdom that affect the extracellular concentration of adenosine by converting it to inosine. The AMP deaminase studied here was first isolated and biochemically characterized from the roman snail *Helix pomatia* in 1983. Determination of the amino acid sequence of the AMP deaminase enabled sequence comparisons to protein databases and revealed it as a member of the ADGF family. Cloning and expression of its cDNA in *Pichia pastoris* allowed the comparison of the biochemical characteristics of the native and recombinant forms of the enzyme and confirmed they correspond to the previously reported

25   activity. Uncharacteristically, the *H. pomatia* AMP deaminase was determined to be dissimilar to the

26   AMP deaminase family by sequence comparison while demonstrating similarity to the ADGFs

27   despite having AMP as its preferred substrate rather than adenosine.

Introduction

Adenosine 5' monophosphate (AMP) deaminases catalyze the conversion of AMP to inosine 5'

monophosphate (IMP) and ammonia (Fig 1). The AMP deaminases (AMPD) are classified as a subgroup of

the adenyl-deaminase family (1) based on their amino acid sequence similarity. The other members of the

adenyl-deaminase family belong to the adenosine deaminase (ADA), adenine deaminase (ADE), adenosine

deaminase-like (ADAL), and adenosine deaminase-related growth factors (ADGF) subgroups.

The foot muscle of the mollusk, *H. pomatia*, is one of the richest sources (units per mg protein) of AMP

deaminase (HPAMPD) (2). HPAMPD was found to convert 5' AMP, 5' ADP, 5' ATP, and NADH to their

respective inosine derivatives with low micromolar K_m values (3). These characteristics are unlike those of

the animal skeletal muscle AMP deaminases, which have a strict specificity for 5' AMP and relatively high

K_m values, >1 mM. Our initial interest in the HPAMPD came from its high activity and its ability to

deaminate a di-adenosine tetraphosphate (AppppA) to di-inosine tetraphosphate (IppppI) (4).

For this report we purified HPAMPD and subsequently cloned, sequenced and expressed the cDNA

encoding the HPAMPD. We found its amino acid sequence to be most similar to an adenosine deaminase

from another mollusk, *Aplysia californica*. The sea slug, *A. californica,* encodes a biologically and

biochemically characterized adenosine deaminase which was initially designated as mollusk derived

growth factor (MDGF) (5) and later, as a member of the ADGF subfamily (1). The ADGF subfamily is

characterized by secreted deaminases which convert the extracellular nucleoside adenosine, a signaling

molecule, to inosine. This subfamily of adenyl deaminases is represented in humans as the enzyme encoded

by the ADA2 (also known as CECR1) gene. ADA2 is involved in a developmental disorder known as Cat Eye

Syndrome (6). We show, by sequence similarity, that the AMPD from *H. pomatia* is a member of the ADGF

subfamily of adenyl deaminases despite the fact that its preferred substrate is AMP. This is the first report

of a member of the ADGF subfamily of adenyl-deaminases to prefer the nucleotide AMP over the nucleoside

adenosine.

Fig 1. AMP deaminase reaction scheme.

Results

57 In furthering the characterization of the AMP deaminase of *H. pomatia* we first isolated the native

58 protein from *H. pomatia* foot muscle and determined the sequences of the peptides that comprise

59 it by 2D LC-MS/MS (7,8). Using these peptide sequences, we identified the ORF with the highest

60 peptide sequence coverage from a *H. pomatia* transcriptome. The cDNA which encoded the protein

61 was identified, expressed and the resulting recombinant enzyme characterized.

Purification and characterization of the AMP deaminase from *H. pomatia*

The native HPAMPD was purified from the foot muscle from *H. pomatia* as described in Materials

and methods. Deamination was measured by change in absorbance at either OD_{265} or OD_{260} (9). The

activity of the HPAMPD for the deamination of the nucleotides AMP, ATP, and adenosine, to their

inosine equivalents is shown in Fig 2. The rate of deamination of adenosine and ATP is 0.04 and

0.25, respectively, relative to the rate of AMP. The specific activity of the native HPAMPD

preparation was 125 units per mg of protein. However, the protein preparation appeared by PAGE

analysis (S1 Fig.) to comprise a protein of 60 kD and contaminating proteins of 32 and 27 kD. This

result is consistent with the monomer size reported by Stankiewicz (2) at 60-64 kD with an

additional two proteolytic degradation products of the monomer observed here. The specific

activity reported by Stankiewicz (10) was similar at 117 units per mg protein.

Fig 2. Relative rates of deamination. Activity of native *H. pomatia* AMP deaminase on 100 μM 5'

AMP, 5' ATP, and adenosine. The reaction volume was 0.4 ml. The change in absorbance at OD_{265}

was followed in a spectrophotometer at 25° C.

Peptide analysis by mass spectrometry

The purified HPAMPD protein was analyzed using a two-dimensional LC MS/MS system (see

Materials and methods). The *de novo* peptides derived from the mass spectrometry were

intersected with open reading frames (ORFs) that were obtained by translating the *H. pomatia*

transcriptome. Peptide scoring validation was based on q<0.01 (11). Using the peptide sequences,

we identified the ORF from the transcriptome with the highest peptide sequence coverage. The

peptides covered 67% of a 536 amino acid ORF encoded by contig 71391 (Fig 3). The list of

identified peptides from analyses of the digests is detailed in Supporting Information (Table S1).

Fig 3. The amino acid sequence of the *H. pomatia* transcriptome contig 71391. The sequence

segments covered by LC-MS/MS derived peptides are shown in blue text.

In order to determine the sequence and obtain a clone for expressing the protein in a recombinant

form we generated cDNA from *H. pomatia* RNA. The cDNA was used as a template for PCR

amplification using primers designed based on the sequence of contig 71391. The resulting 1596 bp

fragment was sequenced and inserted into the pET28c vector (see Material and methods). Several

nucleotide positions in the cDNA differed from the transcriptome which resulted in nine amino acid

differences from contig 71391 presumably because the transcriptome had been derived from a

different population of *H. pomatia* (S2 Fig.).

The contig 71391 encodes for a protein with an amino terminal end consistent with a signal peptide

as predicted by SignalP-4.1 (12) of MSQVGAVTMVSIICVVVLGAVGGPVAG which *in vivo* should result

in a secreted protein with either an amino terminal sequence of AVGGPVAGLAVRFPT or LAVRFPT.

Both alternative amino terminal versions of the processed protein were submitted to structure

prediction by ColabFold (13). The resulting structures indicated that in the protein with the

additional 8 terminal residues, AVGGPVAG, did not appear to be a structural feature of consequence

(S3 Fig.).

HPAMPD is a member of the ADGF subfamily of adenyl-

deaminases.

The generation of a phylogenetic tree based on amino acid sequence reveals HPAMPD to cluster

within the ADGF subfamily of the adenyl-deaminase family (Fig 4). Consistent with the ADGF

subfamily, the signal peptide of HPAMPD is a characteristic indicating the protein is secreted.

InterPro (14) analysis of the HPAMPD amino acid sequence also indicated it is a member of the

adenosine deaminase-related growth factors (ADGF). The determination that the HPAMPD was a

member of the ADGF subfamily of adenosine deaminases rather than a member of the AMPD

subfamily was surprising because of its preference for AMP over that of adenosine as substrate.

Fig 4. Phylogenetic tree of the adenyl-deaminase family of proteins. The tree was generated with FastTree (15) from a MAFFT alignment. The numerical values at each branch represent substitutions per site. ADGF are the adenosine deaminase-related growth factors, AMPD are the AMP deaminases, ADE are the adenine deaminases, ADAL are the adenosine deaminase-like proteins, ADAR are the adenosine deaminases acting on RNA, and the ADA are the adenosine deaminases. The substrate assigned to each subfamily is indicated on the right. The accession numbers for each enzyme follow: ADGF human, AAF65941.1; ADGF Aplysia, AAD13112.1; ADGF Fish, AF384217.1; ADGF Lottia, XP_009053965.1; ADGF Chicken, AAX10953.1; ADGF Frog, AAX10952.1; ADAL Arabidopsis, NP_192397.2; ADAL human iso3, NP_001311295.1; ADE Aspergillus, AL56636.1; ADE Saccharomyces, NP_014258.1; ADA C elegans, NP_872091.1; ADA human iso1, NP_000013.2; AMPD human, NP_000471.1; AMPD Arabidopsis, NP_565886.1; AMPD2 human, NP_001244290.1; AMPD Drosophila, NP_727740.2.

Observing that HPAMPD is a member of the ADGF subfamily, we used MAFFT (16) to align HPAMPD with representatives of the ADGF subfamily (Fig 5) which includes the most similar member, *A. californica* ADGF, with an identity score of 57% and an additional mollusk, *Lottia gigantea*. The other members span the vertebrates. Fortuitously, there is an experimentally derived three-dimensional structure for the human ADGF, with an identity score of 38%. This allowed us to superimpose it with the predicted HPAMPD structure.

Fig 5. Alignment of the HPAMPD to other ADGF subfamily members. Underlined residues are predicted signal sequences (17) and blue outlined boxes are catalytic residue positions defined by (18) in human ADA2. HPAMPD sequence is derived from cDNA.

Structure comparison of HPAMPD to human and *A. californica* ADGF

Structures of the HPAMPD and *A. californica* ADGF were predicted with AlphaFold2 (13,19) and were compared to the reported experimentally derived human ADGF structure, 3LGG, (18) obtained from the Protein Data Bank (PDB) (20). The PDB files were visualized and aligned with Pymol (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.)(Fig 6a). HPAMPD is aligned to *A. californica* ADGF with a RMSD of 0.594 and to human ADGF with an RMSD of 0.777. Structure alignments of the PDB files with Pymol demonstrate the close structural relatedness of HPAMPD to the two ADGFs. The active site of the human ADGF is elucidated because it formed a cocrystal with coformycin, a transition state analog and potent inhibitor of adenosine deaminases. Based on the structure of the human ADGF, which harbors a tightly coordinated zinc ion at the base of its active site (18) it is expected that the HPAMPD also contains a zinc ion as all four of the zinc-coordinating residues of the human sequence are conserved and aligned in the HPAMPD structure (Fig 6b).

Fig 6. Structural Comparison of HPAMPD and human ADGF. (A) A superimposition of the AlphaFold2 predicted HPAMPD monomer and a monomer of the human ADGF (PDB 3LGG). Human ADGF is colored pink and the HPAMPD is colored cream. Side chains of active site residues and coformycin (colored blue) are drawn as sticks. The zinc ion is represented as a sphere and dashed lines depict interactions with coordinating residues from human ADGF. (B) A close-up view of the active site residues, zinc and coformycin. The labeled residues correspond to the 3LGG structure.

Cloning the cDNA encoding HPAMPD for expression in *Pichia pastoris*

Because the ADGFs as a group are extracellular eukaryotic secreted proteins, we utilized the budding yeast *Pichia pastoris* expression system (21) to express and secrete HPAMPD (see Materials and methods). The processed amino terminus of the HPAMPD is engineered to start at residues LAVR by use of the N-terminal secretion signal from *S. cerevisiae* alpha-factor. The carboxy end is terminated with 10 amino acid histidine residues. The amino acid sequence of the protein as secreted from *P. pastoris* is shown in Fig 7. The molecular weight of the protein as expressed in *P. pastoris* is 58676 daltons which is consistent with the value from SDS-PAGE of 60 kD (S1 Fig.). The specific activity of the recombinant HPAMPD preparation was determined to be 550 units/mg of protein.

Fig 7. The amino acid sequence of the HPAMPD cDNA clone expressed in *P. pastoris*.

Characterization of recombinant HPAMPD activity

The recombinant form of the HPAMPD has a similar spectrum of activity on adenyl substrates as the native enzyme preparations from this study and (2) (Table 1). All three 5' nucleotides, AMP, ADP and AMP are actively deaminated while the absence of a phosphate at the 5' position lowers the activity substantially. Adenosine, 3'5'-cyclic AMP and 3' AMP are poor substrates. This profile is consistent with the presumption that the native and recombinant forms of HPAMPD are one and the same enzyme and represent the same enzyme characterized in 1983.

Substrate	Relative activity from (Stankiewicz 1984) 70 µM substrate	Relative activity of native 100 µM substrate	Relative activity of recombinant 100 µM substrate
5' AMP	100	100	100
5' ADP	95	83	74
5' ATP	36	25	ND
3'5'-cyclic AMP	0.35	2	0.5
adenosine	10 (@3 mM adenosine)	4	2.8
3'-5' adenosine diphosphate (PAP)	8 (@ 3mM PAP)	1.6	0.7
3' AMP	ND	0.9	< 0.1

Table 1. Relative rates of deamination. Spectrophotometric assay of deamination of adenyl base

monitored by loss of OD_{260} or OD_{265}. The rates are linear initial rates and expressed as a percentage

of the 5' AMP rate. Reactions were at 25°C in 20 mM Tris-HCl pH 7.5, 50 mM NaCl and the indicated

concentration of each substrate.

Kinetic constants for recombinant HPAMPD and AMP, K_m and K_{cat}, were determined from linear

initial rates of reaction where 57.5 fmoles of enzyme were incubated in 200 µL reactions with

various concentrations of AMP. The K_m of 25 µM and K_{cat} (turnover number) of 170 per second are

derived from the Lineweaver-Burk plot (Fig 8). Stankiewicz (3) reported a K_m value of 31 µM for 5'

AMP.

Fig 8. Lineweaver-Burke plot for determination of V_{max} and K_m values for recombinant HPAMPD with AMP.

Preincubation of the enzyme at 65° C in the absence of substrate in 10 mM Tris-HCl pH 7.5, 50 mM NaCl, 1 mM EDTA for 30 minutes reduces its activity by more than 95%, while 42° C preincubation has no effect on activity (S4 Fig.). Analysis of thermal denaturation with Prometheus (Nanotemper) indicates the HPAMPD initiates denaturation at 47° C and melts (50 % unfolded) at 54 ° C.

Comparison of *A. californica* and human ADGF activities on adenosine and AMP

The determination that the HPAMPD was a member of the ADGF subfamily of adenosine deaminases was surprising because of its preference for AMP over that of adenosine as substrate. This led us to question the annotation of ADGFs, as we could find only one biochemical evaluation of the deamination of AMP (less than 1%) relative to adenosine in the literature (22). Therefore, we directly compared the activities of HPAMPD, A. *californica* and human homologues on both adenosine and AMP. The relative initial rates of the reactions are listed in Table 2. The results substantiate the assertion that the human and *A. californica* ADGFs deaminate adenosine and do not deaminate AMP, whereas HPAMPD deaminates AMP and does not significantly deaminate adenosine.

	AMP	Adenosine
HPAMPD	1 (1)	0.03 (0.04)

Aplysia ADGF	0.001	1
Human ADGF	< 0.01	1

Table 2. Relative initial rates of deamination of adenosine and AMP for each deaminase. The

deamination was measured by loss of OD_{265} in 10 mM Tris-HCl pH 7.5, 1 mM EDTA, 50 mM NaCl

buffer. The substrate concentrations were 100 µM for the HPAMPD and *A. californica* enzymes and

2 mM for the human enzyme. The reaction temperature for *H. pomatia* and *A. californica* ADGFs was

25° and 37° for human ADGF. All values were obtained with recombinant enzymes except for the

value in parentheses which was obtained with the native preparation of HPAMPD.

Discussion

Here, we report for the first time the identification of an ADGF family member whose substrate

preference is adenosine 5' monophosphate. The biochemical characterization indicates that

HPAMPD deaminates AMP with a K_m of 25 µM and a K_{cat} of 170 per second indicating that this is one

of the more efficient reactions among the adenosine and AMP deaminases. We present evidence

that the AMP deaminase reported in 1983 (3) is the same as the one cloned here from *H. pomatia*.

HPAMPD is unlikely to be the only ADGF which prefers AMP. The few ADGFs which have been

biochemically examined have been shown to deaminate adenosine. Here we demonstrate that the

ADGFs most closely related to HPAMPD, the *A. californica* ADGF and the human ADGF prefer

adenosine to AMP as substrate. ADGFs listed in protein databases are annotated as such primarily

on the basis of sequence similarity. Prediction of gene annotations ideally should utilize full

descriptions of the encoded proteins' biochemical properties, that is in the case of an enzyme, its

substrates and those related compounds that are not (23). However, in practice, that is

understandably an extremely difficult requirement to fulfill. Among the 1777 ADGF family

members listed in the UniProtKB database (Uniprot consortium) only 7 members have been

manually annotated by Swiss-Prot. Of the manually annotated members only 4 have been assayed

for activity on adenosine and only one of those on AMP for which there was less than 1% of the

activity as there was on adenosine (22). Based on the results of our study, it is probable that some

fraction of the 1777 ADGF members prefer AMP as substrate and are therefore incorrectly

annotated.

We suggest that the inability of the human and Aplysia ADGFs to deaminate AMP is not due to the

positioning of the catalytic residues but rather may reflect a steric clash and/or charge repulsion of

the AMP phosphate. The HPAMPD has the same positioning of the catalytic residues as human

ADGF. Clearly the same chemistry is taking place for deamination for all the ADGFs. The HPAMPD

preference for AMP could be explained with a binding pocket that can accommodate the phosphate

of AMP which contributes to substrate binding. Further work in this lab will attempt to elucidate

the basis for the adenosine vs. AMP preference.

The ADGFs are involved in purinergic signaling. They are characterized as secreted growth factors

and affect growth by modulating the extracellular levels of adenosine. We have shown that

HPAMPD has an encoded signal peptide. Therefore, it is likely secreted in *H. pomatia,* and its site of

action is extracellular. We propose that HPAMPD affects the level of extracellular AMP and is

involved in purinergic signaling in some as yet unknown role. Skaldin (24) reported a bacterial

ADGF homologue which is secreted and deaminates adenosine and suggested that the ADGFs

evolved from a secreted bacterial homologue with a similar role involved in bacterial

communication.

The occurrence of purinoreceptors in primitive invertebrates suggests that ATP is an early

evolutionary extracellular signaling molecule (25,26). Perhaps *H. pomatia* uses an alternative

pathway for modulating extracellular levels of ATP. For example, rather than converting ATP to

AMP and then AMP to adenosine and deaminating adenosine to inosine as a last step, *H. pomatia*,

removes AMP directly by deamination to IMP by HPAMPD as a last step.

In conclusion, HPAMPD is an AMP deaminase that is assigned by sequence similarity to the

adenosine deaminase-related growth factor family which suggests that there is a biological role of

AMP deaminases as growth factors. Furthermore, HPAMPD is a prime example for illustrating that

the assignment of an enzymatic activity based only on sequence similarity in absence of

biochemical evidence is imprudent.

Materials and methods

Purification of native AMPD from *H. pomatia*

The *H. pomatia* were obtained from My Happy Snails (myhappysnails.com) and fed on romaine

lettuce for 5 days. The foot muscle from 2 snails was dissected. 1 gram was stored at -70° C and was

later used for RNA extraction and cDNA cloning. The remaining 5 grams was homogenized in 20 ml

of 20 mM Tris-HCl pH 7.5, 1 mM EDTA, 1 mM DTT, 1M NaCl. The homogenate was clarified by

centrifugation and NaCl concentration was reduced to 200 mM by dilution with 20 mM Tris-HCl pH

7.5, 1 mM EDTA, 1 mM DTT. The clarified extract contained 600 AMP deaminase units.

The clarified extract was applied to a 5 ml Q sepharose column where 80% of the AMP deaminase

flowed through the column. The Q flow-through was applied to a 5 ml HiTrap™ Heparin column. A

gradient of NaCl was run to 1.2 M and the AMPD activity eluted at about 0.4M NaCl. The peak of

activity was pooled and diluted to 0.1 M NaCl and subsequently applied to a 5 ml HiTrap™ Q

column in 20 mM Tris-HCl pH 7.5, 1 mM EDTA, 1 mM DTT, 0.1 M NaCl. The activity flowed through

the column and was then applied directly onto a 5 ml HiTrap™ SP column. A gradient to 1 M NaCl

was run and the activity eluted at about 0.5 M NaCl. An aliquot (10 μL) of a peak fraction was

subjected to *de novo* peptide sequence analysis. The remainder of the AMP deaminase active

fractions were pooled and represent the native enzyme preparation with a yield of 30% at 200

units with a specific activity of about 160 units/mg protein.

AMPD assay

The assay is based on Kalckar's spectrophotometric method (9) where the decrease in optical

density at 265 nM is measured. The reaction buffer is 20 mM Tris-HCl @ pH 7.5, 50 mM NaCl and

typically the substrate concentration is 100 μM AMP. The reaction volume ranged from 200 to 400

μL. The optical density was determined with a Spectramax M3. One unit of the enzyme activity is

defined as the amount of the enzyme which deaminates 1 μmole of 100 μmolar 5' AMP/min in 20

mM Tris-HCl pH7.5, 50 mM NaCl at 25° C.

H. pomatia transcriptome

The *H. pomatia* snails were obtained from a farmers' market in Germany and stored at -80°C.

Approximately 100 mg of *H. pomatia* body were frozen in liquid nitrogen and ground to a powder.

Total RNA was purified from rehydrated powder with the RNeasy Plant Mini Kit (Qiagen,

Chatworth, CA). An mRNA transcript library for Illumina sequencing was created using the

NEBNext rRNA Depletion Kit (Human/Mouse/Rat) (New England Biolabs, Ipswich, MA). A NextSeq

500 sequencer (Illumina, San Diego, CA) was used to do paired-end deep sequencing (2X 150 bp) of

the library. The utility CutAdapt (27) was used to remove adapter sequences from the raw reads.

The trimmed reads were then assembled using the Trinity software package (28). The assembled

transcriptome data was deposited at DDBJ/EMBL/GenBank under the accession GKIM00000000.

The version described in this paper is the first version, GKIM01000000. Raw reads were deposited

in the NCBI Sequence Read Archive (accession #PRJNA936131).

Peptide sequencing by 2D LC-MS/MS

The purified HPAMPD protein fraction was digested in solution in parallel, with either trypsin at a

ratio of trypsin to protein at 1:50 or subtilisin at a ratio of 1:160. The subtilisin digested fraction

was analyzed twice, to include charge state 1 peptides. The trypsin digest was analyzed with one

injection. Each digest was directly injected onto an analytical C18 column and analyzed with Easy

n1000 nLC-LTQ Orbitrap (Thermo Fisher Scientific, Waltham, MA) at 300 nL/min. All MS/MS

spectra were analyzed with ProteomeDiscoverer version: 2.0.0.802 in one workflow. The fasta

database consisted of open reading frames translated from the *H. pomatia* transcriptome. Peptide

scoring validation was based on q<0.01 (11).

RNA isolation for cDNA

The source of *H. pomatia* for cDNA was the same as for the HPAMPD enzyme preparation. About 1

gram of frozen *H. pomatia* foot tissue was pulverized in liquid nitrogen with mortar and pestle.

About 0.2 grams of powdered tissue was homogenized in 1 ml of TRIzol reagent (Thermo Fisher).

The debris was pelleted, and 0.2 ml of chloroform was added per ml of supernatant vortexed for 30

sec and phases separated by a 15 min centrifugation. 1 volume of ethanol was added to the aqueous

layer and the mixture was applied to and eluted from a Direct-zol column (Zymo RNA miniprep kit)

yielding 25 µg of RNA.

cDNA cloning

First strand cDNA was generated from *H. pomatia* RNA with Protoscript II (NEB) and random

primers. The first strand cDNA was used as a template for PCR amplification using primers A and B

(see below) which were designed based on the transcriptome derived sequence for HPAMPD. The

resulting 1556 bp fragment was inserted into the pET28c vector digested with EcoRI and NcoI

using the NEBuilder HiFi Assembly kit (NEB).

Primer sequences:

A: ACT TTA AGA AGG AGA TAT ACC ATG CTC GCC GTC AGA TTT CCG

B: AAG CTT GTC GAC GGA GCT CGT ACC GAT AAC TTT CTT GTT GGT TAG

Construction of expression cassette for *P. pastoris*

pD912-P$_{GAP}$ plasmid was obtained from ATUM Bio (Newark, CA). A linear expression cassette of

pD912-P$_{GAP}$-Hp AMPD-His$_{10}$ was assembled using the NEBuilder HiFi DNA Assembly Cloning Kit

(NEB #E5520S). Primers were designed using the NEBuilder Assembly Tool. The HPAMPD

fragment was amplified with the forward primer C and reverse primer D (containing a 10X His-tag

sequence) using plasmid pET28c-Hp AMPD as template. The pD912-P$_{GAP}$ fragment containing the

transcriptional terminator (TT), ILV5 promoter, and zeocin resistant gene was amplified with the

forward primer E and reverse primer F using pD912-P$_{GAP}$ plasmid as template. All products were

purified by gel extraction. The linear expression cassette was then amplified with forward primer G

and reverse primer H. The gel extracted product was used to transform electrocompetent *P.*

pastoris D aox1 (MutS) cells (ATUM Bio.) and plated on yeast peptone dextrose agar medium

supplemented with 1 M sorbitol and 500 µg/mL Zeocin (Teknova, Hollister, CA.).

Eight to twelve transformants were patched onto fresh selection plates and incubated for an

additional 1-2 days at 30°C. For the identification of transformants by PCR, genomic DNA was

isolated from each strain using lithium acetate/sodium dodecyl sulfate (LiOAc/SDS) method (29).

PCR was used to identify transformants having an integrated expression cassette.

Primer sequences:

C: CGA GAA AAG AGA GGC CGA AGC TCT CGC CGT CAG ATT TCC G

D: TTG AGC GGC CGC CCC TTC AAC CTC AGT GGT GGT GGT GGT GGT GGT GGT GGT GGT GAC CGA

TAA TTT CTT GTT GGT TAG

E: CAC CAC CAC CAC CAC CAC CAC CAC CAC CAC TGA GGT TGA AGG GGC GGC CGC

F: AGC TTC GGC CTC TCT TTT C

G: GCT CAT TCC AAT TCC TTC TAT TAG

H: GAG CTC CAA TCA AGC CCA ATA AC

P. pastoris clone harboring HPAMPD cDNA

The *P. pastoris* transformants that had integrated the expression cassette coding for the HPAMPD were grown in liquid cultures at 30°C in 5 mL of BMGY-Buffered Glycerol Complex Medium (Teknova) (1% yeast extract, 2% tryptone, 1.34% yeast nitrogen base without amino acids with ammonium sulfate, 0.0004% biotin, 1% glycerol as the carbon source, 100 mM potassium phosphate, pH 6.0). After 48 hours, the spent culture media was harvested. The culture fluid from 4 cultures one of which was a non-transformant control culture were each concentrated about 10-fold and assayed for AMP deaminase activity. One transformant had no detectable deaminase activity while the other two converted about 25% and 90% of the AMP to IMP in 4 minutes at 25° C.

Purification of recombinant HPAMPD

The culture identified with the highest HPAMPD activity was then grown in 2400 ml of Tolner media (30) with 1% glycerol and 100 µg/ml ampicillin at 30° C for 72 hours. The culture was clarified by centrifugation and the supernatant culture fluid was brought 75% ammonium sulfate saturation and centrifuged to clarify. The purification was performed at 4° - 10° C except where noted. The 3.5-liter supernatant was then diluted with 700 ml of water and applied to a 250 ml bed Phenyl Sepharose column. After loading the column with supernatant, the column was washed with 500 ml of 3M ammonium sulfate, 20 mM Tris-HCl pH 7.5, 1mM EDTA and 1 mM DTT. A reverse gradient of 1800 ml from 2M to 0 M ammonium sulfate was applied to the column. The HPAMPD activity that eluted from the column was pooled (120 ml) and diluted with 2 liters of 20 mM Tris-HCl pH 7.5, 25 mM NaCl, 1mM EDTA, 1 mM DTT. This was applied to a 10 ml Heparin HyperD column washed with 20 mM Tris-HCl pH 7.5, 1 mM EDTA, 1 mM DTT, 50 mM NaCl and eluted with a 400 ml gradient from 50 mM to 1 M NaCl in 20 mM Tris-HCl pH 7.5, 1 mM EDTA, 1 mM DTT. The peak of activity was pooled and half of the pool was treated with Endo Hf (NEB) by

first dialyzing the 25 ml against 2 changes of 500 ml of 20 mM Na acetate pH 6.0, 1 mM EDTA, 50

mM NaCl, 1 mM DTT. Five hundred thousand units of EndoHf was added to the 25 ml dialysate and

incubated at 25° C overnight. A small amount of precipitate formed and was pelleted and removed

by centrifugation. In order to remove the EndoHf from the HPAMPD the supernatant was brought

to 100 mM Tris -HCl pH 7.5 and 100 mM NaCl and applied to a one ml amylose resin (NEB) column

at a rate of about 1 drop per second at 25° C. The flow through containing the HPAMPD was then

applied directly to a 5 ml HiTrap™ Q column in line with a 5 ml HiTrap™ SP column and washed

with 20 mM Tris-HCl pH 7.5, 1mM EDTA and 1 mM DTT, 50 mM NaCl. The Q column was removed

and a 50 ml gradient from 50 mM to 1M NaCl was applied to the SP column. The peak of HPAMPD

activity was pooled and dialyzed against 50% glycerol, 10 mM Tris-HCl pH 7.4, 1 mM DTT, 0.1 mM

EDTA. The recombinant HPAMPD preparation is stored at -20° C. The protein is >95%

homogenous by PAGE (S1 Fig.) and protein concentration is 2.6 mg/ml based on OD_{280} with

an extinction coefficient of 85,000 M^{-1} cm^{-1}.

Human and *A. californica* ADGF

403 The human ADGF (Recombinant Human Adenosine deaminase 2/CECR1) #7518-AD-010 was from

404 R&D Systems, Minneapolis, MN and the *A. californica* ADGF was purified from construct described

405 below. The *P. pastoris* clone carrying the *A. californica* ADGF cDNA was grown and purified as

406 described for the *H. pomatia* construct except deglycosylation was not performed.

Yeast culture conditions and expression

Construction of the *A. californica* construct for expression in *P. pastoris* was performed as follows:

The transformation cassette for *A. californica* ADGF, was generated by PCR using primers 1941 and

1942 and as a template the clone in PD912-GAP which was obtained from Genscript, Piscataway, NJ.

The PCR product was confirmed for size by agarose gel electrophoresis and subsequently purified

by QIAquick (Qiagen) columns. The purified product (1 μg) was electroporated into *P. pastoris*

competent cells and positive colonies identified as described above.

Primer sequences:

1941: GCT CAT TCC AAT TCC TTC TAT TAG

1942: GAG CTC CAA TCA AGC CCA ATA AC

Acknowledgments

420　We thank Sean Johnson for a critical reading of the manuscript, the DNA sequencing core lab (NEB),

421　Saulius Vainauskas for advice with *Pichia*, Vladimir Potapov for help with data deposition, Tom

422　Evans, Richard Roberts and Donald Comb for encouraging basic research at New England Biolabs.

References

1.　Maier SA, Galellis JR, McDermid HE. Phylogenetic Analysis Reveals a Novel Protein Family

Closely Related to Adenosine Deaminase. J Mol Evol. 2005 Dec;61(6):776–94.

2.　Stankiewicz AJ. Non-specific Adenylate Deaminase from Snail Foot Muscle. Adva Exp Med Biol.

1984;165 pt B:457–64.

3.　Stankiewicz AJ. Direct deamination of AMP, ADP, ATP and NADH by non-specific adenylate

deaminase in the foot muscle of the snail *Helix pomatia*. Biochem J. 1983 Oct 1;215(1):39–44.

4.   Guranowski A, Starzynska, E, Sillero, MA Gunther, Sillero, A. Conversion of adenosine 5'

oligophospho 5' adenosines into inosine 5" oligophospho 5' inosines by non-specific adenylate

deaminase from the snail Helix pomatia. Biochim Biophys Acta. 1995;1234:78–84.

5.   Akalal DBG, Nagle GT. Mollusk-derived growth factor: cloning and developmental expression in

the central nervous system and reproductive tract of Aplysia. Mol Brain Res. 2001 Jul;91(1–

2):163–8.

6.   Riazi MA, Brinkman-Mills P, Nguyen T, Pan H, Phan S, Ying F, et al. The Human Homolog of

Insect-Derived Growth Factor, CECR1, Is a Candidate Gene for Features of Cat Eye Syndrome.

Genomics. 2000 Mar;64(3):277–85.

7.   Bouzid W, Verdenaud M, Klopp C, Ducancel F, Noirot C, Vétillard A. De Novo sequencing and

transcriptome analysis for Tetramorium bicarinatum: a comprehensive venom gland

transcriptome analysis from an ant species. BMC Genomics. 2014;15:987.

8.   Evans VC, Barker G, Heesom KJ, Fan J, Bessant C, Matthews DA. De novo derivation of

proteomes from transcriptomes for transcript and protein identification. Nat Methods. 2012

Dec;9(12):1207–11.

9.   Kalckar HM. Differential spectrometry of purine compounds by means of specific enzymes. J

Biol Chem. 1947 Feb;167(2):461–75.

10. Stankiewicz A. Nonspecific Snail Muscle Adenylate Deaminase: Simplified Purification,

Characterization and Use for the Preparation of Deamino Derivatives of NAD, NADH and AMP-

P(NH)P. Enzyme. 1986;36(3):187–96.

11. Savitski MM, Wilhelm M, Hahne H, Kuster B, Bantscheff M. A Scalable Approach for Protein

False Discovery Rate Estimation in Large Proteomic Data Sets. Mol Cell Proteomics. 2015

Sep;14(9):2394–404.

12. Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides from

transmembrane regions. Nat Methods. 2011 Sep 29;8(10):785–6.

13. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making

protein folding accessible to all. Nat Methods. 2022 Jun;19(6):679–82.

14. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al. InterPro in 2022.

Nucleic Acids Res. 2023 Jan 6;51(D1):D418–27.

15. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for Large

Alignments. Poon AFY, editor. PLoS ONE. 2010 Mar 10;5(3):e9490.

16. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements

in Performance and Usability. Mol Biol Evol. 2013 Apr 1;30(4):772–80.

17. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, et al. SignalP

6.0 predicts all five types of signal peptides using protein language models. Nat Biotechnol.

2022 Jul;40(7):1023–5.

18. Zavialov AV, Yu X, Spillmann D, Lauvau G, Zavialov AV. Structural Basis for the Growth Factor

Activity of Human Adenosine Deaminase ADA2. J Biol Chem. 2010 Apr 16;285(16):12367–77.

19. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein

structure prediction with AlphaFold. Nature. 2021 Aug 26;596(7873):583–9.

20. Berman H, Henrick K, Nakamura H. Announcing the worldwide Protein Data Bank. Nat Struct

Mol Biol. 2003 Dec;10(12):980–980.

21. Sreekrishna K, Potenz RHB, Cruze JA, McCombie WR, Parker KA, Nelles L, et al. High level

expression of heterologous proteins in methylotrophic yeast Pichia pastoris. J Basic Microbiol.

1988;28(4):265–78.

22. Iijima R, Kunieda T, Yamaguchi S, Kamigaki H, Fujii-Taira I, Sekimizu K, et al. The Extracellular

Adenosine Deaminase Growth Factor, ADGF/CECR1, Plays a Role in Xenopus Embryogenesis

via the Adenosine/P1 Receptor. J Biol Chem. 2008 Jan;283(4):2255–64.

23. Zhao Y, Wang J, Chen J, Zhang X, Guo M, Yu G. A Literature Review of Gene Function Prediction

by Modeling Gene Ontology. Front Genet. 2020 Apr 24;11:400.

24. Skaldin M, Tuittila M, Zavialov AV, Zavialov AV. Secreted bacterial adenosine deaminase is an

evolutionary precursor of adenosine deaminase growth factor. Perna N, editor. Mol Biol Evol.

2018 Oct 16;35(12):2851–61.

25. Vultaggio-Poma V, Falzoni S, Salvi G, Giuliani AL, Di Virgilio F. Signalling by extracellular

nucleotides in health and disease. Biochim Biophys Acta BBA - Mol Cell Res. 2022

May;1869(5):119237.

26. Burnstock G. Purine and purinergic receptors. Brain Neurosci Adv. 2018

Jan;2:239821281881749.

27. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal. 2011;17:10–2.

28. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol. 2011

Jul;29(7):644–52.

29. Lõoke M, Kristjuhan K, Kristjuhan A. Extraction of genomic DNA from yeasts for PCR-based

applications. BioTechniques. 2011 May;50(5):325–8.

30. Tolner B, Smith L, Begent RHJ, Chester KA. Production of recombinant protein in Pichia pastoris

by fermentation. Nat Protoc. 2006 Aug;1(2):1006–21.

5' AMP $\rightarrow$ 5' IMP + NH₃

Figure

Figure

```
RSVVRMSQVGAVTMVSIICVVVLGAVGGPVAGLAVRFPTMDEYTNAREELIGSEQYLRVG   60
GSITLNDKEKKLNQFILREKRAIIENSRLNKTQYIPAVSFFLSKSQMESTPIFKIIKDMP  120
KGAALHLHDTASARIDWIVSNATYRDHVYMCMDRDNFVRLTVSGTGLPADSGCEWKLVET  180
ERANSGDIAAFDHWLKSNISLLTTDPLVTYPSLDKVWGRFDKHFSQLRGIIYHTPIRRDY  240
YRQILEEFRSDNVQYVEVRSSLSGYYDLDGTVHDPEYGLQLYKAVTEEFVRTYPDFSGAK  300
IIKSTARVKPNTDVFNDVKLSMDLYKRYPGFFLGFDLVAQEDPNTPLLGYIDSLLYPSRQ  360
NPPVSLPYYFHAGETNWQGTEVDYNLVDALLLNATRIGHGFALIKHPRVIELVKSRGVAV  420
EVNPVSNQLLGLVKDLRNHAAAPLLAQNVPVVISSDAPGVWEALPMSHDMYVAFMDLVGE  480
DAGLDVLKQLVWNSIQYSSMNATEKRTALKLLQAKWNKFINDSLIKWKLTNKKVIG      536
```

Figure

AMP

adenosine

N⁶mAMP

adenine

adenosine

AMP

RNA

Figure

```
HPAMPD                   MSQVGAVTMVSIICVVVLGAVGGPVAGLAVRFPTMDEYTNAREELIGSEQYLRVGGS
ADGF Aplysia californica MSSFSTHNFVAIATFVCWFCCLATAAPLTSKAAYLLKRNSLIEEDASRKLGAK
MLGF Haliotus discus     MTYSCFKAYFPLLLLNVLFLHIDCMSISAYTLERQRIMKSEESDMFGSS
ADGF Lottia gigantea     MTKGLFIVFLTVWQWNYCVCVPASYQRQRDAIVQKEMDMRIGSH
ADGF Homo sapien         MLVDGPSERPALCFLLLAVAMSFFGSALSIDETRAHLLLKEKMMRLGGR
ADGF Xenopus laevis      MACVSHWLLLLSLASATLSRPLWSERNGLIEMENSIRLGGN
ADGF Gallus gallus       MKPPSGQGIRLACLLGLASLCLPTPLWEDRDSLMQEEEARLTGGN
ADGF Danio rerio         MHVLFLGDLMWIYLLLLCCASCNGRPDPRKRDALIGLEASRRTGGD

                         60          70          80          90          100         110
HPAMPD                   INLNNKEKKLNQFILREKRAIIENSRLNKTQYIPAVSFFLSKSQMESTPIFKIIKDM
ADGF Aplysia californica IVLTNEEKVLDDFILAEKRKLIDDSRLNQTEYMPAASFYRSKDFIDTTFAYKIIQDM
MLGF Haliotus discus     LVLNKQEGLISDILIKAKRAEIEDSRVNGTIFPPAINFFESKKLIDSSKVFQIIKRM
ADGF Lottia gigantea     WNLTDDEKIVNNIFLGAKKNDYFNARVMNSFYPPAQNFFQSKKFIDKSLVFQMIEKM
ADGF Homo sapien         LVLNTKEELANERLMTLKIAEMKEA-MRTLFIFPPSMHFFQAKHLIERSQVFNILRMM
ADGF Xenopus laevis      IILTPSEATANQKLMTVKGAEFKEA-ESTGLFPPSMHFFKARPLIQQSHVFSILHQM
ADGF Gallus gallus       LVLSTQEQQLSMKLISLKKKEVAAA-LNTGQFPPSMHFFRAKSLIEQSAVFSILKRM
ADGF Danio rerio         ITLNEREKLLDGKLQKLKQHDMEA----GQFPPSMHFFKAKRLIDQSPVFNLIQKM

                         120         130         140         150         160         170
HPAMPD                   PKGAALHLHDTASARIDWIVSNATYRDHVYMCMDQ-DNFVRLTVSGTGP--PANSGC
ADGF Aplysia californica PKGGALHLHDLAIASLDWVVKNATYRDNVYMCMDK-DNDVNLRVLQLIP--P-DPFC
MLGF Haliotus discus     PKGGALHTHGSAITSLDWLVKNVTYRANCYMCHDN-QGHLQMHFYDTPP--Q-PSYC
ADGF Lottia gigantea     PKGCALHIHDSSVASLDWFIQNVTYRDNLYICFNSVSGKPSLHFYSKAP--A-NPDC
ADGF Homo sapien         PKGAALHLHDIGIVTMDWLVRNVTYRPHCICFTP-RGIMQFRFAHPTP--RPSEKC
ADGF Xenopus laevis      PKGGALHLHDFAILSVDWLVKNASYMADCYMCLTR-DGGVRFLFAKPAPVGMLPPGC
ADGF Gallus gallus       PKGAVLHLHDYAMLSVEWLVYNATYLPDCYICFTQ-TGTVRFRFSKPHPPHPVPAQC
ADGF Danio rerio         PKGAALHVHDFAMVSVDWLVKNVTYRENCYVCFTD-KQTVQFIFSSGPP--ASSSHC

                         180         190         200         210         220
HPAMPD                   -EWKLVETERANSGDIAAFDHWLKSNISLLTTDPLVTYPSLDKVWGRFDKHFSQLRG
ADGF Aplysia californica -VWKLVATERANSGDVEAFDDWLKKNISYLSTDPVTQYATVDSVWVRFNKYFAQVIG
MLGF Haliotus discus     -QWESVQHVRERSGNALLFDQQLIKNISIIVDDPHSTYVTVNDVWRRFGLYFKIVNG
ADGF Lottia gigantea     -YWELISAVRKRSGDAKVFDEELYKNLSIIVDNPDDVYKTVDDVWQKFNDYFSLLNG
ADGF Homo sapien         SKWILLEDYRKRVQNVTEFDDSLLRNFTLVTQHPEVIYTNQNVWSKFETIFFTISG
ADGF Xenopus laevis      SEWILLETYRKKLGDVTEFDKGLIRNLTLLTDSPEPHIPSQDEIWRRFEGAFITASG
ADGF Gallus gallus       SQWVLLETYRKQLRNVTEFDHSLLRNLTLVTDAPELAYPSQAFIWKRFEEVFLIASG
ADGF Danio rerio         SSWTLLRSLREKIKNTTELDNSFIRNLTLFTEDPDRAYPNQDTVWERFEQVFLVAYG

                         230         240         250         260         270         280
HPAMPD                   IIYHTPIRRDYYRQILEEFRSDNVQYYVEVRSSLSGYYDLDGTVHDPEYGLQLYKAVT
ADGF Aplysia californica LLFYAPIMRDYYRQALEEFRADNVQYIELRSQLFGFFELDGTVHDAEFGLNLYKSVT
MLGF Haliotus discus     LYRYTPIFKDYLQQALREFEEDNVQYIEIRATLSPLYELDGSKHDQEHKLLTYRDTV
ADGF Lottia gigantea     AFQYVPVFQDYFRRALSEFYQDNVQCVELRVTLSPIYELDGSVKSPEWIVNAYRTIA
ADGF Homo sapien         LIHYAPVFRDYVFRSMQEFYEDNVLYMEIRARLLPVYELSGEHHDEEWSVKTYQEVA
ADGF Xenopus laevis      LICYAPVFKEYFYESLRELYEDNIQYLEMRAMLPPVYELDGTVHDQFWSMAIYRDMA
ADGF Gallus gallus       LICYAPVFKAYFYQGLLELYEDNIQYVEIRAILPAVYELDGTQHNKSWSVAAYQEVS
ADGF Danio rerio         LVTYAPVFKDYLYEGLRQFYEDNIMYVEIRALLPQTYELDGRLNDKDWSMAACQEVV

                         290         300         310         320         330         340
HPAMPD                   EEFVRTYPDFSGAKIIKSTARVK--PNTDIFNDVKLSMDLYKRYPGFFLGFDLVAQE
ADGF Aplysia californica EEFQREYPDFIGAKILSGLRFK--SQEEILNEVKIAMDLHKKYPDFFLGYDLVGQE
MLGF Haliotus discus     NDYMAAHPDFIGCKLIQSGIRFF--KRDDILDDIKSYVKYQQKYSDIIRGFDLVGQE
ADGF Lottia gigantea     KEFIQQNPDFLGVKLIYSGIRYVFFPKPDILKDVKTAMSLIQKYKDFVVGFDLVGQE
ADGF Homo sapien         QKFVETHPEFIGKIIYSDHRSK--DVAVIAESIRMAMGLRIKFPTVVAGFDLVGHE
ADGF Xenopus laevis      NKFVGAHPDFLGAKIIYTVHRHE--DLAQVTEAVHLAMKLMEAFPEIMAGFDLVGQE
ADGF Gallus gallus       RRFVQEHPDFLGAKVIFAAHRML--NVSQIKETIITAMELRAHFPDTLAGFDLVGNE
ADGF Danio rerio         NQFKKHHPDFIGARVIFTVHRKI--NATEAVKTVEEAMILRRNFPDVMAGFDFVGQE

                         350         360         370         380         390
HPAMPD                   DPNTSLLGYIDSLLYPSRQNPPVSLPYYFHAGETNWQGTEVDYNLVDALLLNATRIG
ADGF Aplysia californica DPNFSLLHYLDALLYPSIQNPPYRLPYFFHAAETNWQGTEVDYNLADALLLNTTRVG
MLGF Haliotus discus     GPGKPLSYLDDLLYPSTLDPPVSVPYFFHAGETDWEGSQVDLSLVDAVLLNTSRIG
ADGF Lottia gigantea     GLGKPLLYYLDALLYPSQQSPPVNLPYFFHAGETKWQDTIVDYNLANTTRIG
ADGF Homo sapien         DTGHSLRDYKEALMIPAKDG--VKLPYFFHAGETDWQGTSIDRNILDALMLNTTRIG
ADGF Xenopus laevis      DAGHSLYQLSDALNIPSKLG--VKLPYFFHAGETNWQGKDVDENVLDALLLNTTRIG
ADGF Gallus gallus       DEGHSLWEFKDVLTIPYSLR--ANLPYFFHAGETDWQGTSVDNNVLDALILNTSRIG
ADGF Danio rerio         DLGRPLWYFKDALSLPEDKG--FNLPYFFHAGETDSQGTDVDQNLMDALLFNTTRIG

                         400         410         420         430         440         450
HPAMPD                   HGFALIKHPRVIELVKSRGVAVEVNPVSNQLLGLVKDLRNHAAAPLLAQNVPVVISS
ADGF Aplysia californica HGFALIKHPRFTELAKENGVAVEVNPISNQILGLVRDVRNHALVPLIADDYPIVISS
MLGF Haliotus discus     HGFALYKHPEVMKMLKQKDIAIEVNPISNQVLKLVEDLRNHGAITMFANDYPLVVGS
ADGF Lottia gigantea     HGFALTKHPYIKNMVEKRGIAVELNPISNQVLKLVDDIRNHPGAILMTKKFPVVVSS
ADGF Homo sapien         HGFALSKHPAVRTYSWKKDIPIEVCPISNQVLGLVSDLRNHPVATLMATGHPMVISS
ADGF Xenopus laevis      HGYALKHPVARNLSLELNVPIEICPISNQVLGLVSDLRNHPAAVLMAEGHPLVVSS
ADGF Gallus gallus       HGLALTKHPVAKNLSLKRDIPVEVCPISNQVLLLVSDLRSHPAASLMAEGHPVVVSP
ADGF Danio rerio         HGFALARHPVVKEMARKMYVPIEVCPISNQVLKLVSDLRDHPAAVLMAEGHPLVISS

                         460         470         480         490         500         510
HPAMPD                   DDPGVWEALPMSHDMYVAFMDLVGEDAGLDVLKQLVWNSIQYSSMNATEKKTALKLL
ADGF Aplysia californica DDPGAWEASPLSHDFYVALMDLCGRDTALTFLKQLALNSIRYSAMSDTEKVAAKAKW
MLGF Haliotus discus     DDPAAWGALPQSHDFYMAYMAMTREDAGLEVLKQLAINSMQYSAMTSSEKMEAMKKW
ADGF Lottia gigantea     DDPAVWGALPLSHDFYVAFMGMCGADEGIAILKQLAYNSIQYSAMTNAEKTNAFQSW
ADGF Homo sapien         DDPAMFGAKGLSYDFYEVFMGIGGMKADLRTLKQLAMNSIKYSTLLESEKNTFMEIW
ADGF Xenopus laevis      DDPSIFGAQGISYDFYEVFMGIGGAKADLRTLKLKAENSIKYSALSKEGKELTEIW
ADGF Gallus gallus       DDPSIFGAKGVSYDFYEVFMAIGGMNADLRTLKLKIKNSIKYSAMLLDEKAKAREVW
ADGF Danio rerio         DDPAVFGATALSHDFYEAFMGFGGMSFNLGTLKELALNSLRYSTLSSQRKEEAIDAL

                         520         530         539
HPAMPD                   QAKWNNFINDSLIKWKLTNKKVIG
ADGF Aplysia californica TTQWDKFVKTSVEGLKPHINDRS
MLGF Haliotus discus     ERKWNEFIIEILRQHSHDGQLIG
ADGF Lottia gigantea     EQRWNAFIKDLVRQHHSRITSLSIVG
ADGF Homo sapien         KKRWDKFIADVATK
ADGF Xenopus laevis      QKKWDKFIKDLAMNWKKEL
ADGF Gallus gallus       QKKWDQFVADLTDSSLMEL
ADGF Danio rerio         LVKWDKFVWESLL
```

Figure

Figure

```
LAVRFPTMDEYTNAREELIGSEQYLRVGGSINLNNKEKKLNQFILREKRAIIENSRLNKT 60
QYIPAVSFFLSKSQMESTPIFKIIKDMPKGAALHLHDTASARIDWIVSNATYRDHVYMCM 120
DQDNFVRLTVSGTGPPANSGCEWKLVETERANSGDIAAFDHWLKSNISLLTTDPLVTYPS 180
LDKVWGRFDKHFSQLRGIIYHTPIRRDYYRQILEEFRSDNVQYVEVRSSLSGYYDLDGTV 240
HDPEYGLQLYKAVTEEFVRTYPDFSGAKIIKSTARVKPNTDIFNDVKLSMDLYKRYPGFF 300
LGFDLVAQEDPNTSLLGYIDSLLYPSRQNPPVSLPYYFHAGETNWQGTEVDYNLVDALL 360
NATRIGHGFALIKHPRVIELVKSRGVAVEVNPVSNQLLGLVKDLRNHAAAPLLAQNVPVV 420
ISSDDPGVWEALPMSHDMYVAFMDLVGEDAGLDVLKQLVWNSIQYSSMNATEKKTALKLL 480
QAKWNNFINDSLIKWKLTNKKVIGHHHHHHHHHH 514
```

Figure

Figure