**FULL TITLE:** The orphan nuclear receptor Nr2f6 impairs skeletal muscle oxidative capacity and

regulates muscle mass in mice.

**SHORT TITLE:** Nr2f6 regulates muscle biology

**AUTHORS:**

Dimitrius Santiago Passos Simões Fróes Guimarães[1,2,*]

Ninon Melany Flores Barrios[1]

André Gustavo de Oliveira[1]

David Rizo-Roca[2]

Maxence Jollet[3]

Jonathon A.B. Smith[2]

Thiago Reis Araujo[1]

Marcos Vinicius da Cruz[1]

Emilio Marconato Junior[1]

Sandro Massao Hirabara[4]

André Schwambach Vieira[1]

Anna Krook[2]

Juleen R. Zierath[2,3]

Leonardo dos Reis Silveira[1]

**AFFILIATIONS:**

[1]Department of Structural and Functional Biology, University of Campinas, Campinas, Brazil

[2]Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden

[3]Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden

[4]Interdisciplinary Post-Graduate Program in Health Sciences, Cruzeiro do Sul University, São Paulo,

Brazil

**\*Correspondence**

Dimitrius Santiago P.S.F. Guimarães

Centro de Pesquisa em Obesidade e Comorbidades

Universidade Estadual de Campinas - Instituto de Biologia

Rua Carl Von Linnaeus - Bloco Z

CEP 13083-862

Campinas - SP

Brazil

F: +55 (19) 3521-6191.

E-mail: dimitrius.guimaraes@ki.se; d211529@unicamp.br

**ABSTRACT**

The maintenance of skeletal muscle plasticity upon changes in the environment, nutrient supply, and exercise depends on regulatory mechanisms that couple structural and metabolic adaptations. However, the transcriptional control of both processes by nuclear receptors (NR) remains underexplored. Nr2f6 is an orphan NR and a key regulator of metabolism and differentiation. Nonetheless, its role in muscle biology remains elusive. Here, we report, for the first time, the effects of Nr2f6 modulation in skeletal muscle *in vivo* and *in vitro*. Depletion of Nr2f6 increased myocyte's oxidative capacity and sharply attenuated lipid-induced cell death, which was associated with direct derepression of uncoupling protein 3 and PGC-1$\alpha$ promoters' transactivation. Conversely, Nr2f6 overexpression in the *tibialis anterior* induced atrophy and hypoplasia, accompanied by impairment of force production and the establishment of a molecular signature of inflammation, and a decrease in genes involved in oxidative metabolism and contraction. Additionally, global transcriptomics showed that Nr2f6 upregulated core components of the cell division machinery and repressed myogenesis genes, thus decoupling myoblast proliferation from differentiation. Collectively, our findings define a novel role for Nr2f6 as a molecular transducer maintaining the balance between skeletal muscle contractile function and oxidative capacity, with implications for metabolic diseases and myopathies treatment.

**KEYWORDS:** muscle atrophy; Nr2f6; skeletal muscle; transcription; energy metabolism; nuclear receptor;

## INTRODUCTION

Muscle contraction is a highly coordinated process initiated by the transmission of the action potential from the afferent neurons to the muscle. Subsequent depolarization of the muscle fiber triggers an action potential that propagates along the sarcolemma, stimulating the release of calcium ions from the sarcoplasmic reticulum, and ultimately promoting myosin-actin cross-bridge cycling and force production. This process requires an accessory metabolic machinery to generate the energy needed to support contraction, and disruption in either muscle structure or metabolism leads to functional defects, such as in Duchenne syndrome[1], [2], sarcopenia[3], and cachexia[4]. Dynamic crosstalk between energetic status, muscle development, mechanical stress, and transcriptional changes is crucial to maintain muscle function. In this context, the nuclear receptor family of transcription factors (NR) is of particular interest since they are regulated by small molecules, such as metabolites and hormones[5]. Although the transcriptional landscape for metabolic-functional signaling has been extensively studied in pathological and physiological conditions, a broader role of some NRs has only recently been recognized, and thus, the role of many members remains elusive[6].

The NRs have a modular architecture that contains a ligand-binding domain (LBD) and a zinc-finger DNA binding domain (DBD), and can be further grouped in endogenous, orphan, or adopted NRs according to the presence of an endogenous ligand, the absence of a known ligand or if a new ligand for a given NR was just identified, respectively[7]. The classical mechanistic model proposes that a small molecule binds to the LBD, changing its conformation to one of higher affinity for a transcriptional co-regulator[8], such as PGC-1α, which in turn can mediate transcriptional modulation through the recruitment of histone acetylases, the mediator complex, and basal transcriptional apparatus.

The orphan nuclear receptor Nr2f6 (also named Ear2 or COUP-TFIII) has been characterized in a broad range of tissues, such as adipose, thyroid, liver, brain, and the immune system, where it plays different and even antagonistic roles[9]. Nr2f6 can impair adipocyte differentiation, increase cancer cell proliferation, induce both resistance and susceptibility to antitumor drugs, and promote the development

of fatty liver disease[10]. In ovarian cancer cells, Nr2f6 binds to the histone acetylase P300 at the

Notch3 promoter, increasing histone H3 K9 and K27 acetylation to activate transcription, increasing cell

proliferation and chemoresistance[11]. So far, the most extensively defined role of Nr2f6 is in the

immune system, in which it directly and strongly suppresses interleukins 17, 21, 2 and interferon γ

transcription by interacting with the NFAT/AP-1 complex at the promoters of these genes[12], [13].

Curiously, Nr2f6 has been reported both as a transcriptional repressor and activator, but the context that

defines its activity state is unknown. Recently, Nr2f6 was classified as a stripe transcription factor[14],

i.e. it can bind to low-complexity motifs together with other transcription factors in a broad range of

promoters, regulating chromatin accessibility. This indicates that the function of Nr2f6 in diverse

environments is influenced not only by its DNA occupancy but also by the presence and activity of

other transcription factors. Whether the current understanding of the role of Nr2f6's can be applied to

other tissues, such as skeletal muscle is unknown. Therefore, we sought to characterize the molecular

mechanisms and functional roles of Nr2f6 in skeletal muscle biology both in vitro and in vivo. We

discovered that Nr2f6 overexpression in skeletal muscle disrupts oxidative metabolism by directly

repressing PGC-1α and the uncoupling protein 3 (UCP3) gene expression, which impairs mitochondrial

function. Moreover, Nr2f6 induces core genes of cell cycle progression and in skeletal muscle activates

immune cells, increasing inflammation, reducing mass, changing fiber type, and reducing force

production, collectively causing a sarcopenic-like state.

**RESULTS**

**Nr2f6 regulates muscle cell metabolism and differentiation.**

Genetic manipulations of Nr2f6 at the whole-body level and *in vitro* have been conducted[10], [11],

[23], [24], but the role of this NR in muscle models is underexplored. We used siRNA-mediated

depletion of Nr2f6 in C2C12 myocytes to verify the outcomes on the transcriptomic landscape

(Supplementary Figure 1A). The 1849 differentially regulated genes, 920 upregulated and 939

downregulated, could be grouped into five main classes, with increased expression of genes related to

muscle differentiation, contraction, and metabolism and decreased expression of genes with roles in cell cycle and DNA packaging (Figure 1A-D). In fact, among the 20 most significative altered genes, nine are linked to muscle contraction (*RYR1, TTN, MYH3, MYH2, ACNT2, ATP2A1, MYL1, TNNC2, MYOM3*) are upregulated by Nr2f6 knockdown (Figure 1A). Accordingly, a panel of canonical markers of muscle differentiation containing muscle regulatory factors (MRFs) and myosin isoforms (Figure 1E) reveals that the Nr2f6 knockdown enhanced C2C12 differentiation. Indeed, data generated in our transcriptomic analysis was correlated with a publicly available C2C12 myogenesis dataset (Supplementary Figure 1B). Consistently, the proteins coded by the upregulated genes belonged mostly to the sarcomere, contractile fiber, and cytoplasm location ontologies. Since both an increase in cellular oxidative capacity and the activation of the PI3K pathway are required for myogenesis[25], [26], we verified whether genes of the main pathways of regulation of glycolysis and fatty acid oxidation were affected and found that several energy sensors such as *AKT2*, *PRKAG3* subunit of AMPK, and mTOR complex were upregulated by Nr2f6 loss-of-function (Figure 1F). Myogenic differentiation demands the withdrawal of the cell cycle and both processes are regulated in a concerted manner[27], [28]. Accordingly, the downregulated genes were related to different phases of cell division, with the enrichment of proteins related to DNA replication, packaging, and chromosome separations. Of note, essential components of the cell cycle progression such as *CDC25B/C* phosphatases and *CDK1/4* kinase promote quiescence and halt cell cycle progression in muscle progenitors when down-regulated[29]–[31]. Altogether, the changes in the myocyte's global transcriptome indicate that Nr2f6 inhibits myoblast differentiation and metabolism. Considering the differences between mouse and human myogenic cell's transcriptional landscape[32] we verified whether the result of Nr2f6 depletion in differentiation markers would be translatable to human primary human skeletal muscle cells. We found that expression of *MYH1/2/7*, muscle creatine kinase, and myosin light chain kinase 1 were upregulated in human cells (Figure 1G), indicating a conserved role for Nr2f6 as repressor of myogenesis.

**Increase in cell oxidative capacity by Nr2f6 knockdown.**

Given the enrichment of genes of oxidative metabolism, we first sought to verify the functional effects

of Nr2f6 knockdown on metabolism. Oxygen consumption assays (Figure 2A), using palmitate as the

major substrate for energy production, provided evidence that Nr2f6 knockdown promoted an increase

in maximal respiration capacity after the addition of the uncoupler (carbonyl cyanide m-chlorophenyl

hydrazone) CCCP and the spare capacity. Although there was no difference between control and

knockdown respiratory parameters in high-glucose media (Supplementary Figure 2A, B), the

extracellular acidification rates were reduced, without a reduction of total ATP pool, which was

confirmed by lower lactate concentration in knockdown cells (Figure 2B, C, S2C). These results are

supported by alterations in the expression of the pyruvate carboxylase, insulin-dependent glucose

transporter, and fatty acid transporters (Figure 2D, E). Together, these results indicate Nr2f6 knockdown

increases pyruvate and acetyl-CoA flux to the mitochondria, which could be reinforced by upregulation

of the glucose transporter *GLUT4*, the anaplerotic enzyme pyruvate carboxylase (PC), and fatty-acid

transporters. Our analysis of ENCODE chromatin immunoprecipitation-sequencing (ChIP-seq) data for

Nr2f6 in K562 and HepG2 cells indicates an increase in mitochondrial genes related to lipid metabolism

(Supplementary Figure 1C). We also identified significant enrichment of kinases related to the insulin

signaling pathway when analyzing genes affected by Nr2f6 knockdown in the RNA-seq data and genes

with Nr2f6 binding within the promoter region in ENCODE with ChIP-seq data (Supplementary Figure

1C, D). Considering the increase in the efficiency of the Nr2f6 silenced myocytes to oxidize lipids, we

hypothesized that depletion of Nr2f6 would be protected against lipid overload in skeletal muscle.

Stable Nr2f6 knockdown in myotubes protected (50%) against palmitate-induced cell death and reduced

mitochondrial superoxide production (40%) and decreased cytosolic reactive oxygen species (20%)

(Figure 2F, G, H). Since Nr2f6 knockdown can increase lipid handling capacity by upregulating

mitochondrial and cytosolic lipid transporters, mitochondrial proteins, and TCA cycle anaplerotic genes,

thereby increasing oxygen consumption, we verified whether Nr2f6 is modulated by palmitate treatment

*in vitro* and by high-fat diet *in vivo* in rodents[33], both pathophysiological conditions of increased lipid

oxidation and supply (Figure 2I-K, S2D, E). Our findings demonstrate that Nr2f6 expression is

consistently reduced under these conditions, indicating a role as an energy stress response gene that

facilitates metabolic adaptations to lipid oxidation. Collectively, the data provide evidence that Nr2f6

inhibition protects against lipid overload by increasing lipid handling capacity in skeletal muscle.

**Nr2f6 directly represses PGC-1α and UCP3 expression.**

We recently provided evidence that the transcriptional regulation of *UCP3* by the peroxisome

proliferator-activated receptor γ co-receptor 1-α (PGC-1α) is essential for the maintenance of myotube

viability during lipid overload, by preventing the production of reactive oxygen species (ROS) [34],

[35]. Considering a similar phenotype by Nr2f6 knockdown in myotubes, we investigated whether

Nr2f6 regulates the same pathway. We found that Nr2f6 overexpression reduced both *UCP3* and *PGC-*

*1α* mRNA in myotubes, which also translated to both reduced PGC-1α protein content and expression of

its mitochondrial electron transfer chain (ETC) target genes (Figure 3A-B). Using luciferase reporter

assays, we observed Nr2f6 overexpression reduced PGC-1α promoter activity, suggesting direct

repression. Conversely, Nr2f6 knockdown in C2C12 myotubes increased PGC-1α downstream ETC

targets (Figure 3D). Further investigation into the regulation of *UCP3* transcription by Nr2f6 using 7kbp

UCP3 promoter reporter plasmid showed a reduction in the luciferase signal by Nr2f6 overexpression

and an increase in activity following Nr2f6 knockdown (Figure 3E). We scanned the UCP3 promoter

region and found an Nr2f6 response element downstream of the transcription initiation site, which

coincided with the open chromatin region and peaks of known the UCP3 transcription factors, Myod1

and Myogenin (Figure 3F), further supporting the notion that Nr2f6 directly repressed *UCP3* expression.

The effects of Nr2f6 knockdown on *UCP3* and *PGC-1α* expression could be reproduced in human and

mouse primary myotubes (Figure 3G, S3A). UCP3 transcription is regulated by peroxisome proliferator-

activated receptors (PPARs) and estrogen-related receptors (ERRs) in skeletal muscle[36], [37],

however, Nr2f6 silencing did not change the transactivation of responsive elements (Supplementary

Figure 3B). Collectively, our results indicate that Nr2f6 is a bona fide transcription regulator of UCP3

and PGC-1α. Given that UCP3 is a PGC-1α target, these results indicate that Nr2f6 represses *UCP3*

expression indirectly by downregulating *PGC-1α* and by directly binding to the *UCP3* promoter region.

**Nr2f6 activates the cell cycle and represses the expression of genes involved in muscle contraction**

**and oxidative metabolism.**

We next explored the effects of Nr2f6 overexpression *in vivo* by electroporation in the *tibialis anterior*

muscle of mice, using the contralateral muscle as control, and studied the global transcriptomic changes

by microarray. There were 3796 genes within the criteria for differential expression (FDR <0.05, Fold

change >2), among which, 1915 were downregulated and 1781 were upregulated, with Nr2f6

overexpression having a major effect on the hierarchical clustering (Figure 4A). Consistent with earlier

reports that highlight Nr2f6 as a gatekeeper of the immune system[24], gene ontology analysis of the

upregulated genes shows enrichment of biological processes and pathways related to the immune system

(Figure 4B). RT-qPCR was used to validate markers modulated in microarray analysis. We found that

indicators of lymphocyte activation *CD44*, the marker for macrophage/monocyte activation *CD68*, and

the macrophage marker *F4-80* were upregulated in Nr2f6 expressing muscle (Figure 4E). Accordingly,

*TGFb*, a potent inhibitor of hematopoietic cell activation, and the marker for endothelial and non-

differentiated hematopoietic cells were downregulated, consistent with an increase in the number and

activity of immune system-derived cells, indicating that Nr2f6 might activate resident cells of the

immune system and/or promote the invasion of circulating cells. Nr2f6 overexpression also increased

expression of Myogenin and, to a lesser extent, *MYOD*, however the downstream targets genes myosin

heavy chains 1 and 2 decreased (Figure 4D). Downregulated genes were enriched in energetic

metabolism pathways, mitochondria, and muscle contraction terms (Figure 4C), reinforcing the

functional phenotypes described *in vitro*. Importantly, the hereby proposed Nr2f6 targets, namely *UCP3*

and *PGC-1α*, were downregulated by Nr2f6 overexpression (Figure 4F). The lipid transporters *CD36*

and *CPT1B*, as well as subunits of the respiratory chain complexes, which were upregulated by Nr2f6

knockdown *in vitro*, were also downregulated by Nr2f6 overexpression. Additionally, the expression of

reactive oxygen species scavengers *SOD1*, *SOD2*, and catalase genes was decreased (Figure 4F).

Collectively, these findings support our functional results and indicate mitochondrial function was

impaired by Nr2f6 overexpression.

**Nr2f6 inhibits muscle development and contraction.**

Since Nr2f6 overexpression negatively affects the mRNA expression of genes involved in muscle

contraction and development, we next investigated whether the Nr2f6 gain-of-function would impair

muscle morphology and function by performing immunostaining for myosin heavy chain (MHC)

isoforms and *ex vivo* contraction experiments, respectively. Intriguingly, Nr2f6 overexpressing *tibialis*

*anterior* (TA) weighted less and were visually paler compared with control muscle (Figure 5A).

Consistent with these observations, Nr2f6 overexpression reduced the total number of fibers (21%),

which together with the increase in cell death-related genes (Figure 4B), and the increase in the

atrogenes cathepsin and calpain 2[38] (Supplementary Figure 4A), characterizes a state of atrophy

(Figure 5B, C). Stratification by fiber type showed that this reduction is particularly due to the decrease

in type IIB fibers, which were reduced by 23%, and although there was a tendency to decrease IIX fiber

number (Figure 5D), there was no statistical significance in these comparisons or the number of IIA

fibers. We then overexpressed Nr2f6 in the *flexor digitorum brevis* (FDB) and performed *ex vivo*

contractions to verify alterations in muscle force production and fatigability. Consistent with the

immunostaining data, mass-corrected maximal force production was reduced (60%) (Figure 5E), but

time to fatigue was unaltered in Nr2f6 overexpressing muscles (Supplementary Figure 4B). Since the *ex*

*vivo* contraction assay surpasses the neuromuscular system by direct electric stimulation, disregarding

action potential issues, fatigability is mostly induced by detriments in calcium homeostasis, such as

reduced $Ca^{2+}$ sensitivity, sarcoplasmic $Ca^{2+}$ reuptake, and release[39]. Therefore, we cannot exclude the

possibility that the time to fatigue is also affected *in vivo* in Nr2f6 gain-of-function models. So far, these

findings strongly suggest induction of atrophy, worsened by an inflammatory state and an imbalance

between satellite cell proliferation and differentiation.

**Nr2f6 modulates myoblasts' proliferation rates.**

Next, we investigated whether Nr2f6 regulates cell cycle genes and myoblast proliferation. Thus, we compared differentially expressed genes identified in the microarray of the Nr2f6 overexpression in TA muscle with the RNA-seq transcriptomics from C2C12 myocytes after transient Nr2f6 knockdown. We found 706 genes were differentially expressed in both experiments, whereby 446 genes were modulated in opposite directions indicating a direct regulation by Nr2f6 or a conserved effect of Nr2f6 modulation in both models (Figure 6A). We further scanned the promoter regions of these genes consisting of 3 kbp upstream and downstream of the transcription start site, for the Nr2f6 binding motif. We found 206 matches in unique genes, whereby 73 were upregulated and 133 downregulated by Nr2f6 overexpression. The interaction network of these high-confidence targets (Figure 6B) reveals that the most connected genes are upregulated by Nr2f6 overexpression and are mostly related to the cell cycle, which emphasizes the role of Nr2f6 as a promoter of cell division, and further reinforces the dysplastic phenotype observed in the gain-of-function experiments *in vivo*. Investigation of canonical pathways of myogenesis cell proliferation and stemness in Nr2f6 overexpressing TAs (Figure 6C, D) showed that the content of the proliferating cell nuclear antigen (PCNA), an important general marker for cell proliferation[40], was increased by Nr2f6 overexpression. Together with the increase of the muscle-specific satellite cell marker Pax7, and the activation by phosphorylation of the stemness markers GSK3a/b[1], [41], ERK[42], [43], and S6, these results point to an increase in myogenic progenitors and the infiltration of other cell types, such as cells of the immune system. Nr2f6 overexpression and knockdown can promote or inhibit cell proliferation in cancer cells, respectively[44]–[46]. Our doubling time experiments (Figure 6E, G) confirm this effect is also conserved in C2C12 myoblasts, with an increase of 4 hours in the average doubling time in knockdown cells and a decrease of 3.5 hours in Nr2f6 overexpression stable cell lines. Moreover, RT-qPCR validation of major markers of cell cycle progression inhibitors *RB1* and *P21* show an increase in both genes by Nr2f6 knockdown and a tendency towards downregulation of *RB1* by Nr2f6 overexpression (Figure 6F, H). Collectively, these

**11**

262 results indicate that the function Nr2f6 function as an important repressor of cell cycle progression is

263 conserved in muscle models.

**DISCUSSION**

Numerous nuclear receptors are necessary for the maintenance of muscle mass[47], [48]. For example,

whole-body knockout of Nr1d1 (Rev-ERBα, Ear-1) leads to an increase in atrophic genes, a decrease in

muscle mass, and a relative increase in low-diameter fibers[49]. More broadly, muscle-specific

knockout of the nuclear receptor co-repressor 1 (NCoR1) leads to skeletal muscle hypertrophy and

increased oxidative metabolism[50]. Here, we found evidence that a disruption in myogenesis also

follows Nr2f6 overexpression *in vivo* and *in vitro,* and myoblast proliferation rates are increased.

Remarkably, Nr2f2 (COUP-TFII), an Nr2f6 interactor, is among the few nuclear receptors found to

promote muscle wasting[51], [52]. However, Nr2f2 expression in myogenic progenitors impairs muscle

differentiation in mice by directly repressing genes related to myoblast fusion and proliferation[53],

[54], implying that myogenesis is disrupted in a different stage. Future studies should address the

redundancy of these NRs in muscle function.

Most of the transgenic and knockdown models provide evidence to suggest that nuclear receptors are

involved in a general activation of oxidative metabolism[47]. For example, the muscle-specific Nr4a3

transgenic mouse displays a marked increase in mitochondrial density and fast-to-slow fiber switch[55].

This general rule is reinforced by the fact that mice lacking nuclear receptor co-activators, such as PGC-

1α and MED1[56], or overexpressing the co-repressor RIP140[57], have decreased mitochondrial

density and fewer oxidative fibers. Conversely, here we show that Nr2f6 is an exception to this model in

skeletal muscle since this nuclear receptor can directly reduce PGC-1α and UCP3 promoter activity,

thereby increasing the susceptibility of muscle cells to lipid overload by reducing fatty-acid oxidation

and increasing reactive oxygen species production. Transgenic mouse models overexpressing UCP3 in

muscle are consistently reported to have improved glucose homeostasis under chow and high-fat diet

(HFD) conditions, as well as resistance to obesity-induced diabetes[58]–[60]. Moreover, increased

levels of circulating lipids increase UCP3 expression[61], [62]. Here, we demonstrate that direct fatty-

acid exposure in C2C12 myotubes or conditions of increased β-oxidation *in vivo* reduces Nr2f6 mRNA

expression and protein content, further reinforcing the finding that Nr2f6 mediates the positive effects of

UCP3 under physiological conditions. Importantly, we show that the regulation of *UCP3* and *PGC-1α*

expression by Nr2f6 is conserved in human skeletal muscle cells.

Sarcopenia is the age-related loss of muscle mass and function, with the reduction of the number and

size of myofibers, a switch from type II to I fibers[63], and an underlying mitochondrial

dysfunction[64]. This phenotype is closely reproduced by Nr2f6 overexpression in skeletal muscle. In

contrast to most members of the NR family, Nr2f6 overexpression not only induces a sarcopenic-like

phenotype, with loss of muscle mass and inflammation but also reduces muscle strength and affects

energy metabolism. Interestingly, Nr2f6 is upregulated 2-fold in skeletal muscle hereditary spastic

paraplegia[65], a disease characterized by progressive lower limb muscle weakness, sometimes

accompanied by mitochondrial dysfunction and morphological fiber defects[66], [67]. While additional

experiments are warranted to assess the effects of Nr2f6 ablation *in vivo*, Nr2f6 knockdown in myotubes

increases myosin heavy chain expression and improves mitochondrial function, suggesting that

inhibition of Nr2f6 might be efficacious in the treatment of sarcopenia or other myopathies. Nr2f6

agonists have been proposed as a possible treatment for colitis[68], but based on our finding that Nr2f6

gain-of-function alone provokes muscle loss, the use of such agonists should be further evaluated for the

treatment of patients suffering from myopathies and cachexia. Considering the known effects of Nr2f6

in the inflammatory response, the atrophic phenotype may be supported by a role of Nr2f6 in immune

cells. Nonetheless, the antagonistic transcriptional changes induced by Nr2f6 overexpression *in vivo,*

and knockdown *in vitro* strongly indicate a direct action of Nr2f6 in the myofibers as the major driver of

the functional changes.

A conceivable model elucidating the mechanism by which Nr2f6 overexpression culminates in a

reduction of muscle force production (Figure 7), entails the downregulation of genes engaged in various

**13**

facets of muscle contraction, such as muscle structure, calcium cycling, and action potential. Notably, a

few of these genes constitute high-confidence targets. The ryanodine receptor 1 (*RYR1*) is a major

component of the calcium release complex, which permits calcium efflux from the sarcoplasmic

reticulum into the cytosol. As such, *in vivo* knockdown and mutations of *RYR1* can cause severe

myopathies[69]. Other putative targets including myosin light chain kinase 4 (*MYLK4*) and myomesin 1

(*MYOM1*) are downstream of the androgen receptor (AR), which mediates the effects on muscle force

production[70]. Interestingly, we found that the spermine oxidase gene (*SMOX*), another important

target of the AR in muscle[71], [72] is downregulated by Nr2f6 overexpression *in vivo*, upregulated by

Nr2f6 silencing in C2C12 myocytes, and contains Nr2f6 binding motifs at the promoter region. The

direct link between Nr2f6 and the spermine synthesis pathway and a possible interaction with the AR

warrant further studies.

Targeted studies provided evidence that Nr2f6 activates gene expression by tethering to the promoters of

*circRHOT1*[73], *DDA1*[74], and *CD36*[10], and represses the expression of numerous others such as

*IL17*, *IL21*, Renin, and Oxytocin[44], [45], [75], [76]. More broadly, our transcriptomics experiments

display an equilibrated number of genes up- and downregulated, further implying that Nr2f6 is a dual-

function transcription factor. Interestingly, Nr2f6 is a target of MiR-142-3p[77], raising the possibility

that miRNAs, besides protein partners and post-translational modifications[44], might aid in the

regulation of Nr2f6 activity. Further studies should help to elucidate the mechanism by which Nr2f6 acts

as a repressor or activator of gene expression. The case of CD36 illustrates context-dependent regulation

given that this gene is activated in the liver[10], but repressed in skeletal muscle by Nr2f6 in mice and

humans. This finding suggests that Nr2f6 may be bound to DNA but kept in a repressive state by post-

translational modifications or interaction partners, such as RAR Related Orphan Receptor γ (RORγ),

another regulator of *CD36* transcription in muscle[78] and liver[79]. In Th17 lymphocytes, Nr2f6 can

compete with RORγ for binding at the *IL17* promoter, maintaining the repressive state. This relationship

may also be present in skeletal muscle.

**14**

As reported for other cell types[11], Nr2f6 also modulates myoblast proliferation *in* vitro and increases

the expression of proliferation markers such as *PCNA* and *KI67 in vivo*. In Nr2f6 overexpressing

muscle, Myogenin expression is also increased. However, the myosin heavy chains and other indicators

of terminally differentiated myofibers are sharply reduced. These findings raise the possibility that the

Nr2f6-overexpressing myoblasts proliferate but fail to assemble into robust myofibrils due to a

dysregulated temporal modulation of the MRFs during myogenesis, which critically impairs muscle

fiber formation and force production. Key regulators of cell cycle cyclin B1 and Cdk1 are placed among

high-confidence direct targets of Nr2f6. Cyclin B1 interacts with Cdk1 and is necessary for kinase

activity[80] and progression through the G2 mitotic phase. Moreover, Ccnb1 overexpression is

increased in several cancer types and ectopic expression increases cell proliferation rates [81], [82]. The

simultaneous effect of Nr2f6 modulation on cell cycle and differentiation markers might also be

sustained indirectly by the expression of the retinoblastoma protein, which is responsible for cell cycle

arrest through the inhibition of the E2F family of transcription factors. In a feedback loop, E2F TFs

antagonize MyoD1, which induces Rb expression, thereby linking processes of proliferation and

differentiation[83], [84].

Our findings provide evidence that Nr2f6 plays a critical role in the regulation of several aspects of

muscle biology. Nr2f6 modulation alone can determine myoblast proliferation rates, consolidating its

role as a major regulator of cell cycle progression. In summary, we report that Nr2f6 is a novel regulator

of muscle contraction and metabolism, which may hold promise as a possible strategy for the treatment

of muscle wasting and metabolic diseases.

**MATERIALS AND METHODS**

The main reagents, tools, and models necessary for replicating the reported results are listed in

Supplementary Table 2.

**Cell culture** Human primary skeletal muscle cells were isolated from healthy female and male donors

(Al-Khalili et al. 2004), age 55 ±5 years old, BMI 25.6 ±1.5 kg.m$^{-2}$. Myoblasts were maintained in

**15**

Growth Media (DMEM/F12 High Glucose (Gibco, #31331093) supplemented with 10 mM HEPES

(Gibco #15630-056), 16% Fetal calf serum (Sigma, #F7524), and antibiotics (Gibco #15240-062) and

differentiated at the confluence with fusion media (74% DMEM High Glucose (Gibco, 31966-021),

20% 199 Medium (Gibco #31150-022), 20 mM HEPES, antibiotics, 0.03 µg/mL Zinc Sulfate (Sigma

#Z4750), 1.4 mg/mL Vitamin B12 (Sigma #V6629), and 2% Fetal Calf Serum) supplemented with

100ug/mL Apotransferrin (Biotechne #3188-AT-001G) and 1.7 mM Insulin (Actrapid Penfill, Novo

Nordisk #13509) before use. After 5 days of fusion, apo transferrin and insulin were removed from the

media, and cells were incubated for 4 more days. Cells were cultivated in a humidified atmosphere

containing 7.5% $CO_2$ and regularly tested for mycoplasma. C2C12s, MEFs, and HEK cells were

maintained in DMEM High Glucose (Gibco, 31966-021) supplemented with 4 mM L-glutamine, 10%

fetal bovine serum, 1 mM sodium pyruvate, and antibiotics. Fetal bovine serum was substituted by 2%

horse serum to induce myogenesis in C2C12 cells when 90-100% confluence was reached, and

experiments were performed 5 days later.

**Primary mouse skeletal muscle cells** Mice's primary skeletal muscle cells were isolated from wild-type

C57Bl6/JUnib as described (Araujo et al. 2020). After euthanasia, hindlimb muscles were dissected and

digested with collagenase II, trypsin, and DNAse I. Cells were sifted through a 70 µm cell strainer and

plated in 0.1% Matrigel-coated plates. Myoblasts were maintained for 2 days in DMEM High Glucose

supplemented with 2 mM L-glutamine, 10% fetal bovine serum, 10% horse serum, 1 mM sodium

pyruvate, and antibiotics. Myogenesis was induced by removing fetal bovine serum from the media

when confluence was reached, and cells were cultivated for 5 more days. The experiments were

approved by the Ethics Committee on Animal Use (CEUA/Unicamp #5626-1/2020).

**Animals** All electroporation experiments were conducted following the guidelines of animal welfare

and were approved by the Stockholm North Animal Ethical Committee (Stockholm, Sweden). Male

C57Bl6/J mice were acquired from Jackson Labs and maintained at 12/12h light/dark cycle under

controlled temperature and humidity, and *ad libitum* access to food (Specialized Research Diets, #

**16**

801722) and water. The use of animals for high-fat diet experiments was approved by the Ethics

Committee on Animal Use (CEUA/Unicamp #5626-1/2020) and all the welfare guidelines of the

National Council of Control of Animal Experimentation (CONCEA) were followed. Male

C57Bl6/JUnib mice were kept under the same conditions described above. Mice were provided a high-

fat diet (PragSolucoes #0015, 60% kcal from lipids) at 4 weeks of age for 16 weeks; littermates were fed

a standard chow diet as a control.

**Reactive oxygen species measurement** Cells were incubated with 5 nM MitoSOX (Invitrogen,

#M36008) or 5 µM Dihydroethidium (DHE, Invitrogen, #D11347) in DMEM without phenol red

supplemented with 1 mM Sodium Pyruvate, 4 mM L-glutamine, and 25 mM Glucose for 30 min and

washed three times before reading in a plate reader 510/580 nm (ex/em) or 520/610 nm (ex/em) for

DHE. Samples were fixed and stained with Crystal Violet for normalization.

**RT-qPCR** Total RNA was extracted from cells with TRIzol (Invitrogen #15596-018) following the

manufacturer's instruction and cDNA was synthesized with a High-Capacity Reverse Transcription kit

(Applied Biosystems #4368814). cDNA was diluted to 10 ng/µL and 20 ng was used for qPCR

reactions. NormFinder (Andersen, Jensen, and Ørntoft 2004) was used to decide the best combination of

internal controls among RPL39, PPIA, HPRT, 18S, ACTB, and GAPDH. In the *in vivo* electroporation

experiments, gene expression was normalized using HPRT-PPIA geomean with TaqMan probes or

HPRT-RPL39 geomean when SYBER green was used. For other experiments, gene expression was

normalized with multiplexed HPRT when TaqMan probes were used or with RPL39 when SYBER was

used. Relative expression was calculated by the $\Delta\Delta C_T$ method (Livak and Schmittgen 2001) and is

expressed as fold change over the indicated control. The primers and probes used are listed in

Supplementary Table 1.

**RNA-seq** Total RNA was extracted with TRIzol and the upper phase containing RNA was loaded into

RNeasy columns (Qiagen, #74004) after the addition of isopropanol, following the manufacturer's

instructions. cDNA libraries were prepared with TruSeq Illumina Total RNA Stranded (Illumina) with

**17**

Ribo-zero rRNA depletion (Illumina). Sequencing was outsourced to Macrogen Inc. (Seoul, South

Korea) and performed in a HiSeq X (Illumina), producing an average of 50.8 Mreads, 95% above Q30.

Sequence trimming and adapter removal were done with Trimmomatic (Bolger, Lohse, and Usadel

2014) with the following modifications: HEADCROP = 10, MINLEN = 20, AVGQUAL = 20.

Reminiscent reads were aligned to the mouse genome (Ensembl GrCm38) with RNA Star 2.7.2b and

gene-level counts were calculated with featureCounts v1.6.4. Differential expression was performed

with EdgeR with TMM normalization and p-value adjustment using Benjamini and Hochberg

normalization with a 0.05 false discovery rate (FDR) cut-off. The Galaxy platform was used to process

all data. Pathway enrichment analysis was done in g:Profiler with an FDR cutoff of 0.01. Interaction

networks were generated by String.db and analyzed with CytoScape v3.8 using the EnrichmentMap

plugin.

**Palmitate treatment** Palmitate (Sigma, #P5585) in absolute ethanol was conjugated with 1% fatty-acid-

free bovine serum albumin (Sigma, #A7030) in cell media for 15 min at 55 $^{o}$C to a 500 µM final

concentration. Cells were treated with fresh solutions of palmitate or vehicle (1%BSA, 1% ethanol) for

20 hours.

**Promoter transactivation assays** Luciferase reporter assays were performed in MEF cells transfected

with a UCP3 reporter plasmid (Harmancey et al. 2015) (UCP3 EP1, Addgene #71743) or PGC-1α 2kb

promoter (Handschin et al. 2003) (Addgene #8887), normalization plasmid coding for *Renilla* luciferase

(pRL-SV40) and either control empty vector or Nr2f6 coding plasmid (Gift from Dr. Gottfried Baier,

Medical University of Innsbruck, Austria) using Lipofectamine 3000. Luciferase activity was measured

with DualGlo Luciferase Reporter Assay (Promega, #E2920). For knockdown assays, cells were

transfected with siRNAs one day before the transfection of the reporter plasmids.

**siRNA knockdown** C2C12 cells were transfected with 200 nM non-target siRNA (siScr, Qiagen) or

siNr2f6 (Sigma) using Lipofectamine RNAiMax (Invitrogen) following manufacturer's instructions

concomitantly with the myogenesis media switch. Experiments were performed on the third or fiftieth

day of differentiation. Primary human skeletal muscle cells were transfected at the fiftieth day of

differentiation with 5 nM siScr (Ambion) or siNr2f6 (Ambion).

**Stable cell lines** The Nr2f6-myc insert was subcloned from the Nr2f6-myc-flag plasmid into the

pBABE-Puro vector using standard PCR with primers spanning the transcription start site and the myc

tag. The viral particles for generating the overexpression HEK293T cells were transfected with pCMV-

VSVG, pCL-Eco, and the pBABE-Nr2f6-myc or empty vector. For producing viral particles with the

knockdown plasmid, HEK293T cells were transfected with packaging vectors pCMV-dR8.2 dvpr,

pCMV-VSVG, and pLKO.1-shGFP or shNr2f6 (TRCN0000026147). Cell medium containing virions

was collected, filtered at 0.45 µm, and stored at -80 $^{\circ}$C until further use. Virus concentrations were

titrated by the minimal dilution method, C2C12 cells were transduced with 1 MOI, and cells were

selected with 2 µg/mL puromycin for 4 days. The clonal selection was performed in the knockdown

cells and the clones were validated as indicated. The modified cells and their respective controls were

cultivated synchronously under the same conditions.

**Electroporation** Mice were kept under 2% isoflurane-induced anesthesia and the *tibialis anterior*

muscles were injected with 30 µL of 1 mg/mL hyaluronidase (Sigma, #H3506). After 2 hours, the lateral

and contralateral *tibialis anterior* were injected with 30 µg of either control empty vector pCMV6 or

Nr2f6-myc-flag overexpression plasmid (Origene, #MR206083) and 220V/cm were applied in 8 pulses

of 20/200 ms on/off (ECM 830 Electroporator, BTX). Terminal experiments were performed 9 days

after electroporation with 13 weeks old mice. For electroporation of FDB muscles, after anesthesia 10

µL of 1 mg/mL hyaluronidase were injected into the footpads and after 1 hour, 20 µg of the control or

Nr2f6 coding plasmids. Mice rested for 15 minutes and then 75V/cm were applied in 20 pulses of 20/99

ms on/off with the aid of sterile gold acupuncture needles.

**Contraction** Mouse FDB muscles were electroporated as described and dissected 8 days later. With the

muscles still attached to the tendons, contraction threads were tied at the most distal and proximal

tendons, and the muscles were transferred to contraction chambers containing prewarmed and

continuously oxygenated KHB buffer at 30°C. The optimal muscle length was determined, and all

subsequent measurements were performed at this length. For maximal force production mice, FDBs

were stimulated at 10, 30, 50, 80, 100, and 120 Hz for 1 second and with 0.1 ms pulses. Muscles were

left to rest for 5 minutes before starting the fatigue protocol as follows: 0.1 s train duration, 0.3 s train

delay, and pulses of 0.1 ms at 50 Hz. The maximal force was evaluated again 5 min after the end of the

fatigue protocol to check muscle integrity. Muscles were weighed and protein extraction was performed

to normalize. The maximal force was calculated with the difference of the peak force at 120 Hz and the

baseline and time to fatigue taken as the time necessary to reach 50% intensity of the first peak.

**MHC Staining** Electroporated *tibialis anterior* muscles were embedded in O.C.T, immediately frozen

in nitrogen-cooled isopentane, and stored at -80°C until cryosectioning. Muscle slices were blocked (5%

Goat serum, 2% BSA, 0,1% sodium azide in PBS) for 3 hours at room temperature and probed with

primary antibodies overnight at 4°C in a humidified chamber. The slides were washed 3 times with PBS

and incubated with Alexa Fluor conjugated secondary antibodies for 2 hours at room temperature.

Coverslips were mounted with ProLong antifade Diamond and whole sections were imaged with a

fluorescent scanning microscope at 20x magnification.

**Oxygen consumption assays** Oxygen consumption rates (OCR) were measured in a Seahorse XF24

extracellular flux analyzer according to the manufacturer's instructions. The following drugs were used

in the assay: 1 μM oligomycin (Oligo), 2 μM carbonyl cyanate m-chlorophenyl hydrazone (CCCP), and

1 μM rotenone/antimycin (Rot./Ant). ATP-linked OCR was calculated by subtracting OCR post

oligomycin addition from the OCR measured before. Reserve capacity was determined by subtracting

basal from maximal OCR. Non-mitochondrial values were subtracted before all calculations. For fatty-

acid oxidation assays, cell media was switched to low glucose 12 hours before the measurements, and

cells were equilibrated in KHB supplemented with 1g/L glucose, 4 mM L-glutamine, and 1 mM sodium

pyruvate for 1 hour. Immediately before the assay, BSA-conjugated palmitate was added to a final

concentration of 200 μM, and the drugs were added in the same manner. During routine oxygen

consumption assays, cells were maintained in phenol red-free DMEM, supplemented with 4.5g/L

glucose, 4 mM L-glutamine, and 1 mM sodium pyruvate, without sodium bicarbonate.

**Lactate measurement** Cells were grown in 96 well plates and then incubated for 3 h with 50 µL Krebs-

Henseleit Buffer (1.2 mM Na2HPO4, 2 mM MgSO4, 4.7 mM KCl, 111 mM NaCl, pH 7.3)

supplemented with 25 mM glucose, 1 mM pyruvate, and 4 mM Glutamine. Lactate production was

enzymatically quantified as NADH fluorescence (360 nm/460 nm) by the reverse reaction of L-lactate

dehydrogenase (Rabbit muscle, L25005KU, Sigma) in a reaction containing 20 µL cell media, 2 µg

enzyme, 50 mM Tris, and 625 mM Hydrazine in PBS. Following the assay, the cells were fixed and

stained with crystal violet for cell number normalization.

**Western blot** Protein extracts from cells and tissues were obtained with RIPA Buffer (Thermo

Scientific, # 89900) and 30 µg loaded into SDS-PAGE gels. Proteins were then transferred to 0.45 µm

PVDF membranes, probed with the indicated primary antibodies, and detected with ECL. Band

intensities were normalized by Ponceau S intensity and data is shown as fold-change over control.

**Microarray** RNA was extracted with TRIzol and subsequently column-purified using RNeasy Mini Kit

(Qiagen). Sample integrity was assessed, and the library was prepared using Affymetrix Whole

Transcript (WT) Assay kit probed in CGAS cartridge for Clariom S (mouse) following manufacturer's

instructions. Total RNA quality was assessed by Agilent Technologies 2200 Tapestation and

concentrations were measured by NanoDrop ND-1000 Spectrophotometer. Total RNA (150 ng) was

used to generate amplified sense strand cDNA targets using GeneChip® WT Plus Reagent Kit

(ThermoFisher Scientific) followed by fragmentation and labeling. 2.3 µg of ss cDNA target was

hybridized to Clariom™ S Mouse Arrays for 16 hours at 45°C under rotation in Affymetrix Gene Chip

Hybridization Oven 645 (ThermoFisher Scientific). Washing and staining were carried out on

Affymetrix GeneChip® Fluidics Station 450 (ThermoFisher Scientific), according to the manufacturer's

protocol. The fluorescent intensities were determined with Affymetrix GeneChip Scanner 3000 7G

(ThermoFisher Scientific). Transcriptome Analysis Console (TAC) software (v4.0.3, ThermoFisher

Scientific) was used for the analysis of microarray data. Signal values were log2-transformed, and

quantile normalized using the Signal Space Transformation (SST-RMA) method. Paired comparisons of

gene expression levels between sample groups were performed using moderated t-test as implemented

in BioConductor package limma. Gene ontology enrichment tests were performed with g:profiler

excluding electronic annotations.

**Cell-death assays** Cell-death assays were performed as described (Lima and Silveira 2018), with slight

modifications. Propidium iodide was added to a concentration of 5 µg/mL in cell culture media and

incubated for 20 minutes. Hoechst 33342 was then added to a final concentration of 1 µg/mL and

samples were incubated for another 10 minutes. Fluorescence was measured at 530/620 nm (ex./em.)

and 350/460 (ex./em.) nm in a plate reader.

**Cell doubling time** Cells ($10^4$) were plated in four replicates in 12-well plates. Thereafter, cells were

collected every 24 hours using trypsin and counted in a Neubauer chamber. The normalized data of

three independent experiments were used to obtain the doubling-time regression curve with the initial

number constraint.

**ATP measurement** Cells were grown in opaque 96-well white plates and then processed according to

the manufacturer's instructions of the CellTiter-Glo® Luminescent Cell Viability Assay kit (Promega).

The standard curve of ATP was determined in parallel for absolute quantitation.

**Bioinformatic analysis of public datasets** Nr2f6 ChIP-seq bigwig files from the ENCODE project

(GSM2797593 and GSM2534343) available on GEO were used. Using the Galaxy platform, the

anchoring position matrix was generated using the compute matrix command from the deeptools

package, relative to human genome annotations extracted from the UCSC Genome Browser in bed

format. The heatmap was obtained using the plotheatmap tool, also from the deeptools package.

Pathway enrichment was analyzed using the g.profiler program (https://biit.cs.ut.ee/gprofiler/gost), with

a significance threshold of 0.01 using the g:SCS parameter, without considering electronic term

annotations. The correlation in Supplementary Figure 1B was produced with fold-changes of

differentially expressed genes from RNA-seq (FDR <0.05) and fold-change values (expression in

myotube/expression in myoblast) from the C2C12 cell differentiation array (GSE4694) considering a p-

value cut-off of 0.01, according to GEO2R. UCP3 genome locus in Figure 3E was extracted from the

UCSC genome browser with the ChIP-seq tracks of Myogenin (wgEncodeEM002136,

wgEncodeEM002132), MyoD (wgEncodeEM002127, wgEncodeEM002129), H3K4me

(wgEncodeEM001450), H3K27Ac (wgEncodeEM001450) and DNA hypersensitivity track

(wgEncodeEM003399) over NCBI37/mm9 mouse genome assembly.

**Statistical analysis and quantification**

GraphPad Prism v7.0 was used for plotting the data and for statistical analysis. Cell culture experiments

were performed independently several times with 3-4 technical replicates. Ratio paired comparison

using Student's t-test was used for the analysis of human cells and electroporation experiments,

otherwise, an unpaired comparison was chosen, and in both cases, a 0.05 p-value cutoff was used.

Details for microarray and RNA-seq statistics are described in their respective methods section and

further statistical details are presented in figure labels.

**Acknowledgments**

We would like to appreciate the technical support of Ann-Marie Petterson.

**DECLARATIONS**

**Ethics approval and consent to participate**

All animal experiments were approved by the local regulatory entities, either the Stockholm North

Animal Ethical Committee or by the Ethics Committee on Animal Use (CEUA/Unicamp #5626-

1/2020), observing the welfare guidelines of the National Council of Control of Animal Experimentation

(CONCEA). Consent to participate not applicable.

**Consent for publication**

Not applicable.

**Availability of data and materials**

The materials originally produced in this study are available upon request to the lead contact. The

transcriptomic data generated in this study can be accessed at the GEO (GSE229102 and GSE228202).

**Competing interests**

The authors declare no competing interests

**REFERENCES**

[1] Z. Ma, Z. Zhong, Z. Zheng, X. M. Shi, and W. Zhang, "Inhibition of glycogen synthase kinase-3β attenuates glucocorticoid-induced suppression of myogenic differentiation in vitro," *PLoS One*, 2014, doi: 10.1371/journal.pone.0105528.

[2] D. Duan, N. Goemans, S. Takeda, E. Mercuri, and A. Aartsma-Rus, "Duchenne muscular dystrophy," *Nature Reviews Disease Primers*. 2021. doi: 10.1038/s41572-021-00248-3.

[3] P. M. Coen, R. V. Musci, J. M. Hinkley, and B. F. Miller, "Mitochondria as a target for mitigating sarcopenia," *Frontiers in Physiology*. 2019. doi: 10.3389/fphys.2018.01883.

[4] J. L. Brown *et al.*, "Mitochondrial degeneration precedes the development of muscle atrophy in progression of cancer cachexia in tumour-bearing mice," *J. Cachexia. Sarcopenia Muscle*, 2017, doi: 10.1002/jcsm.12232.

[5] K. De Bosscher, S. J. Desmet, D. Clarisse, E. Estébanez-Perpiña, and L. Brunsveld, "Nuclear receptor crosstalk — defining the mechanisms for therapeutic innovation," *Nature Reviews Endocrinology*. 2020. doi: 10.1038/s41574-020-0349-5.

[6] and V. A. N. Kumar, Ashok, "Nuclear receptors as potential therapeutic targets in peripheral arterial disease and related myopathy," *FEBS J.*, 2022, doi: 10.1111/febs.16593.

[7] M. Robinson-Rechavi, H. E. Garcia, and V. Laudet, "The nuclear receptor superfamily," *Journal of Cell Science*. 2003. doi: 10.1242/jcs.00247.

[8] K. W. Nettles and G. L. Greene, "Ligand control of coregulator recruitment to nuclear receptors," *Annual Review of Physiology*. 2005. doi: 10.1146/annurev.physiol.66.032802.154710.

[9] V. Klepsch, K. Siegmund, and G. Baier, "Emerging next-generation target for cancer immunotherapy research: The orphan nuclear receptor NR2F6," *Cancers*. 2021. doi: 10.3390/cancers13112600.

[10] B. Zhou *et al.*, "The Nuclear Orphan Receptor NR2F6 Promotes Hepatic Steatosis through Upregulation of Fatty Acid Transporter CD36," *Adv. Sci.*, 2020, doi: 10.1002/advs.202002273.

[11]    H. Li *et al.*, "Nuclear orphan receptor NR2F6 confers cisplatin resistance in epithelial ovarian

cancer cells by activating the Notch3 signaling pathway," *Int. J. Cancer*, 2019, doi: 10.1002/ijc.32293.

[12]    N. Hermann-Kleiter *et al.*, "Nuclear orphan receptor NR2F6 directly antagonizes NFAT and

RORγt binding to the Il17a promoter," *J. Autoimmun.*, 2012, doi: 10.1016/j.jaut.2012.07.007.

[13]    V. Klepsch *et al.*, "Nuclear receptor NR2F6 inhibition potentiates responses to PD-L1/PD-1

cancer immune checkpoint blockade," *Nat. Commun.*, 2018, doi: 10.1038/s41467-018-04004-2.

[14]    Y. Zhao *et al.*, "'Stripe' transcription factors provide accessibility to co-binding partners in

mammalian genomes.," *Mol. Cell*, vol. 82, no. 18, pp. 3398-3411.e11, Sep. 2022, doi:

10.1016/j.molcel.2022.06.029.

[15]    L. Al-Khalili, D. Krämer, P. Wretenberg, and A. Krook, "Human skeletal muscle cell

differentiation is associated with changes in myogenic markers and enhanced insulin-mediated MAPK

and PKB phosphorylation," *Acta Physiol. Scand.*, 2004, doi: 10.1111/j.1365-201X.2004.01259.x.

[16]    H. N. Araujo *et al.*, "Regulation of Lin28a-miRNA let-7b-5p pathway in skeletal muscle cells

by peroxisome proliferator-activated receptor delta," *Am. J. Physiol. - Cell Physiol.*, 2020, doi:

10.1152/ajpcell.00233.2020.

[17]    C. L. Andersen, J. L. Jensen, and T. F. Ørntoft, "Normalization of real-time quantitative

reverse transcription-PCR data: A model-based variance estimation approach to identify genes suited

for normalization, applied to bladder and colon cancer data sets," *Cancer Res.*, 2004, doi:

10.1158/0008-5472.CAN-04-0496.

[18]    K. J. Livak and T. D. Schmittgen, "Analysis of relative gene expression data using real-time

quantitative PCR and the 2-ΔΔCT method," *Methods*, 2001, doi: 10.1006/meth.2001.1262.

[19]    A. M. Bolger, M. Lohse, and B. Usadel, "Trimmomatic: A flexible trimmer for Illumina

sequence data," *Bioinformatics*, 2014, doi: 10.1093/bioinformatics/btu170.

[20]    R. Harmancey, D. L. Haight, K. A. Watts, and H. Taegtmeyer, "Chronic hyperinsulinemia

causes selective insulin resistance and down-regulates uncoupling protein 3 (ucp3) through the

activation of sterol regulatory element-binding protein (srebp)-1 transcription factor in the mouse heart," *J. Biol. Chem.*, 2015, doi: 10.1074/jbc.M115.673988.

[21]   C. Handschin, J. Rhee, J. Lin, P. T. Tarr, and B. M. Spiegelman, "An autoregulatory loop controls peroxisome proliferator-activated receptor γ coactivator 1α expression in muscle," *Proc. Natl. Acad. Sci. U. S. A.*, 2003, doi: 10.1073/pnas.1232352100.

[22]   T. I. Lima and L. R. Silveira, "A microplate assay for measuring cell death in C2C12 cells," *Biochem. Cell Biol.*, 2018, doi: 10.1139/bcb-2018-0005.

[23]   M. Warnecke, H. Oster, J. Revelli, G. Alvarez-bolado, and G. Eichele, "Abnormal development of the locus coeruleus in," *Genes Dev.*, 2005, doi: 10.1101/gad.317905.mice.

[24]   N. Hermann-Kleiter *et al.*, "The Nuclear Orphan Receptor NR2F6 Is a Central Checkpoint for Cancer Immune Surveillance," *Cell Rep.*, 2015, doi: 10.1016/j.celrep.2015.08.035.

[25]   P. Kaliman, F. Viñals, X. Testar, M. Palacín, and A. Zorzano, "Phosphatidylinositol 3-kinase inhibitors block differentiation of skeletal muscle cells," *J. Biol. Chem.*, 1996, doi: 10.1074/jbc.271.32.19146.

[26]   P. Rochard *et al.*, "Mitochondrial activity is involved in the regulation of myoblast differentiation through myogenin expression and activity of myogenic factors," *J. Biol. Chem.*, 2000, doi: 10.1074/jbc.275.4.2733.

[27]   O. Halevy *et al.*, "Correlation of terminal cell cycle arrest of skeletal muscle with induction of p21 by MyoD," *Science (80-. ).*, 1995, doi: 10.1126/science.7863327.

[28]   I. L. De la Serna, K. Roy, K. A. Carlson, and A. N. Imbalzano, "MyoD Can Induce Cell Cycle Arrest but Not Muscle Differentiation in the Presence of Dominant Negative SWI/SNF Chromatin Remodeling Enzymes," *J. Biol. Chem.*, 2001, doi: 10.1074/jbc.M107281200.

[29]   Y. Kobayashi *et al.*, "Cyclin-Dependent Kinase 1 Is Essential for Muscle Regeneration and Overload Muscle Fiber Hypertrophy," *Front. Cell Dev. Biol.*, 2020, doi: 10.3389/fcell.2020.564581.

[30]   T. Sato, T. Yamamoto, and A. Sehara-Fujisawa, "MiR-195/497 induce postnatal quiescence of

skeletal muscle stem cells," *Nat. Commun.*, 2014, doi: 10.1038/ncomms5597.

[31]  J. M. Zhang, Q. Wei, X. Zhao, and B. M. Paterson, "Coupling of the cell cycle and myogenesis

through the cyclin D1-dependent interaction of MyoD with cdk4," *EMBO J.*, 1999, doi:

10.1093/emboj/18.4.926.

[32]  A. M. Abdelmoez *et al.*, "Comparative profiling of skeletal muscle models reveals

heterogeneity of transcriptome and metabolism," *Am. J. Physiol. - Cell Physiol.*, 2020, doi:

10.1152/ajpcell.00540.2019.

[33]  I. Casimiro, N. D. Stull, S. A. Tersey, and R. G. Mirmira, "Phenotypic sexual dimorphism in

response to dietary fat manipulation in C57BL/6J mice," *J. Diabetes Complications*, 2021, doi:

10.1016/j.jdiacomp.2020.107795.

[34]  T. I. Lima *et al.*, "Role of NCoR1 in mitochondrial function and energy metabolism," *Cell*

*Biology International*. 2018. doi: 10.1002/cbin.10973.

[35]  T. I. Lima *et al.*, "Opposing action of NCoR1 and PGC-1α in mitochondrial redox

homeostasis," *Free Radic. Biol. Med.*, 2019, doi: 10.1016/j.freeradbiomed.2019.08.006.

[36]  V. A. Narkar *et al.*, "AMPK and PPARδ Agonists Are Exercise Mimetics," *Cell*, 2008, doi:

10.1016/j.cell.2008.06.051.

[37]  P. M. Badin *et al.*, "Exercise-like effects by Estrogen-related receptor-gamma in muscle do not

prevent insulin resistance in db/db mice," *Sci. Rep.*, 2016, doi: 10.1038/srep26442.

[38]  D. Taillandier and C. Polge, "Skeletal muscle atrogenes: From rodent models to human

pathologies," *Biochimie*. 2019. doi: 10.1016/j.biochi.2019.07.014.

[39]  D. G. Allen, G. D. Lamb, and H. Westerblad, "Skeletal muscle fatigue: Cellular mechanisms,"

*Physiological Reviews*. 2008. doi: 10.1152/physrev.00015.2007.

[40]  D. R. Dietrich, "Toxicological and pathological applications of proliferating cell nuclear

antigen (PCNA), a novel endogenous marker for cell proliferation," *Crit. Rev. Toxicol.*, 1993, doi:

10.3109/10408449309104075.

[41] N. A. M. Pansters *et al.*, "Muscle-specific GSK-3β ablation accelerates regeneration of disuse-

atrophied skeletal muscle," *Biochim. Biophys. Acta - Mol. Basis Dis.*, 2015, doi:

10.1016/j.bbadis.2014.12.006.

[42] I. Michailovici *et al.*, "Nuclear to cytoplasmic shuttling of ERK promotes differentiation of

muscle stem/progenitor cells," *Dev.*, 2014, doi: 10.1242/dev.107078.

[43] N. C. Jones, Y. V. Fedorov, R. S. Rosenthal, and B. B. Olwin, "ERK1/2 is required for

myoblast proliferation but is dispensable for muscle gene expression and cell fusion," *J. Cell. Physiol.*,

2001, doi: 10.1002/1097-4652(200101)186:1<104::AID-JCP1015>3.0.CO;2-0.

[44] N. Hermann-Kleiter *et al.*, "The Nuclear Orphan Receptor NR2F6 Suppresses Lymphocyte

Activation and T Helper 17-Dependent Autoimmunity," *Immunity*, 2008, doi:

10.1016/j.immuni.2008.06.008.

[45] W. J. Olson *et al.*, "Orphan Nuclear Receptor NR2F6 Suppresses T Follicular Helper Cell

Accumulation through Regulation of IL-21," *Cell Rep.*, 2019, doi: 10.1016/j.celrep.2019.08.024.

[46] S. L. Yang *et al.*, "The expression and biological effect of NR2F6 in non-small cell lung

cancer.," *Front. Oncol.*, vol. 12, p. 940234, 2022, doi: 10.3389/fonc.2022.940234.

[47] B. Kupr, S. Schnyder, and C. Handschin, "Role of nuclear receptors in exercise-induced

muscle adaptations," *Cold Spring Harbor Perspectives in Medicine*. 2017. doi:

10.1101/cshperspect.a029835.

[48] S. A. J. Verbrugge, M. Schönfelder, L. Becker, F. Y. Nezhad, M. H. de Angelis, and H.

Wackerhage, "Genes whose gain or loss-of-function increases skeletal muscle mass in mice: A

systematic literature review," *Frontiers in Physiology*. 2018. doi: 10.3389/fphys.2018.00553.

[49] A. Mayeuf-Louchart *et al.*, "Rev-erb-α regulates atrophy-related genes to control skeletal

muscle mass," *Sci. Rep.*, 2017, doi: 10.1038/s41598-017-14596-2.

[50] H. Yamamoto *et al.*, "NCoR1 is a conserved physiological modulator of muscle mass and

oxidative function," *Cell*, vol. 147, no. 4, pp. 827–839, 2011, doi: 10.1016/j.cell.2011.10.017.

694 [51] N. Shimizu *et al.*, "A muscle-liver-fat signalling axis is essential for central control of adaptive

695 adipose remodelling," *Nat. Commun.*, 2015, doi: 10.1038/ncomms7693.

696 [52] D. Avram *et al.*, "Heterodimeric interactions between chicken ovalbumin upstream promoter-

697 transcription factor family members ARP1 and Ear2," *J. Biol. Chem.*, 1999, doi:

698 10.1074/jbc.274.20.14331.

699 [53] H.-J. Lee *et al.*, "Dysregulation of nuclear receptor COUP-TFII impairs skeletal muscle

700 development OPEN", doi: 10.1038/s41598-017-03475-5.

701 [54] X. Xie, S. Y. Tsai, and M. J. Tsai, "COUP-TFII regulates satellite cell function and muscular

702 dystrophy," *J. Clin. Invest.*, 2016, doi: 10.1172/JCI87414.

703 [55] M. A. Pearen *et al.*, "The nuclear receptor, Nor-1, markedly increases type II oxidative muscle

704 fibers and resistance to fatigue," *Mol. Endocrinol.*, 2012, doi: 10.1210/me.2011-1274.

705 [56] W. Chen, X. Zhang, K. Birsoy, and R. G. Roeder, "A muscle-specific knockout implicates

706 nuclear receptor coactivator MED1 in the regulation of glucose and energy metabolism," *Proc. Natl.*

707 *Acad. Sci. U. S. A.*, 2010, doi: 10.1073/pnas.1005626107.

708 [57] A. Seth *et al.*, "The Transcriptional Corepressor RIP140 Regulates Oxidative Metabolism in

709 Skeletal Muscle," *Cell Metab.*, 2007, doi: 10.1016/j.cmet.2007.08.004.

710 [58] C. Son *et al.*, "Reduction of diet-induced obesity in transgenic mice overexpressing uncoupling

711 protein 3 in skeletal muscle," *Diabetologia*, 2004, doi: 10.1007/s00125-003-1272-8.

712 [59] J. Darcy MacLellan, M. F. Gerrits, A. Gowing, P. J. S. Smith, M. B. Wheeler, and M. E.

713 Harper, "Physiological increases in uncoupling protein 3 augment fatty acid oxidation and decrease

714 reactive oxygen species production without uncoupling respiration in muscle cells," *Diabetes*, 2005,

715 doi: 10.2337/diabetes.54.8.2343.

716 [60] J. C. Clapham *et al.*, "Mice overexpressing human uncoupling protein-3 in skeletal muscle are

717 hyperphagic and lean," *Nature*, 2000, doi: 10.1038/35019082.

718 [61] C. Son *et al.*, "Up-regulation of uncoupling protein 3 gene expression by fatty acids and

agonists for PPARs in L6 myotubes," *Endocrinology*, 2001, doi: 10.1210/endo.142.10.8446.

[62]   D. S. Weigle *et al.*, "Elevated free fatty acids induce uncoupling protein 3 expression in muscle: A potential explanation for the effect of fasting," *Diabetes*, 1998, doi: 10.2337/diab.47.2.298.

[63]   A. J. Cruz-Jentoft and A. A. Sayer, "Sarcopenia," *The Lancet*. 2019. doi: p.

[64]   A. Picca *et al.*, "Update on mitochondria and muscle aging: All wrong roads lead to sarcopenia," *Biological Chemistry*. 2018. doi: 10.1515/hsz-2017-0331.

[65]   M. Bakay *et al.*, "Nuclear envelope dystrophies show a transcriptional fingerprint suggesting disruption of Rb-MyoD pathways in muscle regeneration," *Brain*, 2006, doi: 10.1093/brain/awl023.

[66]   S. Salinas, C. Proukakis, A. Crosby, and T. T. Warner, "Hereditary spastic paraplegia: clinical features and pathogenetic mechanisms," *The Lancet Neurology*. 2008. doi: 10.1016/S1474-4422(08)70258-8.

[67]   J. K. Fink, "Hereditary spastic paraplegia: Clinico-pathologic features and emerging molecular mechanisms," *Acta Neuropathologica*. 2013. doi: 10.1007/s00401-013-1115-8.

[68]   V. Klepsch *et al.*, "Nuclear orphan receptor NR2F6 as a safeguard against experimental murine colitis," *Gut*, 2018, doi: 10.1136/gutjnl-2016-313466.

[69]   L. Pelletier *et al.*, "In vivo RyR1 reduction in muscle triggers a core-like myopathy," *Acta Neuropathol. Commun.*, 2020, doi: 10.1186/s40478-020-01068-4.

[70]   I. Sakakibara *et al.*, "Myofiber androgen receptor increases muscle strength mediated by a skeletal muscle splicing variant of Mylk4," *iScience*, 2021, doi: 10.1016/j.isci.2021.102303.

[71]   N. K. L. Lee and H. E. Maclean, "Polyamines, androgens, and skeletal muscle hypertrophy," *Journal of Cellular Physiology*. 2011. doi: 10.1002/jcp.22569.

[72]   M. Cervelli, A. Leonetti, G. Duranti, S. Sabatini, R. Ceci, and P. Mariottini, "Skeletal Muscle Pathophysiology: The Emerging Role of Spermine Oxidase and Spermidine," *Med. Sci.*, 2018, doi: 10.3390/medsci6010014.

[73]   L. Wang, H. Long, Q. Zheng, X. Bo, X. Xiao, and B. Li, "Circular RNA circRHOT1 promotes

hepatocellular carcinoma progression by initiation of NR2F6 expression," *Mol. Cancer*, 2019, doi:

10.1186/s12943-019-1046-7.

[74]   J. Liu, T. Li, and X. L. Liu, "DDA1 is induced by NR2F6 in ovarian cancer and predicts poor

survival outcome," *Eur. Rev. Med. Pharmacol. Sci.*, 2017.

[75]   E. T. Weatherford, X. Liu, and C. D. Sigmund, "Regulation of renin expression by the orphan

nuclear   receptors   Nr2f2   and   Nr2f6," *Am.   J.   Physiol.   -   Ren.   Physiol.*, 2012, doi:

10.1152/ajprenal.00362.2011.

[76]   K. Chu and H. H. Zingg, "The nuclear orphan receptors COUP-TFII and Ear-2 act as silencers

of the human oxytocin gene promoter," *J. Mol. Endocrinol.*, 1997, doi: 10.1677/jme.0.0190163.

[77]   C. Jin, L. Xiao, Z. Zhou, Y. Zhu, G. Tian, and S. Ren, "MiR-142-3p suppresses the

proliferation, migration and invasion through inhibition of NR2F6 in lung adenocarcinoma," *Hum.

Cell*, 2019, doi: 10.1007/s13577-019-00258-0.

[78]   S. Raichur, P. Lau, B. Staels, and G. E. O. Muscat, "Retinoid-related orphan receptor γ

regulates several genes that control metabolism in skeletal muscle cells: Links to modulation of

reactive oxygen species production," *J. Mol. Endocrinol.*, 2007, doi: 10.1677/jme.1.00010.

[79]   J. Iqbal *et al.*, "Deletion of retinoic acid-related orphan receptor gamma reduces body weight

and hepatic lipids in mice by modulating the expression of lipid metabolism genes," *Vessel Plus*, 2019,

doi: 10.20517/2574-1209.2019.28.

[80]   Y. Fang, H. Yu, X. Liang, J. Xu, and X. Cai, "Chk1-induced CCNB1 overexpression promotes

cell proliferation and tumor growth in human colorectal cancer," *Cancer Biol. Ther.*, 2014, doi:

10.4161/cbt.29691.

[81]   V. Huang *et al.*, "Upregulation of Cyclin B1 by miRNA and its implications in cancer,"

*Nucleic Acids Res.*, 2012, doi: 10.1093/nar/gkr934.

[82]   B. Bao, X. Yu, and W. Zheng, "MiR-139-5p Targeting CCNB1 Modulates Proliferation,

Migration, Invasion and Cell Cycle in Lung Adenocarcinoma," *Mol. Biotechnol.*, 2022, doi:

10.1007/s12033-022-00465-5.

[83]    V. K. Rao *et al.*, "G9a promotes proliferation and inhibits cell cycle exit during myogenic

differentiation," *Nucleic Acids Res.*, 2016, doi: 10.1093/nar/gkw483.

[84]    J. D. R. Knight and R. Kothary, "The myogenic kinome: Protein kinases critical to mammalian

skeletal myogenesis," *Skeletal Muscle*. 2011. doi: 10.1186/2044-5040-1-29.

[85]    H. Tajsharghi and A. Oldfors, "Myosinopathies: Pathology and mechanisms," *Acta

Neuropathologica*. 2013. doi: 10.1007/s00401-012-1024-2.

[86]    S. Schiaffino, "Muscle fiber type diversity revealed by anti-myosin heavy chain antibodies,"

*FEBS Journal*. 2018. doi: 10.1111/febs.14502.

 **FIGURES LEGENDS**

**Figure 1. Nr2f6 knockdown derepresses the expression of genes involved in metabolism and**

**myogenesis. (A)** Volcano plot of Nr2f6 knockdown C2C12 myocytes. Genes upregulated in red and

downregulated in blue (FDR <0.05, $\log_2$FC >0.5). N=4-5.**(B)** Network of ontology terms enriched in the

differentially expressed genes. Groups of similar terms were manually curated and encircled as

indicated. **(C, D)** Gene ontology enrichment of downregulated and upregulated genes. **(E)** Panel of

myogenic differentiation markers differentially regulated by Nr2f6 knockdown with Myogenic

Regulatory Factors (MRFs) and myosin isoforms with their respective fiber expression pattern[85], [86].

**(F)** Insulin signaling pathway schematic displaying differentially expressed genes after Nr2f6

knockdown and other components of the pathway. Metabolites are depicted in yellow borders and

unchanged genes are in orange borders. **(G)** Gene expression measured by RT-qPCR of markers of

myogenic differentiation in primary human skeletal muscle cells transfected with control non-target

RNAi (siScr) or siNr2f6. N = 5-6. Boxplot with whiskers spanning minimum to maximal and box edges

$25^{th}$-$75^{th}$ percentile, the line at the median and + at the mean. * Indicates $p < 0.05$ using unpaired two-

tailed Student's t-test.

**Figure 2. Nr2f6 depletion increases fatty acid oxidation and protects cells against lipid-induced**

**stress. (A)** Fatty-acid-dependent oxygen consumption assay in control siScr and siNr2f6 C2C12

myocytes. Data displayed as mean ±SD. On the right, calculated respiratory parameters are displayed as

a line on the mean and minimum to max bars. N = 3. * Indicates $p < 0.05$ using unpaired two-tailed

Student's t-test. **(B)** Oligomycin-induced extracellular acidification rate during a high-glucose oxygen

consumption assay. N=4. **(C)** Lactate measurement in cell culture media of C2C12 myocytes transfected

with control siScr and siNr2f6. N=3. **(D, E)** Relative gene expression using RT-qPCR in stable Nr2f6

knockdown (shNr2f6) C2C12 cells and control shGFP stable cells. **(F)** Cell death as measured by

propidium iodide in control (shGFP) and shNr2f6 myocytes following treatment with 500 μM palmitate

for 20 hours. N=3. **(G, H)** Mitochondrial and total superoxide production following palmitate treatment

in shGFP and shNr2f6 stable C2C12 cells. N=3-4. (**I**) Relative Nr2f6 mRNA expression in C2C12

myotubes treated with 500 µM palmitate or vehicle for 20 hours. N=3. (**J**) Relative Nr2f6 mRNA

expression in the gastrocnemius of mice undergoing a control chow or high-fat diet for 16 weeks. N=7.

(**K**) Densitometry and representative western blot image of Nr2f6 in gastrocnemius lysates from mice

fed HFD or control chow for 16 weeks. N=6. Boxplot with whiskers spanning minimum to maximal and

box edges $25^{th}$-$75^{th}$ percentile, the line at the median and + at the mean. * Indicates $p < 0.05$ using

unpaired two-tailed Student's t-test.

**Figure 3. Nr2f6 directly regulates PGC1-a and UCP3 gene expression.** (**A**) Relative gene expression

using RT-qPCR in stable Nr2f6-myc overexpression myotubes. N=4-5. (**B**) Densitometry and

representative images of PGC-1α western blot in stable Nr2f6-myc overexpression myotubes. N=5. (**C**)

Relative gene expression using RT-qPCR in stable Nr2f6 knockdown myotubes. N=3-5. (**D**) Luciferase

reporter assay in HEK293 cells with 2 kbp PGC1A promoter overexpressing HA-tagged Nr2f6 or

control empty vector (EV). (**E**) Luciferase activity of UCP3 promoter transactivation assay in cells

overexpressing Nr2f6-myc and siNr2f6 transfected cells. N=3. (**F**) Mouse UCP3 genomic locus

retrieved from UCSC Genome Browser with the Nr2f6 response element highlighted. Top tracks: ChIP-

seq of Myogenin and MyoD at 24h and 60h of differentiation. Middle track: DNAse hypersensitivity

assay, with open sensitive regions in grey. Bottom tracks: histone marks ChIP-seq. (**G**) Relative gene

expression using RT-qPCR in human primary skeletal myotubes transfected with siNr2f6 or siScr. N=6.

Boxplot with whiskers spanning minimum to maximal and box edges $25^{th}$-$75^{th}$ percentile, the line at the

median and + at the mean. * Indicates $p < 0.05$ using unpaired two-tailed Student's t-test. The numbers

above some bars indicate the p-value.

**Figure 4. The molecular signature of Nr2f6 overexpression in skeletal muscle reveals an increase**

**in inflammation and a decrease in muscle contraction and metabolism.** (**A**) Heat-map of top 30

most modulated genes in *tibialis anterior* muscle electroporated with empty vector (control) or an Nr2f6

coding plasmid. N=4. (**B, C**) Gene ontology enrichment of downregulated and upregulated genes. (**D, E,**

**F)** Validation of selected markers modulated in the microarray by RT-qPCR. N=4. Insert on D:

representative western blot for validation of Nr2f6 protein content in *tibialis anterior* samples under

control and electroporated conditions. Boxplot with whiskers spanning minimum to maximal and box

edges 25[th]-75[th] percentile, the line at the median and + at the mean. * Indicates $p < 0.05$ using unpaired

two-tailed Student's t-test. The numbers above some bars indicate the p-value.

**Figure 5. Overexpression of Nr2f6 induces muscle atrophy and impairs muscle force production.**

**(A)** Weight of *tibialis anterior* muscles (TA) electroporated with empty vector or Nr2f6 coding plasmid.

Top: representative photo. N=12. **(B)** Representative images of myosin heavy chain staining in the

electroporated TAs for fiber type determination. In green, MHC IIA; in red, MHC IIB; unstained fibers

as IIX. No significant number of MHCI fibers were stained, therefore the corresponding channel was

omitted. N=7. **(C, D)** Total and type-segmented number of fibers. N=7. **(E)** *Ex-vivo* contraction maximal

force production in FDB muscles electroporated with control empty vector (EV) or Nr2f6 coding

plasmid. N=5. Data are displayed as individual animals and bars at the mean. * Indicates $p < 0.05$ using

ratio paired two-tailed Student's t-test.

**Figure 6. Nr2f6 promotes myoblast proliferation. (A)** Scatter plot of differentially expressed genes in

Nr2f6 knockdown in C2C12 myocytes and Nr2f6 overexpression in mice TA. In red: genes upregulated

by Nr2f6; in blue: genes downregulated by Nr2f6; in grey: genes with the same direction of modulation

by Nr2f6 overexpression and knockdown. **(B)** Interaction network of genes consistently regulated by

Nr2f6 overexpression and knockdown and with detected Nr2f6 binding motif at the promoter region. In

blue: genes downregulated; in red: genes upregulated. The number of connections of each gene

increases clockwise. **(C, D)** Representative images of western blot of the electroporated *tibialis anterior*

and densitometric quantitation of protein bands. N=4. **(E, G)** Proliferation curves of stable Nr2f6

knockdown and overexpression cell lines and the calculated doubling time. **(F, H)** RT-qPCR of cell

cycle arrest markers in Nr2f6 knockdown and overexpression stable cell lines, respectively. N=4-6.

Boxplot with whiskers spanning minimum to maximal and box edges 25[th]-75[th] percentile, the line at the

median and + at the mean. * Indicates p < 0.05 using unpaired two-tailed Student's t-test. The numbers

above some bars indicate the p-value.

**Figure 7. Nr2f6 represses core genes of muscle contraction.** Nr2f6 overexpression reduces the

expression of several genes of the contractile apparatus, myofiber calcium handling, and action potential

transduction. Genes with Nr2f6 binding motif at the promoter are underscored. Differentially expressed

genes following Nr2f6 overexpression in mouse TA were selected according to ontology terms related

to muscle contraction and function. The arrows indicate the up- or downregulation. Sodium Voltage-

Gated Channel Alpha Subunit 4 (Scn4a), Potassium Inwardly Rectifying Channel Subfamily J Member

2 (Kcnj2), Solute Carrier Family 8 Member A3 (Slc8a3), Muscle Associated Receptor Tyrosine Kinase

(Musk), Ryanodine Receptor 1 (Ryr1), Calsequestrin 1 (Casq1), ATPase Sarcoplasmic/Endoplasmic

Reticulum Ca2+ Transporting 2 (SERCA2, Atp2a2), Cholinergic Receptor Nicotinic Alpha

1/delta/gamma subunit Chrna1/d/g), Troponin T1/I1/I2/C1 (Tnnt1/Tnni1/Tnni2/c1), Myom1/2

(Myomesin1/2), Myozenin1/3 (Myoz1/3), Myosin light chain kinase 2/4 (Mylk2/4), Myosin heavy

chain 3 (Myh3), Myosin binding protein C1/2 (Mybpc1/2)

**A** C2C12 Stable Nr2f6 Overexpression PGC1-α targets

**B** PGC1-α Protein content

**C** PGC1-α promoter

**D** C2C12 Stable Nr2f6 Knockdown PGC1-α targets

**E** UCP3 promoter

**F**

**G** HSMC

Labels:
- Nr2f6 knockdown
- Nr2f6 overexpression

1931 TAGGGCAAAGGGGA 1944

**A**

EV    NR2F6

**B**

EV    NR2F6

500 μm    500 μm

■ **Type IIA** ■ **Type IIB** ■ **Type IIX**

**C**

Total fiber number

*

**D**

Fyber type

□ EV ■ Nr2f6

IIA    IIB    IIX

*

0.1

**E**

Specific Max.
Force Production

*

EV Nr2f6

52 ◄    Nr2f6