On-farm hatching and contact with adult hen post hatch induce sex-

dependent effects on performance and welfare in broiler chickens

L. A. Guilloteau[a*], A. Bertin[b], S. Crochet[a], C. Bagnard[c], A. Hondelatte[c], L. Ravon[c], C.

Schouler[d], K. Germain[c], A. Collin[a]

[a] *INRAE, Université de Tours, BOA, 37380 Nouzilly, France*

[b] *CNRS, IFCE, INRAE, Université de Tours, PRC, 37380 Nouzilly, France*

[c] *INRAE, EASM, 17700 Surgères, France*

[d] *INRAE, Université de Tours, ISP, 37380 Nouzilly, France*

*Corresponding author: Laurence Guilloteau. Email: Laurence.guilloteau@inrae.fr

Abstract

To improve the early perinatal conditions of broiler chicks, alternative hatching systems

have been developed. On-farm hatching (OFH) with an enriched microbial and

stimulating environment by the presence of an adult hen is a promising solution. Day-

old certified JA 757 chicks were allotted within different hatching and rearing

conditions: OFH, conventional hatchery (CH), CH and post-hatching treatment with

antibiotics (CH + AB), as well as both hatching systems with an adult hen at hatching

(OFH + H, CH + H). On day (D) 27, chickens were challenged by combining transport

in boxes in a new room at a lower temperature and fasting for 4 h. On their return to

the original room, the chicken density was increased, and birds were orally vaccinated

with the Gumboro vaccine. The impacts of these conditions on hatchability, chick

quality score, performance, health and robustness were determined. The OFH chick

body weights (BWs) were significantly greater than those of CH chicks at hatching.

Whereas there was no effect of hatching conditions, the presence of hens, categorised

according to their behaviour, decreased the hatchability rate, the quality score of OFH

chicks and increased mortality at hatching. Treatment of CH chicks with antibiotics

temporarily decreased chicken BW at D19, but the feed conversion ratio (FCR) was

not modified. At D19, OFH chicks had the best BW compared to the other groups, and

the presence of hens at hatching harmed chicken BW regardless of the hatching

condition and FCR. An interaction between the effect of hatching conditions and

chicken sex was observed later in BW. In males, the OFH chickens were the heaviest

compared to the other groups at D34 but not at D56. The presence of hens eventually

negatively impacted CH chicken BW at D56. In females, there was no effect of hatching

condition on the BWs at D34 and D56, and the presence of hens eventually had a

positive impact on OFH chicken BW. There was no effect of hatching conditions on

health parameters. In conclusion, the OFH system was a hatching system at least

equivalent to the CH system, if not better in this study. The effects of the hen's

presence at hatching and during the chick start-up phase on performance interacted

with the hatching condition and the sex of the chickens. The health status and brooding

behaviour of the hens are essential to ensure the health and welfare of the chicks.

Introduction

In poultry, the perinatal period is a stressful period for broiler chicks, which includes the hatching phase and major physiological changes to adapt to new food resources and environments. In hatcheries, chicks hatch at between 19 and 21 days of incubation. They often stay more than 12 hours in the hatcher, under optimal temperature, without light and usually without access to feed and water until placement in farm buildings. The fasting period of the chicks is further increased by the time needed for hatchery processing, transportation duration and unloading at the farm, which might last up to the first 72 h after hatching. Even though chicks can use energy reserves from their yolk sac (van der Wagt et al., 2020), these conditions induce immediate and long-lasting metabolic changes (Beauclercq et al., 2019; Foury et al., 2020), behavioural impacts by increasing fear responses (Jessen et al., 2021) and consequences on chicken development, performance and welfare (de Jong et al., 2017).

To improve the early perinatal conditions of chicks, alternative hatching systems have been developed. On-farm hatching provides the chicks with immediate access to feed and water according to their needs and avoids the exposure to stressors encountered in conventional hatcheries (van de Ven et al., 2009). Eggs incubated for 18 days are transported to the farm and placed either in trays or in the litter where they hatch. The effects of these on-farm hatching systems on broiler health, welfare and performance were recently studied under commercial or more controlled conditions and had shown effects that are not always beneficial. Total mortality and footpad dermatitis in on-farm hatched (OFH) chicks were lower compared to conventionally hatched (CH) fast-

growing broiler chickens (de Jong et al., 2019; 2020; Giersberg et al., 2021; Jessen et

al., 2021). However, day-old chick quality was worse and breast myopathy prevalence

was higher for OFH than CH chickens (de Jong et al., 2019; Souza da Silva et al.,

2021).

Chicken activity and general behaviour were little affected by the hatching system, with

fast-growing OFH chickens being more fearful and less active (Giersberg et al., 2020).

Slower-growing broiler chickens hatched in organic farms tended to express less

general fearfulness than CH chickens (Jessen et al., 2021). A positive effect on growth

performance was observed during the first week of life until 21 days in OFH and CH

fed at the hatchery compared to CH chickens (de Jong et al., 2020), and longer when

parent flocks were young (Souza da Silva et al., 2021).

Maintaining optimal health, welfare and performance of chickens is highly dependent

on the gut physiology in interaction with the microbiota and mucosal immune system

(Fortun-Lamothe et al., 2023). Antibiotics have been largely used in poultry production

to improve performance by acting on the gut barrier function (Broom, 2018). However,

growing concerns about the increase of antimicrobial resistance in farm animals led to

changes in EU and national legislation governing the use of antibiotics as growth

promoters in poultry feed, which resulted in their suppression in 2006 (Council

Directive 96/22/EC; Axis 2 and measure 19 of the EcoAntibio 2017 plan). This made it

possible to become aware of the crucial role of the gut barrier and of the quality of the

microbiota implanted in the chick's gut at hatching on the physiological and immune

development, its robustness in the face of the hazards encountered during the chicks'

lives, and consequently on performance.

Greater attention to the environment during the chick postnatal period, especially the

microbial environment, is key to optimising the gut barrier function and more broadly

the health and welfare of the chickens and their performance. Naturally, chicks hatch

in contact with an adult hen who is a donor of microbiota and a model of learning and

maternal care (Edgar et al., 2016). Early implantation of adult microbiota into the chick

digestive system accelerates the maturation of the microbiota and immune system

(Volf et al., 2016; Broom & Kogut, 2018; Meijerink et al., 2020). In addition, chicks

reared in the presence of their mothers are less fearful than those raised without their

mothers and develop more behavioural synchrony (Perré et al., 2002), even though

hen genetics has a strong effect on chick behaviour, with commercial lines being less

maternal (Hewlett et al., 2019). The combination of a new hatching system like OFH

with an enriched microbiota and stimulating environment from the presence of an adult

hen is a possible solution for chick conditions to be improved and could contribute to

poultry health and welfare and product quality.

In this study, we analysed the benefits/risks of hatching systems (OFH and CH), with

an adult hen (OFH + H, CH + H) or not, and the combination of CH and post-hatching

treatment with antibiotic growth promoter (CH + AB) on hatchability on chick quality

score, performance, health and robustness.

Material and methods

All experimental procedures were approved by the Ethics Committee COMETHEA

POITOU-CHARENTES n°84 (APAFIS#24474-2020021816237418 v3) and carried out

following current European legislation (EU Directive 2010/63/EU). All steps of hatching,

experimentation and rearing were done at the experimental unit (EASM, Poultry

alternative breeding facility, INRAE, 17700 Surgères, France, DOI:

10.15454/1.5572418326133655E12).

Experimental design

Seven hundred twenty-day-old, certified JA 757 chicks, among which 432 were from a

conventional hatchery (CH) and 288 were hatched on-farm (OFH), were allocated into

five groups: CH, CH + antibiotics treatment (CH + AB), CH + hen (CH + H), OFH, OFH

+ hen (OFH + H). Each group was randomly placed in the room, repeated in eight pens

(18 chicks/pen, 3 m^2). Antibiotic treatment was only applied in chick drinking water for

the CH + AB group: ADJUSOL$^®$ TMP SULF Liquid (25 mg/kg sulfadiazine and 5 mg/kg

trimethoprim, VIRBAC, CARROS, France) for 5 days (D2–D6) and SURAMOX 50 (400

mg/10 kg, i.e. 20 mg/kg amoxicillin, VIRBAC) for 5 days (D19–D23). Sex was

determined on tagged chickens on D19, and the number of chickens was adjusted to

a maximum of 16 per pen, keeping a balanced ratio between males and females. On

D27, chickens were challenged by combining transport in boxes to a new room at a

lower temperature (15 °C instead of 25 °C) and 4 h of feed deprivation. On their return

to the original room, the pen size was reduced from 3 m^2 to 1.5 m^2 to increase chicken

density, and birds were orally vaccinated with the live Gumboro vaccine in drinking

water (HIPRAGUMBORO$^®$ - G97, HIPRA FRANCE, Saint-Herblain, France). Chickens

had ad libitum access to feed without anticoccidial drugs. They were fed with a

standard starter diet (raw energy = 4462 kcal/kg, crude protein = 23.91%) until D19,

then a grower diet from D20 to D34 (4527 kcal/kg, crude protein = 20.51%) and a

finisher diet from D35 to D56 (4600 kcal/kg, crude protein = 19.98%). Faeces were

collected from litter on D14 and D54 for parasite analyses.

Hatching and husbandry

Hatching conditions

Certified JA 757 18-day embryonated eggs (Galina Vendée, Essarts-en-Bocage,

France) were either placed at 37.6°C with 75% relative humidity and no light in a

conventional hatchery (CH) or laid directly in the litter of the pens under infrared heat

lamps to allow on-farm hatching (OFH). The average temperature of the eggs in the

litter was 37.9°C and under 20 h light per day until OFH chick hatching. The ambient

room temperature was maintained at 25 °C with a fan heater. Day-old CH chicks were

transported for one hour in a transport van before placement in pens to simulate

conventional hatchery processing, which has been described to have long-term

deleterious effects on fear response when combined with delayed nutrition (Hollemans

et al., 2018). Both CH and OFH chicks were placed under heat lamps. In pens hosting

hens, 18-day embryonated eggs or chicks were placed in a gridded space under the

heat lamp. Temperatures under heat lamps were decreased from 35–38 °C to 31–32

°C from D0 to D3, then 29–30 °C from D4 to D6 and 26–27 °C from D7 to D13. The

light cycle was 20 h light at the CH chick placement or until hatching time for OFH chick

(D0), 13 h light on D1 (increased dark time to promote maternal behaviour of hens),

18 h on D2 and 16 h on D3 and during the rearing period with minimum 20 lux on 80%

of the lighted surface.

Chicks had ad libitum access to water and food; a wire mesh platform and a perch

were used for environmental enrichment.

Contact with hens

Sixteen Lohmann Brown hens, acting as natural sources of gut microbiota and adult

presence, were obtained from a local commercial egg-laying hen farm (La cabane à

Chiron, Benet, France). The hens were aged 31 weeks, vaccinated against Marek

Disease Virus (MDV), Infectious Bursite Disease Virus (IBDV) and Infectious Bronchitis

Virus (IBV) infections, and were sanitary controlled and declared free of *Mycoplasma*

gallisepticum, *Mycoplasma synoviae*, *Chlamydia psittaci* and *Salmonella pullorum*

gallinarum. Only *Ascaris* and *Heterakis* parasites were detected in hen faeces, and

they were at a very low level.

Each hen was placed separately in a wire-latticed pen (3 m^2) in the experimental pens

described above with a nest box, perch, feed and water ad libitum. Hens were

accustomed to their new environment for 12 days, fed with a standard rearing diet for

laying hens (30099G25, Arrivé Nutrition Animale, Saint-Fulgent, France) and allowed

to deposit faecal and caecal microbiota on litter. The room temperature was 25 °C and

the artificial photoperiod was 16 h L:8 h D before egg deposition, 20 h L:4 h D during

hatching and the same programme as the chicks afterwards. Two days before chick

arrival or egg hatching, a wire-latticed space for chicks was placed in their pen. Eight

hens were used for 8 groups of 18 OFH chicks, and eight hens were used for 8 groups

of 18 CH chicks. On D0, day-old CH chicks were placed under the pen's wire-latticed

space, and OFH chicks were already under this space. Chicks and hens were in visual

and auditory contact for a few hours. Then hens were deprived of feed and water from

the morning. When lights were switched off, the hens were shut up in their nest boxes,

and chicks were placed under each hen as gently as possible for 11 h without any feed

and water. Chicks and hens were put physically together in a closed nest for the night

to promote maternal behaviour and the acceptance of chicks (Richard-Yris &

Leboucher, 1987). The following morning, one hour before the lights were switched on,

the nest-box doors were taken away to allow free access to the whole pen. Free in-

access feed and water were placed under wire-latticed space for chicks and in raised

troughs for hens, not accessible for chicks. Hens were present with chicks for two

weeks, the critical period for chick start, and removed on D15. Weight and clinical

examinations of the hens were recorded the day before they were installed in the pens

and, on D15, when they were removed.

Behavioural observations

The scan sampling method was used to follow the behaviour of hens and chicks on

days 2, 5, 6, 7, 8, 9, 12, 13 and 14 with the following repertoire: resting (the hen is lying

or standing still, eyes closed and without chicks), maintenance (preening, scratching,

stretching), feeding behaviour (the hen is eating or drinking), locomotion, exploration

(the hen is scratching or pecking at the ground or the environment), observation (the

hen is observing the environment with neck movements), maternal behaviour (the hen

is making food offering to the chicks, the hen is expressing maternal calls, the hen is

brooding the chicks by lying down and spreading her wings), fear behaviour (the hen

is flying or running from the experimenter, freezing, alert), agonistic behaviour (the hen

is chasing the chicks, the hen is pecking the chicks, others (punctual behaviours like

vocalisations).

To evaluate the proximity between chicks and hens, the experimenter also recorded

the localisation of four randomly tagged chicks and the hen within the pen. To that end,

the pen was virtually divided into four zones (Figure 1). The observations were

conducted between 10 AM and noon and between 3 and 5 PM by the same

experimenter. The experimenter walked slowly in front of each pen and recorded the

behaviour of the hen and the localisation of the four tagged chicks every eight minutes

(approximately), with a total of 10 scans per hen per day and 177 scans per hen for

the whole period of observation.

Figure 1. Schematic representation of the pen ($3m^2$) with the zones used to locate the

four tagged chicks and the hen during behavioural observations; A: the nest, B and D:

two halves of the pen and C: the wire-latticed space for the chicks (101×50 cm).

Chick quality scores

Chick quality scores were determined at placement in the pen for CH chicks, corresponding to 21 days of incubation for OFH chicks, on 24 to 25 chicks from the three treatments: CH, OFH and OFH + H. They were macroscopically defined according to the grid of Tona (Tona et al., 2003) and modified by adding several other parameters issued from the CASDAR QUALICOUV project (Guinebretière et al., 2022). Briefly, the chicks were scored on a total score of 110, including scores of posture (on 5), down (on 5), legs (on 6), red dot on the beak (on 10), grouped into an "*appearance*" score (on 26); activity (on 6), eyes (on 16), leg joint inflammation (on 5) and leg dehydration (on 5) were grouped into a "*tiredness*" score (32), and finally, retracted yolk (on 12), navel (on 12), remaining membrane (on 12), and remaining yolk (on 16) were grouped in an "*abdomen*" score (on 52).

Health parameters

Droppings deposited on pen litter were collected on D14 and D54 and analysed for parasite detection (*Coccidia*, *Ascaris* and *Heterakis*). Five grams of droppings were homogenised in 70 mL of flotation solution (0.36% of sodium chloride). The mixture was then filtered and pressed through a tea strainer (small mesh) to extract as much of the liquid part as possible. A homogeneous sample was deposited into a McMaster cell counter, and after 5 min of rest, the oocysts and nematode eggs were counted, and their number was expressed per gram of droppings (OPG). Health disorders, mortality and causes of death were registered during the experiment.

Performance

Body weight (BW) was measured at D1, D19, D34 and D55. Feed consumption was measured in each pen for the periods between D1–D19, D19–D34 and D34–D55, and then used to calculate the feed conversion ratio (FCR) as the feed consumption-to-BW gain ratio per pen during both periods and the entire rearing period. At D56, 16 identified males per group were slaughtered, and *pectoralis major* and *pectoralis minor* (breast) muscles were weighed to calculate their yields relative to BW and ultimate pH. Ultimate pH was measured as the pectoralis major pH 24 hours after slaughter.

Statistical analyses

Hatching rates between hatchery and on-farm hatchings were compared using chi-squared tests. Chick quality parameters were analysed by a non-parametric Kruskal-Wallis test, considering the treatment (CH, CH + H, OFH and OFH + H), followed by Mann-Whitney post hoc tests. The normality of residual distribution was checked with the Shapiro-Wilk test for BW, feed intakes and FCR. A 2-way ANOVA was then carried out to test the hatching condition and the sex effect, as well as the two-by-two interactions on performance. When there was an interaction between variables, a Fisher (LSD) test was used to determine the statistical significance of the difference. Differences were considered significant when p-values < 0.05 and a tendency for $0.1 < p < 0.05$. Analyses were performed using XLSTAT software (version 2015, Addinsoft, Paris, France).

Behavioural data did not meet the assumption of normality and homogeneity of variances. Non-parametric Mann-Whitney U-tests were used on the mean percentage

of scans per behavioural category to compare the behaviour of hens in contact with

CH chicks to the hens in contact with OFH chicks. To compare the proximity of CH and

OFH chicks towards the hen, Mann-Whitney U tests were conducted on the mean

number of tagged chicks located in the same area of the pen as the hen over the 177

scans recorded per hen.

Results

Behavioural observations

Because 3 hens (1 OFH + H and 2 CH + H) were very aggressive and injured their

chicks, they were removed from the pens and the later behavioural analysis. There

was no significant difference in the behaviour of the hens, regardless of the hatching

condition of chicks, except for the frequency of the behaviour "*observe*"; OFH hens

tended to observe their environment less than CH hens (Table 1).

Table 1. Behaviour of hens according to the chick hatching conditions

| Hen behaviour | Hatching conditions | | P-value |
	CH	OFH	
Agonistic	2.54 ± 3.74	1.37 ± 0.72	0.550
Rest/Comfort	17.72 ± 7.16	31.16 ± 22.74	0.181
Fear	7.07 ± 3.39	4.92 ± 1.95	0.384
Feeding	18.10 ± 4.52	19.45 ± 11.56	0.731
Locomotion	6.78 ± 4.12	3.39 ± 2.95	0.146
Observation	17.53 ± 7.45	9.52 ± 4.76	0.045
Exploration	22.62 ± 7.62	19.77 ± 10.78	0.656
Maternal	1.32 ± 1.94	3.39 ± 7.98	0.732
Others	6.32 ± 2.31	7.02 ± 7.02	0.470

CH = conventional hatchery (n = 6), OFH = on-farm hatching (n = 7)
Behaviour observations (mean ± SD of scan percentage over 9 days)
P-value < 0.05 = significant difference between hatching conditions (Mann-Whitney U-test)

To characterise hens' behaviour towards the chicks, each hen was categorised

according to the frequencies of agonistic or maternal behaviours (Table 2). We defined

three categories: 1) maternal (M): the hens expressed only maternal behaviours

towards the chicks; 2) tolerant (T): the hens expressed both maternal and agonistic

behaviours towards the chicks or less than 2% of scans with maternal behaviour; 3)

aggressive (A): the hens rejected the chicks and expressed only agonistic behaviour

towards them. Two hens were defined as maternal, six were tolerant, and five were

aggressive among the 13 hens analysed (Table 2).

Table 2. Classification of hen according to the frequencies
of maternal or agonistic behaviours expressed towards chicks.

Hatching conditions	Hen behaviours		Category
	Agonistic	Maternal	
CH1	7.91 ± 0.27	0	A
CH2	0	0.57 ± 0.07	T
CH3	0.56 ± 0.07	0.56 ± 0.07	T
CH4	0	5.08 ± 0.22	M
CH5	0	1.69 ± 0.13	T
CH6	6.78 ± 0.25	0	A
OFH1	1.13 ± 0.11	0	A
OFH2	0	21.47 ± 0.41	M
OFH3	1.69 ± 0.13	0.56 ± 0.07	T
OFH4	1.69 ± 0.13	0.56 ± 0.07	T
OFH5	1.13 ± 0.11	1.13 ± 0.11	T
OFH6	1.69 ± 0.13	0	A
OHF7	2.26 ± 0.15	0	A

Thirteen hens were classified, three agressive hens
(1 OFH+H and 2 CH+H) were removed from the analysis
CH = conventional hatchery; OFH = on-farm hatching
Behaviour observations (mean ± SD of scan percentages over 9 days)
A = Agressive (maternal behaviour = 0)
T = Tolerant (0 < maternal behaviour < 5)
M = Maternal (maternal behaviour > 5 and agonistic behaviour = 0)

The mean number of chicks observed in the same area as the hen did not differ

significantly between CH and OFH chicks ($p > 0.05$). (Figure 2).

Figure 2. Proximity between chicks and hens according to hatching conditions; four

chicks were observed per pen (n ≤ 8 scans per day) per hatching condition

(conventional hatchery, CH or on-farm hatching, OFH)

Hatchability and chick quality

Hatchability

For conventional hatchers, 97.7% of CH fertile eggs hatched at E21 and 97.2% ± 4.2%

of OFH fertile eggs hatched at E21 in pens. The presence of hens had a significant

impact on the OFH condition (p = 0.034). In the presence of hens, 86.8% ± 11.9% of

OFH + H chicks hatched at E21 (Figure 3). Unhatched eggs were mainly found in the

pens with aggressive hens (9/11) or in the OFH pens next to those with aggressive

hens (4/4). No mortality of CH chicks or OFH chicks was observed at hatching,

whereas 5.6% ± 5.9% OFH + H chicks died or were removed at hatching (n = 10); due

to three hens' aggressiveness or another reason. Only 3.6% (2/56) of chicks had

residual yolk sacs at the age of 20 days (one CH and one CH + AB) and no yolk residue

was found at 56 days.

Figure 3. Number of live hatched chicks according to hatching conditions; conventional

hatchery (CH) condition performed in one hatchery (one value); on-farm hatching

(OFH) and on-farm hatching with hen (OFH + Hen) conditions were repeated in eight

pens each, each pen contained 18 embryonated eggs or chicks; values are expressed

as means ± standard error

Quality scores of chicks

Whereas no difference was shown due to the hatching conditions (p > 0.05) on the

total quality scores, with good scores in the three groups considered (OFH: 96.2 ± 1.5,

CH: 97.3 ± 1.5; CH+H: 95.1 ± 1.7), when considering the subtotal scores linked to the

appearance, the tiredness or the abdomens of the chicks, it appeared that the subtotal

of the appearance score changed depending on the treatment (Figure 4), with the two

other subtotals not being significantly changed ($p > 0.05$, data not shown). Indeed,

whereas the subtotal score for appearance was not different between CH chicks or

OFH chicks, it was deteriorated by the presence of the hen within the hatching pen in

OFH chicks ($p = 0.01$).

Figure 4. Chick appearance subtotal score at the placement in the pen according to

hatching conditions; appearance scores noted on 26 included scores of posture (on

5), down (on 5), legs (on 6), and a red dot on the beak (on 10); n = 24 to 25

chicks/hatching condition; conventional hatchery (CH), on-farm hatching (OFH), OFH

+ hen (OFH + H)

Performance

Hatching conditions significantly influenced chick BW from hatching to slaughter age.

The OFH chick BW was significantly greater than that of all CH chicks at hatching,

whether hens were present or not (p ≤ 0.002, Figure 5). Independently of the treatment,

a sex effect was observed from D19 onwards; male chicken BWs were greater than

those of females (males: 503 ± 46g, females: 469 ± 37g, p = 0.0001). Treatment of CH

chicks with antibiotics temporarily decreased chicken BW at D19 (p = 0.035) (Figure

5) due to a decrease in weight gain (CH: 497 ± 38g, CH + AB: 486 ± 37g, p = 0.0001)

compared to CH chickens, while feed intake (data not shown) and FCR were not

different (Table 3). At D19, OFH chickens had the best BW compared to all other

groups of chicks (p ≤ 0.0003) (Figure 5). At this time, the presence of hens at hatching

had a remnant negative impact on chicken BW regardless of the hatching condition (p

< 0.0001), as well as on FCR (Table 3). Both the feed intake per chicken (CH: 624 ±

12g^a, CH + AB: 600 ± 27g^{ab}, CH + H: 603 ± 25g^{bc}, OFH: 652 ± 33^a, OFH + H: 615 ±

34^c, p = 0.001) and the weight gained per chicken (CH: 455 ± 37g^b, CH + AB: 445 ±

37g^c, CH + H: 421 ± 40g^d, OFH: 471 ± 42g^a, OFH + H: 425 ± 47g^d, p = 0.0001)

decreased compared to the other groups, and the FCR increased (Table 3). An

interaction between the effect of the hatching condition and chicken sex on BW was

observed later at D34 (p = 0.012) and D56 (p = 0.022) on BW, even though the FCR

was not affected (Table 3). At D34, a week after the challenge, the OFH male chickens

were the heaviest compared to the other groups (p ≤ 0.033) and the presence of hens

at hatching harmed chicken BW (p ≤ 0.0004), regardless of the hatching condition

(Figure 6A). In females, there was no effect of hatching condition or presence of hens

on the BW at D34 (Figure 6A). At slaughter age (D56), there was no effect of hatching

condition on the male chicken BW, but the presence of hens at hatching harmed CH

chicken BW (p = 0.0008) (Figure 6B). There was a pen effect in CH + H (p = 0.016)

and OFH + H chickens (p = 0.001), the pen with the lightest CH + H males was in the

presence of an aggressive hen, and the heaviest OFH + H males were in a pen in the

presence of a tolerant hen, but all combinations were observed (Figure 7). In females,

there was no effect of the hatching condition on the BW. The presence of hens at

hatching had a positive impact on OFH female chickens compared to CH female

chicken BW (p = 0. 0096), with the OFH + H chickens being the heaviest compared to

the other CH female conditions (Figure 6B). There was no significant pen effect

between CH + H and OFH + H female chickens (p = 0.447).

Table 3. Performance according to the hatching conditions of chicks

Day ranges	Feed conversion ratio (g/g)					
	CH	CH + AB	CH + H	OFH	OFH + H	
D1–D19	1.370 ± 0.024c	1.350 ± 0.066c	1.416 ± 0.049ab	1.388 ± 0.022bc	1.447 ± 0.035a	0.001
D20–D34	1.807 ± 0.030	1.773 ± 0.042	1.769 ± 0.039	1.795 ± 0.035	1.787 ± 0.057	0.355
D35–D55	2.194 ± 0.091	2.213 ± 0.055	2.188 ± 0.054	2.201 ± 0.049	2.141 ± 0.038	0.173
D1–D55	1.904 ± 0.036	1.902 ± 0.025	1.913 ± 0.040	1.910 ± 0.022	1.912 ± 0.015	0.924

Hatching conditions: conventional hatchery (CH), CH + antibiotics treatment (CH + AB), CH + hen (CH + H), on-farm hatching (OFH), OFH + hen (OFH + H)

Values are expressed as mean ± standard error

a,b,c, Different letters correspond to significant differences between treatment groups

381 Figure 5. Body weight at D1 and D19 and according to the hatching conditions:

382 conventional hatchery (CH), CH + antibiotics treatment (CH + AB), CH + hen (CH +

383 H), on-farm hatching (OFH), OFH + hen (OFH + H); values are expressed as means ±

384 standard error; different letters correspond to significant differences between treatment

385 groups

A

B

Figure 6. Weight at D34 (A) and D56 (B) of male and female chickens according to the

hatching conditions: conventional hatchery (CH), CH + antibiotics treatment (CH + AB),

CH + hen (CH + H), on-farm hatching (OFH), OFH + hen (OFH + H); values are

expressed as mean ± standard error: different letters correspond to significant

differences between treatment groups

Figure 7. Body weight at D56 of male chickens according to the behaviour of the hen

present at the starting period, M: maternal, T: tolerant, A: aggressive, AR: aggressive

and removed from the pen; CH + H: chicks hatched in the hatchery and in the presence

of hens; OFH + H: chicks hatched on-farm in the presence of hens; median ± SD (n ≤

9).

Breast weight was not affected by the hatching conditions (6.99 ± 0.06, p = 0.357) and

ultimate pH was not modified either (5.7 ± 0.1, p = 0.951).

Health and robustness

Coccidia was detected in variable amounts in the droppings of all the pens at D54

(200–85500 OPG) without any significant effect of the hatching conditions in the

presence of hen or not (p = 0.606). No clinical signs were observed during the

experiment. In all hatching conditions combined, the viability rate of the chickens was

95.3%. The mortality rate during the whole experiment was 3.19% (23/720). Seventeen

chicks died during the first week of life, 11 OFH + H and 5 CH + H in the presence of

hens and one OFH chick for an unknown reason. Six CH chickens died during the rest

of the experiment, five of which were due to heart problems (2 CH, 1 CH + AB, 2 CH

+ H) and one to unknown causes (CH + H). Eleven chicks were additionally eliminated

after hatching in pens in the presence of hens (4 at D1, 4 at D2, and 1 at D4) and two

later (D33 and D55) for morphological reasons.

Discussion

New hatching systems are being developed in Europe, and the enrichment of the

rearing environment is also in full development, notably by optimising the microbial

environment of the chicks to limit the use of antibiotics. In this study, we analysed the

benefits/risks of hatching systems (OFH and CH, treated with antibiotics or not) and of

the presence of an adult hen or not on hatchability, chick quality score, performance,

health and robustness.

Hatching conditions

The BW of OFH-certified JA757 chicks was significantly greater than that of CH chicks

at hatching, even though the hatchability rate and the quality score of chicks were

comparable between the two conditions, and no mortality was reported. These results

agree with other studies on OFH performed in slow (Jessen et al., 2021) and fast-

growing broilers (de Jong et al., 2020; Souza da Silva et al., 2021) for the BW but not

for other parameters that were reported higher for the hatchability, lower for the quality

score of chicks and lower for the mortality. However, whereas there was no effect of

hatching conditions, the presence of hens, categorised according to their behaviour,

decreased the hatchability rate, the appearance quality score of OFH chicks and
increased mortality at hatching. These degraded indicators could be since in our
experimental design, very few hens expressed a clear maternal behaviour towards the
chicks (n = 2/16), and some even showed agonistic behaviour. This may be explained
by the genetic line of hens used (Lohmann Brown), which is highly selected for laying
and counter-selected for brooding. However, this genetic line was chosen because the
studied practice could favour the possibility to use culled hens in breeding, and
because of their rather tolerant social behaviour, their brooding behaviour could be
optimised. Improvements could be obtained by carrying it out in a season with days
with greater light amplitudes (spring) to facilitate brooding behaviour, which was not
the case in this study (winter), and by selecting hens with brooding behaviour to
facilitate maternal behaviour (Shimmura et al., 2010). In addition, in our experimental
design, the chicks had to feed under the wire-lattice space, which was not accessible
to the hen. As they obtained both food and warmth under this space, the hens probably
did not have enough tactile stimulation from the chicks to fully express their maternal
behaviour with no agonistic behaviour. Indeed, in addition to the physiological state,
tactile stimulations from chicks play an important role in the expression and
maintenance of maternal behaviour in hens (Richard-Yris & Leboucher, 1987).

Starting period

Hatching conditions and the presence of hens for 15 days after placement significantly
influenced chick performance during the starting period. At D19, OFH chicks had the
best BW compared to the other groups. and the presence of hens at hatching harmed
chicken BW regardless of the hatching condition and on FCR. No significant
differences were observed in the behaviour of hens present with OFH and CH chicks,

except for OFH hens, which were found to observe their environment less than CH

hens. With our small sample size, this result could be explained by the behaviour of

one OFH hen, which spent much of the time resting. The CH and OFH chicks did not

differ in their proximity towards the hen. The mean number of chicks observed in the

same area as the hen was very low (less than 1 chick), indicating that they were rarely

in contact with the hen. However, chick performance was affected by the presence of

the hens, including lower feed intake and consequently lower weight gain and higher

FCR. This could be explained by the agonistic behaviour of some hens towards chicks,

the attempt of the hens to eat the chick feed and the stress that this may have caused

the chicks.

Treatment of CH chicks with antibiotics, assessed as growth promoters, temporarily

decreased chicken BW at D19, but FCR was not modified. This effect was not

observed later, but growth promotion was not observed in CH chicks treated with

antibiotics. This result is not in agreement with the use of antibiotics as growth

promoters in farm animals, but the relative lack of published data on chicken

performance limits knowledge of the actual effects of antibiotics on animal performance

(Kumar et al., 2018; Broom, 2018). Their effects also result from their interaction with

the microbiota and the variables chosen in the experimental studies. The effects

observed in farms are dependent on the sanitary conditions present, which are

different from the much more controlled sanitary conditions in the experimental studies

and may contribute to different effects of treatment with antibiotics.

Growth period

An interaction between the effect of hatching conditions and chicken sex was observed

on BW after the challenge on D27. In males, the OFH chicken group was the heaviest

compared to the other groups at D34 but not at D56. These results are consistent with

a previous study that observed the beneficial effects of OFH on BW only until D21 (de

Jong et al., 2020), and not until slaughter time, as reported in various studies when

post-hatching feed deprivation time was at least 36 h (de Jong et al., 2017).

Alternatively, this may also be a result of the response to the challenge experienced

by the chickens at D27, including transport, exposure to low temperature, transient

feed deprivation, vaccination and a change to a higher rearing density, but the fact is

that there is no ultimate positive impact of OFH on BW at slaughter time. Moreover, in

our conditions, the presence of hens eventually negatively impacted male chicken BW,

but only for CH chickens at D56. In females, there was no effect of hatching conditions

on the BW at D34 and D56, and the presence of hens eventually had a positive impact

on OFH female chicken BW. These results were unexpected, but it is known that early

stress induces sex-specific, immediate and life-long effects on the stress response,

behaviour, sex hormones, and hypothalamic and blood gene expression in chickens

(Madison et al., 2008; Elfwing et al., 2015; Foury et al., 2020), with the males being

more reactive than the females. The results observed in this study raise questions

about the consequences of hatching conditions in the presence of a hen according to

the sex of the chicks. It appeared that male OFH chicks developed more fear and

stress responses than females when placed in the presence of a hen that was not their

mother, and this had effects on their growth until slaughter age. For male OFH chicks,

in which the effect of hen presence on their growth was only observed during the

growth phase, there was possibly communication between hens and embryonated

eggs before hatching and with chicks at hatching that may have increased their fear

and stress responses and therefore harmed their growth. This could even have had

negative consequences on hatchability and mortality rates, but the sex of the chicks

was not recorded at that time. The presence of hens with the female OFH chicks did

not affect their performance and even had a beneficial effect on their growth at

slaughter age. These differences observed between treatments and chick sexes for

performance are not likely explained by a difference in proximity between hens and

chicks, which was low in this experiment.

Health and Robustness

There were no effects of hatching conditions on health parameters (parasitic load,

clinical signs, rate of mortality), even after exposure of chickens during their growth

phase to an environmental and vaccine challenge. However, this challenge could have

accentuated the differences in the effects of hatching conditions on performance

parameters between males and females, but we did not perform the unchallenged

rearing conditions to assert this. The implantation of adult microbiota into the chick

digestive system by the presence of hens should be nevertheless beneficial for the

maturation of the chick microbiota and gut immune system and still needs to be

assessed.

Altogether, on-farm hatching of certified broilers was a hatching system at least

equivalent to the hatchery hatching system, if not better, in this study. The effects of

the hens' presence at hatching and during the chick start-up phase on performance

interacted with the hatching condition and the sex of the chickens. In this case, the

health status and brooding behaviour of the hens are essential to ensure the health

and welfare of the chicks. These practices offer possible evolutions of the rearing

conditions to continue to decrease the use of antibiotics.

Ethics approval

All experimental procedures were approved by the Ethics Committee COMETHEA POITOU-CHARENTES n°84 (APAFIS#24474-2020021816237418 v3) and carried out following current European legislation (EU Directive 2010/63/EU).

Author contributions

LAG, AB, CS, KG and AC designed the study with the help of CB. LAG, CB and AC performed the experiment with the technical help of SC for the organisation of the experiment and AH for parasitic analyses. CB and LR collected the performance and health parameters. LAG analysed data with the help of AB and CB for the behaviour data. LAG, AB and CB wrote the paper with the help of KG and AC. All the authors reviewed and approved the manuscript.

Author ORCIDs

LG: https://orcid.org/0000-0001-7089-2196

AB: https://orcid.org/0000-0001-5647-1758

CB: https://orcid.org/0000-0003-1181-992X

CS: https://orcid.org/0000-0002-3480-6278

KG: https://orcid.org/0009-0005-6638-9404

AC: https://orcid.org/0000-0002-3410-6108

Acknowledgements

We are grateful to all the members of the RIMEL network whose shared thinking made the design of this study possible. We thank the staff of the poultry alternative breeding experimental unit (EASM, INRAE, 17700 Surgères, France, DOI: 10.15454/1.5572418326133655E12) for the development of the experimental set-up and the conduct of the experimentation. We are very grateful to the staff of the MOQA team (INRAE, 37380 Nouzilly, France) for their help during the experimentation. The manuscript has been professionally proofread.

Funding

This research was supported by a grant from INRAE, Department of Animal Physiology and Livestock Systems for the RIMEL network.

Data and model availability statement

The datasets used during the current study are available on line:

https://doi.org/10.57745/6INVYL

References

Beauclercq S, Lefèvre A, Montigny F, Collin A, Tesseraud S, Leterrier C, Emond P, Guilloteau LA (2019) A multiplatform metabolomic approach to characterize fecal signatures of negative postnatal events in chicks: a pilot study. J Anim Sci Biotechnol, 10, 21. https://doi.org/10.1186/s40104-019-0335-8

Broom LJ (2018) Gut barrier function: Effects of (antibiotic) growth promoters on key

barrier components and associations with growth performance. Poult Sci, 97,

1572-1578. https://doi.org/10.3382/ps/pey021

Broom LJ, Kogut MH (2018) The role of the gut microbiome in shaping the immune

system of chickens. Vet Immunol Immunopathol, 204, 44-51.

https://doi.org/10.1016/j.vetimm.2018.10.002

de Jong IC, van Riel J, Bracke MBM, van den Brand H (2017) A 'meta-analysis' of

effects of post-hatch food and water deprivation on development, performance

and welfare of chickens. PLoS One, 12, e0189350.

https://doi.org/10.1371/journal.pone.0189350

de Jong IC, Gunnink H, van Hattum T, van Riel JW, Raaijmakers MMP, Zoet ES, van

den Brand H (2019) Comparison of performance, health and welfare aspects

between commercially housed hatchery-hatched and on-farm hatched broiler

flocks. Animal, 13, 1269-1277. https://doi.org/10.1017/S1751731118002872

de Jong IC, van Hattum T, van Riel JW, De Baere K, Kempen I, Cardinaels S, Gunnink

H (2020) Effects of on-farm and traditional hatching on welfare, health, and

performance of broiler chickens. Poult Sci, 99, 4662-4671.

https://doi.org/10.1016/j.psj.2020.06.052.

Edgar J, Held S, Jones C, Troisi C (2016) Influences of maternal care on chicken

welfare. Animals (Basel), 6, 2. https://doi.org/10.3390/ani6010002

Elfwing M, Nätt D, Goerlich-Jansson VC, Persson M, Hjelm J, Jensen P (2015) Early

stress causes sex-specific, life-long changes in behaviour, levels of gonadal

hormones, and gene expression in chickens. PLoS One, 10, e0125808.

https://doi.org/10.1371/journal.pone.0125808

Fortun-Lamothe L, Collin A, Combes S, Ferchaud S, Germain K, Guilloteau L, Gunia

M, Lefloc'h N, Manoli C, Montagne L, Savietto D (2023) Principes, cadre

d'analyse et leviers d'action à l'échelle de l'élevage pour une gestion intégrée de

la santé chez les animaux monogastriques. INRAE Prod Anim, 35, 307–326.

https://doi.org/10.20870/productions-animales.2022.35.4.7225

Foury A, Collin A, Helbling JC, Leterrier C, Moisan MP, Guilloteau LA (2020)

Spontaneous intake of essential oils after a negative postnatal experience has

long-term effects on blood transcriptome in chickens. Sci Rep, 10, 20702.

https://doi.org/10.1038/s41598-020-77732-5

Giersberg M F, Poolen I, de Baere K, Gunnink H, van Hattum T, van Riel J W, de Jong

IC (2020). Comparative assessment of general behaviour and fear-related

responses in hatchery-hatched and on-farm hatched broiler chickens. Appl Anim

Behav Sci, 232, 105100. https://doi.org/10.1016/j.applanim.2020.105100

Giersberg MF, Molenaar R, de Jong IC, Souza da Silva C, van den Brand H, Kemp B,

Rodenburg TB (2020) Effects of hatching system on the welfare of broiler

chickens in early and later life. Poult Sci, 100, 100946.

https://doi.org/10.1016/j.psj.2020.12.043

Guinebretière M, Puterflam J, Keïta A, Réhault-Godbert S, Thomas R, Chartrin P,

Cailleau-Audouin E, Coudert E, Collin A (2022) Storage temperature or thermal

treatments during long egg storage duration influences hatching performance

and chick quality. Front Physiol, 13, 852733.

https://doi.org/10.3389/fphys.2022.852733

Hewlett SE, Nordquist RE (2019) Effects of maternal care during rearing in White

Leghorn and Brown Nick Layer Hens on cognition, sociality and fear. Animals

(Basel), 9, 454. https://doi.org/10.3390/ani9070454

Hollemans MS, de Vries S, Lammers A, Clouard C (2018) Effects of early nutrition and
transport of 1-day-old chickens on production performance and fear response.
Poult Sci, 97, 2534-2542. https://doi.org/10.3382/ps/pey106

Jessen CT, Foldager L, Riber AB (2021) Effects of hatching on-farm on performance
and welfare of organic broilers. Poult Sci, 100, 101292.
https://doi.org/10.1016/j.psj.2021.101292

Kumar S, Chen C, Indugu N, Werlang GO, Singh M, Kim WK, Thippareddi H (2018)
Effect of antibiotic withdrawal in feed on chicken gut microbial dynamics,
immunity, growth performance and prevalence of foodborne pathogens. PLoS
One, 13, e0192450. https://doi.org/10.1371/journal.pone.0192450

Madison FN, Jurkevich A, Kuenzel WJ (2008) Sex differences in plasma corticosterone
release in undisturbed chickens (Gallus gallus) in response to arginine vasotocin
and corticotropin releasing hormone. Gen Comp Endocrinol, 155, 566-573.
https://doi.org/10.1016/j.ygcen.2007.08.014

Meijerink N, Kers JG, Velkers FC, van Haarlem DA, Lamot DM, de Oliveira JE, Smidt
H, Stegeman JA, Rutten VPMG, Jansen CA (2020) Early life inoculation with
adult-derived microbiota accelerates maturation of intestinal microbiota and
enhances NK cell activation in broiler chickens. Front Vet Sci, 7, 584561.
https://doi.org/10.3389/fvets.2020.584561

Perré Y, Wauters A-M, Richard-Yris M-A (2002) Influence of mothering on emotional
and social reactivity of domestic pullets. Appl Anim Behav Sci, 75, 133-146.
https://doi.org/10.1016/S0168-1591(01)00189-7

Richard-Yris MA, Leboucher G (1987) Responses to successive test of induction of
maternal behaviour in hens. Behav Processes, 15, 17-26.
https://doi.org/10.1016/0376-6357(87)90030-1

Shimmura T, Kamimura E, Azuma T, Kansaku N, Uetake K, Tanaka T (2010) Effect of

broody hens on behaviour of chicks. Appl Anim Behav, 126, 125-133.

https://doi.org/10.1016/j.applanim.2010.06.011

Souza da Silva C, Molenaar R, Giersberg MF, Rodenburg TB, van Riel JW, De Baere

K, Van Dosselaer I, Kemp B, van den Brand H, de Jong IC (2021) Day-old chicken

quality and performance of broiler chickens from 3 different hatching systems.

Poult Sci, 100, 100953. https://doi.org/10.1016/j.psj.2020.12.050

Tona K, Bamelis F, De Ketelaere B, Bruggeman V, Moraes VM, Buyse J, Onagbesan

O, Decuypere E (2003) Effects of egg storage time on spread of hatch, chick

quality, and chick juvenile growth. Poult Sci, 82, 736-741.

https://doi.org/10.1093/ps/82.5.736

van de Ven LJ, van Wagenberg AV, Groot Koerkamp PW, Kemp B, van den Brand H

(2009) Effects of a combined hatching and brooding system on hatchability, chick

weight, and mortality in broilers. Poult Sci, 88, 2273-2279.

https://doi.org/10.3382/ps.2009-00112

van der Wagt I, de Jong IC, Mitchell MA, Molenaar R, van den Brand H (2020) A review

on yolk sac utilization in poultry. Poult Sci, 99, 2162-2175.

https://doi.org/10.1016/j.psj.2019.11.041

Volf J, Polansky O, Varmuzova K, Gerzova L, Sekelova Z, Faldynova M, Babak V,

Medvecky M, Smith AL, Kaspers B, Velge P, Rychlik I (2016) Transient and

prolonged response of chicken cecum mucosa to colonization with different gut

microbiota. PLoS One, 11, e0163932.

https://doi.org/10.1371/journal.pone.0163932