The origin, evolution and molecular diversity of the chemokine system

Alessandra Aleotti[1,2,*,#], Matthew Goulty[1,2,*,#], Clifton Lewis[1,2], Flaviano Giorgini[1,2] and Roberto

Feuda[1,2,#]

[*]Equal contribution

[#]Author for correspondence: rf190@leicester.ac.uk; aa1176@leicester.ac.uk; mg478@leicester.ac.uk

[1]Neurogenetics Group

[2]Department of Genetics and Genome Biology University of Leicester, Leicester, United Kingdom

Keywords: Chemokine, Phylogeny, Evolution, Vertebrate, GPCR, Receptor, Ligand, CK, CKL, CKR,

CKLF, TAFA, CYTL

ABSTRACT

Chemokine signalling performs key functions in cell migration via chemoattraction, such as attracting

leukocytes to the site of infection during host defence. The system consists of a ligand, the

chemokine, usually secreted outside the cell, and a chemokine receptor on the surface of a target

cell that recognises the ligand. Several non-canonical components interact with the system. These

include a variety of molecules that usually share some degree of sequence similarity with canonical

components and, in some cases, are known to bind to canonical components and/or to modulate

cell migration (1, 2). While canonical components have been described in vertebrate lineages, the

distribution of the non-canonical components is less clear. Uncertainty over the relationships

between canonical and non-canonical components hampers our understanding of the evolution of

the system. We used phylogenetic methods, including gene-tree to species-tree reconciliation, to

untangle the relationships between canonical and non-canonical components, identify gene

duplication events and clarify the origin of the system. We found that unrelated ligand groups

independently evolved chemokine-like functions. We found non-canonical ligands outside

vertebrates, such as TAFA "chemokines" found in urochordates. In contrast, all receptor groups are

vertebrate-specific and all - except ACKR1 - originated from a common ancestor in early vertebrates.

Both ligand and receptor copy numbers expanded through gene duplication events at the base of

jawed vertebrates, with subsequent waves of innovation occurring in bony fish and mammals.

INTRODUCTION

The chemokine system is responsible for regulating many biological processes, including host defence, neuronal communication and homeostasis (3–5). The system has two components, a ligand, usually a small cytokine called a chemokine, and a receptor. It typically operates through chemoattraction, wherein one cell type produces and secretes chemokines, creating a chemical gradient as these molecules disperse. Cells equipped with the corresponding chemokine receptors on their membranes can recognise and bind to specific chemokines, promoting their migration along the gradient (4). This mechanism allows cells to reach target locations, such as infection sites during inflammation or tissues important for homeostatic functions, e.g., leukocyte maturation and trafficking (3, 6). Chemokines involved in the latter homeostatic functions are usually constitutively expressed, while those involved in inflammatory responses have an inducible expression (7). Chemokine ligands are categorised into four groups, XC, CC, CXC, and CX3C, according to the pattern of cysteine residues in the N-terminal portion of the protein (8). Likewise, the receptors are classified based on the ligands they bind to into four groups, the XCR, CCR, CXCR, and CX3CR, and all of them belong to the GPCR class A superfamily (9). In addition to canonical components, other molecules have been discovered to function similarly to chemokine ligands (1) or receptors (2) (see Table 1). These include: the chemokine-like factor (CKLF) that binds to chemokine receptor CCR4 (10, 11) and drives cell migration *in vivo* (12); TAFA chemokines, expressed mainly in the nervous system, which share structural similarities to canonical chemokines (13, 14) and bind GPCRs related to chemokine receptors, e.g. formyl peptide receptors (15, 16) and GPR1 (17); Cytokine-like 1 (CYTL1) that binds CCR2 (18) and has been suggested to be related to CC ligands based on the presence of a IL8-like chemokine fold (19). There are also non-canonical chemokine receptors, such as: the chemokine-like receptor (CML1, or also CMKLR1) ((20); atypical chemokine receptors (ACKRs) (21); and viral chemokine receptors (22–25). Unlike other chemokine receptors, atypical receptors cannot initiate classical chemokine signaling upon ligand binding (21, 26). The human genome encodes four types of atypical chemokine receptors: the ACKR1 (also known as DARC), ACKR2 (also known as D6), ACKR3

(also known as CXCR7) and ACKR4 (also known as CCRL1) (27, 28). Additionally, several proteins of

viral origins, such as US28 from human cytomegalovirus, have chemokine-receptor/binding activity

(22, 23). These viral proteins can bind a wide array of chemokine ligands (23).

Despite the extensive research on the chemokine system, with over 320,000 papers

available on PubMed, many aspects of its evolution remain unclear. For instance, the homology

between canonical and non-canonical ligands is uncertain and supported by circumstantial evidence,

such as shared specific motifs (12, 13, 19, 29). Furthermore, the relationships between canonical,

atypical, and viral receptors and the outgroup of the canonical chemokine receptors remain

uncertain. Finally, the evolutionary history of the canonical and non-canonical components remains

poorly understood outside a few key model systems (9, 30, 31). These outstanding questions share

common underlying causes, including the use of inadequate inference methods (such as relying

solely on sequence similarities) and limited sampling of species (e.g., focusing mainly on humans,

mice, and zebrafish (7, 32)). Additionally, solving the phylogenetic relationships for short molecules

such as chemokine receptors and ligands is particularly challenging due to the lack of strong

phylogenetic signals (33).

Here, to clarify these outstanding questions, we use state-of-the-art phylogenetic methods,

including those designed for single-gene phylogenies, a large taxonomical sampling comprising both

vertebrate and invertebrate genomes and the entire complement of canonical and non-canonical

components of both receptors and ligands. Our findings substantially clarify the phylogenetic

relationship between canonical and non-canonical ligands and receptors and suggest that unrelated

proteins evolved "chemokine-like" ligand function multiple times independently. Additionally, we

discovered that all the canonical and non-canonical chemokine receptors (except ACKR1) originated

from a single duplication in the vertebrate stem group, which also gave rise to many GPCRs. Lastly,

we characterized the complement of canonical and non-canonical components in the common

ancestor of vertebrates and identified several other ligands and receptors with potential chemokine-

related properties that could be explored in future functional work.

RESULTS

There are five unrelated groups of ligands.

92   Initially, we focused on the ligands, including all the canonical chemokines, the CYTL, the

TAFAs and the CKLF Super Family (CKLFSF) proteins (Table 1). The presence of a four transmembrane

MARVEL domain in the latter proteins (12, 34, 35) distinguishes them from canonical chemokines,

the CYTL and the TAFAs. Therefore, we separated these two groups for further analysis. Using

BLASTP or PSI-BLAST (36–38) (see Materials and Methods for more details) against 64 species from

19 animal phyla (Table S1), we identified 891 putative homologs for chemokines, TAFA and CYTL and

602 putative homologs of the CKLF Super Family.

99   We utilised CLANS (39, 40), a clustering tool based on sequence similarity and local

alignment, to identify homology within these two groups. Unlike traditional phylogenetic methods,

CLANS assigns homology between sequences based on BLAST and customisable stringency levels

defined according to p-values (39). When two (or more) sequences are connected at a lower p-value

(closer to 0), this indicates a high level of homology. Conversely, if two or more sequences only

connect at a higher p-value, this suggests a relatively low level of sequence homology. Our analysis

shows that canonical chemokines form a distinct group with a clear distinction between C-X-C-type

and C-C-type (Figure 1A). Whereas CXCL17, TAFA and CYTL remain separate from canonical

chemokines and from each other even at the loosest p-values tested (Figure 1A). The distinction

between CXCL17 and all other canonical chemokines is consistent with our receptor results, showing

that the potential receptor for CXCL17, GPR35 (41), is also not within the canonical chemokine

receptor group (see below). However, it is important to note that recent studies fail to demonstrate

CXCL17 activity at GPR35 (42, 43). Within the CKLFSF, two large clusters were identified, named CKLF

I and CKLF II, although these ultimately connect to form one large superfamily (Figure 1B). These

clusters are robust to the different stringency thresholds used (Figures S1 and S2 and Materials and

Methods for further details). Our results indicate that even when the stringency level to detect

homology is relaxed, canonical chemokines, TAFA, CYTL, and CXCL17 remain in distinct clusters. This

suggests that, similarly to CKLFs, these proteins are not homologous and convergently evolved

chemokine-like properties. We have thus identified five distinct groups of ligands: i) the canonical

chemokines, ii) TAFA "chemokines", iii) CYTL, iv) CXCL17, and v) CKLF Super Family (Figure 1A and

1B).

The evolution of chemokine and chemokine-like ligands in animals.

To better understand the evolution of both canonical and non-canonical chemokine ligands,

we performed a separate phylogenetic reconstruction for each group (Figure 1C, D and Figures S8-

17) (see methods for details). To evaluate the nodal support, in addition to the UltraFast bootstrap

(UFB) (44, 45), we used Transfer Bootstrap Expectation (TBE), a method that has been developed for

single-gene phylogeny (46). To evaluate ortholog/paralog relationships and overall dynamics of the

ligand complement, we used GeneRax (47). This new method uses maximum likelihood to reconcile

the gene tree with the species tree (47). In brief, given a gene and species tree, GeneRax uses a

maximum likelihood approach to optimise the duplication and loss events (47, 48).

Our analysis initially identified a few invertebrate putative chemokine ligands (Figure 1A),

however, these sequences lacked protein signatures associated with the canonical ligands (Figures

S3-5 and File S3), and they were therefore excluded from further analysis (see Supplementary

Results for further information). The phylogenetic tree for the canonical ligands identifies two major

groups, the CC-type, which also includes the XC- and X3C-types, and the CXC-type (TBE = 0.95, UFB =

92%) (Figure 1C and Figures S8,9), confirming the previous finding obtained using synteny data (49,

50). Next, to clarify the distribution of canonical chemokines, we first reconciled their gene tree with

the species tree and then used the reconciled tree to trace the presence/absence of each chemokine

group throughout all the species (Figure 2A and Figure S18). Our results confirm previous findings

that canonical chemokines are uniquely present in vertebrates (30, 49). Additionally, they indicate

that chemokines are not evenly distributed across vertebrates and can be different even between

closely related species (51). Some are very ancient, e.g., CXCL12 is present in lamprey; CXCL14 and

CCL20 are present in all jawed vertebrates; and CXCL8 is present throughout bony fishes and

tetrapods, with few exceptions, notably mice and rats. However, a large part of the chemokine

diversity evolved within mammals (e.g., CXCL1/2/3, CXCL16, and CCL25), particularly placentals (e.g.,

CXCL5/6 and CCL3/18). The phylogenetic relationships we uncovered in our reconciled tree were

mostly compatible with known syntenic relationships as described in human (7). For example, the

large cluster of CXC-type chemokine genes present in human chromosome 4 contains CXCL1-11 plus

CXCL13 (7), all of which coalesce within a monophyletic group in our tree (Figure 2A). The micro-

synteny within this cluster is also, to some extent, reflected in the phylogenetic relationships.

Similarly, the other large syntenic cluster of chemokines, located on human chromosome 17,

containing most of the CC-type chemokines (7), corresponds, with few exceptions, to a large

monophyletic clade in our tree (Figure 2A). CXCL16 which is on a nearby locus of chromosome 17, is

also phylogenetically related to this CC-type clade (Figure 2A). The complement of the canonical

chemokines undergoes the largest expansion at the base of jawed vertebrates, where there is an

expansion from 4 to 18 genes (Figure 2B). A second expansion occurred at the base of bony fishes

(i.e., Osteichthyes) followed by relative stability until placental mammals, where the total number of

canonical chemokine ligands jumped to 45 genes. Finally, unlike previous works (52) our results

support the presence of orthologs of both CC-type and CXC-type in the common ancestor of all

vertebrates (Figure 2A).

Differently from the canonical chemokines, we identified a *bona fide* TAFA, i.e., with specific

protein motifs, in the urochordates, the sister group to vertebrates (see Supplementary Results and

Figures S6-7). The phylogenetic trees (Supplementary Figures S10,11) identified monophyletic

groups for TAFA5 (TBE=0.98, UFB=98%), TAFA1 (TBE=0.94, UFB=98%), TAFA4 (TBE=0.77, UFB=75%)

and TAFA2/3 (TBE=0.65, UFB=84%). The reconciled tree from GeneRax places the root at the

urochordate sequence (Figure S19), therefore clarifying that the TAFA5 clade is the sister group to

TAFA1-4 (Figure 2A). The family originated in the ancestor of urochordates and vertebrates, and the

first duplications occurred at the base of vertebrates giving rise to the TAFA5 split followed by the

TAFA1 split. Subsequently, at the base of jawed vertebrates, additional duplications bring the

complements from 3 to 10 (Figure 2B), giving rise to the remaining groups so that all jawed

vertebrates possess the full diversity of TAFAs.

The phylogenetic trees for CYTL and CXCL17 mainly reflect the species trees (Figures S12-15),

and the reconciliations revealed very simple complement dynamics (Figure 2B and Figures S20,21).

However, these molecules show a remarkable difference in their distribution. CYTLs are present

throughout gnathostomes, while CXCL17 is found only in placental mammals (Figure 2A).

The phylogenetic analysis for the CKLF super family (Figure 1D and Figures S16,17) recovered

a monophyletic clade for the CKLF I group (TBE=0.96, UFB=80%) that we had already identified

through CLANS. This group contains CKLF, which is known to interact with C-C chemokine receptor 4

(10, 11), as well as CMTM1, 2, 3, 5, and proteolipid protein 2 (PLP2). Other monophyletic clades that

are consistent with the CLANS are CMTM4/6 (TBE=0.90, UFB=61%), CMTM7 (TBE=0.92, UFB=83%)

and a clade containing CMTM8 plus other related molecules such as plasmolipin (PLLP) and myelin

and lymphocyte proteins (MAL) (TBE=0.89, UFB=60%). The latter were all part of a large cluster that

we called CKLF II in the CLANS (Figure 1B). However, the placement of the root of the tree in Figure

1D can affect the interpretation of the relationships among CKLF II subgroups. To address this

problem and clarify the patterns of duplications and the presence/absence of each group

throughout animals, we used GeneRax to reconcile the gene with the species tree (see above and

Material and Methods for details). Our results suggest (Figure 2 and Figure S22) that most CKLFSF

groups, such as CMTM4,6 and 8, originate in the vertebrate stem group from pre-existing CMTM

genes and are widely distributed in animals. The CKLF I subgroups originate from duplications at the

base of jawed vertebrates, except for the split between CKLF and CMTM1 that occurs only within

mammals (Figure 2A). We observe the major two expansions of the CKLFSF genes in the stem group

of vertebrates (from 6 to 10 complements), and then in jawed vertebrates (from 10 to 16

complements). Interestingly, the extent of these expansions are less drastic than those we see for

canonical chemokines (Figure 2B). In total, we have identified that the five distinct ligand groups

have a different origin in the animal tree of life and underwent divergent evolutionary histories.

Canonical and non-canonical chemokine receptors are divided into four groups.

195   Next, we investigated the origin and pattern of duplication for the chemokine receptors and

chemokine-like receptors (Table 1). Using BLASTP against the 64 species, we identified 7,157

putative chemokine receptors (see Materials and Methods for more details) and investigated their

relationships using CLANS (see above for justification). The result (Figure S23C) identifies four main

groups of chemokine receptors and chemokine-like receptors. The first comprises canonical

receptors (i.e., CCR, CXCR, CX3CR1, CX3C, and XCR1), and the second includes atypical receptor 3

and GPR182, which has been recently shown to have chemokine receptor activity (53). The third

group, which we named Chemokine-like plus (CML-plus), contains the chemokine-like receptors

(CML1 also known as chemerin receptor 1), formyl peptide receptors (FPR) that bind the TAFA

ligands (15, 16) and other GPCRs such as GPR1 (chemerin receptor 2), GPR33, PTGDR2. Furthermore,

the CLANS analysis identifies an intermediate group containing angiotensin, apelin and other

receptors and shows sequence similarity to canonical and chemokine-like receptors (Figure S23B).

Finally, our analysis identifies a small cluster composed of only ACKR1 that do not connect to other

GPCRs or other atypical receptors even at loose p-value thresholds. This indicates that their

sequence is either non-homologous or highly divergent from other chemokine receptors and

atypical receptors. Overall, these groups are robust to the stringency threshold used (i.e., different

p-values) (Figure S23). Interestingly, no specific cluster of viral or viral-like receptors was identified,

but 6 of the reference viral receptor sequences clustered with the canonical chemokine receptors.

213   Altogether, these results confirm the homology between the canonical receptors and

atypical receptor 3/GPR182. However, these findings indicate that the other GPCRs, such as the

chemokine-like receptors, formyl peptide receptors, GPR1, and GPR33, are also closely related to the

canonical receptors. Remarkably, these results also indicate that ACKR1 is not homologous to the

canonical chemokine receptors. Furthermore, all clusters of chemokine receptors contained only

vertebrate sequences, except for the receptors of viral origin.

Canonical and chemokine-like receptors derive from single gene duplication in the ancestor of

vertebrates.

Previous studies suggested that the chemokine receptors evolved from a duplication of

angiotensin receptors (54) or adrenomedullin receptors (30, 55). However, these works were based

on error-prone phylogenetic methods such as Neighbour Joining (55). Our CLANS results indicate

that chemokine receptors and chemokine-like receptors have only been observed in vertebrates.

Therefore, we need to focus on invertebrate genomes to clarify the chemokine receptor's outgroup.

To clarify this, we lowered the p-value thresholds of CLANS (to p-value $< 1e^{-50}$) and collected a

combined dataset including all chemokine receptor sequences and outgroups (i.e., sequences that

connect to the chemokine receptor cluster), resulting in 3,026 sequences. We then performed a

phylogenetic tree on this dataset using maximum likelihood methods with UFB and TBE for

evaluating nodal support (see above and Materials and Methods for details).

Our combined phylogenetic analysis shows strong support for the monophyly of canonical

chemokine receptors (UFB=96, TBE=1.0), the CML-plus (UFB=95, TBE=0.99) and the atypical

3/GPR182 (UFB=100, TBE=1) (Figure 3, S24 and S25). In contrast, viral chemokine receptors are

paraphyletic, with three sequences placed within the canonical chemokine receptors and 3 forming

a monophyletic group sister to them (UBF=84 TBE=1.0). Our results also suggest that the

intermediate group, which includes apelin receptors, angiotensin receptors, bradykinin receptors,

and orphan GPCRs (e.g., GPR25; GPR15) forms a monophyletic clade with the canonical chemokine

receptors, CML-plus group and atypical3/GPR182 (UFB=61, TBE=0.91). However, its position changes

between the sister group to canonical chemokine receptors plus atypical3/GPR182 in the TBE tree

(TBE=0.84) and sister to CML plus in the ultrafast bootstrap tree (UFB=38).

All the groups mentioned above form a large clade composed of vertebrate-specific GPCRs

(UBF=100 TBE=0.96) that also includes other GPCRs, such as CLTR and P2RY receptors (Figure 3, S24

and S25). Another orphan GPCR, GPR35, had been proposed as a potential chemokine receptor (41);

however, this was later questioned (42, 43) and GPR35 is still generally considered orphan (56–58).

Our analysis collected GPR35 and placed it within this large vertebrate specific clade indicating that

is it also a vertebrate specific gene but not phylogenetically a 'canonical' chemokine receptor. The

closest outgroup to this clade is composed of a few sequences from urochordates, the sister group

of vertebrates (UFB=49 TBE=0.91) (Figure 3, S24 and S25). Interestingly, as the sister group of this

clade, we identify a group composed of Relaxin receptors, which contain sequences from both

urochordates and vertebrates (UBF=53 TBE=0.95). Finally, as the sister group of these large clades,

we identified a clade of cephalochordate-specific sequences (UBF=44).

To clarify the duplication pattern and origin of the chemokine receptors, we used GeneRax

(47) (see Materials and Methods). Our results indicate (Figure 4A, S26) that all chemokine receptors

(except ACKR1) originated from a duplication in the stem lineage of vertebrates. This duplication of

an unknown GPCR gave rise to the CML-plus, the canonical chemokine receptors and atypical

3/GPR182 groups as well as the intermediate group and other GPCRs (Figure 4A, S26). This result is

consistent with the distribution of the paralogous Relaxin receptors which are present both in

urochordates and vertebrates and the position of the orphan urochordate sequences as sister group

of canonical chemokine receptors, CML-plus group and atypical3/GPR182 and other GPCRs (see

above). Furthermore, the phylogenetic relationships amongst canonical chemokine receptors are

overall consistent with the syntenic gene patterns known in human (7). The largest cluster of

chemokine receptor genes spans 3 closely located loci on human chromosome 3 (7). It includes most

CCRs, XCR, and CX3CR and corresponds to one of the two major monophyletic clades in our tree

(Figure 4A). Another example is the mini cluster of CXCR1 and CXCR2, located on human

chromosome 2 (7), which we also found to form a monophyletic clade (Figure 4A).

We used the reconciliation to better understand the repertoires of receptors present at key

nodes during vertebrate evolution. Our results (Figure 4B) show a substantial difference in the

duplication pattern of different receptor families. For example, the complement of the

atypical3/GPR182 remains constant throughout vertebrate evolution while the canonical and

chemokine-like receptor groups expanded dramatically. The canonical chemokine receptors

expanded from 5 to 20 genes and the CML-plus from 1 to 11 in the ancestor of the jawed

vertebrates (Figure 4B). The expansion of the canonical CKRs is also not evenly distributed across its

subgroups, with the ancestral CC type receptors undergoing a series of duplications in jawed

vertebrates while the CXCR paralogs did not, specifically one (CXCR4) remains in single copy across

all vertebrates. We inferred that in the stem lineage of vertebrates, five canonical chemokine

receptor paralogs had already diverged, representing the two major types of receptors (2 CCR and 3

CXCR paralogs). Also, present in the stem lineage of vertebrates were ACKR3 and GPR182 as well as

a single copy gene which would later diverge to produce all the CML-plus clade.

DISCUSSION

281   This work substantially clarifies the evolutionary assembly of the chemokine system. Our

analysis shows that contrary to the receptors which evolved from a single duplication event in the

vertebrate stem group, several unrelated molecules acquired the ability to interact with chemokine

receptors over the course of evolutionary history. Furthermore, our results (summarized in Figure 5)

suggest that the key components of the chemokine system, including the chemokine receptors

themselves, evolved in the stem group of vertebrates in the Cambrian around 500 million years ago

and then underwent substantial diversification in the stem group of jawed vertebrates. These

findings shed new light on the complex evolutionary history of the chemokine system.

Unrelated molecules converged to chemokine function.

290   Based on the presence of shared protein motifs, TAFA "chemokines" (13, 14), CXCL17 (59,

60) and CYTL (19) have been proposed to be homologous to chemokine ligands. However, our

findings strongly suggest that these molecules are not homologous (Figure 1) and likely acquired the

ability to activate a chemokine-like response through convergent evolution. Our conclusions differ

from those previous studies (13, 19, 59, 60) due to the differences in data completeness and

methodological approach. Specifically, we used a complete set of canonical and non-canonical

ligands and assessed the homology using overall sequence similarity rather than single motifs. Our

results support and expand upon the findings of (29), which suggested that the presence of a CXC or

CC motif is necessary but not sufficient for a protein to be defined as a chemokine ligand. Similarly,

CKLF has been considered a "new member" of the chemokine family based on its function (12) we

argue that classification based solely on function is insufficient and can be misleading. Instead, we

recommend considering the evolutionary relationships among these molecules as the primary

criterion for classification.

Most of the canonical and non-canonical ligands are vertebrate innovations.

Our results clarify the distribution of canonical chemokine ligands in animals (Figure 2) and

confirm that they are present only in vertebrates (30). We identify orthologs of CXCL and CCL ligands

in both extant lineages of cyclostomes (Figure 2A). While chemokines have already been described in

lamprey (52, 61, 62), it is the first time, to our knowledge, that they are described also in hagfish.

Our findings also indicate that both CC and CXC types were present in the common ancestor of all

vertebrates and that few ancestral genes gave rise to the entire diversity of ligands that we know in

current animals. Furthermore, our results indicate that many chemokines, such as CXCL1-7, CXCL16,

as well as CCL25, CCL11/13, and CCL2/7, are uniquely present in mammals, suggesting that the

mammal ligand repertoire is substantially more complex than the one observed in other vertebrates.

Regarding non-canonical chemokine-like families, our findings indicate that the TAFA family

originated in the ancestor of vertebrates and urochordates; CYTL is a novelty of jawed vertebrates;

and CXCL17 is mammal-specific and likely unrelated to canonical chemokines (similar to its

controversial putative receptor, GPR35 (41–43), that is not a canonical chemokine receptor). The

CKLF superfamily has a more complex pattern with the presence of few groups in invertebrates and

then great expansions occurring at the base of vertebrates. The CKLFSF includes a monophyletic

clade (CKLF group I) comprising the original CKLF that binds CCR4, as well as CMTM1,2,3,5, derived

from duplications at the jawed vertebrates stem group. Interestingly, our analysis also revealed that

additional molecules not previously considered part of the CKLF super family are closely related to

classic members and should be included in it. For example, proteolipid protein 2 (PLP2) belongs to

the CKLF I group and is, therefore more closely related to the CKLF with chemokine function than

several other CKLFSF members. Similarly, CMTM8 is more closely related to plasmolipin (PLLP) and

myelin and lymphocyte protein (MAL) than to any of the classic CKLFSF members. Although this

relationship had been proposed based only on sequence similarity (34), our phylogenetic analysis

provides additional evidence for it. Therefore, the potential chemokine function of all these

additional members should be explored *in vitro* and *in vivo* in both vertebrates and invertebrates.

Most receptors derive from a single gene duplication.

Our results clarify the distribution of canonical chemokine receptors in vertebrates (Figure

4), and their evolutionary relationships and identify the pattern of duplication that leads to their

origin (Figure 4A, S26). Unlike previous works (63), we identify that atypical receptors do not form a

monophyletic group. Specifically, atypical 2 and 4 are part of the canonical clade specifically related

to CC-type receptor subclades. Furthermore, we find that the atypical 3 receptors are related to

GPR182, supporting previous functional data suggesting that the latter are atypical chemokine

receptors binding CXCL10, 12, and 13 (53). We attribute these differences to our use of wider GPCR

sampling and improved methods for phylogenetic inference.

Remarkably, our results do not identify ACKR1 as related to the main chemokine receptors

but rather as a divergent clade (Figure S23). To our knowledge, this is the first time this observation

has been made. Our current results do not allow us to clarify the evolutionary origin of ACKR1.

However, the presence of 7TMD domains suggests that they are GPCRs that independently acquired

the ability to bind chemokines. Alternatively, similarly to other genes evolved in the immune system,

ACKR1 may have been subjected to strong selective pressures that substantially changed their

sequence, obscuring their phylogenetic relationships. The case of ACKR1 being the most distantly

related receptor is intriguing as it is one of the most promiscuous chemokine receptors (2, 64) and it

has been shown to bind both CC and CXC chemokines (65, 66).

Viral chemokine receptors represent a cryptic group that can bind multiple chemokines (22,

23). Despite their functional similarity to canonical chemokine receptors, viral chemokine receptors'

evolutionary origin and distribution remain poorly understood. Our results indicate that viral GPCRs

do not form a monophyletic group, suggesting that the ability to encode chemokine-like receptors

has evolved independently in multiple viruses, including cytomegaloviruses and poxviruses. The

placement of viral sourced sequences within an otherwise vertebrate-specific clade supports the

hypothesis that viruses acquired these genes through non-vertical inheritance. Given the paraphyly

of viral receptors, this appears to have occurred multiple times. However, there are significant

uncertainties and further work is needed to untangle details of viral chemokine receptors' evolution.

Our analysis reveals that the clade comprising apelin receptors, angiotensin receptors,

bradykinin receptors, and orphan GPCRs (shown in Figure 3, 4 and S24–26) is closely related to

chemokine receptors. This finding partially supports previous studies (54) that suggested a gene

duplication event gave rise to both chemokine receptors and angiotensin receptors. Interestingly,

we found that single gene duplication in the vertebrate stem group led to the emergence of

canonical receptors and atypical 2,3,4, GPR182, chemokine-like receptors, formyl peptide receptors,

the intermediate group, and many other known and orphan GPCRs including the controversial

putative CXCL17 receptor GPR35. These findings suggest that two rounds of genome duplication (67,

68) played a role in the expansion of GPCR gene families. Future research will focus on investigating

the functions of the orphan genes and many-to-one orthologs discovered in urochordates. This will

provide further insight into the evolution and diversification of GPCR families in vertebrates.

The molecular assembly of the chemokine system

In this work, we explored the evolution of both ligand and receptor components of the

chemokine signaling system, including non-canonical molecules with either chemokine-like function

or sequence similarity. Our analysis suggests that the canonical chemokine signaling evolved in the

vertebrate stem group (about 500 Mya) likely due to the two rounds of genome duplication that

gave rise to many vertebrate novelties (67, 68). We found that the ancestral vertebrate repertoire

included orthologs of both major ligand groups (CXCL and CCL) and both CCR and CXCR receptors

and non-canonical components such as TAFA and CKLFSF ligands, and the receptors Atypical 3 and

GPR182 (Figure 5). The distribution of ligands and receptors in the ancestor of all vertebrates seems

to confirm the hypothesis that the ancestral function of chemokines was homeostatic (e.g., CXCL12,

CXCL14) with inflammatory functions arising from recent duplications (e.g., CXCL5, CXCL6),

potentially reflecting a rapid evolution induced by the selective pressure of new pathogens (7).

Chemokine ligand and receptor genes are known to cluster on specific chromosomes (7), consistent

with the hypothesis that they may be the result of the combination of *en bloc* duplication followed

by tandem duplications (30, 49, 50). Due to limited high-quality genomes, syntenic patterns of

chemokine genes described so far are based primarily on humans and a handful of other species (30,

49, 50), hampering our understanding of the level of conservation of these syntenic patterns.

Conversely, our large-scale phylogenetic analyses encompassed many species. We uncovered

several phylogenetic relationships that are consistent with known syntenic patterns in human,

providing stronger evidence for their evolutionary relationship.

The evolutionary history of canonical components includes several examples of known

ligand-receptor pairs following a corresponding pattern of origin and temporal dynamics of

duplications. This is true for example, for the ancient homeostatic CXCL12 ligand and its

corresponding receptors CXCR4 and ACKR3, that all originated in early vertebrates (7). The early

origin and conservation of CXCR4 and CXCL12 in the ancestor of vertebrates is interesting as this pair

plays a key role in the migration of neural crest cells (69) - a key vertebrate innovation (70). This

combined with the fact that homeostatic chemokine ligands/receptors tend to be restricted to

monogamous pairing (2, 51) suggests that homeostatic chemokine pairings are more ancient and

conserved being in single copy throughout much of the vertebrates. Contrastingly, inflammatory

chemokine pairings are more promiscuous, and this could be linked to the more recent duplications

in the genes, such as for CCL2/7/8/11/13 (Figure 2A) and their receptors CCR1/2/3/4/5 (Figure 4A).

For many of the non-canonical components, however, the ligand-receptor interactions are largely

unclear, and their pattern throughout vertebrate evolution remains to be explored. Overall, our

results indicate that three waves of molecular innovation in the vertebrates, jawed vertebrates,

bony fishes and mammal stem group increased the chemokine system's molecular complexity

(Figure 5), allowing for the fine-tuning present in modern-day animals.

MATERIALS AND METHODS

Data Mining and Dataset Assembly. We collected 64 proteomes from 25 vertebrates, six chordates

and 33 other animals covering the whole animal tree (Table S1). BUSCO v4.0.6 (71, 72) and the

metazoa_odb10 set of 954 genes were used to evaluate their completeness (Table S1).

To identify potential homologs of canonical chemokines, TAFA chemokines and CYTL1, we

used 207 curated sequences that we obtained from SwissProt (73, 74) as seeds for an initial BLASTP

(36, 38) with e-value < e^{-10}. To identify putative chemokines in cyclostomes, the lamprey *Petromyzon*

marinus (75) and the hagfish *Eptatretus burger* (76), we loosened the e-value to 0.05. Where

putative chemokine sequences were found for one cyclostome species but not the other, those

sequences were used to search again the other species. Furthermore, to investigate the presence of

ligands outside vertebrates, we performed an additional BLASTP on invertebrate proteomes with an

even looser e-value (0.1) and collected only up to five hits. This provided 18 initial candidate

homologs spanning multiple invertebrate phyla. Further characterisation of these invertebrate

sequences, through BLASTP versus SwissProt, protein domains search with InterProScan (77, 78),

position in CLANS analysis (see below) and, where necessary, multiple sequence alignments, led us

to retain only one urochordate sequence as a putative TAFA homolog (see Supplementary Results

for details).

To identify homologs for the CKLF superfamily, we used 21 SwissProt-reviewed sequences.

In addition to the BLASTP search, we used a position-specific iterative BLAST (PSI-BLAST) (37) with an

e-value threshold of <e^{-10}. Using this approach, we identified a total of 590 putative homologs,

including 186 from invertebrates.

We used BLASTP using 178 manually annotated receptor sequences from SwissProt as query

sequences for the chemokine receptors. This includes all human canonical and atypical chemokine

receptors (79). We also collected 8 viral sequences with chemokine receptor activity from UniProt

(80) and performed a second BLASTP. We extracted all BLAST hits with e-values $< e^{-10}$ and used

Phobius (81) to predict their transmembrane domain structure. Only sequences with 5-8

transmembrane domains were kept. Hit sequences were annotated by their top 5 BLAST hits against

SwissProt. All hits from both BLASTs were merged and filtered by cd-hit (82, 83) to remove

redundant sequences at the 95% similarity threshold. This resulted in 7,157 putative chemokine

GPCR sequences.

Identification of subgroups with Cluster Analysis of Sequences (CLANS). We utilised CLANS (39, 40)

with default parameters and different p-values (i.e., stringency values) to visualize the relationships

between subgroups of ligands and receptors. We assessed the similarity and interrelationships

between different clusters by gradually relaxing the p-value threshold (Figures S1, S2 and S23).

Additionally, we annotated each cluster using gene annotations for key species *Homo, Mus, Gorilla,*

Gallus, Anolis and *Danio*. In the case of the receptors, to improve the cluster annotation all human

Class-A GPCRs (excluding olfactory receptors) from GPCRdb (84) were added to the dataset as well

as the 8 seed viral chemokine receptors from UniProt (80).

Alignment and phylogenetic analysis

Alignment. All ligand and receptor sequences were aligned using MAFFT (85, 86) with the --auto

setting and using trimAl (87) to remove positions with >70% gaps.

Gene Trees. All gene alignments were analysed using IQTREE2 (88), the model test algorithm (89)

was used to select the best substitution model for each analysis. Best models selected by IQTREE2

for each set are listed in Table S2 (for receptors we manually selected GTR20+F+G4 as the model as

it was a large dataset). Nodal support was estimated using 1,000 ultrafast bootstrap (44, 45)

replicates. All analyses were repeated to run 100 non-parametric bootstrap repeats to calculate

nodal support with transferable bootstrap expectation: which is specifically designed to account for

phylogenetic instability (46).

For the receptors, due to the high computational burden of running TBE analyses on

sequence-dense datasets, we first analysed the full set of 3,026 sequences connected in CLANS at a

p-value of < 1 e^{-50} using UFB (Figure S25). Then, we extracted the chordate-specific clade sequences,

including all chemokine receptor groups and their immediate outgroups, to analyse using TBE.

Gene tree species tree reconciliation. To understand the pattern of duplication and the evolution of

gene complement we used GeneRax (47). GeneRax requires a gene tree that was obtained as

described above and a species tree that we constructed manually using publicly available

information. In the instances where the genes tree contained polytomies, we used ETE3 (90) to solve

them. The undated DL mode and the closest approximation of the best-fitting substitution model

were used for each alignment. To track the evolution of sub-lineages within each group, we used

annotated sequences of key species (e.g., *Homo sapiens* and *Mus musculus*) as reference. For the

receptors, we used only the chordate-specific clade subtree and sequences due to the

computational burden of running GeneRax on a high number of sequences. For species tree-gene

tree reconciliation, we treat the viral sequences as human sequences.

DATA AVAILABILITY

Supplementary material and raw output files for all the analyses described in this paper are available

in the GitHub page: Roberto-Feuda-Lab/Chemokine2023 (github.com).

ACKNOWLEDGEMENTS

This work is supported by a University Research Fellowship (UF160226) to RF. AA is supported by a

Research Grant from the Royal Society to RF (RGF\R1\181012). MG is supported by a PhD

Scholarship from the University of Leicester. CL is supported by a BBRSC MIBPT fellowship. This

research used the ALICE High-Performance Computing Facility at the University of Leicester.

REFERENCES

1.   K. Zhang, S. Shi, W. Han, Research progress in cytokines with chemokine-like
function. *Cellular & Molecular Immunology* **15**, 660–662 (2018).
2.   K. Chen, *et al.*, Chemokines in homeostasis and diseases. *Cell Mol Immunol* **15**, 324–
334 (2018).
3.   X. Blanchet, M. Langer, C. Weber, R. Koenen, P. von Hundelshausen, Touch of
Chemokines. *Frontiers in Immunology* **3**, 175 (2012).
4.   P. López-Cotarelo, C. Gómez-Moreira, O. Criado-García, L. Sánchez, J. L.
Rodríguez-Fernández, Beyond Chemoattraction: Multifunctionality of Chemokine Receptors
in Leukocytes. *Trends in Immunology* **38**, 927–941 (2017).
5.   P. B. Tran, R. J. Miller, Chemokine receptors: signposts to brain development and
disease. *Nature Reviews Neuroscience* **4**, 444–455 (2003).
6.   B. Moser, M. Wolf, A. Walz, P. Loetscher, Chemokines: multiple levels of leukocyte
migration control. *Trends in Immunology* **25**, 75–84 (2004).
7.   A. Zlotnik, O. Yoshie, The Chemokine Superfamily Revisited. *Immunity* **36**, 705–716
(2012).
8.   A. Zlotnik, O. Yoshie, Chemokines: A New Classification System and Their Role in
Immunity. *Immunity* **12**, 121–127 (2000).
9.   H. Nomiyama, N. Osada, O. Yoshie, A family tree of vertebrate chemokine receptors
for a unified nomenclature. *Developmental & Comparative Immunology* **35**, 705–715 (2011).
10.   Y. Wang, *et al.*, Chemokine-like factor 1 is a functional ligand for CC chemokine
receptor 4 (CCR4). *Life Sciences* **78**, 614–621 (2006).
11.   Y. Wang, *et al.*, Two C-terminal peptides of human CKLF1 interact with the
chemokine receptor CCR4. *The International Journal of Biochemistry & Cell Biology* **40**,
909–919 (2008).
12.   D.-D. Liu, *et al.*, Progress in pharmacological research of chemokine like factor 1
(CKLF1). *Cytokine* **102**, 41–50 (2018).
13.   Y. Tom Tang, *et al.*, TAFA: a novel secreted family with conserved cysteine residues
and restricted expression in the brain. *Genomics* **83**, 727–734 (2004).
14.   D. C. Sarver, X. Lei, G. W. Wong, FAM19A (TAFA): An Emerging Family of
Neurokines with Diverse Functions in the Central and Peripheral Nervous System. *ACS*
Chem. Neurosci. **12**, 945–958 (2021).
15.   W. Wang, *et al.*, FAM19A4 is a novel cytokine ligand of formyl peptide receptor 1
(FPR1) and is able to promote the migration and phagocytosis of macrophages. *Cell Mol*
Immunol **12**, 615–624 (2015).
16.   M. Y. Park, *et al.*, FAM19A5, a brain-specific chemokine, inhibits RANKL-induced
osteoclast formation through formyl peptide receptor 2. *Sci Rep* **7**, 15575 (2017).

17.   C. Zheng, *et al.*, FAM19A1 is a new ligand for GPR1 that modulates neural stem-cell
proliferation and differentiation. *The FASEB Journal* **32**, 5874–5890 (2018).
18.   X. Wang, *et al.*, Cytokine-like 1 Chemoattracts Monocytes/Macrophages via CCR2.
The Journal of Immunology **196**, 4090–4099 (2016).
19.   A. Tomczak, M. T. Pisabarro, Identification of CCR2-binding features in Cytl1 by a
CCL2-like chemokine model. *Proteins: Structure, Function, and Bioinformatics* **79**, 1277–
1292 (2011).
20.   T. Yoshimura, J. J. Oppenheim, Chemokine-like receptor 1 (CMKLR1) and
chemokine (C–C motif) receptor-like 2 (CCRL2); Two multifunctional receptors with
unusual properties. *Experimental Cell Research* **317**, 674–684 (2011).
21.   R. Bonecchi, G. J. Graham, Atypical Chemokine Receptors and Their Roles in the
Resolution of the Inflammatory Response. *Frontiers in Immunology* **7**, 224 (2016).
22.   T. N. Kledal, M. M. Rosenkilde, T. W. Schwartz, Selective recognition of the
membrane-bound CX3C chemokine, fractalkine, by the human cytomegalovirus-encoded
broad-spectrum receptor US28. *FEBS Letters* **441**, 209–214 (1998).
23.   T. F. Miles, *et al.*, Viral GPCR US28 can signal in response to chemokine agonists of
nearly unlimited structural degeneracy. *eLife* **7**, e35850 (2018).
24.   M. M. Rosenkilde, M. J. Smit, M. Waldhoer, Structure, function and physiological
consequences of virally encoded chemokine seven transmembrane receptors. *British Journal
of Pharmacology* **153**, S154–S166 (2008).
25.   H. Daiyasu, W. Nemoto, H. Toh, Evolutionary Analysis of Functional Divergence
among Chemokine Receptors, Decoy Receptors, and Viral Receptors. *Frontiers in
Microbiology* **3** (2012).
26.   M. Meyrath, *et al.*, The atypical chemokine receptor ACKR3/CXCR7 is a broad-
spectrum scavenger for opioid peptides. *Nature Communications* **11**, 3033 (2020).
27.   R. J. B. Nibbs, G. J. Graham, Immune regulation by atypical chemokine receptors.
Nat Rev Immunol **13**, 815–829 (2013).
28.   F. Bachelerie, *et al.*, New nomenclature for atypical chemokine receptors. *Nat
Immunol* **15**, 207–208 (2014).
29.   S. S. Denisov, CXCL17: The Black Sheep in the Chemokine Flock. *Frontiers in
Immunology* **12**, 2811 (2021).
30.   M. E. DeVries, *et al.*, Defining the Origins and Evolution of the
Chemokine/Chemokine Receptor System. *The Journal of Immunology* **176**, 401 (2006).
31.   B. Bajoghli, Evolution and function of chemokine receptors in the immune system of
lower vertebrates. *European Journal of Immunology* **43**, 1686–1692 (2013).
32.   H. Nomiyama, *et al.*, Extensive expansion and diversification of the chemokine gene
family in zebrafish: Identification of a novel chemokine subfamily CX. *BMC Genomics* **9**,
222 (2008).
33.   J. F. Fleming, R. Feuda, N. W. Roberts, D. Pisani, A Novel Approach to Investigate
the Effect of Tree Reconstruction Artifacts in Single-Gene Analysis Clarifies Opsin
Evolution in Nonbilaterian Metazoans. *Genome Biology and Evolution* **12**, 3906–3916
(2020).

34.   W. Han, *et al.*, Identification of eight genes encoding chemokine-like factor
superfamily members 1–8 (CKLFSF1–8) by in silico cloning and experimental validation.
Genomics **81**, 609–617 (2003).
35.   H.-J. Duan, X.-Y. Li, C. Liu, X.-L. Deng, Chemokine-like factor-like MARVEL
transmembrane domain-containing family in autoimmune diseases. *Chinese Medical Journal*
133 (2020).
36.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
search tool. *Journal of Molecular Biology* **215**, 403–410 (1990).
37.   S. F. Altschul, *et al.*, Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Research* **25**, 3389–3402 (1997).
38.   C. Camacho, *et al.*, BLAST+: architecture and applications. *BMC Bioinformatics* **10**,
421 (2009).
39.   T. Frickey, A. Lupas, CLANS: a Java application for visualizing protein families
based on pairwise similarity. *Bioinformatics* **20**, 3702–3704 (2004).
40.   F. Gabler, *et al.*, Protein Sequence Analysis Using the MPI Bioinformatics Toolkit.
Current Protocols in Bioinformatics **72**, e108 (2020).
41.   J. L. Maravillas-Montero, *et al.*, Cutting Edge: GPR35/CXCR8 Is the Receptor of the
Mucosal Chemokine CXCL17. *The Journal of Immunology* **194**, 29–33 (2015).
42.   S.-J. Park, S.-J. Lee, S.-Y. Nam, D.-S. Im, GPR35 mediates lodoxamide-induced
migration inhibitory response but not CXCL17-induced migration stimulatory response in
THP-1 cells; is GPR35 a receptor for CXCL17? *British Journal of Pharmacology* **175**, 154–
161 (2018).
43.   N. A. S. B. M. Amir, *et al.*, Evidence for the Existence of a CXCL17 Receptor
Distinct from GPR35. *The Journal of Immunology* **201**, 714–724 (2018).
44.   B. Q. Minh, M. A. T. Nguyen, A. von Haeseler, Ultrafast Approximation for
Phylogenetic Bootstrap. *Molecular Biology and Evolution* **30**, 1188–1195 (2013).
45.   D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution* **35**, 518–
522 (2018).
46.   F. Lemoine, *et al.*, Renewing Felsenstein's phylogenetic bootstrap in the era of big
data. *Nature* **556**, 452–456 (2018).
47.   B. Morel, A. M. Kozlov, A. Stamatakis, G. J. Szöllősi, GeneRax: A Tool for Species-
Tree-Aware Maximum Likelihood-Based Gene Family Tree Inference under Gene
Duplication, Transfer, and Loss. *Molecular Biology and Evolution* **37**, 2763–2774 (2020).
48.   T. A. Williams, *et al.*, The power and limitations of species tree-aware phylogenetics.
2023.03.17.533068 (2023).
49.   H. Nomiyama, N. Osada, O. Yoshie, Systematic classification of vertebrate
chemokines based on conserved synteny and evolutionary history. *Genes to Cells* **18**, 1–16
(2013).
50.   A. Zlotnik, O. Yoshie, H. Nomiyama, The chemokine and chemokine receptor
superfamilies and their molecular evolution. *Genome Biol* **7**, 243–243 (2006).
51.   P. M. Murphy, "15 - Chemokines and Chemokine Receptors" in *Clinical Immunology*
(Sixth Edition), R. R. Rich, *et al.*, Eds. (Elsevier, 2023), pp. 215–227.

52. Z. Sun, *et al.*, The evolution and functional characterization of CXC chemokines and
receptors in lamprey. *Developmental & Comparative Immunology* **116**, 103905 (2021).
53. A. Le Mercier, *et al.*, GPR182 is an endothelium-specific atypical chemokine receptor
that maintains hematopoietic stem cell homeostasis. *Proceedings of the National Academy of*
Sciences **118**, e2021596118 (2021).
54. P. Liò, M. Vannucci, Investigating the evolution and structure of chemokine
receptors. *Gene* **317**, 29–37 (2003).
55. R. Fredriksson, M. C. Lagerström, L.-G. Lundin, H. B. Schiöth, The G-Protein-
Coupled Receptors in the Human Genome Form Five Main Families. Phylogenetic Analysis,
Paralogon Groups, and Fingerprints. *Mol Pharmacol* **63**, 1256–1272 (2003).
56. S. Xiao, W. Xie, L. Zhou, Mucosal chemokine CXCL17: What is known and not
known. *Scandinavian Journal of Immunology* **93**, e12965 (2021).
57. S. P. Giblin, J. E. Pease, What defines a chemokine? – The curious case of CXCL17.
Cytokine **168**, 156224 (2023).
58. J. Duan, et al., Insights into divalent cation regulation and G13-coupling of orphan
receptor GPR35. *Cell Discov* **8**, 1–12 (2022).
59. M. T. Pisabarro, *et al.*, Cutting Edge: Novel Human Dendritic Cell- and Monocyte-
Attracting Chemokine-Like Protein Identified by Fold Recognition Methods. *The Journal of*
Immunology **176**, 2069–2073 (2006).
60. E. J. Weinstein, *et al.*, VCC-1, a novel chemokine, promotes tumor growth.
Biochemical and Biophysical Research Communications **350**, 74–81 (2006).
61. A. M. Najakshin, L. V. Mechetina, B. Y. Alabyev, A. V. Taranin, Identification of an
IL-8 homolog in lamprey (Lampetra fluviatilis): early evolutionary divergence of
chemokines. *European Journal of Immunology* **29**, 375–382 (1999).
62. B. Bajoghli, *et al.*, Evolution of Genetic Networks Underlying the Emergence of
Thymopoiesis in Vertebrates. *Cell* **138**, 186–197 (2009).
63. L. Pan, J. Lv, Z. Zhang, Y. Zhang, Adaptation and Constraint in the Atypical
Chemokine Receptor Family in Mammals. *BioMed Research International* **2018**, 9065181
(2018).
64. S. J. Allen, S. E. Crown, T. M. Handel, Chemokine:Receptor Structure, Interactions,
and Antagonism. *Annual Review of Immunology* **25**, 787–820 (2007).
65. R. Horuk, *et al.*, A Receptor for the Malarial Parasite Plasmodium vivax: the
Erythrocyte Chemokine Receptor. *Science* **261**, 1182–1184 (1993).
66. R. Horuk, The Duffy Antigen Receptor for Chemokines DARC/ACKR1. *Frontiers in*
Immunology **6** (2015).
67. M. Kasahara, The 2R hypothesis: an update. *Current Opinion in Immunology* **19**,
547–552 (2007).
68. O. Simakov, *et al.*, Deeply conserved synteny resolves early events in vertebrate
evolution. *Nat Ecol Evol* **4**, 820–830 (2020).
69. W. Tang, Y. Li, A. Li, M. E. Bronner, Clonal analysis and dynamic imaging identify
multipotency of individual Gallus gallus caudal hindbrain neural crest cells toward cardiac
and enteric fates. *Nat Commun* **12**, 1894 (2021).
70. J. R. York, D. W. McCauley, The origin and evolution of vertebrate neural crest cells.
Open Biology **10**, 190285 (2020).

71.   R. M. Waterhouse, *et al.*, BUSCO Applications from Quality Assessments to Gene
Prediction and Phylogenomics. *Mol Biol Evol* **35**, 543–548 (2018).
72.   M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and*
Evolution **38**, 4647–4654 (2021).
73.   E. Boutet, *et al.*, "UniProtKB/Swiss-Prot, the Manually Annotated Section of the
UniProt KnowledgeBase: How to Use the Entry View" in *Plant Bioinformatics: Methods and*
Protocols, Methods in Molecular Biology., D. Edwards, Ed. (Springer, 2016), pp. 23–54.
74.   S. Poux, *et al.*, On expert curation and scalability: UniProtKB/Swiss-Prot as a case
study. *Bioinformatics* **33**, 3454–3460 (2017).
75.   J. J. Smith, *et al.*, Sequencing of the sea lamprey (Petromyzon marinus) genome
provides insights into vertebrate evolution. *Nat Genet* **45**, 415–421 (2013).
76.   D. Yu, *et al.*, Hagfish genome illuminates vertebrate whole genome duplications and
their evolutionary consequences. 2023.04.08.536076 (2023).
77.   E. M. Zdobnov, R. Apweiler, InterProScan – an integration platform for the signature-
recognition methods in InterPro. *Bioinformatics* **17**, 847–848 (2001).
78.   P. Jones, *et al.*, InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**, 1236–1240 (2014).
79.   F. Bachelerie, *et al.*, Chemokine receptors (version 2020.5) in the IUPHAR/BPS
Guide to Pharmacology Database. *IUPHAR/BPS Guide to Pharmacology CITE* **2020** (2020).
80.   The UniProt Consortium, UniProt: the Universal Protein Knowledgebase in 2023.
Nucleic Acids Research **51**, D523–D531 (2023).
81.   L. Käll, A. Krogh, E. L. L. Sonnhammer, Advantages of combined transmembrane
topology and signal peptide prediction—the Phobius web server. *Nucleic Acids Research* **35**,
W429–W432 (2007).
82.   W. Li, L. Jaroszewski, A. Godzik, Clustering of highly homologous sequences to
reduce the size of large protein databases. *Bioinformatics* **17**, 282–283 (2001).
83.   L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).
84.   G. Pándy-Szekeres, *et al.*, GPCRdb in 2023: state-specific structure models using
AlphaFold2 and new ligand resources. *Nucleic Acids Research* **51**, D395–D402 (2023).
85.   K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**,
3059–3066 (2002).
86.   K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772–
780 (2013).
87.   S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973
(2009).
88.   B. Q. Minh, *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534 (2020).

89.   S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin,
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**, 587–
589 (2017).
90.   J. Huerta-Cepas, F. Serra, P. Bork, ETE 3: Reconstruction, Analysis, and
Visualization of Phylogenomic Data. *Molecular Biology and Evolution* **33**, 1635–1638
(2016).
91.   W. Han, *et al.*, Molecular cloning and characterization of chemokine-like factor 1
(CKLF1), a novel human cytokine with unique structure and potential chemotactic activity.
Biochem J **357**, 127–135 (2001).
92.   L. Wang, *et al.*, Molecular cloning and characterization of chemokine-like factor
super family member 1 (CKLFSF1), a novel human gene with at least 23 alternative splicing
isoforms in testis tissue. *The International Journal of Biochemistry & Cell Biology* **36**, 1492–
1501 (2004).
93.   C. Jin, P. Ding, Y. Wang, D. Ma, Regulation of EGF receptor signaling by the
MARVEL domain-containing protein CKLFSF8. *FEBS Letters* **579**, 6375–6382 (2005).
94.   Z.-Z. Wang, *et al.*, Chemokine-like factor 1, a novel cytokine, induces nerve cell
migration through the non-extracellular Ca2+-dependent tyrosine kinases pathway. *Brain
Research* **1308**, 24–34 (2010).
95.   T. Li, *et al.*, Expression of chemokine-like factor 1 is upregulated during T
lymphocyte activation. *Life Sciences* **79**, 519–524 (2006).
96.   Y. Zhang, *et al.*, C-terminal peptides of chemokine-like factor 1 signal through
chemokine receptor CCR4 to cross-desensitize the CXCR4. *Biochemical and Biophysical
Research Communications* **409**, 356–361 (2011).
97.   H. Li, *et al.*, A novel 3p22.3 gene CMTM7 represses oncogenic EGFR signaling and
inhibits cancer cell growth. *Oncogene* (2014) https:/doi.org/10.1038/onc.2013.282.
98.   S. Zhu, *et al.*, Protein Cytl1: its role in chondrogenesis, cartilage homeostasis, and
disease. *Cell. Mol. Life Sci.* **76**, 3515–3523 (2019).
99.   H. Xue, *et al.*, CYTL1 Promotes the Activation of Neutrophils in a Sepsis Model.
Inflammation **43**, 274–285 (2020).
100.   X. Wang, *et al.*, Tafa-2 plays an essential role in neuronal survival and
neurobiological function in mice. *Acta Biochimica et Biophysica Sinica* **50**, 984–995 (2018).
101.   Y. Wang, *et al.*, Novel Adipokine, FAM19A5, Inhibits Neointima Formation After
Injury Through Sphingosine-1-Phosphate Receptor 2. *Circulation* **138**, 48–63 (2018).
102.   J. Okada, *et al.*, Analysis of FAM19A2/TAFA-2 function. *Physiology & Behavior*
208, 112581 (2019).
103.   B. L. Lokeshwar, G. Kallifatidis, J. J. Hoy, "Chapter One - Atypical chemokine
receptors in tumor cell growth and metastasis" in *Advances in Cancer Research*, GPCR
Signaling in Cancer., A. K. Shukla, Ed. (Academic Press, 2020), pp. 1–27.
104.   H.-Q. He, R. D. Ye, The Formyl Peptide Receptors: Diversity of Ligands and
Mechanism for Recognition. *Molecules* **22**, 455 (2017).
105.   M. Dohrmann, G. Wörheide, Dating early animal evolution using phylogenomic data.
Sci Rep **7**, 3599 (2017).
106.   F. Delsuc, *et al.*, A phylogenomic framework and timescale for comparative studies of
tunicates. *BMC Biol* **16**, 39 (2018).

107.   F. M. Gradstein, J. G. Ogg, "Chapter 2 - The Chronostratigraphic Scale" in *The*
Geologic Time Scale, F. M. Gradstein, J. G. Ogg, M. D. Schmitz, G. M. Ogg, Eds. (Elsevier,
2012), pp. 31–42.
736

	Names	Abbreviations	*H. sapiens* Orthologs	Functions	References
Ligand Groups	**Canonical Chemokines**	CCL, CXCL, XCL, CX3CL	CCL1-3, 3L1, 3L3, 4, 4L1-L2, 5, 7, 8, 11, 13, 14-28; CXCL1-4, 4L1, 5-14, 16,17; XCL1,2; CX3CL1	- Chemokine receptor binding and signalling - Chemoattraction of leukocytes - Homeostasis of leukocytes	(2, 4, 7)
	CKLF-Like MARVEL Transmembrane Domain-Containing Proteins (Chemokine-Like Factor Super Family)	CKLF, CMTM	CKLF1; CMTM1-8 (CKLF, CKLFSF1-8)	- CKLF1 (CKLF) binds to chemokine receptor CCR4 - CKLF1 (CKLF): chemotactic activity for lymphocytes, macrophages, and neutrophils - Other CMTMs: variably expressed in immune system; putative roles in immunity, programmed cell death, regulation of anti-tumour immunity etc.	(1, 10–12, 34, 35, 91–97)
	Cytokine-Like Protein 1 (Protein C17 or C4orf4)	CYTL	CYTL1	- Chemokine receptor binding (CCR2) and signalling - Chemoattraction monocytes/macrophages - Chemotactic activity in neutrophils	(1, 18, 98, 99)
	TAFA Chemokines (Family with sequence similarity 19 (chemokine (C-C motif)-like) member A)	TAFA	TAFA1-5 (FAM19A1-5)	- Formyl-peptide receptor binding and signalling (TAFA4 and 5) - Putative binding to other GPCRs: GPR1 (TAFA1); S1PR2 (TAFA5) - Expressed in central and peripheral nervous system - Implicated in vast diversity of physiological processes	(1, 13–17, 100–102)
Receptor Groups	**Canonical Chemokine Receptors**	CCR, CXCR, XCR, CX3CR	CCR1-10; CXCR1-6; XCR1; CX3CR1	- Chemokine binding and signalling - Chemotaxis of leukocytes - Homeostasis of leukocytes	(2, 4, 7)
	Atypical Chemokine Receptors	ACKR	ACKR1-4 (DARC; D6; CXCR7; CCRL1)	- Chemokine binding, but no signalling - Resolution of inflammatory response	(21, 27, 28)
	Chemokine Receptor-Like (Chemokine (C-C motif) receptor-like 2)	CCRL	CCRL2 (ACKR5)	- Binds CCL5 and CCL19, but no signalling - Binds chemerin and presents it to CMKLR1	(20, 103)
	Chemokine-Like Receptor 1	CML	CML1 (CMKLR1; ChemR23)	- Binds chemerin inducing migration of macrophages and dendritic cells - Binds also other anti-inflammatory molecules (e.g., Resolvin E1 (RvE1))	(20)
	Formyl-peptide Receptors	FPR	FPR 1-3	- TAFA chemokine binding - Chemoattraction, modulation of inflammation	(15, 16, 104)
	Putative Chemokine Receptors	ACKR6, CXCR8	PTITMP3, CXCR8 (GPR35)	- ACKR6/PTITMP3: Binds CCL18 (NB: It is not a GPCR) - CXCR8/GPR35: binds CXCL17	(41, 103)

Table 1: Summary table of all the canonical and non-canonical chemokine components analyzed in

this study.

Figure Legends

Figure 1: Cluster Analysis and Phylogeny of Ligand groups. A) Similarity-based clustering, using

CLANS, of canonical chemokines and related molecules with sequence similarity. Canonical

chemokines are an independent group from other related molecules (TAFA, CYTL and CXCL17).

Canonical chemokines are composed of two large groups (CC-type and CXC-type) within which some

divergent subgroups are highlighted. The clustering and connections shown are at the p-value

threshold of 1E-6. Other p-values tested are shown in Supplementary Figure S1. Candidate

invertebrate sequences are shown as crosses and further information regarding them can be found

in Supplementary Results. **B)** Similarity-based clustering, using CLANS, of the CKLF super family

(CKLFSF). Two major clusters are formed: the smaller "CKLF Group I" and the heterogenous "CKLF

group II" that also includes some invertebrate sequences (shown as crosses). Subclades, including

the known members of the CKLF super family, are highlighted. The clustering and connections

shown are at the p-value threshold of 1E-15, as this is the threshold at which the two major clusters

connect. Other p-values tested are shown in Supplementary Figure S2. **C)** Maximum-Likelihood un-

rooted phylogenetic tree of canonical chemokines. CC-type and CXC-type are split into two separate

clades. Supports for key nodes are indicated in boxes with Transferable Bootstrap Expectation (TBE)

represented by triangles and the Ultrafast Bootstraps (UFB) as circles. A traffic light colour code is

used to indicate the level of support: high (green); intermediate (yellow) and low (red). **D)**

Maximum-Likelihood un-rooted phylogenetic tree of the CKLF super family (CKLFSF). The CKLF group

I is monophyletic, while the CKLF group II is not. Supports for key nodes are indicated in boxes with

Transferable Bootstrap Expectation (TBE) represented by triangles and the Ultrafast Bootstraps

(UFB) as circles. A traffic light colour code is used to indicate the level of support: high (green);

intermediate (yellow) and low (red).

Figure 2: Distribution and duplication patterns of ligand groups. A) Presence of all ligand groups are

mapped onto a species tree. Gene trees and duplication events are based on the gene tree to

species tree reconciliation analyses. The nomenclature for canonical chemokines is primarily based

on known chemokines of human (or mouse). Where human and mouse chemokines do not

correspond, the default name refers to the human gene and the mouse (*Mus musculus*) one is

indicated with "Mm". Chemokines that have been classically described as having either homeostatic

or inflammatory function are indicated with a circle or a star respectively. The classification used

here was based on Zlotnik and Yoshie 2012 (7) with the inflammatory type also including

chemokines they described as plasma/platelet types. Overall, canonical chemokines originated in

vertebrates and expanded a first time in jawed vertebrates and a second time in mammals.

Homeostatic chemokines (e.g., CXCL12) are generally more ancient than inflammatory ones. CXCL17

and CYTL are mammal and jawed vertebrate specific respectively. TAFA originated in the common

ancestor of vertebrates and urochordates, while the CKLF super family is present in invertebrates

although key duplications occurred at the base of vertebrates. **B)** Number of complements for each

ligand group at key species nodes are mapped onto the species tree. The number of complements in

each group reflects the pattern of duplications. The major increase occurred at the level of jawed

vertebrates with canonical chemokines undergoing a second significant increase within placentals.

Silhouette images are by Andreas Hejnol (*Xenopus laevis*); Andy Wilson (*Anas platyrhynchos*,

Taeniopygia guttata); Carlos Cano-Barbacil (*Salmo trutta*); Christoph Schomburg (*Anolis carolinensis*,

Ciona intestinalis, *Eptatretus burgeri*, *Petromyzon marinus*); Christopher Kenaley (*Mola mola*);

Chuanixn Yu (*Latimeria chalumnae*); Daniel Jaron (*Mus musculus*); Daniel Stadtmauer (*Monodelphis*

domestica); Fernando Carezzano (Asteroidea); Ingo Braasch (*Callorhinchus milii*); Jake Warner (*Danio*

rerio); Kamil S. Jaron (*Poecilia formosa*); Mali'o Kodis, photograph by Hans Hillewaert

(*Branchiostoma lanceolatum*, https://www.phylopic.org/images/719d7b41-cedc-4c97-9ffe-

dd8809f85553/branchiostoma-lanceolatum); Margot Michaud (*Canis lupus*, *Physeter*

macrocephalus); NASA (*Homo sapiens sapiens*); Nathan Hermann (*Scophthalmus aquosus*); Ryan

Cupo (*Rattus norvegicus*); seung9park (*Takifugu rubripes rubripes*); Soledad Miranda-Rottmann

(*Pelodiscus sinensis*, https://www.phylopic.org/images/929fd134-bbd7-4744-987f-

1975107029f5/pelodiscus-sinensis); Steven Traver (*Gallus gallus domesticus*, *Ornithorhynchus*

anatinus); Stuart Humphries (*Thunnus thynnus*); T. Michael Keesey (after Colin M. L. Burnett) (*Gorilla*

gorilla gorilla); Thomas Hegna (based on picture by Nicolas Gompel) (*Drosophila (Drosophila)*

mojavensis); and Yan Wong (*Balanoglossus*).

Figure 3: Phylogeny of Receptor groups.

An unrooted maximum likelihood phylogeny of chemokine receptors. The tree shown is the transfer

bootstrap expectation (TBE) tree including just the chordate specific clade from the ultrafast

bootstrap tree (UFB). Node supports from both TBE (triangle) and UFB (circle) shown for equivalent

key nodes in boxes with arrows to indicate node. A traffic light colour code is used to indicate the

level of support: high (green); intermediate (yellow) and low (red). Key clades highlighted: yellow =

chemokine like plus group (CMLplus); blue = intermediate group; green = atypical 3 and GPR182

(ACKR3/GPR182); purple = canonical chemokines (Canonical CKR); and pink = relaxin receptors

(RL3R). Branches scaled by amino acid substitutions per site.

Figure 4: Distribution and duplication patterns of receptor groups.

A) Presence of all receptor groups are mapped onto a species tree. Gene trees and duplication

events are based on the gene tree to species tree reconciliation analyses. The nomenclature for

genes is primarily based on human chemokines. The canonical chemokines had 5 paralogs present in

the vertebrate common ancestor. These undergo a heterogeneous pattern of duplication

throughout vertebrates with different paralogs duplicating different number of times and in

different groups of species. Chemokines that have been classically described as having either

homeostatic or inflammatory function are indicated with a circle or a star respectively. The

classification used here was based on Zlotnik and Yoshie 2012 (7). **B)** Number of complements for

each receptor group at key species nodes are mapped onto the species tree. The number of

complements in each group reflects the pattern of duplications. The chemokine groups diverged in

the vertebrate stem group. The major expansion occurred at the level of jawed vertebrates with

canonical chemokine receptors, the chemokine-like receptor plus group and intermediate groups

increasing in copy number. Canonical chemokine underwent another small subsequent increase

within placentals. Silhouette images are by Andreas Hejnol (*Xenopus laevis*); Andy Wilson (*Anas*

platyrhynchos, Taeniopygia guttata); Carlos Cano-Barbacil (*Salmo trutta*); Christoph Schomburg

(*Anolis carolinensis, Ciona intestinalis, Eptatretus burgeri, Petromyzon marinus*); Christopher Kenaley

(*Mola mola*); Chuanixn Yu (*Latimeria chalumnae*); Daniel Jaron (*Mus musculus*); Daniel Stadtmauer

(*Monodelphis domestica*); Fernando Carezzano (Asteroidea); Ingo Braasch (*Callorhinchus milii*); Jake

Warner (*Danio rerio*); Kamil S. Jaron (*Poecilia formosa*); Mali'o Kodis, photograph by Hans Hillewaert

(*Branchiostoma lanceolatum*, https://www.phylopic.org/images/719d7b41-cedc-4c97-9ffe-

dd8809f85553/branchiostoma-lanceolatum); Margot Michaud (*Canis lupus, Physeter*

macrocephalus); NASA (*Homo sapiens sapiens*); Nathan Hermann (*Scophthalmus aquosus*); Ryan

Cupo (*Rattus norvegicus*); seung9park (*Takifugu rubripes rubripes*); Soledad Miranda-Rottmann

(*Pelodiscus sinensis*, https://www.phylopic.org/images/929fd134-bbd7-4744-987f-

1975107029f5/pelodiscus-sinensis); Steven Traver (*Gallus gallus domesticus, Ornithorhynchus*

anatinus); Stuart Humphries (*Thunnus thynnus*); T. Michael Keesey (after Colin M. L. Burnett) (*Gorilla*

gorilla gorilla); Thomas Hegna (based on picture by Nicolas Gompel) (*Drosophila (Drosophila)*

mojavensis); and Yan Wong (*Balanoglossus*).

Figure 5: Summary of the evolution of ligands and receptors.

A summary diagram of the evolution of the different chemokine system components. A simplified

phylogenetic tree of species is shown, calibrated to time according to Dohrmann and Wörheide 2017

(105) for Deuterostomia and Bilateria nodes and Delsuc et al. 2018 (106) for all other nodes. Circles

represent ligand groups, and 7 transmembrane domain structure icons represent GPCR groups. Icons

are colour coded by group, and placed adjacent to the branch in the species tree where they first

appear. X2 and X5 indicate the number of paralogs present for CXCL ligand group and the canonical

CKR groups respectively, on the branch where they first appear. Question mark refers to the

uncertainty regarding the origin of the CKLF group I in jawed vertebrates or deuterostome stem

group (see Figure 2). Geological column is shown along the bottom, in accordance with the ICS

International Chronostratigraphic Chart (107).

Supplementary Tables:

Table S1. List of the species used in this work and their respective BUSCO values and accession

number.

Table S2. Substitution models used in the phylogenetic analyses.

Supplementary Figure captions:

Figure S1. CLANS clustering of chemokines and related molecules sequences. Initial identification

and annotation of clusters was performed at the strict p-value of 1E-35 **(A)**. Subsequent loosening of

the p-value clarified the relationships across clusters and defined bigger groups. At p-value 1E-15 **(B)**

two major canonical chemokine groups are well-defined: the CCL group, which includes also XCL and

X3CL; and the CXCL group. At this level of stringency, only few canonical chemokines remain

isolated: CCL27/28; CXCL12; CXCL14; CXCL16; as well as CXCL17. TAFA and CYTL are also isolated. At

p-value 1E-10 **(C)** the two major chemokine groups connect to each other. CCL27/28 is connected to

the CCL group and CXCL12 and CXCL14 are connected to the CXCL group, while CXCL16 and CXCL17

are still isolated. At p-value 1E-6 **(D)** all chemokine groups are connected in one big cluster, except

for CXCL17. TAFA and CYTL are also still isolated. Crosses indicate the few invertebrate sequences

that were collected from the BLAST search, more information in Supplementary Results.

Figure S2. CLANS clustering of CKLF Super Family sequences. Initial identification and annotation of

clusters was performed at the strict p-value of 1E-60 **(A)**. Subsequent loosening of the p-value

clarified the relationships across clusters and defined bigger groups. At 1E-20 **(B)** two major clusters

have formed. One, that we called CKLF group I, includes CKLF, CMTM1,2,3,5, and PLP2. The other,

that we called CKLF group II, includes CMTM4/6,7,8, and other groups. At 1E-16 **(C)** more sequences

have joined the two major groups that are still separate. At 1E-15 **(D)** the two major groups connect,

as well as few extra sequences, see Supplementary Results for extra details. Crosses indicate

invertebrate sequences.

Figure S3. Alignment of candidate brachiopod CCL24 sequence with mammalian CCL24s. Our BLAST

searches picked up a sequence from the brachiopod *Lingula unguis* that when re-blasted versus

SwissProt returned a CCL24 as hit. Alignment of the brachiopod sequence with mammalian CCL24

sequences reveals a poor overall conservation, with the brachiopod sequence also being significantly

longer than any of the other sequences. Further details about this sequence can be found in

supplementary file S3 and in Supplementary Results.

Figure S4. Alignment of candidate cnidarian CCL3 sequence with mammalian CCL3s. Our BLAST

searches picked up a sequence from the cnidarian *Clytia hemisphaerica* that when re-blasted versus

SwissProt returned a CCL3 as hit. Alignment of the cnidarian sequence with mammalian CCL3

sequences reveals a poor overall conservation, with the cnidarian sequence being extremely longer

than any of the other sequences. Further details about this sequence can be found in supplementary

file S3 and in Supplementary Results.

Figure S5. Alignment of candidate echinoderm CXCL10 sequence with mammalian CXCL10s. Our

BLAST searches picked up a sequence from the echinoderm *Acanthaster planci* that when re-blasted

versus SwissProt returned a CXCL10 as hit. Alignment of the echinoderm sequence with mammalian

CXCL10 sequences reveals a poor overall conservation, with the brachiopod sequence also being

significantly longer than any of the other sequences. Further details about this sequence can be

found in supplementary file S3 and in Supplementary Results.

Figure S6. Alignment of 4 candidate urochordate TAFA sequences with vertebrate TAFAs. Our

BLAST searches picked up 4 sequences from the urochordate *Ciona intestinalis*, that connected with

the TAFA cluster in the CLANS analysis. One of these sequences when blasted versus SwissProt

returned a TAFA as hit. This sequence was also annotated as TAFA by InterProScan. Alignment of the

urochordate sequences with vertebrate TAFA sequences reveals that only the one annotated as

TAFA aligns well. While the other 3 align poorly and are also significantly longer than any of the

other sequences. Further details about these sequences can be found in supplementary file S3 and

in Supplementary Results.

Figure S7. Alignment of best candidate urochordate TAFA sequence with vertebrate TAFAs. Of the

4 urochordate candidate TAFA sequences, only one was annotated as TAFA with both SwissProt and

InterProScan annotation and appeared to align well with other TAFAs with a preliminary alignment

with all urochordate sequences (Figure S6). Here we removed the other 3 urochordate sequences

and aligned only the best candidate with the vertebrate TAFAs. The sequence conservation is even

more apparent with this alignment. Importantly, 8 of the 10 typical cysteine residues of TAFA1-4 are

conserved, and the two missing cysteines are the same ones missing in TAFA5. Further discussion

can be found in Supplementary Results.

Figure S8. Unrooted phylogenetic tree of canonical chemokines with TBE supports. Phylogenetic

tree under the model GTR20+F+R4. Nodal support is calculated from 100 non-parametric bootstrap

repeats with transferable bootstrap expectation. CCL clade is in orange, CXCL clade in blue.

Figure S9. Unrooted phylogenetic tree of canonical chemokines with UFB supports. Phylogenetic

tree under the model GTR20+F+R4. Nodal support is calculated from 1000 ultrafast bootstrap

repeats. CCL clade is in orange, CXCL clade in blue.

Figure S10. Unrooted phylogenetic tree of TAFA with TBE supports. Phylogenetic tree under the

model JTT+R5. Nodal support is calculated from 100 non-parametric bootstrap repeats with

transferable bootstrap expectation.

Figure S11. Unrooted phylogenetic tree of TAFA with UFB supports. Phylogenetic tree under the

model JTT+R5. Nodal support is calculated from 1000 ultrafast bootstrap repeats.

Figure S12. Unrooted phylogenetic tree of CYTL with TBE supports. Phylogenetic tree under the

model JTT+I+G4. Nodal support is calculated from 100 non-parametric bootstrap repeats with

transferable bootstrap expectation.

Figure S13. Unrooted phylogenetic tree of CYTL with UFB supports. Phylogenetic tree under the

model JTT+I+G4. Nodal support is calculated from 1000 ultrafast bootstrap repeats.

Figure S14. Unrooted phylogenetic tree of CXCL17 with TBE supports. Phylogenetic tree under the

model JTT. Nodal support is calculated from 100 non-parametric bootstrap repeats with transferable

bootstrap expectation.

Figure S15. Unrooted phylogenetic tree of CXCL17 with UFB supports. Phylogenetic tree under the

model JTT. Nodal support is calculated from 1000 ultrafast bootstrap repeats.

Figure S16. Unrooted phylogenetic tree of CKLFSF with TBE supports. Phylogenetic tree under the

model GTR20+F+R7. Nodal support is calculated from 100 non-parametric bootstrap repeats with

transferable bootstrap expectation. Red clade = CMTM4/6; blue clade = CKLF I group; green clade =

CMTM7; turquois clade = MAL/MALL/MAL2.

Figure S17. Unrooted phylogenetic tree of CKLFSF with UFB supports. Phylogenetic tree under the

model GTR20+F+R7. Nodal support is calculated from 1000 ultrafast bootstrap repeats. Red clade =

CMTM4/6; blue clade = CKLF I group; green clade = CMTM7; turquois clade = MAL/MALL/MAL2.

Figure S18. Rooted species tree reconciled gene tree for canonical chemokines. The canonical

chemokines gene tree was reconciled with the species tree using GeneRax. "S" or "D" at the node

indicates a speciation or duplication event respectively. CCL clade is in orange, CXCL clade in blue.

Figure S19. Rooted species tree reconciled gene tree for TAFA. The TAFA gene tree was reconciled

with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or duplication

event respectively.

Figure S20. Rooted species tree reconciled gene tree for CYTL. The CYTL gene tree was reconciled

with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or duplication

event respectively.

Figure S21. Rooted species tree reconciled gene tree for CXCL17. The CXCL17 gene tree was

reconciled with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or

duplication event respectively.

Figure S22. Rooted species tree reconciled gene tree for CKLFSF. The CKLFSF gene tree was

reconciled with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or

duplication event respectively. Red clade = CMTM4/6; blue clade = CKLF I group; green clade =

CMTM7; turquois clade = MAL/MALL/MAL2.

Figure S23. CLANS clustering of receptors and related molecules sequences. A CLANS clustering

layout where shapes indicate sequences and lines are connections indicating similarity between

sequences at or surpassing the p-value similarity threshold. Sequences are positioned in clusters

based on similarity. Initial identification and annotation of clusters was performed using the inbuilt

convex clustering at the p-value of 1E-100. **(A)** clustering was loosened till the canonical receptor

annotated groups formed a cluster at 1E-65. **(B)** loosening of the p-value to 1E-60 identified

relationships between clusters of interest and identified the intermediate group as connecting to

both canonical and chemokine-like plus groups. All sequences connected to groups of interest are

vertebrate sequences. **(C)** further loosening to p-value 1E-50 connects the vertebrate sequences of

interest to a large cluster of sequences which contains vertebrate and invertebrate sequences which

are annotated as opioid and somatostatin receptors and other GPCRs. Crosses indicate invertebrate

sequences and Y-shape indicates the reference viral sequences included. Shapes are colour coded by

group of interest: purple = canonical chemokine receptors; yellow = chemokine-like plus; green =

atypical receptor 3/GPR182; blue = intermediate group; pink = relaxin receptors.

Figure S24. Unrooted phylogenetic tree of receptors with TBE supports. Phylogenetic tree of

receptor sequences of interest and putative outgroups under the model GTR20+F+G4. Sequences

used are a subset of sequences extracted from CLANs, specifically they are those in the chordate

specific clade in the ultrafast bootstrap tree. Nodal support is calculated from 100 non-parametric

bootstrap repeats with transferable bootstrap expectation. Branches colour coded by group of

interest: purple = canonical chemokine receptors; yellow = chemokine-like plus; green = atypical

receptor 3/GPR182; blue = intermediate group; pink = relaxin receptors.

1001

1002 **Figure S25. Unrooted phylogenetic tree of receptors with UFB supports.** Phylogenetic tree of all

1003 receptor sequences of interest and putative outgroups extracted from clans under the model

1004 GTR20+F+G4. Nodal support is calculated from 1000 ultrafast bootstrap repeats. Branches colour

1005   coded by group of interest: purple = canonical chemokine receptors; yellow = chemokine-like plus;

1006   green = atypical receptor 3/GPR182; blue = intermediate group; pink = relaxin receptors.

1007

Figure S26. Rooted species tree reconciled gene tree for receptors. The ultrafast bootstrap receptor

tree was modified to extract the subtree of the chordate specific clade. This gene tree was

reconciled with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or

duplication event respectively. Branches colour coded by group of interest: purple = canonical

chemokine receptors; yellow = chemokine-like plus; green = atypical receptor 3/GPR182; blue =

intermediate group; pink = relaxin receptors.

Fig. 2

Fig. 3

Fig. 4

Fig. 5