The origin, evolution and molecular diversity of the chemokine system

Alessandra Aleotti[1,2] *, Matthew Goulty[1,2] *, Clifton Lewis[1,2], Flaviano Giorgini[1,2] and Roberto Feuda[1,2]

#

*Equal contribution

#Author for correspondence: rf190@leicester.ac.uk

[1] Department of Genetics and Genome Biology University of Leicester, Leicester, United Kingdom

[2]Neurogenetics Group, College of Life Sciences, University of Leicester, Leicester, United Kingdom

Keywords: Chemokine, Phylogeny, Evolution, Vertebrate, GPCR, Receptor, Ligand, CK, CKL, CKR,

CKLF, TAFA, CYTL

ABSTRACT

Chemokine signalling performs key functions in cell migration via chemoattraction, such as attracting leukocytes to the site of infection during host defence. The system consists of a ligand, the chemokine, usually secreted outside the cell, and a chemokine receptor on the surface of a target cell that recognises the ligand. A number of non-canonical components interact with the system. While canonical components have been described in vertebrate lineages, the distribution of the non-canonical components is less clear. Uncertainty over the relationships between canonical and non-canonical components hampers our understanding of the evolution of the system. We used phylogenetic methods, including gene-tree to species-tree reconciliation to untangle the relationships between canonical and non-canonical components, identify gene duplication events and clarify the origin of the system. We found that unrelated ligand groups independently evolved chemokine-like functions. We found non-canonical ligands outside vertebrates, such as TAFA "chemokines" found in urochordates. In contrast, all receptor groups are vertebrate-specific and all - except ACKR1 - originated from a common ancestor in early vertebrates. Both ligand and receptor copy numbers expanded through gene duplication events at the base of jawed vertebrates, with subsequent waves of innovation occurring in bony fish and mammals.

INTRODUCTION

The chemokine system is responsible for regulating many biological processes, including host defence, neuronal communication and homeostasis (1–3). It has two components, a ligand, usually a small cytokine called a chemokine, and a receptor. The ligands are categorized into four groups, XC, CC, CXC, and CX3C, according to the pattern of cysteine residues in the N-terminal portion of the protein (4). Likewise, the receptors are classified based on the ligands they bind to into four groups, the XCR, CCR, CXCR, and CX3CR, and all of them belong to the GPCR class A superfamily (5). In addition to canonical components, other molecules have been discovered to function similarly to chemokine ligands (6) or receptors (7) (see Table S1). These include: the chemokine-like factor (CKLF) that binds to chemokine receptor CCR4 (8, 9) and drives cell migration *in vivo* (10); TAFA chemokines, expressed mainly in the nervous system, which share structural similarities to canonical chemokines (11, 12) and bind GPCRs related to chemokine receptors, e.g. formyl peptide receptors (13, 14) and GPR1 (15); Cytokine-like 1 (CYTL1) that binds CCR2 (16) and has been suggested to be related to CC ligands based on the presence of a IL8-like chemokine fold (17). There are also non-canonical chemokine receptors, such as: the chemokine-like receptor (CML1, or also CMKLR1) (18); atypical chemokine receptors (ACKRs) (19); and viral chemokine receptors (20–23). Unlike other chemokine receptors, atypical receptors cannot initiate classical chemokine signaling upon ligand binding (19, 24). The human genome encodes four types of atypical chemokine receptors: the ACKR1 (also known as DARC), ACKR2 (also known as D6), ACKR3 (also known as CXCR7) and ACKR4 (also known as CCRL1) (25, 26). Additionally, several proteins of viral origins, such as US28 from human cytomegalovirus, have chemokine-receptor/binding activity (20, 21). These viral proteins can bind a wide array of chemokine ligands (21).

Despite the extensive research on the chemokine system, with over 320,000 papers available on PubMed, many aspects of its evolution remain unclear. For instance, the homology between canonical and non-canonical ligands is uncertain and supported by circumstantial evidence, such as shared specific motifs (10, 11, 17, 27). Furthermore, the relationships between canonical,

atypical, and viral receptors and the outgroup of the canonical chemokine receptors remain

uncertain. Finally, the evolutionary history of the canonical and non-canonical components remains

poorly understood outside a few key model systems (5, 28, 29). These outstanding questions share

common underlying causes, including the use of inadequate inference methods (such as relying

solely on sequence similarities) and limited sampling of species (e.g., focusing mainly on humans,

mice, and zebrafish (30, 31)). Additionally, solving the phylogenetic relationships for short molecules

such as chemokine receptors and ligands is particularly challenging due to the lack of strong

phylogenetic signals (32).

Here, to clarify these outstanding questions, we use state-of-the-art phylogenetic methods,

including those designed for single-gene phylogenies, a large taxonomical sampling comprising both

vertebrate and invertebrate genomes and the entire complement of canonical and non-canonical

components of both receptors and ligands. Our findings substantially clarify the phylogenetic

relationship between canonical and non-canonical ligands and receptors and suggest that unrelated

proteins evolved "chemokine-like" ligand function multiple times independently. Additionally, we

discovered that all the canonical and non-canonical chemokine receptors (except ACKR1) originated

from a single duplication in the vertebrate stem group, which also gave rise to many GPCRs. Lastly,

we characterized the complement of canonical and non-canonical components in the common

ancestor of vertebrates and identified several other ligands and receptors with potential chemokine-

related properties that could be explored in future functional work.

RESULTS

There are five unrelated groups of ligands.

Initially, we focused on the ligands, including all the canonical chemokines, the CYTL, the

TAFAs and the CKLF Super Family (CKLFSF) proteins. The presence of a four transmembrane MARVEL

domain in the latter proteins (10, 33, 34) distinguishes them from canonical chemokines, the CYTL

and the TAFAs. Therefore, we separated these two groups for further analysis. Using BLASTP or PSI-

BLAST (35–37) (see Materials and Methods for more details) against 64 species from 19 animal phyla

(Table S2), we identified 891 putative homologs for chemokines, TAFA and CYTL and 602 putative

homologs of the CKLF Super Family.

We utilized CLANS (38, 39), a clustering tool based on sequence similarity and local

alignment, to identify homology within these two groups. Unlike traditional phylogenetic methods,

CLANS assigns homology between sequences based on BLAST and customizable stringency levels

defined according to p-values (38). When two (or more) sequences are connected at a lower p-value

(closer to 0), this indicates a high level of homology. Conversely, if two or more sequences only

connect at a higher p-value, this suggests a relatively low level of sequence homology. Our analysis

shows that canonical chemokines form a distinct group with a clear distinction between C-X-C-type

and C-C-type (Figure 1A). Whereas, CXCL17, TAFA and CYTL remain separate from canonical

chemokines and from each other even at the loosest p-values tested (Figure 1A). The distinction

between CXCL17 and all other canonical chemokines is consistent with our receptor results showing

that the receptor for CXCL17, GPR35 (40), is also not within the canonical chemokine receptor group

(see below). Within the CKLFSF, two large clusters were identified, named CKLF I and CKLF II,

although these ultimately connect to form one large superfamily (Figure 1C). These clusters are

robust to the different stringency thresholds used (Figures S1 and S2 and Materials and Methods for

further details). Our results indicate that even when the stringency level to detect homology is

relaxed, canonical chemokines, TAFA, CYTL, and CXCL17 remain in distinct clusters. This suggests

that, similarly to CKLFs, these proteins are not homologous and convergently evolved chemokine-

like properties. We have thus identified five distinct groups of ligands: i) the canonical chemokines,

ii) TAFA "chemokines", iii) CYTL, iv) CXCL17, and v) CKLF Super Family (Figure 1A and 1C).

The evolution of chemokine and chemokine-like ligands in animals.

To better understand the evolution of both canonical and non-canonical chemokine ligands,

we performed a separate phylogenetic reconstruction for each group (Figure 1B, D and Figures S8-

17) (see methods for details). To evaluate the nodal support, in addition to the UltraFast bootstrap

(UFB) (41, 42), we used Transfer Bootstrap Expectation (TBE), a method that has been developed for

single-gene phylogeny (43). To evaluate ortholog/paralog relationships and overall dynamics of the

ligand complement, we used GeneRax (44). This new method uses maximum likelihood to reconcile

the gene tree with the species tree (44). In brief, given a gene and species tree, GeneRax uses a

maximum likelihood approach to optimize the duplication and loss events (44, 45).

Our analysis initially identified a few invertebrate putative chemokine ligands (Figure 1A),

however, these sequences lacked protein signatures associated with the canonical ligands (Figures

S3-5 and File S3), and they were therefore excluded from further analysis (see Supplementary

Results for further information). The phylogenetic tree for the canonical ligands identifies two major

groups, the CC-type, which also includes the XC- and X3C-types, and the CXC-type (TBE = 0.95, UFB =

92%) (Figure 1B and Figures S8,9), confirming the previous finding obtained using synteny data (46,

47). Next, to clarify the distribution of canonical chemokines, we first reconciled their gene tree with

the species tree and then used the reconciled tree to trace the presence/absence of each chemokine

group throughout all the species (Figure 2A and Figure S18). Our results confirm previous findings

that canonical chemokines are uniquely present in vertebrates (28, 46). Additionally, they indicate

that chemokines are not evenly distributed across vertebrates. Some are very ancient, e.g., CXCL12

is present in lamprey; CXCL14 and CCL20 are present in all jawed vertebrates; and CXCL8 is present

in all jawed vertebrates except in the cartilaginous fishes lineage. However, a large part of the

chemokine diversity evolved within mammals (e.g., CXCL1/2/3, CXCL16, and CCL25), particularly

placentals (e.g., CXCL5/6 and CCL3/18). The complement of the canonical chemokines undergoes the

largest expansion at the base of jawed vertebrates, where there is an expansion from 4 to 18 genes

(Figure 2B). A second expansion occurred at the base of bony fishes (i.e., Osteichthyes) followed by a

relative stability until placental mammals, where the total number of canonical chemokine ligands

jumped to 45 genes. Finally, unlike previous works (48) our results support the presence of orthologs

of both CC-type and CXC-type in the common ancestor of all vertebrates (Figure 2A).

Differently from the canonical chemokines, we identified a *bona fide* TAFA, i.e., with specific

protein motifs, in the urochordates, the sister group to vertebrates (see Supplementary Results and

Figures S6-7). The phylogenetic trees (Supplementary Figures S10,11) identified monophyletic

groups for TAFA5 (TBE=0.98, UFB=98%), TAFA1 (TBE=0.94, UFB=98%), TAFA4 (TBE=0.77, UFB=75%)

and TAFA2/3 (TBE=0.65, UFB=84%). The reconciled tree from GeneRax places the root at the

urochordate sequence (Figure S19), therefore clarifying that the TAFA5 clade is the sister group to

TAFA1-4 (Figure 2A). The family originated in the ancestor of urochordates and vertebrates, and the

first duplications occurred at the base of vertebrates giving rise to the TAFA5 split followed by the

TAFA1 split. Subsequently, at the base of jawed vertebrates, additional duplications bring the

complements from 3 to 10 (Figure 2B), giving rise to the remaining groups so that all jawed

vertebrates possess the full diversity of TAFAs.

The phylogenetic trees for CYTL and CXCL17 mainly reflect the species trees (Figures S12-15),

and the reconciliations revealed very simple complement dynamics (Figure 2B and Figures S20,21).

However, these molecules show a remarkable difference in their distribution. CYTLs are present

throughout gnathostomes, while CXCL17 is found only in placental mammals (Figure 2A).

The phylogenetic analysis for the CKLF super family (Figure 1D and Figures S16,17) recovered

a monophyletic clade for the CKLF I group (TBE=0.96, UFB=80%) that we had already identified

through CLANS. This group contains CKLF, that is known to interact with C-C chemokine receptor 4

(8, 9), as well as CMTM1, 2, 3, 5, and proteolipid protein 2 (PLP2). Other monophyletic clades that

are consistent with the CLANS are CMTM4/6 (TBE=0.90, UFB=61%), CMTM7 (TBE=0.92, UFB=83%)

and a clade containing CMTM8 plus other related molecules such as plasmolipin (PLLP) and myelin

and lymphocyte proteins (MAL) (TBE=0.89, UFB=60%). The latter were all part of a large cluster that

we called CKLF II in the CLANS (Figure 1C). However, the placement of the root of the tree in Figure

1D can affect the interpretation of the relationships among CKLF II subgroups. To address this

problem and clarify the patterns of duplications and the presence/absence of each group

throughout animals, we used GeneRax to reconcile the gene with the species tree (see above and

Material and Methods for details). Our results suggest (Figure 2 and Figure S22) that most CKLFSF

groups, such as CMTM4,6 and 8, originate in the vertebrate stem group from pre-existing CMTM

genes and are widely distributed in animals. The CKLF I subgroups originate from duplications at the

base of jawed vertebrates, except for the split between CKLF and CMTM1 that occurs only within

mammals (Figure 2A). We observe the major two expansions of the CKLFSF genes in the stem group

of vertebrates (from 6 to 10 complements), and then in jawed vertebrates (from 10 to 16

complements). Interestingly the extents of these expansions are less drastic than those we see for

canonical chemokines (Figure 2B). In total, we have identified that the five distinct ligand groups

have a different origin in the animal tree of life and underwent divergent evolutionary histories.

Canonical and non-canonical chemokine receptors are divided into four groups.

Next, we investigated the origin and pattern of duplication for the chemokine and

chemokine-like receptors. Using BLASTP against the 64 species, we identified 7,157 putative

chemokine receptors (see Materials and Methods for more details), and we investigated their

relationships using CLANS (see above for justification). The result (Figure S23C) identifies four main

groups of chemokine and chemokine-like receptors. The first comprises canonical receptors (i.e.,

CCR, CXCR, CX3CR1, CX3C, and XCR1), and the second includes atypical receptor 3 and GPCR182,

which has been recently shown to have chemokine activity (49). The third group, which we named

Chemokine-like plus (CML-plus), contains the chemokine-like receptors (CML1 also known as

chemerin receptor 1), formyl peptide receptors (FPR) that bind the TAFA ligands (13, 14) and other

GPCRs such as GPCR1 (chemerin receptor 2), GPCR33, PTGDR2. Furthermore, the CLANS analysis

identifies an intermediate group containing angiotensin, apelin and other receptors and shows

sequence similarity to canonical and chemokine-like receptors (Figure S23B). Finally, our analysis

identifies a small cluster composed of only ACKR1 that do not connect to other GPCRs or other

atypical receptors even at loose p-value thresholds. This indicates that their sequence is either non-

homologous or highly divergent from other chemokines and atypical receptors. Overall, these

groups are robust to the stringency threshold used (i.e., different p-values) (Figure S23).

Interestingly, no specific cluster of viral or viral-like receptors was identified, but 6 of the reference

viral receptor sequences clustered with the canonical chemokine receptors.

Altogether, these results confirm the homology between the canonical receptors and

atypical receptor 3/GPCR182. However, the results indicate that the other GPCRs, such as the

chemokine-like receptors, formyl peptide receptors, GPCR1, and GPCR33, are also closely related to

the canonical receptors. Remarkably, these results also indicate that ACKR1 is not homologous to the

canonical chemokine receptors. Furthermore, all clusters of chemokine receptors contained only

vertebrate sequences, except for the receptors of viral origin.

Canonical and chemokine-like receptors derive from single gene duplication in the ancestor of

vertebrates.

Previous studies suggested that the chemokine receptors evolved from a duplication of

angiotensin receptors (50) or adrenomedullin receptors (28, 51). However, these works were based

on error-prone phylogenetic methods such as Neighbour Joining (51). Our CLANS results indicate

that chemokine receptors and chemokine-like receptors have only been observed in vertebrates.

Therefore, we need to focus on invertebrate genomes to clarify the chemokine's outgroup. To clarify

this, we lowered the p-value thresholds of CLANS (to p-value < $1e^{-50}$) and collected a combined

dataset including all chemokine sequences and outgroups (i.e., sequences that connect to the

chemokine cluster), resulting in 3,026 sequences. We then performed a phylogenetic tree on this

dataset using maximum likelihood methods with UFB and TBE for evaluating nodal support (see

above and Materials and Methods for details).

Our combined phylogenetic analysis shows strong support for the monophyly of canonical

chemokines (UFB=96, TBE=1.0), the CML-plus (UFB=95, TBE=0.99) and the atypical 3/GPCR182

(UFB=100, TBE=1) (Figure 3, S24 and S25). In contrast, viral chemokines are paraphyletic, with three

sequences placed within the canonical chemokines and 3 forming a monophyletic group sister to

them (UBF=84 TBE=1.0). Our results also suggest that the intermediate group, which includes apelin

receptors, angiotensin receptors, bradykinin receptors, and orphan GPCRs (e.g., GPCR25; GPCR15)

forms a monophyletic clade with the canonical chemokines, CML-plus group and atypical3/GPCR182

(UFB=61, TBE=0.91). However, its position changes between the sister group to canonical

chemokines plus atypical3/GPCR182 in the TBE tree (TBE=0.84) and sister to CML plus in the ultrafast

bootstrap tree (UFB=38).

All the groups mentioned above form a large clade composed of vertebrate-specific GPCRs

(UBF=100 TBE=0.96) that also includes other GPCRs, such as CLTR and P2RY receptors (Figure 3, S24

and S25). Another orphan GPCR, GPR35, had been proposed as a potential chemokine receptor

(putative CXCR8) (40). Our analysis collected GPR35 and placed it within this large vertebrate specific

clade indicating that is it also a vertebrate specific gene but not phylogenetically a 'canonical'

chemokine. The closest outgroup to this clade is composed of a few sequences from urochordates,

the sister group of vertebrates (UFB=49 TBE=0.91) (Figure 3, S24 and S25). Interestingly, as the sister

group of this clade, we identify a group composed of Relaxin receptors which contain sequences

from both urochordates and vertebrates (UBF=53 TBE=0.95). Finally, as the sister group of these

large clades, we identified a clade of cephalochordate-specific sequences (UBF=44).

To clarify the duplication pattern and origin of the chemokine receptors, we used GeneRax

(44) (see Materials and Methods). Our results indicate (Figure 4A, S26) that all chemokine receptors

(except ACKR1) originated from a duplication in the stem lineage of vertebrates. This duplication of

an unknown GPCR gave rise to the CML-plus, the canonical chemokines and atypical 3/GPCR182

groups as well as the intermediate group and other GPCRs (Figure 4A, S26). This result is consistent

with the distribution of the paralogous Relaxin receptors which are present both in urochordates

and vertebrates and the position of the orphan urochordate sequences as sister group of canonical

chemokines, CML-plus group and atypical3/GPCR182 and other GPCRs (see above).

We used the reconciliation to better understand the repertoires of receptors present at key

nodes during vertebrate evolution. Our results (Figure 4B) show a substantial difference in the

duplication pattern of different receptor families. For example, the complement of the

atypical3/GPCR182 remains constant throughout vertebrate evolution while the canonical and

chemokine-like groups expanded dramatically. The canonical chemokine expanded from 5 to 20

genes and the CML-plus from 1 to 11 in the ancestor of the jawed vertebrates (Figure 4B). The

expansion of the canonical CKRs is also not evenly distributed across its subgroups, with the

ancestral CC type receptors undergoing a series of duplications in jawed vertebrates while the CXCR

paralogs did not, specifically one (CXCR4) remains in single copy across all vertebrates. We inferred

that in the stem lineage of vertebrates, five canonical chemokine paralogs had already diverged,

representing the two major types of receptors (2 CCR and 3 CXCR paralogs). Also, present in the

stem lineage of vertebrates were ACKR3 and GPCR182 as well as a single copy gene which would

later diverge to produce all the CML-plus clade.

DISCUSSION

This work substantially clarifies the evolutionary assembly of the chemokine system. Our

analysis shows that, contrary to the receptors which evolved from a single duplication event in the

vertebrate stem group, several unrelated molecules acquired the ability to interact with chemokine

receptors over the course of evolutionary history. Furthermore, our results (summarized in Figure 5)

suggest that the key components of the chemokine system, including the chemokine receptors

themselves, evolved in the stem group of vertebrates in the Ediacaran/Cambrian boundary around

540 million years ago and then underwent substantial diversification in the stem group of jawed

vertebrates. These findings shed new light on the complex evolutionary history of the chemokine

system.

Unrelated molecules converged to chemokine function.

Based on the presence of shared protein motifs, TAFA "chemokines" (11, 12), CXCL17 (52,

53) and CYTL (17) have been proposed to be homologous to chemokine ligands. However, our

findings strongly suggest that these molecules are not homologous (Figure 1) and likely acquired the

ability to activate a chemokine-like response through convergent evolution. Our conclusions differ

from those previous studies (11, 17, 52, 53) due to the differences in data completeness and

methodological approach. Specifically, we used a complete set of canonical and non-canonical

ligands, and assessed the homology using overall sequence similarity rather than single motifs. Our

results support and expand upon the findings of (27), which suggested that the presence of a CXC or

CC motif is necessary but not sufficient for a protein to be defined as a chemokine ligand. Similarly,

CKLF has been considered a "new member" of the chemokine family based on its function (10) we

argue that classification based solely on function is insufficient and can be misleading. Instead, we

recommend considering the evolutionary relationships among these molecules as the primary

criterion for classification.

Most of the canonical and non-canonical ligands are vertebrate innovations.

Our results clarify the distribution of canonical chemokine ligands in animals (Figure 2) and

confirm that they are present only in vertebrates (28). We identify orthologs of CXCL and CCL ligands

in both extant lineages of cyclostomes (Figure 2A). While chemokines have already been described in

lamprey (48, 54, 55), it is the first time, to our knowledge, that they are described also in hagfish.

Our findings also indicate that both CC and CXC types were present in the common ancestor of all

vertebrates and that few ancestral genes gave rise to the entire diversity of ligands that we know in

current animals. Furthermore, our results indicate that many chemokines, such as CXCL1-7, CXCL16,

as well as CCL25, CCL11/13, and CCL2/7, are uniquely present in mammals suggesting that the

mammal ligand repertoire is substantially more complex than the one observed in other vertebrates.

Regarding non-canonical chemokine-like families, our findings indicate that the TAFA family

originated in the ancestor of vertebrates and urochordates; CYTL is a novelty of jawed vertebrates;

and CXCL17 is mammal-specific and likely unrelated to canonical chemokines (similar to its receptor,

GPR35 (40), that is not a canonical chemokine receptor). The CKLF super family has a more complex

pattern with the presence of few groups in invertebrates and then great expansions occurring at the

base of vertebrates. The CKLFSF includes a monophyletic clade (CKLF group I) comprising the original

CKLF that binds CCR4, as well as CMTM1,2,3,5, derived from duplications at the jawed vertebrates

stem group. Interestingly, our analysis also revealed that additional molecules not previously

considered part of the CKLF super family are closely related to classic members and should be

included in it. For example, proteolipid protein 2 (PLP2) belongs to the CKLF I group and is, therefore

more closely related to the CKLF with chemokine function than several other CKLFSF members.

Similarly, CMTM8 is more closely related to plasmolipin (PLLP) and myelin and lymphocyte protein

(MAL) than to any of the classic CKLFSF members. Although this relationship had been proposed

based only on sequence similarity (33), our phylogenetic analysis provides additional evidence for it.

Therefore, all these additional members should also be explored for potential chemokine-like

functions in both vertebrates and invertebrates.

All but one receptor derive from a single gene duplication.

Our results clarify the distribution of canonical chemokine receptors in vertebrates (Figure

4), and their evolutionary relationships and identify the pattern of duplication that leads to their

origin (Figure 4A, S26). Unlike previous works (56), we identify that atypical receptors do not form a

monophyletic group. Specifically, atypical 2 and 4 are part of the canonical clade specifically related

to CC-type receptor subclades. Furthermore, we find that the atypical 3 receptors are related to

GPCR182, supporting previous functional data suggesting that the latter are atypical chemokine

receptors binding CXCL10, 12, and 13 (49). We attribute these differences to our use of wider GPCR

sampling and improved methods for phylogenetic inference.

Remarkably, our results do not identify ACKR1 as related to the main chemokine receptors

but rather as a divergent clade (Figure S23). To our knowledge, this is the first time this observation

has been made. Our current results do not allow us to clarify the evolutionary origin ofACKR1.

However, the presence of 7TMD domains suggests that they are GPCRs that independently acquired

the ability to bind chemokines. Alternatively, similarly to other genes evolved in the immune system,

ACKR1 may have been subjected to strong selective pressures that substantially changed their

sequence, obscuring their phylogenetic relationships.

Viral chemokines represent a cryptic group that can bind multiple chemokines (20, 21).

Despite their functional similarity to canonical chemokines, viral chemokines' evolutionary origin and

distribution remain poorly understood. Our results indicate that viral GPCRs do not form a

monophyletic group, suggesting that the ability to encode chemokine-like receptors has evolved

independently in multiple viruses, including cytomegaloviruses and poxviruses. However, further

research is needed to clarify the distribution and origins of viral GPCRs.

Our analysis reveals that the clade comprising apelin receptors, angiotensin receptors,

bradykinin receptors, and orphan GPCRs (shown in Figure 3, 4 and S24–26) is closely related to

chemokine receptors. This finding partially supports previous studies (50) that suggested a gene

duplication event gave rise to both chemokines and angiotensin receptors. Interestingly, we found

that single gene duplication in the vertebrate stem group led to the emergence of canonical

receptors and atypical 2,3,4, GPCR182, chemokine-like, formyl peptide receptors, the intermediate

group, and many other known and orphan GPCRs including putative chemokine receptor GPR35.

These findings suggest that two rounds of genome duplication (57, 58) played a role in the expansion

of GPCR gene families. Future research will focus on investigating the functions of the orphan genes

and many-to-one orthologs discovered in urochordates. This will provide further insight into the

evolution and diversification of GPCR families in vertebrates.

The molecular assembly of the chemokine system

In this work, we explored the evolution of both ligand and receptor components of the

chemokine signaling system, including non-canonical molecules with either chemokine-like function

or sequence similarity. Our analysis suggests that the canonical chemokine signaling evolved in the

vertebrate stem group (about 540 Mya) likely due to the two rounds of genome duplication that

gave rise to many vertebrate novelties (57, 58). We found that the ancestral vertebrate repertoire

included orthologs of both major ligand groups (CXCL and CCL) and both CCR and CXCR receptors

and non-canonical components such as TAFA and CKLFSF ligands, and the receptors Atypical 3 and

GPCR182 (Figure 5). The distribution of ligands and receptors in the ancestor of all vertebrates,

seems to confirm the hypothesis that the ancestral function of chemokines was homeostatic (e.g.,

CXCL12, CXCL14) with inflammatory functions arising from recent duplications (e.g., CXCL5, CXCL6),

potentially reflecting a rapid evolution induced by the selective pressure of new pathogens (30).

The evolutionary history of canonical components includes several examples of known

ligand-receptor pairs following a corresponding pattern of origin and temporal dynamics of

duplications. This is true for example, for the ancient homeostatic CXCL12 ligand and its

corresponding receptors CXCR4 and ACKR3, that all originated in early vertebrates (30). The early

origin and conservation of CXCR4 and CXCL12 in the ancestor of vertebrates is interesting as this pair

plays a key role in the migration of neural crest cells (59) - a key vertebrate innovation (60). For

many of the non-canonical components, however, the ligand-receptor interactions are largely

unclear, and their pattern throughout vertebrate evolution remains to be explored. Overall, our

results indicate that three waves of molecular innovation in the vertebrates, jawed vertebrates,

bony fishes and mammal stem group increased the chemokine system's molecular complexity

(Figure 5), allowing for the fine-tuning present in modern-day animals.

MATERIALS AND METHODS

Data Mining and Dataset Assembly. We collected 64 proteomes from 25 vertebrates, six chordates

and 33 other animals covering the whole animal tree (Table S2). BUSCO v4.0.6 (61, 62) and the

metazoa_odb10 set of 954 genes were used to evaluate their completeness (Table S2).

To identify potential homologs of canonical chemokines, TAFA chemokines and CYTL1, we

used 207 curated sequences that we obtained from SwissProt (63, 64) as seeds for an initial BLASTP

(35, 37) with e-value < e^{-10}. To identify putative chemokines in cyclostomes, the lamprey *Petromyzon*

marinus (65) and the hagfish *Eptatretus burger* (66), we loosened the e-value to 0.05. Where

putative chemokine sequences were found for one cyclostome species but not the other, those

sequences were used to search again the other species. Furthermore, to investigate the presence of

ligands outside vertebrates, we performed an additional BLASTP on invertebrate proteomes with an

even looser e-value (0.1) and collected only up to five hits. This provided 18 initial candidate

homologs spanning multiple invertebrate phyla. Further characterisation of these invertebrate

sequences, through BLASTP versus SwissProt, protein domains search with InterProScan (67, 68),

position in CLANS analysis (see below) and, where necessary, multiple sequence alignments, lead us

to retain only one urochordate sequence as a putative TAFA homolog (see Supplementary Results

for details).

To identify homologs for the CKLF superfamily, we used 21 SwissProt-reviewed sequences.

In addition to the BLASTP search, we used a position-specific iterative BLAST (PSI-BLAST) (36) with an

e-value threshold of $<e^{-10}$. Using this approach, we identified a total of 590 putative homologs,

including 186 from invertebrates.

We used BLASTP using 178 manually annotated receptor sequences from SwissProt as query

sequences for the chemokine receptors. This includes all human canonical and atypical chemokine

receptors (69). We also collected 8 viral sequences with chemokine receptor activity from UniProt

(70) and performed a second BLASTP. We extracted all BLAST hits with e-values $< e^{-10}$ and used

Phobius (71) to predict their transmembrane domain structure. Only sequences with 5-8

transmembrane domains were kept. Hit sequences were annotated by their top 5 BLAST hits against

SwissProt. All hits from both BLASTs were merged and filtered by cd-hit (72, 73) to remove

redundant sequences at the 95% similarity threshold. This resulted in 7,157 putative chemokine

GPCR sequences.

Identification of subgroups with Cluster Analysis of Sequences (CLANS). We utilized CLANS (38, 39)

with default parameters and different p-values (i.e., stringency values) to visualize the relationships

between subgroups of ligands and receptors. We assessed the similarity and interrelationships

between different clusters by gradually relaxing the p-value threshold (Figures S1, S2 and S23).

Additionally, we annotated each cluster using gene annotations for key species *Homo, Mus, Gorilla,*

Gallus, Anolis and *Danio*. In the case of the receptors, to improve the cluster annotation all human

Class-A GPCRs (excluding olfactory receptors) from GPCRdb (74) were added to the dataset as well

as the 8 seed viral chemokine receptors from UniProt (70).

Alignment and phylogenetic analysis

Alignment. All ligand and receptor sequences were aligned using MAFFT (75, 76) with the --auto

setting and using trimAl (77) to remove positions with >70% gaps.

Gene Trees. All gene alignments were analysed using IQTREE2 (78), the model test algorithm (79)

was used to select the best substitution model for each analysis. Best models selected by IQTREE2

for each set are listed in Table S3 (for receptors we manually selected GTR20+F+G4 as the model as

it was a large dataset). Nodal support was estimated using 1,000 ultrafast bootstrap (41, 42)

replicates. All analyses were repeated to run 100 non-parametric bootstrap repeats to calculate

nodal support with transferable bootstrap expectation: which is specifically designed to account for

phylogenetic instability (43).

409   For the receptors, due to the high computational burden of running TBE analyses on

sequence-dense datasets, we first analysed the full set of 3,026 sequences connected in CLANS at a

p-value of $< 1\ e^{-50}$ using UFB (Figure S25). Then, we extracted the chordate-specific clade sequences,

including all chemokine groups and their immediate outgroups, to analyse using TBE.

Gene tree species tree reconciliation. To understand the pattern of duplication and the evolution of

gene complement we used GeneRax (44). GeneRax requires a gene tree that was obtained as

described above and a species tree that we constructed manually using publicly available

information. In the instances where the genes tree contained polytomies, we used ETE3 (80) to solve

them. The undated DL mode and the closest approximation of the best-fitting substitution model

were used for each alignment. To track the evolution of sub-lineages within each group, we used

annotated sequences of key species (e.g., *Homo sapiens* and *Mus musculus*) as reference. For the

receptors, we used only the chordate-specific clade subtree and sequences due to the

computational burden of running GeneRax on a high number of sequences. For species tree-gene

tree reconciliation, we treat the viral sequences as human sequences.

DATA AVAILABILITY

Supplementary material and raw output files for all the analyses described in this paper are available

in the GitHub page: [Roberto-Feuda-Lab/Chemokine2023 (github.com)](github.com).

ACKNOWLEDGEMENTS

This work is supported by a University Research Fellowship (UF160226) to RF. AA is supported by a

Research Grant from the Royal Society to RF (RGF\R1\181012). MG is supported by a PhD

Scholarship from the University of Leicester. CL is supported by a BBRSC MIBPT fellowship. This

research used the ALICE High-Performance Computing Facility at the University of Leicester.

REFERENCES

1.   X. Blanchet, M. Langer, C. Weber, R. Koenen, P. von Hundelshausen, Touch of Chemokines.

Frontiers in Immunology **3**, 175 (2012).

2.   P. López-Cotarelo, C. Gómez-Moreira, O. Criado-García, L. Sánchez, J. L. Rodríguez-

Fernández, Beyond Chemoattraction: Multifunctionality of Chemokine Receptors in Leukocytes.

Trends in Immunology **38**, 927–941 (2017).

3.   P. B. Tran, R. J. Miller, Chemokine receptors: signposts to brain development and disease.

Nature Reviews Neuroscience **4**, 444–455 (2003).

4.   A. Zlotnik, O. Yoshie, Chemokines: A New Classification System and Their Role in Immunity.

Immunity **12**, 121–127 (2000).

5.   H. Nomiyama, N. Osada, O. Yoshie, A family tree of vertebrate chemokine receptors for a

unified nomenclature. *Developmental & Comparative Immunology* **35**, 705–715 (2011).

6.   K. Zhang, S. Shi, W. Han, Research progress in cytokines with chemokine-like function.

Cellular & Molecular Immunology **15**, 660–662 (2018).

7.   K. Chen, *et al.*, Chemokines in homeostasis and diseases. *Cell Mol Immunol* **15**, 324–334

(2018).

8.   Y. Wang, *et al.*, Chemokine-like factor 1 is a functional ligand for CC chemokine receptor 4

(CCR4). *Life Sciences* **78**, 614–621 (2006).

9.   Y. Wang, *et al.*, Two C-terminal peptides of human CKLF1 interact with the chemokine

receptor CCR4. *The International Journal of Biochemistry & Cell Biology* **40**, 909–919 (2008).

10.   D.-D. Liu, *et al.*, Progress in pharmacological research of chemokine like factor 1 (CKLF1).

Cytokine **102**, 41–50 (2018).

11.   Y. Tom Tang, *et al.*, TAFA: a novel secreted family with conserved cysteine residues and

restricted expression in the brain. *Genomics* **83**, 727–734 (2004).

12.   D. C. Sarver, X. Lei, G. W. Wong, FAM19A (TAFA): An Emerging Family of Neurokines with

Diverse Functions in the Central and Peripheral Nervous System. *ACS Chem. Neurosci.* **12**, 945–958

(2021).

13.   W. Wang, *et al.*, FAM19A4 is a novel cytokine ligand of formyl peptide receptor 1 (FPR1) and

is able to promote the migration and phagocytosis of macrophages. *Cell Mol Immunol* **12**, 615–624

(2015).

14.   M. Y. Park, *et al.*, FAM19A5, a brain-specific chemokine, inhibits RANKL-induced osteoclast

formation through formyl peptide receptor 2. *Sci Rep* **7**, 15575 (2017).

15.   C. Zheng, *et al.*, FAM19A1 is a new ligand for GPR1 that modulates neural stem-cell

proliferation and differentiation. *The FASEB Journal* **32**, 5874–5890 (2018).

16.   X. Wang, *et al.*, Cytokine-like 1 Chemoattracts Monocytes/Macrophages via CCR2. *The*

Journal of Immunology **196**, 4090–4099 (2016).

17.   A. Tomczak, M. T. Pisabarro, Identification of CCR2-binding features in Cytl1 by a CCL2-like

chemokine model. *Proteins: Structure, Function, and Bioinformatics* **79**, 1277–1292 (2011).

18.   T. Yoshimura, J. J. Oppenheim, Chemokine-like receptor 1 (CMKLR1) and chemokine (C–C

motif) receptor-like 2 (CCRL2); Two multifunctional receptors with unusual properties. *Experimental*

Cell Research **317**, 674–684 (2011).

19.   R. Bonecchi, G. J. Graham, Atypical Chemokine Receptors and Their Roles in the Resolution

of the Inflammatory Response. *Frontiers in Immunology* **7**, 224 (2016).

20.   T. N. Kledal, M. M. Rosenkilde, T. W. Schwartz, Selective recognition of the membrane-

bound CX3C chemokine, fractalkine, by the human cytomegalovirus-encoded broad-spectrum

receptor US28. *FEBS Letters* **441**, 209–214 (1998).

21.   T. F. Miles, *et al.*, Viral GPCR US28 can signal in response to chemokine agonists of nearly

unlimited structural degeneracy. *eLife* **7**, e35850 (2018).

22.   M. M. Rosenkilde, M. J. Smit, M. Waldhoer, Structure, function and physiological

consequences of virally encoded chemokine seven transmembrane receptors. *British Journal of*

Pharmacology **153**, S154–S166 (2008).

23.   H. Daiyasu, W. Nemoto, H. Toh, Evolutionary Analysis of Functional Divergence among

Chemokine Receptors, Decoy Receptors, and Viral Receptors. *Frontiers in Microbiology* **3** (2012).

24.   M. Meyrath, *et al.*, The atypical chemokine receptor ACKR3/CXCR7 is a broad-spectrum

scavenger for opioid peptides. *Nature Communications* **11**, 3033 (2020).

25.   R. J. B. Nibbs, G. J. Graham, Immune regulation by atypical chemokine receptors. *Nat Rev*

Immunol **13**, 815–829 (2013).

26.   F. Bachelerie, *et al.*, New nomenclature for atypical chemokine receptors. *Nat Immunol* **15**,

207–208 (2014).

27.   S. S. Denisov, CXCL17: The Black Sheep in the Chemokine Flock. *Frontiers in Immunology* **12**,

2811 (2021).

28.   M. E. DeVries, *et al.*, Defining the Origins and Evolution of the Chemokine/Chemokine

Receptor System. *The Journal of Immunology* **176**, 401 (2006).

29.   B. Bajoghli, Evolution and function of chemokine receptors in the immune system of lower

vertebrates. *European Journal of Immunology* **43**, 1686–1692 (2013).

30.   A. Zlotnik, O. Yoshie, The Chemokine Superfamily Revisited. *Immunity* **36**, 705–716 (2012).

31.   H. Nomiyama, *et al.*, Extensive expansion and diversification of the chemokine gene family in

zebrafish: Identification of a novel chemokine subfamily CX. *BMC Genomics* **9**, 222 (2008).

32.   J. F. Fleming, R. Feuda, N. W. Roberts, D. Pisani, A Novel Approach to Investigate the Effect

of Tree Reconstruction Artifacts in Single-Gene Analysis Clarifies Opsin Evolution in Nonbilaterian

Metazoans. *Genome Biology and Evolution* **12**, 3906–3916 (2020).

33.   W. Han, *et al.*, Identification of eight genes encoding chemokine-like factor superfamily

members 1–8 (CKLFSF1–8) by in silico cloning and experimental validation. *Genomics* **81**, 609–617

(2003).

34.   H.-J. Duan, X.-Y. Li, C. Liu, X.-L. Deng, Chemokine-like factor-like MARVEL transmembrane

domain-containing family in autoimmune diseases. *Chinese Medical Journal* **133** (2020).

35.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool.

Journal of Molecular Biology **215**, 403–410 (1990).

36.   S. F. Altschul, *et al.*, Gapped BLAST and PSI-BLAST: a new generation of protein database

search programs. *Nucleic Acids Research* **25**, 3389–3402 (1997).

37.   C. Camacho, *et al.*, BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

38.   T. Frickey, A. Lupas, CLANS: a Java application for visualizing protein families based on

pairwise similarity. *Bioinformatics* **20**, 3702–3704 (2004).

39.   F. Gabler, *et al.*, Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. *Current

Protocols in Bioinformatics* **72**, e108 (2020).

40.   J. L. Maravillas-Montero, *et al.*, Cutting Edge: GPR35/CXCR8 Is the Receptor of the Mucosal

Chemokine CXCL17. *The Journal of Immunology* **194**, 29–33 (2015).

41.   B. Q. Minh, M. A. T. Nguyen, A. von Haeseler, Ultrafast Approximation for Phylogenetic

Bootstrap. *Molecular Biology and Evolution* **30**, 1188–1195 (2013).

42.   D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2: Improving the

Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution* **35**, 518–522 (2018).

43.   F. Lemoine, *et al.*, Renewing Felsenstein's phylogenetic bootstrap in the era of big data.

Nature **556**, 452–456 (2018).

44.   B. Morel, A. M. Kozlov, A. Stamatakis, G. J. Szöllősi, GeneRax: A Tool for Species-Tree-Aware

Maximum Likelihood-Based Gene Family Tree Inference under Gene Duplication, Transfer, and Loss.

Molecular Biology and Evolution **37**, 2763–2774 (2020).

45.   T. A. Williams, *et al.*, The power and limitations of species tree-aware phylogenetics.

2023.03.17.533068 (2023).

46.   H. Nomiyama, N. Osada, O. Yoshie, Systematic classification of vertebrate chemokines based

on conserved synteny and evolutionary history. *Genes to Cells* **18**, 1–16 (2013).

47.   A. Zlotnik, O. Yoshie, H. Nomiyama, The chemokine and chemokine receptor superfamilies

and their molecular evolution. *Genome Biol* **7**, 243–243 (2006).

48.   Z. Sun, *et al.*, The evolution and functional characterization of CXC chemokines and receptors

in lamprey. *Developmental & Comparative Immunology* **116**, 103905 (2021).

49.   A. Le Mercier, *et al.*, GPR182 is an endothelium-specific atypical chemokine receptor that

maintains hematopoietic stem cell homeostasis. *Proceedings of the National Academy of Sciences*

118, e2021596118 (2021).

50.   P. Liò, M. Vannucci, Investigating the evolution and structure of chemokine receptors. *Gene*

317, 29–37 (2003).

51.   R. Fredriksson, M. C. Lagerström, L.-G. Lundin, H. B. Schiöth, The G-Protein-Coupled

Receptors in the Human Genome Form Five Main Families. Phylogenetic Analysis, Paralogon Groups,

and Fingerprints. *Mol Pharmacol* **63**, 1256–1272 (2003).

52.   M. T. Pisabarro, *et al.*, Cutting Edge: Novel Human Dendritic Cell- and Monocyte-Attracting

Chemokine-Like Protein Identified by Fold Recognition Methods. *The Journal of Immunology* **176**,

2069–2073 (2006).

53.   E. J. Weinstein, *et al.*, VCC-1, a novel chemokine, promotes tumor growth. *Biochemical and*

Biophysical Research Communications **350**, 74–81 (2006).

54.   A. M. Najakshin, L. V. Mechetina, B. Y. Alabyev, A. V. Taranin, Identification of an IL-8

homolog in lamprey (Lampetra fluviatilis): early evolutionary divergence of chemokines. *European*

Journal of Immunology **29**, 375–382 (1999).

55.   B. Bajoghli, *et al.*, Evolution of Genetic Networks Underlying the Emergence of Thymopoiesis

in Vertebrates. *Cell* **138**, 186–197 (2009).

56.   L. Pan, J. Lv, Z. Zhang, Y. Zhang, Adaptation and Constraint in the Atypical Chemokine

Receptor Family in Mammals. *BioMed Research International* **2018**, 9065181 (2018).

57.   M. Kasahara, The 2R hypothesis: an update. *Current Opinion in Immunology* **19**, 547–552

(2007).

58.   O. Simakov, *et al.*, Deeply conserved synteny resolves early events in vertebrate evolution.

Nat Ecol Evol **4**, 820–830 (2020).

59.   W. Tang, Y. Li, A. Li, M. E. Bronner, Clonal analysis and dynamic imaging identify

multipotency of individual Gallus gallus caudal hindbrain neural crest cells toward cardiac and

enteric fates. *Nat Commun* **12**, 1894 (2021).

60.   J. R. York, D. W. McCauley, The origin and evolution of vertebrate neural crest cells. *Open*

Biology **10**, 190285 (2020).

61.   R. M. Waterhouse, *et al.*, BUSCO Applications from Quality Assessments to Gene Prediction

and Phylogenomics. *Mol Biol Evol* **35**, 543–548 (2018).

62.   M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO Update: Novel and

Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of

Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution* **38**, 4647–4654 (2021).

63.   E. Boutet, *et al.*, "UniProtKB/Swiss-Prot, the Manually Annotated Section of the UniProt

KnowledgeBase: How to Use the Entry View" in *Plant Bioinformatics: Methods and Protocols*,

Methods in Molecular Biology., D. Edwards, Ed. (Springer, 2016), pp. 23–54.

64.   S. Poux, *et al.*, On expert curation and scalability: UniProtKB/Swiss-Prot as a case study.

Bioinformatics **33**, 3454–3460 (2017).

65.   J. J. Smith, *et al.*, Sequencing of the sea lamprey (Petromyzon marinus) genome provides

insights into vertebrate evolution. *Nat Genet* **45**, 415–421 (2013).

66.   D. Yu, *et al.*, Hagfish genome illuminates vertebrate whole genome duplications and their

evolutionary consequences. 2023.04.08.536076 (2023).

67.   E. M. Zdobnov, R. Apweiler, InterProScan – an integration platform for the signature-

recognition methods in InterPro. *Bioinformatics* **17**, 847–848 (2001).

68.   P. Jones, *et al.*, InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

69.   F. Bachelerie, *et al.*, Chemokine receptors (version 2020.5) in the IUPHAR/BPS Guide to

Pharmacology Database. *IUPHAR/BPS Guide to Pharmacology CITE* **2020** (2020).

70.   The UniProt Consortium, UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic*

Acids Research **51**, D523–D531 (2023).

71.   L. Käll, A. Krogh, E. L. L. Sonnhammer, Advantages of combined transmembrane topology

and signal peptide prediction—the Phobius web server. *Nucleic Acids Research* **35**, W429–W432

(2007).

72.   W. Li, L. Jaroszewski, A. Godzik, Clustering of highly homologous sequences to reduce the

size of large protein databases. *Bioinformatics* **17**, 282–283 (2001).

73.   L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-generation

sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

74.   G. Pándy-Szekeres, *et al.*, GPCRdb in 2023: state-specific structure models using AlphaFold2

and new ligand resources. *Nucleic Acids Research* **51**, D395–D402 (2023).

75.   K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple sequence

alignment based on fast Fourier transform. *Nucleic Acids Research* **30**, 3059–3066 (2002).

76.   K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772–780 (2013).

77.   S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

78.   B. Q. Minh, *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference

in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534 (2020).

79.   S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin, ModelFinder:

fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**, 587–589 (2017).

80.   J. Huerta-Cepas, F. Serra, P. Bork, ETE 3: Reconstruction, Analysis, and Visualization of

Phylogenomic Data. *Molecular Biology and Evolution* **33**, 1635–1638 (2016).

81.   D. H. Erwin, *et al.*, The Cambrian Conundrum: Early Divergence and Later Ecological Success

in the Early History of Animals. *Science* **334**, 1091–1097 (2011).

82.   F. Delsuc, *et al.*, A phylogenomic framework and timescale for comparative studies of

tunicates. *BMC Biol* **16**, 39 (2018).

83.   K. M. Cohen, S. C. Finney, P. L. Gibbard, J.-X. Fan, The ICS International Chronostratigraphic

Chart. *Episodes Journal of International Geoscience* **36**, 199–204 (2013).

Figure Legends

Figure 1: Cluster Analysis and Phylogeny of Ligand groups. A) Similarity-based clustering, using

CLANS, of canonical chemokines and related molecules with sequence similarity. Canonical

chemokines are an independent group from other related molecules (TAFA, CYTL and CXCL17).

Canonical chemokines are composed of two large groups (CC-type and CXC-type) within which some

divergent subgroups are highlighted. The clustering and connections shown are at the p-value

threshold of 1E-6. Other p-values tested are shown in Supplementary Figure S1. Candidate

invertebrate sequences are shown as crosses and further information regarding them can be found

in Supplementary Results. **B)** Maximum-Likelihood un-rooted phylogenetic tree of canonical

chemokines. CC-type and CXC-type are split into two separate clades. Supports for key nodes are

indicated in boxes with Transferable Bootstrap Expectation (TBE) represented by triangles and the

Ultrafast Bootstraps (UFB) as circles. A traffic light colour code is used to indicate the level of

support: high (green); intermediate (yellow) and low (red). **C)** Similarity-based clustering, using

CLANS, of the CKLF super family (CKLFSF). Two major clusters are formed: the smaller "CKLF Group I"

and the heterogenous "CKLF group II" that also includes some invertebrate sequences (shown as

crosses). Subclades, including the known members of the CKLF super family, are highlighted. The

clustering and connections shown are at the p-value threshold of 1E-15, as this is the threshold at

which the two major clusters connect. Other p-values tested are shown in Supplementary Figure S2.

D) Maximum-Likelihood un-rooted phylogenetic tree of the CKLF super family (CKLFSF). The CKLF

group I is monophyletic, while the CKLF group II is not. Supports for key nodes are indicated in boxes

with Transferable Bootstrap Expectation (TBE) represented by triangles and the Ultrafast Bootstraps

(UFB) as circles. A traffic light colour code is used to indicate the level of support: high (green);

intermediate (yellow) and low (red).

Figure 2: Distribution and duplication patterns of ligand groups. A) Presence of all ligand groups are

mapped onto a species tree. Gene trees and duplication events are based on the gene tree to

species tree reconciliation analyses. The nomenclature for canonical chemokines is primarily based

on known chemokines of human (or mouse). Where human and mouse chemokines do not

correspond, the default name refers to the human gene and the mouse (*Mus musculus*) one is

indicated with "Mm". Chemokines that have been classically described as having either homeostatic

or inflammatory function are indicated with a circle or a star respectively. The classification used

here was based on Zlotnik and Yoshie 2012 (30) with the inflammatory type also including

chemokines they described as plasma/platelet types. Overall, canonical chemokines originated in

vertebrates and expanded a first time in jawed vertebrates and a second time in mammals.

Homeostatic chemokines (e.g., CXCL12) are generally more ancient than inflammatory ones. CXCL17

and CYTL are mammal and jawed vertebrate specific respectively. TAFA originated in the common

ancestor of vertebrates and urochordates, while the CKLF super family is present in invertebrates

although key duplications occurred at the base of vertebrates. **B)** Number of complements for each

ligand group at key species nodes are mapped onto the species tree. The number of complements in

each group reflects the pattern of duplications. The major increase occurred at the level of jawed

vertebrates with canonical chemokines undergoing a second significant increase within placentals.

Figure 3: Phylogeny of Receptor groups.

An unrooted maximum likelihood phylogeny of chemokine receptors. The tree shown is the transfer

bootstrap expectation (TBE) tree including just the chordate specific clade from the ultrafast

bootstrap tree (UFB). Node supports from both TBE (triangle) and UFB (circle) shown for equivalent

key nodes in boxes with arrows to indicate node. A traffic light colour code is used to indicate the

level of support: high (green); intermediate (yellow) and low (red). Key clades highlighted: yellow =

chemokine like plus group (CMLplus); blue = intermediate group; green = atypical 3 and GPR182

(ACKR3/GPR182); purple = canonical chemokines (Canonical CKR); and pink = relaxin receptors

(RL3R). Branches scaled by amino acid substitutions per site.

Figure 4: Distribution and duplication patterns of receptor groups.

A) Presence of all receptor groups are mapped onto a species tree. Gene trees and duplication

events are based on the gene tree to species tree reconciliation analyses. The nomenclature for

genes is primarily based on human chemokines. The canonical chemokines had 5 paralogs present in

the vertebrate common ancestor. These undergo a heterogeneous pattern of duplication

throughout vertebrates with different paralogs duplicating different number of times and in

different groups of species. Chemokines that have been classically described as having either

homeostatic or inflammatory function are indicated with a circle or a star respectively. The

classification used here was based on Zlotnik and Yoshie 2012 (30). **B)** Number of complements for

each receptor group at key species nodes are mapped onto the species tree. The number of

complements in each group reflects the pattern of duplications. The chemokine groups diverged in

the vertebrate stem group. The major expansion occurred at the level of jawed vertebrates with

canonical chemokine receptors, the chemokine-like receptor plus group and intermediate groups

increasing in copy number. Canonical chemokine underwent another small subsequent increase

within placentals.

Figure 5: Summary of the evolution of ligand and receptors.

A summary diagram of the evolution of the different chemokine system components. A simplified

phylogenetic tree of species is shown, calibrated to time according to Erwin et al. 2011 (81), Delsuc

et al. 2018 (82)and Cohen et al. 2013 (83). Circles represent ligand groups and 7 transmembrane

domain structure icon represents GPCR groups. Icons are colour coded by group and placed adjacent

to the branch in the species tree where they first appear. X2 and X5 shown to indicate the number of

paralogs present for CXCL ligand group and the canonical CKR groups respectively, on the branch

where they first appear. Question mark refers to the uncertainty regarding the origin of the CKLF

group I in jawed vertebrates or deuterostome stem group (see Figure 2). Geological column is shown

along the bottom, in accordance with the ICS International Chronostratigraphic Chart.

Supplementary Tables:

Table S1. Summary of canonical and non-canonical chemokine system components.

Table S2. List of the species used in this work and their respective BUSCO values and accession

number.

Table S3. Substitution models used in the phylogenetic analyses.

Supplementary Figure captions:

Figure S1. CLANS clustering of chemokines and related molecules sequences. Initial identification

and annotation of clusters was performed at the strict p-value of 1E-35 **(A)**. Subsequent loosening of

the p-value clarified the relationships across clusters and defined bigger groups. At p-value 1E-15 **(B)**

two major canonical chemokine groups are well-defined: the CCL group, which includes also XCL and

X3CL; and the CXCL group. At this level of stringency, only few canonical chemokines remain

isolated: CCL27/28; CXCL12; CXCL14; CXCL16; as well as CXCL17. TAFA and CYTL are also isolated. At

p-value 1E-10 **(C)** the two major chemokine groups connect to each other. CCL27/28 is connected to

the CCL group and CXCL12 and CXCL14 are connected to the CXCL group, while CXCL16 and CXCL17

are still isolated. At p-value 1E-6 **(D)** all chemokine groups are connected in one big cluster, except

for CXCL17. TAFA and CYTL are also still isolated. Crosses indicate the few invertebrate sequences

that were collected from the BLAST search, more information in Supplementary Results.

Figure S2. CLANS clustering of CKLF Super Family sequences. Initial identification and annotation of

clusters was performed at the strict p-value of 1E-60 **(A)**. Subsequent loosening of the p-value

clarified the relationships across clusters and defined bigger groups. At 1E-20 **(B)** two major clusters

have formed. One, that we called CKLF group I, includes CKLF, CMTM1,2,3,5, and PLP2. The other,

that we called CKLF group II, includes CMTM4/6,7,8, and other groups. At 1E-16 **(C)** more sequences

have joined the two major groups that are still separate. At 1E-15 **(D)** the two major groups connect,

as well as few extra sequences, see Supplementary Results for extra details. Crosses indicate

invertebrate sequences.

Figure S3. Alignment of candidate brachiopod CCL24 sequence with mammalian CCL24s. Our BLAST

searches picked up a sequence from the brachiopod *Lingula unguis* that when re-blasted versus

SwissProt returned a CCL24 as hit. Alignment of the brachiopod sequence with mammalian CCL24

sequences reveals a poor overall conservation, with the brachiopod sequence also being significantly

longer than any of the other sequences. Further details about this sequence can be found in

supplementary file S3 and in Supplementary Results.

Figure S4. Alignment of candidate cnidarian CCL3 sequence with mammalian CCL3s. Our BLAST

searches picked up a sequence from the cnidarian *Clytia hemisphaerica* that when re-blasted versus

SwissProt returned a CCL3 as hit. Alignment of the cnidarian sequence with mammalian CCL3

sequences reveals a poor overall conservation, with the cnidarian sequence being extremely longer

than any of the other sequences. Further details about this sequence can be found in supplementary

file S3 and in Supplementary Results.

Figure S5. Alignment of candidate echinoderm CXCL10 sequence with mammalian CXCL10s. Our

BLAST searches picked up a sequence from the echinoderm *Acanthaster planci* that when re-blasted

versus SwissProt returned a CXCL10 as hit. Alignment of the echinoderm sequence with mammalian

CXCL10 sequences reveals a poor overall conservation, with the brachiopod sequence also being

significantly longer than any of the other sequences. Further details about this sequence can be

found in supplementary file S3 and in Supplementary Results.

Figure S6. Alignment of 4 candidate urochordate TAFA sequences with vertebrate TAFAs. Our

BLAST searches picked up 4 sequences from the urochordate *Ciona intestinalis*, that connected with

the TAFA cluster in the CLANS analysis. One of these sequences when blasted versus SwissProt

returned a TAFA as hit. This sequence was also annotated as TAFA by InterProScan. Alignment of the

urochordate sequences with vertebrate TAFA sequences reveals that only the one annotated as

TAFA aligns well. While the other 3 align poorly and are also significantly longer than any of the

other sequences. Further details about these sequences can be found in supplementary file S3 and

in Supplementary Results.

Figure S7. Alignment of best candidate urochordate TAFA sequence with vertebrate TAFAs. Of the

4 urochordate candidate TAFA sequences, only one was annotated as TAFA with both SwissProt and

InterProScan annotation and appeared to align well with other TAFAs with a preliminary alignment

with all urochordate sequences (Figure S6). Here we removed the other 3 urochordate sequences

and aligned only the best candidate with the vertebrate TAFAs. The sequence conservation is even

more apparent with this alignment. Importantly, 8 of the 10 typical cysteine residues of TAFA1-4 are

conserved, and the two missing cysteines are the same ones missing in TAFA5. Further discussion

can be found in Supplementary Results.

Figure S8. Unrooted phylogenetic tree of canonical chemokines with TBE supports. Phylogenetic

tree under the model GTR20+F+R4. Nodal support is calculated from 100 non-parametric bootstrap

repeats with transferable bootstrap expectation. CCL clade is in orange, CXCL clade in blue.

Figure S9. Unrooted phylogenetic tree of canonical chemokines with UFB supports. Phylogenetic

tree under the model GTR20+F+R4. Nodal support is calculated from 1000 ultrafast bootstrap

repeats. CCL clade is in orange, CXCL clade in blue.

Figure S10. Unrooted phylogenetic tree of TAFA with TBE supports. Phylogenetic tree under the

model JTT+R5. Nodal support is calculated from 100 non-parametric bootstrap repeats with

transferable bootstrap expectation.

Figure S11. Unrooted phylogenetic tree of TAFA with UFB supports. Phylogenetic tree under the

model JTT+R5. Nodal support is calculated from 1000 ultrafast bootstrap repeats.

Figure S12. Unrooted phylogenetic tree of CYTL with TBE supports. Phylogenetic tree under the

model JTT+I+G4. Nodal support is calculated from 100 non-parametric bootstrap repeats with

transferable bootstrap expectation.

Figure S13. Unrooted phylogenetic tree of CYTL with UFB supports. Phylogenetic tree under the

model JTT+I+G4. Nodal support is calculated from 1000 ultrafast bootstrap repeats.

Figure S14. Unrooted phylogenetic tree of CXCL17 with TBE supports. Phylogenetic tree under the

model JTT. Nodal support is calculated from 100 non-parametric bootstrap repeats with transferable

bootstrap expectation.

Figure S15. Unrooted phylogenetic tree of CXCL17 with UFB supports. Phylogenetic tree under the

model JTT. Nodal support is calculated from 1000 ultrafast bootstrap repeats.

Figure S16. Unrooted phylogenetic tree of CKLFSF with TBE supports. Phylogenetic tree under the

model GTR20+F+R7. Nodal support is calculated from 100 non-parametric bootstrap repeats with

transferable bootstrap expectation. Red clade = CMTM4/6; blue clade = CKLF I group; green clade =

CMTM7; turquois clade = MAL/MALL/MAL2.

Figure S17. Unrooted phylogenetic tree of CKLFSF with UFB supports. Phylogenetic tree under the

model GTR20+F+R7. Nodal support is calculated from 1000 ultrafast bootstrap repeats. Red clade =

CMTM4/6; blue clade = CKLF I group; green clade = CMTM7; turquois clade = MAL/MALL/MAL2.

Figure S18. Rooted species tree reconciled gene tree for canonical chemokines. The canonical

chemokines gene tree was reconciled with the species tree using GeneRax. "S" or "D" at the node

indicates a speciation or duplication event respectively. CCL clade is in orange, CXCL clade in blue.

Figure S19. Rooted species tree reconciled gene tree for TAFA. The TAFA gene tree was reconciled

with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or duplication

event respectively.

Figure S20. Rooted species tree reconciled gene tree for CYTL. The CYTL gene tree was reconciled

with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or duplication

event respectively.

Figure S21. Rooted species tree reconciled gene tree for CXCL17. The CXCL17 gene tree was

reconciled with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or

duplication event respectively.

Figure S22. Rooted species tree reconciled gene tree for CKLFSF. The CKLFSF gene tree was

reconciled with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or

duplication event respectively. Red clade = CMTM4/6; blue clade = CKLF I group; green clade =

CMTM7; turquois clade = MAL/MALL/MAL2.

Figure S23. CLANS clustering of receptors and related molecules sequences. A CLANS clustering

layout where shapes indicate sequences and lines are connections indicating similarity between

sequences at or surpassing the p-value similarity threshold. Sequences are positioned in clusters

based on similarity. Initial identification and annotation of clusters was performed using the inbuilt

convex clustering at the p-value of 1E-100. **(A)** clustering was loosened till the canonical receptor

annotated groups formed a cluster at 1E-65. **(B)** loosening of the p-value to 1E-60 identified

relationships between clusters of interest and identified the intermediate group as connecting to

both canonical and chemokine-like plus groups. All sequences connected to groups of interest are

vertebrate sequences. **(C)** further loosening to p-value 1E-50 connects the vertebrate sequences of

interest to a large cluster of sequences which contains vertebrate and invertebrate sequences which

are annotated as opioid and somatostatin receptors and other GPCRs. Crosses indicate invertebrate

sequences and Y-shape indicates the reference viral sequences included. Shapes are colour coded by

group of interest: purple = canonical chemokine receptors; yellow = chemokine-like plus; green =

atypical receptor 3/GPR182; blue = intermediate group; pink = relaxin receptors.

Figure S24. Unrooted phylogenetic tree of receptors with TBE supports. Phylogenetic tree of

receptor sequences of interest and putative outgroups under the model GTR20+F+G4. Sequences

used are a subset of sequences extracted from CLANs, specifically they are those in the chordate

specific clade in the ultrafast bootstrap tree. Nodal support is calculated from 100 non-parametric

bootstrap repeats with transferable bootstrap expectation. Branches colour coded by group of

interest: purple = canonical chemokine receptors; yellow = chemokine-like plus; green = atypical

receptor 3/GPR182; blue = intermediate group; pink = relaxin receptors.

Figure S25. Unrooted phylogenetic tree of receptors with UFB supports. Phylogenetic tree of all

receptor sequences of interest and putative outgroups extracted from clans under the model

GTR20+F+G4. Nodal support is calculated from 1000 ultrafast bootstrap repeats. Branches colour

coded by group of interest: purple = canonical chemokine receptors; yellow = chemokine-like plus;

green = atypical receptor 3/GPR182; blue = intermediate group; pink = relaxin receptors.

Figure S26. Rooted species tree reconciled gene tree for receptors. The ultrafast bootstrap receptor

tree was modified to extract the subtree of the chordate specific clade. This gene tree was

reconciled with the species tree using GeneRax. "S" or "D" at the node indicates a speciation or

duplication event respectively. Branches colour coded by group of interest: purple = canonical

chemokine receptors; yellow = chemokine-like plus; green = atypical receptor 3/GPR182; blue =

intermediate group; pink = relaxin receptors.

Fig. 1

Fig. 2

Fig. 3

A

TBE
△ > 0.9
△ 0.75 – 0.9
△ < 0.75

UFB
● > 90%
● 75 – 90%
● < 75%

Cephalochordates   Urochordates

RL3R

Chemokine-like receptors
CML1; FPR; GPR1 (CML2); C3AR;
C5AR

Canonical CKRs
CC, CXC, CX3C and XC type
receptors; ACKR2; ACKR4; CCRL2

Intermediate group
AGTR; APJ; BKRB2; GPR12; GPR25

0.6

ACKR3/GPR182

Fig. 4

Fig. 5