Title: A Novel Pan-RAS Inhibitor with a Unique Mechanism of Action Blocks Tumor

Growth in Mouse Models of GI Cancer

Authors: [1]Jeremy B. Foote, [2]Tyler E. Mattox, [3,12]Adam B. Keeton,[3,12]Xi Chen, [3]Forrest

Smith, [3]Kristy L. Berry, [3]Thomas Holmes, [3]Junwei Wang, [3]Chung-Hui Huang, [2]Antonio

B. Ward, [3]Amit K. Mitra, [2]Veronica Ramirez-Alcantara, [1]Cherlene Hardy, [4]Karrianne G.

Fleten; [4]Kjersti Flatmark; [5]Karina J. Yoon, [6]Sujith Sarvesh, [6]Ganji Purnachandra

Nagaraju, [6]Dhana Sekhar Reddy Bandi, [3]Yulia Y. Maxuitenko, [3]Jacob Valiyaveettil,

[6]Julienne L. Carstens, [7]Donald J. Buchsbaum, [8]Jennifer Yang, [9]Gang Zhou, [10]Elmar

Nurmemmedov, [10]Ivan Babic, [11]Vadim Gaponenko, [11]Hazem Abdelkarim, [12]Michael R.

Boyd, [13]Gregory S. Gorman, [14]Upender Manne, [15]Sejong Bae, [6]Bassel F. El-Rayes, and

[3,12]Gary A. Piazza

Affiliations: [1]Department of Microbiology, University of Alabama at Birmingham,

Birmingham AL, [2]University of South Alabama, Mobile, AL; [3]Department of Drug

Discovery and Development, Harrison College of Pharmacy, Auburn University, Auburn,

AL;[4] Department of Gastroenterological Surgery, Oslo University Hospital, The Radium

Hospital, Oslo, Norway and Institute of Clinical Medicine, Faculty of Medicine, University

of Oslo, Oslo, Norway; [5]Department of Pharmacology and Toxicology, University of

Alabama at Birmingham, Birmingham, AL; [6]Department of Hematology and Oncology,

University of Alabama at Birmingham, Birmingham, AL; [7]Department of Obstetrics and

Gynecology, University of Alabama at Birmingham, Birmingham, AL; [8]University of

Connecticut, Mansfield, CT; [9]Georgia Cancer Center, University of Augusta, Augusta,

GA; [10]Celaris Bio LLC, San Diego, CA; [11]Department of Biochemistry and Molecular

Genetics, University of Illinois, Chicago, IL; [12]ADT Pharmaceuticals LLC, Orange

Beach, AL; [13]Department of Pharmaceutical, Social and Administrative Sciences,

McWhorter School of Pharmacy, Samford University; Birmingham, AL; [14]Department of

Pathology, University of Alabama at Birmingham, Birmingham, AL; [15]Division of

Preventive Medicine, University of Alabama at Birmingham, Birmingham, AL.

Running Title: *A novel pan-RAS inhibitor for RAS-driven GI cancers*

Key Words: KRAS, pan-RAS inhibitor, pancreatic cancer, colorectal cancer, tumor

immune microenvironment

Funding Sources: NIH/NCI R01CA254197, R01CA238514, P30CA01314, and

Norwegian Cancer Society 197837.

Corresponding Author: Gary A. Piazza, Ph.D., W.W. Walker Endowed Professor,

Department Head, Drug Discovery and Development; Director, Cancer Research

Center, Harrison College of Pharmacy, Auburn University, Auburn, AL, Walker Building,

362 W. Thach Ave, Room 3306F, 334-844-7409, gap0034@auburn.edu

Disclosure of Potential Conflicts of Interest: G.A. Piazza, A.B. Keeton, X. Chen, and

M.R. Boyd are co-founders of ADT Pharmaceuticals Inc. and consultants. All other

authors declare no potential conflicts of interest.

ABSTRACT

Here, we describe a novel pan-RAS inhibitor, ADT-007, that potently inhibited the growth of RAS mutant cancer cells irrespective of the RAS mutation or isozyme. RAS^{WT} cancer cells with GTP-activated RAS from upstream mutations were equally sensitive. Conversely, RAS^{WT} cancer cells harboring downstream BRAF mutations and normal cells were essentially insensitive to ADT-007. Sensitivity of cancer cells to ADT-007 required activated RAS and dependence on RAS for proliferation, while insensitivity was attributed to metabolic deactivation by UDP-glucuronosyltransferases expressed in RAS^{WT} and normal cells but repressed in RAS mutant cancer cells. ADT-007 binds nucleotide-free RAS to block GTP activation of effector interactions and MAPK/AKT signaling, resulting in mitotic arrest and apoptosis. ADT-007 displayed unique advantages over mutant-specific KRAS and pan-KRAS inhibitors, as well as other pan-RAS inhibitors that could impact *in vivo* antitumor efficacy by escaping compensatory mechanisms leading to resistance. Local administration of ADT-007 showed robust antitumor activity in syngeneic immune-competent and xenogeneic immune-deficient mouse models of colorectal and pancreatic cancer. The antitumor activity of ADT-007 was associated with the suppression of MAPK signaling and activation of innate and adaptive immunity in the tumor immune microenvironment. Oral administration of ADT-007 prodrug also inhibited tumor growth, supporting further development of this novel class of pan-RAS inhibitors for RAS-driven cancers.

SIGNIFICANCE

ADT-007 has unique pharmacological properties with distinct advantages over other RAS inhibitors by circumventing resistance and activating antitumor immunity. ADT-007 prodrugs and analogs with oral bioavailability warrant further development for RAS-driven cancers.

INTRODUCTION

Colorectal cancer (CRC) and pancreatic ductal adenocarcinoma (PDA) are projected to cause 53,010 and 51,750 deaths in the US during 2024, respectively [1]. The 5-year survival rates for CRC and PDA are estimated to be 64% and 13%, respectively [1]. Over 50% of CRC and 90% of PDA patients harbor gain-in-function mutations in KRAS genes that are associated with poor prognosis, making the development of novel RAS inhibitors an urgent unmet medical need [2].

For many cancers, RAS mutations initiate tumorigenesis in cooperation with the acquisition of mutations in other oncogenes or inactivating mutations in tumor suppressor genes such as p53 [3], SMAD4 [4], APC [5], or p16 [6], driving the progression from precancerous lesions to malignant disease. Signaling through mutant RAS or activated wild-type RAS (RAS^{WT}) is critical for transformation [7], cancer cell proliferation and survival [7, 8], chemoresistance [9], and metastasis [5, 10]. RAS mutations also promote immunosuppression within the tumor immune microenvironment (TiME) through various mechanisms, including increased acute inflammatory cytokines (IL-6, IL-8) [11], priming of immunosuppressive T-regs [12], MDSCs [13], downregulation of MHCI [14], increased

expression of programmed cell death ligand 1 (PD-L1) [15], and suppression of intra-tumoral interferon (IFN) signaling [16]. Cumulatively, oncogenic RAS-mediated reprograming of the TiME results in weakened recognition and reduced cytotoxic T and NK cell functions supporting tumor progression.

The recent development of mutant-specific KRAS inhibitors directed against cancers harboring KRAS mutations at the G12C and G12D codons renewed optimism for selective targeting of KRAS-driven malignancies [17]. Clinical trials evaluating sotorasib (AMG-510) and adagrasib (MRTX-849) selective for the KRAS[G12C] mutation have reported efficacy for NSCLC patients with mutant KRAS[G12C] but a lower response rate for patients diagnosed with KRAS[G12C] mutant CRC [18], possibly resulting from treatment related adaptive resistance [19] involving new activating mutations in RAS isozymes (K/N/HRAS) or activation of RAS[WT] isozymes from upstream receptor tyrosine kinases (RTK) [20]. In addition, the presence of different RAS mutations in the same tumor and the role of RAS[WT] proteins in oncogenesis [7] expands the potential efficacy advantages of RAS inhibitors capable of inhibiting both mutant RAS and hyper-activated RAS[WT].

Here, we characterize a novel pan-RAS inhibitor, ADT-007, with potential advantages over other RAS inhibitors to circumvent resistance and activate antitumor immunity. ADT-007 emerged from screening a compound library that shares the indene scaffold of the nonsteroidal anti-inflammatory drug (NSAID), sulindac, previously reported to selectively inhibit the formation of RAS mutant tumors in a rat model of chemical-induced mammary tumorigenesis [21] and directly bind RAS to inhibit MAPK signaling, albeit with low potency and confounding cyclooxygenase inhibitory activity [22, 23]. We determined the anticancer activity of ADT-007 using cancer cell lines representative of

the complex mutational landscape of RAS-driven cancers and immune modulatory

potential with mouse tumor models having different immune compositions representative

of the heterogeneity in patients with RAS-driven cancers.

MATERIAL AND METHODS:

<u>Chemical synthesis:</u> ADT-007, (*Z*)-2-(5-fluoro-1-(4-hydroxy-3,5-dimethoxybenzylidene)-

2-methyl-1H-inden-3-yl)-N-(furan-2-ylmethyl)acetamide, and ADT-316, (*Z*)-4-((5-fluoro-

3-(2-((furan-2-ylmethyl)amino)-2-oxoethyl)-2-methyl-1H-inden-1-ylidene)methyl)-2,6-

dimethoxyphenyl 4-methylpiperazine-1-carboxylate hydrochloride, were synthesized as

follows: To a 250-mL round-bottom flask equipped with a gas outlet, (Z)-2-(1-(4-

acetoxy-3,5-dimethoxybenzylidene)-5-fluoro-2-methyl-1H-inden-3-yl) acetic acid (10.3 g,

25.00 mMol) was added to 180 mL of anhydrous dichloromethane, followed by carbonyl

diimidazole (CDI) (4.25 g, 26.25 mMol). The mixture was stirred for 30 min followed by

adding furfuryl amine (2.68g, 26.25 mMol) and anhydrous pyridine (20 mL). The solution

was stirred at room temperature for 48 h and washed with 0.2N HCl (120 mL) and water

(120 mL x 2). The organic layer was concentrated, and the residue dissolved in 300 mL

of methanol, followed by adding ammonium in methanol (7N, 100 mL). The red mixture

was stirred vigorously overnight and filtrated. The solid was dissolved in 400 mL of ethyl

acetate, treated with 1N HCl, washed with brine (150 mL x 3), and dried over anhydrous

sodium sulfate. After filtration, the solution was concentrated and recrystallized from

acetone/ethyl acetate to make ADT-007 as a yellow crystal. The mother liquor was

concentrated and purified by silica gel column chromatography, followed by the above

recrystallization procedure to yield the second batch (9.54 g combined, 85% yield).

Treatment of ADT-007 with 4-methylpiperazine-1-carbonyl chloride in anhydrous

pyridine at 50°C overnight, followed by chromatography purification and acidification,

afforded ADT-316 HCl as a yellow crystal.

Mice: Eight-week-old female C57BL/6 mice (000664), male and female NOD.Cg-

$Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/SzJ (NSG, 00557) and female B6.129S7-$Rag1^{tm1Mom}$/J (Rag 1$^{-/-}$,

02216) mice were purchased from the Jackson Laboratory. Eight-week-old female

Balb/c mice were purchased from Charles River Laboratories (028). Female

B6.Ifng/Thy1.1 KI (IFN-γ.Thy1.1) and B6.Cg-Tg-IL2$^{tm1(eGFP)/Weav}$ (IL-2.GFP) knock-in

cytokine reporter mice were kindly provided by Dr. Casey Weaver (University of

Alabama at Birmingham, Birmingham, AL) [24]. Six-eight-week-old female athymic

foxn1 nude mice were bred at the Department of Comparative Medicine, Oslo University

Hospital (OUH).

Cell lines and patient-derived xenografts: Murine PDA cell lines were derived from

tumor-bearing Kras$^{LSL-G12D/+}$; Trp53$^{LSL-R172H/+}$; Pdx1-Cre; Rosa26$^{YFP/YFP}$ (7160c2,

2838c3) and Kras$^{LSL-G12D/+}$; Trp53$^{L/+}$; Pdx1-Cre (5363) mice and cultured in DMEM

complete media as previously indicated [25]. Human tumor cell lines (BxPC-3, AsPC-1,

CFPAC-1, MIA PaCa-2, PANC-1, SW-1990, HT-29, HCT-116, A549, SW480, MDA-MB-

231, Hs 578T, SK-MEL-2, NCI-H1299, OV-90, H-1975, U-87 MG, SK-OV-3), and the

murine tumor lines (CT26 colon, B16 melanoma, and Lewis lung cancer) were

purchased from ATCC and maintained according to ATCC guidelines. Human OVCAR-

8 cell line was purchased from DCTD Tumor Repository (National Cancer Institute at

Frederick) and maintained according to Tumor Repository guidelines. MKN-1 cells were

purchased from Accegen Biotech (Paoli, PA). Normal human airway epithelial cells

(NHAEC) were purchased from Lonza Bioscience and cultured using bronchial epithelial

growth medium per the supplier's protocol. Normal human colon mucosal (NCM460)

cells were obtained from InCell Corporation and maintained in medium/nutrient mixture

with 10% fetal bovine serum (FBS) for less than 12 passages, according to the

supplier's protocol. The human gall bladder adenocarcinoma expressing KRAS^{G12V} was

derived from an 80-year-old male patient with a history of pancreatic cancer. Diagnosed

initially as PDA, the tumor was reclassified as a gall bladder adenocarcinoma. This

patient-derived xenograft (PDX) model was serially passaged in NSG mice. The CRC

PDX models were established by implanting peritoneal metastasis samples collected at

the time of cytoreductive surgery – hyperthermic intraperitoneal chemotherapy (CRS-

HIPEC) into the peritoneal cavity of nude mice. The PMP-1 model was established from

a patient with an appendiceal primary lesion, while PMCA-1 was from a patient with a

primary mucinous rectal carcinoma [26, 27]. The PC244 model was established from a

peritoneal metastasis specimen from a 69-year-old female with a poorly differentiated

mucinous adenocarcinoma of the cecum. Targeted next-generation sequencing of

PC244 was performed as previously described [28].

Tumor cell transfections: BxPC-3 and HT-29 cells were initially exposed to lentiviral

particles containing a mutant HRAS^{G12V} or *KRAS*^{G12C} construct for 48 h in a 6-well plate

at 37°C in a humidified incubator with 5% carbon dioxide. The media was then removed

and replenished with fresh media containing HA-*HRAS*^{G12V} or HA-*KRAS*^{G12C} lentiviral

particles. After another 48 h, the media was removed, and fresh media containing 5

μg/mL puromycin was added to the well. Surviving cells were expanded in a 10 cm dish,

and expression of the construct was confirmed by western blotting (WB) with an HA tag

antibody (Cell Signaling Technology).

HEK 293FT (Invitrogen, # R70007) cell line was transfected with either pBabe-

puro-HA-KRAS^{G12C} (Addgene, #58901) or pBabe-puro-Empty (Addgene, #1764)

retroviral constructs along with pCL-10A1 retrovirus packaging vector (Novus

Biologicals, # NBP2-29542). Following a 48-h incubation at 37°C, culture supernatants

containing viral particles were collected and mixed with PEG-itTM (SBI, #LV810A-1)

virus precipitation reagent (5X) followed by overnight incubation at 4°C. Supernatants

were centrifuged at 1500 x g for 30 min, and white pellets of retroviral particles

containing either pBabe-puro-HA-KRAS^{G12C} or pBabe-puro-Empty retroviral construct

were collected and resuspended in sterile phosphate-buffered saline (PBS). BxPC-3

cells were transduced with retroviral particles in a 6-well plate using polybrene

transfection reagent (EMD Millipore, TR-1003-G) and incubated for 48 h at 37°C. The

media was replaced with fresh media containing 5 µg/mL puromycin. Surviving cells

were selected, expanded, and confirmed for KRAS^{G12C} expression by WB using an HA

tag antibody (Cell Signaling Technology).

In vitro drug treatments and EGF stimulation: For *in vitro* ADT-007 studies, cell cultures

were incubated with increasing concentrations (0.1, 0.5, 1, 5, 10, 50, 100, 250, 500,

1000, 10000 nM) of ADT-007, AMG-510 (Selleck Chemicals), BI-2865 or RMC-6236

(MedChemExpress), MRTX849 (Selleck Chemicals) or vehicle (DMSO) for 24 – 72 h

depending on the assay. In some experiments, recombinant mouse EGF (Gibco) was

added 10 min before harvesting protein lysates from treated tumor cell lines to assess

the impact of upstream receptor tyrosine kinase signaling on RAS-MAPK and RAS-PI3K

pathways.

<u>Cell growth assays:</u> To determine 2D growth IC_{50} values, 5 x 10^3 tumor cells were

plated in triplicate using 96-well flat-bottom plates. To determine 3D growth IC_{50} values,

1 x 10^3 tumor cells were plated in triplicate using 96-well round bottom non-binding

plates. Cultures were incubated with various concentrations of ADT-007 (0.1. 0.5. 1, 5,

10, 50, 100, 250, 500, 1000, 10000 nM) and cultured for an appropriate period

depending on cellular growth kinetics as indicated in figure legends. At 72-96 h (2D) or

8-10 days (3D spheroid), cells were processed according to the manufacturer's

recommendations to assess intra-cellular ATP levels as a measure of cellular

proliferation and viability using CellTiter-Glo ATP content assay (CTG assay)

(Promega). Nonlinear dose-response curves showing the effects of compound

concentration on colony count were generated using GraphPad Prism (GraphPad

Software).

<u>Mouse liver microsome stability assays:</u> ADT-007 (1 µM) was incubated with mouse

liver microsomes along with a NADPH regeneration solution and UGT cofactors

(glucuronidation) in a phosphate buffer (pH 7.4) at 37°C in a shaking water bath.

Samples were taken at 0, 5, 15, 20, 60 and 120 min, quenched with acetonitrile and

analyzed by LC-tandem mass spectrometry to measure the remaining amount of ADT-

007 as a function of time. Peak area ratios of ADT-007 to an internal standard

(terfenadine) were calculated at each time point and data were expressed as percent of

time zero. Samples at each time point were also analyzed in a Q1 and triggered

independent data acquisition scan to determine the identity of the metabolites. The data

were processed using LightSight metabolite ID software. The primary metabolite was

found to be a mono-glucuronide conjugate of ADT-007.

In vitro intracellular compound analysis: Triplicate samples of 60% confluent HT-29 and

HCT-116 cells were treated with 100 nM ADT-007 for 16h in 15 cm plates. Treatment

medium was collected and snap frozen. Cell pellets were collected by scraping and

weighed in tared tubes. Concentration of ADT-007 and relative ratio of ADT-007

glucuronide (m/z 446+193) were quantified by LC-MS/MS as described above.

Gene expression database analysis: Transcriptomic (RNAseq data) for most of the cell

lines in Table 1 is publicly available in Cancer Cell Line Encyclopedia accessed via the

DepMap Portal [29]. The Student's t-test was used to identify significant differences in

19,221 TPM (transcripts per million) +1 normalized transcripts between the two

categories of cell lines. Due to the modest number of resistant lines available, a

threshold of $p < 0.01$ was chosen, which identified 719 transcripts and were further

reduced to 254 unique transcripts which expressed at least a 5-fold difference.

NMR Spectroscopy: Recombinant KRAS[WT] was dialyzed 2x for 24 h (12 h each in 1L 50

mM Tris-Citrate pH=6.5, 50 mM NaCl, 10 mM β-mercaptoethanol) at 4°C following

treatment with 10 mM EDTA to remove Mg^{2+} and the bound nucleotide. To record [1]H-

[15]N heteronuclear single quantum coherence (HSQC) NMR spectra of nucleotide-free

KRAS (nf-KRAS), 900 MHz and 600 MHz Avance spectrometers (Bruker) equipped with

cryogenic probes were used. Each experiment was performed at 25°C in 50 mM Tris-

Citrate pH=6.5, 50 mM NaCl, 10 mM β-mercaptoethanol containing 10% $^{2}H_{2}O$. The

chemical shift assignments were taken from the Biological Magnetic Resonance Data

Bank (BMRB) database (http://www.bmrb.wisc.edu) using the BMRB accession

numbers 18529, 25114, and 26635. Chemical shift perturbations higher than the

average sum and one standard deviation (SD) were considered statistically significant.

<u>Nucleotide Exchange Assays:</u> The fluorescent GTP analog MANT-GTP was purchased

from Millipore Sigma. For each reaction, 5 µL of 5 µM purified GDP-bound KRAS in

reaction buffer (40 mM Tris-HCl, 100 mM NaCl, 20 mM MgCl$_2$, 1 mg/mL BSA, pH 7.5)

and 2.5 µL of 200 mM EDTA were added to a microcentrifuge tube to make a master

mix. In some reactions, water was used in place of EDTA to control the release of GDP.

7.5 µL of master mix per well was added to a 384-well plate for each reaction.

Compound (5 µL) was added to each respective well at five times the desired

concentration or vehicle in the reaction buffer. After 2.5 min for equilibration at 26°C in a

Bio-Tek Synergy H4 microplate reader, 12.5 µL of 1.5 µM MANT-GTP in reaction buffer

was added to each well for a final concentration of 750 nM MANT-GTP, 1 µM RAS, 20

mM EDTA, and the desired concentration of compound or vehicle. In the case of

compound preincubation, the plate was incubated for 1 h at 26°C before MANT-GTP

addition. Fluorescence was measured at an excitation wavelength of 360 nm and an

emission wavelength of 460 nm in a Bio-Tek Synergy H4 microplate reader. Kinetic

curves were generated with GraphPad Prism.

<u>Cellular thermal stability assays:</u> Experiments to determine ADT-007 binding to RAS in

cells were performed by Cellaris Bio LLC using HCT-116 and HT-29 cells cultured in

RPMI-1640 medium supplemented with 10% FBS. To determine the melting profile of

KRAS, fresh cell lysates were prepared in non-denaturing buffer and cleared *via*

centrifugation at 5,000 rpm for 15 min. Cleared lysates were dispensed into 96-well

PCR (equivalent of approx. 6,000 cells/well/50 µl), then subjected to the temperature

gradient (38-65°C) for 10 min. Subsequently, centrifugation was performed at 14,000

rpm to sediment the unstable protein content. Following WB of the soluble fraction,

bands of stable KRAS were quantified, normalized to DMSO, and plotted using

GraphPad Prism. Nonlinear regression analysis was used to fit a sigmoidal dose-

response curve with variable slope to generate EC_{50} values for ADT-007 binding to

KRAS.

Additional cellular thermal stability experiments used a Micro-Tag enzyme

complementation method. A $KRAS^{G12C}$ Micro-Tag construct was engineered with a 15-

amino acid tag at the N-terminus. The construct was transiently expressed in HEK293

human embryonic kidney cells. Two days post-transfection, cells were suspended,

washed with Tris Buffered Saline (TBS: 20mM Tris HCl pH 7.4, 150mM NaCl), and

resuspended at 1000 cells/μl in TBS and 50,000 cells aliquoted to wells of a PCR plate.

Increasing concentrations of ADT-007 were added and incubated with cells for 45 min

followed by heating for 10 min at 55°C and a 1 min cool-down on ice. An equal volume

of 0.5% Triton-X-100 in TBS was added, and cells lysed for 10 min on ice. Fluorescence

was measured and plotted with the concentration of ADT-007 on a semi-log scale using

GraphPad Prism 9.0 software. Nonlinear regression analysis was used to fit a sigmoidal

dose-response curve with variable slope to generate EC_{50} values for ADT-007 binding

to KRAS.

RAS activation assays (RAS-RBD pulldown assays): Cells were grown to 80 percent

confluence using complete growth medium, then washed with 1 mL of PBS and lysed

on ice for 5 min (25 mM Tris HCl, 150 mM NaCl, 5 mM $MgCl_2$, 1% NP-40, 5% glycerol,

pH 7.2) with Halt Protease and Phosphatase Inhibitor Cocktail (Thermo Fisher

Scientific), then clarified at 16,000 RCF 15 min at 4°C. Protein concentration of the

supernatant was normalized using the DC protein assay (Bio-Rad). 200 µg of whole cell

lysate was incubated with GST-RAF-RBD (Addgene) bound to glutathione agarose

beads at 4°C for 1 h. Bound active RAS was collected by centrifugation at 6,000 RCF

for 30 seconds at 4°C, washed thrice with lysis buffer, then eluted with SDS sample

buffer and subsequently detected by WB with a pan-RAS antibody or isozyme-specific

antibody as described below.

Western blotting: Tumor or cell lysates were generated after treatment with 500 µL

RIPA lysis buffer (Thermo Scientific, 89900) with protease and phosphatase inhibitors

(Thermo Scientific, 78440), boiled for 5 min, and equal protein concentrations were

loaded, and run on a 12% SDS-PAGE. Gels were transferred to a PVDF membrane,

which was blocked with either 5% non-fat dry milk in TBS-T (TBS with 0.1% Tween 20)

or 5% BSA in TBS-T, depending on the blocking agent used for the primary antibodies,

for 1 h at room temperature. Membranes were incubated overnight at 4°C on a rocker

with primary antibodies **(Supplemental Table 1)** according to the manufacturer's

instructions and then developed with HRP-conjugated secondary antibodies (Cell

Signaling Technology) for 1 h at room temperature on an orbital shaker. Membranes

were developed using SuperSignal West Enhanced Chemiluminescent substrate

(Thermo Fisher Scientific) and imaged using a ChemiDoc Imaging System (Bio-Rad).

In vivo efficacy experiments:

Murine tumor models: A single cell suspension of murine CT26 CRC or 2838c3,

7160c2, and 5363 PDA KRAS^{G12D} cells was prepared in PBS and kept on ice until

injection. Syngeneic mouse CRC and PDA tumor cells ($1x10^6$/mouse) were injected

subcutaneously (SQ) into 6-8-week-old Balb/c or C57BL/6 mice, respectively. When

tumors reached an average tumor volume of 150 mm^3, mice began receiving daily intra-

tumoral (10 mg/kg, QD, 7 days/week for 13 days) or peri-tumoral (5 mg/kg, BID, 7

days/week for 13-24 days) treatment with ADT-007. In all experiments tumor growth

was assessed by caliper measurements (2-3/week), and tumor volumes (TV = W x L^2/2)

were calculated throughout the study. No overt signs of toxicity (body weight loss or

organ [lungs, heart, liver, GI tract] pathology) were noted with local administration.

PDX CRC tumor models: mucinous PMCA-1, PMP1, and PC244 PDX colorectal

adenocarcinomas were implanted intraperitoneally (IP) in six-eight-week-old female

athymic foxn1 nude mice. Treatment with ADT-007 (IP, 0.33, 0.67, 1.25 and/or 2.5

mg/kg, BID, 5 days/week for 18-23 total days of treatment) was initiated 24 h after

implantation (n = 5-6 mice/group) and mice were followed by MRI for 24 to 50 days after

implantation (6-31 days post-treatment cessation).

PDX GI tumor model: KRAS^{G12V} PDX gall bladder adenocarcinoma was implanted SQ

in eight-week-old male NSG mice. When tumors reached an average tumor volume of

150 mm^3, mice began receiving daily peri-tumoral (5 mg/kg, BID, 7 days/week for 13

days, n = 5 mice/group) treatment with ADT-007. Tumor growth was assessed by

caliper measurements (2-3/week), and tumor volumes were calculated throughout the

study.

All experiments were randomized but not blinded. For animal studies performed at the

University of Alabama at Birmingham (UAB) and the University of South Alabama all

procedures were performed under approved protocols according to the Institutional

Animal Care & Use Committee (IACUC) guidelines and Animal Resources Program of

UAB. All cell lines and PDX models were tested against an extended panel of excluded

pathogens, including mycoplasma, before implantation into mice. For animal studies

performed at OUH Institute for Cancer Research (Norwegian Radium Hospital, Oslo,

Norway) all procedures and experiments involving animals were approved by the

Norwegian Food Safety Authority (application ID #18209) and were conducted

according to the recommendations of the European Laboratory Animals Science

Association and the ARRIVE guidelines.

Flow Cytometry: Mouse tumors were digested in a 1 mg/mL collagenase IV and 0.1

mg/mL DNAase 1 (Worthington Biochemical) in HBSS for 45 min at 37 °C with

intermittent shaking. Samples were washed with media and filtered through a 70 μM

strainer generating suspensions of single cells. Cells were labeled with primary

fluorophore-conjugated antibodies and a live/dead stain (**Supplemental Table 2**) for 30-

60 min at 4°C, washed, and resuspended in flow buffer for downstream immune cell

analyses (**Supplemental Figures 9A and 10A**). Cytokine expression from single-cell

suspensions of spleen and tumor was quantified as follows: cells were plated at a

concentration of 10^6 cells in 1 mL of RP-10 media and stimulated for 5 h at 37°C using a

1x solution of a Cell Activation Cocktail containing PMA ionomycin, and Brefeldin A

(BioLegend). Conversely, cells were cultured for 5 h at 37°C using a solution containing

5 μg of Brefeldin A (BioLegend) as an unstimulated control. After cell surface staining,

cells were fixed in 4% paraformaldehyde for 45 min at 4°C then washed with 1x

Perm/Wash (BD, 554723). For intracellular staining in tumor and spleen samples, cells

were stained with an antibody cocktail-specific intracellular cytokines (IL-2, IFNγ, TNFα,

and granzyme B) in Perm/Wash solution for 60 min at 4°C. Data were acquired on a

Symphony A5 flow cytometer, with analysis performed using FlowJo version 10.7.2.

Statistical Analyses: GraphPad Prism (version 9.1.2) was used for statistical analyses

and graphical representation with data presented as either means $\pm$ standard deviation

(SD) or standard error of the mean (SEM). Two-tailed Student's or Welch's t-test and

two-way Analysis of Variance (ANOVA) with multiple corrections (Tukey's or Sadik's

method) were performed to determine the statistical significance between groups. To

calculate IC_{50} values, compound concentrations were log-transformed, and data

normalized to controls to generate a *log*(inhibitor) *vs*. response (four-parameter

nonlinear dose-response). A $p < 0.05$ was considered statistically significant.

Data Availability: Data generated in this study will be made available upon request from

the corresponding author.

RESULTS

ADT-007 potently and selectively inhibits the growth of cancer cells with mutated

or activated RAS

An indene library was synthesized and screened for differential growth inhibitory

activity by comparing sensitivity of CRC HCT-116 cells harboring a KRAS[G13D] mutation

vs. HT-29 cells with RAS[WT] but mutant BRAF[V600E]. A lead compound, ADT-007 (**Figure**

1A), and several analogs emerged following chemical optimization for potency and

selectivity. ADT-007 inhibited HCT-116 growth with a 5 nM IC_{50}, while HT-29 cells were

~100-fold less sensitive (**Figure 1B**). RAS-RBD pulldown assays confirmed high

activated RAS (RAS-GTP) levels in HCT-116 cells, while RAS-GTP was undetectable in

HT-29 cells. Despite differences in RAS-GTP levels, HCT-116 and HT-29 cells

depended on MAPK signaling for growth, as both were sensitive RAF, MEK, and ERK

inhibitors (not shown). The requirement of mutant RAS for sensitivity to ADT-007 was

determined by transfecting HT-29 cells with HRAS^{G12V}. ADT-007 treatment of RAS-

transfected HT-29 cells resulted in a 24 nM IC$_{50}$ in growth assays compared with 549

and 730 nM IC$_{50}$ values for parental and vector-control cells, respectively (**Figure 1C**).

A panel of histologically diverse cancer cell lines was also screened for sensitivity

to ADT-007 (**Table 1**). Cancer cell lines with mutations in K/N/HRAS isozymes or

upstream signaling components (e.g., EGFR, PDGFR, NF1) were uniformly sensitive to

ADT-007 with low nM IC$_{50}$ values. Like HT-29 cells, other cancer cell lines with

downstream BRAF mutations were essentially insensitive to ADT-007, as were normal

colonocytes and airway epithelium cells.

KRAS^{G12C} MIA PaCa-2 PDA cells were highly sensitive to ADT-007 with a 2 nM

IC$_{50}$ in growth assays while RAS^{WT} BxPC-3 PDA cells had a 2500 nM IC$_{50}$ (**Table 1 and**

Supplemental Figure 1A). RAS-RBD pulldown assays confirmed high RAS-GTP in

MIA PaCa-2 and three other PDA cell lines that were comparably sensitive to ADT-007,

which harbor KRAS^{G12V} and KRAS^{G12D} (**Supplemental Figure 1B**) and represent the

most prevalent KRAS mutations in PDA patients [30]. By comparison, RAS-GTP levels

in BxPC-3 cells were undetectable. Consistent with results from HT-29 cells,

transfecting BxPC-3 cells with KRAS^{G12C} induced sensitivity to ADT-007 in growth

assays (**Supplemental Figure 1C-D**).

ADT-007 treatment of MIA PaCa-2 cells blocked growth in 2D monolayer or 3D

spheroid cultures with comparable low nM IC_{50} values, resulting in unusually steep

dose-response curves that was a characteristic feature of ADT-007 and related analogs

(**Supplemental Figure 1E-F**). ADT-007 also potently blocked colony formation of

KRAS mutant PDA cell lines without impacting colony formation of RAS^{WT} BxPC-3 cells

(**Supplemental Figure 1G**). In addition, ADT-007 potently and selectively inhibited

colony formation of HCT-116 cells without effects on HT-29 cells (**Supplemental Figure**

1H).

ADT-007 inhibits activated RAS and MAPK/AKT signaling

ADT-007 reduced RAS-GTP and phosphorylated MAPK/AKT components in

HCT-116 cells (**Figure 1D**). ADT-007 also inhibited EGF-stimulated MAPK/AKT

signaling in MIA PaCa-2 cells (**Figure 1E**). ADT-007 reduction of RAS-GTP occurred if

cells were grown in serum, serum-starved, or EGF stimulated (**Supplemental Figures**

2A-C). ADT-007 also decreased RAS-GTP by treating lysates from HCT-116 cells or

$HRAS^{G12V}$ transfected HT-29 cells, suggesting a direct effect of ADT-007 on RAS

protein rather than on RAS signaling (not shown). ADT-007 treatment of MIA PaCa-2

cells serum-starved and stimulated with EGF also reduced levels of GTP-$HRAS^{WT}$ and

GTP-$NRAS^{WT}$ isozymes co-expressed with GTP-$KRAS^{G12C}$ (**Figure 1F**). As expected,

EGF did not impact GTP-$KRAS^{G12C}$ levels but activated GTP-$HRAS^{WT}$ and GTP-

$NRAS^{WT}$ isozymes. Nonetheless, ADT-007 reduced GTP-$KRAS^{G12C}$ to the same extent

as EGF-activated RAS^{WT} isozymes.

ADT-007 induces mitotic arrest and apoptosis

ADT-007 induced G2/M arrest in HCT-116 and MIA PaCa-2 cells following 24 h

treatment at concentrations comparable to growth IC_{50} values while concentrations at

least 100x higher were required to have a similar effect in HT-29 or BxPC-3 cells

(**Supplemental Figure 3A-D**). Time-course experiments revealed that ADT-007

induced apoptosis of MIA PaCa-2 cells following G2/M arrest (**Supplemental Figure**

3E). ADT-007 also induced apoptosis in HCT-116 cells but not HT-29 cells

(**Supplemental Figure 3F**). Mitotic arrest was confirmed by experiments showing that

ADT-007 treatment of HCT-116 cells increased phospho-histone 3B levels, which

coincided with reduced cell number and occurred at concentrations within growth IC_{50}

values (**Supplemental Figure 3G**).

Given previous reports that other indenes bind tubulin and induce mitotic arrest

[31], the effect of ADT-007 on tubulin polymerization was determined using recombinant

α-tubulin. ADT-007 exhibited a weak inhibitory effect on tubulin polymerization at

concentrations ~100-fold above growth IC_{50} values (**Supplemental Figure 3H**). Similar

weak effects on tubulin polymerization were observed by treating RAS mutant cancer

cells with ADT-007 (not shown). Moreover, colchicine, a known tubulin inhibitor, lacked

selectivity to inhibit the growth of KRAS mutant cell lines.

ADT-007 induces cytotoxicity; other RAS inhibitors result in cytostasis

ADT-007 was compared with the $KRAS^{G12C}$ inhibitor, AMG-510, and the pan-

KRAS inhibitor, BI-2865, in growth assays involving MIA PaCa-2 cells. Both AMG-510

and BI-2865 were significantly less potent and caused less growth inhibition than ADT-

007 (**Figure 1G**). Unlike ADT-007, neither compound induced mitotic arrest as

determined by phospho-histone 3B levels (**Supplemental Figure 3G**). Another key

difference that may be relevant to resistance was apparent in experiments using LLC

cells harboring KRAS^{G12C} and NRAS^{Q61H} mutations. LLC cells were found to be

insensitive to AMG-510 and BI-2865 in growth assays but retained full sensitivity to

ADT-007 (**Figure 1H-I**).

We also compared ADT-007 with another pan-RAS inhibitor, RMC-6236. While

RMC-6236 displayed comparable potency as ADT-007 in growth assays involving MIA

PaCa-2 cells, viable cells were observed microscopically after RMC-6236, as well as BI-

2865, treatment, while only rounded condensed cells were present after ADT-007

treatment (**Supplemental Figure 4A**). Consistent with experiments described above

(**Supplemental Figure 3E-F**), ADT-007 induced apoptosis of MIA PaCa-2 cells with a 4

nM EC$_{50}$. However, neither RMC-6236 nor BI-2865 induced apoptosis (**Figure 1J**).

Moreover, colony formation assays revealed that ADT-007 caused near complete

inhibition of colony formation, while BI-2865 and RMC-6236 were appreciably less

effective (**Supplemental Figure 4B-C**). These results show fundamental differences

between ADT-007, which selectively induces apoptosis (cytotoxicity) of RAS mutant

cancer cells, while the growth inhibitory activity of other RAS inhibitors appears to be

limited to cytostasis.

Chemical and biological specificity of ADT-007

The chemical specificity of ADT-007 was evaluated using a closely related

chemical analog, ADT-106, containing a methoxy group in place of the hydroxyl moiety.

ADT-106 lacked the selectivity of ADT-007 to inhibit the growth of MIA PaCa-2 cells

compared with BxPC-3 cells (**Figure 2A-B**). ADT-106 also had reduced potency with a

150 nM IC_{50}. It is also noteworthy from these experiments that the potency of ADT-007

to inhibit MIA PaCa-2 growth paralleled its potency to reduce RAS-GTP levels, while

ADT-106 did not impact RAS-GTP levels. These results show the requirement of the

phenolic hydroxyl moiety for both the selectivity and potency of ADT-007, as well as the

association between growth inhibition by ADT-007 and its ability to suppress RAS-GTP.

Metabolic mechanisms impacting sensitivity to ADT-007

Ex vivo treatment of mouse liver microsomes with ADT-007 followed by LC-

MS/MS analysis of metabolites revealed that the phenolic hydroxyl moiety is rapidly

glucuronidated by UDP-glucuronosyltransferases (UGTs) for which multiple UGT

isozymes exist in various cell types (**Figure 2C-D**). These results suggest that UGTs

could contribute to the sensitivity of cells to ADT-007 both *in vitro* and *in vivo*.

Measurement of UGT1 revealed higher protein levels in RAS^{WT} HT-29 and BxPC-3 cells

compared with KRAS mutant HCT-116 and MIA PaCa-2 cells (**Figure 2E-F**).

Transcriptomic data (RNAseq) analysis (Cancer Cell Line Encyclopedia) for mRNA

levels of six distinct UDP-glucuronosyltransferase isozymes (UGT1A1, 1A6, 1A10, 2A1,

2B10, and 2B15) in most cancer cell lines from **Table 1** [29] provided additional

evidence that RAS^{WT} cancer cell lines express higher levels of UGT isozymes

compared with RAS mutant cancer cell lines (**Table 1, Supplemental Table 3**).

Consistent with these differences, measurement of ADT-007 levels by LC-

MS/MS from cells treated overnight with ADT-007 resulted in 350-fold higher

intracellular levels in HCT-116 cells than in HT-29 cells (**Figure 2G**). We confirmed that

UGTs selectively deactivated ADT-007 in RAS^{WT} cancer cells but not in RAS mutant

cells by detection of glucuronide-conjugated ADT-007 in HT-29 cells as none was

detected in HCT-116 cells. In other experiments, we determined if exposure of BxPC-3

cells with high UGTs to ADT-007 could metabolize ADT-007 by collecting "conditioned

medium" (CM) following 24 h treatment with ADT-007. CM from BxPC-3 cells did not

inhibit MIA PaCa-2 growth, while CM from MIA PaCa-2 cells caused comparable growth

inhibition as freshly prepared ADT-007 (**Figure 2H**). Similar results were obtained using

CM from HT-29 cells and testing for growth inhibition of HCT-116 cells (not shown).

Additional experiments showed that pretreatment with propofol, a UGT inhibitor,

increased the sensitivity of BxPC-3 and HT-29 cells to ADT-007 in growth assays

without significantly affecting the sensitivity of MIA PaCa-2 or HCT-116 cells

(**Supplemental Table 4**).

ADT-007 binds RAS in cells

The ability of ADT-007 to directly bind KRAS in cells was measured using cellular

thermal stability assays based on the principle that a protein maintains thermal stability

when in a conformation favorably for binding to a small molecule inhibitor. Initial

experiments measured the optimal temperature to denature 75% (T_{agg75}) KRAS in whole

cell lysates, followed by WB quantification of non-denatured KRAS levels. Treatment of

HCT-116 lysates with ADT-007 increased KRAS thermal stability (T_{agg75}), resulting in a

0.45 nM EC_{50} (**Figure 3A**). By comparison, ADT-007 did not affect thermal stability of

KRAS in HT-29 cells. ADT-007 also did not impact thermal stability of an unrelated

protein, PDE10A, in either HCT-116 or HT-29 cells known to be expressed in both cell

lines [32] (not shown). Additional experiments involving ADT-007 treatment of intact

HEK293 embryonic kidney cells transfected with epitope-tagged mutant KRAS^{G12C} and

detection of thermal stability by WB analysis with anti-Micro-Tag antibody resulted in an

8.49 nM EC$_{50}$ (**Figure 3B**). Consistent with experiments described above (**Figure 2A-**

B) showing the importance of the phenolic hydroxyl moiety for potency and RAS

selectivity, a trimethoxy ADT-007 analog did not impact thermal stability of KRAS (not

shown).

Molecular modeling studies

Molecular docking studies using human GDP-bound KRAS^{WT} (PDB ID 4OBE)

[33] were conducted using molecular dynamics and induced-fit to identify potential

binding sites for ADT-007 to bind RAS. In simulations of Mg^{+2} present or absent, the

best GLIDE docking scores (-11.08 and -12.06, respectively) were calculated with ADT-

007 binding to a site partially overlapping the nucleotide-binding domain that was

associated with an altered conformation of the switch-1 region **(Supplemental Figure**

5A-D). These results include the use of explicit water molecules that were especially

critical for ADT-007 binding to 4OBE (KRAS^{WT} Mg^{+2} present). When Mg^{+2} was present,

the phenolic hydroxyl moiety of ADT-007 was predicted to H-bond with Ser17 through a

water bridge and with Glu31 through the amide side chain projecting into solvent space.

If Mg^{+2} was absent, the phenolic hydroxyl was predicted to interact with Asp119 via a

water bridge and Ala146 by a direct H-bond. Whether Mg^{+2} was present or absent, the

GLIDE docking score of GDP to bind RAS (-9.6 and -8.80, respectively) was

considerably less favorable than ADT-007 (not shown). These results suggest that a

conformational change induced by the binding of ADT-007 may decrease the affinity of

RAS to bind nucleotide since in the docking score for GDP to bind native RAS is -15.15.

The results also indicate that there may be a difference in how ADT-007 binds RAS

depending on whether RAS is in a Mg^{+2} bound (nucleotide-loaded) conformation or

Mg^{+2} free conformation as an apoenzyme referred to as nucleotide-free RAS (nf-RAS).

546 ADT-007 inhibits GTP activation of recombinant nf-RAS

547  We next determined if ADT-007 disrupts RAS-GTP effector binding using

recombinant full-length KRAS^{WT}. KRAS was treated with EDTA to chelate Mg^{+2},

simulating GEF removal of GDP, to generate nf-RAS. ADT-007 treatment of nf-RAS

before GTP addition inhibited RAS-RAF binding as determined by RAS-RBD pulldown

assays at concentrations comparable to growth IC_{50} values and to an extent

approaching the RAS-GDP control (**Figure 3C**). There was a negligible effect of ADT-

007 on binding if RAS was preloaded with GTP (**Figure 3D**). A fluorescent GTP analog

(Mant-GTP) nucleotide exchange assay was used to determine if ADT-007 disrupts

GTP binding to KRAS^{WT}. ADT-007 reduced the rate of GTP binding to RAS if

pretreated with EDTA to chelate Mg^{2+} (**Figure 3E**), but not after adding non-

hydrolysable GTPγS (**Figure 3F**). The pan GTPase inhibitor, CID1067700, inhibited

Mant-GTP binding to RAS to the same extent as ADT-007 (not shown) [34].

559 ADT-007 binds recombinant nf-RAS

560  Nuclear magnetic resonance (NMR) spectroscopy was used to confirm binding of

ADT-007 to recombinant full-length KRAS^{WT}. These experiments were conducted by

chelating Mg^{2+} with EDTA to disassociate GDP and generate nf-RAS, producing a

sample with multiple NMR spectral differences from GDP-KRAS (**Figure 4A**). Removal

of GDP included attenuated signals for G13, a hotspot mutation site in the P-loop [35],

and K16 (adjacent to S17) critical for Mg^{2+} binding [36], which could be readily assigned

due to their unique resonance frequencies. Treatment of nf-RAS with ADT-007 resulted

in several discrete changes in HN signals, including attenuated signals from G13 and

K16 residues (**Figure 4B**). Consistent with biochemical experiments, the NMR spectra

of GDP-KRAS^{WT} in the presence of Mg^{2+} showed no significant differences by ADT-007

treatment, indicating a low likelihood of ADT-007 binding to nucleotide-loaded RAS (not

shown).

ADT-007 inhibits tumor growth and RAS signaling *in vivo*

As described above, *ex vivo* treatment of mouse liver microsome with ADT-007

identified a metabolic vulnerability of ADT-007 to be rapidly metabolized by

glucuronidation (**Figure 2C-D**) prompting experiments to evaluate the antitumor activity

of ADT-007 by localized delivery to minimize liver glucuronidation. Localized delivery

(peri-tumorally) of ADT-007 completely arrested tumor growth and induced partial

regression in the KRAS^{G12V} gall bladder adenocarcinoma PDX model (**Figure 5A**) and

reduced tumor growth in a KRAS^{G12C} PDA PDX model (not shown). Intraperitoneal

injection of ADT-007 significantly reduced growth in three peritoneal metastasis PDX

models bearing mutant KRAS^{G12A} (PMCA-1) (**Figure 5B**), KRAS^{G12D} (PMP1) (**Figure**

5C), or KRAS^{WT} and PDGFR^{T922I} (PC244) (**Figure 5D**). PMCA-1 was the most sensitive

to ADT-007 with the growth inhibition persisting long after treatment discontinuation,

and two mice tumor-free on day 100. Consistent with *in vitro* experiments described

above showing that RAS^{WT} cancer cell lines with upstream mutations are sensitive to

ADT-007 (**Table 1**), the PC244 PDX model expressing RAS^{WT} isozymes, but a

suspected activating mutation in PDGFR, was sensitive to ADT-007 *in vivo.* Reductions

in mucin production were notable by MRI (**Supplemental Figure 6A**) and tumor volume

measurements in each model (**Supplemental Figure 6B-D**).

The antitumor activity of ADT-007 was also determined against an array of

syngeneic CRC and PDA tumor models bearing KRAS^{G12D} mutations grown SQ in

immune-competent mice. When administered locally by intra-tumor (IT) injection, ADT-

007 (10 mg/kg, QD) strongly inhibited SQ tumor growth (T/C$_{max}$ of 27%) in the KRAS

mutant CT26 colon tumor model (**Figure 5E, left panel**) with an 80% reduction in mean

tumor volume at the end of the treatment compared to sham-treated mice (**Figure 5E,**

right panel). The extent of tumor growth inhibition by ADT-007 in each mouse was

closely associated with the extent of reduction of phosphorylated RAF, MEK, and ERK

(**Supplemental Figure 7**). Peritumoral administration of ADT-007 (5 mg/kg, BID)

induced growth arrest in previously characterized syngeneic mouse PDA cell lines

2838c3 (**Figure 5F**) and 7160c2 (**Figure 5G**) [25], and tumor regressions in a KPC-

derived PDA cell line 5363 (**Figure 5H**).

Although ADT-007 was rapidly metabolized by the liver, the compound was

stable in mouse plasma (**Supplemental Figure 8A**). We therefore synthesized a

carbamate prodrug of ADT-007, ADT-316, to protect the vulnerable hydroxyl group from

glucuronidation (**Supplemental Figure 8B**). Oral administration of ADT-316 generated

plasma levels of ADT-007 appreciably higher than growth inhibitory IC$_{50}$ values for at

least 8 h following a single dose (**Supplemental Figure 8C**). While PK analysis of

plasma levels of ADT-316 and ADT-007 revealed modest conversion of the prodrug to

ADT-007, ADT-316 significantly inhibited tumor growth in the CT26 murine CRC model

at a well-tolerated dosage (100 mg/kg, bid) (**Supplemental Figure 8D**). Given that the

incomplete conversion of ADT-316 to ADT-007 likely accounts for its modest antitumor activity, these results support further development of prodrugs to generate higher or more sustained systemic ADT-007 levels.

ADT-007-mediated inhibition of KRAS^{G12D} alters the tumor immune microenvironment (TiME)

To determine the impact of ADT-007 on tumor immunity in mutant KRAS^{G12D} tumors, we assessed the activity of ADT-007 in C57BL/6 recipient mice using 2 PDA tumor lines bearing KRAS^{G12D}. Bot cell lines (2838c3 and 7160c2) has been previously characterized as tumor models with increased T cell infiltrate, yet exhibiting different abundances of myeloid subsets (**Supplemental Figure 9A-B**), basal levels of T cell activation, and function [25]. At treatment termination, ADT-007 decreased the proportion of YFP$^+$ tumor cells in both models (**Figures 6A and 6I**). ADT-007 treatment increased proportions of gMDSC and mMDSC subsets in 7160c2 (**Figure 6B**), but not in 2838c3 tumor-bearing mice, which exhibited a significant decrease in macrophages (**Figure 6J**) in response to treatment. No changes in dendritic cell subset numbers were noted in either model (**Figures 6C**,*7160c2* and **6K**,*2838c3*). Analysis of macrophage phenotypes in both models revealed trends for decreasing MHCII and CD206 expression (**Supplemental Figures 9C**,*2838c3* and **9D**,7160c2).

Efficacy ADT-007 was partially dependent upon T and B cells, as its efficacy was reduced in immune-deficient C57BL/6 Rag 1$^{-/-}$ *vs.* C57BL/6 mice (**Figures 6D**,*7160c2* and **6L**,*2838c3*). Both models contained similar numbers of T cells prior to treatment (**Supplemental Figure 10A-B**). ADT-007 treatment increased numbers of γδ, NKT, and NK cells, whereas CD4$^+$ and CD8$^+$ T cell numbers remained unchanged within the TiME

in both models (**Figures 6E-F***,7160c2* **and 6M-N***,2838c3*). The activation status of T cell

subsets (**Supplemental Figure 10A**) in the 7160c2, but not 2838c3 TiME, shifted to a

CD44^{hi} CD62L^{-}effector phenotype with ADT-007 treatment (**Supplemental Figure 10C-**

D) and PD-1 increased on CD4^{+} and CD8^{+} T cells in 7610c2, but not 2838c3 TiME

(**Supplemental Figure 11A***,7160c2* and **11B***,2838c3*), despite higher baseline levels of

PD-1 expression (**Supplemental Figure 11C**). Co-expression of PD-1 and TIM-3 on

tumor infiltrating T cells was modestly increased only in 7160c2 model after ADT-007

treatment (**Supplemental Figure 11D-E**).

In the 7160c2 TiME both CD4^{+} and CD8^{+} T cells exhibited increased expression

of IL-2, granzyme B, and IFNγ (**Figure 6G-H**) after stimulation with PMA/ionomycin,

whereas minimal changes in cytokine expression were noted in T cells within the

2838c3 TiME (**Figure 6O-P**). Using an IFNγ-Thy1.1 reporter model system [24], we

assessed IFNγ production marked by surface Thy1.1 on innate and adaptive T and NK

subsets. Consistent with intracellular cytokine staining (ICS) results, ADT-007 increased

IFNγ production in CD4^{+} and CD8^{+} T cell subsets in the 7160c2, but not in 2838c3

TiME. In NKT and γδ T cells, IFNγ was unchanged (**Supplemental Figure 12A***,7160c2*

and 12B*,2838c3*), whereas a decrease was noted in NK cells from ADT-007 treated

tumors (**Supplemental Figure 12A***,7160c2* **and 12B***,2838c3*).

DISCUSSION

ADT-007 potently inhibited the growth of cancer cells with mutant RAS

regardless of the RAS mutation or isozyme. Cancer cells with RAS^{WT} but mutations

resulting in constitutive activation of RTK, loss-of-function in GAPs were comparably

sensitive to ADT-007 as was a gastric cancer cell line harboring gene amplification of

the KRAS^{WT} gene. Despite broad activity, ADT-007 displayed exquisite selectivity as

cancer cells with BRAF mutations, immediately downstream from RAS, and cells from

normal tissues were essentially insensitive to ADT-007. ADT-007 sensitivity was

attributed to high RAS-GTP level and dependence on RAS signaling for proliferation

(e.g., RAS addiction), while normal cells or cancer cell lines with RAS^{WT} have a

metabolic mechanism protecting them from ADT-007 cytotoxicity involving

glucuronidation.

Concentrations of ADT-007 effective for binding RAS in cells or recombinant

RAS were within the same range as concentrations required to inhibit growth, colony

formation, and induce mitotic arrest and apoptosis, as well as concentrations that block

RAS activation and MAPK/AKT signaling. The inhibitory effect of ADT-007 on RAS-GTP

occurred by treating KRAS mutant cancer cells under standard growth conditions,

serum-free conditions, or following EGF stimulation. ADT-007 reduced RAS-GTP from

mutant KRAS and co-expressed HRAS^{WT} and NRAS^{WT} isozymes that were activated by

EGF. This activity may be a key advantage over mutant-specific KRAS or pan-KRAS

inhibitors to circumvent resistance from compensatory activity of unchecked RAS^{WT}

NRAS or HRAS isozymes, although as discussed below ADT-007 may have additional

advantages such as its unique ability to induce mitotic arrest and apoptosis that may

also avert resistance.

Localized delivery of ADT-007 strongly inhibited tumor growth in immunodeficient

and syngeneic murine models of CRC and PDA with partial dependence on the

adaptive immune system. ADT-007 activated T cells and enhanced their cytotoxic

functions in KRAS^{G12D} mutated CRC and PDA, immune-competent mouse models. Oral

delivery of a prototypic prodrug, ADT-316, significantly inhibited tumor growth but with

modest activity likely because of incomplete conversion to ADT-007. Recognizing the

limitations of ADT-316, we designed an improved series of prodrugs that are currently

under study, one of which, ADT-1004, shows robust antitumor activity and exquisite

RAS selectivity in mouse models involving mouse or human PDA cell lines or PDX

models of PDA. ADT-1004 also shows superior antitumor activity over sotorasib and

adagrasib in experiments comparing equal dosages of each compound administer orally

in mice implanted with resistant MIA PaCa-2 PDA cells (bioRxiv 2024.

10.1101/2024.10.04.616725).

ADT-007 appears to bind RAS within the nucleotide binding domain to block

GTP-activation of effector interactions and downstream MAPK/AKT signaling.

Molecular modeling studies provided initial evidence that ADT-007 binds RAS within the

nucleotide-binding domain, which were supported by biochemical experiments showing

that ADT-007 inhibits GTP binding to recombinant RAS and GTP-dependent RAS-RAF

interactions. These effects were only observed under nucleotide-free conditions, but

not under nucleotide loading conditions. The selectivity of ADT-007 to bind nf-RAS was

confirmed by NMR spectroscopy experiments, which provided additional evidence for

binding within the nucleotide-binding domain, although further studies are needed to

define the mechanism of binding more precisely.

ADT-007 binding to nf-RAS could negatively impact its antitumor activity, given

this is a transitional conformation and the presumed temporal nature of target exposure.

While *in vitro* experiments show near complete killing of RAS mutant cancer cells by

ADT-007 given adequate time, an important consideration for *in vivo* antitumor activity is

if ADT-007 has a sufficiently long plasma half-life for target occupancy. Indeed, oral

administration of ADT-316 and ADT-1004 generated sustained plasma levels of ADT-007 in

mice that were appreciably greater than *in vitro* growth IC_{50} values and significantly inhibited

tumor growth *in vivo*. In addition, previous reports that a significant percentage of RAS

exists in cells as an apoenzyme and that a nf-RAS monobody inhibits tumor growth [37],

suggest that ADT-007 binding to nf-RAS can overcome this perceived liability.

Metabolic deactivation of ADT-007 by UGT-mediated glucuronidation presents

implications for the interpretation of both *in vitro* and *in vivo* experiments. Liver

microsome experiments showed that ADT-007 is rapidly glucuronidated by UGT

enzymes ubiquitously expressed in normal tissues and well known to metabolize

compounds with phenolic hydroxyl moieties. Unexpectedly, RAS^{WT} cancer cells with

mutant BRAF had high levels of major UGT isozymes commonly expressed in normal

cells. In contrast, cancer cells with mutant KRAS had low levels of UGT isozymes.

Consistent with these differences, low ADT-007 levels and glucuronide-conjugated

ADT-007 were measured in RAS^{WT} cells, while high ADT-007 levels but no glucuronide-

conjugated ADT-007 were measured in KRAS mutant HCT-116 cells by LC-MS/MS.

Moreover, conditioned medium obtained from RAS^{WT} cells exposed to ADT-007, but not

KRAS mutant cells, completely lost growth inhibitory activity, reflecting metabolic

deactivation of ADT-007. Pretreatment with a known competitive UGT inhibitor,

propofol, also increased sensitivity of RAS^{WT} cells to ADT-007 but did not significantly

impact sensitivity of mutant KRAS cells to ADT-007. These results suggest that UGT-

mediated glucuronidation protects bystander cells from normal tissues from pan-RAS

inhibition by ADT-007. Despite acting as a pan-RAS inhibitor, which otherwise would be

considered lethal based on genomic knock-out experiments in mice [38, 39], ADT-007

has a unique metabolic mechanism of RAS selectivity. While UGT isozymes are well

known to be expressed in normal tissues, including blood cells, and regulated by

multiple signaling pathways and transcription factors [40], further study is needed to

understand how UGT expression is suppressed by RAS mutations.

Other sulindac derivatives were previously reported to inhibit cancer cell growth

by a dual mechanism involving inhibition of cyclic guanine monophosphate

phosphodiesterase (cGMP PDE) activity and tubulin polymerization [31]. The former

was based on reports that derivatives with a trimethoxy substitution inhibited cGMP

PDE activity at concentrations that inhibit cancer cell growth [21, 41], while the latter

was based on their chemical resemblance with colchicine, which is known to bind

tubulin and disrupt microtubules [42]. We, therefore, determined if ADT-007 inhibits

cGMP PDE activity by measuring effects on cGMP hydrolysis using recombinant

PDE10, a cGMP-degrading PDE isozyme (not shown), or capacity to bind PDE10 in

cells using cellular thermal stability assays **(Figure 3A)**, but found no evidence of

PDE10 inhibition or binding, respectively. In addition, ADT-007 only inhibited tubulin

polymerization at concentrations ~100-fold higher than growth IC_{50} values

(**Supplemental Figure 3H**). These observations lead us to conclude that neither cGMP

PDE nor tubulin polymerization inhibition account for the growth inhibitory activity of

ADT-007. Reports that tubulin inhibitors induce MAPK pathway activity [43], rather than

inhibit MAPK signaling, provide additional evidence that ADT-007 acts by targeting

RAS. Moreover, colchicine lacked selectivity in growth assays involving RAS mutant vs.

RAS[WT] cancer cells (not shown). We also tested the topoisomerase inhibitor,

etoposide, because it has the same phenolic hydroxyl moiety as ADT-007, which we

show is essential for RAS binding but found no evidence of RAS selectivity in growth

assays.

Unlike KRAS[G12C] inhibitors that have limited use for patients with this relatively

rare mutation [30, 44], ADT-007 has broad anticancer activity with potential to be

developed as a generalizable monotherapy for the majority of patients with RAS mutant

cancers. In addition, a significant proportion of patients with RTK fusions that hyper-

activate RAS[WT] isozymes [45] would be expected to be sensitive to ADT-007. Evidence

indicates that ADT-007 exhibits high potency in tumor cells with activated RAS

signaling, which may provide greater therapeutic utility in patients with metastatic

disease in which the proliferation of cancer cells depends upon activated KRAS

signaling for survival in the metastatic niche [46].

Equally significant to the broad antitumor activity of ADT-007 is its potential to

provide a second-line therapy for patients with KRAS mutant cancers that develop

resistance to other RAS inhibitors, for example, from co-occurring RAS mutations or

compensation from RAS[WT] isozymes [7, 20]. We show here that ADT-007 maintains

growth inhibitory activity in LLC cells with co-occurring KRAS[G12C] and NRAS[Q61H]

mutations [47], while the KRAS[G12C] inhibitor, sotorasib, and the pan-KRAS inhibitor, BI-

2865 lost growth inhibitory activity. Other experiments showed that ADT-007 induces

mitotic arrest and apoptosis resulting in cytotoxicity, while BI-2865 or AMG-510 did not,

and another pan-RAS inhibitor, RMC-2865, did not induce apoptosis, appearing to be

limited to a cytostatic mechanism. This difference was apparent in clonogenic colony

formation assays in which ADT-007 treatment caused near complete inhibition and to a

much greater extent than BI-2865 or RMC-6236 when compared side-by-side. These

advantages of ADT-007 over other RAS inhibitors as observed *in vitro* awaits further *in*

vivo studies but suggests the potential to escape intrinsic and acquired resistance.

ADT-007-mediated *in vivo* tumor growth inhibition also appears to be partially

dependent upon the adaptive immune response as evidenced by reduced antitumor

activity of ADT-007 in 2838c3 and 7160c2 tumors grown in immune-deficient Rag $1^{-/-}$

mice lacking B and T cells *vs.* in immune-competent C57BL/6. Similar findings were

observed with $KRAS^{G12C}$ [48, 49] and $KRAS^{G12D}$-specific inhibitors [50], demonstrating

that T cells contribute to the efficacy of other RAS inhibitors. The presence of T cells

predicted synergy with PD-1 immune checkpoint blockade in preclinical models of

NSCLC and PDA [49, 51]. However, the importance of B cells to anti-tumor immunity of

ADT-007 requires further study.

In T cell inflamed PDA tumor models (2838c3 and 7160c2), ADT-007 treatment

increased numbers of innate-like T lymphocytes (NK T ($CD3^+$ $NK1.1^+$), γδT, and $CD3^+$

$CD4^-$/$CD8^-$ T, and NK ($CD3^-$ $NK1.1^+$) cells), whereas conventional $CD4^+$ and $CD8^+$ T cell

numbers did not change. Analysis of IFNγ levels using Thy1.1-IFNγ reporter mice

revealed that most adaptive and innate-like T cells were $Thy1.1^+$, consistent with high

levels of IFNγ production *in vivo*, whereas, differences in ADT-007 on T cell activation

and cytokine secretion in both models, may be explained by the higher baseline

activation and cytokine secretion of T cells isolated from the 2838c3 model system [25],

suggesting that differences in baseline T cell activation and function may contribute to

differences in outcomes with ADT-007 treatment in different tumor models. Overall,

these findings suggest role(s) for innate and adaptive immune response in the antitumor

activity of ADT-007 that can potentially be exploited with relevant immune therapy

combinations.

In conclusion, ADT-007 represents a 1st-in-class pan-RAS inhibitor with potential

to address the complex RAS mutational landscape of many human cancers and

properties superior to other RAS inhibitors to circumvent resistance. These results

support IND-enabling studies of an ADT-007 analog or prodrug as a generalizable

monotherapy or combined with other RAS pathway-directed inhibitors or immunotherapy

for treating RAS-driven cancers.

References

1.   Siegel RL, Giaquinto AN, Jemal A: **Cancer statistics, 2024**. *CA Cancer J Clin* 2024, **74**(1):12-49.

2.   Yang Y, Zhang H, Huang S, Chu Q: **KRAS Mutations in Solid Tumors: Characteristics, Current**
Therapeutic Strategy, and Potential Treatment Exploration. *J Clin Med* 2023, **12**(2).

3.   Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, Ross S, Conrads TP,
Veenstra TD, Hitt BA *et al*: **Preinvasive and invasive ductal pancreatic cancer and its early**
detection in the mouse. *Cancer Cell* 2003, **4**(6):437-450.

4.   Kojima K, Vickers SM, Adsay NV, Jhala NC, Kim HG, Schoeb TR, Grizzle WE, Klug CA: **Inactivation**
of Smad4 accelerates Kras(G12D)-mediated pancreatic neoplasia. *Cancer Res* 2007,
67(17):8121-8130.

5.   Boutin AT, Liao WT, Wang M, Hwang SS, Karpinets TV, Cheung H, Chu GC, Jiang S, Hu J, Chang K
et al: **Oncogenic Kras drives invasion and maintains metastases in colorectal cancer**. *Genes Dev*
2017, **31**(4):370-382.

6.   Bardeesy N, Aguirre AJ, Chu GC, Cheng KH, Lopez LV, Hezel AF, Feng B, Brennan C, Weissleder R,
Mahmood U *et al*: **Both p16(Ink4a) and the p19(Arf)-p53 pathway constrain progression of**
pancreatic adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A* 2006, **103**(15):5947-5952.

7.   Sheffels E, Kortum RL: **The Role of Wild-Type RAS in Oncogenic RAS Transformation**. *Genes*
(Basel) 2021, **12**(5).

8.   Ying H, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sananikone E, Locasale JW, Son J,
Zhang H, Coloff JL *et al*: **Oncogenic Kras maintains pancreatic tumors through regulation of**
anabolic glucose metabolism. *Cell* 2012, **149**(3):656-670.

9.   Mukhopadhyay S, Goswami D, Adiseshaiah PP, Burgan W, Yi M, Guerin TM, Kozlov SV, Nissley
DV, McCormick F: **Undermining Glutaminolysis Bolsters Chemotherapy While NRF2 Promotes**
Chemoresistance in KRAS-Driven Pancreatic Cancers. *Cancer Res* 2020, **80**(8):1630-1643.

10.   Collins MA, Brisset JC, Zhang Y, Bednar F, Pierre J, Heist KA, Galban CJ, Galban S, di Magliano MP:
Metastatic pancreatic cancer is dependent on oncogenic Kras in mice. *PLoS One* 2012,
7(12):e49707.

11.   Zhang Y, Yan W, Collins MA, Bednar F, Rakshit S, Zetter BR, Stanger BZ, Chung I, Rhim AD, di
Magliano MP: **Interleukin-6 is required for pancreatic cancer progression by promoting MAPK**
signaling activation and oxidative stress resistance. *Cancer Res* 2013, **73**(20):6359-6374.

12.   Zdanov S, Mandapathil M, Abu Eid R, Adamson-Fadeyi S, Wilson W, Qian J, Carnie A, Tarasova N,
Mkrtichyan M, Berzofsky JA *et al*: **Mutant KRAS Conversion of Conventional T Cells into**
Regulatory T Cells. *Cancer Immunol Res* 2016, **4**(4):354-365.

13.   Bayne LJ, Beatty GL, Jhala N, Clark CE, Rhim AD, Stanger BZ, Vonderheide RH: **Tumor-derived**
granulocyte-macrophage colony-stimulating factor regulates myeloid inflammation and T cell
immunity in pancreatic cancer. *Cancer Cell* 2012, **21**(6):822-835.

14.   Ischenko I, D'Amico S, Rao M, Li J, Hayman MJ, Powers S, Petrenko O, Reich NC: **KRAS drives**
immune evasion in a genetic model of pancreatic cancer. *Nat Commun* 2021, **12**(1):1482.

15.   Coelho MA, de Carne Trecesson S, Rana S, Zecchin D, Moore C, Molina-Arcas M, East P, Spencer-
Dene B, Nye E, Barnouin K *et al*: **Oncogenic RAS Signaling Promotes Tumor Immunoresistance**
by Stabilizing PD-L1 mRNA. *Immunity* 2017, **47**(6):1083-1099 e1086.

16.   Muthalagu N, Monteverde T, Raffo-Iraolagoitia X, Wiesheu R, Whyte D, Hedley A, Laing S,
Kruspig B, Upstill-Goddard R, Shaw R *et al*: **Repression of the Type I Interferon Pathway**
Underlies MYC- and KRAS-Dependent Evasion of NK and B Cells in Pancreatic Ductal
Adenocarcinoma. *Cancer Discov* 2020, **10**(6):872-887.

17.   Stalnecker CA, Der CJ: **RAS, wanted dead or alive: Advances in targeting RAS mutant cancers**.
Sci Signal 2020, **13**(624).
18.   Kwan AK, Piazza GA, Keeton AB, Leite CA: **The path to the clinic: a comprehensive review on**
direct KRAS(G12C) inhibitors. *J Exp Clin Cancer Res* 2022, **41**(1):27.
19.   Ryan MB, Fece de la Cruz F, Phat S, Myers DT, Wong E, Shahzade HA, Hong CB, Corcoran RB:
Vertical Pathway Inhibition Overcomes Adaptive Feedback Resistance to KRAS(G12C)
Inhibition. *Clin Cancer Res* 2020, **26**(7):1633-1643.
20.   Awad MM, Liu S, Rybkin, II, Arbour KC, Dilly J, Zhu VW, Johnson ML, Heist RS, Patil T, Riely GJ *et*
al: **Acquired Resistance to KRAS(G12C) Inhibition in Cancer**. *N Engl J Med* 2021, **384**(25):2382-
2393.
21.   Thompson HJ, Jiang C, Lu J, Mehta RG, Piazza GA, Paranka NS, Pamukcu R, Ahnen DJ: **Sulfone**
metabolite of sulindac inhibits mammary carcinogenesis. *Cancer Res* 1997, **57**(2):267-271.
22.   Herrmann C, Block C, Geisen C, Haas K, Weber C, Winde G, Moroy T, Muller O: **Sulindac sulfide**
inhibits Ras signaling. *Oncogene* 1998, **17**(14):1769-1776.
23.   Karaguni IM, Glusenkamp KH, Langerak A, Geisen C, Ullrich V, Winde G, Moroy T, Muller O: **New**
indene-derivatives with anti-proliferative properties. *Bioorg Med Chem Lett* 2002, **12**(4):709-
713.
24.   Harrington LE, Janowski KM, Oliver JR, Zajac AJ, Weaver CT: **Memory CD4 T cells emerge from**
effector T-cell progenitors. *Nature* 2008, **452**(7185):356-360.
25.   Li J, Byrne KT, Yan F, Yamazoe T, Chen Z, Baslan T, Richman LP, Lin JH, Sun YH, Rech AJ *et al*:
Tumor Cell-Intrinsic Factors Underlie Heterogeneity of Immune Cell Infiltration and Response
to Immunotherapy. *Immunity* 2018, **49**(1):178-193 e177.
26.   Flatmark K, Reed W, Halvorsen T, Sorensen O, Wiig JN, Larsen SG, Fodstad O, Giercksky KE:
Pseudomyxoma peritonei--two novel orthotopic mouse models portray the PMCA-I
histopathologic subtype. *BMC Cancer* 2007, **7**:116.
27.   Flatmark K, Davidson B, Kristian A, Stavnes HT, Førsund M, Reed W: **Exploring the peritoneal**
surface malignancy phenotype—a pilot immunohistochemical study of human
pseudomyxoma peritonei and derived animal models. *Human Pathology* 2010, **41**(8):1109-
1119.
28.   Fleten KG, Lund-Andersen C, Waagene S, Abrahamsen TW, Mørch Y, Boye K, Torgunrud A,
Flatmark K: **Experimental Treatment of Mucinous Peritoneal Metastases Using Patient-Derived**
Xenograft Models. *Transl Oncol* 2020, **13**(8):100793.
29.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, Wilson CJ, Lehar J,
Kryukov GV, Sonkin D *et al*: **The Cancer Cell Line Encyclopedia enables predictive modelling of**
anticancer drug sensitivity. *Nature* 2012, **483**(7391):603-607.
30.   Waters AM, Der CJ: **KRAS: The Critical Driver and Therapeutic Target for Pancreatic Cancer**.
Cold Spring Harb Perspect Med 2018, **8**(9).
31.   Xiao D, Deguchi A, Gundersen GG, Oehlen B, Arnold L, Weinstein IB: **The sulindac derivatives**
OSI-461, OSIP486823, and OSIP487703 arrest colon cancer cells in mitosis by causing
microtubule depolymerization. *Mol Cancer Ther* 2006, **5**(1):60-67.
32.   Lee KJ, Chang WL, Chen X, Valiyaveettil J, Ramirez-Alcantara V, Gavin E, Musiyenko A, Madeira
da Silva L, Annamdevula NS, Leavesley SJ *et al*: **Suppression of Colon Tumorigenesis in Mutant**
Apc Mice by a Novel PDE10 Inhibitor that Reduces Oncogenic β-Catenin. *Cancer Prev Res*
(Phila) 2021, **14**(11):995-1008.
33.   Hunter JC, Gurbani D, Ficarro SB, Carrasco MA, Lim SM, Choi HG, Xie T, Marto JA, Chen Z, Gray
NS *et al*: **In situ selectivity profiling and crystal structure of SML-8-73-1, an active site inhibitor**
of oncogenic K-Ras G12C. *Proc Natl Acad Sci U S A* 2014, **111**(24):8895-8900.

34.   Agola JO, Hong L, Surviladze Z, Ursu O, Waller A, Strouse JJ, Simpson DS, Schroeder CE, Oprea TI,
Golden JE *et al*: **A competitive nucleotide binding inhibitor: in vitro characterization of Rab7**
GTPase inhibition. *ACS Chem Biol* 2012, **7**(6):1095-1108.
35.   Hunter JC, Manandhar A, Carrasco MA, Gurbani D, Gondi S, Westover KD: **Biochemical and**
Structural Analysis of Common Cancer-Associated KRAS Mutations. *Mol Cancer Res* 2015,
13(9):1325-1335.
36.   John J, Rensland H, Schlichting I, Vetter I, Borasio GD, Goody RS, Wittinghofer A: **Kinetic and**
structural analysis of the Mg(2+)-binding site of the guanine nucleotide-binding protein p21H-
ras. *J Biol Chem* 1993, **268**(2):923-929.
37.   Khan I, Koide A, Zuberi M, Ketavarapu G, Denbaum E, Teng KW, Rhett JM, Spencer-Smith R,
Hobbs GA, Camp ER *et al*: **Identification of the nucleotide-free state as a therapeutic**
vulnerability for inhibition of selected oncogenic RAS mutants. *Cell Rep* 2022, **38**(6):110322.
38.   Esteban LM, Vicario-Abejon C, Fernandez-Salguero P, Fernandez-Medarde A, Swaminathan N,
Yienger K, Lopez E, Malumbres M, McKay R, Ward JM *et al*: **Targeted genomic disruption of H-**
ras and N-ras, individually or in combination, reveals the dispensability of both loci for mouse
growth and development. *Mol Cell Biol* 2001, **21**(5):1444-1452.
39.   Johnson L, Greenbaum D, Cichowski K, Mercer K, Murphy E, Schmitt E, Bronson RT, Umanoff H,
Edelmann W, Kucherlapati R *et al*: **K-ras is an essential gene in the mouse with partial**
functional overlap with N-ras. *Genes Dev* 1997, **11**(19):2468-2481.
40.   Allain EP, Rouleau M, Lévesque E, Guillemette C: **Emerging roles for UDP-**
glucuronosyltransferases in drug resistance and cancer progression. *Br J Cancer* 2020,
122(9):1277-1287.
41.   Tinsley HN, Gary BD, Thaiparambil J, Li N, Lu W, Li Y, Maxuitenko YY, Keeton AB, Piazza GA:
Colon tumor cell growth-inhibitory activity of sulindac sulfide and other nonsteroidal anti-
inflammatory drugs is associated with phosphodiesterase 5 inhibition. *Cancer Prev Res (Phila)*
2010, **3**(10):1303-1313.
42.   Lu Y, Chen J, Xiao M, Li W, Miller DD: **An overview of tubulin inhibitors that interact with the**
colchicine binding site. *Pharm Res* 2012, **29**(11):2943-2971.
43.   Waters AM, Khatib TO, Papke B, Goodwin CM, Hobbs GA, Diehl JN, Yang R, Edwards AC, Walsh
KH, Sulahian R *et al*: **Targeting p130Cas- and microtubule-dependent MYC regulation sensitizes**
pancreatic cancer to ERK MAPK inhibition. *Cell Rep* 2021, **35**(13):109291.
44.   Coley AB, Ward A, Keeton AB, Chen X, Maxuitenko Y, Prakash A, Li F, Foote JB, Buchsbaum DJ,
Piazza GA: **Pan-RAS inhibitors: Hitting multiple RAS isozymes with one stone**. *Adv Cancer Res*
2022, **153**:131-168.
45.   Luchini C, Paolino G, Mattiolo P, Piredda ML, Cavaliere A, Gaule M, Melisi D, Salvia R, Malleo G,
Shin JI *et al*: **KRAS wild-type pancreatic ductal adenocarcinoma: molecular pathology and**
therapeutic opportunities. *J Exp Clin Cancer Res* 2020, **39**(1):227.
46.   di Magliano MP, Logsdon CD: **Roles for KRAS in pancreatic tumor development and**
progression. *Gastroenterology* 2013, **144**(6):1220-1229.
47.   Molina-Arcas M, Moore C, Rana S, van Maldegem F, Mugarza E, Romero-Clavijo P, Herbert E,
Horswell S, Li LS, Janes MR *et al*: **Development of combination therapies to maximize the**
impact of KRAS-G12C inhibitors in lung cancer. *Sci Transl Med* 2019, **11**(510).
48.   Canon J, Rex K, Saiki AY, Mohr C, Cooke K, Bagal D, Gaida K, Holt T, Knutson CG, Koppada N *et al*:
The clinical KRAS(G12C) inhibitor AMG 510 drives anti-tumour immunity. *Nature* 2019,
575(7781):217-223.
49.   Mugarza E, van Maldegem F, Boumelha J, Moore C, Rana S, Llorian Sopena M, East P, Ambler R,
Anastasiou P, Romero-Clavijo P *et al*: **Therapeutic KRAS(G12C) inhibition drives effective**

interferon-mediated antitumor immunity in immunogenic lung cancers. *Sci Adv* 2022,
8(29):eabm8780.
50.   Kemp SB, Cheng N, Markosyan N, Sor R, Kim IK, Hallin J, Shoush J, Quinones L, Brown NV,
Bassett JB *et al*: **Efficacy of a Small-Molecule Inhibitor of KrasG12D in Immunocompetent**
Models of Pancreatic Cancer. *Cancer Discov* 2023, **13**(2):298-311.
51.   Mahadevan KK, McAndrews KM, LeBleu VS, Yang S, Lyu H, Li B, Sockwell AM, Kirtley ML, Morse
SJ, Moreno Diaz BA *et al*: **KRAS(G12D) inhibition reprograms the microenvironment of early**
and advanced pancreatic cancer to promote FAS-mediated killing by CD8(+) T cells. *Cancer Cell*
2023, **41**(9):1606-1620 e1608.

Figures and Figure Legends

Figure 1. ADT-007 selectively inhibits growth of RAS mutant cancer cells by

blocking activated RAS and MAPK/AKT signaling and superior in comparison

with other RAS inhibitors. (A) Chemical structure of ADT-007. **(B)** ADT-007 inhibition

of KRAS^{G13D} HCT-116 vs. KRAS^{WT}, BRAF^{V600E} HT-29 CRC cell growth following 72 h of

treatment (CTG assay)**.** Quadruplicate samples ± SEM. Inset: Activated RAS-GTP-RBD

pulldown evaluated by WB using pan-RAS antibodies. Total RAS and GAPDH served

as loading controls. **(C)** Ectopic expression of HRAS^{G12V} sensitized HT-29 cells to ADT-

007 growth inhibition (quadruplicates ± SEM). Inset: Activated RAS in untreated

parental, vector control (retro), and transfected cells (H-Ras). (**D**) HCT-116 KRAS^{G13D}

CRC cells were treated with increasing concentrations of ADT-007 overnight in

complete medium (10% FBS). Activated RAS-GTP pulldowns, phospho(P)-cRAF, P-

MEK, P-ERK, P-AKT, and total RAS and GAPDH as loading controls were detected by

WB. Data represent 3 independent experiments. (**E, F**) Serum-starved MIA PaCa-2

KRAS^{G12C} PDA cells were treated overnight with vehicle or increasing concentrations of

ADT-007, then EGF stimulated (30 ng/mL, 10 min). (**E**) P-cRAF, P-MEK, P-ERK, P-AKT

and GAPDH were detected by WB. Data represent two independent experiments. (**F**)

Activated RAS-GTP was isolated by RAS-RBD pulldown followed by WB with pan-RAS,

KRAS, NRAS, or HRAS specific antibodies and GAPDH. Data represent at least two

independent experiments. (**G**) Growth inhibition of MIA PaCa-2 cells by ADT-007 *vs.* BI-

2865 or AMG-510 as determined by CTG assay following 72-h treatment (**H, I**) Growth

inhibition of LLC cells (KRAS^{G12C} / NRAS^{Q61H}) by ADT-007 *vs.* BI-2865 or AMG-510

following 72-h treatment. (**J**) Apoptosis induction of MIA PaCa-2 cells following 48-h

treatment with ADT-007, BI-2865 or RMC-6236 as measured by cleaved caspase-3/7

using a luminescence assay (Promega). Error bars represent SEM.

Type	Tissue Origin	Cell Line	Mutation Status	IC_{50} (nM)	UGT1A10
Mutant RAS, sensitive	Pancreatic	MiaPaCa2	KRAS G12C	2.1	0.263
	Pancreatic	Panc1	KRAS G12D	2.4	0.043
	Pancreatic	SW1990	KRAS G12D	6.3	0.138
	Lung	A549	KRAS G12S	6.8	0.070
	Pancreatic	CFPAC1	KRAS G12V	5.8	2.63
	Colon	SW480	KRAS G12V	6.5	0.58
	Breast	MDA231	KRAS G13D	3.6	0.043
	Colon	HCT116	KRAS G13D	4.7	0
	Ovarian	OVCAR8	KRAS P121H	5.8	0.057
	Breast	HS578	HRAS G12D	3.9	0.098
	Skin	SKMEL2	NRAS Q61K	2.1	0
	Lung	H1299	NRAS Q61K	2.4	0
WT RAS, resistant	Normal Lung	NHAEC	WT RAS	19100	*
	Normal Colon	NCM460	WT RAS	50000	*
	Gastric	MKN-1	Amplified WT KRAS	4.4	0.49
Downstream mutation (WT RAS) resistant	Ovarian	OV90	WT RAS/mut BRAF	350	4.57
	Colon	HT29	WT RAS/mut BRAF	512	6.97
	Pancreatic	BXPC3	WT RAS/mut BRAF	2500	7.75
Upstream RAS activation sensitive	Lung	H1975	WT RAS/mut EGFR	3.9	*
	Skin (mouse)	B16	WT RAS/mut PDGFR	5.8	*
	Brain	U87MG	WT RAS/mut NF1	1.6	0
	Ovarian	SKOV3	WT RAS/mut NF1	6.7	0.202
Amplified WT RAS, sensitive	Gastric	MKN-1	Amplified WT KRAS	4.4	0.490

Table 1. ADT-007 sensitivity is associated with high activated RAS and low UGT

superfamily enzyme levels. Growth inhibitory (IC_{50}) values following 72 h of treatment

(CTG assay). TPM normalized expression of the most significant differentially

expressed UGT1A10 enzyme from published data (Cancer Cell Line Encyclopedia;

DepMap Portal; Broad Institute). * Expression data not available, † TPM normalized

UGT2B7 expression in SK-MEL-28.

Figure 2. Phenolic moiety is critical for RAS selectivity of ADT-007 and renders sensitivity to metabolic deactivation. (**A**) Left panel: ADT-007 selectively inhibits the growth of KRAS^{G12C} MIA PaCa-2 vs. RAS^{WT} BxPC-3 PDA cells (72 h, CTG assay; triplicates ± SEM). (**B**) Left panel: ADT-106 lacking a phenolic OH moiety was less potent and nonselective (72 h, CTG assay; triplicates ± SEM). Right panels (**A, B**): MIA PaCa-2 cells were treated for 18 h with increasing concentrations of ADT-007 (**A**) or ADT-106 (**B**) and activated RAS-GTP-RBD pulldown evaluated by WB using pan-RAS antibodies. Total RAS and GAPDH served as loading controls. (**C**) ADT-007 is rapidly degraded by mouse liver microsomes with a corresponding increase in ADT-007-glucuronide (ADT-007-Glu) levels (**D**). (**E**) UGT1 protein expression in ADT-007 sensitive *vs.* resistant tumor cells by WB. (**F**) UGT1 protein expression and control GAPDH (WB) from (**E**) were quantified and normalized using Image J. Representative of two independent experiments. (**G**) ADT-007 and ADT-007-glucuronide levels in HCT-116 and HT-29 cell pellets following 24 h treatment with ADT-007. Levels were quantified in triplicate by LC-MS/MS (mean ± SEM), * = p < 0.05. (**H**) Conditioned medium (CM) from MIA PaCa-2 (low UGT expression) and BxPC-3 (high UGT expression) PDA cells pretreated with ADT-007 at 100 nM for 24 h was collected for growth inhibition assays involving MIA PaCa-2 cells. CM from MIA PaCa-2 cells showed similar growth inhibitory activity as the control group using freshly prepared ADT-007. CM from BxPC-3 cells resulted in complete loss of potency of ADT-007 to inhibit the growth of MIA PaCa-2 cells (96 h, CTG assay; triplicates ± SEM).

1011

1012

Figure 3. ADT-007 binds KRAS in cells and inhibits nucleotide loading and

effector binding using recombinant RAS. Cellular target engagement assays show:

(**A**) ADT-007 binds KRAS with high affinity in KRAS^{G13D} HCT-116 but not KRAS^{WT} HT-

29 CRC cells. Graphs show levels of compound stabilized KRAS detected by WB

compared with vehicle control. Curves are graphed as the average of two replicates ±

SEM. (**B**) Binding affinity of ADT-007 for mutant KRAS^{G12C} was assessed by 45 min

treatment of HEK293 cells expressing KRAS^{G12C}-Micro-Tag construct compared with

DMSO vehicle. Curve is graphed as the average of two replicates ± SEM. (**C-D**) The

inhibitory activity of ADT-007 to block recombinant KRAS^{WT} activation (effector binding)

in a cell-free system was evaluated by RAS-RBD pulldown assays. (**C**) ADT-007

inhibited KRAS^{WT}-RAF activation before GTP addition. (**D**) ADT-007 did not significantly

inhibit KRAS^{WT} activation when added after GTP loading. (**E-F**) Kinetics of GTP binding

to KRAS were measured using a fluorescent GTP analog (Mant-GTP). ADT-007

reduced the rate of Mant-GTP binding to KRAS if pretreated with EDTA to chelate Mg2+

(**E**), but not if added after non-hydrolysable GTPγS before incubation with ADT-007 (**F**).

Data are representative of at least two independent experiments.

1029

1030

1031

1032

1033

1034

1035

Figure 4. NMR analysis of ADT-007 binding to KRAS. (A) NMR spectrum of full

length KRAS in the presence (blue) or absence of Mg^{2+} (red) representing nucleotide-

loaded or nf-RAS, respectively. (B) The addition of ADT-007 (red) to nf-KRAS (blue)

(5:1 ratio) resulted in statistically significant NMR signal attenuation at G13 and K16 and

other residues (described in methods). Assignments for G13 and K16, due to their

unique resonance frequencies, are shown to the left.

1042

1043

1044

Figure 5. ADT-007 induces growth arrest and regression in mouse tumor models

(**A**) Left panel: Growth inhibition of SQ KRAS^{G12V} PDX gall bladder adenocarcinoma (PT 5 mg/kg, BID, 13 days, n = 5 mice/group, mean $\pm$ SEM). Right panel: % change in tumor volume. (**B-D**) Left panels: Growth inhibition of mucinous PDX colorectal adenocarcinomas (**B**) PMCA-1, (**C**) PMP1, and (**D**) PC244 in nude mice (IP, 2.5 mg/kg, BID, treatment periods as indicated, n = 5-6 mice/group, mean $\pm$ SEM). Right panels: % change in tumor volume. (**E**) Left panel: Growth inhibition of SQ CT26 CRC tumors (IT, 10 mg/kg, QD, 13 days, mean $\pm$ SEM, n=12). Right panel: % change in tumor volume. (**F-H**) Left panels: Growth inhibition of SQ (**F**) 2838c3, (**G**) 7160c2, (**H**) 5363 PDA tumors in C57BL/6 mice (PT, 5 mg/kg, BID, treatment periods as indicated) (n = 7 - 10 mice/group, mean $\pm$ SEM). Right Panels: % change in tumor volume. All animal studies are representative of 2-3 independent experiments. Left panels: Statistical significance was assessed using 2-way ANOVA with Sadik's multiple comparisons test. Right panels: Statistical significance was assessed with Welch's t-test.

Figure 6

Figure 6. ADT-007 modulates tumor immunity in the PDA TME. (A-H) 7160c2 PDA

model (20 day of treatment) and (**I-P**) 2838c3 PDA model (17 day of treatment). (**A and**

I) % CD45$^-$ YFP$^+$ tumor cells in ADT-007 vs. sham (n = 8). (**B and J**) Numbers/mg

tumor of total myeloid (CD11b$^{+/-}$CD11c$^{+/-}$, macrophage, gMDSC, and mMDSC

(**Supplemental Figure 8A**-*gating scheme*) in ADT-007 treated vs. sham (n = 8). (**C and**

K) Numbers/mg tumor of cDC1 and cDC2 subsets in ADT-007 vs. sham (n = 8). (**D and**

L) Tumor growth in 7160c2 PDA tumor-bearing C57BL/6 WT vs. C57BL/6 Rag 1$^{-/-}$ mice

(SQ, 5 mg/kg, BID, days 7-28, n = 6-22 mice/group, mean $\pm$ SEM). (**E and M**)

Numbers/mg tumor of CD4$^+$ Foxp3$^+$, CD4$^+$ Foxp3$^-$, CD8$^+$ T, γδ$^+$ (TCRγδ$^+$), NK T (CD3$^+$

NK1.1$^+$), and CD4/CD8$^-$ / γδ$^-$ T cells and (**F and N**) NK (CD3$^-$ NK1.1$^+$) cells

(**Supplemental Figure 9A**-*gating scheme*) in ADT-007 vs. sham (n = 8). (**G-H and O-P**)

% intracellular cytokine (IL-2, granzyme B, and IFNγ) production (ICS) from (**G and O**)

CD4$^+$ and (**H and P**) CD8$^+$ T cells were evaluated after 5-h stimulation with

PMA/ionomycin + brefeldin A in ADT-007 vs. sham (n = 8). Column graphs - 2

independent experiments, statistical significance: Welch's t-test, error bars - SD. Tumor

growth curves - aggregate of 2 independent experiments, statistical significance: 2-way

ANOVA with Tukey's multiple comparison, error bars - SEM.