Title: A Novel Pan-RAS Inhibitor with a Unique Mechanism of Action Blocks Tumor Growth in

Mouse Models of GI Cancer

Authors: [1]Jeremy B. Foote, [2]Tyler E. Mattox, [3,12]Adam B. Keeton,[3,12]Xi Chen, [3]Forrest Smith,

[3]Kristy L. Berry, [3]Thomas Holmes, [3]Junwei Wang, [3]Chung-Hui Huang, [2]Antonio B. Ward, Amit

K. Mitra[3], [2]Veronica Ramirez-Alcantara, [1]Cherlene Hardy, [4]Karrianne G. Fleten; [4]Kjersti

Flatmark; [5]Karina J. Yoon, [6]Sujith Sarvesh, [6]Ganji Purnachandra Nagaraju, [3]Yulia Maxuitenko,

[3]Jacob Valiyaveettil, [6]Julienne L. Carstens, [7]Donald J. Buchsbaum, [8]Jennifer Yang, [9]Gang Zhou,

[10]Elmar Nurmemmedov, [10]Ivan Babic, [11]Vadim Gaponenko, [11]Hazem Abdelkarim, [12]Michael R.

Boyd, [13]Gregory S. Gorman, [14]Upender Manne, [15]Sejong Bae, [6]Bassel F. El-Rayes, and [3,12]Gary

A. Piazza

Affiliations: [1]Department of Microbiology, University of Alabama at Birmingham, Birmingham

AL, [2]University of South Alabama, Mobile, AL; [3]Department of Drug Discovery and

Development, Harrison College of Pharmacy, Auburn University, Auburn, AL;[4] Department of

Gastroenterological Surgery, Oslo University Hospital, The Radium Hospital, Oslo, Norway and

Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway;

[5]Department of Pharmacology and Toxicology, University of Alabama at Birmingham,

Birmingham AL; [6]Department of Hematology and Oncology, University of Alabama at

Birmingham, Birmingham, AL; [7]Department of Obstetrics and Gynecology, University of

Alabama at Birmingham, Birmingham, AL; [8]University of Connecticut, Mansfield, CT; [9]Georgia

Cancer Center, University of Augusta, Augusta, GA; [10]Celaris Bio LLC, San Diego, CA;

[11]Department of Biochemistry and Molecular Genetics, University of Illinois, Chicago, IL; [12]ADT

Pharmaceuticals LLC, Orange Beach, AL. [13]Department of Pharmaceutical, Social and

Administrative Sciences, McWhorter School of Pharmacy, Samford University; Birmingham, AL,

[14]Department of Pathology, University of Alabama at Birmingham, Birmingham, AL; [15]Division of

Preventive Medicine, University of Alabama at Birmingham, Birmingham, AL.

Running Title: *A novel pan-RAS inhibitor for RAS-driven GI cancers*

Key Words: KRAS, pan-RAS inhibitor, pancreatic cancer, colorectal cancer, tumor immune

microenvironment

Funding Sources: NIH/NCI R01CA254197, R01CA238514, P30CA01314, and Norwegian

Cancer Society 197837.

Corresponding Author: Gary A. Piazza, Ph.D. W.W. Walker Endowed Professor, Department

Head, Drug Discovery and Development; Director, Cancer Research Center, Harrison College

of Pharmacy, Auburn University, Auburn, AL, Walker Building, 362 W. Thach Ave, Room 3306F,

334-844-7409, gap0034@auburn.edu

Disclosure of Potential Conflicts of Interest: G.A. Piazza, A.B. Keeton, X. Chen, and M.R.

Boyd are co-founders of ADT Pharmaceuticals Inc. and consultants. All other authors declare

no potential conflicts of interest.

ABSTRACT

Here, we describe a novel pan-RAS inhibitor, ADT-007, that potently inhibited the growth of RAS mutant cancer cells irrespective of the RAS mutation or isozyme. RAS^{WT} cancer cells with activated RAS from upstream mutations were equally sensitive. Conversely, cells from normal tissues or RAS^{WT} cancer cells harboring downstream BRAF mutations were insensitive. Insensitivity to ADT-007 was attributed to low activated RAS levels and metabolic deactivation by UDP-glucuronosyltransferases expressed in normal cells but repressed in RAS mutant cancer cells. Cellular, biochemical, and biophysical experiments show ADT-007 binds nucleotide-free RAS to block GTP activation of RAS and MAPK/AKT signaling. Local administration of ADT-007 strongly inhibited tumor growth in syngeneic immune-competent and xenogeneic immune-deficient mouse models of colorectal and pancreatic cancer while activating innate and adaptive immunity in the tumor immune microenvironment. Oral administration of ADT-007 prodrug inhibited tumor growth, supporting further development of this novel class of pan-RAS inhibitors for treating RAS-driven cancers.

SIGNIFICANCE

ADT-007 is a 1st-in-class pan-RAS inhibitor with ultra-high potency and unique selectivity for cancer cells with mutant or activated RAS capable of circumventing resistance and activating antitumor immunity. Further development of ADT-007 analogs or prodrugs with oral bioavailability as a generalizable monotherapy or combined with immunotherapy is warranted.

INTRODUCTION

Colorectal cancer (CRC) and pancreatic ductal adenocarcinoma (PDA) are projected to cause 53,010 and 51,750 deaths in the US in 2024, respectively [1]. The 5-year survival rates for CRC and PDA are 63% and 13%, respectively [1]. Over 50% of CRC and 90% of PDA

patients harbor gain-in-function mutations in KRAS genes that are associated with poor

prognosis, making the development of novel RAS inhibitors an urgent unmet medical need [2].

For many cancers, RAS mutations initiate tumorigenesis that in cooperation with the

acquisition of mutations in other oncogenes or inactivating mutations in tumor suppressor genes

such as p53 [3], SMAD4 [4], APC [5], or p16 [6], drive the progression from precancerous lesions

to malignant disease. Signaling through mutant or non-mutant wild-type RAS (RAS^{WT}) is critical

for transformation [7], cancer cell proliferation and survival [7, 8], chemoresistance [9], and

metastasis [5, 10]. RAS mutations also promote immunosuppression within the tumor immune

microenvironment (TiME) through a broad array of mechanisms, including increased production

of acute inflammatory cytokines (IL-6, IL-8) [11], priming of immunosuppressive T-regs [12],

MDSCs [13], downregulation of MHCI [14], increased expression of programmed cell death ligand

1 (PD-L1) [15], and suppression of intra-tumoral interferon (IFN) signaling [16]. Cumulatively,

oncogenic RAS-mediated reprograming of the TiME results in weakened recognition and reduced

cytotoxic T and NK cell functions that support tumor progression.

The recent development of drugs directed against cancers harboring KRAS mutations at

the G12C and G12D codons has set the stage for selective targeting of KRAS-driven

malignancies [17]. Clinical trials evaluating sotorasib (AMG-510) and adagrasib (MRTX-849)

selective for the KRAS^{G12C} mutation show promising efficacy for NSCLC patients with mutant

KRAS^{G12C} but a lower response rate in patients with KRAS^{G12C} mutant CRC [18]. Preclinical data

shows much promise of KRAS^{G12C} and newly developed KRAS^{G12D} inhibitors to rewire the

immunosuppressive machinery that limits T cell recognition and cytotoxic activity [19]. However,

clinical trials indicate that combining such drugs with checkpoint immunotherapy may not be well

tolerated [20]. Adaptive resistance is another limitation of mutant-specific KRAS inhibitors [21].

Resistance may result from new activating mutations involving RAS isozymes (KRAS, HRAS, and

NRAS) or activation of RAS^{WT} isozymes from upstream receptor tyrosine kinases (RTK) [22],

making pan-RAS inhibitors an attractive therapeutic option to dampen the RAS-related adaptive

resistance circuitry. In addition, the presence of different RAS mutations in the same tumor and

the role of RAS[WT] proteins in oncogenesis [7] expands the potential efficacy advantages of RAS

inhibitors capable of inhibiting both mutant RAS and RAS[WT] activated from upstream signaling

molecules.

Here, we characterize a novel pan-RAS inhibitor, ADT-007, with the potential to

circumvent resistance and activate antitumor immunity. ADT-007 was discovered from screening

a library of compounds that share the indene scaffold of the nonsteroidal anti-inflammatory drug

(NSAID), sulindac, previously reported to preferentially inhibit the formation of RAS mutant tumors

in a rat model of chemical-induced mammary tumorigenesis [23]. Subsequent studies reported

that sulindac binds RAS and inhibits RAS-induced transformation and signaling, albeit with

modest potency and confounding inhibition of cyclooxygenase (COX) [24, 25]. The current studies

determined the mechanism of action of ADT-007 that lacks COX inhibitory activity yet is ~10,000

fold more potent than sulindac to inhibit cancer cell growth *in vitro*. We also assessed the

anticancer activity of ADT-007 using cancer cell lines representative of the complex mutational

landscape of RAS-driven cancers and immune modulatory potential using mouse tumor models

with different immune compositions representative of the diversity seen in patients with CRC and

PDA.

Material and Methods:

Chemical synthesis: ADT-007, (*Z*)-2-(5-fluoro-1-(4-hydroxy-3,5-dimethoxybenzylidene)-2-

methyl-1H-inden-3-yl)-N-(furan-2-ylmethyl)acetamide, and ADT-316, (*Z*)-4-((5-fluoro-3-(2-

((furan-2-ylmethyl)amino)-2-oxoethyl)-2-methyl-1H-inden-1-ylidene)methyl)-2,6-

dimethoxyphenyl 4-methylpiperazine-1-carboxylate hydrochloride, were synthesized as follows:

To a 250-mL round-bottom flask equipped with a gas outlet, (Z)-2-(1-(4-acetoxy-3,5-

dimethoxybenzylidene)-5-fluoro-2-methyl-1H-inden-3-yl) acetic acid (10.3 g, 25.00 mMol) was

added to 180 mL of anhydrous dichloromethane, followed by carbonyl diimidazole (CDI) (4.25 g,

26.25 mMol). The mixture was stirred for 30 minutes, followed by the addition of furfuryl amine

(2.68g, 26.25 mMol) and anhydrous pyridine (20 mL). The solution was stirred at room

temperature for 48 hours and washed with 0.2N HCl (120 mL) and water (120 mL x 2). The

organic layer was concentrated. The residue was dissolved in 300 mL of methanol, followed by

addition of ammonium in methanol (7N, 100 mL). The red mixture was stirred vigorously

overnight, then filtrated. The solid was dissolved in 400 mL of ethyl acetate, treated with 1N HCl,

and washed with brine (150 mL x 3) and dried over anhydrous sodium sulphate. After filtration

the solution was concentrated. The residue was recrystallized from acetone/ethyl acetate to

afford the ADT-007 as a yellow crystal. The mother liquor was concentrated. Purification of the

residue with silica gel column chromatography, followed by the above recrystallization

procedure gave the second crop (9.54 g combined, 85%). Treatment of ADT-007 with 4-

methylpiperazine-1-carbonyl chloride in anhydrous pyridine at 50 °C overnight, followed by

chromatography purification and acidification afforded ADT-316 HCl as a yellow crystal.

<u>Mice</u>: Eight-week-old female C57BL/6 mice (000664), male and female NOD.Cg-

$Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/SzJ (NSG, 00557) and female B6.129S7-$Rag1^{tm1Mom}$/J (Rag 1$^{-/-}$, 02216)

were purchased from Jackson Laboratory. Female B6.Ifng/Thy1.1 KI (IFN-γ.Thy1.1) and B6.Cg-

Tg-IL2$^{tm1(eGFP)/Weav}$ (IL-2.GFP) knock-in cytokine reporter mice were kindly provided by Dr. Casey

Weaver (University of Alabama at Birmingham, Birmingham, AL) [39]. 6–8-week-old female

athymic foxn1 nude mice were bred at Department of Comparative Medicine, Oslo University

Hospital (OUH). Experiments were randomized, but not blinded. All animal experiments and

procedures were performed under approved protocols in accordance with guidelines from the

University's Institutional Animal Care & Use Committee (IACUC). For animal studies performed

at OUH Institute for Cancer Research (Norwegian Radium Hospital, Oslo, Norway) all

procedures and experiments involving animals were approved by the Norwegian Food Safety

Authority (application ID #18209) and were conducted according to the recommendations of the

European Laboratory Animals Science Association and the ARRIVE guidelines.

Cell lines and patient derived xenografts: Murine PDA cell lines were derived from tumor-

bearing $Kras^{LSL-G12D/+}$; $Trp53^{LSL-R172H/+}$; Pdx1-Cre; $Rosa26^{YFP/YFP}$ (7160c2, 2838c3) and $Kras^{LSL-G12D/+}$;

$Trp53^{L/+}$; Pdx1-Cre (5363) mice and cultured in DMEM complete media as previously

indicated [36]. Human tumor cell lines (BxPC-3, AsPC-1, CFPAC1, MIA PaCa-2, PANC-1, SW-

1990, HT-29, HCT-116, A-549, SW-480, MDA-MB-231, OVCAR-8, Hs578T, SK-MEL2, H-1299,

OV-90, H-1975, U87-MG, SK-OV3), and the murine tumor lines (B16 melanoma and Lewis lung

cancer) were purchased from ATCC and maintained according to ATCC guidelines. The normal

lung epithelial cells (NHAEC) were purchased from Lonza and cultured using bronchial epithelial

growth medium per the supplier's protocol. NCM-460 cells were obtained from InCell (San

Antonio) and maintained in medium/nutrient mixture with 10% FBS for less than 12 passages,

according to the supplier's protocol. The human gall bladder adenocarcinoma expressing

$KRAS^{G12V}$ was derived from an 80-year-old male patient, with previous history of pancreatic

cancer. Originally diagnosed as PDA, the tumor was reclassified as a gall bladder

adenocarcinoma. This PDX model was serially passaged in NSG mice. The CRC PDX models

were established by implanting peritoneal metastasis samples collected at the time of

cytoreductive surgery – hyperthermic intraperitoneal chemotherapy (CRS-HIPEC) into the

peritoneal cavity of nude mice. The PMP-1 model was established from a patient with an

appendiceal primary lesion, while PMCA-1 was from a patient with a primary mucinous rectal

carcinoma [51, 52]. The PC244 model was established from a peritoneal metastasis specimen

from a 69-year-old female with a poorly differentiated mucinous adenocarcinoma of the cecum.

Targeted next-generation sequencing of PC244 was performed as previously described [53]

For studies performed at UAB and University of South Alabama (USA), all cell lines and PDX

models were tested against an extended panel of excluded pathogens including mycoplasma

prior to *in vivo* studies as mandated by UAB and USA IACUC and ARP guidelines.

<u>Tumor cell transfections:</u> BxPC-3 and HT-29 cells were initially exposed to lentiviral particles

containing a mutant HRAS^{G12V} or *KRAS*^{G12C} construct for 48 hours in a 6-well plate at 37°C in a

humidified incubator with 5% carbon dioxide. The media was then removed and replenished

with fresh media containing HA-*HRAS*^{G12V} or HA-*KRAS*^{G12C} lentiviral particles. After another 48

hours, the media was removed, and fresh media containing 5 µg/mL puromycin was added to

the well. Surviving cells were expanded in a 10 cm dish, and expression of the construct was

confirmed by WB with an HA tag antibody (Cell Signaling Technology).

HEK 293FT (Invitrogen, # R70007) cell line was transfected with either pBabe-puro-HA-

KRASG12C (Addgene, #58901) or pBabe-puro-Empty (Addgene, #1764) retroviral constructs

along with pCL-10A1 retrovirus packaging vector (Novus Biologicals, # NBP2-29542). Following

a 48-h incubation at 37°C, culture supernatants containing viral particles were collected and

mixed with PEG-itTM (SBI, #LV810A-1) virus precipitation reagent (5X) followed by overnight

incubation at 4°C. Thereafter, supernatants were centrifuged at 1500 x g for 30 min and white

pellets of retroviral particles containing either pBabe-puro-HA-KRAS G12C or pBabe-puro-

Empty retroviral construct were collected and resuspended in sterile phosphate buffered saline

(PBS). BxPC-3 cells were transduced with retroviral particles in a 6-well plate using polybrene

transfection reagent (EMD MIllipore, TR-1003-G) and incubated for 48 h at 37°C. Thereafter,

media was replaced with fresh media containing 5 µg/mL puromycin. Surviving cells were

selected, expanded, and confirmed for KRAS^{G12C} expression by WB using an HA tag antibody

(Cell Signaling Technology).

<u>ADT-007 treatment and EGF stimulation:</u> For *in vitro* ADT-007 studies, cell cultures were

incubated with increasing concentrations (0.1, 0.5, 1, 5, 10, 50, 100, 250, 500, 1000, 10000 nM)

of ADT-007, AMG-510 (Selleck Chemicals, Houston, TX) or MRTX849 (Selleck Chemicals,

Houston, TX) or vehicle (DMSO) for a period of 24 – 72 hours depending on the assay. In some

experiments, recombinant mouse EGF (Gibco, Jenks, OK) was added 10 minutes prior to

harvesting protein lysates from treated tumor cell lines to assess the impact of upstream

receptor tyrosine kinase signaling on RAS-MAPK and RAS-PI3K pathways.

<u>Cell growth assays:</u> To determine 2D growth IC_{50} values, 5×10^3 tumor cells were plated in

triplicate in 96 well flat bottom plate. To determine 3D growth IC_{50} values, 1×10^3 tumor cells

were plated in triplicate in 96 well round bottom non-binding plate. Cultures were incubated with

various concentrations of ADT-007 (0.1. 0.5. 1, 5, 10, 50, 100, 250, 500, 1000, 10000 nM) and

cultured for an appropriate period depending on cellular growth kinetics as indicated in figure

legends. At 72-96 hours (2D) or 8-10 days (3D spheroid), cells were processed according to the

manufacturer's recommendations to assess intra-cellular ATP levels as a measure of cellular

proliferation and viability using CellTiter-Glo (Promega, Madison, WI). Nonlinear dose-response

curves showing the effects of compound concentration on colony count were generated using

GraphPad Prism.

<u>Gene expression database analysis</u>: Transcriptomic (RNAseq data) for most of the cell lines in

Table 1 is publicly available in Cancer Cell Line Encyclopedia accessed via the DepMap Portal)

[30]. The Student's T-test was used to identify significant differences in 19,221 TPM+1

normalized transcripts between the two categories of cell lines. Due to the modest number of

resistant lines available, a threshold of $p < 0.01$ was chosen, which identified 719 transcripts

and was reduced to 254, unique transcripts expressed at least a 5-fold difference.

<u>NMR Spectroscopy:</u> Recombinant WT KRAS was dialyzed 2X for 24 hours (12 hours each in 1L

50 mM Tris-Citrate pH=6.5, 50 mM NaCl, 10 mM β-mercaptoethanol) at 4 °C following treatment

with 10 mM EDTA to remove Mg^{2+} and the bound nucleotide. To record ^1H-^{15}N heteronuclear

single quantum coherence (HSQC) NMR spectra of nfKRAS we used 900 MHz and 600 MHz

Avance spectrometers (Bruker, Billerica, MA, USA), equipped with cryogenic probes.

Each *in vitro* experiment was performed at 25°C in 50 mM Tris-Citrate pH=6.5, 50 mM NaCl, 10

mM β-mercaptoethanol containing 10% ^2H_2O. The chemical shift assignments were taken from

the Biological Magnetic Resonance Data Bank (BMRB) database (http://www.bmrb.wisc.edu) by

use of the BMRB accession numbers: 18529, 25114, and 26635. Chemical shift perturbations

higher than the sum of the average and 1 SD were considered to be statistically significant.

Nucleotide Exchange Assays: The fluorescent GTP analog MANT-GTP was purchased from

Millipore Sigma. For each reaction, 5 µL of 5 µM purified GDP-bound K-RAS in reaction buffer

(40 mM Tris-HCl, 100 mM NaCl, 20 mM MgCl, 1 mg/mL BSA, pH 7.5) and 2.5 µL of 200 mM

EDTA was added to a microcentrifuge tube to make a master mix. Water was used in place of

EDTA in some reactions as a control for the release of GDP. 7.5 µL of master mix per well was

added to a 384-well plate for each respective reaction. 5 µL of compound at five times the

desired concentration or vehicle in reaction buffer was added to each respective well. After 2.5

minutes for equilibration at 26°C in a Bio-Tek Synergy H4 microplate reader, 12.5 µL of 1.5 µM

MANT-GTP in reaction buffer was added to each well for a final concentration of 750 nM MANT-

GTP, 1 µM RAS, 20 mM EDTA, and the desired concentration of compound or vehicle. In the

case of compound preincubation, the plate was incubated for 1 hour at 26°C prior to MANT-

GTP addition. Fluorescence was measured at an excitation wavelength of 360 nm and an

emission wavelength of 460 nm in a Bio-Tek Synergy H4 microplate reader. Kinetic curves were

generated with GraphPad Prism.

Cellular thermal stability assays: Experiment to determine ADT-007 binding to RAS in cells was

performed by Cellaris Bio LLC (San Diego, CA) using HCT-116 and HT-29 cells cultured in

RPMI medium supplemented with 10% FBS. For an initial determination of the melting profile of

KRAS, fresh cell lysates were prepared in non-denaturing buffer was cleared via centrifugation

at 5.000 rpm for 15 min. Cleared lysates were dispensed into 96-well PCR (equivalent of

approx. 6000 cells/well/50 µl), then subjected to temperature gradient (38-65°C) for 10 min.

Subsequently, centrifugation was performed at 14.000 rpm to sediment the unstable protein

content. Following WB of the soluble fraction, bands of stable KRAS were quantified,

normalized to DMSO, and plotted using GraphPad Prism software. Non-linear regression

analysis was used to fit a Sigmoidal dose-response curve with variable slope to generate EC_{50}

values for ADT-007 binding to KRAS.

Additional cellular thermal stability experiments used a Micro-Tag enzyme

complementation method. A KRAS G12C Micro-Tag construct was engineered having a 15-

amino acid tag at the N-terminus. The construct was transiently expressed in HEK293 cells.

Two days post-transfection cells were lifted, washed with Tris Buffered Saline (TBS: 20mM Tris

HCl pH 7.4, 150mM NaCl) and re-suspended at 1000 cells/µl in TBS and 50,000 cells aliquoted

to wells of a PCR plate. Increasing doses of ADT-007 and a control compound (ADT-652) were

added and incubated with cells for 45min followed by heating for 10min at 55°C and a 1min cool

down on ice. An equal volume of 0.5% Triton-X-100 in TBS was added, and cells lysed for

10min on ice followed by addition of the Tag binding protein and substrate and immediately

proceeded to measurement of fluorescence signal. Fluorescence signal was plotted with

concentration of the compounds on a semi-log scale using GraphPad Prism 9.0 software. Non-

linear regression analysis was used to fit a Sigmoidal dose-response curve with variable slope

to generate EC_{50} values for ADT-007 binding to KRAS.

RAS Activation Assays: Cells were grown to 80 percent confluence using complete growth

medium then washed with 1 mL of PBS and lysed on ice for 5 min, (25 mM Tris HCl, 150 mM

NaCl, 5 mM $MgCl_2$, 1% NP-40, 5% glycerol, pH 7.2) with Halt Protease and Phosphatase

Inhibitor Cocktail (Thermo Fisher Scientific), then clarified at 16,000 RCF 15 minutes at 4°C.

Protein concentration of the supernatant was normalized using the DC protein assay (Bio-Rad).

200 µg of whole cell lysate was incubated with GST-RAF-RBD (Addgene) bound to glutathione

agarose beads at 4°C for 1 hour. Bound active RAS was collected by centrifugation at 6,000

RCF for 30 seconds at 4°C, washed thrice with lysis buffer, then eluted with SDS sample buffer

and subsequently detected by WB with a Pan-RAS antibody or isotype specific antibody as

described below.

<u>Western blotting:</u> Tumor or cell lysates were generated after treatment with 500 uL RIPA lysis

buffer (Thermo Scientific, 89900) with protease and phosphatase inhibitors (Thermo Scientific,

78440), boiled for 5 minutes and equal protein concentrations were loaded and run on a 12%

SDS-PAGE. Gels were transferred to a PVDF membrane, which were blocked with either 5%

non-fat dry milk in TBS-T (TBS with 0.1% Tween 20) or 5% BSA in TBS-T, depending on the

blocking agent used for the primary antibodies, for 1 hour at room temperature. Membranes

were incubated overnight at 4°C on a rocker with primary antibodies **(Supplemental Table 2)**

according to the manufacturer's instructions and then developed with HRP-conjugated

secondary antibodies (Cell Signaling Technology) for 1 hour at room temperature on an orbital

shaker. Membranes were developed using SuperSignal West Enhanced Chemiluminescent

substrate (Thermo Fisher Scientific) and imaged using a ChemiDoc Imaging System (Bio-Rad).

<u>Tumor cell implantation and ADT-007 treatment:</u> A single cell suspension of murine CRC or

PDA cells was prepared in PBS and kept on ice until injection. 1×10^6 syngeneic mouse CRC

and PDA tumor cells were injected subcutaneously into 6–8-week-old BALB/C or C57BL/6 mice,

respectively. When tumors reached an average tumor volume of 150 mm^3 mice began receiving

daily intra-tumoral (10 mg/kg, QD, 7 days/week for 17-21 days) or peri-tumoral (5 mg/kg, BID, 7

days/week for 17-21 days). Tumor growth was assessed by caliper measurements (2-3/week)

and tumor volumes (TV = W x L^2/2) were calculated throughout the study. No overt signs of

toxicity (body weight loss or organ [lungs, heart, liver, GI tract] pathology) were noted with local

administration.

Flow Cytometry: Mouse tumors were digested in a 1 mg/mL collagenase IV and 0.1 mg/mL

DNAase 1 (Worthington Biochemical, Lakewood, NJ) in HBSS for 45 minutes at 37 °C with

intermittent shaking. Samples were then washed with RPMI containing 10% FBS and filtered

through a 70µM strainer generating single cell suspensions. Cells were labeled with primary

fluorophore-conjugated antibodies and a live/dead stain (**Supplemental Table 3**) for 30-60 min

at 4 °C, washed and re-suspended in flow buffer. Cytokine expression from single cell

suspensions of spleen and tumor was quantified as follows: cells were plated at a concentration

of 10^6 cells in 1 mL of RP-10 media and stimulated for 5 hours at 37 °C using a 1x solution of a

Cell Activation Cocktail containing PMA ionomycin, and Brefeldin A (Biolegend, San Diego, CA).

Conversely, cells were cultured for 5 hours at 37°C using a solution containing 5 ug of Brefeldin

A (BioLegend, San Diego, CA) to serve as an unstimulated control. After cell surface staining,

cells were fixed in 4% paraformaldehyde for 45 minutes at 4 °C then washed with 1x

Perm/Wash (BD, 554723). For intracellular staining in tumor and spleen samples, cells were

stained with an antibody cocktail specific intracellular cytokines (IL-2, IFNγ, TNFα, and

granzyme B) in Perm/Wash solution for 60 minutes at 4 °C. Data were acquired on a Symphony

A5 flow cytometer, with analysis performed using FlowJo version 10.7.2.

Statistical Analyses: GraphPad Prism (version 9.1.2) was used for statistical analyses and

graphical representation with data are presented as either means $\pm$ standard deviation (SD) or

standard error of the mean (SEM). Two-tailed Student's t-test and Two-way Analysis of

Variance (ANOVA) with multiple corrections (Bonnferoni method) were performed for

determination of statistical significance between groups. For IC_{50} generation, concentrations

were log transformed, data were then normalized to control and *log*(inhibitor) vs. response (four

parameter nonlinear dose-response) test was used. A $p < 0.05$ was considered statistically

significant.

<u>Data Availability:</u> The data generated in this study are available upon request from the

corresponding author.

RESULTS

ADT-007 potently and selectively inhibits the growth of cancer cells with mutated or

activated RAS

A library of indenes was synthesized and screened for differential growth inhibitory

activity using HCT-116 CRC cells harboring a $KRAS^{G13D}$ mutation vs. HT-29 CRC cells with

RAS^{WT} but mutant $BRAF^{V600E}$. A lead compound, ADT-007 (**Figure 1A**), and a series of analogs

emerged following extensive chemical optimization for potency and selectivity. ADT-007

inhibited the growth of HCT-116 cells with an IC_{50} of 5 nM, while HT-29 cells were

approximately 100-fold less sensitive with an IC_{50} of 493 nM (**Figure 1B**). As expected, HCT-

116 cells expressed high levels of activated RAS (GTP-RAS) as measured by RAS-RBD

pulldown assays that were undetectable in HT-29 cells (**Figure 1B**). Despite differences in

activated RAS levels, both cell lines depended on MAPK signaling for growth, given comparable

sensitivity to inhibitors targeting downstream kinases, RAF, MEK, and ERK in growth assays

(data not shown). The requirement of mutant or activated RAS for sensitivity to ADT-007 was

determined by transfecting HT-29 cells with $HRAS^{G12V}$. ADT-007 treatment of transfected cells

resulted in an IC_{50} of 24 nM in growth assays compared with IC_{50} values of 549 and 730 nM for

parental and vector control cells, respectively (**Figure 1C**).

We further evaluated the growth inhibitory activity of ADT-007 using a large panel of

histologically diverse cancer cell lines with various mutations in RAS isozymes or upstream

signaling components that activate or regulate RAS such as RTK or GTPase activating proteins

(GAPS), respectively (**Table 1**). Cancer cell lines with mutations in K/N/HRAS isozymes or

upstream signaling components (e.g., PDGF, EGFR, or NF1) were uniformly sensitive to ADT-

007 with growth IC_{50} values in the single-digit nM range. In contrast, cancer cell lines with

RAS[WT] but with downstream BRAF mutations were essentially insensitive to ADT-007. Normal

colonocytes (NCM460) or airway epithelium cells (NHAEC) were also insensitive to ADT-007

treatment.

KRAS[G12C] MIA PaCa-2 PDA cells were highly sensitive to ADT-007 with IC_{50} values as

low as 2 nM in growth assays, resulting in a selectivity index approaching 1200-fold relative to

RAS[WT] BxPC-3 PDA cells (**Table 1 and Supplemental Figure 1A**). RAS-RBD pulldown assays

confirmed high levels of GTP-RAS in MIA PaCa-2 and three other PDA cell lines that were

sensitive to ADT-007 and harbor KRAS[G12V] and KRAS[G12D] mutations (**Supplemental Figure**

1B), representing the most prevalent KRAS mutations in PDA patients [26]. By comparison,

activated RAS levels in RAS[WT] BxPC-3 cells were undetectable in RAS-RBD pulldown assays.

As observed for HT-29 cells, transfecting BxPC-3 cells with mutant KRAS[G12C] induced sensitivity

to ADT-007 in growth assays (**Supplemental Figure 1C**). The introduction of activated RAS

BxPC-3 cells by transfection was confirmed by RBD pulldown assays (**Supplemental Figure**

1D).

Other experiments showed that ADT-007 treatment of MIA PaCa-2 cells comparably

blocked growth in 2D monolayer and 3D spheroid cultures, resulting in unusually steep dose-

response curves, which was a characteristic feature of ADT-007 and related analogs, indicating

a highly uniform response of the cells to treatment (**Supplemental Figure 1E-F**). ADT-007 also

completely inhibited the colony formation of KRAS mutant PDA cell lines without significantly

affecting the colony formation of BxPC-3 cells (**Supplemental Figure 1G**).

ADT-007 inhibits activated RAS and MAPK/AKT signaling in human CRC and PDA cell

lines

We next evaluated the effect of ADT-007 on RAS activation and RAS-dependent

signaling in RAS mutant cancer cell lines. ADT-007 caused a concentration-dependent

reduction of RAS-GTP levels and phosphorylated c-RAF, MEK, ERK, and AKT components of

MAPK and PI3K signaling in HCT-116 cells (**Figure 1D**). Similar effects of ADT-007 treatment

on levels of phosphorylated MAPK and PI3K signaling components were observed in MIA

PaCa-2 cells that were serum-starved and treated with ADT-007, followed by EGF stimulation

(**Figure 1E**). ADT-007 reduction of RAS-GTP levels occurred under multiple conditions when

MIA PaCa-2 cells were grown in the presence of serum, serum-starved, or serum-starved and

EGF stimulated (**Supplemental Figures 2A-C**). Pan-RAS inhibitory activity of ADT-007 was

confirmed by experiments showing that ADT-007 treatment of MIA PaCa-2 cells reduced levels

of GTP-HRAS^{WT} and GTP-NRAS^{WT} isozymes that were co-expressed with GTP-KRAS^{G12C}

(**Figure 1F**). As expected, EGF did not increase GTP-KRAS^{G12C} levels but only activated GTP-

HRAS^{WT} and GTP-NRAS^{WT} isozymes. Nonetheless, ADT-007 reduced mutant KRAS isozyme

levels to the same extent as EGF-activated RAS^{WT} isozymes.

Comparison with mutant-specific KRAS inhibitors

We also compared the anticancer activity of ADT-007 with the KRAS^{G12C} inhibitor, AMG-

510 (sotorasib), and the pan-KRAS inhibitor, BI-2865. AMG-510 and BI-2865 were significantly

less potent and caused much less growth inhibition than ADT-007 in growth assays involving

MIA PaCa-2 cells (Figure 1**G**). In addition, the growth of LLC cells harboring dual mutations,

KRAS^{G12C} and NRAS^{Q61H}, was only partially inhibited by BI-2865, whereas ADT-007 retained

activity (**Figure 1H**). LLC cells were insensitive to AMG-510 (**Figure 1I**), as previously reported

[27], but retained sensitivity to ADT-007. These results show that ADT-007 has appreciably

greater intrinsic anticancer activity and maintains potency in the presence of two independent

activating RAS mutations compared with sotorasib, which is known from clinical trials to have

modest efficacy and prone to resistance from the acquisition of additional cis- or trans-mutations

in RAS alleles [21, 28].

ADT-007 blockade of RAS activation and MAPK/AKT signaling induced cell cycle arrest and apoptosis

We also examined the effect of ADT-007 treatment on cell cycle distribution and apoptosis. ADT-007 induced G2/M cell cycle arrest in HCT-116 and MIA PaCa-2 cells at concentrations within the range of growth IC_{50} values, while HT-29 and BxPC-3 cells displayed profoundly reduced sensitivity (**Supplemental Figure 3A-D**). ADT-007 also increased levels of active, cleaved caspase-3 protein in MIA PaCa-2 cells, which followed effects on cell cycle arrest (**Supplemental Figure 3E**). Similar findings were noted with HCT-116 cells and by measuring apoptosis using Annexin as a marker of apoptosis (data not shown). Given the effect of ADT-007 on G2/M arrest and previous studies suggesting that certain other indenes non-selectively bind tubulin [29], the impact of ADT-007 on tubulin polymerization was determined using recombinant α-tubulin. ADT-007 exhibited a weak inhibitory effect on tubulin polymerization at a concentration 100-fold above its growth IC_{50} value, making a direct link between effects on tubulin polymerization and cell cycle arrest unlikely (**Supplemental Figure 3F**). Similar weak effects on tubulin polymerization were observed by treating RAS mutant cancer cells with ADT-007 at growth IC_{50} values.

Metabolic mechanisms impacting resistance and sensitivity to ADT-007

As described above, ADT-007 reduced activated RAS at concentrations effective for inhibiting growth. The potency and selectivity of ADT-007 to inhibit RAS-dependent growth and reduce activated RAS was significantly diminished when the hydroxyl moiety on the phenolic ring was substituted with a methoxy moiety as shown with a trimethoxy analog, ADT-106 (**Figure 2A-B**). The results show exquisite chemical selectivity for inhibiting RAS-dependent growth and biological selectivity linked with reducing activated RAS.

As discussed below*, ex vivo* treatment of mouse liver microsomes with ADT-007 followed by LC-MS/MS analysis of metabolites revealed that the hydroxyl moiety is rapidly

glucuronidated by UDP-glucuronosyltransferases (UGTs) (**Supplemental Figure 5A-B**). These

results suggested that high UGT levels known to be expressed in normal cells could account for

insensitivity to ADT-007. Measurement of UGT protein levels by WB using a pan-UGT1

antibody revealed lower UGT expression in sensitive RAS mutant HCT-116 and MIA PaCa-2

cells lines compared with insensitive RAS[WT] HT-29 and BxPC-3 cell lines (**Figure 2C-D**).

Evaluation of transcriptomic data (RNAseq) of most of the cell lines in **Table 1** (Cancer Cell Line

Encyclopedia) [30] describing mRNA expression of six distinct UDP-glucuronosyltransferase

isoforms (UGT1A1, 1A6, 1A10, 2A1, 2B10, and 2B15) provided additional evidence for lower

UGT expression RAS mutant sensitive vs. RAS[WT] sensitive cell lines to ADT-007 (**Table 1,**

Supplemental Table 1). Measurement of ADT-007 levels by LC-MS/MS from cells treated

overnight with ADT-007 revealed 350-fold higher intracellular levels in sensitive RAS mutant

HCT-116 cells than insensitive RAS[WT] HT-29 cells. Consistent with the ability of UGTs to

deactivate ADT-007, a higher ratio of the glucuronide-conjugated form of ADT-007 was

measured in HT-29 cells, while none was detected in HCT-116 cells (**Figure 2E-F**).

Experiments using a competitive UGT inhibitor, propofol, supported these observations by

showing that pretreatment increased the sensitivity of RAS[WT] HT-29 and BxPC3 cells to ADT-

007 in growth assays, decreasing the IC_{50} value by 26- and 18-fold, respectively, without

significantly affecting the sensitivity of RAS mutant HCT-116 or MIA PaCa-2 cells to ADT-007

(**Supplemental Table 4**).

ADT-007 binds RAS

The ability of ADT-007 to directly bind KRAS in cells was measured using two different

cellular thermal stability assays based on the principle that a protein maintains thermal stability

when in a conformation favorably for binding to a small molecule inhibitor. Initial experiments

measured the optimal temperature to denature 75% (T_{agg75}) KRAS in whole cell lysates,

followed by WB quantification of non-denatured KRAS levels. Treating HCT-116 lysates with

ADT-007 increased KRAS^{G13D} thermal stability (T$_{agg75}$), resulting in an EC$_{50}$ of 0.45 nM for ADT-

007 to bind KRAS. By comparison, ADT-007 did not affect the thermal stability of KRAS^{WT} in

HT-29 cells **(Figure 3A)**. As a control, ADT-007 did impact the thermal stability of an unrelated

protein (PDE10) in either HCT-116 or HT-29 cells (data not shown), previously reported to be

comparably expressed in both cell lines [31]. Additional experiments were conducted involving

ADT-007 treatment of intact HEK293 embryonic kidney cells transfected with epitope-tagged

mutant KRAS ^{G12C} and detection of thermal stability by WB analysis with anti-Micro-Tag

antibody. These experiments resulted in an EC$_{50}$ value of 8.5 nM for ADT-007 to bind KRAS

(**Figure 3B**). A trimethoxy analog of ADT-007 (ADT-652) lacking RAS selectivity in growth

assays used as a control compound did not significantly bind KRAS in transfected KRAS ^{G12C}

HEK293 embryonic kidney cells. These results indicate direct binding of ADT-007 to mutant

RAS in cells with an affinity consistent with potency to inhibit growth.

Molecular docking studies using KRAS^{WT} (PDB ID 4OBE) [32] were conducted to

understand how ADT-007 may bind RAS. The studies used molecular dynamics and induced fit

docking to identify potential binding sites for ADT-007 on KRAS. Simulations were performed

with either Mg^{+2} present or Mg^{+2} removed since it was uncertain which conformation of RAS

would be the more favorable for ADT-007 binding **(Supplemental Figure 4A-B)**. A difference

was observed in the orientation of ADT-007 within its predicted binding sites when Mg^{+2} was

present versus when absent **(Supplemental Figure 4C-D)**. When Mg^{+2} was present, the

phenolic OH group of ADT-007 was predicted to H-bond with S17 through a water bridge and

with E31 through a direct H-bond. When Mg^{+2} was absent, the phenolic OH was predicted to

interact with D119 via a water bridge and A146 by a direct H-bond. In simulations of Mg^{+2} being

present and absent, the best GLIDE docking scores (-11.08 and -12.06, respectively) were

calculated with ADT-007 binding to a site partially overlapping the nucleotide binding site that

was associated with an altered conformation of the switch-1 region **(Supplemental Figure 4A-**

B). Regardless if Mg^{+2} was present or absent, the GLIDE docking score of GDP to bind RAS (-

9.6 and -8.80, respectively) was considerably less favorable than ADT-007. These results

suggested that a conformational change induced by the binding of ADT-007 may decrease the

affinity for nucleotide. The results also indicate there may be a preference for ADT-007 binding

to RAS in a conformation when Mg^{+2} is absent as nucleotide-free RAS (nf-RAS) and that ADT-

007 bound better to this site than nucleotide.

ADT-007 binds nucleotide-free RAS and inhibits GTP activation

Next, we determined whether ADT-007 could inhibit GTP loading to nf-RAS and GDP-

bound RAS using a recombinant full-length $KRAS^{WT}$ protein. To confirm the binding of ADT-007

to nf-RAS, recombinant full-length $KRAS^{WT}$ was treated with EDTA to chelate the Mg^{+2}, thereby

simulating GEF removal of GDP, producing the apoenzyme bearing the nucleotide-free

conformation of RAS. ADT-007 treatment of RAS before GTP addition inhibited RAS activation

as determined by measuring GTP-RAS levels with RAS-RBD pulldown assays. This effect

occurred at concentrations comparable to growth IC_{50} values and to a similar extent as GDP-

RAS controls (**Figure 3C**). However, there was a negligible effect of ADT-007 on activated RAS

levels when RAS was preloaded with GTP (**Figure 3D**). These results are consistent with

molecular modeling observations suggesting that nf-RAS is the preferred conformation for ADT-

007 to bind RAS. In addition to RAS-RBD pulldown assays, we utilized a nucleotide exchange

assay to determine the effect of ADT-007 on the kinetics of GTP binding to $KRAS^{WT}$ using a

fluorescent GTP analog (Mant-GTP). These experiments showed that ADT-007 reduced the

rate of GTP binding to RAS if pretreated with EDTA to chelate Mg^{2+} (**Figure 3E**) but not after

adding non-hydrolyzable GTPγS (**Figure 3F**). A tool compound, CID1067700, previously

reported as a pan GTPase inhibitor, also inhibited Mant-GTP binding to RAS to a comparable

extent as ADT-007 (data not shown) [33].

We also used protein nuclear magnetic resonance (NMR) spectroscopy to define the

molecular determinants for these biochemical findings. First, we compared ^{15}N-HSQC spectra of

GDP-KRAS^{WT} protein alone and GDP-KRAS^{WT} in the presence of ADT-007. The lack of spectral

differences under these conditions suggests a low likelihood of direct binding of ADT-007 to

inactive GDP-KRAS (data not shown). To further investigate the binding of ADT-007 to RAS,

Mg^{2+} was chelated with EDTA to generate nf-RAS. These conditions allow the dissociation of

GDP from KRAS, producing a sample with NMR spectral features distinct from GDP-KRAS.

Treatment of nf-RAS with ADT-007 led to attenuation of several HN signals, including signals for

G13, a hotspot mutation site in the P-loop [34], and at K16 (adjacent to S17) (**Figure 4**), which

is critical for Mg^{2+} binding [35]. These discrete signals can readily be assigned due to their

unique resonance frequencies. NMR signal attenuation in the presence of ADT-007 indicates

direct binding of ADT-007 to KRAS, leading to an intermediate exchange between the

compound-occupied and free KRAS or altered backbone dynamics in the compound binding

site.

ADT-007 inhibits mouse colon tumor growth and RAS signaling *in vivo*

As mentioned above, *ex vivo* treatment of mouse liver microsome with ADT-007 followed

by LC-MS/MS analysis of metabolites identified a metabolic vulnerability of ADT-007 to be

rapidly metabolized by glucuronidation (**Supplemental Figure 5A-B**). ADT-007 was, however,

stable in mouse plasma (**Supplemental Figure 5C**), which prompted studies to evaluate

antitumor activity from localized delivery to minimize hepatic glucuronidation. Antitumor activity

of ADT-007 was initially evaluated at a dose of 5 mg/kg for 14 days (BID) by peri-tumoral

injection. In the KRAS^{G12V} gall bladder adenocarcinoma (GBA) PDX model, ADT-007 completely

arrested tumor growth and induced partial regression (**Figure 5A, left and right panels**). ADT-

007 also caused growth arrest in a KRAS^{G12C} PDA PDX model by peritumoral injection (data not

shown). Localized treatment by intraperitoneal injection of ADT-007 was also determined using

three peritoneal metastasis PDX models bearing mutant KRAS^{G12A} (PMCA-1) (**Figure 5B**),

KRAS^{G12D} (PMP1) (**Figure 5C**), or KRAS^{WT} and PDGFR^{T922I} (PC244) (**Figure 5D**).

Representative growth curves (**Figure 5B-D, left panels**) and tumor volumes at the end of

treatment (right panels) are displayed for each model system (**Figure 5B-D, right panels**).

ADT-007 significantly reduced tumor growth in all three models. Of the three models, PMCA-1

was the most sensitive, where the growth inhibitory activity persisted for a substantial time after

treatment discontinuation, and two mice were deemed as cured or tumor-free on day 100.

Consistent with experiments described above involving cancer cell lines (**Table 1**), the PC244

PDX model expressing RAS^{WT} isozymes, but a suspected activating mutation in PDGFR, was

comparably sensitive to ADT-007 *in vivo*. Furthermore, reductions in mucin production were

notable by MRI (**Supplemental Figure 6A**) with tumor volumes exhibiting a dose dependency

in each model (**Supplemental Figure 6B-D**). These findings indicate that ADT-007 induces

tumor growth arrest and, in some cases, tumor regression of established and metastatic PDX

PDA and CRC tumors in immune-deficient recipient mouse models.

The antitumor activity of ADT-007 was also determined against an array of syngeneic,

CRC and PDA tumor models bearing KRAS^{G12D} mutations using immune-competent mice.

When administered locally by intratumor (IT) injection, ADT-007 (10 mg/kg, qd) strongly

inhibited SC tumor growth (T/C$_{max}$ of 27%) in the KRAS mutant CT26 colon tumor model

(**Figure 5E, left panel**) with an 80% reduction in mean tumor growth compared to sham-treated

mice (**Figure 5E, right panel**), which correlated with reduced levels of activated RAS and

phosphorylated MAPK components (RAF, MEK, and ERK) compared with control mice

(**Supplemental Figure 7**). Peritumoral administration of ADT-007 (5 mg/kg, BID, administered

for 17-24 days) induced growth arrest in syngeneic mouse PDA models including previously

characterized cell lines 2838c3 (**Figure 5F, left panel**) and 7160c2 (**Figure 5G, left panel**) [36],

and a novel KPC cell line 5363 (**Figure 5H, left panel**) with variable patterns of growth arrest to

regression noted (**Figure 5F-H, right panels**). These findings indicate that ADT-007 inhibits *in*

vivo growth of mouse and human tumor models bearing different KRAS mutations or those

exhibiting hyperactivation of WT KRAS.

We next developed a prodrug approach to achieve systemic levels of ADT-007 by oral

administration. A carbamate prodrug of ADT-007 coded as ADT-316 was synthesized to protect

the vulnerable hydroxyl group from glucuronidation (**Supplemental Figure 5D**). Oral

administration of ADT-316 provided plasma levels of ADT-007 approximately 10-fold higher

than growth inhibitory IC_{50} values (**Supplemental Figure 5E**) and displayed antitumor activity in

a SC CT26 murine CRC model at a dosage that was well tolerated (**Supplemental Figure 5F**).

However, PK analysis of plasma levels of ADT-316 and ADT-007 revealed incomplete

conversion of the prodrug to ADT-007 (**Supplemental Figure 5E**), which likely accounts for its

modest antitumor activity and supports the need for further development of prodrugs or analogs

as discussed below.

ADT-007-mediated inhibition of KRAS^{G12D} alters the tumor immune microenvironment

(TiME)

Given the impact of mutant-specific KRAS^{G12C} [19, 37] and KRAS^{G12D} inhibitors [38] on

tumor-specific immunity, as previously reported, we assessed the activity of ADT-007 in

syngeneic, immune-competent mouse models of CRC and PDA-bearing KRAS^{G12D} mutation. To

evaluate the effect of ADT-007 on the PDA TiME, we studied two genetically characterized cell

lines bearing increased T cell infiltrate (2838c3 and 7160c2) but exhibiting different abundances

of myeloid cells, basal levels of T cell activation, and function [36]. At the termination of

treatment, 2838c3 and 7160c2 tumor-bearing mice were euthanized, and tumors were collected

to evaluate changes in YFP⁺ tumor and CD11b⁺ CD11c⁺ myeloid cell subsets within the TiME.

In both models, ADT-007 decreased the proportion of YFP⁺ tumor cells (**Figure 6A and I**).

Using multiparameter flow cytometry (**Supplemental Figure 8A**), proportions of myeloid

subsets with the 7160c2 and 2838c3 PDA TiME were evaluated. A direct comparison between

myeloid abundances in each model (sham-treated) revealed that 2838c3 had an increase in

overall proportions and abundance of myeloid cells compared to 7160c2 with significant

increases in Ly6G$^+$ Ly6C$^+$ granulocytes (**Supplemental Figure 8B**). After treatment, an

increase in proportions of CD11b$^+$ Ly6G$^+$ Ly6C$^+$ gMDSC and CD11b$^+$ Ly6G$^-$ Ly6C^{hi} mMDSC

subsets was noted in 7160c2, but in ADT-007 treated 2838c3 tumor-bearing mice, which

exhibited a decrease in macrophages (**Figure 6B***,7160c2* and **5J***,2838c3*). No changes in

dendritic cell subset (cDC1, cDC2, CD103$^+$ DC) numbers were noted in either model (**Figures**

6C*,7160c2* and **5I-K***,2838c3*). Analysis of CD11b$^+$ F4/80$^+$ macrophage phenotypes in both

models revealed trends for decreasing MHCII and CD206 expression (**Supplemental Figure**

8C*,7160c2* and **Supplemental Figure 8D**,2838c3).

Next we evaluated the importance of T and B cells in ADT-007 efficacy in 2838c3 and

7160c2 PDA models using both immune-competent C57BL/6 mice or immune-deficient

C57BL/6 Rag 1$^{-/-}$ mice. Treatment with ADT-007 reduced growth in both immune-competent

C57BL/6 and immune-deficient C57BL/6 Rag 1$^{-/-}$ mice, however, efficacy was reduced in

C57BL/6 Rag 1$^{-/-}$ recipient mice with both models (**Figure 6D***,7160c2* and **6L***,2838c3*). The

effect of ADT-007 on T and NK cell numbers, activation, and cytokine production in 7160c2 and

2838c3 TiME was evaluated using multiparameter flow cytometry (**Supplemental Figure 9A**).

An initial comparison between sham treatment groups for 2838c3 and 7160c2 revealed a

modest but statistically insignificant increase in overall CD3$^+$ T cell numbers in the 2838c3

model (**Supplemental Figure 9B**). However, after ADT-007 treatment numbers of γδ, NKT, and

NK cells increased within the TiME in both models (**Figure 6E-F***, 7160c2* **and 6M-N***, 2838c3*).

Additionally, in the 2838c3 TiME, an increase in CD3$^+$ CD4/8$^-$ TCRγδ$^-$ DN T cells was noted

(**Figure 6M**). The activation/memory status of T cell subsets was evaluated by CD44 and

CD62L expression (**Supplemental Figure 9A**). T cell phenotypes in the 7160c2 TiME shifted to

a CD44^{hi} CD62L$^-$ effector memory phenotype with ADT-007 treatment (**Supplemental Figure**

9C). In comparison, ADT-007 treatment of 2838c3 tumors increased proportions of pre-effector

and effector memory phenotypes in the CD4$^+$ Foxp3$^-$ and CD3$^+$ TCRγδ$^+$ T cell subsets

(**Supplemental Figure 9D** and data not shown). Expression of PD-1 increased on CD4 and

CD8$^+$ T cells in 7610c2 (**Supplemental Figure 10A**, *7160c2* and **B***, 2838c3*), but not 2838c3

TiME, despite higher baseline levels of PD-1 on T cells (**Supplemental Figure 10C**). Evaluation

of PD-1 and TIM-3 co-expression as an indicator of an exhausted T cell state revealed a modest

trend for co-expression in T cells in 7160c2 but not 2838c3 TiME (**Supplemental Figure 10D-**

E). Analysis of T cell infiltrates in the CT-26 TiME revealed trends for increased numbers, PD-1

expression (not statistically significant), and reduced CTLA-4 expression in both CD4$^+$ and CD8$^+$

T cell subsets (data not shown).

Functional assessments of IL-2, IFNγ, TNFα, and granzyme B production in T cell

subsets from 7160c2 and 2838c3 TiME was performed by intracellular staining after PMA and

ionomycin stimulation. In the 7160c2 TiME both CD4$^+$ and CD8$^+$ T cells exhibited increased

expression of IL-2, granzyme B, IFNγ, and TNFα (**Figure 6G-H**), whereas minimal changes in

cytokine expression were noted in T cells within the 2838c3 TiME (**Figure 6O-P**). Functional

differences as evidenced by the production of IL-2, IFNγ, TNFα, and granzyme B, were not

apparent in CD4$^+$ and CD8$^+$ T cells within the CT-26 TiME (data not shown). Using an IFNγ-

Thy1.1 reporter model system [39] we assessed vivo IFNγ production marked by surface Thy1.1

on innate and adaptive T and NK subsets as an indicator of cytotoxic activity. Consistent with

ICS resulst, ADT-007 increased IFNγ production in T cell subsets in the 7160c2, but not 2838c3

PDA TiME. In NK, NKT, and γδ T cells, IFNγ was unchanged (**Supplemental Figure 11A***,*

7160c2 **and 11B***, 2838c3*) whereas, a significant decrease in Thy1.1 was noted in NK cells from

ADT-007 treated tumors (**Supplemental Figure 11A***, 7160c2* **and 11B***, 2838c3*). Overall, these

findings show that ADT-007 modulates the innate and adaptive immunity in CRC and PDA TiME

and suggests a role for the adaptive immune system in enhancing ADT-007 efficacy.

Discussion

ADT-007 potently and selectively inhibited the growth of cancer cell lines with mutant

RAS or activated RAS^{WT} from mutations in upstream signaling molecules by binding RAS to

block GTP-activation of MAPK/AKT signaling, resulting in cell cycle arrest and apoptosis.

Despite its broad anticancer activity, ADT-007 displayed exquisite selectivity as cancer cell lines

with RAS^{WT} and downstream mutation in BRAF, as well as cells from normal tissues, were

essentially insensitive in growth assays. ADT-007 displayed high affinity to bind RAS in cells

with EC_{50} values in the same range as concentrations required to inhibit growth, colony

formation, and induce cell cycle arrest and apoptosis, which were also in the same range as

concentrations effective for reducing RAS-GTP levels and blocking MAPK/AKT signaling. The

inhibitory effect of ADT-007 on RAS-GTP levels was apparent under standard growth

conditions, serum-free conditions, or following EGF stimulation. As a pan-RAS inhibitor, ADT-

007 reduced RAS-GTP levels of mutant KRAS and co-expressed WT HRAS and NRAS

isozymes, which we suspect will be advantageous over mutant-specific or pan-KRAS inhibitors.

Localized delivery of ADT-007 strongly inhibited tumor growth in immunodeficient and

syngeneic murine models of CRC and PDA with partial dependence on the adaptive immune

system. ADT-007 activated T cells and enhanced their cytotoxic functions in $KRAS^{G12D}$ mutated

CRC and PDA, immune-competent mouse models. Oral delivery of a prototypic prodrug, ADT-

316, generated adequate plasma levels to inhibit tumor growth but with modest efficacy, which

we attribute to incomplete hydrolysis of the prodrug moiety to ADT-007. Recognizing the

limitations of ADT-316, we designed a new series of prodrugs that are currently under study,

one of which shows robust antitumor activity in mouse, human cell line, and patient-derived

models of PDA and is in further development by the National Cancer Institute Stepping Stones

Program.

ADT-007 specificity for RAS binding

ADT-007 appears to preferentially bind nf-RAS to block GTP activation of effector

interactions, which, to our knowledge, is the first-ever reported small molecule to inhibit RAS by

this mechanism. Molecular modeling studies revealed that ADT-007 generated the highest

docking score when RAS is in a nucleotide-free conformation. Biochemical experiments using

recombinant RAS showed that ADT-007 inhibited GTP binding to RAS and GTP-dependent

RAS-RAF binding under nucleotide-free conditions. NMR spectroscopy confirmed these

observations by showing binding to nf-RAS but not nucleotide-loaded RAS. Previous studies

reporting that a fraction of RAS exists in cells in a nucleotide-free state [40] support the ability of

ADT-007 to inhibit RAS-dependent growth by binding nf-RAS within the nucleotide-binding

domain to block GTP activation of effector interactions.

Cellular thermal stability assays provided additional evidence for direct binding of ADT-

007 to RAS in cells, showing that exposure of lysates from $KRAS^{G13D}$ mutant HCT-116 cells to

ADT-007 resulted in sub-nanomolar EC_{50} values to bind RAS but not in RAS^{WT} HT-29 cells. In

addition, ADT-007 treatment of intact $KRAS^{WT}$ HEK293 cells transfected with a $KRAS^{G12C}$

mutant allele resulted in EC_{50} values for binding RAS comparable to growth IC_{50} values. The

lack of binding of ADT-007 in HT-29 cells may be attributed to low levels of activated RAS or, as

discussed below, also attributed to metabolic deactivation by UGT isozymes remaining active in

cell lysates. Nonetheless, results from experiments involving transfection of mutant RAS into

RAS^{WT} cancer cells led us to conclude that activated RAS from either a mutation or upstream

signaling is necessary and sufficient for the sensitivity of cancer cells to ADT-007.

Metabolic deactivation of ADT-007 by UGT-mediated glucuronidation is also a significant

factor impacting sensitivity to treatment that has implications for the interpretation of *in vitro* and

in vivo experiments. Liver microsome experiments revealed that ADT-007 is rapidly

glucuronidated by UGT enzymes, ubiquitously expressed in normal tissues, and known to

metabolize compounds with phenolic OH moieties. Unexpectedly, cancer cells with WT RAS

had high mRNA and protein levels of major UGT isozymes expressed in normal cells. In

contrast, cancer cells with mutant RAS had low levels of UGT isozymes. Measurement of

cellular levels of ADT-007 following overnight treatment with ADT-007 using LC-MS/MS

confirmed high ADT-007 levels and undetectable ADT-007 glucuronide levels in HCT-116 cells,

but low ADT-007 levels and high ADT-007 glucuronide levels in HT-29 cells. In addition,

pretreatment with a known competitive UGT inhibitor, propofol, increased sensitivity in resistant

HT-29 and BxPC-3 cancer cell lines with low activated RAS to ADT-007. Propofol pretreatment

did not significantly impact cells with mutant KRAS and low UGT levels. These results suggest

that UGT-mediated glucuronidation allows bystander cells from normal tissues to be protected

from pan-RAS inhibition by ADT-007 and contributes to its unique selectivity despite previous

reports that knock-out of all three RAS isozymes is lethal to mice [41, 42]. While UGT isozymes

are expressed in metabolic tissues such as the liver, colon, and kidney, they are also expressed

in other normal tissues, including blood cells, and regulated by multiple signaling pathways and

transcription factors [43]. These results support further study of the relationship between RAS

mutational status and UGT isozyme expression.

Other indenes chemically related to ADT-007 were previously reported to inhibit cancer

cell growth by a dual mechanism involving inhibition of cyclic guanine monophosphate

phosphodiesterase (cGMP PDE) activity and tubulin polymerization [29]. The former was based

on reports that certain sulindac derivatives inhibit cGMP PDE activity at concentrations that

inhibit CRC cell growth [23, 44]. The latter was hypothesized due to the chemical resemblance

of certain indenes containing a trimethoxy substitution with colchicine, which is well known to

bind tubulin and disrupt microtubule assembly [45]. We, therefore, determined if ADT-007

inhibited cGMP PDE activity by measuring effects on cGMP hydrolysis using recombinant

PDE10, a cGMP-degrading PDE isozyme, or capacity to bind PDE10 in cells using cellular

thermal stability assays. However, we found no evidence of PDE10 inhibition or binding at

concentrations within a range that effectively inhibited growth. In addition, ADT-007 only

inhibited tubulin polymerization at concentrations ~100-fold higher than growth IC_{50} values.

These observations indicate that neither cGMP PDE nor tubulin polymerization inhibition likely

accounts for the anticancer activity of ADT-007. Moreover, reports that tubulin inhibitors induce

MAPK pathway activity [46], rather than inhibit MAPK signaling provide further evidence that

ADT-007 does not target tubulin. However, the likelihood that ADT-007 inhibits RAS and, at

higher concentrations, also inhibits tubulin polymerization suggests additional therapeutic

advantages, given that both RAS and tubulin are well-validated anticancer targets.

ADT-007 inhibits a wide range of RAS-addicted tumors and circumvents resistance

involving reactivation of the RAS pathway

Unlike mutant-specific KRAS or pan-KRAS inhibitors [47], ADT-007 exhibited broad

anticancer activity for mutant K/H/NRAS isozymes inhibiting downstream MAPK/AKT signaling,

resulting in cell cycle arrest and apoptosis. The broad anticancer activity of ADT-007 supports

its potential to be developed as a generalizable monotherapy for patients with PDA, given that

94% of patients diagnosed with this malignancy have multiple KRAS mutations [26] and a

significant proportion of patients with WT KRAS have RTK fusions that would be predicted to

have hyper-activated RAS^{WT} isozymes [48]. Furthermore, evidence indicates that ADT-007

exhibits high potency in tumor cells with activated RAS signaling, which may provide greater

therapeutic utility in patients with metastatic disease in which the proliferation of cancer cells

depends upon activated KRAS signaling for survival in the metastatic niche [49]. ADT-007

exhibits potent growth inhibition *in vitro* and *in vivo* with syngeneic KRAS^{G12D} mutated CRC and

PDA cell lines and human KRAS^{G12C} and KRAS^{G12V} mutated PDA and CC PDX models

consistent with its pan-RAS inhibitory activity.

Equally important to the broad activity of this class of compounds is the potential to

provide a second-line therapy for patients with RAS mutant cancers that develop resistance to

mutant-specific KRAS inhibitors. In NSCLC patients who go on to develop resistance to

KRAS[G12C] mutant-specific inhibitors, the majority of resistance pathways involve further

modulation of RAS signaling, including polyclonal mutations in RAS or upstream receptor

tyrosine kinases or GTPase that result in dysregulated RAS[WT] signaling [7, 22]. Our findings

indicate that ADT-007 fully maintains activity in MIA PaCa-2 cell lines that develop resistance to

sotorasib and adagrasib (not shown) and in LLC cells with KRAS[G12C] NRAS[Q61H] dual mutations

[27].

ADT-007 modulates the tumor immune microenvironment

ADT-007-mediated *in vivo* tumor growth inhibition appears to be partially dependent

upon the adaptive immune response as evidenced by reduced growth inhibitory activity of ADT-

007 on 2838c3 tumor cell growth in immune-deficient Rag 1[-/-] mice lacking B and T cells. Similar

findings have been observed with KRAS[G12C] [19, 37] and KRAS[G12D]-specific inhibitors [38],

which demonstrated that T cells either contributed to the efficacy of targeted mutation-specific

KRAS inhibitors or are critical for their long-term control of regrowth from microscopic disease

[38] in different syngeneic mouse PDA models. Furthermore, the presence of T cells predicted

the response to PD-1 immune checkpoint blockade in preclinical models of NSCLC and PDA.

[37, 50].

In T cell inflamed PDA tumor models (2838c3 and 7160c2), ADT-007 treatment

increased numbers of innate-like T lymphocytes (NK T, γδT, and DN T) and NK cells, whereas

conventional CD4[+] and CD8[+] T cell numbers did not change. In the CRC model, CT-26, trends

for increases in conventional CD4[+] and CD8[+] T cells were consistently measured, although

variable in statistical significance. In the 7160c2 TiME, ADT-007 treatment increased

proportions of CD44[hi] CD62L[-] effector memory T cells and production of T_H1 cytokines,

including IL-2, TNFα, IFNγ, and the cytotoxic protein granzyme B. Although increased

production of these cytokines was not observed in the CD4[+] and CD8[+] T cells in the 2838c3

model, increased proportions of effector T cells and higher levels of PD-1 expression were

observed when compared to CD4$^+$ and CD8$^+$ T cells within the 7160c2 TiME consistent with

previous findings [36]. Analysis of *in vivo* production of IFNγ using Thy1.1-IFNγ reporter mice

revealed that most adaptive and innate-like T cells were Thy1.1$^+$, consistent with IFNγ

production. These findings suggest role(s) for innate and adaptive immune response in

contributing to the antitumor activity of ADT-007.

In conclusion, we believe this novel class of pan-RAS inhibitors has strong potential to

address the complex genetic landscape of RAS-driven cancers. The results support further

development of ADT-007 analogs and prodrugs with oral bioavailability as a generalizable

monotherapy or combined with other RAS pathway-directed inhibitors or immunotherapy.

References

1. Siegel RL, Giaquinto AN, Jemal A. Cancer statistics, 2024. *CA Cancer J Clin* **2024**;74(1):12-49.
2. Yang Y, Zhang H, Huang S, Chu Q. KRAS Mutations in Solid Tumors: Characteristics, Current Therapeutic Strategy, and Potential Treatment Exploration. *J Clin Med* **2023**;12(2):709.
3. Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, *et al*: Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell* **2003**; 4(6):437-450.
4. Kojima K, Vickers SM, Adsay NV, Jhala NC, Kim HG, Schoeb TR, et al. Inactivation of Smad4 accelerates Kras(G12D)-mediated pancreatic neoplasia. *Cancer Res* **2007**;67(17):8121-8130.
5. Boutin AT, Liao WT, Wang M, Hwang SS, Karpinets TV, Cheung H, *et al*: Oncogenic Kras drives invasion and maintains metastases in colorectal cancer. *Genes Dev* **2017**;31(4):370-382.
6. Bardeesy N, Aguirre AJ, Chu GC, Cheng KH, Lopez LV, Hezel AF, *et al*. Both p16(Ink4a) and the p19(Arf)-p53 pathway constrain progression of pancreatic adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A* **2006**;103(15):5947-5952.
7. Sheffels E, Kortum RL. The Role of Wild-Type RAS in Oncogenic RAS Transformation. *Genes (Basel)* **2021**;12(5):662.
8. Ying H, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sananikone E, *et al*: Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* **2012**; 149(3):656-670.
9. Mukhopadhyay S, Goswami D, Adiseshaiah PP, Burgan W, Yi M, Guerin TM, et al. Undermining Glutaminolysis Bolsters Chemotherapy While NRF2 Promotes Chemoresistance in KRAS-Driven Pancreatic Cancers. *Cancer Res* **2020**;80(8):1630-1643.
10. Collins MA, Brisset JC, Zhang Y, Bednar F, Pierre J, Heist KA, et al. Metastatic pancreatic cancer is dependent on oncogenic Kras in mice. *PLoS One* **2012**;7(12):e49707.
11. Zhang Y, Yan W, Collins MA, Bednar F, Rakshit S, Zetter BR, et al. Interleukin-6 is required for pancreatic cancer progression by promoting MAPK signaling activation and oxidative stress resistance. *Cancer Res* **2013**;73(20):6359-6374.
12. Zdanov S, Mandapathil M, Abu Eid R, Adamson-Fadeyi S, Wilson W, Qian J, *et al*. Mutant KRAS Conversion of Conventional T Cells into Regulatory T Cells. *Cancer Immunol Res* **2016**;4(4):354-365.
13. Bayne LJ, Beatty GL, Jhala N, Clark CE, Rhim AD, Stanger BZ, et al. Tumor-derived granulocyte-macrophage colony-stimulating factor regulates myeloid inflammation and T cell immunity in pancreatic cancer. *Cancer Cell* **2012**;21(6):822-835.
14. Ischenko I, D'Amico S, Rao M, Li J, Hayman MJ, Powers S, et al. KRAS drives immune evasion in a genetic model of pancreatic cancer. *Nat Commun* **2021**;12(1):1482.
15. Coelho MA, de Carne Trecesson S, Rana S, Zecchin D, Moore C, Molina-Arcas M, *et al*. Oncogenic RAS Signaling Promotes Tumor Immunoresistance by Stabilizing PD-L1 mRNA. *Immunity* **2017**; 47(6):1083-1099 e1086.
16. Muthalagu N, Monteverde T, Raffo-Iraolagoitia X, Wiesheu R, Whyte D, Hedley A, *et al*. Repression of the Type I Interferon Pathway Underlies MYC- and KRAS-Dependent Evasion of NK and B Cells in Pancreatic Ductal Adenocarcinoma. *Cancer Discov* **2020**;10(6):872-887.

793   17.   Stalnecker CA, Der CJ. RAS, wanted dead or alive: Advances in targeting RAS mutant
794   cancers. *Sci Signal* **2020**;13(624).

795   18.   Kwan AK, Piazza GA, Keeton AB, Leite CA. The path to the clinic: a comprehensive
796   review on direct KRAS(G12C) inhibitors. *J Exp Clin Cancer Res* **2022**;41(1):27.

797   19.   Canon J, Rex K, Saiki AY, Mohr C, Cooke K, Bagal D, *et al*: The clinical KRAS(G12C)
798   inhibitor AMG 510 drives anti-tumour immunity. *Nature* **2019**;575(7781):217-223.

799   20.   Li BT, Falchook GS, Durm GA, Burns TF, Skoulidis F, Ramalingam SS, *et al*.
800   CodeBreaK 100/101: First Report of Safety/Efficacy of Sotorasib in Combination with
801   Pembrolizumab or Atezolizumab in Advanced KRAS p.G12C NSCLC. *J Thorac Oncol*
802   **2022**;17(9S):S10.

803   21.   Ryan MB, Fece de la Cruz F, Phat S, Myers DT, Wong E, Shahzade HA, *et al.* Vertical
804   Pathway Inhibition Overcomes Adaptive Feedback Resistance to KRAS(G12C)
805   Inhibition. *Clinical cancer research : an official journal of the American Association for*
806   *Cancer Research* **2020**;26(7):1633-1643.

807   22.   Awad MM, Liu S, Rybkin, II, Arbour KC, Dilly J, Zhu VW, *et al*: Acquired Resistance to
808   KRAS(G12C) Inhibition in Cancer. *N Engl J Med* **2021**;384(25):2382-2393.

809   23.   Thompson HJ, Jiang C, Lu J, Mehta RG, Piazza GA, Paranka NS, *et al.* Sulfone
810   metabolite of sulindac inhibits mammary carcinogenesis. *Cancer Res* **1997**;57(2):267-
811   271.

812   24.   Herrmann C, Block C, Geisen C, Haas K, Weber C, Winde G, *et al.* Sulindac sulfide
813   inhibits Ras signaling. *Oncogene* **1998**;17(14):1769-1776.

814   25.   Karaguni IM, Glüsenkamp KH, Langerak A, Geisen C, Ullrich V, Winde G, *et al.* New
815   indene-derivatives with anti-proliferative properties. *Bioorg Med Chem Lett* 2002,
816   12(4):709-713.

817   26.   Waters AM, Der CJ. KRAS: The Critical Driver and Therapeutic Target for Pancreatic
818   Cancer. *Cold Spring Harb Perspect Med* **2018**;8(9):a031435.

819   27.   Molina-Arcas M, Moore C, Rana S, van Maldegem F, Mugarza E, Romero-Clavijo P, *et*
820   *al*. Development of combination therapies to maximize the impact of KRAS-G12C
821   inhibitors in lung cancer. *Sci Transl Med* **2019**;11(510):eaaw7999.

822   28.   Tanaka N, Lin JJ, Li C, Ryan MB, Zhang J, Kiedrowski LA, *et al.* Clinical Acquired
823   Resistance to KRAS(G12C) Inhibition through a Novel KRAS Switch-II Pocket Mutation
824   and Polyclonal Alterations Converging on RAS-MAPK Reactivation. *Cancer Discov*
825   **2021**;11(8):1913-1922.

826   29.   Xiao D, Deguchi A, Gundersen GG, Oehlen B, Arnold L, Weinstein IB. The sulindac
827   derivatives OSI-461, OSIP486823, and OSIP487703 arrest colon cancer cells in mitosis
828   by causing microtubule depolymerization. *Mol Cancer Ther* **2006**;5(1):60-67.

829   30.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, *et al*. The
830   Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug
831   sensitivity. *Nature* **2012**; 483(7391):603-607.

832   31.   Lee KJ, Chang WL, Chen X, Valiyaveettil J, Ramirez-Alcantara V, Gavin E, *et al*.
833   Suppression of Colon Tumorigenesis in Mutant Apc Mice by a Novel PDE10 Inhibitor
834   that Reduces Oncogenic β-Catenin. *Cancer Prev Res (Phila)* **2021**;14(11):995-1008.

835   32.   Hunter JC, Gurbani D, Ficarro SB, Carrasco MA, Lim SM, Choi HG, *et al*. In situ
836   selectivity profiling and crystal structure of SML-8-73-1, an active site inhibitor of
837   oncogenic K-Ras G12C. *Proc Natl Acad Sci U S A* **2014**;111(24):8895-8900.

838   33.   Agola JO, Hong L, Surviladze Z, Ursu O, Waller A, Strouse JJ, *et al*. A competitive
839   nucleotide binding inhibitor: in vitro characterization of Rab7 GTPase inhibition. *ACS*
840   *chemical biology* **2012**; 7(6):1095-1108.

841   34.   Hunter JC, Manandhar A, Carrasco MA, Gurbani D, Gondi S, Westover KD. Biochemical
842   and Structural Analysis of Common Cancer-Associated KRAS Mutations. *Molecular*
843   *cancer research : MCR* **2015**;13(9):1325-1335.

35. John J, Rensland H, Schlichting I, Vetter I, Borasio GD, Goody RS, *et al.* Kinetic and
structural analysis of the Mg(2+)-binding site of the guanine nucleotide-binding protein
p21H-ras. *The Journal of biological chemistry* **1993**;268(2):923-929.
36. Li J, Byrne KT, Yan F, Yamazoe T, Chen Z, Baslan T, *et al.* Tumor Cell-Intrinsic Factors
Underlie Heterogeneity of Immune Cell Infiltration and Response to Immunotherapy.
Immunity **2018**; 49(1):178-193 e177.
37. Mugarza E, van Maldegem F, Boumelha J, Moore C, Rana S, Llorian Sopena M, *et al.*
Therapeutic KRAS(G12C) inhibition drives effective interferon-mediated antitumor
immunity in immunogenic lung cancers. *Sci Adv* **2022**;8(29):eabm8780.
38. Kemp SB, Cheng N, Markosyan N, Sor R, Kim IK, Hallin J, *et al.* Efficacy of a Small-
Molecule Inhibitor of KrasG12D in Immunocompetent Models of Pancreatic Cancer.
Cancer Discov **2023**; 13(2):298-311.
39. Harrington LE, Janowski KM, Oliver JR, Zajac AJ, Weaver CT. Memory CD4 T cells
emerge from effector T-cell progenitors. *Nature* **2008**;452(7185):356-360.
40. Khan I, Koide A, Zuberi M, Ketavarapu G, Denbaum E, Teng KW, *et al.* Identification of
the nucleotide-free state as a therapeutic vulnerability for inhibition of selected
oncogenic RAS mutants. *Cell Rep* **2022**;38(6):110322.
41. Esteban LM, Vicario-Abejon C, Fernandez-Salguero P, Fernandez-Medarde A,
Swaminathan N, Yienger K, *et al.* Targeted genomic disruption of H-ras and N-ras,
individually or in combination, reveals the dispensability of both loci for mouse growth
and development. *Mol Cell Biol* **2001**; 21(5):1444-1452.
42. Johnson L, Greenbaum D, Cichowski K, Mercer K, Murphy E, Schmitt E, *et al*: K-ras is
an essential gene in the mouse with partial functional overlap with N-ras. *Genes Dev*
1997;11(19):2468-2481.
43. Allain EP, Rouleau M, Lévesque E, Guillemette C. Emerging roles for UDP-
glucuronosyltransferases in drug resistance and cancer progression. *Br J Cancer* **2020**;
122(9):1277-1287.
44. Tinsley HN, Gary BD, Thaiparambil J, Li N, Lu W, Li Y,*et al.* Colon tumor cell growth-
inhibitory activity of sulindac sulfide and other nonsteroidal anti-inflammatory drugs is
associated with phosphodiesterase 5 inhibition. *Cancer Prev Res (Phila)*
2010;3(10):1303-1313.
45. Lu Y, Chen J, Xiao M, Li W, Miller DD. An overview of tubulin inhibitors that interact with
the colchicine binding site. *Pharm Res* **2012**;29(11):2943-2971.
46. Waters AM, Khatib TO, Papke B, Goodwin CM, Hobbs GA, Diehl JN, *et al*: Targeting
p130Cas- and microtubule-dependent MYC regulation sensitizes pancreatic cancer to
ERK MAPK inhibition. *Cell Rep* **2021**;35(13):109291.
47. Coley AB, Ward A, Keeton AB, Chen X, Maxuitenko Y, Prakash A, *et al.* Pan-RAS
inhibitors: Hitting multiple RAS isozymes with one stone. *Adv Cancer Res* **2022**;153:131-
168.
48. Luchini C, Paolino G, Mattiolo P, Piredda ML, Cavaliere A, Gaule M, *et al.* KRAS wild-
type pancreatic ductal adenocarcinoma: molecular pathology and therapeutic
opportunities. *J Exp Clin Cancer Res* **2020**;39(1):227.
49. di Magliano MP, Logsdon CD. Roles for KRAS in pancreatic tumor development and
progression. *Gastroenterology* **2013**;144(6):1220-1229.
50. Mahadevan KK, McAndrews KM, LeBleu VS, Yang S, Lyu H, Li B, *et al.* KRAS(G12D)
inhibition reprograms the microenvironment of early and advanced pancreatic cancer to
promote FAS-mediated killing by CD8(+) T cells. *Cancer Cell* **2023**;41(9):1606-1620
e1608.
51. Flatmark K, Reed W, Halvorsen T, Sorensen O, Wiig JN, Larsen SG, *et al.*
Pseudomyxoma peritonei--two novel orthotopic mouse models portray the PMCA-I
histopathologic subtype. *BMC Cancer* **2007**;7:116.

52.   Flatmark K, Davidson B, Kristian A, Stavnes HT, Førsund M, Reed W. Exploring the
896   peritoneal surface malignancy phenotype—a pilot immunohistochemical study of human
897   pseudomyxoma peritonei and derived animal models. *Human Pathology*
898   **2010**;41(8):1109-1119.
899   53.   Fleten KG, Lund-Andersen C, Waagene S, Abrahamsen TW, Mørch Y, Boye K, *et al.*
900   Experimental Treatment of Mucinous Peritoneal Metastases Using Patient-Derived Xenograft
901   Models. *Translational oncology* **2020**;13(8):100793.

    Figures and Figure Legends:

Figure 1. ADT-007 inhibits activation of mutant and RAS^WT isozymes and downstream

MAPK signaling circumventing resistance. (A) Chemical structure of ADT-007. **(B)** ADT-007

inhibition of KRAS^{G13D} HCT-116 vs. KRAS^{WT}, BRAF^{V600E} HT-29 growth following 72 hours of

treatment (CellTiter-Glo ATP content Assay; (CTG))**.** Quadruplicate samples ± standard error of

the mean (SEM). Inset: Activated RAS-GTP-RBD pulldown evaluated by pan-RAS western

(WB). Total RAS and GAPDH were evaluated by WB. **(C)** Ectopic expression of HRAS^{G12V}

sensitized HT-29 cells to ADT-007 growth inhibition (quadruplicates ± SEM). Inset: Activated

RAS in untreated parental, vector control (retro), and transfected (HRAS^{G12V}). (**D**) HCT-116

KRAS^{G13D} cells were treated with increasing concentrations of ADT-007 overnight in complete

medium (10% FBS). Activated RAS-GTP pulldowns, phospho(P)-cRAF, P-MEK, P-ERK, total

RAS and GAPDH as loading controls were detected by WB. Data represent 3 independent

experiments. (**E,F**) Serum-starved MIA PaCa-2 KRAS^{G12C} PDA cells were treated overnight with

vehicle or increasing concentrations of ADT-007, then EGF stimulated (10 min). (**E**) P-cRAF, P-

MEK, P-ERK, P-AKT and GAPDH as loading controls were detected by WB. Data represent 2

independent experiments. (**F**) Activated RAS-GTP was isolated by RAS-RBD pulldown followed

by WB with pan-RAS, KRAS, HRAS, or NRAS specific antibodies and GAPDH control. Data

represent at least two independent experiments. Growth inhibition of LLC cells (**G, H**; KRAS^{G12C}

/ NRAS^{Q61H}) or MIA PaCa-2 cells (**I**; KRAS^{G12C}) by ADT-007 vs BI-2865 or AMG-510.

Type	Tissue Origin	Cell Line	Mutation Status	IC$_{50}$ (nM)	UGT1A10
Mutant RAS, sensitive	Pancreatic	MIA PaCa-2	KRAS G12C	2.1	0.263
	Pancreatic	PANC-1	KRAS G12D	2.4	0.043
	Pancreatic	SW-1990	KRAS G12D	6.3	0.138
	Lung	A549	KRAS G12S	6.8	0.070
	Pancreatic	CFPAC-1	KRAS G12V	5.8	2.63
	Colon	SW480	KRAS G12V	6.5	0.58
	Breast	MDA-MB-231	KRAS G13D	3.6	0.043
	Colon	HCT-116	KRAS G13D	4.7	0
	Ovarian	OVCAR-8	KRAS P121H	5.8	0.057
	Breast	Hs 578T	HRAS G12D	3.9	0.098
	Skin	SK-MEL-2	NRAS Q61K	2.1	0
	Lung	NCI-H1299	NRAS Q61K	2.4	0
WT RAS, resistant	Normal Lung	NHAEC	WT RAS	19100	*
	Normal Colon	NCM460	WT RAS	50000	*
Downstream mutation (WT RAS) resistant	Ovarian	OV-90	WT RAS/mut BRAF	350	4.57
	Colon	HT-29	WT RAS/mut BRAF	512	6.97
	Skin	SK-MEL-28	WT RAS/mut BRAF	> 25,000	0.14 (3.53†)
	Pancreatic	BxPC-3	WT RAS/mut BRAF	2500	7.75
Upstream RAS activation sensitive	Lung	NCI-H1975	WT RAS/mut EGFR	3.9	*
	Skin (mouse)	B16	WT RAS/mut PDGFR	5.8	*
	Brain	U-87 MG	WT RAS/mut NF1	1.6	0
	Ovarian	SK-OV-3	WT RAS/mut NF1	6.7	0.202

Table 1. ADT-007 sensitivity is associated with high activated RAS and low UGT superfamily enzyme levels. Growth inhibitory (IC$_{50}$) values following 72 hours of treatment (CTG assay). TPM normalized expression of the most significant differentially expressed UGT1A10 enzyme from published data (Cancer Cell Line Encyclopedia; DepMap Portal; Broad Institute). * expression data not available, † TPM normalized UGT2B7 expression in SK-MEL-28.

Figure 2. Phenolic moiety is critical for RAS selectivity of ADT-007-. (A) Left panel: ADT-007 selectively inhibits growth of mutant KRAS[G12C] MIA PaCa-2 PDA cells vs. KRAS[WT] BxPC-3 (72h, CTG Assay; triplicates ± SEM).   **(B)** Left panel: ADT-106 lacking a phenolic OH moiety was less potent (72h, CTG Assay; triplicates ± SEM). Right panels (**A, B**): Cells were treated for 18 hours, followed by activated RAS RAS-RBD pulldowns and WB analysis. **(C)** UGT1 protein expression in ADT-007 sensitive versus resistant cells by WB. (**D**) UGT1 protein expression and

946   control GAPDH (WB) were quantified and normalized (**E**) in image J. Representative of two

947   independent experiments. **(E, F)** Mean ADT-007 and ADT-007-glucuronide were quantified in

948   triplicate cell pellets by LC-MS/MS ± SEM), * = $p < 0.05$.

Figure 3. ADT-007 binds KRAS, inhibits nucleotide loading and effector binding. Cellular

target engagement assays show: **(A)** ADT-007 binds KRAS with high affinity in KRAS[G13D] HCT-

116 but not KRAS[WT] HT-29 CRC cells. Graphs show levels of compound stabilized KRAS

detected by WB compared with vehicle control. (**B**) Binding affinity of ADT-007 for mutant

KRAS^{G12C} was assessed by 45 min treatment of HEK293 cells expressing KRAS^{G12C}-Micro-Tag

construct compared with a nonselective negative control compound. Curves are graphed as the

average of two replicates ± SEM. **(C-D)** The inhibitory activity of ADT-007 to block recombinant

KRAS^{WT} activation (effector binding) in a cell-free system was evaluated by RAS-RBD pulldown

assays. (**C**) ADT-007 inhibited KRAS^{WT}-RAF activation before GTP addition. **(D)** ADT-007 did

not significantly inhibit KRAS^{WT} activation / pulldown when added after GTP loading. (**E-F**)

Kinetics of GTP binding to KRAS were measured using a fluorescent GTP analog (Mant-GTP).

ADT-007 reduced the rate of Mant-GTP binding to KRAS if pretreated with EDTA to chelate

Mg2+ (**E**), but not if added after non-hydrolysable GTPγS prior to incubation with ADT-007 (**F**).

Data are representative of at least two independent experiments.

970   **Figure 4. NMR analysis of ADT-007 binding to KRAS.** Addition of ADT-007 (red) to Mg^{2+}-free

971   KRAS (blue) (5:1 ratio) resulted in statistically significant NMR signal attenuation at G13 and

972   K16 and other residues (described in methods). Assignments for G13 and K16 due to their

973   unique resonance frequencies are shown as a larger image to the left.

Figure 5. ADT-007 induces growth arrest and regression of CRC, GBA, and PDA cells *in*

vivo. (**A**) Left panel: Growth inhibition of subcutaneous patient derived xenograft (PDX) gall

bladder adenocarcinoma (GBA) bearing KRAS^G12V by peritumoral administration (5 mg/kg, BID,

13 days, n = 5 mice/group, mean ± SEM). Right panel: Percent change in tumor volume at

endpoint compared to volume at start of treatment. (**B-D**) Left panels: Growth inhibition of

peritoneal metastasis in mucinous colorectal adenocarcinoma models (**B**) PMCA1, (**C**) PMP1,

and (**D**) PC244 in nude mice by IP administration of ADT-007 (2.5 mg/kg, BID) for treatment

periods of 18 (PMCA1), 23 (PMP1), and 18 days (PC244). Treatment was initiated 24 hours

after implantation (n = 5-6 mice/group, mean ± SEM) and mice were followed by MRI for a

period of 24 to 50 days (6 – 31 days post treatment cessation) after implantation. Right panels:

tumor sizes at end of treatment. (**E**) Left panel: Growth inhibition of subcutaneous CT26 colon

tumors in BALB/c mice by IT administration of ADT-007 (10 mg/kg, QD, 13 days, mean ± SE,

n=12). Right panel: Percent change in tumor volume at endpoint compared to volume at start of

treatment. (**F-H**) Left panels: Growth inhibition studies of subcutaneous (**F**) 2838c3, (**G**) 7160c2,

(**H**) 5363 PDA tumors in C57BL/6 mice by peritumoral administration of ADT-007 (5 mg/kg, BID)

for treatment periods of 17 (2838c3), 21 (7160c2), and 24 (5363) days (n = 7 - 10 mice/group,

mean ± SEM). Right Panels: Percent change in tumor volume at endpoint compared to volume

at start of treatment. All animal studies are representative of 2-3 independent experiments. Left

panels: Error bars are representative of SEM. Right panels: error bars are representative of SD.

Left panels: Statistical significance was assessed using 2-way ANOVA with Sadik's multiple

comparisons test. Right panels: Statistical significance was assessed with Welch's t-test. **

$p<0.001$, **** $p<0.0001$,

7160c2, SQ, 20 days of treatment

2838c3, SQ, 17 days of treatment

Figure 6. ADT-007 modulates myeloid subsets, induces T cell activation, and enhances

cytotoxic function in PDA mouse models. (**A-H**) Data is from the 7160c2 PDA model. At 21

days of post treatment the proportions of tumor and myeloid cells were evaluated by

multiparameter flow cytometry. (**A**) The percentages of CD45$^-$ YFP$^+$ tumor cells were decreased

in ADT-007 treated mice (n = 8) compared to sham (n = 8). (**B**) The numbers/mg tumor of

CD11b$^{+/-}$ CD11c$^{+/-}$ myeloid, CD11b$^+$ Ly6G$^+$ Ly6C$^+$ (gMDSC), and CD11b$^+$ Ly6G$^-$ Ly6C^{hi}

(mMDSC) in ADT-007 treated (n = 8) compared to sham (n = 8) treated mice. (**C**) The

numbers/mg tumor of CD11b$^-$ CD11c^{hi} MHCII^{hi} XCR1$^+$ cDC1 and CD11b$^+$ CD11c^{hi} MHCII^{hi}

CD172α$^+$ cDC2 subsets in ADT-007 (n = 8) compared to sham (n = 8) treated mice. (**D**) Tumor

growth in 7160c2 PDA tumor bearing C57BL/6 WT vs. C57BL/6 Rag 1$^{-/-}$ mice by peritumoral

administration (5 mg/kg, BID, days 7-28, n = 6-22 mice/group, mean $\pm$ SE). Data is an

aggregate of 2 separate experiments. (**E**) The numbers/mg of tumor of CD4$^+$ Foxp3$^-$, CD4$^+$

Foxp3$^+$, CD8$^+$ T cells, CD3$^+$ γδTCR$^+$(γδT), CD3$^+$ NK1.1$^+$(NKT) and CD3$^+$ CD4/8$^-$ γδTCR$^-$ (DN) T

cells and (**F**) CD3$^-$ NK1.1$^+$ NK cells in ADT-007 (n = 8) and sham (n = 8) treatment. (**G-H**)

Intracellular cytokine (IL-2, granzyme B, and IFNγ) production (ICS) from (**G**) CD4$^+$ and (**H**)

CD8$^+$ T cells isolated from the 7160c2 TME were evaluated after 5-hour stimulation with

PMA/ionomycin in the presence of brefeldin A in ADT-007 (n = 8) or sham (n = 8) treated mice.

(**I-P**) Data is from the 2838c3 PDA model. At 17 days post treatment the proportions of tumor

and myeloid cells were evaluated by multiparameter flow cytometry. (**I**) The percentages of

CD45$^-$ YFP$^+$ tumor cells were decreased in ADT-007 treated mice (n = 8) compared to sham (n

= 8). (**J**) The numbers/mg tumor of myeloid, gMDSC, and mMDSC in ADT-007 treated (n = 8)

and sham (n = 8) treated mice. (**K**) The numbers/mg tumor of cDC1 and cDC2 cells were

evaluated in ADT-007 (n = 8) and sham (n = 8) treated mice. (**L**) Tumor growth in 2838c3 PDA

tumor bearing C57BL/6 WT vs. C57BL/6 Rag 1$^{-/-}$ mice by peritumoral administration (5 mg/kg,

BID, days 7-24, n = 6-22 mice/group, mean $\pm$ SE). Data is an aggregate of 2 separate

experiments. (**M**) Numbers/mg tumor of CD4$^+$ Foxp3$^-$, CD4$^+$ Foxp3$^+$, CD8$^+$, γδ, and DN T and

(**N**) NK cells in ADT-007 (n = 8) and sham (n = 8) treatment. (**O-P**) ICS assay results as

described above on (**O**) CD4$^+$ and (**P**) CD8$^+$ T cells isolated from the 2838c3 TiME in ADT-007

(n = 8) or sham (n = 8) treated mice. For all column graphs, data is representative of 2

independent experiments. Statistic significance was calculated by Welch's t-test, ns indicates

not significant. Error bars indicate SD. For evaluation of statistical significance in tumor growth

curves 2way ANOVA with Tukey's multiple comparison test was used to calculate statistical

significance. Error bars indicate SEM.

****$p < 0.0001$

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044