The small acid-soluble proteins of *Clostridioides difficile* regulate sporulation in a
SpoIVB2-dependent manner

Hailee N. Nerber, Marko Baloh, Joshua N. Brehm, and Joseph A. Sorg[*]

Department of Biology, Texas A&M University, College Station, TX 77845

[*]corresponding author

PH: 979-845-6299

Email: jsorg@bio.tamu.edu

Abstract

Clostridioides difficile is a pathogen whose transmission relies on the formation of dormant

endospores. Spores are highly resilient forms of bacteria that resist environmental and chemical

insults. In recent work, we found that *C. difficile* SspA and SspB, two small acid-soluble proteins

(SASPs), protect spores from UV damage and, interestingly, are necessary for the formation of

mature spores. Here, we build upon this finding and show that *C. difficile sspA* and *sspB* are

required for the formation of the spore cortex layer. Moreover, using an EMS mutagenesis

selection strategy, we identified mutations that suppressed the defect in sporulation of *C. difficile*

SASP mutants. Many of these strains contained mutations in *CDR20291_0714* (*spoIVB2*)

revealing a connection between the SpoIVB2 protease and the SASPs in the sporulation

pathway. This work builds upon the hypothesis that the small acid-soluble proteins can regulate

gene expression.

Importance

C. difficile is easily spread through the production of highly resistant spores. Understanding how

spores are formed could yield valuable insight into how the sporulation process can be halted to

render spores that are sensitive to cleaning methods. Here, we identify another protein involved

in the sporulation process that is seemingly controlled by the small acid-soluble proteins

(SASPs). This discovery allows us to better understand how the *C. difficile* SASPs may bind to

specific sites on the genome to regulate gene expression.

Introduction

Clostridioides difficile is a Gram-positive pathogen that causes approximately 220,000 cases of infection and nearly 13,000 deaths annually [1]. *C. difficile* vegetative cells produce toxins that disrupt the colonic epithelium, resulting in diarrhea and colonic inflammation [2, 3]. These toxin-producing vegetative cells are strictly anaerobic and cannot survive outside of a host for extended periods [4]. However, *C. difficile* produces endospores that are shed into the environment, can withstand oxygen and other environmental insults, and serve as the transmissive form of the organism [5-7].

Endospores are highly structured forms of bacteria. Residing in the spore core are the DNA, RNA, ribosomes, calcium dipicolinic acid (Ca-DPA), small acid-soluble proteins (SASPs), and other proteins that are necessary for the spore to outgrow into a vegetative cell [8-13]. Surrounding the core is a phospholipid membrane, cell wall, and a specialized cortex peptidoglycan layer. In the cortex, many of the N-acetylmuramic acid residues are converted into muramic-δ-lactam residues, which are recognized by the spore cortex lytic enzymes during germination [8, 14-16]. Outside of the cortex is a phospholipid outer membrane, a proteinaceous spore coat, and an exosporium [8, 17-21].

Generally, endospores are formed in response to nutrient deprivation. Upon initiation of sporulation, the vegetative cell asymmetrically divides into the larger mother cell and the smaller forespore compartments [22, 23]. The forespore becomes engulfed by the mother cell so that it can be matured into the dormant endospore. Once the endospore is fully formed, the mother cell lyses and releases the spore into the environment [24].

Like all known endospore-forming bacteria, the *C. difficile* sporulation program initiates upon phosphorylation of the sporulation master transcriptional activator, Spo0A [5, 25, 26]. After

asymmetric division, each compartment begins a cascade of sigma factor activation [27, 28]. In

the mother cell compartment, σ^E becomes activated and leads to σ^K expression. In the

forespore compartment, σ^F is activated and leads to σ^G activation [8, 23]. Loss of σ^F results in a

strain that does not complete engulfment or form the cortex layer [28]. The loss of σ^G results in a

strain that forms a localized coat layer but does not fully complete engulfment (i.e. no

membrane fission) or form the cortex layer [27]. Loss of σ^E results in a strain that is blocked at

asymmetric septation. Loss of σ^K results in a strain that fully engulfs the forespore and forms a

correctly localized cortex layer, but no visible coat layer [28]. Thus, cortex assembly occurs

through σ^G regulated genes and coat production is dependent on σ^K genes.

The small acid-soluble proteins (SASPs) are very abundant in spores and have high

sequence similarity across spore-forming species [29]. In many organisms, including *Bacillus*

subtilis and *Clostridium perfringens,* the SASPs protect DNA against UV damage and damage

from genotoxic chemicals [30-33]. In *B. subtilis*, the SASPs are considered non-specific DNA

binding proteins that coat the DNA and change the conformation to a more rigid, intermediate, B

to A form [29, 34-37]. This conformation leads to difficulty in forming UV-induced thymidine-

dimers and, instead, promotes the formation of spore photoproducts; a repair mechanism is

present in the spore to correct these lesions [38-40]. In *in vitro* transcription assays, addition of

SASPs to DNA reduced transcription of some, but not all, genes, further illustrating their ability

to bind DNA [37]. Moreover, the absence of transcription in mutant strains whose spores cannot

degrade SASPs, suggest that SASPs could regulate gene expression [37, 41].

In prior work, we found that the *C. difficile* SASPs are important for spore UV resistance

but do not strongly contribute to chemical resistances [42]. Surprisingly, a *C. difficile* Δ*sspA*

Δ*sspB* double mutant strain could not complete spore formation, a phenotype not observed in

other endospore-forming bacteria. This led us to hypothesize that the *C. difficile* SASPs are

involved, somehow, in regulating sporulation. We hypothesize that SASPs have regions of high

affinity on DNA where they bind to influence the transcription of genes. As the concentration of

SASPs increases, they nonspecifically coat the DNA to provide the protection normally

associated with SASPs. In the *C. difficile* Δ*sspA* Δ*sspB* strain, we hypothesize that sporulation

is reduced due to altered gene expression of important sporulation genes.

Using a strategy that selected for the generation of mature spores from the sporulation

deficient *C. difficile* Δ*sspA* Δ*sspB* strain, we identified mutations in *spoIVB2* that suppressed the

mutant sporulation phenotype. SpoIVB2 is a protease that is recently characterized in *C. difficile*

and the *C. difficile* Δ*spoIVB2* mutant strain has a phenotype similar to the *C. difficile* Δ*sspA*

Δ*sspB* strain. Based upon the data in this manuscript, we hypothesize that the σ^G-dependent

expression of the *C. difficile* SASPs activates the σ^F-dependent expression of *spoIVB2*, and that

low levels of SpoIVB2 in a *C. difficile* Δ*sspA* Δ*sspB* mutant halts sporulation by an unknown

mechanism.

Results

C. difficile sspA and sspB regulate sporulation in the C. difficile CD630Δerm strain

In prior work, we discovered that *C. difficile* SspA and SspB were, individually, important

for UV resistance [42]. Surprisingly, we found that the combinatorial deletion of the *sspA* and

sspB genes, or a deletion in *sspB* and an *sspA*$_{G52V}$ missense mutation (referred to as *C. difficile*

Δ*sspB** hereafter), in the *C. difficile* R20291 strain resulted in the drastic reduction of mature

spore formation and, instead, resulted in phase gray spores [42]. To confirm that this phenotype

was strain independent, we generated the single and double mutants of *sspA* and *sspB* in the

C. difficile CD630Δ*erm* strain. Unsurprisingly, the CD630Δ*erm* Δ*sspA* Δ*sspB* double mutant

also produced phase gray spores that were trapped within mother cells (Figure 1A). Though the

single mutants did not affect spore yield, the double mutant had a 5-$\log_{10}$ decrease in spore

formation. This defect could be restored to near wildtype levels by expression of *sspA* and *sspB*,

in trans, from a plasmid (Figure 1B). When UV resistance was assessed, the *C. difficile*

CD630Δ*erm* Δ*sspA* and the Δ*sspB* single mutant strains both had an approximate 1-$\log_{10}$ loss

in viability after 10 minutes of UV exposure (Figure 1C). Though consistent with the findings we

observed for the *C. difficile* R20291 Δ*sspB* mutant strain, the impact on viability for the *sspA*

mutant was less in the *C. difficile* CD630Δ*erm* background than in *C. difficile* R20291

background. SspA and SspB appear to regulate sporulation in both the *C. difficile* R20291 and

CD630Δ*erm* strains, hence, this is likely a conserved function in *C. difficile*.

B. subtilis sspA complements UV and sporulation phenotypes of C. difficile R20291 mutants.

Due to the high sequence similarity of SASPs, and their ability to cross-complement in

other organisms, we assessed whether *sspA* from *B. subtilis* would complement the phenotypes

observed in the *C. difficile* SASP mutants [29, 43]. Sporulation was complemented to varying

degrees by the expression of *B. subtilis sspA* from the *C. difficile sspA* promoter in *C. difficile*

Δ*sspB** and *C. difficile* Δ*sspA* Δ*sspB*. In *C. difficile* Δ*sspB**, expression of *B. subtilis sspA*

increased spore yield by approximately 10-fold. In the *C. difficile* Δ*sspA* Δ*sspB* strain,

expression of *B. subtilis sspA* increased spore yield by approximately 100-fold (Figure 2A).

Spores derived from a *C. difficile* Δ*sspA* mutant strain with a plasmid expressing *B.*

subtilis sspA, under the *C. difficile sspA* native promoter, were exposed to UV light for 10

minutes and their viability assessed. *B. subtilis sspA* could partially restore UV resistance to the

C. difficile Δ*sspA* mutant strain, although not to wild type levels (Figure 2B). These data show

that *B. subtilis* and *C. difficile* SspA could function in similar ways due to the ability of *B. subtilis*

sspA to complement phenotypes found in *C. difficile* SASP mutants.

Visualizing the impact of SASP mutations on C. difficile spores

To visualize the impact of the *C. difficile ΔsspA ΔsspB* deletions on spore structure, we

used transmission electron microscopy (TEM). Strains generated in the *C. difficile* R20291

background were cultured for 6 days and then prepared for TEM. As expected, the *C. difficile*

R20291 wild type strain generated fully formed and mature spores. The *C. difficile ΔsspA* and

C. difficile ΔsspB single mutant strains also formed spores with the expected spore structures

(*e.g.*, cortex and coat layers). However, *C. difficile ΔsspB** and *C. difficile ΔsspA ΔsspB* strains

generated spores that did not form cortex layers, and had a visible, but anomalous, coat layer

(Figure 3). Expression of the SASPs *in trans* under their native promoter regions complemented

the mutant phenotypes by restoring the cortex layer and overall spore morphology.

Isolating suppressor mutations of the SASP mutant phenotypes

To gain insight into how the SASPs are involved in spore formation, we used

ethylmethane sulphonate (EMS) to introduce random mutations into the *C. difficile* genome, as

we have previously done (Supplement Figure 1) [44, 45]. The *C. difficile ΔsspB** or the *C.*

difficile ΔsspA ΔsspB strains were treated with EMS, washed, and then incubated for 5 days to

generate potential spores. Subsequently, the samples were heat-treated to kill vegetative cells

and immature spores. After removing cellular debris, the cultures were plated on a medium

supplemented with germinant [46]. Afterwards, we isolated strains and confirmed that they

generated spores. After confirmation, gDNA was extracted and sequenced to reveal the location

of mutations. From independent EMS mutageneses, we identified 4 suppressor strains

generated from the *C. difficile ΔsspB** strain and 11 from the *C. difficile ΔsspA ΔsspB* strain.

Unsurprisingly, there were many mutations in each strain, but mutations that potentially

contributed to suppression of the phenotype are listed in Table 1 (a full list of mutations can be

found in Table S3). As expected, due to the strong selection for spore dormancy, 2 out of 4 of

the isolates from *C. difficile* Δ*sspB** had a reversion mutation in *sspA*. We identified mutations in

different RNA polymerase subunits in 6 of 15 strains. These mutations could potentially affect

transcription rates of various genes. Mutations within the *sigG* and *spoVT* genes were also

present in some strains. *sigG* and *spoVT* mutants have a similar phenotype to the *C. difficile*

Δ*sspA* Δ*sspB* strain [27, 28, 47, 48]. Interestingly, 7 out of 15 isolates (from separate

mutagenesis experiments) contained mutations in *CDR20291_0714*. Among these strains, we

observed one strain with an A20T missense mutation and six with a synonymous mutation

(F37F). The *C. difficile* CD630Δ*erm* genome encodes a gene homologous to *CDR20291_0714*

and is annotated as *spoIVB2*. SpoIVB2 is a paralog of the SpoIVB protease, and we refer to

CDR20291_0714 as SpoIVB2 from here on.

We first tested if *in trans* expression of the identified *spoIVB2* alleles could restore

sporulation to the SASP mutant by generating merodiploid strains. When wild type *spoIVB2* was

expressed in *C. difficile* R20291 or *C. difficile* Δ*sspB** the spore yield did not change from their

respective phenotypes while the spore yield in the *C. difficile* Δ*sspA* Δ*sspB* strain increased by

1-$\log_{10}$ (Figure 4A). We also tested if catalytic activity impacted restoration. The catalytic site

was identified by aligning *C. difficile* SpoIVB / SpoIVB2 to *B. subtilis* SpoIVB. The three catalytic

residues found in *B. subtilis* are conserved in both SpoIVB and SpoIVB2 of *C. difficile* and we

have used *spoIVB2*$_{S301A}$ as a catalytically dead mutant [49]. In the wildtype *C. difficile* R20291

strain, the spore yield was not impacted when the *spoIVB2*$_{A20T}$ or *spoIVB2*$_{F37F}$ alleles were

combined with S301A (Figure 4B).

When the *spoIVB2* alleles were introduced into the *C. difficile* Δ*sspB** strain, the

spoIVB2$_{S301A}$ allele did not restore sporulation, but the *spoIVB2*$_{A20T}$ and *spoIVB2*$_{F37F}$ alleles

increased sporulation by approximately 2 and 3-$\log_{10}$, respectively. When these alleles were

combined with the *spoIVB2*$_{S301A}$ allele, sporulation was not restored (Figure 4C). These results

were similar to when the *spoIVB2* alleles were expressed in the *C. difficile* Δ*sspA* Δ*sspB* strain

[the expression of *spoIVB2*$_{A20T}$ and *spoIVB2*$_{F37F}$ resulted in an approximate 2-log$_{10}$ increase in

spore yield] (Figure 4D). The catalytically dead allele in combination with the identified alleles

from EMS was again unable to restore sporulation. These results suggest that the catalytic

activity of *C. difficile* SpoIVB2 is important for its function.

From these data, we hypothesized that SspA and SspB are activating the expression of

spoIVB2. We hypothesize that in the *C. difficile* Δ*sspA* Δ*sspB* and the *C. difficile* Δ*sspB**

mutants the levels of SpoIVB2 are reduced, which leads to an unprocessed target that is

essential for sporulation. The expression of wild type *spoIVB2* (Figure 4A) does not greatly

restore the sporulation deficient phenotype, likely because it is not expressed during σ^G gene

activation. The suppressor strains potentially increase the amount of SpoIVB2 present,

bypassing the need for σ^G expression. To further evaluate, we expressed wild type *spoIVB2*

(from a plasmid) in the suppressor strains that have *spoIVB2*$_{A20T}$ or *spoIVB2*$_{F37F}$ and quantified

spore formation. We found no significant difference in spore yield between the suppressor

strains with an empty vector and those expressing wild type *spoIVB2* from its native promoter

(Supplementary Figure 2A). Furthermore, we generated clean strains containing *spoIVB2*$_{A20T}$ or

spoIVB2$_{F37F}$ in the wild type or the *C. difficile* Δ*sspA* Δ*sspB* strains to eliminate from analysis the

outside mutations from EMS treatment. The *spoIVB2* alleles in the wild type background, with

an empty vector or a vector expressing wild type *spoIVB2*, did not impact the spore yield.

However, the strains containing the identified *spoIVB2* alleles, with an empty vector or a vector

expressing wild type *spoIVB2*, in the *C. difficile* Δ*sspA* Δ*sspB* strain increase spore yield 3-log$_{10}$,

compared to *C. difficile* Δ*sspA* Δ*sspB* alone. Again, the addition of wild type *spoIVB2* did not

further rescue the spore yield, indicating that the suppression is due to the altered *spoIVB2*

alleles (Supplementary Figure 2B).

The C. difficile spoIVB2 mutant is phenotypically similar to the C. difficile ΔsspA ΔsspB strain

To further evaluate the role of SpoIVB2 during sporulation, we generated a deletion of

spoIVB2 in the *C. difficile* R20291 strain. The *C. difficile* Δ*spoIVB2* strain generated phase gray

spores, similar to our observations for the *C. difficile* Δ*sspA* Δ*sspB* strain (Figure 5A). This

phenotype could be complemented by expression of *spoIVB2*$_{WT}$, *spoIVB2*$_{A20T}$, or *spoIVB2*$_{F37F}$

alleles from a plasmid. However, restoration did not occur when the catalytically dead

spoIVB2$_{S301A}$ was expressed (Figure 5A). The spore yield of the *C. difficile* Δ*spoIVB2* strain was

6-log$_{10}$ lower than wild type. The *C. difficile* Δ*spoIVB2* mutant supplemented with a plasmid

expressing *spoIVB2* wild type, A20T or F37F alleles restored the spore yield to wild type levels

(Figure 5B). However, when the S301A allele was present or in combination with the A20T or

F37F alleles, sporulation was not restored, again highlighting the importance of catalytic activity

in the function of SpoIVB2 (Figure 5B).

Analysis of the *C. difficile* Δ*spoIVB2* strain by TEM revealed many problems with the

sporulating cells (Figure 6). As seen in the field of view image, it was difficult to locate whole

cells for imaging. When a sporulating cell was found, there were structural issues within the

forespore. The cortex was missing and, with the lack of its constraint around the core, allowed

for expansion of the core contents. Though the coat was present, it appears anomalous. When

spoIVB2$_{WT}$, *spoIVB2*$_{A20T}$, or *spoIVB2*$_{F37F}$ were expressed from a plasmid, the structural

appearance of the spore was restored to wild type. However, when *spoIVB2*$_{S301A}$ was

expressed, it remained difficult to locate any sporulating cells.

Testing the impact of the suppressor alleles on spoIVB2 expression

To understand how the SASPs influence *spoIVB2* and / or other gene transcripts, RNA

was extracted from *C. difficile* wild type, *sspA* and *sspB* single and double mutants, the *sspB**

and *spoIVB2* mutant strains, as well as two representative suppressor strains from EMS

mutagenesis at 11 hours post plating on sporulation medium and RT-qPCR was performed.

Overall, at this time point, there were few differences in transcript levels. The *spoIVB2*$_{A20T}$

isolate (HNN19) was variable between extractions despite testing more biological replicates,

potentially due to other mutations from the EMS treatment. Though *sspA* or *sspB* transcripts

levels were largely unchanged, there was a slight increase in transcript levels in comparison to

wild type for *spoIVA* which encodes a protein involved in spore coat localization (Supplementary

Figure 3A-C) [50]. Transcripts for *sleC, pdaA,* and *spoVT* remained similar to wild type levels

(Supplementary Figure 3D-F). *spoIVB* transcript levels did not have a concise trend while, for

spoIVB2, the general trend was towards slightly reduced transcripts in the mutant strains with a

larger fold change in the EMS identified alleles (Supplementary Figure 4A-B). *spoIIP* transcripts

were slightly elevated in the mutant strains, except for the EMS isolates (Supplementary Figure

4C). For the DPA synthesis and packaging protein transcripts (*dpaA*, *spoVAC*, *spoVAD*, and

spoVAE), there were minimal differences for the mutant strains besides a slight increase in

spoVAC (Supplementary Figure 5A-D).

Manipulation of the F37 and F36 codons impact suppression

Next, we manipulated the F37 codon to see if other changes would allow for sporulation

to be restored in the mutant strains. We also changed the F36 codon from UUU to UUC

(generating an F36F silent mutation and the opposite codon change that occurred in the F37F

allele). These constructs were expressed from a plasmid under the *spoIVB2* native promoter

region, and the spore yield was assessed. When wild type *spoIVB2* (UUC codon) was

expressed in the *C. difficile* Δ*sspB** strain, sporulation was not restored to wild type levels

(Figure 7A). However, sporulation was partially restored with the *spoIVB2*$_{F37F}$ (UUU codon), the

spoIVB2$_{F37L}$ (UUA codon), the *spoIVB2*$_{F37L}$ (UUG codon) and the *spoIVB2*$_{F36F}$ (UUC codon)

alleles (Figure 7A). This suggests that multiple *spoIVB2* variants were sufficient to restore

sporulation in an otherwise sporulation deficient strain.

Expression of these plasmids in the *C. difficile* Δ*sspA* Δ*sspB* double mutant strains

showed variation from the previously assessed strain. First, expression of the wild type *spoIVB2*

allele resulted in an approximate 1-$\log_{10}$ increase in spore yield compared to the mutant strain

with an empty vector (Figure 7B). However, expression of the *spoIVB2*$_{F37F}$ (UUU codon) or the

spoIVB2$_{F37L}$ (UUA or UUG codons) restored the spore yield to a higher level than the wild type

spoIVB2 allele. Interestingly, in the *C. difficile* Δ*sspA* Δ*sspB* strain, *spoIVB2*$_{F36F}$ did not

complement the sporulation phenotype as it did in the *C. difficile* Δ*sspB** strain (Figure 7B).

Finally, expression of any of the *spoIVB2* alleles restored sporulation in the *C. difficile* Δ*spoIVB2*

mutant strain (Figure 7C). These data suggest that either altering the F37 codon in either of the

sporulation deficient strains or expressing additional SpoIVB2 can restore sporulation.

spoIVB2$_{A20T}$ *and spoIVB2*$_{F37F}$ *have increased abundance*

We next wanted to determine if the suppressor alleles restore sporulation through

translational differences, rather than transcriptional, we designed a luciferase-based assay [10,

51, 52]. SpoIVB2 is a single span transmembrane protein whose C-terminus is located outside

of the forespore cytoplasm. To the *spoIVB2* gene, we engineered a *ssrA* tag to the 3' end of the

gene. This will tag the protein for degradation by the ClpP protease if the protein is in the

cytoplasm but ClpP will not have access to the C-terminus if it is localized properly [53]. This

assay will allow us to quantify differences in properly-localized SpoIVB2. As a control, the *bitLuc*

gene with and without the *ssrA* tag was put under control of the native *spoIVB2* promoter. We

also coupled the native *spoIVB2* promoter to the *bitLuc* gene and either wild type *spoIVB2,*

spoIVB2$_{A20T}$, or *spoIVB2*$_{F37F}$ and tagged the construct for degradation with a *ssrA* tag. These

constructs were introduced into the wild type and, as a negative control, the *C. difficile* Δ*spo0A*

strain. When in *C. difficile* Δ*spo0A*, all constructs had minimal RLU/OD$_{600}$ values. After

expression in the wild type strain, the control construct containing the *ssrA* tag had significantly

lower normalized luminescence / OD_{600} than the construct without the tag (Figure 8). This shows

that the *ssrA* tag successfully reduced luciferase abundance. After 48 hours of incubation in the

wild type strain, the *spoIVB2*$_{A20T}$ construct had 800x greater levels of luminescence / OD_{600}

compared to wild type and the *spoIVB2*$_{F37F}$ construct had approximately 1,300x greater levels

(Figure 8). These data suggest that sporulation is restored in the suppressor strains because

the identified alleles increased the levels of SpoIVB2 that were present in the sporulating cell.

Restoration of sporulation using different promoters to drive spoIVB2 expression.

To understand if the SASPs allow for continued *spoIVB2* expression during σ^G gene

activation, we generated plasmids containing *spoIVB2* expressed by various promoters. The

spoIVB promoter region served as a lower activity σ^G promoter while the *sspA* promoter region

served as a higher activity σ^G promoter. The spore yield of the *C. difficile* Δ*spoIVB2* strain was

rescued when *spoIVB2* was expressed under the *spoIVB2*, *spoIVB,* or the combined *spoIVB* /

spoIVB2 promoters, suggesting that SpoIVB2 can be present during later stage sporulation

(Figure 9B). However, *spoIVB2* expressed under the *sspA* or the combined *spoIVB2* / *sspA*

promoters did not restore sporulation. Interestingly, the *spoIVB2* / *sspA* promoter combination

when in wild type cells also reduced spore yield 5-$\log_{10}$ (Figure 9A). Similarly, spore yield in *C.*

difficile Δ*sspA* Δ*sspB* was restored when *spoIVB2* was expressed under the *spoIVB, spoIVB2,*

or the *spoIVB* and *spoIVB2* combined promoters (Figure 9C). When expressed under the *sspA*

or the combined *spoIVB2* and *sspA* promoters, restoration did not occur. We hypothesize that

the highly active *sspA* promoter leads to overproduction of SpoIVB2, which is then detrimental

to the sporulating cells.

Discussion

The formation of endospores in *C. difficile* is vital for transmission of disease and the

mechanisms involving spore formation are complex [8]. In prior work, we determined that the *C.*

difficile Δ*sspA* Δ*sspB* strain was halted during sporulation suggesting that the *C. difficile* SASPs

are important for regulating late-stage sporulation, somehow [42]. Here, we built upon our

findings by further exploring the SASP mutant strain using TEM and a selection strategy to

identify potential suppressor mutants.

Oddly, during the course of the prior work, we identified a mutation in the *C. difficile sspA*

gene during the generation of the *sspB* mutant using CRISPR-Cas9 editing. This strain, *C.*

difficile Δ*sspB*; *sspA*$_{G52V}$ (*C. difficile* Δ*sspB**), had a phenotype similar to the *C. difficile* Δ*sspA*

Δ*sspB* strain. This phenotype was likely due to the missense mutation within a conserved

glycine residue. Prior work in *B. subtilis* found that SspC^{G52A} poorly bound DNA [37, 54]. Oddly,

we have since observed a similar off-target effect in the *sspB* gene when targeting *sspA* using

CRISPR-Cas9 mutagenesis. During the process of targeting *sspA* in a *C. difficile* Δ*gpr* strain, an

sspB$_{E64stp}$ allele was also observed upon confirmation of the mutant's DNA sequence. The two

genes are not located in close proximity nor do the constructs for deletion encode this

sequence. We hypothesize that there may be some selective pressure to mutate *sspA* or *sspB*

within a deletion strain.

With further evaluation of SASP mutant strains by TEM, we found that the *C. difficile*

Δ*sspA* Δ*sspB* strain produces forespores that are blocked after the engulfment step, and do not

contain cortex. Cortex is synthesized under SpoVD and potentially SpoVE and is modified by

PdaA, GerS, and CwlD proteins [55-60]. These proteins modify the peptidoglycan to generate

muramic-δ-lactam residues [58-60]. The cortex provides a physical constraint around the spore

core, maintaining size and preventing water from hydrating the Ca-DPA-rich core [8, 61]. In the

absence of cortex, it is likely that some contents in the spore core leak out. This likely explains

our previous findings that the few spores that could be purified from the *C. difficile* Δ*sspA* Δ*sspB*

and the *C. difficile* Δ*sspB** strains contained little CaDPA [42]. Because our RT-qPCR data

showed that *dpaA* and *spoVAC/D/E* transcript levels are similar to wild type levels in these

mutant strains, it is likely that DPA is being synthesized and transported into the spore but

cannot be concentrated into the core without a mature cortex layer.

An EMS mutagenesis strategy to find suppressors of the defect in sporulation of the *C.*

difficile Δ*sspA* Δ*sspB* strain identified mutations in *spoIVB2*. SpoIVB2 is homologous to *B.*

subtillis SpoIVB. Though *B. subtilis* contains the same spore layers and sigma factors that

regulate sporulation in *C. difficile*, the process is more complex. The *B. subtilis* SpoIVB protease

is produced under σ^G control. Located in the *B. subtilis* outer forespore membrane are the

SpoIVFB, SpoIVFA, and BofA proteins [62]. BofA is an inhibitor of the SpoIVFB protease, and

SpoIVFA keeps the proteins localized in the membrane. SpoIVB is secreted through the inner

forespore membrane and processes SpoIVFA, thereby relieving BofA inhibition of SpoIVFB.

Activated SpoIVFB cleaves the pro-peptide from σ^K, resulting in σ^K activation [62].

In addition to its role in σ^K activation, SpoIVB has other functions in *B. subtilis, e.g.*

cleavage of SpoIIQ [63]. SpoIIQ is required for σ^G synthesis and contributes to the formation of

a feeding tube between the mother cell and the forespore compartments. SpoIVB cleaves

SpoIIQ upon completion of engulfment, however, this cleavage is not necessary for spore

formation or any later-stage gene expression [63, 64]. A *spoIVB* null mutant blocks the

formation of fully formed, heat resistant spores [65]. Spores derived from this strain form the

forespore but lack the germ cell wall layer and do not generate mature spores. Interestingly, this

phenotype was independent of SpoIVB's role in the activation of σ^K [65]. An alternative role for

SpoIVB may be in germ cell wall biosynthesis or as a DNA binding regulatory protein.

Even though the *C. difficile* sporulation program does not contain the cross-talk sigma

factor activation or homologs to *bofA*, *spoIVFA*, or *spoIVFB*, it does contain the SpoIVB and

SpoIVB2 paralogs [28]. SpoIVB2 is σ^F-regulated while SpoIVB is σ^G-regulated [47]. *C. difficile*

SpoIVB and SpoIVB2 contain 31% identity to each other and have 36% and 37% identity to *B.*

subtilis SpoIVB, respectively [47].

In our sporulation assays, *spoIVB2*$_{A20T}$ and *spoIVB2*$_{F37F}$ can rescue the mutant

phenotype and form mature, dormant spores. We hypothesize that the A20T and F37F alleles

suppress the phenotype through translational changes. Interestingly, in the identified

spoIVB2$_{F37F}$ strain, the wildtype UUC codon is used in 5.9 / 1000 codons but the UUU codon in

the suppressor strain is used 37.4 / 1000 codons [66]. Also, out of the 18 phenylalanine

residues found in the SpoIVB2 protein, only F37 uses the UUC codon. Even though the codon

changes to one that is used more frequently, this data is based on codon usage across the

whole *C. difficile* genome and not just spore specific genes.

When the wobble position of *spoIVB2*$_{F37}$ was manipulated, sporulation was restored in

both, the *C. difficile* Δ*sspA* Δ*sspB* mutant and the *C. difficile* Δ*sspB** mutant strains (even

though the F37F allele was only identified in the former strain). Also of note, it is likely that the

mutation to the UUU codon was the only identified change after EMS treatment, instead of the

UUA or UUG codons, due to the nature of EMS mutagenesis which results in transition

mutations. However, it is likely that the specific manipulations do not matter as long as the

codon increases translation efficiency compared to the UUU codon. We analyzed transcript

variation among strains for various genes, including *spoIVB2.* In this data, the transcripts for

spoIVB2 in representative EMS strains, for both the *spoIVB2*$_{A20T}$ and *spoIVB2*$_{F37F}$ alleles,

trended toward being downregulated, though this difference was only ~4 fold and did not meet

statistical significance. These results support the hypothesis that SASPs are necessary to

further activate *spoIVB2* transcription. Unfortunately, *C. difficile* sporulation is asynchronous and

samples from any time point contain cells in every stage of sporulation. This could explain why

the fold changes are small and variable across all strains and all transcripts analyzed. Because

of this noise, it is difficult to draw definitive conclusions from the RT-qPCR data.

In our working model for how the *C. difficile* SASPs influence *spoIVB2*, *w*e hypothesize

that SASP binding could activate gene expression by enhancing interactions with RNA

polymerase. Because *spoIVB2* is under σ^F-control, σ^G-produced SASPS could further activate

spoIVB2 expression to maintain SpoIVB2 abundance in the spore (Figure 10). In the absence

of *C. difficile sspA* and *sspB,* SpoIVB2 activity is reduced (Figure 10). Since the SASPs are not

present in the suppressor strains to lead to activated *spoIVB2* transcription, we hypothesized

that the *spoIVB2* alleles identified altered translation rates and, thus, increased SpoIVB2 levels

(Figure 10). Supporting this, our BitLuc data showed a significant increase in RLU for translation

of the A20T and F37F alleles compared to wild type.

Separate from how the *spoIVB2* alleles restore sporulation to the SASP mutant strains,

what is the function of SpoIVB2 during sporulation? While it is possible that the *C. difficile*

SpoIVB and SpoIVB2 proteins retain a function in SpoIIQ cleavage, in *C. difficile,* SpoIIQ does

not appear to be cleaved during *C. difficile* sporulation [67]. However, unlike in *B. subtilis*, *C.*

difficile SpoIIP has a cleaved form that is only present in cells that have completed engulfment

[67]. SpoIIP is an amidase and endopeptidase that works in concert with SpoIID to restructure

peptidoglycan during forespore engulfment. In a *C. difficile* Δ*spoIIP* strain, the leading edge of

engulfment does not progress, so engulfment does not occur [67]. In recent collaborative work,

we found that *C. difficile* SpoIVB2 does cleave SpoIIP *in vitro* and *in vivo* [68]. We thought it

possible that SpoIIP needs to be cleaved post engulfment completion to allow the following

stages of sporulation to continue. However, we found that strains containing *spoIIP* with an

altered cleavage site (SpoIVB2 is unable to cleave this form *in vitro*) could still form mature

spores [68]. While it is possible that this altered SpoIIP could still be processed by SpoIVB2, just

highly inefficiently, it is likely that there are uncharacterized targets / functions of SpoIVB2

during sporulation.

To better understand if the SASPs function similarly among other organisms, we also

tested the ability of the *B. subtilis sspA* gene to complement sporulation and UV phenotypes in

C. difficile mutants. When expressed under the *C. difficile sspA* native promoter region, *B.*

subtilis sspA can partially restore sporulation and UV resistance. In prior work from the Setlow

lab [69], the authors suggested that SASPs may affect forespore transcription, likely by

physically blocking RNA polymerase when binding in high concentrations. Furthermore, *in vitro*

transcription assays in *B. subtilis* show that less *in vitro* transcription occurs when SASPs are

incubated with DNA. However, transcription occurs in the absence of SASPs or in the presence

of a SASP variant with poor DNA binding ability [37]. However, they also found evidence of

some genes (mainly later stage sporulation genes) having higher / lower transcription in the *B.*

subtilis Δ*sspA* Δ*sspB* strain [69]. These data indicate that the SASPs could be regulating

sporulation in the forespore of both *C. difficile* and *B. subtilis*, suggesting that the different

phenotypes observed between the Δ*sspA* Δ*sspB* double mutants in the two organisms lie in the

differences between the mechanism of compartmental signaling during sporulation. This leads

to further questions about whether the genes / regions of DNA that are influenced by the SASPs

and how the SASPs may potentiate these affects differs between the two organisms.

Overall, this study gives insight into the sporulation process and regulation in *C. difficile.*

It is likely that the SASPs have binding "hotspots" where in low concentrations they

preferentially bind to influence transcription. Although the RT-qPCR data did not show many

transcriptional changes, we hypothesize that the SASPs are influencing transcription of target

genes. It is possible that the time of extraction was not ideal for capturing transcriptional

changes, the change is small enough that the variability in data due to sporulation being

asynchronous could be enough to hide the effects, or, though unlikely based on our data, the

SASPs have different targets than those tested. This study also highlights the importance of *C.*

difficile SpoIVB2 during sporulation even though its exact role is still unknown. Further work

needs to be completed to understand the influence of SpoIVB2 during sporulation and to

determine other potential targets for the SASPs.

Materials and Methods

Bacterial growth conditions: *C. difficile* strains were grown in a Coy anaerobic chamber at

~4% H_2, 5% CO_2, and balanced N_2 at 37 °C [70]. Strains were grown in / on brain heart infusion

(BHI), BHI supplemented with 5 g / L yeast extract (BHIS), 70:30 (70% BHIS, 30% SMC) or

tryptone yeast (TY) medium. 0.1% L-cysteine was added to BHI and BHIS while 0.1%

thioglycolate was added to TY. Media was supplemented with thiamphenicol (10 µg / mL),

taurocholate [TA] (0.1%), cycloserine (250 µg / mL), kanamycin (50 µg / mL), lincomycin (20 µg

/ mL), rifampicin (20 µg / mL), ethylmethane sulfonate [EMS] (1%), or D-xylose (0.5% or 1%)

where indicated. *E. coli* strains were grown on LB at 37 °C and supplemented with

chloramphenicol (20 µg / mL) or ampicillin (100 µg / mL). *B. subtilis* BS49 was grown on LB

agar plates or in BHIS broth at 37 °C and supplemented with 2.5 µg / mL chloramphenicol or 5

µg / mL tetracycline.

Plasmid construction: All cloning was performed in *E. coli* DH5α.

For construction of the *C. difficile* CD630Δ*erm sspA*-targeting CRISPR vector, pHN120,

500 bp of upstream homology was amplified from CD630Δ*erm* genomic DNA with primers

5'sspA_MTL and 3'sspA_UP while the downstream homology arms were amplified with

5'sspA_down and 3' sspA_xylR. These were inserted into pKM197 at the *Not*I and *Xho*I sites

using Gibson assembly [71]. The gRNA gBlock (Integrated DNA Technologies, Coralville, IA)

CRISPR_sspA_165 was inserted at the *Kpn*I and *Mlu*I sites. pHN120 was then used as the

base plasmid to change the *Tn916* oriT for the *traJ* oriT at the *Apa*I sites, resulting in pHN131.

traJ was amplified from pMTL84151 with primers 5'traJ and 3'traJ. The gRNA was then

replaced with gBlock CRISPR_sspA_135 at the *Kpn*I and *Mlu*I restriction sites, generating

pHN138, which was used to make the deletion.

For generating the *C. difficile* CD630Δ*erm sspB* targeting CRISPR vector, pHN121, the

upstream homology was amplified from CD630Δ*erm* genomic DNA with 5' sspB UP and 3' sspB

UP and the downstream homology with 5' sspB DN and 3' sspB_xylR. These homology arms

were inserted into pKM197 at *Not*I and *Xho*I restriction sites using Gibson assembly [71]. The

gRNA gBlock CRISPR_sspB_144 was also inserted into the *Kpn*I and *Mlu*I sites. The oriT was

changed from *Tn916* to *traJ* by amplifying *traJ* from pMTL84151 with 5'traJ and 3'traJ and

inserting in the *apa*I sites to generate pHN132.

For generating the *C. difficile* R20291 *spoIVB2* targeted CRISPR plasmid, the upstream

homology arm was amplified from R20291 with primers 5' CDR20291_0714 UP and 3'

CDR20291_0714 UP while the downstream was amplified with 5' CDR20291_0714 DN and 3'

CDR20291_0714 DN. These were inserted into pKM197 at the *Not*I and *Xho*I sites using

Gibson assembly [71]. The gRNA was amplified from pKM197 using primers CDR20291_0714

gRNA 3 and 3' gRNA_change. This fragment was inserted into the *Kpn*I and *Mlu*I sites using

Gibson assembly, generating pHN157 [71].

Plasmid pHN149 was generated by amplifying the *traJ* oriT from pMTLYN4 with primers

5' tn916.traJ and 3'traJ and the *Tn916* oriT from pJS116 with 5'Tn916ori_gibson and 3'

tn916.traJ. These were inserted into the A*pa*I site of pMTL84151.

For the generation of pHN122, pHN123, and pHN127, the *spoIVB2* gene and promoter

regions were amplified from HNN37, HNN19, and R20291, respectively, using 5'

CDR20291_0714 and 3' CDR20291_0714. These fragments were inserted into pJS116 (for

pHN122 and pHN123) or pHN149 (for pHN127) at the *Not*I and *Xho*I sites using Gibson

assembly [71].

For pHN145, pHN146, and pHN147, the first segment of DNA was amplified from

R20291, pHN122, or pHN123, respectively, using 5' CDR20291_0714 and 3' 0714_S301A. The

second segment of DNA was amplified from pHN127 for all 3 plasmids using 5' 0714_S301A

and 3' CDR20291_0714. These two fragments were inserted using Gibson assembly into

pJS116 at the *Not*I and *Xho*I sites [71].

The CD630Δ*erm sspA* gene and promoter region were amplified from CD630Δ*erm* with

the primers 5'sspA_MTL and 3' sspA.pJS116. This fragment was inserted into pMTL84151 at

the *Not*I and *Xho*I sites using Gibson assembly, generating pHN152 [71].

For pHN153, the CD630Δ*erm sspA* gene and promoter region were amplified with

5'sspA_MTL and 3' sspAsspB. The CD630Δ*erm sspB* gene and promoter region were amplified

with 5' sspAsspB and 3'sspBpJS116. These fragments were inserted into pMTL84151 at the

*Not*I and *Xho*I sites using Gibson assembly [71].

The CD630Δ*erm sspB* gene and promoter region were amplified using 5' sspB UP and

3'sspBpJS116. This fragment was inserted into pHN149 at the *Not*I and *Xho*I sites using Gibson

assembly to generate pHN176 [71].

pHN271 and pHN272, the *spoIVB2*$_{A20T}$ / *spoIVB2*$_{F37F}$ theophylline allelic exchange

plasmids, respectively, were generated by amplifying the *spoIVB2* region with homology from

pHN123 for pHN271 and pHN122 from pHN272 with primers 5' spoIVB2_theo and 3'

spoIVB2_theo. These fragments were inserted into pJB94 at the *Not*I and *Xho*I sites using

Gibson assembly [71, 72].

pJB96 was generated by amplifying the *sacB* gene from pJB94 with primers 5'sacB_UP

and sacB_3'_XhoI and inserted into pHN149 at the *Not*I and *Xho*I restriction sites by Gibson

assembly [71].

The *spoIVB2* under *sspA* promoter control plasmid, pHN312, was generated by

amplifying the *sspA* promoter from R20291 with the primers 5'sspA_MTL and 3'

PsspA_spoIVB2. The *spoIVB2* gene was amplified from R20291 with the primers 5'

spoIVB2_PsspA and 3' CR20291_0714. These fragments were cloned into pJB96 at the *Not*I

and *Xho*I sites by Gibson assembly. pHN329, *spoIVB2* under the *spoIVB* promoter, was made

by amplifying the *spoIVB* promoter from R20291 with primers 5' spoIVB.pHN149 and 3'

PspoIVB_spoIVB2. The spoIVB2 gene was amplified with 5' spoIVB2_PspoIVB and 3'

CDR20291_0714 from R20291 template DNA. The fragments were assembled with Gibson

assembly into pJB96 at *Not*I and *Xho*I cut sites [71]. The plasmid pHN330 containing *spoIVB2*

driven by the *spoIVB* and *spoIVB2* promoters was generated by amplifying the *spoIVB2*

promoter from R20921 DNA with primers 5' PspoIVB2_pHN149 and 3'

PspoIVB(100)_PspoIVB2. The *spoIVB* promoter was amplified with primers 5'

PspoIVB2_PspoIVB and 3' PspoIVB_spoIVB2 from R20291 DNA. The *spoIVB2* gene was

amplified from R20291 with primers 5' spoIVB2_PspoIVB and 3' CDR20291_0714. These

fragments were assembled into pJB96 at *Not*I and *Xho*I cut sites using Gibson assembly [71].

The plasmid pHN331 containing *spoIVB2* driven by the *sspA* and *spoIVB2* promoters was

generated by amplifying the *spoIVB2* promoter from R20921 DNA with primers 5'

PspoIVB2_pHN149 and 3' PsspA_PspoIVB2. The *sspA* promoter was amplified with primers 5'

PsspA_PspoIVB2 and 3' PsspA_spoIVB2 from R20291 DNA. The *spoIVB2* gene was amplified

from R20291 with primers 5' spoIVB2_PsspA and 3' CDR20291_0714. These fragments were

assembled into pJB96 at *Not*I and *Xho*I cut sites using Gibson assembly [71].

The luciferase plasmids pHN335 through pHN337 were generated by amplifying the

spoIVB2 promoter region from R20291 using primers 5' CDR20291_0714 and 3'

spoIVB2_homol. The *spoIVB2* gene fragments were amplified with primers 5'

spoIVB2_gene_homol and 3' spoIVB2end_lrgBit from R20291 for pHN335, pHN123 for

pHN336, and pHN122 for pHN337. The *bitLuc* gene fragment with a *ssrA* tag was amplified

from pMB81 with primers 5' lrgBit_spoIVB2end and 3' luciferase_ssrA_pHN149. These 3

fragments were cloned into pJB96 at the *Not*I and *Xho*I restriction sites using Gibson assembly

[71]. For the control luciferase plasmids, pHN338-339, the *spoIVB2* promoter region was

amplified from R20291 using primers 5 CDR20291_0714 and 3' spoIVB2_bitLuc. For pHN338,

the *bitLuc* gene portion with a *ssrA* tag was amplified from pMB81 with primers 5'

bitLuc_PspoIVB2 and 3' luciferase_ssrA_pHN149. For pHN339, the *bitLuc* gene was amplified

from pMB81 with primers 5' bitLuc_PspoIVB2 and 3' luciferase_pHN149. The *spoIVB2* promoter

fragment and the luciferase fragments were cloned into pJB96 at the *Not*I and *Xho*I restriction

sites using Gibson assembly [71].

pHN220 was generated by amplifying the *sspA* promoter region from *C. difficile* R20291

with primers 5'sspA_MTL and 3'PsspA_BS49. The *sspA* gene was amplified from *B. subtillis*

BS49 with primers 5' sspA_BS49 and 3' sspA_BS49. These fragments were put into the

pHN149 backbone at the *Not*I and *Xho*I sites by Gibson assembly [71].

To generate pHN208, the promoter region through F36 of *CDR20291_0714* was

amplified with 5' CDR20291_0714 and 3' spoIVB2 F36F, while the F36 through the end of

CDR20291_0714 was amplified with 5' spoIVB2 F36F and 3' CDR20291_0714, both using

pHN127 as the DNA template. These fragments were inserted by Gibson assembly into the

pHN149 plasmid backbone at the *Not*I and *Xho*I sites [71]. pHN218 was generated by

amplifying from the pHN127 template DNA *CDR20291_0714* promoter region through F37 with

primers 5' CDR20291_0714 and 3' spoIVB2 F37.UUA and the fragment with F37 through the

end of the gene was amplified by 5' spoIVB2 F37.UUA and 3' CDR20291_0714. These

fragments were inserted in pHN149 at the *Not*I and *Xho*I sites through Gibson assembly [71].

Similarly, pHN219 was generated but used 5' CDR20291_0714 with 3' spoIVB2 F37.UUG for

the first fragment and 5' spoIVB2 F37.UUG and 3' CDR0291_0714 for the second fragment,

both with pHN127 as the template DNA. These were also inserted by Gibson assembly into

pHN149 at the *Not*I and *Xho*I sites [71].

All plasmids were sequenced to confirm the correct sequence of the inserts.

Conjugations: For conjugations between *C. difficile* and *E. coli* HB101 pRK24, 5 mL of LB

supplemented with chloramphenicol and ampicillin was inoculated with a colony from the HB101

pRK24 transformation. Concurrently, *C. difficile* strains were cultured in 5 mL BHIS broth. After

approximately 16 hours of incubation, 1 mL of *C. difficile* culture was heat shocked at 52 °C for

5 minutes, in the anaerobic chamber, and then allowed to cool. While heat shocking, 1 mL of *E.*

coli culture was centrifuged at 2,348 x g for 5 minutes and the supernatant poured off. The *E.*

coli pellets were passed into the chamber and resuspended with the cooled *C. difficile* culture.

20 µL spots were plated onto BHI. The next day, growth was scraped into 1 mL BHIS broth and

distributed onto BHIS plates supplemented with thiamphenicol, kanamycin, and cycloserine

(TKC) or TKC plus lincomycin (TKLC) for the 2-plasmid CRISPR system.

For conjugations between *C. difficile* and *B. subtilis* BS49, the plasmids generated in

DH5α were used to transform *E. coli* MB3436 (a *recA+* strain of *E. coli*) and plasmid purified.

This plasmid preparation was then used to transform BS49. *C. difficile* was cultured in 5 mL

BHIS broth overnight. After approximately 16 hours, the *C. difficile* culture was back diluted 1:20

and grown for 4 hours. *B. subtilis* was grown for 4 hours in 5 mL BHIS broth supplemented with

chloramphenicol and tetracycline. After incubation, the *B. subtilis* cultures were passed into the

chamber and 100 µL of BS49 and 100 µL of *C. difficile* were spread onto TY plates. The next

day, the growth was resuspended in 1 mL of BHIS broth and distributed between BHIS plates

with thiamphenicol and kanamycin. Colonies were screened by streaking colonies onto BHIS

supplemented with thiamphenicol and kanamycin and to BHIS supplemented with tetracycline to

identify isolates that were tet-sensitive (do not contain the *Tn*916 transposon).

PCR was used to confirm strains and plasmids in each conjugate.

CRISPR induction: For induction, colonies were passaged on TY agar supplemented with 1%

xylose and thiamphenicol [42, 73]. Mutants were detected by PCR and the plasmid was cured

by heat shocking overnight cultures and isolating colonies that had lost their antibiotic

resistance.

Induction of R20291 pHN138 resulted in the *C. difficile* CD630Δ*erm* Δ*sspA* mutant,

HNN45. R20291 pHN132 induction resulted in the *C. difficile* CD630Δ*erm* Δ*sspB* mutant,

HNN43. To generate the *C. difficile* CD630Δ*erm* Δ*sspA* Δ*sspB* strain, the pHN132 vector was

induced in HNN45, resulting in HNN46. The *C. difficile CDR20291_0714* (*spoIVB2*) mutant

HNN49 was produced from induction of R20291 pHN157.

Theophylline allelic exchange: Strains were generated as previously described [72]. Briefly,

transconjugants were passaged on medium with thiamphenicol and no theophylline to

encourage integration of the plasmid into the chromosome. Once integration occurred, the

isolates were passaged on plates containing theophylline to encourage excision. HNN57 was

generated from the passaging of R20291 pHN272. HNN60 was generated from the passaging

of R20291 pHN271. HNN64 was generated from the passaging of *C. difficile* Δ*sspA* Δ*sspB*

pHN271. HNN57 was generated from the passaging of *C. difficile* Δ*sspA* Δ*sspB* pHN272.

Phase contrast imaging: The strains were inoculated onto 70:30 sporulation media and

incubated for 6 days. After, the samples were fixed in a 4% paraformaldehyde and 2%

glutaraldehyde solution in 1x PBS. The samples were imaged on a Leica DM6B microscope at

the Texas A&M University Microscopy and Imaging Center Core Facility (RRID:SCR_022128).

TEM: For sporulating cells, the relevant strains were incubated in the anaerobic chamber on

sporulation media for 6 days, and then the growth scraped with an inoculation loop into 1,950

μL of fixative (5% glutaraldehyde, 2% acrolein in 0.05 M HEPES, pH 7.4) in a 2.0 mL

microcentrifuge tube. The samples were incubated in the fixative overnight at 4 °C. The

following day, the fixed samples were centrifuged for 5 min at >14,000 × g, and post-fixed and

stained for 2 hours with 1% osmium tetroxide in 0.05 M HEPES, pH 7.4.

The samples were then centrifuged and washed with water 5 times, and dehydrated with

acetone, using the following protocol: 15 minutes in 30%, 50%, 70%, 90% acetone each, then

100% acetone 3 changes, 30 minutes each. At the final wash step, a small amount of acetone,

barely covering the pellets, was retained to avoid rehydration of the samples. The samples were

then infiltrated with modified Spurr's resin (Quetol ERL 4221 resin, Electron Microscopy

Sciences, RT 14300) in a Pelco Biowave processor (Ted Pella, Inc.), as follows: 1:1

acetone:resin for 10 minutes at 200 W – no vacuum, 1:1 acetone:resin for 5 minutes at 200 W –

vacuum 20'' Hg (vacuum cycles with open sample container caps), 1:2 acetone:resin for 5

minutes at 200 W – vacuum 20'' Hg, 4 x 100% resin for 5 minutes at 200 W – vacuum 20'' Hg.

The resin was then removed, and the sample fragments transferred to BEEM conical tip

capsules prefilled with a small amount of fresh resin, resin added to fill the capsule, and

capsules left to stand upright for 30 minutes to ensure that the samples sank to the bottom. The

samples were then polymerized at 65 °C for 48 hours in the oven, then left at RT for 24 hours

before sectioning. 70-80 nm samples were sectioned by Leica UC / FC7 ultra-microtome (Leica

Microsystems), deposited onto 300 mesh copper grids, stained with uranyl acetate / lead citrate

and imaged. All ultrathin TEM sections were imaged on JEOL 1200 EX TEM (JEOL, Ltd.) at 100

kV, images were recorded on SIA-15C CCD (Scientific Instruments and Applications) camera at

the resolution of 2721 x 3233 pixels using MaxImDL software (Diffraction Limited). Images were

subsequently adjusted for brightness / contrast using Fiji [74]. All equipment used is located at

Texas A&M University Microscopy and Imaging Center Core Facility (RRID:SCR_022128).

Sporulation Assay: Sporulation assays were completed as previously described [42]. Briefly,

70:30 plates were inoculated with the indicated strains and grown for 48 hours. 1/3 of the plate

was harvested into 1 mL of PBS. 500 µL of the culture was treated for 20 minutes with 300 µL of

100% EtOH and 200 µL of dH_2O to make a 30% final solution. After incubation, the samples

were serially diluted in PBS + 0.1% TA and plated onto BHIS supplemented with TA to

enumerate spores. The CFUs derived from spores were $\log_{10}$ transformed.

Spore Purification: Spores were purified as previously described [46, 75]. Briefly, the cultures

from 70:30 agar medium were scraped into 1 mL of dH_2O and left overnight at 4°C. The next

day, the pellets were resuspended and centrifuged for 1 minute at max speed. The upper layer

of cell debris was removed and the sample was resuspended in 1 mL dH_2O. Again, the tubes

were centrifuged and the upper layer removed. This was repeated approximately 5 times until

the spore pellet was relatively free of debris. The 1 mL of spores in dH_2O was loaded onto 50%

sucrose and centrifuged at 4,000 x g for 20 minutes 4°C. The spore pellet was then washed as

described above approximately 5 times and then stored at 4°C until future use.

UV exposure: UV experiments were performed as previously described [42]. Briefly, 1×10^8

spores / mL in PBS were treated for 10 minutes with constant agitation. The T_0 and T_{10} samples

were serially diluted and plated onto rich medium containing germinant taurocholic acid (TA).

Treated spore counts were normalized to untreated and then this ratio was normalized to the

ratio for wild type spores.

EMS treatment: For EMS treatment, the HNN04 or HNN05 strains with pJS116 were used to

help prevent contamination by providing antibiotic selection. Overnight cultures were back

diluted to OD_{600} 0.05 in 15 mL of BHIS + Tm [44, 45]. The cultures were grown to an OD_{600} of

0.5. The culture was split into 2 tubes of 5 mL, each. One tube served as the negative control

and one tube was treated with 1% EMS. The cultures were grown for 3 hours with vigorous

shaking every 30 minutes (to keep the EMS in solution). The cultures were passed out of the

chamber and centrifuged at 3,000 x g for 10 minutes, passed into the chamber, decanted,

resuspended with 10 mL BHIS to wash and then passed out and centrifuged again. This wash

step was repeated 1 more time for a total of 2 washes. After the second wash, the cell pellet

was resuspended with 1 mL of BHIS and deposited into 39 mL BHIS + Tm to recover overnight.

The next day, to determine mutagenesis rates, 10 µL, 25 µL, and 50 µL volumes were

each plated onto BHIS rifampicin agar and CFUs were counted after 24 - 48 hours. From the

EMS (+) culture, 50 µL was plated onto 20 BHIS Tm5 agar plates and left in the chamber to

incubate for 5 days. For the EMS (-) culture, a whole genome prep was performed as described

below.

After the incubation period, the plates were scraped into individual tubes with 1 mL dH_2O and

left overnight at 4 °C. The tubes were purified to remove cell debris as done with spore

purification described above. The samples were combined to one tube and heated at 65 °C for 1

hour, with intermittent vortexing. The sample was then distributed between 20 BHIS Tm5 plates

for another round of incubation. This enrichment step was completed 3 times before isolates

were selected and PCR was used to confirm the genotype (to confirm that wildtype

contamination did not occur during the selection). After confirmation, the samples were plated

onto 70:30 Tm5 and incubated for 5 days. These samples were then checked under a phase

contrast microscope for spores. Genomic DNA was purified from samples that had spores and

sent for whole genome resequencing at Microbial Genome Sequencing Center (MiGS;

Pittsburgh, PA).

Whole genome preparation: 4 tubes of 10 mL each were inoculated overnight for

approximately 18 hours (or the 40 mL of culture from EMS (-) strains were used). The next day,

the samples were centrifuged at 4,000 x g for 10 minutes, 4 °C. They were decanted, then

resuspended with 1 mL TE buffer (10 mM Tris-HCl, 1 mM EDTA). Samples were centrifuged

again, decanted, and resuspended with 200 µL of genomic DNA solution (34% sucrose in TE

buffer) and transferred to a 2 mL Eppendorf tube (for each strain, the 4 tubes are kept

separate). The tubes were incubated at 37°C for 2 hours. Then, 100 µL of 20% Sarkosyl and 15

µL of 10 mg / mL RNase A were added to the sample and incubated at 37 °C for 30 minutes.

After this incubation, 15 µL of proteinase K solution was added and incubated 37 °C for 30

minutes at. The samples were brought up to 600 µL with TE buffer.

600 µL of 25:24:1 phenol/chloroform/isoamyl alcohol was added to the samples and

were rocked gently for 20 minutes. After the incubation, the samples were centrifuged for 10

minutes at max speed. The upper layer was transferred to a new tube with a cut pipette tip (so

as not to shear the DNA) and 600 µL chloroform was added to the sample and rocked for

another 20 minutes. The centrifugation, sample transfer, and chloroform treatment were

repeated for a total of 3 times. After which, the upper phase was transferred to a new tube and

precipitated at -20 °C overnight with 50 µL of 3 M sodium acetate, pH 5.2, and 3 volumes of

cold 95% ethanol.

After precipitation, one tube from each strain was centrifuged 15 minutes at max speed,

4 °C. The supernatant was discarded and the solution from the second tube was transferred to

the tube with the DNA pellet and centrifuged again. This was repeated until the DNA from all 4

tubes was combined into one pellet. The DNA pellet was washed with 500 µL of 70% ethanol

and centrifuged again. The samples were decanted and allowed to dry at room temperature

until all of the ethanol was evaporated (approximately 60 – 90 minutes). After drying, 500 µL of

either dH_2O or TE buffer was added, and the samples were rocked overnight to allow the pellets

to dissolve.

RNA extraction and processing: Strains were plated onto 70:30 media for 11 hours before

extraction. RNA extraction was performed using the FastRNA Pro Blue Kit (MP Biomedicals,

Solon, OH). Briefly, the culture was scraped into 1 mL of PBS and centrifuged 2,348 x g for 5

minutes. The pellet was resuspended in 1 mL of RNApro solution and transferred to the

provided tubes with lysing Matrix B. The cells were lysed in an MP FastPrep-24 bead beater for

40 seconds on and 20 seconds off for a total of 2 rounds. Further processing followed the

FastRNA Pro Blue Kit protocol except that the RNA was precipitated overnight, and the

remainder of the protocol was continued the next day.

Contaminating DNA was removed using the TURBO DNA-free kit (Invitrogen, Waltham,

MA). 10 µg of RNA was treated 3 times with DNase following the protocol provided in the kit.

The RNA was precipitated at -20 °C overnight with 0.1 volume of 3 M sodium acetate, 5 µg of

glycogen, and an equal volume of 100% ethanol. RNA was recovered by centrifuging at 13,000

x g at 4 °C for 30 minutes. The pellet was washed 2 times with 70% cold ethanol. The pellet was

air-dried at room temperature and then resuspended in dH$_2$O.

cDNA was generated using the Superscript III First-Strand Synthesis System (Invitrogen,

Waltham, MA) reagents and protocol.

RT-qPCR: qPCR was performed with PowerUP SYBR Green Master Mix (Applied Biosystems,

Waltham, MA) according to provided protocol on an Applied Biosystems QuantStudio 6 Flex

Real-Time PCR system. Primers used are as follows: *rpoA*: 5' rpoA & 3' rpoA; *sspA*: 5'

sspA_qPCR & 3' sspA_qPCR; *sspB*: 5' sspB_qPCR & 3' sspB_qPCR; *sleC*: 5'sleC_qPCR &

3'sleC_qPCR; *spoVT*: 5' spoVT_qPCR & 3' spoVT_qPCR; *pdaA*: 5' pdaA_qPCR & 3'

pdaA_qPCR; *spoIVA*: 5'spoIVA_qPCR & 3'spoIVA_qPCR; *spoIVB*: 5' spoIVB_qPCR_1 & 3'

spoIVB_qPCR_1; *spoIVB2*: 5' spoIVB2_qPCR_1 & 3' spoIVB2_qPCR_1; *spoIIP*: 5'

spoIIP_qPCR_1 & 3' spoIIP_qPCR_1; *dpaA*: 5' dpaA_qPCR & 3' dpaA_qPCR; *spoVAC*: 5'

spoVAC_qPCR & 3' spoVAC_qPCR; *spoVAD*: 5' spoVAD_qPCR & 3' spoVAD_qPCR; *spoVAE*:

5' spoVAE_qPCR & 3' spoVAE_qPCR.

Analysis was performed by the ΔΔCT method with comparison to internal control *rpoA*

and then mutant strains compared to WT (R20291) [76].

Luciferase Assays: Overnight cultures were back diluted to OD$_{600}$ = 0.05 in BHIS

supplemented with thiamphenicol. The cultures were grown for 48 hours. Post incubation, the

OD$_{600}$ was recorded and the cultures were used for the Nano-Glo Luciferase assay (Promega,

Madison, WI). Briefly, 100 μL of culture was put into a standard Optiplate White bottom 96 well

plate. 20 μL of buffer/substrate mixture, prepared as per the kit instructions, was added to the

culture. The plate was shaken for 3 minutes before the RLU was determined. The RLU was

normalized to the OD_{600} [51, 52].

For each trial, 2 technical replicates were measured in different positions in the 96 well

plate, due to some variation in measurements based on location within the plates.

Table 1: **Shortened list of potential suppressor mutations:** The mutations that could

potentially suppress the sporulation defect in EMS treated isolates. The underlined strains were

derived from the *C. difficile ΔsspB** strain while the nonunderlined strains were derived from the

C. difficile ΔsspA ΔsspB strain.

Gene	Function	Isolates	Mutation
rpoB	RNA polymerase beta subunit	HNN37	P893S
rpoC	RNA polymerase beta' subunit	HNN40	H94Y
rpoC	RNA polymerase beta' subunit	HNN32	W225 stp
rpoC	RNA polymerase beta' subunit	HNN40, HNN22	S315F
rpoA	RNA polymerase alpha subunit	HNN51, HNN19	S281F
CDR20291_0714	Stage IV sporulation protein	HNN19	A20T

CDR20291_0714	Stage IV sporulation protein	HNN33, HNN35, HNN37, HNN38, HNN41, HNN48	F37F ; Synonymous
sigG	Forespore sporulation sigma factor	HNN39	M97I
sspA	Small acid soluble protein	<u>HNN26</u>, <u>HNN28</u>	V52G ; Reversion
spoVT	Stage V sporulation protein	HNN51	P39S

Figure Legends

Figure 1. Impact of *C. difficile* CD630Δ*erm sspA* and *sspB* mutations on sporulation and UV resistance. A) Day 6 sporulating cultures were fixed in 4% formaldehyde and 2% glutaraldehyde in PBS and imaged on a Leica DM6B microscope. The red arrow represents an immature spore. B) Strains were grown on 70:30 sporulation medium for 48 hours and the cultures then were treated with 30% ethanol and plated onto rich medium supplemented with TA to enumerate spores. Spore yield was calculated by log_{10} transformation of the CFUs derived from spores. C) Spores were exposed to UV for 10 minutes with constant agitation. After treatment, they were serially diluted and plated onto rich medium supplemented with TA. The ratio of treated to untreated CFUs of the mutant strains was then compared to the ratio from WT. pEV indicates an empty plasmid within the strain. All data points represent the average of

three independent experiments. Statistical analysis by one-way ANOVA with Šídák's multiple

comparisons test. * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Figure 2. *B. subtilis sspA* complements *C. difficile* SASP mutant sporulation and UV

defects. A) Spore yield of the indicated strain was determined as described in Figure 1. B)

Spores were exposed to UV for 10 minutes with constant agitation. After treatment, they were

serially diluted and plated onto rich medium supplemented with TA. The ratio of treated to

untreated CFUs of the mutant strains was then compared to the ratio from WT. pEV indicates

an empty plasmid within the strain. All data points represent the average of three independent

experiments. Statistical analysis by two-way ANOVA with Šídák's multiple comparisons test. *

P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Figure 3. SASP mutants do not form the cortex layer. Day 6 sporulating cultures of wild type

and mutant strains containing an empty vector (pEV) or the indicated plasmids were prepared

for TEM. The coat layer is indicated with a white arrow, while the cortex layer is indicated with a

black arrow.

Figure 4. Mutations in *spoIVB2* suppress the *C. difficile sspA sspB* mutant phenotype.

Sporulation was completed over 48 hours and the spore yield was determined. A) Sporulation of

indicated strains with an empty vector (pEV) or wild type *spoIVB2* B) Sporulation of *C. difficile*

R20291 with the indicated *spoIVB2* allele expressed from a plasmid. C) Sporulation assay of *C.*

difficile R20291 Δ*sspB** with the indicated *spoIVB2* allele expressed from a plasmid. D)

Sporulation assay of R20291 Δ*sspA* Δ*sspB* with the indicated *spoIVB2* allele expressed from a

plasmid. All data points represent the average from at least three independent experiments.

Statistical analysis by two-way ANOVA with Šídák's multiple comparisons test* $P<0.05$, **

$P<0.01$, *** $P<0.001$, **** $P<0.0001$.

Figure 5. *C. difficile* Δ*spoIVB2* has a sporulation defect. A) Day 6 cultures were fixed in 4%

formaldehyde and 2% glutaraldehyde in PBS and imaged on a Leica DM6B microscope. B)

Spore yield of the indicated strain was determined as described in Figure 1. pEV indicates an

empty plasmid within the strain. All data points represent the average from three independent

experiments. Statistical analysis by one-way ANOVA with Šídák's multiple comparisons test. *

$P<0.05$, ** $P<0.01$, *** $P<0.001$, **** $P<0.0001$.

Figure 6. SpoIVB2 is required for cortex synthesis. Day 6 sporulating cultures were

prepared for TEM. A field of view image is shown for the *C. difficile* Δ*spoIVB2* mutant while the

remainder of the images are zoomed into the sporulating cell / spore. The coat layer is indicated

with a white arrow, while the cortex layer is indicated with a black arrow. pEV indicates an

empty plasmid within the strain.

Figure 7. Manipulation of *spoIVB2*$_{F36 / F37}$ restores sporulation. Spore yield of the indicated

strain was determined as described in Figure 1. The plasmids expressing the *spoIVB2*$_{F37}$ alleles

and the *spoIVB2*$_{F36F}$ allele were assessed in the strains A) *C. difficile* Δ*sspB**, B) *C. difficile*

Δ*sspA* Δ*sspB*, C) *C. difficile* Δ*spoIVB2.* pEV indicates an empty plasmid within the strain. All

data represents the average of five independent experiments. Statistical analysis by ANOVA

with Šídák's multiple comparison test. * $P<0.05$, ** $P<0.01$, *** $P<0.001$, **** $P<0.0001$.

Figure 8. Luciferase assays show minimal differences between *spoIVB2* protein levels.

Steady-state protein levels of the indicated strain were determined using alleles with an

engineered *ssrA* tag (as described in the materials and methods). The cultures were grown for

48 hours and the OD_{600} and the RLUs were determined. The RLUs were normalized to the

OD_{600} of each culture. All data represents the average of six independent experiments. One

data point in the R20291 *spoIVB2*$_{F37F}$_*bitLuc_ssrA* data set was found to be an outlier by the

ROUT test and was removed from analysis. Statistical analysis by ANOVA with Šídák's multiple

comparison test. * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Figure 9. Alternative promoters driving *spoIVB2* rescues the sporulation phenotype.

Plasmids expressing *spoIVB2* under various promoters were expressed in either wild type, the

C. difficile Δ*spoIVB2*, or the *C. difficile* Δ*sspA* Δ*sspB* strains. Spore yield was determined after

48 hours of incubation by treatment with 30% EtOH and plating on medium containing

germinants. The CFUs were log_{10} transformed. All data represents the average of three

independent experiments. Statistical analysis by ANOVA with Šídák's multiple comparison test.

* P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Figure 10. Model for *spoIVB2* regulation by the *C. difficile* SASPs. In a working model for

how *C. difficile* SASPs regulate sporulation, we hypothesize that upon σ^G activation in the

forespore of wildtype *C. difficile* cells, SspA and SspB become expressed, and they bind to

spoIVB2 to prevent prolonged accumulation of SpoIVB2. In the *C. difficile* Δ*sspA* Δ*sspB* or in

the *C. difficile* Δ*sspB** strains, the SASPs do not accumulate and SpoIVB2 retains prolonged

activity in the forespore compartment. In the *C. difficile* Δ*sspA* Δ*sspB* or in the *C. difficile*

Δ*sspB** suppressor strains, the SpoIVB2^{A20T} or the *spoIVB2*$_{F37F}$ alleles lead to lower amounts of

SpoIVB2 activity (either through changes in protease activity or due to translational changes,

respectively) later in sporulation. Created with BioRender.com.

Acknowledgments

831 This project was supported by awards R01AI116895 and R01AI172043 from the National

832 Institute of Allergy and Infectious Diseases. The content is solely the responsibility of the

833 authors and does not necessarily represent the official views of the NIAID. The funders had no

834 role in study design, data collection and interpretation, or the decision to submit the work for

835 publication.

References

1.   McDonald LC, Gerding DN, Johnson S, Bakken JS, Carroll KC, Coffin SE, et al. Clinical
Practice Guidelines for *Clostridium difficile* Infection in Adults and Children: 2017 Update by the
Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of
America (SHEA). Clin Infect Dis. 2018;66(7):987-94. doi: 10.1093/cid/ciy149. PubMed PMID:
29562266.
2.   Voth DE, Ballard JD. *Clostridium difficile* toxins: mechanism of action and role in
disease. Clin Microbiol Rev. 2005;18(2):247-63. doi: 10.1128/CMR.18.2.247-263.2005. PubMed
PMID: 15831824; PubMed Central PMCID: PMCPMC1082799.
3.   Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. *Clostridium difficile* infection. Nat
Rev Dis Primers. 2016;2:16020. doi: 10.1038/nrdp.2016.20. PubMed PMID: 27158839; PubMed
Central PMCID: PMCPMC5453186.
4.   Jump RL, Pultz MJ, Donskey CJ. Vegetative *Clostridium difficile* survives in room air on
moist surfaces and in gastric contents with reduced acidity: a potential mechanism to explain
the association between proton pump inhibitors and *C. difficile*-associated diarrhea? Antimicrob
Agents Chemother. 2007;51(8):2883-7. doi: 10.1128/AAC.01443-06. PubMed PMID: 17562803;
PubMed Central PMCID: PMCPMC1932506.
5.   Deakin LJ, Clare S, Fagan RP, Dawson LF, Pickard DJ, West MR, et al. The *Clostridium
difficile spo0A* gene is a persistence and transmission factor. Infect Immun. 2012;80(8):2704-11.
Epub 2012/05/23. doi: 10.1128/IAI.00147-12. PubMed PMID: 22615253; PubMed Central
PMCID: PMCPMC3434595.
6.   Seekatz AM, Young VB. *Clostridium difficile* and the microbiota. J Clin Invest.
2014;124(10):4182-9. Epub 20140718. doi: 10.1172/JCI72336. PubMed PMID: 25036699;
PubMed Central PMCID: PMCPMC4191019.
7.   Lawley TD, Clare S, Walker AW, Goulding D, Stabler RA, Croucher N, et al. Antibiotic
treatment of *Clostridium difficile* carrier mice triggers a supershedder state, spore-mediated
transmission, and severe disease in immunocompromised hosts. Infect Immun.
2009;77(9):3661-9. doi: 10.1128/IAI.00558-09. PubMed PMID: 19564382; PubMed Central
PMCID: PMCPMC2737984.
8.   Paredes-Sabja D, Shen A, Sorg JA. *Clostridium difficile* spore biology: sporulation,
germination, and spore structural proteins. Trends Microbiol. 2014;22(7):406-16. doi:

10.1016/j.tim.2014.04.003. PubMed PMID: 24814671; PubMed Central PMCID:
PMCPMC4098856.
9.   Permpoonpattana P, Tolls EH, Nadem R, Tan S, Brisson A, Cutting SM. Surface layers
of *Clostridium difficile* endospores. J Bacteriol. 2011;193(23):6461-70. Epub 2011/09/29. doi:
10.1128/JB.05182-11

JB.05182-11 [pii]. PubMed PMID: 21949071; PubMed Central PMCID: PMC3232898.
10.   Baloh M, Sorg JA. *Clostridioides difficile* SpoVAD and SpoVAE interact and are required
for dipicolinic acid uptake into spores. J Bacteriol. 2021;203(21):e0039421. Epub 2021/08/24.
doi: 10.1128/JB.00394-21. PubMed PMID: 34424035.
11.   Setlow B, Magill N, Febbroriello P, Nakhimovsky L, Koppel DE, Setlow P. Condensation
of the forespore nucleoid early in sporulation of Bacillus species. J Bacteriol.
1991;173(19):6270-8. doi: 10.1128/jb.173.19.6270-6278.1991. PubMed PMID: 1917859;
PubMed Central PMCID: PMCPMC208380.
12.   Setlow P. Spore resistance properties. Microbiol Spectr. 2014;2(5). doi:
10.1128/microbiolspec.TBS-0003-2012. PubMed PMID: 26104355.
13.   Setlow P. I will survive: DNA protection in bacterial spores. Trends Microbiol.
2007;15(4):172-80. doi: 10.1016/j.tim.2007.02.004. PubMed PMID: 17336071.
14.   Burns DA, Heap JT, Minton NP. SleC is essential for germination of *Clostridium difficile*
spores in nutrient-rich medium supplemented with the bile salt taurocholate. J Bacteriol.
2010;192(3):657-64. doi: 10.1128/jb.01209-09.
15.   Gutelius D, Hokeness K, Logan SM, Reid CW. Functional analysis of SleC from
Clostridium difficile: an essential lytic transglycosylase involved in spore germination.
Microbiology. 2014;160(Pt 1):209-16. doi: 10.1099/mic.0.072454-0. PubMed PMID: 24140647;
PubMed Central PMCID: PMCPMC3917228.
16.   Coullon H, Candela T. *Clostridioides difficile* peptidoglycan modifications. Curr Opin
Microbiol. 2022;65:156-61. Epub 20211206. doi: 10.1016/j.mib.2021.11.010. PubMed PMID:
34883390.
17.   Driks A, Eichenberger P. The spore coat microbiol Spectr. 2016;4(2). doi:
10.1128/microbiolspec.TBS.
18.   Driks A. Surface appendages of bacterial spores. Molecular Microbiology.
2007;63(3):623-5.
19.   Permpoonpattana P, Phetcharaburanin J, Mikelsone A, Dembek M, Tan S, Brisson MC,
et al. Functional characterization of *Clostridium difficile* spore coat proteins. J Bacteriol.
2013;195(7):1492-503. Epub 2013/01/22. doi: 10.1128/JB.02104-12. PubMed PMID: 23335421;
PubMed Central PMCID: PMCPMC3624542.
20.   Pizarro-Guajardo M, Calderon-Romero P, Castro-Cordova P, Mora-Uribe P, Paredes-
Sabja D. Ultrastructural variability of the exosporium layer of *Clostridium difficile* spores Appl
Environ Microbiol. 2016;82(7):2202-9. doi: 10.1111/mmi.12611. PubMed PMID: 24720767.
21.   Zeng J, Wang H, Dong M, Tian GB. *Clostridioides difficile* spore: coat assembly and
formation. Emerg Microbes Infect. 2022;11(1):2340-9. doi: 10.1080/22221751.2022.2119168.
PubMed PMID: 36032037; PubMed Central PMCID: PMCPMC9542656.
22.   de Hoon MJ, Eichenberger P, Vitkup D. Hierarchical evolution of the bacterial
sporulation network. Current biology : CB. 2010;20(17):R735-45. Epub 2010/09/14. doi:
10.1016/j.cub.2010.06.031. PubMed PMID: 20833318; PubMed Central PMCID:
PMCPMC2944226.
23.   Talukdar PK, Olguin-Araneda V, Alnoman M, Paredes-Sabja D, Sarker MR. Updates on
the sporulation process in *Clostridium* species. Res Microbiol. 2015;166(4):225-35. doi:
10.1016/j.resmic.2014.12.001. PubMed PMID: 25541348.

24. Paredes-Sabja D, Torres JA, Setlow P, Sarker MR. *Clostridium perfringens* spore
germination: characterization of germinants and their receptors. J Bacteriol. 2008;190(4):1190-
201. doi: 10.1128/jb.01748-07.
25. Lee CD, Rizvi A, Edwards AN, DiCandia MA, Vargas Cuebas GG, Monteiro MP, et al.
Genetic mechanisms governing sporulation initiation in *Clostridioides difficile*. Curr Opin
Microbiol. 2022;66:32-8. Epub 20211218. doi: 10.1016/j.mib.2021.12.001. PubMed PMID:
34933206; PubMed Central PMCID: PMCPMC9064876.
26. DiCandia MA, Edwards AN, Jones JB, Swaim GL, Mills BD, McBride SM. Identification
of Functional Spo0A Residues Critical for Sporulation in *Clostridioides difficile*. J Mol Biol.
2022;434(13):167641. Epub 20220518. doi: 10.1016/j.jmb.2022.167641. PubMed PMID:
35597553; PubMed Central PMCID: PMCPMC9327077.
27. Fimlaid KA, Bond JP, Schutz KC, Putnam EE, Leung JM, Lawley TD, et al. Global
analysis of the sporulation pathway of *Clostridium difficile*. PLoS Genet. 2013;9(8):e1003660.
Epub 2013/08/21. doi: 10.1371/journal.pgen.1003660

PGENETICS-D-12-03167 [pii]. PubMed PMID: 23950727; PubMed Central PMCID:
PMC3738446.
28. Fimlaid KA, Shen A. Diverse mechanisms regulate sporulation sigma factor activity in
the Firmicutes. Curr Opin Microbiol. 2015;24:88-95. doi: 10.1016/j.mib.2015.01.006. PubMed
PMID: 25646759; PubMed Central PMCID: PMCPMC4380625.
29. Setlow P. Small, acid-soluble spore proteins of *Bacillus* species: structure, synthesis,
genetics, function, and degradation. Ann Rev Microbial 1988;42.
30. Mason J, Setlow P. Essential role of small, acid-soluble spore proteins in resistance of
Bacillus subtilis spores to UV light. Journal of Bacteriology 1986;167:174-8.
31. Raju D, Setlow P, Sarker MR. Antisense-RNA-mediated decreased synthesis of small,
acid-soluble spore proteins leads to decreased resistance of *Clostridium perfringens* spores to
moist heat and UV radiation. Appl Environ Microbiol. 2007;73(7):2048-53. Epub 2007/01/30. doi:
10.1128/AEM.02500-06. PubMed PMID: 17259355; PubMed Central PMCID:
PMCPMC1855649.
32. Li J, McClane BA. A novel small acid soluble protein variant is important for spore
resistance of most *Clostridium perfringens* food poisoning isolates. PLoS Pathog.
2008;4(5):e1000056. Epub 20080502. doi: 10.1371/journal.ppat.1000056. PubMed PMID:
18451983; PubMed Central PMCID: PMCPMC2323104.
33. Li J, Paredes-Sabja D, Sarker MR, McClane BA. Further characterization of *Clostridium*
perfringens small acid soluble protein-4 (Ssp4) properties and expression. PLoS One.
2009;4(7):e6249. Epub 20090717. doi: 10.1371/journal.pone.0006249. PubMed PMID:
19609432; PubMed Central PMCID: PMCPMC2706996.
34. Mohr SC, Sokolov NVHA, He C, Setlow P. Binding of small acid-soluble spore proteins
from *Bacillus subtilis* changes the conformation of DNA from B to A. Proc Natl Acad Sci.
1991;88:77-81.
35. Seog Lee K, Bumbaca D, Kosman J, Setlow P, Jedrzejas MJ. Structure of a protein–
DNA complex essential for DNA protection in spores of *Bacillus* species. PNAS. 2007;105(8).
36. Griffith J, Makhov A, Santiago-Larat L, Setlow P. Electron microscopic studies of the
interaction between a *Bacillus subtilis* α/β-type small, acid-soluble spore protein with DNA:
protein binding is cooperative, stiffens the DNA, and induces negative supercoiling. Proc Natl
Acad Sci. 1994;91:8224-8.
37. Setlow B, Sun D, Setlow P. Interaction between DNA and α/β-type small, acid-soluble
spore proteins: a new class of DNA-binding protein. J Bacteriol. 1992.
38. Setlow P. Resistance of spores of *Bacillus* species to ultraviolet light. Environmental and
Molecular Mutagenesis 2001;38:97-104.

39.   Munakata N, Ruper CS. Genetically controlled removal of "spore photoproduct" from
deoxyribonucleic acid of ultraviolet-irradiated *Bacillus subtilis* spores. J Bacteriol.
1972;111(1):192-8.
40.   Yang L, Li L. Spore photoproduct lyase: the known, the controversial, and the unknown.
J Biol Chem. 2015;290(7):4003-9. Epub 2014/12/06. doi: 10.1074/jbc.R114.573675. PubMed
PMID: 25477522; PubMed Central PMCID: PMCPMC4326811.
41.   Sanchez-Salas J-L, Santiago-Lara ML, Sussman MD, Setlow P. Properties of *Bacillus*
megaterium and *Bacillus subtilis* mutants which lack the protease that degrades small, acid-
soluble proteins during spore germination. J Bacteriol. 1992;174(3):804-14.
42.   Nerber HN, Sorg JA. The small acid-soluble proteins of *Clostridioides difficile* are
important for UV resistance and serve as a check point for sporulation. PLoS Pathog.
2021;17(9):e1009516. Epub 20210908. doi: 10.1371/journal.ppat.1009516. PubMed PMID:
34496003; PubMed Central PMCID: PMCPMC8452069.
43.   Leyva-Illades JF, Setlow B, Sarker MR, Setlow P. Effect of a small, acid-soluble spore
protein from *Clostridium perfringens* on the resistance properties of *Bacillus subtilis* spores. J
Bacteriol. 2007;189(21):7927-31. Epub 2007/09/04. doi: 10.1128/JB.01179-07. PubMed PMID:
17766414; PubMed Central PMCID: PMCPMC2168745.
44.   Shrestha R, Cochran AM, Sorg JA. The requirement for co-germinants during
Clostridium difficile spore germination is influenced by mutations in *yabG* and *cspA*. PLoS
Pathog. 2019;15(4):e1007681. doi: 10.1371/journal.ppat.1007681. PubMed PMID: 30943268;
PubMed Central PMCID: PMCPMC6464247.
45.   Francis MB, Allen CA, Shrestha R, Sorg JA. Bile acid recognition by the *Clostridium*
difficile germinant receptor, CspC, is important for establishing infection. PLoS Pathog.
2013;9(5):e1003356. doi: 10.1371/journal.ppat.1003356. PubMed PMID: 23675301; PubMed
Central PMCID: PMCPMC3649964.
46.   Sorg JA, Sonenshein AL. Bile salts and glycine as cogerminants for *Clostridium difficile*
spores. J Bacteriol. 2008;190(7):2505-12. doi: 10.1128/JB.01765-07. PubMed PMID: 18245298;
PubMed Central PMCID: PMCPMC2293200.
47.   Saujet L, Pereira FC, Serrano M, Soutourina O, Monot M, Shelyakin PV, et al. Genome-
wide analysis of cell type-specific gene transcription during spore formation in *Clostridium*
difficile. PLoS Genet. 2013;9(10):e1003756. doi: 10.1371/journal.pgen.1003756. PubMed PMID:
24098137; PubMed Central PMCID: PMCPMC3789822.
48.   Eijlander RT, Holsappel S, de Jong A, Ghosh A, Christie G, Kuipers OP. SpoVT: From
Fine-Tuning Regulator in *Bacillus subtilis* to Essential Sporulation Protein in *Bacillus cereus*.
Front Microbiol. 2016;7:1607. Epub 2016/10/30. doi: 10.3389/fmicb.2016.01607. PubMed PMID:
27790204; PubMed Central PMCID: PMCPMC5061766.
49.   Hoa NT, Brannigan JA, Cutting SM. The *Bacillus subtilis* signaling protein SpoIVB
defines a new family of serine peptidases. J Bacteriol. 2002;184(1):191-9. Epub 2001/12/14.
doi: 10.1128/JB.184.1.191-199.2002. PubMed PMID: 11741860; PubMed Central PMCID:
PMCPMC134772.
50.   Putnam EE, Nock AM, Lawley TD, Shen A. SpoIVA and SipL are *Clostridium difficile*
spore morphogenetic proteins. J Bacteriol. 2013;195(6):1214-25. Epub 2013/01/08. doi:
10.1128/JB.02181-12

JB.02181-12 [pii]. PubMed PMID: 23292781; PubMed Central PMCID: PMC3592010.
51.   Oliveira Paiva AM, Friggen AH, Hossein-Javaheri S, Smits WK. The Signal Sequence of
the Abundant Extracellular Metalloprotease PPEP-1 Can Be Used to Secrete Synthetic
Reporter Proteins in *Clostridium difficile*. ACS Synth Biol. 2016;5(12):1376-82. doi:
10.1021/acssynbio.6b00104. PubMed PMID: 27333161.
52.   Oliveira Paiva AM, Friggen AH, Qin L, Douwes R, Dame RT, Smits WK. The Bacterial
Chromatin Protein HupA Can Remodel DNA and Associates with the Nucleoid in *Clostridium*

difficile. J Mol Biol. 2019;431(4):653-72. Epub 2019/01/12. doi: 10.1016/j.jmb.2019.01.001.
PubMed PMID: 30633871.
53. Lavey NP, Shadid T, Ballard JD, Duerfeldt AS. *Clostridium difficile* ClpP Homologues are
Capable of Uncoupled Activity and Exhibit Different Levels of Susceptibility to Acyldepsipeptide
Modulation. ACS Infect Dis. 2019;5(1):79-89. Epub 20181126. doi:
10.1021/acsinfecdis.8b00199. PubMed PMID: 30411608; PubMed Central PMCID:
PMCPMC6497155.
54. Tovar-Rojo F, Setlow P. Effects of Mutant Small, Acid-Soluble Spore Proteins from
Bacillus subtilis on DNA In Vivo and In Vitro. J Bacteriol. 1991;173(15):4827-35.
55. Alabdali YAJ, Oatley P, Kirk JA, Fagan RP. A cortex-specific penicillin-binding protein
contributes to heat resistance in *Clostridioides difficile* spores. Anaerobe. 2021;70:102379.
Epub 20210430. doi: 10.1016/j.anaerobe.2021.102379. PubMed PMID: 33940167; PubMed
Central PMCID: PMCPMC8417463.
56. Srikhanta YN, Hutton ML, Awad MM, Drinkwater N, Singleton J, Day SL, et al.
Cephamycins inhibit pathogen sporulation and effectively treat recurrent *Clostridioides difficile*
infection. Nat Microbiol. 2019;4(12):2237-45. Epub 20190812. doi: 10.1038/s41564-019-0519-1.
PubMed PMID: 31406331.
57. Popham DL, Bernhards CB. Spore Peptidoglycan. Microbiol Spectr. 2015;3(6). doi:
10.1128/microbiolspec.TBS-0005-2012. PubMed PMID: 27337277.
58. Diaz OR, Sayer CV, Popham DL, Shen A. *Clostridium difficile* lipoprotein GerS is
required for cortex modification and thus spore germination. mSphere. 2018;3(3). doi:
10.1128/mSphere.00205-18. PubMed PMID: 29950380; PubMed Central PMCID:
PMCPMC6021603.
59. Coullon H, Rifflet A, Wheeler R, Janoir C, Boneca IG, Candela T. N-Deacetylases
required for muramic-delta-lactam production are involved in *Clostridium difficile* sporulation,
germination, and heat resistance. J Biol Chem. 2018;293(47):18040-54. Epub 2018/09/30. doi:
10.1074/jbc.RA118.004273. PubMed PMID: 30266804; PubMed Central PMCID:
PMCPMC6254358.
60. Feliciano CA, Eckenroth BE, Diaz OR, Doublié S, Shen A. A lipoprotein allosterically
activates the CwlD amidase during *Clostridioides difficile* spore formation. PLoS Genet.
2021;17(9). doi: 10.1101/2021.06.21.449279.
61. Francis MB, Sorg JA. Dipicolinic acid release by germinating *Clostridium difficile* spores
occurs through a mechanosensing mechanism. mSphere. 2016;1(6). doi:
10.1128/mSphere.00306-16. PubMed PMID: 27981237; PubMed Central PMCID:
PMCPMC5156672.
62. Dong TC, Cutting SM. SpoIVB-mediated cleavage of SpoIVFA could provide the
intercellular signal to activate processing of Pro-s^K in *Bacillus subtilis.* Mol Microbiol. 2003;49 (5
):1425-34.
63. Crawshaw AD, Serrano M, Stanley WA, Henriques AO, Salgado PS. A mother cell-to-
forespore channel: current understanding and future challenges. FEMS Microbiol Lett.
2014;358(2):129-36. doi: 10.1111/1574-6968.12554. PubMed PMID: 25105965.
64. Chiba S, Coleman K, Pogliano K. Impact of membrane fusion and proteolysis on SpoIIQ
dynamics and interaction with SpoIIIAH. J Biol Chem. 2007;282(4):2576-86. Epub 20061122.
doi: 10.1074/jbc.M606056200. PubMed PMID: 17121846; PubMed Central PMCID:
PMCPMC2885159.
65. Oke V, Shchepetov M, Cutting SM. SpoIVB has two distinct functions during spore
formation in *Bacillus subtilis.* Mol Microbiol. 1997;23(2):223-30.
66. Nakamura Y, Gojobori T, Ikemura T. Codon usage tabulated from the international DNA
sequence databases: status for the year 2000. . Nucleic Acids Res. 2000;28(292).
67. Ribis JW, Fimlaid KA, Shen A. Differential requirements for conserved peptidoglycan
remodeling enzymes during *Clostridioides difficile* spore formation. Mol Microbiol.

2018;110(3):370-89. doi: 10.1111/mmi.14090. PubMed PMID: 30066347; PubMed Central
PMCID: PMCPMC6311989.
68.   Martins D, Nerber HN, Roughton CG, Fasquelle A, Barwinska-Sendra A, Vollmer D, et
al. Cleavage of an engulfment peptidoglycan hydrolase by a sporulation signature protease in
Clostridioides difficile. Molecular Microbiology. n/a(n/a). doi: https://doi.org/10.1111/mmi.15291.
69.   Setlow B, McGinnis KA, Ragkousi K, Setlow P. Effects of Major Spore-Specific DNA
Binding Proteins on *Bacillus subtilis* Sporulation and Spore Properties J Bacteriol.
2000;182(24):6906-12.
70.   Sorg JA, Dineen SS. Laboratory maintenance of *Clostridium difficile*. Curr Protoc
Microbiol. 2009;Chapter 9:Unit9A 1. doi: 10.1002/9780471729259.mc09a01s12. PubMed PMID:
19235151.
71.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. Enzymatic
assembly of DNA molecules up to several hundred kilobases. Nat Methods. 2009;6(5):343-5.
doi: 10.1038/nmeth.1318. PubMed PMID: 19363495.
72.   Brehm JN, Sorg JA. Theophylline-based control of *repA* on a *Clostridioides difficile*
plasmid for use in allelic exchange. Anaerobe. 2024:102858. Epub 20240429. doi:
10.1016/j.anaerobe.2024.102858. PubMed PMID: 38692475.
73.   Brehm JN, Sorg JA. Plasmid Sequence and Availability for an Improved *Clostridioides*
difficile CRISPR-Cas9 Mutagenesis System. Microbiol Resour Announc.
2022;11(12):e0083322. Epub 20221107. doi: 10.1128/mra.00833-22. PubMed PMID:
36342279; PubMed Central PMCID: PMCPMC9753633.
74.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub
20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID:
PMCPMC3855844.
75.   Francis MB, Allen CA, Sorg JA. Spore cortex hydrolysis precedes dipicolinic acid release
during *Clostridium difficile* spore germination. J Bacteriol. 2015;197(14):2276-83. doi:
10.1128/JB.02575-14. PubMed PMID: 25917906; PubMed Central PMCID: PMCPMC4524186.
76.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T)
method. Nature protocols. 2008;3(6):1101-8. doi: 10.1038/nprot.2008.73. PubMed PMID:
18546601.

1097

1098

A

CD630Δ*erm* pEV | CD630Δ*erm* Δ*sspA* Δ*sspB* pEV | CD630Δ*erm* Δ*sspA* Δ*sspB* p*sspA,sspB*

B

R20291 Δ*spo0A*
Δ*sspA* Δ*sspB* p*sspA,sspB*
Δ*sspA* Δ*sspB* pEV
Δ*sspB* p*sspB*
Δ*sspB* pEV
Δ*sspA* p*sspA*
Δ*sspA* pEV
CD630Δ*erm* pEV

Spore yield (log$_{10}$)

****
****

C

Δ*sspB* p*sspB*
Δ*sspB* pEV
Δ*sspA* p*sspA*
Δ*sspA* pEV
CD630Δ*erm* pEV

UV Resistance
(% of Wild Type)

****
****

R20291 pEV   0.2 μm

Δ*sspA* pEV   0.5 μm

Δ*sspB* pEV   0.5 μm

Δ*ssp*B* pEV   2 μm

Δ*ssp*B*
p*sspB*   0.2 μm

Δ*sspA* Δ*ssp*B pEV   0.5 μm

Δ*sspA* Δ*ssp*B
p*sspA,sspB*   0.5 μm

Wildtype	ΔsspA ΔsspB	ΔsspA ΔsspB suppressors
σ^F	σ^F	σ^F
spoIVB2	*spoIVB2*	*spoIVB2*$_{F37F \text{ or } A20T}$
Initial Expression	Initial Expression	Initial Expression
σ^G	σ^G	σ^G
sspA / sspB	*sspA / sspB*	*sspA / sspB*
spoIVB2	*spoIVB2*	*spoIVB2*$_{F37F \text{ or } A20T}$
Continued Expression mediated through SspA/B	Decreased SpoIVB2 transcription without SspA/B	Suppressors increase SpoIVB2 abundance to restore sporulation