Increased plasma lipopolysaccharide-binding protein and altered inflammatory mediators in overweight women suggest a state of subclinical endotoxemia

Christine N. Metz[1,2]*, Xiangying Xue[1], Prodyot K Chatterjee[1], Robert P. Adelson[1], Jesse Roth[1,2] Michael Brines[1], Kevin J. Tracey[1,2], Peter K. Gregersen[1,2], Valentin A. Pavlov[1,2]*

[1]The Feinstein Institutes for Medical Research, Northwell Health, Manhasset, NY 11030, USA

[2]Zucker School of Medicine at Hofstra/Northwell-Hofstra University, Hempstead, NY 11550, USA

*Corresponding authors:

Christine Metz, PhD - Email: cmetz@northwell.edu

Valentin Pavlov, PhD – Email: vpavlov@northwell.edu

Abstract

Chronic low-grade inflammation has been recognized as an underlying event linking obesity to cardiovascular disease (CVD). However, inflammatory alterations in individuals who are overweight remain understudied. To provide insight, we determined the levels of key circulating biomarkers of endotoxemia and inflammation, including lipopolysaccharide-binding protein (LBP), CRP, IL-6, leptin, and adiponectin in adult female subjects (n=40) who were lean or overweight and had high cholesterol and/or high blood pressure - two important conventional risk factors for CVD. Plasma levels of LBP were significantly higher in the overweight group compared with the lean group (P=0.005). The levels of CRP were also significantly higher in overweight subjects (P=0.01), as were IL-6 (P=0.02) and leptin (P=0.002), pro-inflammatory mediators associated with cardiovascular risk. Levels of adiponectin, an adipokine with anti-inflammatory and anti-atherogenic functions, were significantly lower in the overweight group (P=0.002). The leptin/adiponectin ratio, a preferential atherogenic marker was significantly increased in women who are overweight (P=0.02). LBP, CRP, leptin, and adiponectin levels significantly correlated with BMI, but not with age and there was a significant correlation between LBP and IL-6 levels. These results reveal the presence of subclinical endotoxemia and a pro-inflammatory state in overweight women and are of interest for further studies with the goal for improved understanding of cardiovascular health risks in women.

Keywords: inflammation; overweight; women; endotoxemia; cytokines; adipokines; cardiovascular disease

Introduction

Obesity and the closely related metabolic syndrome are associated with an increased risk for cardiovascular disease (CVD) and other debilitating and lethal disorders (1-6). Publicly available information on the World Health Organization (WHO) website states that in 2016 around 1.9 billion adults (people over 18 years of age) were overweight, and more than 600 million were obese and the expectations are that more than 2.16 billion people will be overweight and 1.12 billion will be obese by 2030. A major underlying factor driving the pathogenesis in obesity and metabolic syndrome is the presence of a chronic low-grade inflammation, which is characterized by increased circulating IL-6 and other cytokines, as well as altered levels of adipokines, such as leptin and adiponectin (3, 7-11). An important driver of the inflammatory state in obesity is *metabolic endotoxemia*, manifested by increased gut lipopolysaccharide (LPS)-containing microbiota and the consequent compromising of intestinal permeability that leads to increased circulatory LPS levels (12-15). Obesity-associated metabolic endotoxemia and chronic inflammation promote metabolic derangements and are associated with increased cardiovascular risk (9, 15-21).

In addition to obesity, there is evidence that overweight individuals may be at increased risk for CVD and other diseases (22, 23). CVD is the leading cause of death among women in the United States (6, 24). Increased cholesterol levels (hypercholesterolemia) and high blood pressure (hypertension) are important risk factors for CVD (6, 25). Elevated total cholesterol, hypertension, and excessive body weight have been linked to age-dependent increases of coronary heart disease incidence and mortality in both men and women, but to a larger extent in women (26). However, the underlying explanation for sex-specific differences in the CVD pathophysiology remain poorly understood (6). As recently summarized, "Cardiovascular disease in women remains understudied, under-recognized, underdiagnosed, and undertreated globally" (6).

While metabolic endotoxemia and inflammation have been documented in people with obesity and linked to CVD and other diseases, endotoxemia and inflammatory alterations in overweight individuals remain to be characterized. This is of specific interest for improved understanding of women's cardiovascular health. To generate insight, we profiled a panel of plasma biomarkers of endotoxemia and inflammation, previously associated with CVD in obesity, in a cohort of women who were overweight compared to those who were lean. As dyslipidemia and hypertension are recognized leading traditional risk factors for CVD in women (6, 25), we enrolled subjects having high cholesterol and/or high blood pressure in both groups. We observed increased circulating levels of LBP, a marker of metabolic endotoxemia in parallel with elevated CRP, leptin, and IL-6 levels, and decreased adiponectin levels in overweight women.

Materials and Methods

Human subjects and samples

All methods were carried out in accordance with relevant guidelines and regulations. Frozen plasma was obtained from research subjects who participated in the Institutional Review Board (IRB)-approved Genotype and Phenotype (GaP) registry (http://www.gapregistry.org), a research program at the Feinstein Institutes for Medical research, Northwell Health. All research subjects completed an informed consent prior to study participation. The consent permits the use of specimens for future research. The study was approved by the Northwell Health IRB - IRB #09-081A. Participants gave random blood samples and were chosen based on gender, BMI (lean: 18-24.9 kg/m^2 (N=20) vs. overweight: 25-29.9 kg/m^2 (N=20)), age, demographic information, and health/medical information (**Supplementary Table 1**). Subjects in both groups were relatively healthy, except they had self-reported hypertension (systolic $\geq$130mm Hg and diastolic >80 mm Hg) and/or high cholesterol (200 mg/dL) and minor conditions, including acne, eczema, gastroesophageal reflux disease (GERD), drug allergies, and other allergies, as well as

osteoarthritis, osteopenia, osteoporosis. Excluded conditions were Lyme disease, cancer (solid and blood [leukemia, lymphoma, etc.]), anemia, pancreatitis, emphysema, asthma, chronic obstructive pulmonary disease (COPD), inflammatory bowel disease (ulcerative colitis, Crohn's), lupus, rheumatoid arthritis, valvular disease, heart failure, HIV, excess alcohol use, diabetes (types 1 and 2) and Alzheimer's disease and other neurological conditions that would impair the subjects' ability to consent, as well as those using steroids, insulin, metformin, or glyburide and those who smoke or vape.

Plasma sample analyses

All plasma samples were collected from consented GaP participants prior to the COVID-19 pandemic, aliquoted, and stored at -80°C in the Boas Center Biorepository. Just prior to analysis plasma samples were thawed and then assayed for numerous analytes (using dilutions optimized in prior studies) according to the manufacturer's guidelines: adiponectin using the adiponectin/Acrp30 ELISA (DY1065, R&D System, lower limit of detection [LLoD] 15.6pg/ml), C-reactive protein or CRP by ELISA (DY1707, R&D Systems, LLoD 15.6pg/ml); leptin by ELISA (DY398-05, R&D Systems, LLoD 31.2pg/ml); LPS binding protein or LBP by ELISA (DY870-05, R&D Systems, LLoD 0.8ng/ml); and IL-6 using the V-plex MSD platform (K151QXD-2, Meso Scale Discovery, LLoD 0.06pg/ml).

Statistical analysis

Data were analyzed using GraphPad Prism 9.5.1 software and applying an unpaired Student's t test with Welch's correction; $P<0.05$ was considered significant. The linear correlation for the data was analyzed using the Pearson's correlation coefficient (or Pearson's r). The strength of the correlations was assessed based on the r values and considered to be weak (0.2-0.39), moderate (0.40-0.59), or strong (0.6-0.79). Graphical representations were created using GraphPad Prism 9.5.1.

Results

Basic demographics of the study population

The average age and BMI for the lean and overweight cohorts is shown in **Table 1**. There was no statistically significant difference in subjects' age. The average BMI of the subjects in the overweight group was significantly higher compared to the average BMI of the lean group (**Table 1**).

Table 1. Study participants' age and BMI.

Subjects	Lean	Overweight	p value
Age (years)	55.65±9.1	59.7± 6.1	P=0.12
BMI (kg/m^2)	22.4±1.6	27.0±1.5	P<0.0001

Circulating markers of endotoxemia and inflammation are altered in overweight women

LBP is an important mediator of LPS interactions with immune cells and the LPS-induced transcription of pro-inflammatory cytokines (27). Because of the documented difficulties in measuring LPS in biological fluids (28), LBP has been proposed as a useful marker for activation of innate immune responses to microbial products, such as LPS (29-31). Increased levels of circulating LBP have been determined in obesity and the metabolic syndrome and associated with increased circulating IL-6 levels, impaired insulin resistance and cardiovascular risk (29, 31-33). In the present study we observed significantly increased plasma LBP in the overweight group compared with the lean group (**Figure 1A**). In addition, plasma levels of CRP, a general inflammatory marker, were significantly higher in the overweight women (**Figure 1B**), as were the cytokine IL-6 and the adipokine leptin (**Figure 1C, D).** In contrast, the levels of adiponectin were

significantly lower in the overweight group **(Figure 1E).** In addition, the leptin/adiponectin ratio values were significantly increased in the overweight compared with the lean group (**Figure 1F).**

Figure 1. Levels of circulating markers of inflammation are altered in overweight women compared with lean women. Plasma samples of overweight and lean subjects were analyzed for (**A**) LPB, (**B**) CRP, (**C**) IL-6, (**D**) leptin, and (**E**) adiponectin as described in Materials and Methods, and leptin/adiponectin ratios (**F)** were calculated. Data are shown as mean ± SD (*P=0.01 (CRP; *P=0.02 (IL-6 and leptin/adiponectin ratio; **P=0.002 (leptin and adiponectin); **P=0.005 (LBP))

Plasma markers of endotoxemia and inflammation correlate with BMI

Additional data evaluation revealed that plasma inflammatory marker alterations correlated with BMI of the study subjects (**Figure 2).** A moderate, but significant correlation was observed between plasma LBP and BMI (**Figure 2A**). Of note, it appeared that there was a breakpoint in the BMI vs. LBP relationship at the lower "overweight" limit. Similarly, we observed a moderate, but significant correlation between plasma CRP and BMI (**Figure 2B**). There was a weak, non-significant correlation between plasma IL-6 levels and BMI (**Figure 2C**). A moderate and significant correlation was observed between plasma leptin and BMI (**Figure 2D**) and between plasma adiponectin and BMI (**Figure 2E**).

Figure 2. Correlation of plasma inflammatory indices and BMI. Plasma LBP (**A**), CRP (**B**), leptin (**D**), and adiponectin (**E**) levels significantly correlate with BMI as indicated by P values and Pearson correlation coefficients (r). Plasma IL-6 levels do not correlate with BMI (**C**).

Plasma LBP level correlate with plasma IL-6 levels

Inter-correlation analysis of the analytes revealed that LBP was significantly correlated with IL-6 As shown in **Figure 3**. In addition, LBP, and CRP, as well as LBP and leptin correlational analyses generated P values of 0.053 and 0.085 respectively.

Figure 3. Correlation of LBP and IL-6. Plasma LBP levels significantly correlate with IL-6 levels as indicated by the P value and Pearson correlation coefficient.

Of note, no significant correlations were observed between the plasma inflammatory analytes and the age of the study participants as shown in **Supplementary Figure 1.**

Discussion

Here, in women matched for age and having self-declared hypercholesterolemia and/or hypertension, we show that overweight individuals exhibited higher plasma concentrations of LBP and a pro-inflammatory state, indicated by increased levels of IL-6 and other proinflammatory markers and decreased adiponectin levels compared to lean controls. These differences are

relatively small, but notable for the existence of a significant correlation with BMI as well as a positive correlation between LBP and the pro-inflammatory cytokine, IL-6.

Previously, the results of one large study evaluating 500 apparently healthy lean individuals, a majority being female, has shown that LBP is weakly, but highly significantly, correlated with BMI (34). In this study, LBP was also significantly positively correlated with blood pressure, HDL and LDL cholesterol, and triglycerides. Additionally, the results of a much larger (n=2,568 men and women) longitudinal study assessing the relationship between circulating LBP concentrations (the majority of which were within the normal range of less than 15 µg/mL) and cardiovascular risk factors show that the relationship between LBP and CRP, an accepted inflammatory marker, has a highly significant positive correlation (35). The current results conform to these findings and together indicate that markers of endotoxemia and a pro-inflammatory state vary continuously with the BMI, suggesting that adiposity of any degree, even if considered to be within the normal range, is associated with detectable levels of pro-inflammatory molecules. Whether these have clinical significance will require additional large, prospective clinical studies, but it stands to reason that a sustained low grade inflammatory milieu likely has clinical consequences.

Although there is a lack of data regarding the risk of subclinical endotoxemia and chronic inflammation in overweight individuals, previous studies in obese individuals have clearly indicated the presence of metabolic endotoxemia (based on LBP levels) and a chronic inflammatory state and their role in promoting further metabolic dysfunction and pathogenesis (3, 7-9, 36-42). Metabolic endotoxemia in obesity has been specifically linked to the pathogenesis of CVD (15, 21) and increased LBP levels have been directly associated with an increased risk of CVD (21). Endotoxemia increases the production of IL-6 and other cytokines and significantly contributes to a pro-inflammatory state. IL-6 and the adipokine leptin are key mediators of inflammation in obesity (3, 43). IL-6 has been characterized as an important link between obesity

and coronary heart disease (44). Importantly, in a large prospective study, increased IL-6 levels were associated with a higher risk of CVD, specifically coronary heart disease, as strongly as major established risk factors, such as blood pressure and blood cholesterol levels (45) Of note, a significant correlation between plasma LBP and IL-6 levels has been previously documented in a large population spanning a large range of BMI and serum LBP concentrations (35), as well as subjects (males and females) with frank obesity (29). Leptin is an adipokine with an essential role in energy balance through a variety of functions, some of which are related to cardiovascular health (7, 18, 20). Increased leptin levels in obesity are associated with activation of pro-inflammatory signaling and increased thrombosis and arterial distensibility in obese patients (11, 46, 47). Elevated leptin levels arising from leptin resistance in obesity are associated with insulin resistance and CVD (18). In contrast, adiponectin is an adipokine with anti-inflammatory and antithrombotic properties (7, 48). Decreased plasma adiponectin levels were associated with an increased risk of myocardial infarction (49). Plasma levels of CRP (high-sensitive C-reactive protein), a general marker of chronic subclinical inflammation, have been positively correlated with plasma leptin levels and inversely with plasma adiponectin (50-52). The leptin/adiponectin ratio is indicated as a more reliable marker in CVD assessment compared with individual leptin and adiponectin measures (53-55) and proposed as a better marker of a first cardiovascular event in men than plasma leptin and adiponectin levels alone (56).

Until longitudinal data concerning subclinical endotoxemia are available, it may be prudent to institute proactive measures to monitor and reduce the circulating levels of LBP, as a surrogate biomarker for endotoxemia. As composition of the diet has been shown to be a critical driver of metabolic endotoxemia (reviewed in (57)), with high saturated fat ingestion causing postprandial endotoxemia with increases in IL-6 even in lean subjects (58, 59), dietary intervention would be a reasonable initial step. Other possibilities include pharmacological interventions, including the

potential development of anti-LPS peptides which neutralize LPS signaling of immune system activation (57).

In addition to the small number of subjects assessed, there are several limitations of this study. An important one is that the degree of abdominal adiposity (in contrast to subcutaneous fat deposits) has been implicated in the pathogenesis of a systemic inflammatory response and correlates well with the production of pro-inflammatory cytokines (60). Of note, as a single measure BMI cannot fully capture this variable as it is quite insensitive to changes in regional body composition. That is, a smaller waist circumference for any given BMI is indicative of subcutaneous fat deposits, in contrast to an abdominal location in an individual possessing a larger waist circumference. A consensus has appeared that including the waist circumference as a measured variable provides information independent of the BMI, and when both are considered together the predictive accuracy of cardiometabolic risk is significantly increased (61). Additional limitations include other potentially contributing factors that were not assessed, including the existence of abnormal glucose tolerance, HOMA-IR information, lipid profile, degree of sedentary behavior, and the confirmation, severity and pharmacological treatment status of self-reported hypercholesterolemia and hypertension, among others. Lastly, as dietary factors are a major driver of the appearance of LPS into the circulation, in future studies plasma samples should be obtained under standardized fasting conditions.

Conclusion

Our results indicate that despite the presence of hypertension and/or high cholesterol levels in two groups of women characterized using BMI as overweight versus lean, circulating biomarkers of innate immune activation and inflammation, including LBP, CRP, IL-6, and leptin are increased and adiponectin is decreased in the overweight group. Previous studies have focused on evaluating LBP and inflammatory markers in subjects with obesity and alterations of these molecules observed in obese individuals have been linked to an increased cardiometabolic risk.

Our study is among the first to provide insight into the overweight category and more specifically in overweight women. While the number of subjects in this study was small, the presence of subclinical endotoxemia and inflammation, the degree of which varied continuously with BMI encourage performing larger studies to better characterize individuals who are overweight, but not yet classified as obese. These individuals may benefit from therapy to alleviate chronic, low-grade inflammation as an additional risk factor for the development of CVD and other disorders.

Acknowledgements: The authors wish to thank Gila Klein of the Genotype and Phenotype Registry, and the participants who provided blood for this study, as well as Michael Ryan and Bibu Jacob and staff members of the Boas Center Biorepository at the Feinstein Institutes for isolating and storing the plasma samples.

Funding: This work was supported by the National Institutes of Health (NIH), National Institute of General Medical Sciences grants: RO1GM128008 and RO1GM121102 (to VAP) and R35GM118182 (to KJT).

Data availability statement: All data generated or analyzed during this study are included in this published article and its supplementary information files.

Declaration of interests: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author contributions: CNM and VAP proposed the experimental concept and designed the experiments. XX and PKC performed the analyses. CNM, VAP, XX, PKC, and RPA analyzed data. CNM, VAP, and MB wrote and revised the manuscript. JR, PG, and KJT provided additional comments to finalize the paper.

References

1.   Heymsfield SB, Wadden TA. Mechanisms, Pathophysiology, and Management of Obesity. The New England journal of medicine. 2017;376(3):254-66.

2.   Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. The Lancet. 2002;360(9331):473-82.

3.   Pavlov VA. The evolving obesity challenge: targeting the vagus nerve and the inflammatory reflex in the response. Pharmacology & therapeutics. 2021;222:107794.

4.   Grundy SM. Obesity, Metabolic Syndrome, and Cardiovascular Disease. The Journal of Clinical Endocrinology & Metabolism. 2004;89(6):2595-600.

5.   Ritchie SA, Connell JMC. The link between abdominal obesity, metabolic syndrome and cardiovascular disease. Nutrition, Metabolism and Cardiovascular Diseases. 2007;17(4):319-26.

6.   Vogel B, Acevedo M, Appelman Y, Bairey Merz CN, Chieffo A, Figtree GA, et al. The Lancet women and cardiovascular disease Commission: reducing the global burden by 2030. The Lancet. 2021;397(10292):2385-438.

7.   Pavlov VA, Tracey KJ. The vagus nerve and the inflammatory reflex--linking immunity and metabolism. Nature reviews Endocrinology. 2012;8(12):743-54.

8.   Shoelson SE, Herrero L, Naaz A. Obesity, inflammation, and insulin resistance. Gastroenterology. 2007;132(6):2169-80.

9.   Esser N, Legrand-Poels S, Piette J, Scheen AJ, Paquot N. Inflammation as a link between obesity, metabolic syndrome and type 2 diabetes. Diabetes Research and Clinical Practice. 2014;105(2):141-50.

10.   Gregor MF, Hotamisligil GS. Inflammatory mechanisms in obesity. Annual review of immunology. 2011;29:415-45.

11.   Ouchi N, Parker JL, Lugus JJ, Walsh K. Adipokines in inflammation and metabolic disease. Nature reviews Immunology. 2011;11(2):85-97.

12.   Cani PD, Delzenne NM. The role of the gut microbiota in energy metabolism and metabolic disease. Current pharmaceutical design. 2009;15(13):1546-58.

13.   Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, et al. Metabolic endotoxemia initiates obesity and insulin resistance. Diabetes. 2007;56(7):1761-72.

14.   Mohammad S, Thiemermann C. Role of metabolic endotoxemia in systemic inflammation and potential interventions. Frontiers in immunology. 2021;11:594150.

15.   Neves AL, Coelho J, Couto L, Leite-Moreira A, Roncon-Albuquerque R. Metabolic endotoxemia: a molecular link between obesity and cardiovascular risk. 2013;51(2):R51.

16.   de Gusmao Correia ML, Haynes WG. Leptin, obesity and cardiovascular disease. Current Opinion in Nephrology and Hypertension. 2004;13(2):215-23.

17.   Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. The Journal of clinical investigation. 2011;121(6):2111-7.

18.   Martin SS, Qasim A, Reilly MP. Leptin Resistance. Journal of the American College of Cardiology. 2008;52(15):1201-10.

19.   Mathieu P, Lemieux I, Després JP. Obesity, inflammation, and cardiovascular risk. Clinical pharmacology & therapeutics. 2010;87(4):407-16.

20.   Nakamura K, Fuster JJ, Walsh K. Adipokines: a link between obesity and cardiovascular disease. Journal of cardiology. 2014;63(4):250-9.

21.   Roberts LM, Buford TW. Lipopolysaccharide binding protein is associated with CVD risk in older adults. Aging Clin Exp Res. 2021;33(6):1651-8.

22.   Kivimäki M, Kuosma E, Ferrie JE, Luukkonen R, Nyberg ST, Alfredsson L, et al. Overweight, obesity, and risk of cardiometabolic multimorbidity: pooled analysis of individual-level data for 120 813 adults from 16 cohort studies from the USA and Europe. The Lancet Public Health. 2017;2(6):e277-e85.

23.   Wilson PW, D'Agostino RB, Sullivan L, Parise H, Kannel WB. Overweight and obesity as determinants of cardiovascular risk: the Framingham experience. Archives of internal medicine. 2002;162(16):1867-72.

24.   Garcia M, Mulvagh SL, Merz CNB, Buring JE, Manson JE. Cardiovascular Disease in Women. Circulation Research. 2016;118(8):1273-93.

25.   Schenck-Gustafsson K. Risk factors for cardiovascular disease in women. Maturitas. 2009;63(3):186-90.

26.   Jousilahti P, Vartiainen E, Tuomilehto J, Puska P. Sex, Age, Cardiovascular Risk Factors, and Coronary Heart Disease. Circulation. 1999;99(9):1165-72.

27.   Hailman E, Lichenstein HS, Wurfel MM, Miller DS, Johnson DA, Kelley M, et al. Lipopolysaccharide (LPS)-binding protein accelerates the binding of LPS to CD14. The Journal of experimental medicine. 1994;179(1):269-77.

28.   Cohen J. The detection and interpretation of endotoxaemia. Intensive care medicine. 2000;26:S051-S6.

29.   Gonzalez-Quintela A, Alonso M, Campos J, Vizcaino L, Loidi L, Gude F. Determinants of serum concentrations of lipopolysaccharide-binding protein (LBP) in the adult population: the role of obesity. PloS one. 2013;8(1):e54600.

30.   Awoyemi A, Trøseid M, Arnesen H, Solheim S, Seljeflot I. Markers of metabolic endotoxemia as related to metabolic syndrome in an elderly male population at high cardiovascular risk: a cross-sectional study. Diabetol Metab Syndr. 2018;10:59.

31.   Moreno-Navarrete J, Ortega F, Serino M, Luche E, Waget A, Pardo G, et al. Circulating lipopolysaccharide-binding protein (LBP) as a marker of obesity-related insulin resistance. International journal of obesity. 2012;36(11):1442-9.

32.   Trøseid M, Nestvold TK, Rudi K, Thoresen H, Nielsen EW, Lappegård KT. Plasma lipopolysaccharide is closely associated with glycemic control and abdominal obesity: evidence from bariatric surgery. Diabetes care. 2013;36(11):3627-32.

33.   Lepper PM, Schumann C, Triantafilou K, Rasche FM, Schuster T, Frank H, et al. Association of lipopolysaccharide-binding protein and coronary artery disease in men. Journal of the American College of Cardiology. 2007;50(1):25-31.

34.   Sun L, Yu Z, Ye X, Zou S, Li H, Yu D, et al. A marker of endotoxemia is associated with obesity and related metabolic disorders in apparently healthy Chinese. Diabetes Care. 2010;33(9):1925-32.

35.   Asada M, Oishi E, Sakata S, Hata J, Yoshida D, Honda T, et al. Serum Lipopolysaccharide-Binding Protein Levels and the Incidence of Cardiovascular Disease in a General Japanese Population: The Hisayama Study. J Am Heart Assoc. 2019;8(21):e013628.

36.   Nathan C. Epidemic inflammation: pondering obesity. Mol Med. 2008;14(7-8):485-92.

37.   Chang EH, Chavan SS, Pavlov VA. Cholinergic Control of Inflammation, Metabolic Dysfunction, and Cognitive Impairment in Obesity-Associated Disorders: Mechanisms and Novel Therapeutic Opportunities. Frontiers in neuroscience. 2019;13:263.

38.   Saltiel AR, Olefsky JM. Inflammatory mechanisms linking obesity and metabolic disease. The Journal of clinical investigation. 2017;127(1):1-4.

39.   Donath MY, Shoelson SE. Type 2 diabetes as an inflammatory disease. Nature Reviews Immunology. 2011;11(2):98-107.

40.   Tilg H, Moschen AR. Adipocytokines: mediators linking adipose tissue, inflammation and immunity. Nature reviews Immunology. 2006;6(10):772-83.

41.   Esser N, Paquot N, Scheen AJ. Anti-inflammatory agents to treat or prevent type 2 diabetes, metabolic syndrome and cardiovascular disease. Expert opinion on investigational drugs. 2015;24(3):283-307.

42.   Kenchaiah S, Evans JC, Levy D, Wilson PW, Benjamin EJ, Larson MG, et al. Obesity and the risk of heart failure. The New England journal of medicine. 2002;347(5):305-13.

43.   Elks CM, Francis J. Central adiposity, systemic inflammation, and the metabolic syndrome. Current hypertension reports. 2010;12(2):99-104.

44.   Yudkin JS, Kumari M, Humphries SE, Mohamed-Ali V. Inflammation, obesity, stress and coronary heart disease: is interleukin-6 the link? Atherosclerosis. 2000;148(2):209-14.

45.   Danesh J, Kaptoge S, Mann AG, Sarwar N, Wood A, Angleman SB, et al. Long-term interleukin-6 levels and subsequent risk of coronary heart disease: two new prospective studies and a systematic review. PLoS Med. 2008;5(4):e78.

46.   Konstantinides S, Schäfer K, Koschnick S, Loskutoff DJ. Leptin-dependent platelet aggregation and arterial thrombosis suggests a mechanism for atherothrombotic disease in obesity. The Journal of clinical investigation. 2001;108(10):1533-40.

47.   Singhal A, Farooqi IS, Cole TJ, O'Rahilly S, Fewtrell M, Kattenhorn M, et al. Influence of leptin on arterial distensibility: a novel link between obesity and cardiovascular disease? Circulation. 2002;106(15):1919-24.

48.   Katsiki N, Mantzoros C, Mikhailidis DP. Adiponectin, lipids and atherosclerosis. Current opinion in lipidology. 2017;28(4):347-54.

49.   Pischon T, Girman CJ, Hotamisligil GS, Rifai N, Hu FB, Rimm EB. Plasma adiponectin levels and risk of myocardial infarction in men. Jama. 2004;291(14):1730-7.

50.   Shetty GK, Economides PA, Horton ES, Mantzoros CS, Veves A. Circulating adiponectin and resistin levels in relation to metabolic factors, inflammatory markers, and vascular reactivity in diabetic patients and subjects at risk for diabetes. Diabetes care. 2004;27(10):2450-7.

51.   Ble A, Windham BG, Bandinelli S, Taub DD, Volpato S, Bartali B, et al. Relation of plasma leptin to C-reactive protein in older adults (from the Invecchiare nel Chianti study). The American journal of cardiology. 2005;96(7):991-5.

52.   Ouchi N, Kihara S, Funahashi T, Nakamura T, Nishida M, Kumada M, et al. Reciprocal association of C-reactive protein with adiponectin in blood stream and adipose tissue. Circulation. 2003;107(5):671-4.

53.   Satoh N, Naruse M, Usui T, Tagami T, Suganami T, Yamada K, et al. Leptin-to-adiponectin ratio as a potential atherogenic index in obese type 2 diabetic patients. Diabetes care. 2004;27(10):2488-90.

54.   Kotani K, Sakane N, Saiga K, Kurozawa Y. Leptin: adiponectin ratio as an atherosclerotic index in patients with type 2 diabetes: relationship of the index to carotid intima–media thickness. Diabetologia. 2005;48:2684-6.

55.   Norata GD, Raselli S, Grigore L, Garlaschelli K, Dozio E, Magni P, et al. Leptin: adiponectin ratio is an independent predictor of intima media thickness of the common carotid artery. Stroke. 2007;38(10):2844-6.

56.   Kappelle PJWH, Dullaart RPF, van Beek AP, Hillege HL, Wolffenbuttel BHR. The plasma leptin/adiponectin ratio predicts first cardiovascular event in men: A prospective nested case–control study. European Journal of Internal Medicine. 2012;23(8):755-9.

57.   Mohammad S, Thiemermann C. Role of Metabolic Endotoxemia in Systemic Inflammation and Potential Interventions. Front Immunol. 2020;11:594150.

58.   Laugerette F, Vors C, Alligier M, Pineau G, Drai J, Knibbe C, et al. Postprandial Endotoxin Transporters LBP and sCD14 Differ in Obese vs. Overweight and Normal Weight Men during Fat-Rich Meal Digestion. Nutrients. 2020;12(6).

59.   Vors C, Pineau G, Drai J, Meugnier E, Pesenti S, Laville M, et al. Postprandial Endotoxemia Linked With Chylomicrons and Lipopolysaccharides Handling in Obese Versus Lean Men: A Lipid Dose-Effect Trial. J Clin Endocrinol Metab. 2015;100(9):3427-35.

60.   Rexrode KM, Pradhan A, Manson JE, Buring JE, Ridker PM. Relationship of total and abdominal adiposity with CRP and IL-6 in women. Ann Epidemiol. 2003;13(10):674-82.

61.   Ross R, Neeland IJ, Yamashita S, Shai I, Seidell J, Magni P, et al. Waist circumference as a vital sign in clinical practice: a Consensus Statement from the IAS and ICCR Working Group on Visceral Obesity. Nat Rev Endocrinol. 2020;16(3):177-89.

Supplementary Table 1. Demographics of the study subjects.

Subject ID	BMI (kg/m²)	Age (years)	Ethnicity	Condition
Lean Group				
1	19.97	57	Caucasian	HC*
2	20.17	56	Caucasian	Hypertension
3	23.18	60	Caucasian	HC
4	22.31	56	Caucasian	Hypertension, Other
5	22.48	58	Black/West Indian	HC
6	19.53	63	Caucasian	HC, Osteoporosis
7	22.27	60	Caucasian	HC, Other
8	21.63	58	Caucasian	HC
9	22.50	61	Caucasian	HC, Osteoarthritis
10	23.49	62	Caucasian	HC, Osteopenia
11	23.96	63	Black/African American	HC, Osteoarthritis
12	24.56	47	Caucasian	HC
13	22.14	30	Caucasian	HC
14	23.18	46	Caucasian/Latino/Hispanic	HC, Other
15	22.59	58	Caucasian	Hypertension, Other
16	23.41	60	Asian	HC, Hypertension, Other
17	19.48	62	Caucasian	HC
18	23.24	61	American Indian/Alaska Native	HC
19	24.80	34	Caucasian	HC
20	23.04	61	Caucasian	HC, Other
Overweight group				
21	28.34	68	Caucasian	HC, Hypertension
22	29.12	67	Caucasian	HC, Hypertension
23	28.83	53	Black/West Indian	Hypertension
24	25.84	52	Caucasian	Hypertension
25	26.54	66	Caucasian	HC, Hypertension, Other
26	29.86	56	Caucasian	Hypertension
27	25.32	57	Caucasian	HC, Hypertension
28	29.45	61	Caucasian	HC, Hypertension
29	25.82	66	Caucasian	HC, Osteoporosis
30	28.19	45	Caucasian	HC, Other
31	27.43	58	Caucasian	HC
32	25.88	66	Caucasian	HC, Osteoarthritis
33	25.84	68	Caucasian	HC, Hypertension, Osteoarthritis, Other
34	25.84	59	Black/African American	HC
35	26.15	64	Caucasian	HC
36	25.79	61	Caucasian	Hypertension

37	27.43	56	Asian	HC
38	25.01	53	Caucasian	Hypertension
39	25.85	61	Caucasian	HC, Hypertension
40	27.37	57	Caucasian	HC, Hypertension

*HC (High cholesterol levels)

Supplementary Figure 1. Correlation of plasma inflammatory indices and age. Plasma LBP (**A**), CRP (**B**), IL-6 (**C**), leptin (**D**), and adiponectin (**E**) levels do not significantly correlate with age as indicated by Pearson correlation coefficients (r) and p values.