The Extrafoveal Preview Effect is More Pronounced

at Polar Angle Locations Where Perception is Poor

Xiaoyi Liu[1,5], David Melcher[1,2], Marisa Carrasco[3], and Nina M. Hanning[3,4]

[1]New York University Abu Dhabi, Division of Science, Psychology Program

[2]New York University Abu Dhabi, Center for Brain and Health, NYUAD Research Institute

[3]New York University, Department of Psychology and Center for Neural Science

[4]Humboldt-Universität zu Berlin, Institut für Psychologie

[5]Princeton University, Department of Psychology

Author Note

Xiaoyi Liu https://orcid.org/0000-0001-5129-0683

Correspondence concerning this article should be addressed to Xiaoyi Liu. E-mail: xiaoyi.x.liu@gmail.com

Abstract

The pre-saccadic preview of a peripheral target enhances the efficiency of its post-saccadic processing, termed the extrafoveal preview effect. Peripheral visual performance –and thus the quality of the preview– varies around the visual field, even at iso-eccentric locations. To investigate whether these polar angle asymmetries influence the preview effect, we asked human participants (N=14) to preview four tilted Gabors at the cardinals, until a central cue indicated to which one to saccade. During the saccade, the target orientation either remained or was flipped (valid/invalid preview). After saccade landing, participants discriminated the orientation of the (briefly presented) second Gabor. Gabor contrast was titrated with adaptive staircases. Valid previews increased participants' post-saccadic contrast sensitivity. This preview effect was inversely related to polar angle perceptual asymmetries; largest at the upper, and smallest at the horizontal meridian. Our finding reveals that the visual system compensates for peripheral asymmetries when integrating information across saccades.

Keywords: contrast sensitivity, eye movement programming, peripheral vision, trans-saccadic perception, visual field asymmetries, visual stability

Statement of Relevance

Even though visual sensitivity degrades with farther distance from the fovea, we make use of peripheral information to monitor or preview the surroundings, for instance when driving (pedestrians at eye level, instrument panel at the lower visual field, farther things in the upper visual field). When we make saccadic eye movements to foveate relevant items, the peripheral information "previewed" before the saccade supports post-saccadic vision. Because our vision differs around the visual field –at the same eccentricity it is best along the horizontal meridian and worst at the upper vertical meridian– investigating whether peripheral information at different polar angles equally facilitates post-saccadic perception has implications for everyday life. Our study reveals that peripheral preview exerts a larger influence on subsequent foveal processing at locations where vision is worse. This finding suggests that the visual system compensates for differences in peripheral vision when integrating information across eye movements.

The Extrafoveal Preview Effect is More Pronounced Where Perception is Poor

When exploring their surroundings, humans typically make fast saccadic eye

movements to bring different parts of the scene onto the fovea, the central region of the

retina with the highest visual acuity. Despite its low resolution, peripheral visual information

plays a critical role in preparing and guiding saccades, a process termed *preview* (Melcher &

Colby, 2008; Rayner et al., 1980). After a saccade, inputs previewed before the saccade are

integrated with the post-saccadic percepts, to facilitate subsequent foveal processing

(review: Huber-Huber et al., 2021). This *extrafoveal preview effect* occurs with a variety of

stimuli, including low-level features (Cont & Zimmermann, 2021; Zimmermann et al., 2016;),

words (Schotter et al., 2012), and objects (Buonocore et al., 2020; Huber-Huber et al., 2019;

Liu et al., 2022). It is reflected both at the behavioral level –e.g., higher accuracy and faster

reaction times (e.g., Huber-Huber et al., 2019)– and at the neural level –e.g., faster

decoding time (Edwards et al., 2018).

Preview effect studies have focused on the horizontal meridian (HM), but peripheral

vision varies greatly with polar angle: it is better along the horizontal than vertical meridian

(HM>VM: horizontal-vertical anisotropy; HVA) and along the lower than upper vertical

meridian (LVM>UVM: vertical meridian asymmetry; VMA) (Abrams et al., 2012; Carrasco et

al., 2001; Greenwood et al., 2017). Visual sensitivity can be two to three times higher for

stimuli on the HM than VM (Himmelberg et al., 2020; Jigo et al., 2023). These polar angle

asymmetries are present in many visual tasks, such as contrast sensitivity (Baldwin et al.,

2012; Carrasco et al., 2001; Himmelberg et al., 2020; Rijskijk et al., 1980), visual acuity

(Barbot et al., 2021; Kwak et al., 2023), spatial resolution (Montaser-Kouhsari & Carrasco,

2009; Talgar & Carrasco, 2002), and crowding (Greenwood et al., 2017; Kurzawski et al.,

2023). (For a review on polar angle asymmetries, see Himmelberg et al., 2023). However, it

is unknown whether and how our heterogeneous peripheral vision affects the integration of

pre- and post-saccadic information for different saccade directions.

Here we consider two opposite hypotheses. According to the optimal integration

framework (Ernst & Banks, 2002; Ernst & Bülthoff, 2004; Körding & Wolpert, 2004), trans-

saccadic integration is similar to other types of sensory integration across modalities, during

which the brain integrates information by weighing the sensory evidence based on relative

precision. Estimated reliability is maximized when stronger sensory signals (e.g., higher

signal-to-noise ratio, reduced uncertainty) receive higher weights (Ernst & Bülthoff, 2004).

Correspondingly, preview benefits in reading are reduced or extinguished when the preview

word is located at a farther eccentricity, e.g., two versus one word from the current fixation

(Angele & Rayner, 2011; McDonald, 2005). More evidence comes from gaze-contingent

tasks that present stimuli according to the observers' gaze position and systematically vary

the relative visibility of peripheral and foveated targets by changing either the stimulus

contrast (Wolf & Schütz, 2015) or eccentricity (Liu et al., 2022), or by adding noise (Hutzler

et al., 2019; Liu et al., 2022). These studies have revealed that with incongruent pre- and

post-saccadic percepts, trans-saccadic perceptual judgements are increasingly biased

towards the peripheral preview as its visibility increases. Because object visibility also varies

with polar angle, one could expect the magnitude of the preview effect around the visual field

to follow the same pattern: a larger preview effect on the HM, where peripheral vision is best,

followed by the lower- and then the upper- vertical meridian.

In contrast, the visual system might compensate for polar angle asymmetries in

peripheral vision when integrating information across saccades, enhancing post-saccadic

vision more at locations where it is needed the most. Endogenous attention, the voluntary

allocation of processing resources based on one's goals (Carrasco, 2011), is an example of

this process. Depending on the current task demand, endogenous attention can exert

different perceptual effects (e.g., Barbot & Carrasco, 2017; Sharp et al., 2018). For example,

it either increases or decreases the temporal integration window when observers are

instructed to segregate or integrate two closely presented stimuli. This effect is more

pronounced when baseline performance is worse (Sharp et al., 2018). A similar

compensation mechanism during trans-saccadic integration would give higher weights to the

pre-saccadic preview signals at locations where peripheral vision is worse, leading to equal

preview effects around the visual field, or even a reversed pattern of preview asymmetries.

However, polar angle asymmetries in peripheral vision are resilient; they are not

alleviated by covert temporal (Fernández et al., 2019) or spatial attention: Both exogenous

attention –which is involuntary, fast and transient (Carrasco et al., 2001; Roberts et al.,

2018)– and endogenous attention –which is voluntary, slower and sustained (Purokayastha

et al., 2021)– similarly improve performance around the visual field, thereby preserving the

polar angle asymmetries. Pre-saccadic attention –which is automatically deployed to the

target of an upcoming eye movement during saccade preparation (Deubel & Schneider,

1996; Deubel, 2008; Hanning et al., 2019; Kowler et al., 1995; Rolfs & Carrasco, 2012) –

enhances perception similarly around the visual field (Kwak et al., 2023) or even exacerbates

polar angle asymmetries, by benefitting performance less where it is already worse (Hanning

et al., 2022; in press). Thus, neither covert nor pre-saccadic attention overcome polar angle

asymmetries in peripheral vision.

Pre-saccadic attention contributes to trans-saccadic integration, but the latter entails

more processes (Huber-Huber et al., 2019; Huber-Huber et al., 2021) and relies on a

prolonged interaction among the oculomotor network and the visual system before and after

saccades (Caruso et al., 2018; Schall & Hanes, 1993). Therefore, to investigate whether and

how the extrafoveal preview effect, a measure for trans-saccadic integration, interacts with

polar angle asymmetries in peripheral vision, we first assessed participants' baseline contrast

sensitivity at four isoeccentric cardinal locations during fixation, and then measured their preview

effect at the same locations with an integration task, as the normalized difference in post-saccadic

contrast sensitivity following valid vs. invalid previews.

Our findings are consistent with the compensation hypothesis. Unlike previous reports

that polar angle asymmetries are preserved or even exacerbated by covert and pre-saccadic

attention (e.g., Carrasco et al., 2001; Hanning et al., 2022, in press; Kwak et al., 2023;

Purokayastha et al., 2021), we found the largest preview effect at the UVM, followed by the

LVM, and lastly the HM –inversely correlated with peripheral polar angle asymmetries during

fixation.

Open Practices Statement

85   Behavioral and eye tracking data as well as analysis scripts will be publicly accessible

86   upon manuscript acceptance on OSF (https://osf.io/fsx7v/). The study was not preregistered.

Method

Participants

We recruited 15 participants at NYU-NY. All participants gave informed consent before

the experiment, received allowance, and had normal or corrected-to-normal vision. All but two

participants (authors XL and NH) were naïve to the purpose of the experiment. One participant

was excluded after data collection for having a threshold estimate three SD away from the

group average in the valid preview condition (see Procedure). The remaining 14 participants
(9 females; all right-handed) had an average age of 28.14 (± 4.74) (M ± SD) years. The
experiment was approved by the NYU Abu Dhabi ethical committee and all experimental
procedures were in agreement with the Declaration of Helsinki.

Apparatus

The experiment was coded with MATLAB PsychToolbox-3 (Brainard 1997), Eyelink
toolbox (Cornelissen et al., 2002), as well as the Palamedes toolbox (Prins & Kingdom, 2018)
for the PEST staircase procedure (Pentland, 1980), and was run on an Apple iMac Intel Core
2 Duo computer (Cupertino, CA, USA). Stimuli were displayed on a gamma-linearized 20-
inch ViewSonic G220fb CRT screen (Brea, CA, USA) (1280×960 screen resolution; 100 Hz
refresh rate) with a gray background (luminance: 128 cd/m^2). Participants sat at a viewing
distance of 57 cm with their head stabilized on a chin-forehead rest. Eye position of the
dominant eye was recorded by an EyeLink 1000 Plus eye tracker (SR Research, Ontario,
Canada) with the desktop mount, at a sampling rate of 1000 Hz.

Procedure

The main study included three experimental tasks performed in separate experimental
sessions with overall four *viewing conditions*: Participants first completed a baseline task, in
which peripheral stimuli were viewed during *fixation* (**Fig. 1a**)**,** followed by an integration task
(with viewing conditions *valid* and *invalid preview* intermixed) and a *postview task*. In all tasks,
participants judged the orientation of a Gabor patch (clockwise/counterclockwise relative to
vertical, 2-alternative forced choice task). To measure contrast sensitivity, stimulus contrast
per experimental condition (4 locations × 4 viewing conditions: fixation, valid preview, invalid
preview, postview) was separately adjusted by an adaptive staircase procedure (PEST; see
Supplemental Material) based on the correctness of participants' tilt direction responses.

In all tasks, a white central fixation dot (0.3 dva diameter; dva: degree of visual angle)
and four black placeholders centered at four isoeccentric locations (8 dva) along the upper,
lower, left, and right meridian of the fixation dot were presented throughout the trial. Each
placeholder consisted of four corners (0.3 dva in length, 0.1 dva in width), indicating the
possible locations of upcoming stimuli and potential saccade targets. Each trial started with
participants fixating on the central fixation dot. Once we detected continuous fixation (within
a radius of 1.5 dva) for 700 ms, the trial continued depending on the specific task.

Baseline fixation task. After the initial fixation, four Gabor patches (spatial
frequency = 4 cpd; diameter = 2.5 dva; σ = 0.625 dva) were presented at the center of the
placeholders for 100 ms while participants kept fixating on the central fixation dot. Three
Gabors were vertically oriented distractors, one Gabor was a target tilted ±15° relative to
vertical. The location of the target was randomly selected and unbeknownst to the
participants until its presentation. Four hundred ms after Gabor offset, the fixation dot color
changed to grey and the placeholder at the target location turned white, prompting
participants to indicate the tilt direction of the target by pressing a button on a standard
keyboard, without time pressure. Audio feedback was provided for 200 ms immediately
following the response.

Integration task. After the initial fixation, participants (while still fixating at the central
fixation dot) peripherally viewed four Gabors, which were presented within the placeholders
and independently tilted slightly clockwise or counterclockwise relative to vertical (**Fig. 1b**).
The tilt angles of the Gabors were determined by a thresholding procedure beforehand (see
Supplemental Material). After 400 ms of preview, a saccade cue (red line of 0.45 dva in
length, 0.05 dva in width) superimposed on the fixation dot, indicated to which Gabor
participants had to make an immediate saccade. We monitored gaze positions online. As
soon as saccade onset was detected (i.e., when two consecutive gaze samples ~1 ms apart
deviated more than 0.18 dva), a visual mask was displayed at each location for 10 ms. This
mask (bandpass filtered random noise, same contrast and size as the Gabor patch) served
to equalize the visual input changes in the valid and invalid conditions (**Fig. 1c**). Note that
participants reported to be unaware of the mask due to saccadic suppression (Vallines &
Greenlee, 2006). The offline saccade detection algorithm used to more accurately detect
saccadic eye movements (see Data Analysis section) shows an average interval between
actual saccade onset and mask onset of 15.23 (± 5.52) ms (**Fig. 1d).**

Immediately after the mask, the target stimulus –either identical to the preview
stimulus (*valid* condition) or with a flipped tilt angle (e.g., from +1° to -1°) (*invalid* condition;
Fig. 1d) was presented at the saccade target. The target stimulus disappeared 80 ms after
we detected participants' gaze within 2 dva from saccade target center (i.e., the online
approximation of a saccade offset). On 98.52% trials, the saccade landed after target
stimulus onset, with an average target-saccade interval of 21.42 (± 11.22) ms **(Fig. 1e).** For
the remaining <1.5% of trials, target stimulus onset occurred within 20 ms after saccade
offset, i.e., within the time range of saccadic suppression (Vallines & Greenlee, 2006). After

target stimulus disappearance, participants indicated the tilt orientation of the target stimulus.

They were explicitly instructed to base their judgement on the post-saccadic target stimulus,

which they foveated after the saccade. To prevent participants from impulsively responding

to the preview, participants completed at least 36 practice trials with prolonged post-saccadic

target duration (300 ms) followed by another 36 practice trials with 80 ms post-saccadic

target duration before the integration task. We would only start the main task after participants

reached 80% correct orientation discrimination performance in the invalid condition of the

first practice.

Fig 1. *Illustration of the task.* **a.** *Fixation task. Each trial started with a central fixation dot, followed by the stimuli. Participants indicated the tilt direction of a tilted Gabor (±15° relative to vertical) presented among distractors.* **b.** *Integration task. After fixation, participants previewed four slightly tilted Gabors, followed by a saccade cue for an immediate eye movement. Saccade onset, detected by an online algorithm (see Methods), triggered a 10ms mask, followed by the target. The target disappeared ~80ms after saccade offset. Participants then indicated the tilt direction of the second, foveated target. The tilt angles of Gabor stimuli were determined by a thresholding task (see Supplemental Material)* **c.** *Task conditions in the integration task.* **d.** *Temporal delay between offline detected saccade onset and mask onset.* **e.** *Temporal delay between target onset and offline detected saccade offset.*

Postview task. To measure performance in a neutral, no-preview condition, in this task

no Gabors were presented for preview. Participants merely fixated on the central fixation until

they saw a saccade cue (the same overall fixation duration as in the integration task). Once

a saccade was detected, we presented bandpass filtered noise at the target location for 10

ms, followed by a target Gabor at the saccade target. Again, the target Gabor disappeared

80 ms after we detected participants' gaze within 2 dva from saccade target center. The rest

of the task was identical to the integration task.

The study comprised 3 experimental sessions completed over 3-4 days; each lasted

90 minutes. In the first session, participants completed the *fixation task* followed by the first

part of the *integration task*. The second session consisted of the *postview task* and the

second part of the *integration task* (order counterbalanced among participants). In the third

session, participants completed the last part of the *integration task*. The first session of the

integration task was preceded by a thresholding procedure (see Supplemental Material) to

determine the tilt angle of the Gabors used in the integration and *postview tasks*. In total,

participants performed 2592 trials: 432 trials in both the *fixation* and *postview tasks*, and 1728

trials in the *integration task*. Trials were aborted and repeated at the end of the session if

participants 1) did not maintain central fixation during the preview, 2) made a saccade too

early (< 100 ms) or too late (> 550 ms) after cue onset, or 3) looked toward a wrong stimulus

in an eye-movement task.

Data analysis

Contrast sensitivity and saccade parameters were analyzed with R and MATLAB.

Contrast thresholds and staircase exclusion were determined with a similar method as in

Hanning et al. (2022a; see Supplemental Material). On average, 12% of staircases were

excluded for each participant. Our dependent variable contrast sensitivity was computed as

the reciprocal of the average threshold estimate. The difference in contrast sensitivity

between the valid and invalid preview conditions divided by their average (i.e., the normalized

difference) was then used as an index for the preview effect. We analyzed the recorded gaze

samples with an offline algorithm that detects saccades based on gaze velocity, amplitude,

and duration (Engbert & Mergenthaler, 2006; see Supplemental Material).

Unless otherwise noted, contrast sensitivity and the preview effect were analyzed with

repeated-measures analyses of variance (ANOVA) in R (package *ez*, version 4.4.0), with

meridian (horizontal, vertical) and location (left and right, or upper and lower) as within-

subject factors. Eye movement data –saccade latency, amplitude, precision (i.e., the

Euclidean distance between the target and saccade endpoint), and post-saccadic eye

movements– were analyzed with 4 (saccade direction: left, right, upper, lower) × 3 (viewing

condition: *valid preview*, *invalid preview*, *postview*; note the *fixation* condition is not

considered here, since no eye movements were executed) repeated-measures ANOVAs. For

all repeated-measures ANOVAs in which the sphericity assumption was not met, we report

Greenhouse-Geisser corrected *p*-values. Post-hoc tests for any significant effects were run

with the package *emmeans* (version 1.7.4-1 in R). Multiple comparisons were corrected by

controlling the false discovery rate. For effect size, we report η_p^2 for ANOVAs, along with

95% confidence interval (CI) for variance explained by each effect (CI_η). For post-hoc tests

and the correlation analysis, we report the 95% CIs for difference and correlation coefficient.

We also provide Bayes factors (package *BayesFactor,* version 0.9.12-4.4) for the main

findings (i.e., peripheral polar angle asymmetries and preview asymmetries) as a measure

of graded evidence.

Power analysis

We determined the sample size based on previous studies on polar angle asymmetries

(e.g., Anton-Erxleben et al., 2012; Barbot et al., 2021; Hanning et al., 2022; in press) or on the

preview effect (e.g., Buonocor et al., 2020; De Pisapia et al., 2010; Liu et al., 2022). A post-hoc

power analysis confirmed that our sample size (n = 14) is associated with 100% power to detect

an existing HVA and VMA, and 98% power to detect the preview effect (Cohen's *d* = 1.18, α =

.005), and thus adequate to provide reliable results.

24   **Results**

Baseline sensitivity during fixation varies with polar angle

To establish the existence of the typical polar angle asymmetries (HVA and VMA), we

analyzed contrast sensitivity using a 2 (meridian: horizontal, vertical) × 2 (location: left/lower,

right/upper) repeated-measures ANOVA. We found a main effect of meridian (**Fig. 2a**), $F(1,$

$13) = 11.56$, $p = .0047$, BF > 100, $\eta_p^2 = .47$, 95%, $CI_\eta = [.063, .68]$. Consistent with the HVA,

sensitivity along the HM (33.13 ± 4.08) (M ± SE) was significantly higher than that along

the VM (21.73 ± 2.39). There was a significant interaction between meridian and location,

$F(1, 13) = 4.97$, $p = .044$, BF > 100, $\eta_p^2 = .28$, 95% $CI_\eta = [.00, .56]$. Post-hoc contrasts

showed no difference between the left (32.47 ± 4.09) and right (33.79 ± 4.4) HM (BF = .31),

whereas sensitivity at the lower VM (24.96 ± 3.1) was significantly higher than the upper

VM (18.49 ± 2.19), $t(25.9) = 2.70$, $p = .012$, BF = 3.19, 95% CI = [1.54, 11.40], consistent

with the VMA.

To visualize the individual differences in HVA and VMA, we plotted each participant's

contrast sensitivity along the HM against the VM (**Fig. 2b**), and along the LVM against the

UVM (**Fig. 2c**). Diagonal lines indicate equivalent performance. HVA and VMA were found

in all participants, except in two for HVA and four for VMA. The strength of the asymmetries

was computed as the sensitivity difference between the two respective meridians divided by

their average, with a positive number representing a typical HVA/VMA. The correlation

between HVA and VMA was not significant ($r = .20$, $p = .50$), suggesting (at least partially)

different origins (Himmelberg et al., 2020; Barbot et al., 2021).

The preview effect is inversely correlated with peripheral sensitivity

To test whether the preview effect, i.e., the difference in contrast sensitivity

following valid and invalid previews, was consistent at all four locations, we performed a

repeated-measures ANOVA with preview validity (valid, invalid) and target location (left, right,

upper, lower) as within-subjects factors. We found a main effect of preview validity, $F(1, 13)$

$= 44.00$, $p < .004$, $\eta_p^2 = .77$, 95% $CI_\eta = [.43, .86]$. Planned contrasts showed significant

preview effects at all four locations (all $p < .001$) (**Fig. 2d;** see **Fig. S1** for sensitivity in the

valid and invalid conditions separately).

Fig. 2. *Behavioral results in the fixation and integration tasks. **a.d**. Contrast sensitivity in the fixation task computed as the reciprocal of the staircase threshold estimate (**a**) or preview effect in the integration task computed as the normalized difference in sensitivity between the valid and invalid conditions (**d**) as a function of target location. Boxes depict the 95% CIs around the mean, represented by the black lines. Horizontal brackets indicate comparisons between the horizontal vs. vertical meridian (HVA) and upper vs. lower vertical meridian (VMA); error bars on the brackets represent 95% CIs of the difference between compared conditions. * p < .05, ** p < .01 (after correcting for multiple comparisons). Polar plots depict the same data. **b.e**. Individual participants' contrast sensitivity (**b**) or preview effect (**e**) along the horizontal meridian plotted against the vertical meridian. Red error bars represent 95% Cis of group averages. **c.f**. Individual participants' contrast sensitivity (**c**) or preview effect (**f**) along the lower against the upper vertical meridian.*

Next, we quantified the magnitude of the preview effect as the normalized difference

between the valid and invalid conditions, which we then analyzed with a 2 (meridian:

horizontal, vertical) × 2 (location: left/lower, right/upper) repeated-measures ANOVA. We

found a significant main effect of meridian, $F(1, 13) = 5.88$, $p = .031$, BF = 6.28, $\eta_p^2 =$

.31, 95% CI$_\eta$ = [.00, .58]. Interestingly, post-hoc contrasts revealed that the preview

asymmetry was opposite to the polar angle asymmetry during fixation, with a larger preview

effect at the VM (1.07 ± 0.16) than the HM (0.65 ± 0.12), $t(13) = -2.43$, $p = .031$, BF = 3.69,

95% CI = [-.79, -.045] (**. 2d**). The reversed HVA in the preview effect was also evident at

the individual level (**Fig. 2e**). Although the interaction between meridian and location was not

significant ($p = .27$), the corresponding BF = 1.36 provides anecdotal evidence supporting the

interaction. On average, the preview effect was slightly larger at the UVM (1.22 ± 0.16) than

the LVM (0.92 ± 0.17), which was consistent in all but two participants (**Fig. 2f**).

To further evaluate the relation between the peripheral contrast sensitivity during

fixation and the magnitude of the preview effect at the individual level, we ran a repeated-

measures correlation (Bakdash & Marusich, 2017), which accounts for the violation of

independence in data due to repeated within-participant data points. This way we measured

the variability across target locations by removing the inter-individual variability. Aligned with

the reversed pattern of asymmetries indicated by the ANOVA, the repeated-measures

correlation showed a significant negative intra-individual correlation between peripheral

contrast sensitivity and the preview effect, $r_m = -.37$, $p = .014$, BF = 11.59, 95% CI = [-0.61, -

.074] (**Fig. 3**), i.e., within individuals, the preview effect was larger where peripheral sensitivity

was smaller (e.g., UVM).

Fig. 3. *Individual participants' preview effect as a function of peripheral contrast sensitivity. Data centered to each participants' average across target locations. Dots represent individual data at the four cardinal locations. The regression line represents the intra-individual (repeated-measures) correlation.*

Lowest post-saccadic sensitivity at the UVM

We analyzed post-saccadic foveal contrast sensitivity (measured in the *postview task*)
with a 2 (meridian: horizontal, vertical) × 2 (location: left/lower, right/upper) repeated-measures
ANOVA. We expected to observe equivalent performance at the four cardinal locations,
because in this task, participants made perceptual judgements about a brief, foveated
target (i.e., polar angle asymmetries should not affect perception). Surprisingly, we found
a main effect of location, $F(1, 13) = 5.46$, $p = .036$, $\eta_p^2 = .30$, 95% $CI_\eta = [.00, .57]$, and an
interaction between meridian and location, $F(1, 13) = 10.08$, $p = .007$, $\eta_p^2 = .44$, 95% CI_η
$= [.044, .66]$. The effects were driven by a lower post-saccadic contrast sensitivity after
upward saccades (15.00 ± 3.16) than after rightward (29.49 ± 4.76; $p = .015$, 95% CI =
[0.91, 28.05]), leftward (26.23 ± 3.64; $p = .054$, 95% CI = [-2.34, 24.80]; marginally
significant) or downward saccades (30.85 ± 2.77; $p = .014$, 95% CI = [2.28, 29.42]) (**Fig.**
4). Other post-hoc contrasts did not reach significance.

*Fig. 4. Contrast sensitivity in the postview task as a function of target location. **a.** Boxes depicts the 95% CIs around the mean, represented by black lines. Horizontal brackets indicate comparisons between the respective target location, with the error bars on the brackets representing 95% CIs of the difference between the compared conditions. * p < .05 (after correcting for multiple comparisons). b. Polar plots showing the same data.*

We further ran a linear regression model (M1) with peripheral contrast sensitivity
and post-saccadic foveal sensitivity as predictors for the preview effect. Consistent with
the repeated-measures correlation, peripheral sensitivity predicted the preview effect in
a negative way ($p = .021$). Post-saccadic sensitivity, however, was not associated with
the preview effect. We then ran a second regression model with peripheral sensitivity as
the sole predictor (M2). By comparing M1 and M2 with a likelihood ratio test, we found
that peripheral and post-saccadic sensitivity (M1) did not explain more variance in the

preview effect than peripheral sensitivity alone (M2; p = .37).

Saccade parameters cannot explain the preview effect asymmetries

To evaluate whether differences in saccade parameters could explain the observed

preview asymmetries around the visual field, we analyzed the offline-computed saccade

parameters (see Data Analysis section).

Longer saccade latencies (and thus a longer preview duration) as well as less precise

saccades or higer post-saccadic movement rates (and thus a lower quality postview) could lead

participants to rely more on the preview information, thereby giving rise to a larger preview effect.

We found main effects of saccade direction, $F(3, 39)$ = 32.14, p < .001, η_p^2 = .71, 95% CI$_\eta$

= [.51, .79] and viewing condition, $F(2, 26)$ = 15.94, p < .001, η_p^2 = .55, 95% CI$_\eta$ = [.24,

.69] (**Fig. 5a**) on saccade latency. Saccade latency was significantly longer for downward

saccades (215.05 ± 2.6 ms) than saccades in the three other directions (left: 194.91 ±

1.98, 95% CI = [13.60, 26.69]; right: 197.81 ± 1.88, 95% CI = [10.64, 23.73]; upper:

196.22 ± 2.13 ms, 95% CI = [12.29, 25.38]; all p < .001), consistent with previous studies

(Hanning et al., 2022; in press; Kwak et al., 2023; Tzelepi et al., 2010). Participants made

earlier saccades in the invalid (195.85 ± 1.87 ms) than in the valid preview (203.99 ± 2.21

ms; p < .001, 95% CI = [-12.21, -4.07]) and postview (203.20 ± 2.24 ms; p < .001, 95% CI =

[-11.42, -3.28]) viewing conditions. Note that as the perceptual task was relatively harder for the

invalid preview condition, Gabor contrast (adjusted by the staircase) accordingly was higher

(invalid preview: 21.24% ± 3.42%; valid preview: 4.58% ±0.56%; postview: 9.49% ±2.45%), which

might explain the latency difference. Neither saccade amplitudes (7.86 ± 0.03 dva; **Fig. 5b**)

nor saccade precision (0.97 ± 0.01 dva; **Fig. 5c**) differed as a function of target location or

viewing condition.

Post-saccadic eye movements. Shortly after a saccade, gaze is unstable. The eyes may

drift or participants may perform tiny (corrective) saccades. As any post-saccadic movement shifts

the image on the retina and thereby directly affects foveal perception (Rucci & Victor, 2015), we

evaluated participants' post-saccadic eye movements and its potential effect on the preview

effect. To consider any ocular movements after saccade landing, including post-saccadic

oscillations (i.e., ocular instability due to pupil movements; Nyström et al., 2013), fixational

eye movements (micro-saccades) and ocular drift, we analyzed eye movement rate and

distance gaze covered (see Supplemental Material) during the post-saccadic target

presentation with repeated-measures ANOVAs. Post-saccadic movement distance did not

differ between saccade directions or viewing conditions, with gaze samples covering an

average range of 0.51 dva (± 0.03 dva) (**Fig. 6a**).

There was a main effect of saccade direction on post-saccadic eye movement rate,

$F(3, 39) = 3.11$, $p = .037$, $\eta_p^2 = .19$, 95% $CI_\eta = [.00, .35]$, driven by a higher movement rate

after upward (6.02 ± 0.07 Hz) than leftward (5.56 ± 0.08 Hz, $p = .025$, 95% CI = [.040, .87])

saccades (**Fig. 6b**). Because other post-hoc contrasts did not reach significance,

differences in post-saccadic ocular instability can only partially explain the observed

asymmetries in preview effect and post-saccadic sensitivity.

Fig. 5. *Group averaged saccade latency (**a**) and saccade amplitude (**b**) as a function of saccade direction and target location. Error bars indicate 95% CIs. **c**. Saccade endpoint frequency maps averaged across participants depicting saccade landing error. The outlined squares represent the placeholders.*

12   *Post-saccadic target viewing duration.* Another potential source of the observed preview

13 asymmetries could be post-saccadic target viewing duration. A shorter viewing time could bias

14 trans-saccadic integration toward the preview, leading to a larger preview effect. Even though

we timed stimulus offset based on the estimated saccade offset (using an online criterion,

see Methods), the actual target viewing time could have varied as a function of target location

due to differences in saccade velocity or post-saccadic instability that could not be detected

online. Indeed, there was a main effect of saccade direction, $F(3, 39) = 4.31$, $p = .010$, η_p^2

= .25, 95% CI_n = [.018, .41]. However, target viewing duration was slightly but significantly

longer after upward saccades (69.78 ± 0.63ms) than leftward (67.18 ± 0.77ms; $p = .039$,

95% CI = [-.17, 5.37]) and downward (66.34 ± 0.73ms; $p = .0082$, 95% CI = [.67, 6.21])

saccades (**Fig. 6c**), ruling out the possibility that a *shorter* post-saccadic target duration

time contributed to the larger preview effect at the UVM. There was also a main effect of

viewing condition, $F(2, 26) = 3.58$, $p = .042$, $\eta_p^2 = .22$, 95% CI_n = [.00, .42], with a shorter

target viewing found in the invalid preview (67.18 ± 0.64ms) than postview condition (68.25

± 0.69ms; $p = .041$, 95% CI = [.033, 2.11]).

Fig. 6. Group averaged post-saccadic gaze distance (a), saccade rate (b), and target viewing duration (c) as a function of task condition and saccade direction. Error bars indicate 95% CIs.

14
15
16   **Discussion**

We investigated trans-saccadic integration around the visual field, by testing whether and

how the extrafoveal preview effect is modulated by polar angle asymmetries in peripheral vision.

We replicated the typical polar angle asymmetries during fixation –higher contrast sensitivity

at the horizontal than vertical meridian (HVA) and at the lower than upper vertical meridian

(VMA) (Abrams et al., 2012; Carrasco et al., 2001; Greenwood et al., 2017)– and found a

robust preview effect at all cardinals: Post-saccadic foveal sensitivity was higher after a valid

than invalid preview. Importantly, the magnitude of this preview effect was inversely related

to the polar angle asymmetries: larger at locations with lower peripheral sensitivity. These

preview asymmetries could not be explained by differences in saccade parameters. Our novel

findings suggest that the visual system compensates for anticipated asymmetries in

peripheral vision by assigning different weights to the previewed information across saccadic

eye movements.

What could give rise to these preview asymmetries? Right before saccade onset,

contrast sensitivity rapidly increases at the saccade target (Hanning et al., 2022; 2023; in

press; Li et al., 2021; Rolfs & Carrasco, 2012) and renders the peripheral information more

fovea-like already before the eyes start moving (Li et al., 2016, 2019; 2022; Kwak et al., 2023;

Ohl et al., 2017). This *pre-saccadic shift of attention* allows a smooth handoff of pre-saccadic

information to the post-saccadic processing area, which is necessary for continuity in trans-

saccadic perception (De Pisapia et al., 2010; Fabius et al., 2020; Melcher, 2007; Stewart et

al., 2020). However, previous studies have found an absent or reduced presaccadic benefit

before upward saccades compared to saccades to other directions (Hanning et al., 2022; in

press). This could explain the worse post-saccadic foveal discrimination in the *postview task*

after upward saccades than saccades in other directions: Without attention shifting

prospectively to the saccade target –easing trans-saccadic integration by rendering the

peripheral visual input more fovea-like– visual sensitivity might recover more slowly after

saccade offset. Similarly, in the integration task, because the target disappeared briefly after

saccade offset (before vision could fully recover), participants needed to rely more on the

peripheral preview (Baumann et al., 2023), leading to a larger preview effect at the UVM.

Regardless of the mechanism underlying the worsened post-saccadic sensitivity following

upward saccades, the visual system might have given more decision weight to the peripheral

preview when a less reliable post-saccadic percepts was expected, consistent with the

optimal integration framework (Ganmor et al., 2015; Steward et al., 2020; Wolf & Schütz,

2015). This differential weight assignment, however, cannot explain the reverse VMA in the

preview effect, as the here observed post-saccadic foveal sensitivity was equivalent between

the LVM and HM. Our regression analysis further verified that it could not explain the variance

in the preview effect, arguing against the optimal integration hypothesis.

Another potential mechanism that could have cause the observed preview

asymmetries is pre-saccadic "*foveal remapping*": Neuroimaging studies have found that

before a saccade, neurons that process foveal information shift their receptive fields to also

process detailed information about the peripheral saccade target (e.g., Fan et al., 2016;

Neupane et al., 2020). Likewise, psychophysical evidence has shown that foveal processing

of the saccade target begins already before saccade onset (Kroell & Rolfs, 2022). Akin to

presaccadic-attention, this mechanism facilitates visual processing at the saccade target,

and thereby a smooth transition from pre- to post-saccadic vision. Our visual system might

strategically maximize foveal remapping for saccade targets with comparatively lower visual

sensitivity (e.g., the LVM and UVM) to compensate for polar angle asymmetries in peripheral

vision, leading to the observed reversed preview asymmetries.

Alternatively, the preview asymmetries might arise from the very first stage of trans-

saccadic integration – when a prediction about the saccadic target is made based on

peripheral inputs long before saccade onset (Fabius et al., 2020; Huber-Huber et al., 2019;

Jensen et al., 2021). A stronger prediction about the peripheral saccade target could cause

its (relative) up-weighting, leading to a larger preview effect. According to predictive

processing, the preview effect is a measure of how much post-saccadic perception is biased

toward a pre-saccadic prediction about the target. Although stronger prediction is intuitively

associated with a higher peripheral sensitivity, strength and content of pre-saccadic

predictions are learned over time: The repeated exposure to consistently invalid previews

(e.g., an orange always changes to a basketball during a saccade), causes participants to

make predictions about the saccadic target in the learnt direction, reducing the preview

effect in response to statistical regularity (Huber-Huber et al., 2021; Osterbrink & Herwig,

2021). Future research is needed to investigate how pre-saccadic prediction strength and

content change around the visual field and how it is related to the preview asymmetries.

Our study is the first to investigate how heterogeneously perceived peripheral

information is integrated across saccadic eye movements and differs from previous studies

testing how object visibility affects trans-saccadic integration (Liu et al., 2022; Wolf & Schütz,

2015) in an important way: Instead of external manipulation on stimulus attributes, the

perceptual quality of peripheral stimuli in the current study was constrained by our visual

system -an internal factor. On the one hand, these asymmetries in peripheral vision are so

robust that they cannot be overridden by any attention mechanism studied so far (e.g.,

Carrasco et al., 2001; Fernández et al., 2019; Hanning et al., 2022; in press; Kwak et al.,

2023; Purokayastha et al., 2021). On the other hand, the current results indicate that our

brain compensates for these asymmetries during trans-saccadic integration.

These results have significant implications in real-world applications. When driving,

for example, people tend to fixate on one point while covertly monitoring the periphery. Polar
angle asymmetries in peripheral vision suggest that drivers will be the most sensitive to
objects along the HM (e.g., pedestrians), followed by the LVM (e.g., signals on the instrument
panel), and lastly the UVM (objects in the distance/the sky). The observed preview
asymmetries show that when drivers make saccadic eye movements to different locations,
integrating information across these saccades results in more homogenous performance for
locations around the visual field.

This study highlights the need in future work to study pre-, post-, and trans-saccadic
perception for different saccade directions. Most studies investigating perceptual
modulations around saccades, such as pre-saccadic attention (Buonocore et al., 2017; De
Pisapia et al., 2010; Li et al., 2016; 2019; Melcher, 2007; Ohl et al., 2017), predictive
remapping (Duhamel et al., 1992; Jonikaitis et al., 2013; Wang et al., 2016), and trans-
saccadic integration (Ganmor et al., 2015; Huber-Huber et al., 2019; Huber-Huber &
Melcher, 2021; Liu et al., 2022; Wolf & Schütz, 2015) are based on horizontal eye
movements. The here observed preview asymmetries, in line with polar angle differences in
pre-saccadic attention (Hanning et al., 2022; in press) call into question the generalizability
of these findings from one saccade angle to another.

In summary, the extrafoveal preview effect, as a measure of trans-saccadic
integration, follows a reverse pattern of peripheral polar angle asymmetries during fixation.
This finding suggests that the visual system actively compensates for polar angle asymmetries in
peripheral visual perception by selectively assigning higher weights to the less-well perceived
preview information around the time of a saccade. Future neurophysiological studies should
investigate how and where such compensation is implemented in the human brain.

Acknowledgements

This research was supported by a Marie Skłodowska-Curie individual fellowship (MSCA-IF 898520) by the European Commission to NMH, a National Institutes of Health – National Eye Institute grant R01 EY019693 to MC, and by the NYUAD Center for Brain and Health, funded by Tamkeen under NYU Abu Dhabi Research Institute grant CG012.

Author Contributions

Xiaoyi Liu: Conceptualization; Methodology; Software; Formal Analysis; Investigation; Data Curation; Writing – Original Draft; Writing – Review & Editing; Visualization

David Melcher: Conceptualization; Writing – Review & Editing; Supervision; Funding Acquisition.

Marisa Carrasco: Conceptualization; Writing – Review & Editing; Validation; Supervision; Funding Acquisition.

Nina M. Hanning: Conceptualization; Methodology; Investigation; Writing – Review & Editing; Visualization; Supervision; Funding Acquisition.

References

Abrams, J., Nizam, A., & Carrasco, M. (2012). Isoeccentric locations are not equivalent: The extent of the vertical meridian asymmetry. *Vision Research*, *52* (1), 70–78.

Angele, B., & Rayner, K. (2011). Parafoveal processing of word n+ 2 during reading: Do the preceding words matter? *Journal of Experimental Psychology: Human Perception and Performance*, *37* (4), 1210.

Anton-Erxleben, K., Herrmann, K., & Carrasco, M. (2013). Independent effects of adaptation and attention on perceived speed. *Psychological science, 24*(2), 150-159.

Bakdash, J. Z., & Marusich, L. R. (2017). Repeated measures correlation. *Frontiers in Psychology*, *8*, 456.

Baldwin, A. S., Meese, T. S., & Baker, D. H. (2012). The attenuation surface for contrast sensitivity has the form of a witch's hat within the central visual field. *Journal of Vision*, *12* (11):23, 1–17.

Barbot, A., & Carrasco, M. (2017). Attention modifies spatial resolution according to task demands. *Psychological Science*, *28* (3), 285–296.

Barbot, A., Xue, S., & Carrasco, M. (2021). Asymmetries in visual acuity around the visual field. *Journal of Vision*, *21* (1):2, 1–23.

Baumann, M. P., Bogadhi, A. R., Denninger, A. F., & Hafed, Z. M. (2023). Sensory tuning in neuronal movement commands. *Proceedings of the National Academy of Sciences*, *120* (38), e2305759120.

Brainard, D. H. (1997). The psychophysics toolbox. *Spatial vision, 10*(4), 433-436.

Buonocore, A., Dimigen, O., & Melcher, D. (2020). Post-saccadic face processing is modulated by pre-saccadic preview: Evidence from fixation-related potentials. *Journal of Neuroscience*, *40* (11), 2305–2313.

Buonocore, A., Fracasso, A., & Melcher, D. (2017). Pre-saccadic perception: Separate time courses for enhancement and spatial pooling at the saccade target. *PLoS One*,*12* (6), e0178902.

Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research*, *51* (13), 1484–1525.

Carrasco, M., Talgar, C. P., & Cameron, E. L. (2001). Characterizing visual performance fields: Effects of transient covert attention, spatial frequency, eccentricity, task and set size. *Spatial Vision*, *15* (1), 61–75.

Caruso, V. C., Pages, D. S., Sommer, M. A., & Groh, J. M. (2018). Beyond the labeled line: variation in visual reference frames from intraparietal cortex to frontal eye fields and the superior colliculus. *Journal of Neurophysiology, 119*(4), 1411–1421.

Cornelissen, F. W., Peters, E. M., & Palmer, J. (2002). The eyelink toolbox: Eye tracking with matlab and the psychophysics toolbox. *Behavior Research Methods, Instruments, & Computers*, *34* (4), 613–617.

Cont, C., & Zimmermann, E. (2021). The motor representation of sensory experience. *Current Biology, 31*(5), 1029–1036.

De Pisapia, N., Kaunitz, L., & Melcher, D. (2010). Backward masking and unmasking across saccadic eye movements. *Current Biology*, *20*(7), 613–617.

Deubel, H., & Schneider, W. X. (1996). Saccade target selection and object recognition: Evidence for a common attentional mechanism. *Vision Research, 36*(12), 1827–1837.

Deubel, H. (2008). The time course of presaccadic attention shifts. *Psychological Research*, *72* (6), 630–640.

Duhamel, J.-R., Colby, C. L., & Goldberg, M. E. (1992). The updating of the representation of visual space in parietal cortex by intended eye movements. *Science*, *255* (5040), 90–92.

Edwards, G., VanRullen, R., & Cavanagh, P. (2018). Decoding trans-saccadic memory. *Journal of Neuroscience*, 38 (5), 1114–1123.

Engbert, R., & Mergenthaler, K. (2006). Microsaccades are triggered by low retinal image slip. *Proceedings of the National Academy of Sciences, 103*(18), 7192–7197.

Ernst, M. O., & Banks, M. S. (2002). Humans integrate visual and haptic information in a statistically optimal fashion. *Nature*, *415* (6870), 429–433.

Ernst, M. O., & Bülthoff, H. H. (2004). Merging the senses into a robust percept. *Trends in cognitive Sciences*, *8* (4), 162–169.

Fabius, J. H., Fracasso, A., Acunzo, D. J., Van der Stigchel, S., & Melcher, D. (2020). Low-level visual information is maintained across saccades, allowing for a postsaccadic handoff between visual areas. *Journal of Neuroscience*, *40* (49), 9476–9486.

Fan, X., Wang, L., Shao, H., Kersten, D., & He, S. (2016). Temporally flexible feedback signal to foveal cortex for peripheral object recognition. *Proceedings of the National Academy of Sciences, 113*(41), 11627-11632.

Fernández, A., Denison, R. N., & Carrasco, M. (2019). Temporal attention improves perception similarly at foveal and parafoveal locations. *Journal of Vision, 19*(1):12, 1–10.

Ganmor, E., Landy, M. S., & Simoncelli, E. P. (2015). Near-optimal integration of orientation information across saccades. *Journal of Vision*, *15* (16):8, 1–12.

Greenwood, J. A., Szinte, M., Sayim, B., & Cavanagh, P. (2017). Variations in crowding, saccadic precision, and spatial localization reveal the shared topology of spatial vision. *Proceedings of the National Academy of Sciences*, *114* (17), E3573–E3582.

Hanning, N. M., Deubel, H., & Szinte, M. (2019). Sensitivity measures of visuospatial attention. *Journal of Vision, 19*(12):17, 1–13.

Hanning, N. M., Himmelberg, M. M., & Carrasco, M. (2022). Presaccadic attention enhances contrast sensitivity, but not at the upper vertical meridian. *iScience*, *25* (2), 103851.

Hanning, N. M., Himmelberg, M. M., & Carrasco, M. (in press). Presaccadic attention depends

on eye movement direction and is related to V1 cortical magnification. *Journal of Neuroscience.*

Hanning, N. M., Fernández, A., & Carrasco, M. (2023). Dissociable roles of human frontal eye fields and early visual cortex in presaccadic attention. *Nature Communications*, 14(1), 5381.

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2020). Stimulus-dependent contrast sensitivity asymmetries around the visual field. *Journal of Vision*, *20*(9):18, 18–19.

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2023). Polar angle asymmetries in visual perception and neural architecture. *Trends in Neurosciences, 46*(6), 445-458.

Huber-Huber, C., Buonocore, A., Dimigen, O., Hickey, C., & Melcher, D. (2019). The peripheral preview effect with faces: Combined eeg and eye-tracking suggests multiple stages of trans-saccadic predictive and non-predictive processing. *NeuroImage*, *200*, 344–362.

Huber-Huber, C., Buonocore, A., & Melcher, D. (2021). The extrafoveal preview paradigm as a measure of predictive, active sampling in visual perception. *Journal of Vision*, *21* (7):12, 1–13.

Huber-Huber, C., & Melcher, D. (2021). The behavioural preview effect with faces is susceptible to statistical regularities: Evidence for predictive processing across the saccade. *Scientific Reports*, *11* (1), 942.

Hutzler, F., Schuster, S., Marx, C., & Hawelka, S. (2019). An investigation of parafoveal masks with the incremental boundary paradigm. *PLoS One*, *14* (2), e0203013.

Jensen, O., Pan, Y., Frisson, S., & Wang, L. (2021). An oscillatory pipelining mechanism supporting previewing during visual exploration and reading. *Trends in Cognitive Sciences*, *25* (12), 1033–1044.

Jigo, M., Tavdy, D., Himmelberg, M. M., & Carrasco, M. (2023). Cortical magnification eliminates differences in contrast sensitivity across but not around the visual field. *Elife*, *12*, e84205.

Jonikaitis, D., Szinte, M., Rolfs, M., & Cavanagh, P. (2013). Allocation of attention across saccades. *Journal of Neurophysiology, 109*(5), 1425–1434.

Körding, K. P., & Wolpert, D. M. (2004). Bayesian integration in sensorimotor learning. *Nature*, *427* (6971), 244–247.

Kowler, E., Anderson, E., Dosher, B., & Blaser, E. (1995). The role of attention in the programming of saccades. *Vision Research, 35*(13), 1897–1916.

Kroell, L. M., & Rolfs, M. (2022). Foveal vision anticipates defining features of eye movement targets. *Elife, 11*, e78106.

Kurzawski, J. W., Burchell, A., Thapa, D., Winawer, J., Majaj, N. J., & Pelli, D. G. (2023). The Bouma law accounts for crowding in fifty observers. *Journal of Vision*, 23(8), 1–34.

Kwak, Y., Hanning, N. M., & Carrasco, M. (2023). Presaccadic attention sharpens visual acuity. *Scientific Reports*, *13* (1), 2981.

Li, H.-H., Barbot, A., & Carrasco, M. (2016). Saccade preparation reshapes sensory tuning. *Current Biology*, *26*(12), 1564–1570.

Li, H.-H., Hanning, N. M., & Carrasco, M. (2021). To look or not to look: dissociating presaccadic and covert spatial attention. *Trends in Neurosciences*, 44(8), 669–686.

Li, H.-H., Pan, J., & Carrasco, M. (2019). Presaccadic attention improves or impairs performance by enhancing sensitivity to higher spatial frequencies. *Scientific Reports*, 9(1), 1–9.

Li, H.-H., Pan, J., & Carrasco, M. (2021). Different computations underlie overt presaccadic and covert spatial attention. *Nature Human Behaviour*, *5*(10), 1418–1431.

Liu, X., Huber-Huber, C., & Melcher, D. (2022). The trans-saccadic extrafoveal preview effect is modulated by object visibility. *2022 Symposium on Eye Tracking Research and Applications*, 1–7.

McDonald, S. A. (2005). Parafoveal preview benefit in reading is not cumulative across multiple saccades. *Vision Research*, *45*(14), 1829–1834.

Melcher, D. (2007). Predictive remapping of visual features precedes saccadic eye movements. *Nature Neuroscience*, *10*(7), 903–907.

Melcher, D., & Colby, C. L. (2008). Trans-saccadic perception. *Trends in Cognitive Sciences*, *12*(12), 466–473.

Montaser-Kouhsari, L., & Carrasco, M. (2009). Perceptual asymmetries are preserved in short-term memory tasks. *Attention, Perception, & Psychophysics*, *71*(8), 1782–1792.

Neupane, S., Guitton, D., & Pack, C. C. (2020). Perisaccadic remapping: What? how? why?. *Reviews in the Neurosciences, 31*(5), 505-520.

Nyström, M., Hooge, I., & Holmqvist, K. (2013). Post-saccadic oscillations in eye movement data recorded with pupil-based eye trackers reflect motion of the pupil inside the iris. *Vision Research*, *92*, 59–66.

Osterbrink, C., & Herwig, A. (2021). Prediction of complex stimuli across saccades. *Journal of Vision*, *21*(2):10, 1–15.

Ohl, S., Kuper, C., & Rolfs, M. (2017). Selective enhancement of orientation tuning before saccades. *Journal of Vision*, 17(13):2, 1-11.

Pentland, A. (1980). Maximum likelihood estimation: The best pest. *Percept. Psychophys.*, *28*, 377–379.

Prins, N., & Kingdom, F. A. (2018). Applying the model-comparison approach to test specific research hypotheses in psychophysical research using the palamedes toolbox. *Frontiers in Psychology*, *9*, 1250.

Purokayastha, S., Roberts, M., & Carrasco, M. (2021). Voluntary attention improves performance similarly around the visual field. *Attention, Perception, & Psychophysics*, *83*(7), 2784–2794.

Rayner, K., McConkie, G. W., & Zola, D. (1980). Integrating information across eye movements. *Cognitive Psychology*, *12*(2), 206–226.

Rijsdijk, J., Kroon, J., & Van der Wildt, G. (1980). Contrast sensitivity as a function of position on the retina. *Vision Research*, *20*(3), 235–241.

Roberts, M., Ashinoff, B. K., Castellanos, F. X., & Carrasco, M. (2018). When attention is intact in adults with ADHD. *Psychonomic Bulletin & Review*, 25, 1423–1434.

Rolfs, M., & Carrasco, M. (2012). Rapid simultaneous enhancement of visual sensitivity and perceived contrast during saccade preparation. *Journal of Neuroscience*, *32*(40), 13744–13752a.

Rucci, M., & Victor, J. D. (2015). The unsteady eye: an information-processing stage, not a bug. *Trends in Neurosciences, 38*(4), 195–206.

Sadeh, M., Sajad, A., Wang, H., Yan, X., & Crawford, J. D. (2015). Spatial transformations between superior colliculus visual and motor response fields during head-unrestrained gaze shifts. *European Journal of Neuroscience*, *42*(11), 2934–2951.

Schall, J. D., & Hanes, D. P. (1993). Neural basis of saccade target selection in frontal eye field during visual search. *Nature*, *366*(6454), 467–469.

Sharp, P., Melcher, D., & Hickey, C. (2018). Endogenous attention modulates the temporal window of integration. *Attention, Perception, & Psychophysics*, *80*, 1214–1228.

Schotter, E. R., Angele, B., & Rayner, K. (2012). Parafoveal processing in reading. *Attention, Perception, & Psychophysics, 74*, 5–35.

Stigliani, A., Weiner, K. S., & Grill-Spector, K. (2015). Temporal processing capacity in high-level visual cortex is domain specific. *Journal of Neuroscience*, *35*(36), 12412–12424.

Stewart, E. E., Valsecchi, M., & Schütz, A. C. (2020). A review of interactions between peripheral and foveal vision. *Journal of Vision, 20*(12):2, 1–35.

Talgar, C. P., & Carrasco, M. (2002). Vertical meridian asymmetry in spatial resolution: Visual and attentional factors. *Psychonomic Bulletin & Review*, *9*, 714–722.

Tzelepi, A., Laskaris, N., Amditis, A., & Kapoula, Z. (2010). Cortical activity preceding vertical saccades: A meg study. *Brain Research*, *1321*, 105–116.

Vallines, I., & Greenlee, M. W. (2006). Saccadic suppression of retinotopically localized blood oxygen level-dependent responses in human primary visual area v1. *Journal of Neuroscience*, *26*(22), 5965–5969.

Wang, X., Fung, C. A., Guan, S., Wu, S., Goldberg, M. E., & Zhang, M. (2016). Perisaccadic receptive field expansion in the lateral intraparietal area. *Neuron*, *90*(2), 400–409.

Wolf, C., & Schütz, A. C. (2015). Trans-saccadic integration of peripheral and foveal feature information is close to optimal. *Journal of Vision*, *15*(16):1, 1–18.

Zimmermann, E., Weidner, R., Abdollahi, R. O., & Fink, G. R. (2016). Spatiotopic adaptation in

visual areas. *Journal of Neuroscience, 36*(37), 9526–9534.