Neural Tracking Measures of Speech Intelligibility: Manipulating Intelligibility while Keeping Acoustics Unchanged

I.M Dushyanthi. Karunathilake[1], Joshua P. Kulasingham[2], and Jonathan Z. Simon[1,3,4,*]

[1]Department of Electrical and Computer Engineering, University of Maryland, College Park, MD, 20742, USA, [2]Department of Electrical Engineering, Linköping University, SE, [3]Department of Biology, University of Maryland, College Park, MD 20742, USA, [4]Institute for Systems Research, University of Maryland, College Park, MD 20742, USA

[*]Corresponding author: Jonathan Z. Simon

Email: jzsimon@umd.edu

Author Contributions: I.M.D.K., J.P.K, and J.Z.S. conceived and designed the research; I.M.D.K. collected data; I.M.D.K. and J.Z.S. analyzed data; I.M.D.K. drafted manuscript; I.M.D.K., J.P.K and J.Z.S. edited and revised manuscript.

Competing Interest Statement: The authors declare no competing interests

Keywords: Speech Intelligibility, MEG, TRF, neural tracking, vocoded speech

Abstract

Neural speech tracking has advanced our understanding of how our brains rapidly map an acoustic speech signal onto linguistic representations and ultimately meaning. It remains unclear, however, how speech intelligibility is related to the corresponding neural responses. Many studies addressing this question vary the level of intelligibility by manipulating the acoustic waveform, but this makes it difficult to cleanly disentangle effects of intelligibility from underlying acoustical confounds. Here, using magnetoencephalography (MEG) recordings, we study neural measures of speech intelligibility by manipulating intelligibility while keeping the acoustics strictly unchanged. Acoustically identical degraded speech stimuli (three-band noise vocoded, ~20 s duration) are presented twice, but the second presentation is preceded by the original (non-degraded) version of the speech. This intermediate priming, which generates a 'pop-out' percept, substantially improves the intelligibility of the second degraded speech passage. We investigate how intelligibility and acoustical structure affects acoustic and linguistic neural representations using multivariate Temporal Response Functions (mTRFs). As expected, behavioral results confirm that perceived speech clarity is improved by priming. TRF analysis reveals that auditory (speech envelope and envelope onset) neural representations are not affected by priming, but only by the acoustics of the stimuli (bottom-up driven). Critically, our findings suggest that segmentation of sounds into words emerges with better speech intelligibility, and most strongly at the later (~400 ms latency) word processing stage, in prefrontal cortex (PFC), in line with engagement of top-down mechanisms associated with priming. Taken together, our results show that word representations may provide some objective measures of speech comprehension.

Significance Statement

Electrophysiological studies have shown that brain tracks different speech features. How these neural tracking measures are modulated by speech intelligibility, however, remained elusive. Using noise-vocoded speech and a priming paradigm, we disentangled the neural effects of intelligibility from the underlying acoustical confounds. Neural intelligibility effects are analyzed at

both acoustic and linguistic level using multivariate Temporal Response Functions. Here, we find evidence for an effect of intelligibility and engagement of top-down mechanisms, but only in responses to lexical structure of the stimuli, suggesting that lexical responses are strong candidates for objective measures of intelligibility. Auditory responses are not influenced by intelligibility but only by the underlying acoustic structure of the stimuli.

Introduction

When we listen to speech, our brains rapidly map the acoustic sounds into linguistic representations while recruiting complex cognitive processes to derive the intended meaning (1, 2). A fundamental goal in auditory neurophysiology is to understand how the brain transforms the acoustic signal into meaningful content. Along this avenue, a large body of research has demonstrated that neural responses time lock to different features of the speech signal ("neural speech tracking") (2, 3). These features have primarily included acoustic features like the speech envelope and envelope onset, but more recently also include linguistic units such as word onsets, phoneme onsets and context-based measures along different levels of the linguistic hierarchy. However, it remains unclear how these neural tracking measures are affected by intelligibility.

Neural tracking measures of intelligibility have often been investigated using experimental designs that manipulate intelligibility by altering the underlying acoustical structure, such as through time compression (4, 5), disruption of spectro-temporal details (6–8) and time reversal (4, 9). While these studies successfully demonstrate differences in cortical tracking responses, interpreting these findings is not straightforward, as the observed changes in the neural response may arise from the alterations in the acoustic waveform (bottom-up), rather than the intelligibility change itself (top-down). Intelligibility related neuro-markers derived from neural responses play a crucial role in advancing our understanding of the neurophysiology of the speech understanding. They would contribute to the clinical evaluation of auditory function across diverse clinical populations and aid in the hearing device evaluation (10). In situations where obtaining subjective measure is challenging, such as infants or individuals with cognitive disabilities, as well as when subjective estimates are affected by individual differences, neuro-markers would offer a non-invasive and objective means to investigate underlying neural processes.

In this study, we disentangle intelligibility from any underlying acoustical confounds by using single instances of noise-vocoded speech combined with a priming paradigm. Noise-vocoded speech greatly reduces intelligibility by removing spectral details but still preserving the slow temporal envelope, and is often used as a surrogate for speech perception by cochlear implant patients (11). It is generated by processing the original speech signal through a multi-frequency-channel vocoder, where higher numbers of channels retain more intelligibility due to the retention of more spectral information. With a sufficiently low number of channels, most vocoded speech is largely unintelligible without practice. In the current study, we used three-band noise-vocoded speech. Magnetoencephalography (MEG) data were recorded from young adult participants (N = 25) as they listened to a passage of noise-vocoded speech, first before any priming (PRE), followed by listening to the original, non-degraded version of the same passage to invoke priming (CLEAN), and then finally listening to the same noise-vocoded speech passage as before (POST), repeated for 36 trials (see Figure 1(A)). At the end of each vocoded speech passage, participants were asked to rate the perceived speech clarity on a scale from 0 to 5. Compared to previous studies (12–15) using a similar paradigm with shorter sentences (<5 s), the current study uses much longer passages (~20 s), making it challenging to rely solely on short-term memory to understand POST vocoded speech.

To characterize how different speech features are tracked in the cortical response, we utilized multivariate temporal response function (mTRF) analysis (16–18), as illustrated in Figure 1(C). Similar to conventional event-related potentials (ERPs), TRFs are used to assess how the brain reacts to different speech features over time. In contrast to ERPs, which rely on averaging numerous short responses to determine the brain's reaction to given stimulus, TRF enables the

examination of brain's responses to continuous speech, as well as simultaneous responses to multiple speech features (1). We included three classes of speech features for which to extract responses: acoustic, sub-lexical, and lexical. The specific features employed were speech envelope, envelope onset, phoneme onset, and word onset, to cover a range of neural responses from acoustic processing to lexical level processing.

We first determined whether each of these features are represented in the cortical response by evaluating the explained response variability for each of the speech conditions (PRE, POST and CLEAN). Then we investigated how the cortical representations of the different speech features are modulated by intelligibility (PRE vs POST) and by acoustics (vocoded PRE and POST vs CLEAN) at different auditory processing stages, by comparing the TRF peak amplitudes and explained variability. All analysis were performed on source-localized brain responses, and were restricted to temporal, frontal and parietal brain regions.

Results and Discussion

Behavioral performance increases with speech priming

We first determined the extent to which the speech priming (perceptual learning) affected the speech intelligibility between PRE and POST vocoded conditions using perceived speech clarity ratings. A linear mixed effect model (LMEM) was modelled with perceived clarity rating as the dependent variable, condition as a fixed effect, and random intercept and slopes for condition by subject as random effects (*clarity rating ~ 1 + condition + (1+condition|Subject)*). As illustrated in Figure 1(B), the fixed effects of condition revealed that rated speech clarity in the POST vocoded condition is improved compared to PRE (increase = 1.81, *SE* = 0.16, *p* < 0.001).

This result supports the idea that presentation of the clear speech, which provides information regarding both the linguistic content (i.e., words, content) and physical acoustical structure (i.e., rhythm, pace) of the degraded speech, facilitates (a top-down influence on the) understanding of POST vocoded speech. Thus, in agreement with previous studies, speech that is acoustically identical but initially unintelligible can be made intelligible through perceptual learning (12, 14, 15, 19–23)

Neural responses to acoustic features do not index speech intelligibility, only acoustics

To test the extent to which each of the acoustic features are represented in the brain and for each condition, we first compared the predictive power, measured as the explained variability (R^2) of the full model against a reduced model that excluded the predictor of interest (see Figure (B,C) right column, brain plots). This prediction accuracy analysis revealed that both acoustic envelope and envelope onset significantly contribute to the model's absolute predictive power for all three conditions, PRE (envelope: t_{max} = 6.5, *p* < 0.001, onset: t_{max} = 6.2, *p* < 0.001), POST (envelope: t_{max} = 7.3, *p* < 0.001, onset: t_{max} = 5.9, *p* < 0.001) and CLEAN (envelope: t_{max} = 6.3, *p* < 0.001, onset: t_{max} = 7.24, *p* < 0.001). These findings suggest that acoustic features are processed irrespective of the stimuli intelligibility. The anatomical distribution of significant acoustic feature processing for each condition was observed in locations spreading spatially from Heschl's gyrus (HG) to superior temporal gyrus (STG) and much of temporal lobe. This distribution was bilateral, and also dominantly in the right hemisphere except for clean speech envelope and vocoded speech envelope onset processing ((left vs right hemisphere) envelope: PRE_t_{max} = -5.35, *p* = 0.02, POST_t_{max} = -5.25, *p* = 0.004, CLEAN_t_{max} = -3.3, *p* = 0.41, envelope onset: PRE_t_{max} = -4.27, *p* = 0.02, POST_t_{max} = -3.46, *p* = 0.39, CLEAN_t_{max} = -4.68, *p* = 0.006). This pattern of source localization, including right-hemisphere dominance, suggests that the processing of these speech features processing relies heavily on bottom-up driven mechanisms (24, 25).

In order to determine whether the processing of acoustic features differed based on speech intelligibility (PRE vs POST) or acoustics of the stimuli (vocoded vs clean), we compared the

model improvements between speech conditions for each acoustic feature individually. Pairwise mass-univariate related samples t-test revealed that there is no significant difference between PRE vs POST vocoded speech with respect to both envelope (t_{max} = -3.9, p = 0.57) and envelope onset (t_{max} = 4.39, p = 0.09). In contrast, the variance explained due to envelope processing was significantly stronger for vocoded speech compared to clean speech (t_{max} = 6.8, p < 0.001), and the opposite for envelope onset processing (t_{max} = -7.44, p < 0.001), suggesting that auditory responses are mainly driven by the manipulations in the stimuli.

We then investigated how the brain responds to acoustic speech representation at different cortical processing stages with an amplitude analysis of the TRF waveforms. In analogy to the EEG P1-N1 peaks at the corresponding latencies, the envelope TRF showed two main peaks (Figure 2(A)), a positive polarity peak at ~50 ms latency followed by a negative polarity peak ~100 ms, known as the $M50_{TRF}$ and $M100_{TRF}$ respectively (1, 17). Analogous to these envelope peaks, envelope onset responses also showed two main peaks at ~75 ms and ~130 ms, consistent with previous studies (1, 26). The $M50_{TRF}$ and $M100_{TRF}$ peaks can be ascribed to different auditory cortical processing stages with the corresponding latencies (27). It has been suggested that the early $M50_{TRF}$ peak dominantly reflects the neural encoding of low-level processing, e.g., physical acoustic of the stimuli (28, 29), whereas the $M100_{TRF}$ peak reflects additional higher-level processing, e.g., selective attention (17, 28, 30). Using a paired-samples t-test, we investigated the extent to which these peak amplitudes are affected by the intelligibility and acoustic characteristics of the stimuli (Figure 2(B,C)). In line with the prediction accuracy results above, we failed to find an effect of intelligibility for the neural responses to acoustic features (both envelope and envelope onset) for both the $M50_{TRF}$ (envelope: p = 0.63, envelope onset: p = 0.80) and $M100_{TRF}$ peaks (envelope: p = 0.71, envelope onset: p = 0.29)), but TRF peak amplitudes were significantly affected by the differences in acoustics for both the $M50_{TRF}$ (PRE vs CLEAN: envelope p < 0.001, envelope onset p < 0.001) and $M100_{TRF}$ peaks (PRE vs CLEAN: envelope p = 0.03, envelope onset p = 0.003); effect sizes are reported in Supplemental Table S1. These results again support the previous finding that such auditory cortical responses are not necessarily driven by the intelligibility or the linguistic content of the stimuli, but rather reflects the sensitivity to differences in sensory input (12, 15, 22, 23, 31). Other studies, in contrast, which reported that increased intelligibility can improve cortical speech tracking measures, did so only in settings where the corresponding underlying acoustics were changed as well, creating a confound (6, 13).

The $M50_{TRF}$ peak of the envelope response was significantly stronger for vocoded speech compared to clean speech, whereas this effect was reversed for the envelope onset response (Table S1), suggesting distinct mechanisms are involved in envelope and envelope onset processing. Analysis of the ratio of $M50_{TRF}$ between envelope and envelope onset comparison across conditions provided additional support for the reliability of this reversing effect (PRE vs POST: t_{24} = 0.53, p = 0.56, PRE vs CLEAN: t_{24} = -6.25, p < 0.001). This finding for the early envelope TRF peak is consistent with previous studies (8, 22, 32, 33), where it was proposed that this effect is modulated by task demand (22) or sensory gain of acoustic properties (8); this early peak is too early to be modulated by attention (17). Instead, we propose that the higher envelope TRF amplitudes observed for vocoded speech are a result of its low spectral variability, which leads to higher levels of synchronization along the tonotopic axes, resulting in stronger MEG responses. In contrast, reduced vocoded speech envelope onset tracking can be attributed to the loss of salient acoustic onsets in vocoded speech, a result of the loss of spectrotemporal details intrinsic to the process of vocoding (see Methods, predictor variables).

Compared to vocoded speech, both envelope and envelope onset in clean speech exhibited stronger $M100_{TRF}$ peak amplitudes. This suggests that the observed differences in vocoded vs clean for the $M50_{TRF}$ and $M100_{TRF}$ are due to different underlying brain processes, where the $M50_{TRF}$ depends more strongly on the physical acoustics while $M100_{TRF}$ also reflects higher-level processing (17, 28, 30). However, to the extent $M100_{TRF}$ peak is indicative of higher-level

processing, it is a striking finding that it is not influenced by intelligibility.

Responses to Phoneme onsets

Although phoneme onsets (see Figure 2(D)) significantly contributed to the model's predictive power in each speech condition (PRE: t_{max} = 4.9, p < 0.001, POST: t_{max} = 5.4, p < 0.001, CLEAN: t_{max} = 6.8, p < 0.001), no significant differences were detected between the conditions. Consistent with previous studies(1), the phoneme onset TRF showed two main peaks, an early positive polarity peak ~ 80 ms and a late negative polarity peak ~150 ms, which were comparable to the envelope onset peaks. The amplitude of the early peak showed no difference between PRE vs POST (p = 0.68) nor vocoded vs clean speech (p = 0.28), indicating that these early responses are not modulated by either acoustics or intelligibility. However, the late negative polarity peak was stronger in clean speech compared to vocoded speech (CLEAN vs PRE: p = 0.003, CLEAN vs POST: p < 0.001, PRE vs POST: p = 0.31), indicating that this late processing stage is affected by the physical acoustics. Phoneme onset processing originated bilaterally from areas including primary auditory cortex and was right lateralized for clean speech (t_{max} = -4.39, p < 0.02), supporting previous results that phoneme onset processing occurs early in the auditory processing hierarchy and may reflect more acoustic processing than linguistic. These results are aligned with a previous EEG study (23) that investigated the impact of perceived intelligibility on phoneme level processing (a more complex measure does show effects of priming in the delta band). Thus, despite the sub-lexical or linguistic nature of the phoneme onset feature, the neural responses suggest that, as a neural measure, it functions more like an auditory (or intermediate auditory-linguistic) measure.

Neural responses to lexical segmentation indexes speech intelligibility

Novel in the current study, we incorporated a lexical feature to extend the investigation of the effects of intelligibility with respect to neural speech representation: the word onset response. Prediction accuracy analysis revealed that the word onset responses significantly explain additional variability in the measured neural response over and beyond acoustic and phoneme onset features, across all three speech conditions (PRE: t_{max} = 4.6, p < 0.001, POST: t_{max} = 4.6, p < 0.001, clean: t_{max} = 5.1, p < 0.001). The word onset TRFs (see Figure 3(A)) showed two main peaks, an early positive polarity peak (~100 ms) and a substantially later negative polarity peak (~400 ms). The late peak is comparable to the latency and polarity of classical N400 responses (9, 34, 35), a potential marker of complex language processing. Interestingly, the peak amplitude comparison revealed that the intelligibility of the speech modulates both early (p = 0.02; Cohen's d = 0.39) and late peak amplitudes (p < 0.001; d = 0.82), with a substantially greater effect size observed for the late peak. Comparing clean vs vocoded speech, we found that the clean speech TRF amplitudes are stronger compared to vocoded speech for both early (p < 0.001) and late peak responses (PRE vs CLEAN: p < 0.001; d = 0.43, POST vs CLEAN: p = 0.03; d = 1.20). Additionally, the late peak amplitude of POST is significantly closer to that of CLEAN than the corresponding amplitude of PRE is to CLEAN, suggesting that the late word onset responses of POST are more similar to CLEAN than those of PRE.

Because prediction accuracy, as implemented above, integrates over a longer time window (–200 - 800 ms), it is not able to disentangle specific contributions to the prediction accuracy and source localizations of each processing stage. To address this, we conducted a separate analysis on the explained variability, focusing on early (50-300 ms) and late (330-650 ms) processing stages separately (see Figure 3(B,C)). As can be seen from Figure 3(C), prediction accuracy comparisons between PRE vs POST revealed that, during the early processing stage, neural processing is significantly stronger in POST compared to PRE in the STG (p = 0.01). Notably, at the late processing stage this effect was extended to the left prefrontal cortex (PFC) (p < 0.001). Additionally, the comparison between vocoded and clean speech revealed that, at the early processing stage, clean speech elicited stronger neural responses across much of the temporal

lobe ($p < 0.001$) and towards the late stage, significant differences were confined primarily to both left STG and PFC ($p < 0.001$).

This key finding suggests that responses to word onset can serve as an index of speech intelligibility independent of the acoustics. Neural tracking of word onsets represent both bottom-up and top-down processes (18, 36, 37). The early word onset peak may dominantly reflect bottom-up driven mechanisms such as acoustics at word boundaries and automatic word segmentation, but nevertheless does show significant changes after priming. The ~400 ms latency peak, however, is too late to be solely modulated by acoustics and could incorporate higher order word segmentation, semantic integration and other top-down processes (5, 37). When the vocoded speech is unintelligible, neither the words nor word boundaries are clear, resulting in weaker synchronized neural responses at the word onsets. As a result of priming, however, the brain has been provided with additional information (perhaps in the form of new priors) regarding the cues for words, enabling higher intelligibility and the concomitant word boundaries, enabling the emergence of word onset responses. These responses are smaller compared to those of clean speech, as might be expected due to less precise time-locking associated with still-present word boundary uncertainty. The anatomical distribution of the early processing stage, bilateral STG, may reflect a mix of both unaccounted-for auditory responses and also the expected higher-level lexical processing, while the late processing stage's activation of left lateralized PFC may link to the engagement of top-down mechanisms. Such prefrontal activation is consistent with functional magnetic resonance imaging (fMRI) studies using a similar paradigm that found more activation in prefrontal and cingulate cortices with increased speech intelligibility (21). The current study expands on this result by leveraging MEG's superior temporal resolution to show that these processes specifically occur in a time-locked manner, with a ~400 ms post-word-onset latency, and the same polarity as the N400.

One additional analysis was conducted, leveraging context-based speech representations (including those from large language models) that have recently gained popularity in the field of neural speech and language processing and may represent aspects of semantic integration and speech comprehension (2, 37–41). Contextual word surprisal was additionally included as a separate predictor, estimated using a generative pre-trained large language model (GPT-2) (42), which quantifies how surprising a word is given the previous context. Analysis revealed that contextual word surprisal responses show similar effects of intelligibility to those observed in word onset responses for the late processing stage (see Supplemental Figure S1) ($p = 0.03$). The early processing stage, however, was not significantly stronger in POST compared to PRE ($p = 0.93$). It is therefore plausible that the neural mechanisms driving both the word onsets and context-based word surprisal responses may be derived from similar underlying sources, especially at the late stage.

Finally, to strengthen support for the idea that the observed differences in PRE vs POST word onset responses are indeed linked to intelligibility itself and are not just a side effect of passage repetition increasing familiarity, we repeated the analysis on a control study involving nine subjects who listened to the same passages but without any priming, (i.e., PRE1, PRE2, CLEAN). The perceived speech clarity rating analysis did show significant improvement in clarity even without the clean speech priming (increase = 0.58, $SE = 0.12$, $p = 0.001$). However, this effect was considerably smaller than the priming effect (PRE2-PRE1 vs POST-PRE = -1.23, $p_{perm} <$ 0.001), suggesting that priming has a critical impact on speech clarity compared to acoustical learning without priming. The comparison of TRF peak amplitudes in the control study (Supplemental Table S2) demonstrates that acoustic and phoneme onset early (envelope: $p =$ 0.01, onset: $p = 0.43$, phoneme onset: $p = 0.40$) and late peaks (envelope: $p = 0.12$, onset: $p =$ 0.13, phoneme onset: $p = 0.53$) do not differ between PRE1 vs PRE2, but are modulated by the acoustic characteristics for envelope and envelope onset (PRE1 or PRE2 vs CLEAN) (envelope: $p = 0.003$, onset: $p = 0.006$, phoneme onset: $p = 0.63$), similar to the trends observed in the main study. Critically, there was no word onset response change from PRE1 to PRE2 for either the

early (p = 0.15) or late peaks (p = 0.53). Word onset TRF peak amplitudes were significantly larger for CLEAN speech compared to PRE1 or PRE2 (early: p = 0.001, late: p = 0.005). Additionally, the increase in word onset TRF peak amplitudes from PRE to POST in main study were significantly larger than if merely repeated in the control study (PRE2-PRE1 vs POST-PRE: early = -0.30, p_{perm} = 0.02, late = -0.22, p_{perm} = 0.03). These results add further support to the idea that, improvements in intelligibility also generate increased neural responses to lexical segmentation feature over and beyond any acoustical learning.

Even though our analysis revealed effects of intelligibility on the word onset responses, no significant trend was observed between those neural measures and behaviorally-measured perceived speech clarity ratings within subjects. This might have been expected since the neural measures must be calculated across multiple trials, and minimizing the effects of individual trials because of the need to employ cross-validation, whereas the speech clarity did vary substantially from trial to trial (Supplemental Figure S2). It is unfortunate that estimating reliable TRFs from a single trial (in particular, estimating peaks with reliable amplitude and latency) is not feasible for only 20 seconds of data, either for acoustic or linguistic stimulus features. Similarly, the same limitation affects any analysis of the relationship between learning over trials and the word onset processing (i.e., early trails vs late trials).

In conclusion, we investigated the extent to which neural measures of lexical processing correspond to speech intelligibility while keeping the acoustical structure fixed. The neural measures associated with word onset processing, especially those time-locked at N400-like latencies (with N400-like polarity), increased substantially between first exposure and after intelligibility-increasing priming. In contrast, auditory and phoneme onset responses are influenced only by the acoustics of the sensory input, not by intelligibility-boosting priming. It is crucial to exercise caution when interpreting auditory neural responses in contexts where the acoustics of the sensory input differ. Our key finding suggests that lexical segmentation responses increase in the same context as speech intelligibility and show engagement of top-down mechanisms. Together, these suggest that time locked neural responses associated with lexical segmentation may serve as an objective measure of speech intelligibility.

Materials and Methods

Participants

A total of 25 native English speakers (age range 18-32 yr, 15 males, 5 left-handed) participated in the main study. Data from two participants were excluded from the analysis due to excessive artifacts in the neural data. 9 native English speakers (age range19-27 yr, 4 males, 1 left-handed) participated in the control study. All participants reported normal hearing and no history of neurological impairments. The experimental procedures were approved by the University of Maryland institutional review board and all participants provided written informed consent before the experiment and were compensated for their time.

Stimuli and Procedure

The stimuli were ~20 s long excerpts (18 s – 26 s), sampled at 44.1 kHz, from the audiobook "The Botany of Desire" by Michael Pollan, narrated by a male speaker. Talker pauses greater than 400 ms were shortened to 400 ms and the excerpts were then low pass filtered below 4 kHz using a third order elliptic filter.

Noise-vocoded speech segments for each excerpt were generated using a custom python script. First, the frequency range 70-4000 Hz was divided into logarithmically spaced three bands (70 Hz - 432 Hz, 432 Hz - 1402 Hz, and 1402 Hz - 4000 Hz). For each band, an envelope modulated noise band was generated from band-limited white noise modulated by the envelope of the band passed speech signal. The envelope of the band passed speech signal was extracted using the

half-wave rectification of the band passed signal followed by low pass filtering with a cutoff of 30 Hz. Finally, the modulated noise bands were summed, and the normalized volume was adjusted to match that of the original speech stimulus. Figure 1(A) shows example spectrogram plots for the original speech stimulus and vocoded speech stimulus, with the corresponding envelope amplitudes overlaid.

Subjects listened to a total of 36 trials that preserved the storyline. A trial consisted of noise-vocoded speech (PRE), followed by the same speech but in the original clear form (CLEAN) and then the second presentation of noise-vocoded speech (POST). All stimuli were presented diotically. At the end of each noise-vocoded passage, participants were asked to rate the perceived speech clarity ("How much could you follow the passage on a scale of 0 – 5?"; 0 - no words, 1 – a few words, 2 - definitely some words, 3 - lots of words but not most, 4 - more than half of all words, 5 - almost all words). The subjects were specifically instructed not to consider the CLEAN speech passage while making their decisions. This rating was used as a subjective measure of speech clarity. Intermittently, they were also asked to repeat back some of the words they could follow from the vocoded speech passage to ensure that the words they understood were actually from the passage. As a means of motivating the subjects to engage in the task, a comprehension-based question was asked at the end of the POST vocoded speech passage.

In the control study, stimuli, procedures and analyses were identical to those used in the main study, with the exception that the order of each CLEAN and POST vocoded speech were swapped. This resulted in a presentation order of PRE, POST and CLEAN for the control study, which are henceforth described as PRE1, PRE2, and CLEAN, respectively, to emphasize that the clean speech was not presented until after the second presentation of the vocoded speech.

MEG data acquisition and preprocessing

Non-invasive neuromagnetic responses were recorded using a 160 channel whole head MEG system (KIT, Kanazawa, Japan), of which 157 channels are axial gradiometers and 3 magnetometers are employed as environment reference channels, inside a dimly lit, magnetically shielded room (Vacuumschmelze GmbH & Co. KG, Hanau, Germany) at the Maryland Neuroimaging Center. The data were sampled at 1 kHz along with an online low-pass filter with cut off frequency at 200 Hz and a 60 Hz notch filter.

During the task, subjects lay in supine position and were asked to minimize body movements as they listened and to keep their eyes open and fixate on a cross at the center of screen. Sound level was calibrated to ~70 dB sound pressure level (SPL) using 500 Hz tones and equalized to be approximately flat from 40 Hz to 4 kHz. The stimuli were delivered using Presentation software (http://www.neurobs.com), E-A-RTONE 3 A tubes (impedance 50 Ω) which strongly attenuate frequencies above 4 kHz and E-A-RLINK (Etymotic Research, Elk Grove Village, United States) disposable earbuds inserted into the ear canals.

All data analyses were performed in mne-python 0.23.0 (43, 44) and Eelbrain 0.36 (45). Flat channels were excluded and the data were denoised using temporal signal space separation (TSSS) (46). Then the MEG data were filtered between 1 and 60 Hz using a zero-phase FIR filter (mne-python 0.23.0 default settings). Independent component analysis (ICA) (47) was then applied to manually remove artifacts such as eye movements, heart beats, muscle artifacts and singular artifacts. The cleaned data were low pass filtered between 1 and 10 Hz and downsampled to 100 Hz for further analysis.

Neural source localization

The head shape of each participant was digitized using Polhemus 3SPACE FASTRAK three-dimensional digitizer. The position of the participant's head relative to the sensors was determined before and after the experiment using five head-position indicator coils attached to

the scalp surface and the two measurements were averaged. The digitized head shape and the marker coils locations were used to co-register the template FreeSurfer "fsaverage" (48) brain to each participant's head shape using rotation, translation and uniform scaling.

A source space was formed by four-fold icosahedral subdivision of the white matter surface of the fsaverage brain, with all source dipoles oriented perpendicularly to the cortical surface. The source space data and the noise covariance estimated from empty room data were used to compute the inverse operator using minimum norm current estimation (49, 50). The analysis were restricted to frontal, temporal and parietal brain regions based on the 'aparc' FreeSurfer parcellation (51). Excluded brain regions are shaded in dark gray in the brain plots (Figure 2, Figure 3).

Predictor variables

The speech signal was transformed into unique feature spaces to represent different levels of the language hierarchy. These feature-based model predictors can be categorized into three main groups: 1. Acoustic (acoustic envelope and acoustic onsets), 2. sub-lexical (phoneme onset), 3. lexical (word onset, contextual word surprisal). All predictor variables were downsampled to 100 Hz.

Acoustic Properties

The acoustic envelope predictor reflects instantaneous acoustic power, and the acoustic onset reflects the salient transients, of the speech signal. Both of these continuous representations were computed via a simple model of the human auditory system using gammatone filters with Gammatone Filterbank Toolkit 1.0 (52). A filterbank-based broad-band envelope extraction method was used (instead of the conventional broad-band envelope extraction method: absolute value of the Hilbert-transformed signal), as it has been shown that the filterbank-based broad-band envelope increases the neural tracking of the speech envelope (53). First, the gammatone spectrogram was generated with cut-off frequencies from 20 to 5000 Hz, 256 filter channels and 0.01 s window length. Each frequency band was then resampled to 1000 Hz and transformed to log scale. Then the envelope spectrogram was averaged across 256 channels, resulting in a broad-band temporal acoustic envelope predictor.

The acoustic onset representations were computed on the gammatone acoustic envelope spectrogram, by applying an auditory edge detection algorithm (54). Similar to the acoustic envelope, the onset spectrogram was also averaged across frequency bands.

In order to estimate differences in the acoustic feature predictors between clean and vocoded speech, we used the linear-correlation coefficient r. We observed a strong positive correlation between vocoded and clean passages for the envelope ($r = 0.92$, $p < 0.001$), as expected from the method used to construct the vocoded speech. However, this correlation was smaller for the envelope onset ($r = 0.46$, $p < 0.001$), since acoustic onsets are less well preserved than the explicitly copied envelope temporal modulations in the noise-vocoded speech (55).

Sub-lexical Properties

The Montreal Forced Aligner (56) was used to align the speech acoustics with the words and phonological forms from a pronunciation dictionary. The CMU Pronouncing Dictionary (http://www.speech.cs.cmu.edu/cgi-bin/cmudict), excluding the stress information, was used as the pronunciation lexicon. The pronunciation lexicon, transcriptions and audio file were aligned using the 'english' pretrained acoustic model. The annotations for phoneme and word onsets were visualized in PRAAT (57) and manually adjusted appropriately. The phoneme onsets predictor was represented as the impulses at the onset of each phoneme.

Lexical Properties

The word onsets were represented as unit impulses at the onset of each word.

Contextual word surprisal was estimated using an open source transformer-based (58) large language model (GPT-2) (42). We used gpt2-large pretrained models implemented in the Hugging Face environment (59). The transcripts for each passage were preprocessed by removing punctuation and capitalization (places and names were retained). The model first tokenizes the words, where the tokens could represent either words or sub-words, and then fed into the model. The model outputs activation at each of the 36 layers in the network and we used the final layer for the word surprisal calculation. The final layer includes prediction scores given the previous context for each word in the passage over the vocabulary token. Here the 'context' refers to all the preceding tokens or sequence of tokens at least 1024 tokens long. The prediction scores were SoftMax transformed to compute the probability. The current word probability was computed by the probability of the corresponding token and for the words spanning over multiple tokens, the word probability was computed by the joint probability of the tokens. Contextual word surprisal was computed as the $-\log_2(P_{word}|context)$ and represented as an impulse at each word onset scaled by the corresponding model surprisal of that word.

Despite the differences in acoustical features between the vocoded and clean speech passages, sub-lexical and lexical feature time series were kept unchanged.

Computational model

A linear forward modeling approach using Temporal response functions (TRFs) was used to analyze the phase-locked neural responses to various speech features simultaneously (17, 60). Analogous to the conventional event related potentials (ERP), TRFs estimate how the brain responds to speech features over time, or from the signal processing viewpoint, the brain's impulse response to any given speech feature. In contrast to ERPs, that rely on averaging (perhaps) hundreds of short responses to estimate the brain responses to a given stimulus, TRF analysis allows to determine the brain responses to long-duration continuous speech. Critically, TRFs can also model simultaneous responses to multiple speech features (1), multivariate form of the TRFs (mTRF).

Here, TRFs were estimated using the boosting algorithm, which minimizes the $l1$ error between the measured and predicted source current time course, over the time lags -200 ms to 800 ms (using a basis of 50 ms width Hamming windows). Fourfold cross-validation (two training sets, one testing set and one validation set) was used to prevent overfitting and improve the generalized performance. Prediction accuracy was estimated as the explained variance of the TRF model. The subsequent statistical analysis was performed using each subject's average TRFs and prediction accuracies across all cross-validation folds. In sum, optimal TRFs were estimated for each subject, speech condition, and each source current dipole including multiple speech features simultaneously.

Prior estimating TRFs, predictors (speech features) and neural responses were z-score normalized, so that they can be compared between subjects and conditions. To visualize the TRFs over ROI (temporal, frontal and parietal) as a single time series with the current direction, TRFs over ROIs were simplified using principal component analysis (PCA). Examples of the first principal component are shown in Figure 2(A) and Figure 3(A).

TRF peak amplitude extraction

The TRFs showed prominent peaks at different latencies. Based on the TRFs across subjects, we identified time windows of interest and polarity for each peak; envelope TRF: P1 (30-140 ms) and N1 (90-200 ms), envelope onset TRF: P1 (30-140 ms) and N1 (90-200 ms), phoneme onset

TRF: P1 (30-140 ms) and N1 (90-240 ms), word onset and contextual word surprisal TRFs: P1 (50-300 ms) and N1 (330-650 ms), where P1 and N1 represent the polarity of the current estimate, positive and negative respectively, in the time windows specified above. The average TRF response for each subject and condition were obtained as the sum of absolute current dipoles across the ROI. TRF peaks for each subject and condition were picked by searching for the maximum peak in the average TRF, that aligned with the current direction from the original source TRFs. The polarity of the source TRFs were determined by the current direction relative to the cortical surface at the transverse temporal region. If none of the peaks satisfied the polarity constraint, the minimum of the average TRFs in the given time window was used as the peak amplitude.

To aid the TRF peak evaluations, TRF peak amplitude noise floor was measured on noise model TRFs. The noise model TRFs were generated by mismatching the predictors and neural data and estimating the TRFs with the same parameters used in the TRF estimation. These noise model TRFs were then subjected to the same peak picking algorithm as described above to measure the noise model TRF peak amplitudes for each subject. The TRF peak amplitude noise floor for each peak and condition is calculated as the mean noisy peak amplitude across subjects, and represented by dashed lines in the bar plots (See Figure 2 and Figure 3). This noise floor serves as a reference for evaluating the significance of the observed TRF peaks.

Quantification and Statistical Analysis

Statistical analysis was performed in R version 4.0 (61) and Eelbrain. The significance level was set at $\alpha = 0.05$.

Linear mixed effect model (LMEM) analysis was performed to evaluate the trends in behavioral measures. For the LMEM analysis, the *lme4* (version 1.1-30) (62), *lmerTest* (version 3.1-30) (63) and *buildmer* (version 2.4) (64) packages in R were used. The best fit model from the full models were determined using the *buildmer* function. The assumptions of mixed effect modelling, linearity, homogeneity of variance, and normality of residuals, were checked per each best fit model based on the residual plots. Reported effect sizes represent the changes in the dependent measure when comparing one level of an independent variable to its reference level. *p*-values were calculated using Satterthwaite approximation for degrees of freedom (65, 66).

To compare the predictive power of two models, such as the full model vs reduced or PRE vs POST, the difference in explained variability at each source dipole was calculated. Significant differences between the two models were tested while controlling for multiple comparisons, using one-tailed paired-sample *t*-test with threshold-free cluster enhancement (TFCE) (67) and with a null distribution based on 10000 random permutations of the condition labels. The largest *t*-value and the corresponding *p* value are reported in results. To visualize the model fits in meaningful units, they were scaled by the largest explanatory power of the full model across subjects, allowing expression as a % of the full model. The model comparison plots (predictive power of each feature) reported in Figure 2 and Figure 3 are masked by significance to emphasize how each feature contributes to the model.

Lateralization tests were performed to estimate any hemispheric asymmetry in speech feature processing. To accomplish this, the predictive power was first transferred to a common space by morphing the source data to the symmetric "fsaverage_sym" brain, followed by morphing the right hemisphere to the left hemisphere. Once the data were put in the common space, a two-tailed paired sample *t*-test with TFCE was used to test for significant differences between the left and right hemispheres.

TRF peak amplitudes between conditions were compared using paired-sample *t*-test. TRF peak amplitude comparisons between main study and control study were performed using two-sample randomization (permutation) test using *EnvStats* (Version 2.7) (68) R package. The effect sizes

for paired or independent sample *t*-tests were calculated using Cohen's d (69) (d), where d = 0.2 indicates a small effect, d = 0.5 indicates a moderate effect, and d = 0.8 indicates a large effect.

Acknowledgments

This work This work was supported by the National Institutes of Health grant R01-DC019394 and National Science Foundation grant SMA 1734892. We thank Philip Resnik and Shohini Bhattasali for their assistance in choosing context-based linguistic feature and Ciaran Stone for excellent technical assistance.

References

1. C. Brodbeck, L. E. Hong, J. Z. Simon, Rapid Transformation from Auditory to Linguistic Representations of Continuous Speech. *Current Biology* **28**, 3976-3983.e5 (2018).

2. M. Gillis, J. Vanthornhout, J. Z. Simon, T. Francart, C. Brodbeck, Neural Markers of Speech Comprehension: Measuring EEG Tracking of Linguistic Speech Representations, Controlling the Speech Acoustics. *J. Neurosci.* **41**, 10316–10329 (2021).

3. C. Brodbeck, J. Z. Simon, Continuous speech processing. *Current Opinion in Physiology* **18**, 25–31 (2020).

4. E. Ahissar, *et al.*, Speech comprehension is correlated with temporal response patterns recorded from auditory cortex. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 13367–13372 (2001).

5. E. Verschueren, M. Gillis, L. Decruy, J. Vanthornhout, T. Francart, Speech Understanding Oppositely Affects Acoustic and Linguistic Neural Tracking in a Speech Rate Manipulation Paradigm. *J. Neurosci.* **42**, 7442–7453 (2022).

6. J. E. Peelle, J. Gross, M. H. Davis, Phase-Locked Responses to Speech in Human Auditory Cortex are Enhanced During Comprehension. *Cerebral Cortex* **23**, 1378–1387 (2013).

7. H. Luo, D. Poeppel, Phase Patterns of Neuronal Responses Reliably Discriminate Speech in Human Auditory Cortex. *Neuron* **54**, 1001–1010 (2007).

8. Y.-P. Chen, *et al.*, Speech intelligibility changes the temporal evolution of neural speech tracking. *NeuroImage* **268**, 119894 (2023).

9. M. P. Broderick, A. J. Anderson, G. M. Di Liberto, M. J. Crosse, E. C. Lalor, Electrophysiological Correlates of Semantic Dissimilarity Reflect the Comprehension of Natural, Narrative Speech. *Current Biology* **28**, 803-809.e3 (2018).

10. Y. Feng, F. Chen, Nonintrusive objective measurement of speech intelligibility: A review of methodology. *Biomedical Signal Processing and Control* **71**, 103204 (2022).

11. R. V. Shannon, F.-G. Zeng, V. Kamath, J. Wygonski, M. Ekelid, Speech Recognition with Primarily Temporal Cues. *Science* **270**, 303–304 (1995).

12. A. Kösem, B. Dai, J. M. McQueen, P. Hagoort, Neural tracking of speech envelope does not unequivocally reflect intelligibility. *NeuroImage*, 120040 (2023).

13. A. W. Corcoran, *et al.*, Expectations boost the reconstruction of auditory features from electrophysiological responses to noisy speech. *Cerebral Cortex* **33**, 691–708 (2023).

14. E. Sohoglu, M. H. Davis, Perceptual learning of degraded speech by minimizing prediction error. *Proc. Natl. Acad. Sci. U.S.A.* **113** (2016).

15. R. E. Millman, S. R. Johnson, G. Prendergast, The Role of Phase-locking to the Temporal Envelope of Speech in Auditory Perception and Speech Intelligibility. *Journal of Cognitive Neuroscience* **27**, 533–545 (2015).

16. E. C. Lalor, B. A. Pearlmutter, R. B. Reilly, G. McDarby, J. J. Foxe, The VESPA: A method for the rapid estimation of a visual evoked potential. *NeuroImage* **32**, 1549–1561 (2006).

17. N. Ding, J. Z. Simon, Emergence of neural encoding of auditory objects while listening to competing speakers. *Proceedings of the National Academy of Sciences* **109**, 11854–11859 (2012).

18. C. Brodbeck, A. Presacco, J. Z. Simon, Neural source dynamics of brain responses to continuous stimuli: Speech processing from acoustics to comprehension. *NeuroImage* **172**, 162–174 (2018).

19. M. H. Davis, I. S. Johnsrude, A. Hervais-Adelman, K. Taylor, C. McGettigan, Lexical Information Drives Perceptual Learning of Distorted Speech: Evidence From the Comprehension of Noise-Vocoded Sentences. *Journal of Experimental Psychology: General* **134**, 222–241 (2005).

20. C. R. Holdgraf, *et al.*, Rapid tuning shifts in human auditory cortex enhance speech intelligibility. *Nat Commun* **7**, 13654 (2016).

21. M. Hakonen, *et al.*, Predictive processing increases intelligibility of acoustically distorted speech: Behavioral and neural correlates. *Brain Behav* **7**, e00789 (2017).

22. L. S. Baltzell, R. Srinivasan, V. M. Richards, The effect of prior knowledge and intelligibility on the cortical entrainment response to speech. *Journal of Neurophysiology* **118**, 3144–3151 (2017).

23. G. M. Di Liberto, M. J. Crosse, E. C. Lalor, Cortical Measures of Phoneme-Level Speech Encoding Correlate with the Perceived Clarity of Natural Speech. *eNeuro* **5**, ENEURO.0084-18.2018 (2018).

24. M. F. Assaneo, *et al.*, The Lateralization of Speech-Brain Coupling Is Differentially Modulated by Intrinsic Auditory and Top-Down Mechanisms. *Front. Integr. Neurosci.* **13**, 28 (2019).

25. L. Shuai, T. Gong, Temporal relation between top-down and bottom-up processing in lexical tone perception. *Front. Behav. Neurosci.* **8** (2014).

26. L. Fiedler, *et al.*, Single-channel in-ear-EEG detects the focus of auditory attention to concurrent tone streams and mixed speech. *J. Neural Eng.* **14**, 036020 (2017).

27. J. J. Lister, N. D. Maxfield, G. J. Pitt, V. B. Gonzalez, Auditory evoked response to gaps in noise: Older adults. *International Journal of Audiology* **50**, 211–225 (2011).

28. R. Näätänen, I. Winkler, The concept of auditory stimulus representation in cognitive neuroscience. *Psychological Bulletin* **125**, 826–859 (1999).

29. R. Ceponiene, P. Alku, M. Westerfield, M. Torki, J. Townsend, ERPs differentiate syllable and nonphonetic sound processing in children and adults. *Psychophysiology* **42**, 391–406 (2005).

30. N. Ding, J. Z. Simon, Adaptive Temporal Encoding Leads to a Background-Insensitive Cortical Representation of Speech. *J. Neurosci.* **33**, 5728–5735 (2013).

31. B. Zoefel, R. VanRullen, EEG oscillations entrain their phase to high-level features of speech sound. *NeuroImage* **124**, 16–23 (2016).

32. N. Ding, M. Chatterjee, J. Z. Simon, Robust cortical entrainment to the speech envelope relies on the spectro-temporal fine structure. *NeuroImage* **88**, 41–46 (2014).

33. F. Kraus, S. Tune, A. Ruhe, J. Obleser, M. Wöstmann, Unilateral Acoustic Degradation Delays Attentional Separation of Competing Speech. *Trends in Hearing* **25**, 233121652110132 (2021).

34. A. Strauß, S. A. Kotz, J. Obleser, Narrowed Expectancies under Degraded Speech: Revisiting the N400. *Journal of Cognitive Neuroscience* **25**, 1383–1395 (2013).

35. M. Kutas, K. D. Federmeier, Thirty Years and Counting: Finding Meaning in the N400 Component of the Event-Related Brain Potential (ERP). *Annu. Rev. Psychol.* **62**, 621–647 (2011).

36. N. Ding, J. Z. Simon, Cortical entrainment to continuous speech: functional roles and interpretations. *Front. Hum. Neurosci.* **8** (2014).

37. H. Weissbart, K. D. Kandylaki, T. Reichenbach, Cortical Tracking of Surprisal during Continuous Speech Comprehension. *Journal of Cognitive Neuroscience* **32**, 155–166 (2020).

38. M. Heilbron, K. Armeni, J.-M. Schoffelen, P. Hagoort, F. P. de Lange, A hierarchy of linguistic predictions during natural language comprehension. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2201968119 (2022).

39. P. W. Donhauser, S. Baillet, Two Distinct Neural Timescales for Predictive Speech Processing. *Neuron* **105**, 385-393.e9 (2020).

40. J. R. Brennan, J. T. Hale, Hierarchical structure guides rapid linguistic predictions during naturalistic listening. *PLoS ONE* **14**, e0207741 (2019).

41. C. Brodbeck, *et al.*, Parallel processing in speech perception with local and global representations of linguistic context. *eLife* **11**, e72056 (2022).

42. A. Radford, K. Narasimhan, T. Salimans, I. Sutskever, Improving Language Understanding by Generative Pre-Training.

43. A. Gramfort, MEG and EEG data analysis with MNE-Python. *Front. Neurosci.* **7** (2013).

44. A. Gramfort, *et al.*, MNE software for processing MEG and EEG data. *NeuroImage* **86**, 446–460 (2014).

45. C. Brodbeck, P. Das, jpkulasingham, S. Reddigari, T. L. Brooks, Eelbrain 0.36 (2021) https:/doi.org/10.5281/zenodo.5152554.

46. S. Taulu, J. Simola, Spatiotemporal signal space separation method for rejecting nearby interference in MEG measurements. *Phys. Med. Biol.* **51**, 1759–1768 (2006).

47. A. J. Bell, T. J. Sejnowski, An Information-Maximization Approach to Blind Separation and Blind Deconvolution. *Neural Computation* **7**, 1129–1159 (1995).

48. B. Fischl, FreeSurfer. *NeuroImage* **62**, 774–781 (2012).

49. A. M. Dale, M. I. Sereno, Improved Localizadon of Cortical Activity by Combining EEG and MEG with MRI Cortical Surface Reconstruction: A Linear Approach. *Journal of Cognitive Neuroscience* **5**, 162–176 (1993).

50. M. S. Hämäläinen, R. J. Ilmoniemi, Interpreting magnetic fields of the brain: minimum norm estimates. *Med. Biol. Eng. Comput.* **32**, 35–42 (1994).

51. R. S. Desikan, *et al.*, An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* **31**, 968–980 (2006).

52. J. Heeris, Gammatone Filterbank Toolkit (2018).

53. W. Biesmans, N. Das, T. Francart, A. Bertrand, Auditory-Inspired Speech Envelope Extraction Methods for Improved EEG-Based Auditory Attention Detection in a Cocktail Party Scenario. *IEEE Trans. Neural Syst. Rehabil. Eng.* **25**, 402–412 (2017).

54. A. Fishbach, I. Nelken, Y. Yeshurun, Auditory Edge Detection: A Neural Model for Physiological and Psychoacoustical Responses to Amplitude Transients. *Journal of Neurophysiology* **85**, 2303–2323 (2001).

55. S. Rosen, P. Souza, C. Ekelund, A. A. Majeed, Listening to speech in a background of other talkers: Effects of talker number and noise vocoding. *The Journal of the Acoustical Society of America* **133**, 2431–2443 (2013).

56. M. McAuliffe, M. Socolof, S. Mihuc, M. Wagner, M. Sonderegger, Montreal Forced Aligner: Trainable Text-Speech Alignment Using Kaldi in *Interspeech 2017*, (ISCA, 2017), pp. 498–502.

57. P. Boersma, D. Weenink, Praat: doing phonetics by computer (2021).

58. A. Vaswani, *et al.*, Attention is All you Need in *Advances in Neural Information Processing Systems*, I. Guyon, *et al.*, Eds. (Curran Associates, Inc., 2017).

59. T. Wolf, *et al.*, HuggingFace's Transformers: State-of-the-art Natural Language Processing. *arXiv:1910.03771 [cs]* (2020) (April 15, 2022).

60. E. C. Lalor, A. J. Power, R. B. Reilly, J. J. Foxe, Resolving Precise Temporal Processing Properties of the Auditory System Using Continuous Stimuli. *Journal of Neurophysiology* **102**, 349–359 (2009).

61. R Core Team, R: A Language and Environment for Statistical Computing (2020).

62. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using **lme4**. *J. Stat. Soft.* **67** (2015).

63. A. Kuznetsova, P. B. Brockhoff, R. H. B. Christensen, **lmerTest** Package: Tests in Linear Mixed Effects Models. *J. Stat. Soft.* **82** (2017).

64. C. C. Voeten, buildmer: Stepwise Elimination and Term Reordering for Mixed-Effects Regression (2020).

65. S. G. Luke, Evaluating significance in linear mixed-effects models in R. *Behav Res* **49**, 1494–1502 (2017).

66. F. E. Satterthwaite, Synthesis of variance. *Psychometrika* **6**, 309–316 (1941).

67. S. Smith, T. Nichols, Threshold-free cluster enhancement: Addressing problems of smoothing, threshold dependence and localisation in cluster inference. *NeuroImage* **44**, 83–98 (2009).

68. S. P. Millard, *EnvStats: an R package for environmental statistics* (Springer, 2013).

69. J. Cohen, *Statistical power analysis for the behavioral sciences*, 2nd ed (L. Erlbaum Associates, 1988).

Figures and Tables

Figure 1. Experimental methodology, spectro-temporal characteristics of the speech stimuli, behavioral results, and analysis framework. **(A)** (top) Participants listened to 36 audiobook passages in vocoded and clear speech conditions. Each trial consisted of presentation of the same passage in vocoded format, clear format and identical vocoded format. At the end of each vocoded speech passage, participants rated the speech clarity on a scale from 0 to 5. (bottom) Spectrograms and temporal envelopes (overlaid in pink) of vocoded and clear speech. Most spectral and fine temporal details of the vocoded speech are lost (e.g., the pronounced vertical striping), but the broad temporal envelopes of the clear and vocoded speech are very similar. **(B)** Linear Mixed Effects Model (LMEM) predicted speech clarity ratings (0-5) for PRE and POST vocoded conditions. Perceived clarity of the vocoded speech is significantly improved after the clear speech priming. **(C)** Multivariate temporal response function (mTRF) analysis of MEG and predictor variables illustrated with sample of stimuli. Individual TRFs represent the brain's responses to the corresponding speech representations at different time lags. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$

Figure 2. Neural responses to envelope, envelope onset and phoneme onset. **(A)** TRFs from a representative subject in source space, visualized as a single time series using principal component analysis (PCA). The TRFs exhibit an early peak (positive polarity peak marked by ∘) and late peak (negative polarity peak marked by ⊗). **(B, C, D)** (left) Bar plots compare the measured peak amplitudes in the denoted time window (for left and right hemispheres separately) for envelope, envelope onset and phoneme onset respectively. Peak amplitudes were extracted as the maximum peak of the sum of absolute current dipole strengths across sources with a specific polarity, where the polarity was determined from the current directions from the original

source TRFs. (right) brain plots show cortical regions where the given speech feature significantly improves the model fit over and beyond other speech features in the model. Vertical significance brackets indicate significant differences between conditions, and horizontal significance brackets below brain plots indicate lateralization differences. The dashed lines within the bars represent the noise floor, where peaks for a noise model were extracted using the same steps as above.

Figure 3. Neural responses to word onset. **(A)** The word onset TRFs for a representative subject shows two main peaks: an early positive peak (50-300 ms) and a late negative peak (330-650 ms). **(B)** Peak amplitude comparison by speech condition for early and late peaks and by hemisphere. **(C)** Word onset contributions to the prediction accuracy, but separated for the early and late processing stages. Other details as in Figure 2. TRF peak amplitude comparison and prediction accuracy comparisons show that both early and late word onset responses are modulated by intelligibility. At the early processing stage neural responses in POST compared to PRE are stronger in STG and towards late processing stage this effect was extended to left PFC.

Supplementary Information

Table S1. Summary Statistics for TRF peak amplitude comparisons between PRE vs POST, PRE vs CLEAN, and POST vs CLEAN and for both early and late peak (paired t values, p values, and Cohen's d for effect size) in the main study. Differences between CLEAN and either PRE or POST are significant except for phoneme onset early peak. Differences between PRE and POST

		Envelope	Envelope Onset	Phoneme Onset	Word onset
PRE vs POST	Early peak	t_{24} = 0.48, p = 0.63; Cohen's d = 0.03	t_{24} = 0.25, p = 0.80; d = 0.02	t_{24} = -0.52, p = 0.61; d = 0.04	t_{24} = -2.37, p = **0.02;** d = 0.39
	Late Peak	t_{24} = -0.37, p = 0.72; d = 0.04	t_{24} = 1.09, p = 0.29; d = 0.21	t_{24} = -1.03, p = 0.31; d = 0.29	t_{24} = -3.87, p < **0.001;** d = 0.87
PRE vs CLEAN	Early peak	t_{24} = 6.04, p < **0.001;** d = 1.04	t_{24} = -4.28, p < **0.001;** d = 0.70	t_{24} = 1.1, p = 0.28; d = 0.11	t_{24} = -4.77, p < **0.001;** d = 1.34
	Late Peak	t_{24} = -2.37, p = **0.03;** d = 0.54	t_{24} = -3.37, p = **0.003;** d = 0.74	t_{24} = -3.24, p = **0.003;** d = 0.93	t_{24} = -5.21, p < **0.001;** d = 1.20
POST vs CLEAN	Early peak	t_{24} = 6.63, p < **0.001;** d = 1.09	t_{24} = -4.35, p < **0.001;** d = 0.70	t_{24} = -1.16, p = 0.25; d = 0.13	t_{24} = -4.59, p = **0.001;** d = 1.14
	Late Peak	t_{24} = -2.29, p = **0.03;** d = 0.5	t_{24} = -3.32, p = **0.003;** d = 0.83	t_{24} = -3.89, p < **0.001;** d = 0.53	t_{24} = -2.18, p = **0.04;** d = 0.40

are significant but only for word onset responses.

Table S2. Summary Statistics for TRF peak amplitude comparisons between PRE1 vs PRE2, PRE1 vs CLEAN, and PRE2 vs CLEAN and for both early and late peak (paired t values, p values, and Cohen's d for effect size) in the control study. Differences between CLEAN and either PRE1 or PRE2 are significant for early peaks except for phoneme onset. Almost all differences

		Envelope	Envelope Onset	Phoneme Onset	Word onset
PRE1 vs PRE2	Early peak	t_8 = -3.36, p = **0.01;** Cohen's d = 0.30	t_8 = 0.84, p = 0.43; d = 0.16	t_8 = -0.95, p = 0.40; d = 0.12	t_8 = 1.61, p = 0.15; d = 0.36
	Late Peak	t_8 = -1.24, p = 0.12; d = 0.23	t_8 = 1.66, p = 0.13; d = 0.54	t_8 = 0.61, p = 0.53; d = 0.16	t_8 = -0.66, p = 0.53; d = 0.14
PRE1 vs CLEAN	Early peak	t_8 = 4.21, p = **0.003;** d = 0.93	t_8 = -3.75, p = **0.006;** d = 0.90	t_8 = -0.44, p = 0.63; d = 0.05	t_8 = -4.86, p = **0.001;** d = 1.35
	Late Peak	t_8 = -1.57, p = 0.15; d = 0.69	t_8 = -1.21, p = 0.26; d = 0.46	t_8 = -1.50, p = 0.17; d = 0.43	t_8 = -3.60, p = **0.007;** d = 1.15
PRE2 vs CLEAN	Early peak	t_8 = 4.13, p = **0.003;** d = 1.08	t_8 = -3.49, p = **0.008;** d = 1.00	t_8 = 0.53, p = 0.61; d = 0.08	t_8 = -4.94, p = **0.001;** d = 1.46
	Late Peak	t_8 = -1.15, p = 0.28; d = 0.48	t_8 =-1.93, p = 0.08; d = 0.69	t_8 = -1.54, p = 0.16; d = 0.58	t_8 = -3.80, p = **0.005;** d = 1.10

between PRE1 and PRE2 are not significant.

Figure S1. TRF peak amplitude comparison by speech condition for model including contextual word surprisal for envelope, envelope onset, phoneme onset, word onset and contextual word surprisal, for early and late peaks, and by hemisphere. The dashed lines within the bars represent the noise floor. Similar to the main model, almost all differences between CLEAN and either PRE or POST are significant, except for phoneme onset early peak. Differences between PRE and POST are significant only for word onset responses and contextual word surprisal late responses in the left hemisphere. Adding contextual word surprisal into the model does not substantially change significant intelligibility effects observed in word onset responses. *p<0.05, **p<0.01, ***p<0.001

Figure S2. Speech clarity rating change over trials. Trials were combined using a tumbling window of 6 trials. Linear Mixed Effects Model (clarity rating ~ 1 + condition + trial group (1+condition|Subject)) predicted clarity rating ± SE change over trial group. As no significant interaction was found between condition and trial group, the model's trends over trials for both PRE and POST are parallel. Overall, speech clarity improves over trials for both PRE and POST vocoded speech (trials 1-6 as the reference, trials 7-12 = 0.53, SE = 0.06, p < 0.001, trials 13-18 = 0.66, SE = 0.06, p < 0.001, trials 19-24 = 0.68, SE = 0.06, p < 0.001, trials 25-30 = 0.91, SE = 0.06, p < 0.001, trials 31-36 = 0.79, SE = 0.06, p < 0.001) and then level off (trials 19-24 to trials 25-30 = 0.15, SE = 0.06, p = 0.33). *p<0.05, **p<0.01, ***p<0.001