Neurophysiological trajectories in Alzheimer's disease progression

Kiwamu Kudo[1,2†*]**, Kamalini G. Ranasinghe**[3†]**, Hirofumi Morise**[1,2]**, Faatimah Syed**[3]**,
Kensuke Sekihara[4]**, Katherine P. Rankin**[3]**, Bruce L. Miller**[3]**, Joel H. Kramer**[3]**,
Gil D. Rabinovici[3,5]**, Keith Vossel**[3,6]**, Heidi E. Kirsch**[1]**, Srikantan S. Nagarajan**[1]

*For correspondence:
kiwamu.kudo@jp.ricoh.com (KK)

[†]These authors contributed
equally to this work

[1] Biomagnetic Imaging Laboratory, Department of Radiology and Biomedical Imaging,
University of California, San Francisco, CA, 94143, USA ; [2] Medical Imaging Business
Center, Ricoh Company, Ltd., Kanazawa, 920-0177, Japan ; [3] Memory and Aging Center,
UCSF Weill Institute for Neurosciences, University of California, San Francisco, CA,
94158, USA. ; [4] Signal Analysis Inc., Hachioji, Tokyo, 192-0031, Japan ; [5] Department of
Radiology and Biomedical Imaging, University of California, San Francisco, CA, 94143,
USA ; [6] Mary S. Easton Center for Alzheimer's Research and Care, Department of
Neurology, David Geffen School of Medicine, University of California Los Angeles, Los
Angeles, CA 90095, USA

Abstract   Alzheimer's disease (AD) is characterized by the accumulation of amyloid-β and
misfolded tau proteins causing synaptic dysfunction and progressive neurodegeneration and
cognitive decline. Altered neural oscillations have been consistently demonstrated in AD.
However, the trajectories of abnormal neural oscillations in AD progression and their relationship
to neurodegeneration and cognitive decline are unknown. Here, we deployed robust event-based
sequencing models (EBMs) to investigate the trajectories of long-range and local neural
synchrony across AD stages, estimated from resting-state magnetoencephalography. Increases
in neural synchrony in the delta-theta band and decreases in the alpha and beta bands showed
progressive changes along the EBM stages. Decreases in alpha and beta-band synchrony
preceded both neurodegeneration and cognitive decline, indicating that frequency-specific
neuronal synchrony abnormalities are early manifestations of AD pathophysiology. The
long-range synchrony effects were greater than the local synchrony, indicating a greater
sensitivity of connectivity metrics involving multiple regions of the brain. These results
demonstrate the evolution of functional neuronal deficits along the sequence of AD progression.

Introduction

Alzheimer's disease (AD) is a neurodegenerative disease characterized by amyloid-β (Aβ) and neu-
rofibrillary tangles of abnormally phosphorylated tau (***DeTure and Dickson*** (***2019***)). Clinical and
epidemiological studies have suggested that Aβ accumulation occurs early in the timeline of neu-
ropathological changes in AD, likely preceding the accumulation of tau, and subsequent neurode-
generation and cognitive decline (***Jack Jr et al., 2010***; ***Sperling et al., 2011***). The neuropathological
changes of AD are therefore described as a continuum, starting from the presymptomatic stage of
proteinopathy and continuing to progress during the symptomatic stage with increasing stages of
disease severity (***Sperling et al., 2011***; ***Jack Jr et al., 2018***). Transgenic mouse models of AD have
shown that AD proteinopathy of Aβ and tau is associated with synaptic and circuit dysfunctions
in neural networks (***Busche et al., 2008***; ***Ahnaou et al., 2017***; ***Busche et al., 2019***). However, the

temporal change in synaptic and circuit dysfunction along disease progression in patients with AD remains largely unknown.

Functional deficits in neural networks, especially in the presymptomatic stage, have attracted attention in recent years with the rapidly evolving landscape of plasma-biomarkers of early detection and the novel therapeutics showing the benefits of early intervention (*Dubois et al., 2016*). In fact, abnormal synchrony of neural oscillations has been reported not only in patients along the clinical spectrum of AD, including mild cognitive impairment (MCI) due to AD and AD-dementia (*Jeong, 2004*; *Fernández et al., 2006*; *Stam et al., 2006*; *Koelewijn et al., 2017*; *Nakamura et al., 2018*; *Hughes et al., 2019*; *Ranasinghe et al., 2020*; *Meghdadi et al., 2021*; *Schoonhoven et al., 2022*) but also during the preclinical stages of AD (*Nakamura et al., 2018*; *Ranasinghe et al., 2022a*). Neuronal oscillations observed by noninvasive electrophysiological measures, such as electroencephalography (EEG) and magnetoencephalography (MEG), represent the synchronized activity of excitatory and inhibitory neurons and thus provide sensitive indices of altered neuronal and circuit functions in AD. As synaptic dysfunction is strongly associated with AD proteinopathy, altered neural oscillation synchrony may capture early functional deficits of neural networks even before clinical symptoms appear. However, it remains unknown which neurophysiological signature changes capture such deficits and the temporal evolution of these changes along the timeline of preclinical to MCI to AD dementia stages in clinical populations.

In this study, we investigated the trajectories of neurophysiological changes along the course of clinical AD progression, by examining long-range and local neural synchrony patterns in the resting brain. We hypothesized that frequency-specific long-range and local synchrony abnormalities in neuronal oscillations may precede both neurodegeneration and cognitive deficits. To examine the temporal relationship amongst altered neural synchrony, neurodegeneration, and cognitive deficits, we used data-driven disease progression models, specifically event-based sequencing models (EBMs), which have been successfully used to predict AD progression from cross-sectional biomarker data (*Fonteijn et al., 2012*; *Young et al., 2014*, *2018*). In an EBM, disease progression is described as a series of discrete events, defined as the occurrence of a particular biomarker reaching a threshold abnormal value, and the estimated likelihood of temporal sequence of events defines the disease trajectory. Modifying the conventional EBMs, to find the neurophysiological trajectories, we developed a robust EBM framework that is less sensitive to the thresholds for determination of abnormality thereby resulting in unbiased estimation of disease stage probability for each study participant.

We considered two representative neuronal oscillatory synchrony metrics: amplitude-envelope correlation (AEC) and regional spectral power. The AEC and spectral power quantify long-range and local neural synchrony, respectively. Recent test-retest studies of MEG resting-state metrics have revealed that both metrics are highly reliable (*Colclough et al., 2016*; *Wiesman et al., 2021a*). To evaluate the frequency specificity of neurophysiological trajectories, three canonical frequency bands, delta-theta, alpha, and beta bands, were considered. For a metric of global cognitive ability, we used the mini-mental state examination (MMSE) score. Neurodegeneration, which is related to neuronal loss as well as synaptic loss and synapse dysfunction (*Selkoe, 2002*; *Spires-Jones and Hyman, 2014*), is detectable as brain atrophy on structural MRI, and therefore we evaluated neurodegeneration as loss of grey matter (GM) volume, specifically volume loss of the parahippocampal gyrus (PHG), extracted from individual T1 MRIs. We first deployed an Atrophy-Cognition EBM (AC-EBM) with only the neurodegeneration and cognitive decline measures, and then quantitatively examined metrics of long-range and local synchrony of neuronal oscillations corresponding to each estimated disease stage. Next, we deployed two separate Synchrony-Atrophy-Cognition EBMs (SAC-EBMs) which respectively included long-range or local neural synchrony measures along with PHG volume and global cognition, and investigated how the synchrony metrics stratify AD progression. Consistent with our hypothesis, we found that long-range and local neural synchrony in the alpha and beta bands, but not in the delta-theta band, becomes abnormal at the earliest preclinical stages of AD, preceding both neurodegeneration and cognitive deficits.

Materials and methods

Participants

The present study included 78 patients who met National Institute of Aging–Alzheimer's Association (NIA-AA) criteria (*McKhann et al., 2011*; *Albert et al., 2011*; *Jack Jr et al., 2018*), and 70 cognitively-unimpaired older adults. All participants were recruited from research cohorts at UCSF Alzheimer's Disease Research Center (UCSF-ADRC). Diagnosis of AD patients was established by consensus in a multidisciplinary team. Amongst the 78 AD patients, 20 had autopsy confirmed AD neuropathology, another 41 patients were positive on the Aβ-PET scans, plus another 9 patients showed cerebrospinal fluid (CSF) assays of amyloid and tau levels consistent with an AD diagnosis. The remaining 8 patients were clinically diagnosed, based on clinical evaluations and the characteristic pattern of cortical atrophy on MRI. The control participants were recruited from an ongoing longitudinal study of healthy aging at UCSF-ADRC. The eligibility criteria for cognitively normal controls included normal cognitive performance, normal MRI, and absence of neurological, psychiatric, or other major medical diseases. Forty-seven (out of 70) controls were evaluated with Aβ-PET and 8 were read as positive (39 as negative). The remaining 23 control participants were not evaluated with Aβ-PET. All participants underwent Mini Mental State exam (MMSE) and a structured caregiver interview to assess the clinical dementia rating scale (CDR). All control participants were identified at CDR 0 indicating cognitively unimpaired status on the CDR scale. Patients with AD ranged from 0.5 to 2 on the CDR scale. The results from demographic, functional, and cognitive assessments are shown in Appendix 1—table 1. Informed consent was obtained from all participants or their assigned surrogate decision makers. The study was approved by the Institutional Review Board of UCSF.

MRI acquisition and analyses

Structural brain images were acquired using a unified MRI protocol on 3T Siemens MRI scanners (MAGNETOM Prisma or 3T TIM Trio) at the Neuroscience Imaging Center (NIC) at UCSF, within an average of 1.05 years (range: −6.91–0.78) and 0.29 years (range: −2.13–1.29) of the MEG evaluation for controls and patients, respectively. The acquired MRI was used to generate the head model for source reconstructions of MEG sensor data and to evaluate GM volumes. The region-based GM volumes corresponding to the 94 anatomical regions included in the automated anatomical labelling 3 (AAL3) atlas (*Rolls et al., 2020*) (Appendix 1—table 2) were computed using the Computational Anatomy Toolbox [CAT12 version 12.8.1 (1987)] (*Gaser et al., 2022*), which is an extension of SPM12 (*Penny et al., 2011*). The regional GM volumes were evaluated by the morphometry pipeline implemented in CAT12 with default parameters. The total intracranial volume (TIV), the sum of all segments classified as gray and white matter, and CSF, was also calculated for each subject.

Resting-state MEG

Data acquisition

Each participant underwent 10–60-minutes resting-state MEG at the UCSF Biomagnetic Imaging Laboratory (BIL). MEG was recorded with a 275 channel full head CTF Omega 2000 system (CTF MEG International Services LP, Coquitlam, British Columbia, Canada). Three fiducial coils including nasion and left and right pre-auricular points were placed to localize the position of head relative to the sensor array and later co-registered with individual MRI to generate an individualized head shape. Data collection was optimized to minimize head movements within the session and to keep it below 0.5 cm. For analysis, a 10-min continuous recording was selected from each subject lying supine and awake with the eyes closed (sampling rate $f_s = 600$ Hz). From the continuous recordings, we further selected a 1-min continuous segment with minimal artifacts (i.e., minimal excessive scatter at signal amplitude) for each subject.

Pre-processing

Each 1-min sensor signal was digitally filtered using a bandpass filter of $0.5–55\,\text{Hz}$. The power spectral density (PSD) of each sensor signal was computed, and artifacts were confirmed by visual inspections. Channels with excessive noise within individual subjects were removed prior to the next process. When environmental noises larger than a few $\text{pT}/\sqrt{\text{Hz}}$ were observed around 1–5-Hz range in a PSD, the dual signal subspace projection (DSSP) (*Sekihara et al., 2016*) with the lead-field vectors computed for each individual subject's head model was applied to the filtered sensor signal for the removal of environmental noises. As a parameter, we chose the dimension of pseudo-signal subspace μ as 50. DSSPs were needed to be applied to thirteen of the total 148 subject's signals. For the 13 data, the resulting dimension of the spatio-temporal intersection, i.e., the degree of freedom to be removed, was 3 or 4. We also applied a preconditioned independent component analysis (ICA) (*Ablin et al., 2018*) to the signal to identify cardiac components and remove them. In each data set, one or two clear cardiac ICA-component waveforms with approximately 1 Hz rhythms were observed, which were easily identified by visual inspections.

Atlas-based source reconstruction

Isotropic voxels (5 mm) were generated in a brain region of a template MRI, resulting in $15,448$ voxels within the brain region. The generated voxels were spatially normalized to individual MRI space, and subject-specific magnetic lead field vectors were computed for each voxel with a single-shell model approximation (*Nolte, 2003*). The voxels for each subject were indexed to 94 cortical/ sub-cortical regions included in the AAL3 atlas.

Array-gain scalar beamformer (*Sekihara et al., 2004*) was applied to the 60-sec cleaned sensor time series to obtain source-localized brain activity at the voxel level, i.e., voxel-level time courses. Lead field vectors were normalized to avoid the center-of-the-head artifact, and a generalized eigenvalue problem was solved to determine the optimal source orientation (*Sekihara and Nagarajan, 2008*). The beamformer weights were calculated in the time domain; a data covariance matrix was calculated using a whole 60-sec time series, and a singular value truncation (threshold of $10^{-6} \times$ maximum singular value) was performed when inverting the covariance matrix. Ninety-four regional time courses were extracted with alignment with the AAL3 atlas by performing principal component analysis (PCA) across voxel-level time courses within each of the regions and taking a time course of the first principal component. These pre-processing and source reconstructions were performed using in-house Matlab scripts utilizing Fieldtrip toolbox functions (*Oostenveld et al., 2011*). We also used BrainNet Viewer toolbox (*Xia et al., 2013*) to obtain brain rendering images of regional MEG metrics and GM atrophy.

MEG resting-state metrics

Based on the regional time courses derived from MEG, we evaluated two measures of neural synchrony: the amplitude-envelope correlation (AEC) and the spectral power, which describe long-range and local neural synchrony, respectively. Three canonical frequency bands were considered: $2–7\,\text{Hz}$ (delta-theta), $8–12\,\text{Hz}$ (alpha), and $15–29\,\text{Hz}$ (beta) bands (Appendix 1—figure 1).

Amplitude-envelope correlation

The AECs are defined as Pearson's correlation coefficients (PCCs) between any two amplitude envelopes of regional time courses (total $94 \times 93/2 = 4,371$ pairs). Regional time courses were first processed by a band-pass filtering and then their envelopes were extracted by Hilbert transform. To discount spurious correlations caused by source leakages, we orthogonalized any two band-limited time courses before computing their envelopes by employing a pairwise orthogonalization (*Hipp et al., 2012*; *Sekihara and Nagarajan, 2015*). The AEC with leakage correction is often expressed as AEC-c and is known as a robust measure (*Briels et al., 2020b*). The pairwise orthogonalization provides asymmetric values between two time courses; the value depends on which time course is taken as a seed. Therefore, PCCs between orthogonalized envelopes for both di-

187 rections were averaged, resulting in a symmetric AEC matrix. Regional AECs, that represent the
188 connectivity strengths of each ROI, were computed by averaging over row/column components of
189 the symmetric AEC matrix.

Spectral power

191 The spectral power of a given band, which has often been used as a metric to discriminate patients
192 with AD from controls (*Jeong, 2004*; *Engels et al., 2016*; *Wiesman et al., 2021b*), is defined by the
193 ratio of a band power to total power and was calculated from regional PSDs. Regional PSDs were
194 calculated from the 94 regional time courses using Welch's method (50% overlap) with 0.293-Hz
195 ($= f_\mathrm{s}/2048$) steps.

Scalar neural synchrony metrics

197 To identify general trends in changes in the long-range and local synchrony with the severity of AD,
198 we performed group comparisons of the regional synchrony metrics between AD patients and con-
199 trols. Based on the group contrasts of regional metrics observed, we introduced scalar synchrony
200 metrics by calculating the averages within several regions where large region-level group contrasts
201 were identified. The scalar MEG metrics were used in the SAC-EBMs.

Metric trajectory analyses

Event-based sequencing modeling

Imaging and neuropsychological biomarkers for AD are continuous quantities taking values from
normal to severe, while the stages of the disease are discrete and are identified by estimating the
values of the biomarkers (*Sperling et al., 2011*). As a data-driven disease progression model, an
event-based sequencing model (EBM) has been proposed that allows us to make inferences about
disease progression from cross-sectional data (*Fonteijn et al., 2012*; *Young et al., 2014*, *2018*). In
an EBM, disease progression is described as a series of metric events, where events are defined
as the occurrences of abnormal values of metrics, and the values of events act as thresholds to
determine discrete stages of disease (*Fonteijn et al., 2012*). The model infers temporal sequences
of the events from cross-sectional data.

It is also possible to set multiple events per metric by defining them as occurrences of taking
certain z-scores within the range from initial to final z-scores ([z_initial z_final]), in which z-scores for
each metric linearly increase between all consecutive events and stages are located at temporal
midpoints between the two consecutive event occurrence times (*Young et al., 2018*). In this lin-
ear z-score event model, a metric trajectory is described as a series of metric values evaluated at
estimated stages.

We developed a robust EBM framework to quantify metric trajectories on the basis of the linear
z-score model, employing the following form of a data likelihood:

$$P(Z|S) = \prod_{j=1}^{J} \sum_{k=1}^{N+1} p(t_j = k)p(Z_j|S, t_j = k), \qquad (1)$$

where N denotes a total number of events, and i, j, and k are the indices of metric, subject, and
stage, respectively. J is the number of subjects ($J = 148$). I is the number of metrics: $I = 2$ for
an AC-EBM and $I = 3$ for an SAC-EBM, respectively. The symbol t_j denotes stages for each subject
j, and a conditional probability, $p(Z_j|S, t_j = k)$, describes the probability that a subject j takes
biomarker values of Z_j given a sequence of events S and that $t_j = k$ (i.e., the subject j is in a stage
k). The symbol $Z_j = [z_{1j}, z_{2j}, \ldots, z_{Ij}]^\mathrm{T}$, where z_{ij} denotes the z-score of a metric i for a subject j, and
the symbol $Z = [Z_1, Z_2, \ldots, Z_J]$ describing the data matrix with the $I \times J$ dimension. Since there
are $N + 2$ event occurrence times including initial and final times, $N + 1$ stages are provided. When
employing *Equation 1*, we assumed that the prior distribution that the subject j is in a stage k is
uniform: $p(t_j = k) = (N + 1)^{-1}$. We also assumed that the prior probability of $p(Z_j|S, t_j = k)$ arises

from independent Gaussian distributions for each metric i resulting in a factorized multivariate prior. Hence,

$$p(Z_j|S, t_j = k) \propto \prod_{i=1}^{I} \exp\left(-\frac{(z_{ij} - \mu_i(k))^2}{2}\right).$$ (2)

The symbol $\mu_i(k)$ denotes a value of the z-score of a metric i at a stage k and is given by a linearly interpolated midpoint z-score between two z-scores evaluated at consecutive event occurrence times. The goal of this formulation is to evaluate the posterior distribution that a subject j belongs to a stage k, $p(t_j = k|Z_j, \bar{S})$, with the most likely order of events $\bar{S}$.

The most likely order of events is given by the sequence of events, S, which maximizes the posterior distribution $P(S|Z) = P(S)P(Z|S)/P(Z)$. On the assumption that the prior $P(S)$ is uniformly distributed (*Fonteijn et al., 2012*), the most likely sequence is obtained by solving the maximum likelihood problem of maximizing *Equation 1*. To solve the problem, for a given set of events, we performed Markov chain Monte Carlo (MCMC) sampling on sequences and chose the maximum likelihood sequence from $50,000$ MCMC samples. In the generation of the MCMC samples, we initialized the MCMC algorithm with an initial sequence close to or equal to the maximum likelihood solution by running a ascent algorithm 10 times from different initialization points, i.e., randomly generated event sequences (*Fonteijn et al., 2012*).

z-scoring of metrics

We computed z-scores of the PHG volume, MMSE score, and scalar neural synchrony metrics to utilize them in the EBM frameworks. Since a linear z-score model assumes a monotonous increase in z-scored metrics along disease progression (i.e., higher stage denotes more severity), "sign-inverted" z-scores were introduced to the metrics with decreasing trends along disease progression. Specifically, for GM volumes, MMSE score, and neural synchrony metrics in the alpha and beta bands, the z-score of a metric i for a subject j was defined by $z_{ij} = (\bar{x}_i^C - x_{ij})/\sigma_i^C$, where x_{ij} denotes a value of a metric i for a subject j, and $\bar{x}_i^C$ and σ_i^C denote the mean and standard deviation (SD) of the metric values of the controls, respectively. For the delta-theta-band neural synchrony metrics, z-scores were defined in a standard way as $z_{ij} = (x_{ij} - \bar{x}_i^C)/\sigma_i^C$. Using these z-scored metrics, the initial and final events, z_{initial} and z_{final}, for each metric were set as the bottom and top 10% average z-scores, respectively.

Events-setting optimization

In addition to the initial and final events of the z-score, we set three events for each metric because various possible curves of the metric trajectories were supposed to be well expressed by three variable points with two fixed points. For example, in an AC-EBM analysis, that is, a two-metric trajectory analysis for PHG volume loss and MMSE decline, a total of six events were considered ($N = 6$). The metric trajectory as a series of stage values $\mu_i(k)$ is sensitive to event settings because predefined events do not necessarily capture appropriate boundaries between disease stages. To determine disease stages less sensitive to specifications of the z-score events, we tried several sets of events and selected the set of events with the largest data likelihood amongst the trials. Specifically, we searched for the set of events that better fits the data Z by trying all combinations of three z-scores from $\{0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8\}$-quantiles for each metric. The number of combinations of events for each metric was accordingly ($_7C_3 =$)35. Therefore, MCMC samplings ($50,000$ samples for each set of events) were performed $1,225$ times for an AC-EBM and $42,875$ times for an SAC-EBM, respectively, to find the set of events and their sequence $\bar{S}$ with the highest data likelihood. This exhaustive search of optimal event settings, which was not implemented in a conventional linear z-score EBM (*Young et al., 2018*), is diagrammatically shown in Appendix 1—figure 2.

While it is tractable to directly evaluate $P(Z|S)$ for all ordered arrangements of z-score events when the number of the permutations for each set of z-score events is just $20(= {_6C_3})$ for $I = 2$ (Appendix 1—table 3) and $1,680(= {_9C_3} \times {_6C_3})$ for $I = 3$, this procedure is not tractable when $I > 3$ and requires MCMC sampling procedures. Furthermore, MCMC enables computation of the sequence

278 statistics. Therefore, we use MCMC sampling from which we could compute the positional variance
279 estimates for each event (see Appendix 1—figure 2 and Appendix 1—figure 3).

Trajectory computations

281 Given the most likely sequence $\bar{S}$ as a result of the exhaustive search, the probabilities that a
282 subject j falls into a stage k are evaluated by the posterior distribution:

$$p(t_j = k|Z_j, \bar{S}) = \frac{p(Z_j|\bar{S}, t_j = k)}{\sum_{k'} p(Z_j|\bar{S}, t_j = k')}. \tag{3}$$

283 These probabilities, $p_j(k) \equiv p(t_j = k|Z_j, \bar{S})$, describe the contribution of a subject j to stage k, which
284 allows us to evaluate a stage value of any metric x of i at a stage k as a weighted mean:

$$\bar{x}_i(k) = \frac{\sum_{j=1}^{J} p_j(k) \cdot x_{ij}}{\sum_{j=1}^{J} p_j(k)} \tag{4}$$

285 Then, we represented the trajectory of the metric i by a series of the stage values, $\bar{x}_i(k)$. The stan-
286 dard error (SE) of the weighted mean at stage k was evaluated by

$$\mathrm{SE}(k) = \sigma_i \cdot \sqrt{\frac{\sum_{j=1}^{J} p_j(k)^2}{\left(\sum_{j=1}^{J} p_j(k)\right)^2}}, \tag{5}$$

where σ_i is a standard deviation of a metric i. This definition of SE provides an usual expression of
the standard error of the mean, $\sigma_i/\sqrt{J}$, if all subjects contributed equally to all stages.
These formulations of trajectories were applied to several metrics. In the AC-EBM, the metrics
i denote the PHG volume loss z-score and the MMSE scores. In the SAC-EBM, they denote each
scalar neural synchrony metric in addition to the PHG volume loss z-score and the MMSE score.
We also used *Equation 4* to evaluate the progressions of the regional neural synchrony metrics
and the GM volume loss z-scores along the estimated EBM stages. When evaluating the ratio of
subjects categorized to each stage, we treated $(x_{i1}, x_{i2}, \ldots, x_{iJ})$ as an one-hot vector, where a metric
i represents a category of subjects provided by CDR scale. For example, when evaluating the ratio
of subject with CDR 0.5, $x_{ij} = 1$ only when a subject j has CDR scale of 0.5, otherwise $x_{ij} = 0$.

Statistical analyses

To test demographic differences between AD patients and controls, unpaired t-test was used for
age, and chi-square test was used for sex. Age was defined at the time of the MEG scan date. In
the statistical analyses, p-values below 0.05 were considered statistically significant. For group com-
parisons of GM volumes, MMSE scores, and neural synchrony metrics, two-sided significance tests
(against a null value of zero) were performed using the general linear model (GLM). For statistical
tests on GM volumes, TIV, age, and the difference between MRI and MEG dates were included as
covariates. For statistical tests on MMSE scores, age and the difference between the MMSE and
MEG dates were included as covariates. For statistical tests on neural synchrony metrics, age was
included as a covariate. The problem of multiple comparisons between 94 regions was solved
by controlling the Benjamini-Hochberg false discovery rate (FDR) (*Benjamini and Hochberg, 1995*).
The FDR adjusted p-value (i.e., q-value) below 0.05 or 0.01 was considered statistically significant.
A nonparametric test was performed to statistically compare metrics between stages, i.e., to
test statistical significance of the difference between stage values represented by weighted means
[e.g., stage k vs. k': $\delta x = \bar{x}_i(k) - \bar{x}_i(k')$]. For a metric i, we used bootstrap resampling (50, 000 samples)
of an original data set, $\boldsymbol{x}_i = (x_{i1}, x_{i2}, \ldots, x_{iJ})$, to generate new data sets, $(x_{i1}^{\star}, x_{i2}^{\star}, \ldots, x_{iJ}^{\star})$, using a
random number generator, where each $x_{ij}^{\star}$ is one of the components of the original data set $\boldsymbol{x}_i$.
We then calculated the weighted means $\bar{x}_i^{\star}(k)$ (*Equation 4*) for each sample. The same procedures
were performed for stage k', obtaining weighted means $\bar{x}_i^{\star}(k')$ for each sample. We then tested
the null hypothesis that a weighted mean in stage k, $\bar{x}_i(k)$, is equal to a weighted mean in stage k',

$\bar{x}_i(k')$, evaluating the null distribution of differences in weighted mean values, $\delta x^{\star} = \bar{x}_i^{\star}(k) - \bar{x}_i^{\star}(k')$. The problem of multiple comparisons across stages was solved by controlling the FDR. The q-value below 0.05 was considered statistically significant.

Results

Participant demographics

This study included a cohort of 78 patients with AD (50 female; 28 male) including 35 patients with AD dementia and 43 patients with MCI due to AD, and also included 70 cognitively-unimpaired older adults as controls (41 female; 29 male). The CDR scales were 0 for the cognitively-unimpaired controls, 0.5 for patients with MCI, and 1 ($n = 27$) or 2 ($n = 8$) for patients with AD dementia. There was no difference in sex distribution between the AD and control groups [$\chi^2(1) = 0.477$; $p = 0.49$]. Average age at the time of the MEG was slightly higher in the control group than patients with AD (controls, mean$\pm$SE: 70.5 ± 0.99, range: 49.5–87.7; AD, mean$\pm$SE: 63.9 ± 1.01, range: 49.0–84.4) [unpaired t-test: $t(146) = -4.708$; $p < 0.001$]. Average MMSE in patients with AD was 22.7 ± 0.43 (mean$\pm$SE) while the average MMSE in the controls 29.2 ± 0.48. The MMSE scores were adjusted for age and the time differences between MMSE administration and MEG scan using a GLM (Appendix 1—figure 4B). MMSE-decline z-scores, z_{MMSE}, were standardized by the adjusted MMSE scores of the control group and sign-inverted (*Figure 1*B).

Group comparisons of GM volumes for each of the anatomical regions included in the AAL3 atlas showed that GM volumes in the temporal regions are significantly smaller in AD patients than in controls (Appendix 1—figure 5; Appendix 1—table 4). Amongst the temporal GM volumes, we focused on a volume of PHG as a key indicator of neurodegeneration in AD progression. The PHG includes perirhinal and entorhinal cortices of the medial temporal lobe (MTL), and MRI-based studies have reported that the volume of MTL reduces, especially in the perirhinal and entorhinal cortices, in early stages of typical AD (*Teipel et al., 2006*; *Echávarri et al., 2011*; *Matsuda, 2016*). In this study, PHG volume was defined as a summation of the volumes of left- and right-hemisphere PHGs. The average PHG volume in patients with AD ($7.99 \, \mathrm{ml} \pm 0.09$) was significantly smaller than in the controls ($9.28 \, \mathrm{ml} \pm 0.11$) [unpaired t-test: $t(143) = -9.508$; $^{***}p < 0.001$] (Appendix 1—figure 4A); the PHG volumes were adjusted for TIV, age, and the difference between MRI and MEG dates by including them in a GLM as covariates. PHG volume loss z-scores, z_{PHG}, were standardized by the adjusted PHG volumes of the control group and sign-inverted (*Figure 1*A).

Abnormal frequency specific long-range and local neural synchrony in AD

For regional long-range synchrony (AEC), increases in delta-theta-band synchrony in patients with AD were identified in frontal regions, and reductions in alpha- and beta-band synchrony were identified in the whole brain (Appendix 1—figure 6C, E; Appendix 1—table 5). These regional contrasts were similar to those observed between AD dementia and subjective cognitive decline (SCD) in MEG/EEG studies (*Schoonhoven et al., 2022*; *Briels et al., 2020a*). For regional local synchrony (spectral power), increases in delta-theta-band power in patients with AD were identified in the whole brain, and reductions in alpha- and beta-band power were identified in temporal regions and the whole brain, respectively (Appendix 1—figure 6D, F; Appendix 1—table 6). These regional contrasts were similar to those observed between MCI and controls in a multicenter study of MEG (*Hughes et al., 2019*).

Based on the group contrasts of regional metrics observed, we introduced six scalar metrics to quantify long-range and local synchrony: [i] frontal delta-theta-band AEC, [ii] whole-brain alpha-band AEC, [iii] whole-brain beta-band AEC, [iv] whole-brain delta-theta-band spectral power, [v] temporal alpha-band spectral power, and [vi] whole-brain beta-band spectral power. We computed the average within several regions where large region-level group contrasts were identified (the temporal and frontal regions of interest are illustrated in Appendix 1—figure 7). Consistent with the regional-level group comparisons, the long-range and local synchrony scalar metrics in

Figure 1. Atrophy-cognition EBM staging of AD progression. (A) Histogram of PHG volume loss z-scores, z_{PHG}. (B) Histogram of MMSE-decline z-scores, z_{MMSE}. The z-scores for PHG volume loss and MMSE were standardized by the adjusted scores of the control group, and sign-inverted so that higher z-scores denote more severity. (C) Posterior probabilities, $p_j(k)$, that a subject j belongs to a stage k evaluated by the AC-EBM. (D) The ratio of subjects classified to each stage; blue: Control (CDR 0), orange: MCI due to AD (CDR 0.5), pink: mild AD dementia (CDR 1), and red: moderate AD dementia (CDR 2). (E) Distribution of the stages in the space spanned by PHG volume loss and MMSE score, in which each subject j was distinctly assigned to one of the stages with the highest posterior probability, $\mathrm{argmax}_k\, p_j(k)$. The colors of the dots denote the seven stages. A star symbol denotes the probability-based weighted means of z_{PHG} and MMSE scores at stage 4: $z_{\mathrm{PHG}} = 1.33(\pm 0.258)$ and MMSE$= 26.3(\pm 0.82)$. The values in parentheses denote the standard error (SE; *Equation 5*) of the weighted means. (F) Trajectories of PHG volume loss and MMSE score as a function of the seven stages. Probability-based weighted means ($\pm$ SE) are shown. The initial and final z-scores used in the AC-EBM were: $(z_{\mathrm{initial}}, z_{\mathrm{final}}) = (-1.372, 3.804)$ for PHG volume loss and $(-0.902, 12.712)$ for MMSE decline, respectively. (G) Progression of GM volume loss (z-scores) from stage 1 to 7. Regional GM atrophy in the predicted MCI stage (stage 4) was circled with dotted lines.

the delta-theta band were increased in AD patients compared to controls, and the long-range and local synchrony scalar metrics in the alpha and beta bands were reduced in AD patients compared to controls (Appendix 1—figure 6A, B). We also calculated the z scores, z_{MEG}, of each scalar metric that were used in the SAC-EBMs.

PHG volume loss precedes the MMSE decline in AD progression

An AC-EBM analysis with the two metrics, PHG volume loss, z_{PHG}, and MMSE decline, z_{MMSE}, was performed for six events ($N = 6$; three events for each metric). Robust event thresholds were determined by the exhaustive search of multiple event thresholds (z-score thresholds) and choosing the set of event thresholds that maximize the data likelihood (*Equation 1*). The AC-EBM provided seven stages, each located between consecutive event occurrence times. The resulting posterior probabilities, $p_j(k)$, that a subject j belongs to a stage k are shown in *Figure 1*C. Based on the probabilities, the ratio of subjects classified to each stage was calculated as the probability-based weighted mean (*Figure 1*D). The ratio of subjects with CDR 0.5 was highest in stage 4, and the ratio of controls with CDR 0 at stage 4 was small compared to those in less severe stages of 1–3, indicating that stage 4 corresponds best to clinical MCI due to AD.

The trajectory of PHG volume loss preceded that of MMSE decline (*Figure 1*F), consistent with the relationship between brain atrophy and cognitive decline described in a hypothetical model of biomarker trajectories (*Jack Jr et al., 2010*; *Sperling et al., 2011*). *Figure 1*E visualizes the distribution of the seven stages in the PHG volume loss versus MMSE score. At stage 4, the value of z_{PHG} of 1.33 ± 0.258 was in the range of 1–2. This z-score range of PHG volume loss corresponds to a mild-atrophy range representing approximately MCI stage, e.g., in the voxel-based specific regional analysis system for Alzheimer's disease (VSRAD) software (*Hirata et al., 2005*; *Matsuda et al., 2012*). Furthermore, the MMSE score of 26.3 ± 0.82 at stage 4 was in the range of 23–27. This range of MMSE scores is considered to be typical for MCI due to AD (*Tsoi et al., 2015*). Stage 4 therefore corresponds to MCI stage, whereas stages 3 and 5 correspond to preclinical-AD and mild AD-dementia stages, respectively.

The GM volume z-scores as a function of the seven stages showed that prominent atrophy with $z > 1$ is observed in the temporal regions starting at stage 4 (*Figure 1*G). This trajectory of GM volume approximated the evolution of brain atrophy in the typical progression of AD reported in MRI-based studies; GM volume loss in AD starts in the MTL in the MCI stage, spreads to the lateral temporal and parietal lobes in the mild AD-dementia stage, and spreads further to the frontal lobe in moderate AD-dementia (*Scahill et al., 2002*; *Tondelli et al., 2012*; *Jack Jr et al., 2013*).

These results of the AC-EBM indicate that the PHG volume loss precedes the MMSE decline, and their metric changes track the stages of AD from preclinical AD to moderate AD-dementia. The order of events for GM volume loss and cognitive decline was consistent with the observation that cognitive decline in the early stage of AD progression reflects neuronal loss in the medial temporal regions (*Jack Jr et al., 2018*; *DeTure and Dickson, 2019*).

Neural synchrony progressively changes throughout AD stages estimated by AC-EBM

For the seven stages determined by the AC-EBM (*Figure 1*E–G), profiles of long-range and local neural synchrony were estimated (*Figure 2*). Along the EBM stages, the delta-theta-band synchrony was consistently increased and the alpha and beta-band synchrony was consistently decreased. Neural synchrony showed prominent changes around stage 4 (clinical stage of MCI due to AD). Alpha- and beta-band long-range synchrony decreased steadily across stages 1–3 and then decreased further in stage 4 (*Figure 2*A). Beta-band local synchrony also decreased by half from 1 to 4 (*Figure 2*B). In contrast, there were little changes in delta-theta-band long-range synchrony and delta-theta and alpha-band local synchrony from stage 1 to 3 but these changes became prominent after stage 3.

Figure 2. Profiles of neural synchrony as a function of the AD stages estimated by AC-EBM. (A,B) Profiles of AEC (A) and spectral power (B) as a function of the seven stages, showing probability-based weighted means (± SE). Neural synchrony increased monotonously with AD progression in the delta-theta band and decreased monotonously in the alpha and beta bands. (C,D) Regional AEC (C) and spectral power (D) as a function of the seven stages. Deviations from the neural synchrony spatial patterns averaged over the controls are displayed. The deviations were calculated by using probability-based weighted means of the z-scores standardized by the controls. Spatial patterns in the MCI stage (stage 4) were circled with dotted lines. (E,F) Changes in neural synchrony during the preclinical stages. Regional comparisons between two stages (stages 4 vs. 1) are shown based on non-parametric tests of the weighted mean differences δz. Differences that exceed the threshold ($q < 0.05$) are displayed. There were no significant differences in long-range synchrony in the delta-theta band.

Regional patterns of long-range and local synchrony as a function of the seven stages indicated that prominent changes manifest themselves at stage 4 (*Figure 2*E, F; Appendix 1—table 7 and Appendix 1—table 8). The regions with prominent deviations overlapped with the regions where significant increase and decrease in the neural synchrony were observed in the group comparisons (Appendix 1—figure 6E, F).

The changing patterns of neural synchrony metrics with AD progression indicate that neural synchrony are sensitive indicators of functional change along the AD progression. To further investigate the temporal association of functional deficits with neurodegeneration and cognitive decline, we included neural synchrony in addition to the PHG volume loss and MMSE decline into the EBM frameworks, performing SAC-EBMs.

Long-range synchrony changes in the alpha and beta bands precede PHG volume loss and MMSE decline

SAC-EBMs that include PHG volume loss, z_{PHG}, MMSE decline, z_{MMSE}, and long-range synchrony metric z-scores, z_{MEG}, were performed setting a total of nine events ($N = 9$). SAC-EBMs separately included long-range neural synchrony metrics in the delta-theta, alpha, and beta bands. Each EBM determined the order of nine events, thus defining ten stages (Appendix 1—figure 8; for the corresponding positional variance diagrams of the optimal set of z-score events in the SAC-EBMs, see Appendix 1—figure 9 and Appendix 1—figure 10). The resulting posterior probabilities, $p_j(k)$, that a subject j belongs to a stage k are shown in Appendix 1—figure 11.

For all frequency bands, around stages 5 and 6, the weighted means of PHG volume loss z-scores were in the range of 1–2 and the MMSE scores were in the range of 23–27 (*Figure 3*B, F, J). Furthermore, the ratio of subjects with CDR 0.5 was large around stage 5 (*Figure 3*A, E, I). These indicated that stage 5 best represents the onset of clinical MCI stage, and stages 1-4, where MMSE scores remain almost constant at or near 30, correspond to the preclinical stages of AD. The changes in long-range synchrony during the preclinical stages are shown as statistical bars, and the region-level changes are shown in *Figure 3*C–D, G–H, and K–L.

Long-range synchrony in the alpha and beta bands decreased markedly during the preclinical stages of AD, preceding both PHG volume loss and MMSE decline. Specifically, between stages 1 and 4, the alpha- and beta-band long-range synchrony decreased by more than 80 % of the total drop seen from stage 1 to 10. The whole brain regions, but especially the temporal regions, were involved in these prominent preclinical changes (*Figure 3*H, L). In contrast, the trajectory of delta-theta-band long-range synchrony (*Figure 3*B, C) was almost identical to the evolution of PHG volume loss throughout the stages, but a large variation occurred around the MCI stages (stages 5 and 6) as was found in the AC-EBM (*Figure 2*A). There were no significant synchrony region-level increases in delta-theta band during preclinical stages (*Figure 3*D), consistent with an observation seen in the AC-EBM (*Figure 2*F).

The trajectory shapes of the PHG volume loss (almost linear) and MMSE scores (half parabola) were similar to those obtained in the AC-EBM (*Figure 1*G). This indicates that prominent changes in alpha- and beta-band long-range synchrony during preclinical stages can be utilized to stratify the preclinical stages determined only by neurodegeneration and cognitive deficits.

Local synchrony changes in the alpha and beta bands precede PHG volume loss and MMSE decline

SAC-EBMs including PHG volume loss, MMSE decline, and local synchrony metric z-scores were performed, separately considering delta-theta-, alpha-, and beta-band local synchrony metrics. When considering delta-theta and alpha bands, around stages 6 and 7, the PHG volume loss z-scores were in the range of 1–2 and the MMSE scores were in the range of 23–27 (*Figure 4*B, F), indicating that stage 6 best represents the onset of the MCI stage. Furthermore, the ratios of the subjects with CDR 0.5 were high in stages 6 and 7 (*Figure 4*A, E). For the beta band, based on similar observations, stage 6 best represented the MCI stage (*Figure 4*I, J). For all frequency bands, stages 1-5,

Figure 3. Trajectories of long-range neural synchrony in delta-theta, alpha, and beta-bands from SAC-EBMs. (A,E,I) The ratio of subjects classified to each stage. The ratio was evaluated on the basis of the probabilities that each subject is assigned to each of the ten stages. (B,F,J) Trajectories of long-range synchrony, PHG volume loss, and MMSE score as a function of the ten stages, showing probability-based weighted means (± SE). The asterisks (*$q < 0.05$ and ***$q < 0.001$, FDR corrected) denote statistical significance in comparisons between stages 5 vs. 1. All pairs of stages with significant weighted-mean differences are listed in Appendix 1—table 9. Initial and final z-scores of long-range synchrony used in the SAC-EBMs were: $(z_{\text{initial}}, z_{\text{final}}) = (-1.083, 2.811), (-1.542, 1.605)$, and $(-1.624, 1.641)$ in the delta-theta, alpha, and beta bands, respectively. (C,G,K) Regional AEC along the stages. The deviations from the regional patterns of the control group are displayed. The regional patterns at the onset of the MCI stage were circled with dotted lines. (D,H,L) Changes in regional patterns during the preclinical stages. Regional comparisons between two stages based on nonparametric tests of weighted mean differences δz are shown. Differences exceeding threshold ($q < 0.05$, FDR corrected) are displayed. The top 10 regions with significant differences are listed in Appendix 1—table 10.

Figure 4. Trajectories of local neural synchrony in delta-theta, alpha and beta bands from SAC-EBMs. (A,E,I) The ratio of subjects classified to each stage. (B,F,J) Trajectories of local synchrony, PHG volume loss, and MMSE score as a function of the 10 stages, showing the weighted mean (± SE). Asterisks (*** $q < 0.001$, FDR corrected) denote statistical significance in comparisons between stages 6 vs. 1. All pairs of stages with significant weighted-mean differences are listed in Appendix 1—table 11. Initial and final z-scores of local synchrony used in the SAC-EBMs were: $(z_{initial}, z_{final}) = (-1.329, 6.097)$, $(-1.461, 2.866)$, and $(-1.810, 2.784)$ in the delta-theta, alpha, and beta bands, respectively. (C,G,K) Regional spectral power along the stages. Departures from the regional patterns of the control group are shown. The regional patterns at the onset of the MCI stages were circled with dotted lines. (D,H,L) Changes in regional patterns during the preclinical stages. Regional comparisons between two stages are shown based on nonparametric tests of weighted mean differences δz. Differences exceeding threshold ($q < 0.05$, FDR corrected) are displayed. The top 10 regions with significant differences are listed in Appendix 1—table 12.

where MMSE scores remain almost constant at or near 30, corresponded to the preclinical stages of AD. The changes in local synchrony during the preclinical stages are shown as statistical bars, and the corresponding region-level changes are shown in *Figure 4*C–D, G–H, and K–L.

Local synchrony in the alpha and beta bands decreased during the preclinical stages of AD, preceding both PHG volume loss and MMSE decline (*Figure 4*F, G and *Figure 4*J, K). On the contrary, local synchrony in the delta-theta band increased, lagging the evolution of PHG volume loss (*Figure 4*B, C). Specifically, the alpha-band local synchrony decreased considerably by the onset of the MCI stage, showing significant reductions in the temporal regions (*Figure 4*H) during the preclinical stages (stages 6 vs 1). It is noted that these trends were inconsistent with those found in the AC-EBM (*Figure 2*B), especially within the preclinical stages, where there was little change found in the alpha-band local synchrony. This can be interpreted as evidence that early stages in AD progression may be better characterized by including neurophysiological markers as AD indicators. Beta-band local synchrony also decreased during preclinical stages, preceding PHG volume loss and MMSE decline; by stage 5, the beta-band power decreased by approximately 55 % of the total drop seen throughout the stages, and the reductions were observed in the whole brain (*Figure 4*L). In contrast to the local synchrony trajectories in the alpha and beta bands, the local synchrony in the delta-theta band increased and the *hyper*-synchrony lagged the evolution of the loss of PHG volume in the preclinical stages and made a large jump around the stages 6 and 7 (*Figure 4*D).

As shown in the previous section, large alpha- and beta-band *hypo*-synchrony during the preclinical stages was also observed in long-range synchrony (*Figure 3*F, J). Notably, the decreases in the long-range metrics were much greater than those in the local metrics, especially in the early stages during the phase of the preclinical AD (stages 1–3).

Discussion

We demonstrated that functional deficits of frequency-specific neural synchrony show progressive changes across AD stages. Both long-range and local neural synchrony in the alpha and beta bands, but not in the delta-theta band, was found to decrease in preclinical stages of AD, preceding neurodegeneration and cognitive decline, with more robust findings for long-range neural synchrony. These findings highlight the frequency-specific manifestations of neural synchrony in AD and that synchrony reductions in the alpha and beta bands are sensitive indices reflecting functional deficits in the earliest stages of disease progression.

Electrophysiological metrics of neural synchrony precede volume loss and cognitive decline

A key finding from the current study is that functional deficits as depicted by reduced neural synchrony precede structural volume loss and cognitive deficits. The EBMs on cross-sectional data clearly demonstrated that alpha- and beta-band synchrony within the inferior temporal and posterior parieto-occipital regions show significant deficits in the early disease stages–stages where volumetric and clinical deficits are still not significantly deviated from their baseline trajectory. This is consistent with the finding that functional changes occur earlier in the time course than structural changes in AD (*Jack Jr et al., 2010*; *Sperling et al., 2011*).

Previous functional MRI studies have demonstrated disrupted connectivity especially between the hippocampus and several cortical default mode network (DMN) areas in subjects with amyloid deposition but without cognitive impairment (*Sperling et al., 2014*). Such a disruption in the DMN has also been observed in clinically normal older individuals without prominent brain atrophy in MTL preserving hippocampal activity (*Miller et al., 2008*; *Hedden et al., 2009*), indicating altered functional connectivity during the period of preclinical AD. In contrast to such fMRI data reflecting the cascade of neural, metabolic, hemodynamic events in AD, our findings from MEG, which captures the synaptic physiology as the collective oscillatory spectra, demonstrate direct observations of AD-related altered neuronal activity.

Frequency-specific manifestations of neural synchrony deficits along the progression of the disease

We demonstrated that the oscillatory deficits and their temporal association to neurodegeneration and cognitive decline are frequency specific. In particular, it is alpha and beta hyposynchrony that precedes PHG atrophy and MMSE decline, whereas delta-theta hypersynchrony does not seem to show such precedence. This is consistent with previous findings that alpha and beta hyposynchrony is more tightly associated with tau accumulation which is closely allied to neurodegeneration and cognitive decline (*Pusil et al., 2019*; *Ranasinghe et al., 2020*, *2021*). Neural hyposynchrony in the alpha and beta bands may represent harbingers of altered synaptic physiology associated with tau accumulation. In fact, in human postmortem studies, the strongest correlate of cognitive deficits in AD patients is loss of synapse (*DeKosky and Scheff, 1990*; *Terry et al., 1991*). A study using transgenic AD mice has also shown that synaptotoxicity is an early phenomenon in AD pathophysiology (*Zhou et al., 2017*). In the context of fluid biomarkers to detect plasma amyloid, alpha and beta hyposynchrony may detect and quantify tau-associated neurodegenerative mechanisms and hence may provide crucial information for early therapeutic interventions.

Previous studies have also shown that delta-theta oscillatory activity is increased in AD and is strongly associated with amyloid accumulation (*Ranasinghe et al., 2020*, *2022b*). In particular, increased delta-theta activity is a robust signal in individuals who are amyloid positive and cognitively unimpaired as well as those who harbor APOE-$\epsilon4$ allele and an increased risk of AD (*Cuesta et al., 2015*; *Nakamura et al., 2018*). These previous findings indicate that delta-theta hypersynchrony is an early change in AD spectrum and may even precede the neurodegeneration and cognitive deficits. However, in the current results, the trajectory of the delta-theta hypersynchrony was identical to or lagged that of the PHG volume loss. This apparent controversy may be due to the possibility that oscillatory changes in the delta-theta band are more closely related to amyloid accumulations in AD, which become saturated early in the disease course and have a poor association with neurodegeneration and cognitive trajectories. It would be worth exploring how the trajectory of early saturated variables may be captured by EBM approaches.

Distinction between long-range and local synchrony deficits in disease progression

The decrease in alpha and beta-band long-range metrics in the preclinical stages was much greater than that in the local metrics. This is consistent with the fact that AD-related abnormal brain activities are observed as disruptions of functional networks. Long-range cross-regional metrics, such as AECs, directly capture network disruptions involving all brain regions, while local metrics capture features of individual regions. From the definition, local synchrony describes collective neuronal oscillations in each local region, and thus the change along AD progression may depend mainly on long-term, slowly changing regional neuronal loss. On the other hand, long-range synchrony describes temporal coherence amongst regional collective neuronal oscillations and is vulnerable to altered neuronal oscillations. Therefore, long-range metrics are more sensitive to abnormal rhythms, gathering local abnormalities.

Preclinical neurophysiological markers that indicate the pathophysiology of AD are clinically important but have not been established. Aβ accumulation in preclinical stages is just a necessary condition for AD, and additional preclinical markers are required to fully predict the progression of AD. From this point of view, the present study indicates that alpha and beta-band MEG metrics, especially long-range-synchrony metrics (AEC), which were found to be sensitive to preclinical stages, might be promising candidates as such additional markers.

Limitations

A limitation of the current study is that there were differences in age between controls and AD patients. Although we adjusted the age of each metric employing GLMs, age trajectories in neurophysiological measures have been reported to be nonlinear even in healthy aging (*Sahoo et al., 2020*). Age-related changes in brain atrophy have also been reported to follow a nonlinear time

course depending on the brain areas (*Coupé et al., 2019*). These studies indicate that it may be better to employ a non-linear method beyond GLM to perfectly correct aging effects.

Another limitation is that we have not performed independent validations of the predicted trajectories and also have not examined the heterogeneity in AD progression, although we clarified for the first time the time courses of MEG neurophysiological metrics in AD progression. In fact, AD is a heterogeneous multifactorial disorder with various pathobiological subtypes (*Jellinger, 2022*). In this context, an EBM called *Subtype and Stage Inference* (SuStaIn) capable of capturing spatio-temporal heterogeneity of diseases (*Young et al., 2018*) has been proposed for subtyping of neurodegenerative diseases including typical AD and has been applied to find different spatio-temporal trajectories of longitudinal tau-PET data in AD (*Vogel et al., 2021*). Since oscillatory rhythms are thought to depend on AD subtypes (*Ranasinghe et al., 2017*, *2022a*), an extended trajectory analysis considering both spatial and temporal variations of MEG/EEG metrics is warranted in the future and such analyses would provide distinct neurophysiological trajectories depending on AD subtypes. As a validation of the predicted trajectories, it would be necessary to investigate whether the predicted EBM stages are reliable and predictive of conversions (e.g., from control to MCI) while taking the AD subtypes into account.

Data availability

The informed consent did not include a declaration regarding the public availability of the data, and the data for this study will not be made publicly available. Anonymized data may be shared on request from qualified investigators for the purposes of replicating procedures and results within the limits of participants' consent. Matlab scripts used in this study are available from the corresponding author upon reasonable requests.

Acknowledgements

The authors thank all study participants for their support for our research.

Funding

This study was supported by the following grants: the National Institutes of Health grants: R01AG062196 (S.S.N.), R01NS100440 (S.S.N.), R01DC017091 (S.S.N.), P50DC019900 (S.S.N.), P30AG062422 (G.D.R., B.L.M.), K23AG038357 (K.V.), K08AG058749 (K.G.R.), R21AG077498 (K.G.R.); University of California grant: UCOP-MRP-17-454755 (S.S.N); John Douglas French Alzheimer's Foundation (K.V.); S.D. Bechtel Jr. Foundation (K.V.); Alzheimer's Association grant: PCTRB-13-288476 mmade possible by Part the Cloud™ (K.V.), AARG-21-849773 (K.G.R.); Larry L. Hillblom Foundation grant: 2015-A-034-FEL (K.G.R.); Larry L. Hillblom Foundation grant: 2019-A-013-SUP (KGR); Research contract from Ricoh MEG Inc (S.S.N., H.E.K.).

Competing interests

K.K. and H.M. are employees of Ricoh Company, Ltd. The authors declare that no other competing interests exist. The other authors declare no competing financial conflicts of interest to disclose.

References

Ablin P, Cardoso JF, Gramfort A. Faster independent component analysis by preconditioning with Hessian approximations. IEEE Transactions on Signal Processing. 2018; 66(15):4040–4049.

Ahnaou A, Moechars D, Raeymaekers L, Biermans R, Manyakov N, Bottelbergs A, Wintmolders C, Van Kolen K, Van De Casteele T, Kemp JA, Drinkenburg WH. Emergence of early alterations in network oscillations and functional connectivity in a tau seeding mouse model of Alzheimer's disease pathology. Scientific Reports. 2017; 7(1):1–14.

Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, Gamst A, Holtzman DM, Jagust WJ, Petersen RC, Snyder PJ, Carrillo MC, Thies B, Phelps CH. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's & Dementia. 2011; 7(3):270–279.

Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological). 1995; 57(1):289–300.

Briels CT, Schoonhoven DN, Stam CJ, de Waal H, Scheltens P, Gouw AA. Reproducibility of EEG functional connectivity in Alzheimer's disease. Alzheimer's Research & Therapy. 2020; 12:1–14.

Briels CT, Stam CJ, Scheltens P, Bruins S, Lues I, Gouw AA. In pursuit of a sensitive EEG functional connectivity outcome measure for clinical trials in Alzheimer's disease. Clinical Neurophysiology. 2020; 131(1):88–95.

Busche MA, Eichhoff G, Adelsberger H, Abramowski D, Wiederhold KH, Haass C, Staufenbiel M, Konnerth A, Garaschuk O. Clusters of hyperactive neurons near amyloid plaques in a mouse model of Alzheimer's disease. Science. 2008; 321(5896):1686–1689.

Busche MA, Wegmann S, Dujardin S, Commins C, Schiantarelli J, Klickstein N, Kamath TV, Carlson GA, Nelken I, Hyman BT. Tau impairs neural circuits, dominating amyloid-β effects, in Alzheimer models in vivo. Nature Neuroscience. 2019; 22(1):57–64.

Colclough GL, Woolrich MW, Tewarie P, Brookes MJ, Quinn AJ, Smith SM. How reliable are MEG resting-state connectivity metrics? Neuroimage. 2016; 138:284–293.

Coupé P, Manjón JV, Lanuza E, Catheline G. Lifespan changes of the human brain in Alzheimer's disease. Scientific Reports. 2019; 9(1):1–12.

Cuesta P, Garcés P, Castellanos NP, López ME, Aurtenetxe S, Bajo R, Pineda-Pardo JA, Bruña R, Marín AG, Delgado M, Barabash A, Ancín I, Cabranes JA, Fernandez A, , Pozo FD, Sancho M, Marcos A, Nakamura A, Maestú F. Influence of the APOE ϵ4 allele and mild cognitive impairment diagnosis in the disruption of the MEG resting state functional connectivity in sources space. Journal of Alzheimer's Disease. 2015; 44(2):493–505.

DeKosky ST, Scheff SW. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society. 1990; 27(5):457–464.

DeTure MA, Dickson DW. The neuropathological diagnosis of Alzheimer's disease. Molecular Neurodegeneration. 2019; 14(1):1–18.

Dubois B, Hampel H, Feldman HH, Scheltens P, Aisen P, Andrieu S, Bakardjian H, Benali H, Bertram L, Blennow K, Broich K, Cavedo E, Crutch S, Dartigues JF, Duyckaerts C, Epelbaum S, Frisoni GB, Gauthier S, Genthon R, Gouw AA, et al. Preclinical Alzheimer's disease: definition, natural history, and diagnostic criteria. Alzheimer's & Dementia. 2016; 12(3):292–323.

Echávarri C, Aalten P, Uylings HB, Jacobs H, Visser PJ, Gronenschild E, Verhey F, Burgmans S. Atrophy in the parahippocampal gyrus as an early biomarker of Alzheimer's disease. Brain Structure and Function. 2011; 215(3):265–271.

Engels M, Hillebrand A, van der Flier WM, Stam CJ, Scheltens P, van Straaten EC. Slowing of hippocampal activity correlates with cognitive decline in early onset Alzheimer's disease. An MEG study with virtual electrodes. Frontiers in Human Neuroscience. 2016; 10:238.

Fernández A, Hornero R, Mayo A, Poza J, Gil-Gregorio P, Ortiz T. MEG spectral profile in Alzheimer's disease and mild cognitive impairment. Clinical Neurophysiology. 2006; 117(2):306–314.

Fonteijn HM, Modat M, Clarkson MJ, Barnes J, Lehmann M, Hobbs NZ, Scahill RI, Tabrizi SJ, Ourselin S, Fox NC, Alexander DC. An event-based model for disease progression and its application in familial Alzheimer's disease and Huntington's disease. NeuroImage. 2012; 60(3):1880–1889.

Gaser C, Dahnke R, Thompson PM, Kurth F, Luders E. CAT-a computational anatomy toolbox for the analysis of structural MRI data. BioRxiv. 2022; p. 2022–06.

Hedden T, Van Dijk KR, Becker JA, Mehta A, Sperling RA, Johnson KA, Buckner RL. Disruption of functional connectivity in clinically normal older adults harboring amyloid burden. Journal of Neuroscience. 2009; 29(40):12686–12694.

Hipp JF, Hawellek DJ, Corbetta M, Siegel M, Engel AK. Large-scale cortical correlation structure of spontaneous
oscillatory activity. Nature Neuroscience. 2012; 15(6):884–890.

Hirata Y, Matsuda H, Nemoto K, Ohnishi T, Hirao K, Yamashita F, Asada T, Iwabuchi S, Samejima H. Voxel-
based morphometry to discriminate early Alzheimer's disease from controls. Neuroscience Letters. 2005;
382(3):269–274.

Hughes LE, Henson RN, Pereda E, Bruña R, López-Sanz D, Quinn AJ, Woolrich MW, Nobre AC, Rowe JB, Maestú
F. Biomagnetic biomarkers for dementia: A pilot multicentre study with a recommended methodological
framework for magnetoencephalography. Alzheimer's & Dementia: Diagnosis, Assessment & Disease Moni-
toring. 2019; 11:450–462.

Jack Jr CR, Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, Holtzman DM, Jagust W, Jessen F, Kar-
lawish J, Liu E, Molinuevo JL, Montine T, Phelps C, Rankin KP, Rowe CC, Scheltens P, Siemers E, Snyder HM,
Sperling R. NIA-AA research framework: toward a biological definition of Alzheimer's disease. Alzheimer's &
Dementia. 2018; 14(4):535–562.

Jack Jr CR, Knopman DS, Jagust WJ, Petersen RC, Weiner MW, Aisen PS, Shaw LM, Vemuri P, Wiste HJ, Weigand SD,
Lesnick TG, Pankratz VS, Donohue MC, Trojanowski JQ. Tracking pathophysiological processes in Alzheimer's
disease: an updated hypothetical model of dynamic biomarkers. The Lancet Neurology. 2013; 12(2):207–216.

Jack Jr CR, Wiste HJ, Vemuri P, Weigand SD, Senjem ML, Zeng G, Bernstein MA, Gunter JL, Pankratz VS, Aisen
PS, Weiner MW, Petersen RC, Shaw LM, Trojanowski JQ, Knopman DS. Brain beta-amyloid measures and
magnetic resonance imaging atrophy both predict time-to-progression from mild cognitive impairment to
Alzheimer's disease. Brain. 2010; 133(11):3336–3348.

Jellinger KA. Recent update on the heterogeneity of the Alzheimer's disease spectrum. Journal of Neural
Transmission. 2022; 129(1):1–24.

Jeong J. EEG dynamics in patients with Alzheimer's disease. Clinical Neurophysiology. 2004; 115(7):1490–1505.

Koelewijn L, Bompas A, Tales A, Brookes MJ, Muthukumaraswamy SD, Bayer A, Singh KD. Alzheimer's disease
disrupts alpha and beta-band resting-state oscillatory network connectivity. Clinical Neurophysiology. 2017;
128(11):2347–2357.

Matsuda H, Mizumura S, Nemoto K, Yamashita F, Imabayashi E, Sato N, Asada T. Automatic voxel-based mor-
phometry of structural MRI by SPM8 plus diffeomorphic anatomic registration through exponentiated lie
algebra improves the diagnosis of probable Alzheimer Disease. American Journal of Neuroradiology. 2012;
33(6):1109–1114.

Matsuda H. MRI morphometry in Alzheimer's disease. Ageing Research Reviews. 2016; 30:17–24.

McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack Jr CR, Kawas CH, Klunk WE, Koroshetz WJ, Manly
JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carrillo MC, Thies B, Weintraub S, Phelps CH.
The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on
Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's &
Dementia. 2011; 7(3):263–269.

Meghdadi AH, Stevanović Karić M, McConnell M, Rupp G, Richard C, Hamilton J, Salat D, Berka C. Resting
state EEG biomarkers of cognitive decline associated with Alzheimer's disease and mild cognitive impairment.
PLOS ONE. 2021; 16(2):e0244180.

Miller SL, Celone K, DePeau K, Diamond E, Dickerson BC, Rentz D, Pihlajamäki M, Sperling RA. Age-related
memory impairment associated with loss of parietal deactivation but preserved hippocampal activation. Pro-
ceedings of the National Academy of Sciences. 2008; 105(6):2181–2186.

Nakamura A, Cuesta P, Fernández A, Arahata Y, Iwata K, Kuratsubo I, Bundo M, Hattori H, Sakurai T, Fukuda
K, Washimi Y, Endo H, Takeda A, Diers K, Bajo R, Maestú F, Ito K, Kato T. Electromagnetic signatures of the
preclinical and prodromal stages of Alzheimer's disease. Brain. 2018; 141(5):1470–1485.

Nolte G. The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalog-
raphy forward calculation in realistic volume conductors. Physics in Medicine & Biology. 2003; 48(22):3637.

Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: open source software for advanced analysis of MEG,
EEG, and invasive electrophysiological data. Computational Intelligence and Neuroscience. 2011; 2011.

Penny WD, Friston KJ, Ashburner JT, Kiebel SJ, Nichols TE. Statistical parametric mapping: the analysis of functional brain images. Elsevier; 2011.

Pusil S, López ME, Cuesta P, Bruna R, Pereda E, Maestu F. Hypersynchronization in mild cognitive impairment: the 'X'model. Brain. 2019; 142(12):3936–3950.

Ranasinghe KG, Cha J, Iaccarino L, Hinkley LB, Beagle AJ, Pham J, Jagust WJ, Miller BL, Rankin KP, Rabinovici GD, Vossel KA, Nagarajan SS. Neurophysiological signatures in Alzheimer's disease are distinctly associated with TAU, amyloid-β accumulation, and cognitive decline. Science Translational Medicine. 2020; 12(534):eaaz4069.

Ranasinghe KG, Hinkley LB, Beagle AJ, Mizuiri D, Honma SM, Welch AE, Hubbard I, Mandelli ML, Miller ZA, Garrett C, La A, Boxer AL, Houde JF, Miller BL, Vossel KA, Gorno-Tempini ML, Nagarajan SS. Distinct spatiotemporal patterns of neuronal functional connectivity in primary progressive aphasia variants. Brain. 2017; 140(10):2737–2751.

Ranasinghe KG, Kudo K, Hinkley L, Beagle A, Lerner H, Mizuiri D, Findlay A, Miller BL, Kramer JH, Gorno-Tempini ML, Rabinovici GD, Rankin KP, Garcia PA, Kirsch HE, Vossel K, Nagarajan SS. Neuronal synchrony abnormalities associated with subclinical epileptiform activity in early-onset Alzheimer's disease. Brain. 2022; 145(2):744–753.

Ranasinghe KG, Petersen C, Kudo K, Mizuiri D, Rankin KP, Rabinovici GD, Gorno-Tempini ML, Seeley WW, Spina S, Miller BL, Vossel K, Grinberg LT, Nagarajan SS. Reduced synchrony in alpha oscillations during life predicts post mortem neurofibrillary tangle density in early-onset and atypical Alzheimer's disease. Alzheimer's & Dementia. 2021; 17(12):2009–2019.

Ranasinghe KG, Verma P, Cai C, Xie X, Kudo K, Gao X, Lerner H, Mizuiri D, Strom A, Iaccarino L, La Joie R, Miller BL, Gorno-Tempini ML, Rankin KP, Jagust WJ, Vossel K, Rabinovici GD, Raj A, Nagarajan SS. Altered excitatory and inhibitory neuronal subpopulation parameters are distinctly associated with tau and amyloid in Alzheimer's disease. Elife. 2022; 11:e77850.

Rolls ET, Huang CC, Lin CP, Feng J, Joliot M. Automated anatomical labelling atlas 3. Neuroimage. 2020; 206:116189.

Sahoo B, Pathak A, Deco G, Banerjee A, Roy D. Lifespan associated global patterns of coherent neural communication. Neuroimage. 2020; 216:116824.

Scahill RI, Schott JM, Stevens JM, Rossor MN, Fox NC. Mapping the evolution of regional atrophy in Alzheimer's disease: unbiased analysis of fluid-registered serial MRI. Proceedings of the National Academy of Sciences. 2002; 99(7):4703–4707.

Schoonhoven DN, Briels CT, Hillebrand A, Scheltens P, Stam CJ, Gouw AA. Sensitive and reproducible MEG resting-state metrics of functional connectivity in Alzheimer's disease. Alzheimer's Research & Therapy. 2022; 14(1):1–19.

Sekihara K, Kawabata Y, Ushio S, Sumiya S, Kawabata S, Adachi Y, Nagarajan SS. Dual signal subspace projection (DSSP): a novel algorithm for removing large interference in biomagnetic measurements. Journal of Neural Engineering. 2016; 13(3):036007.

Sekihara K, Nagarajan SS. Electromagnetic brain imaging: a bayesian perspective. Springer; 2015.

Sekihara K, Nagarajan SS, Poeppel D, Marantz A. Asymptotic SNR of scalar and vector minimum-variance beamformers for neuromagnetic source reconstruction. IEEE Transactions on Biomedical Engineering. 2004; 51(10):1726–1734.

Sekihara K, Nagarajan SS. Adaptive spatial filters for electromagnetic brain imaging. Springer Science & Business Media; 2008.

Selkoe DJ. Alzheimer's disease is a synaptic failure. Science. 2002; 298(5594):789–791.

Sperling R, Mormino E, Johnson K. The evolution of preclinical Alzheimer's disease: implications for prevention trials. Neuron. 2014; 84(3):608–622.

Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM, Iwatsubo T, Jack Jr CR, Kaye J, Montine TJ, Park DC, Reiman EM, Rowe CC, Siemers E, Stern Y, Yaffe K, Carrillo MC, Thies B, Morrison-Bogorad M, Wagster MV, et al. Toward defining the preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's & Dementia. 2011; 7(3):280–292.

Spires-Jones TL, Hyman BT. The intersection of amyloid beta and tau at synapses in Alzheimer's disease. Neu-
ron. 2014; 82(4):756–771.

Stam CJ, Jones B, Manshanden I, Van Walsum AvC, Montez T, Verbunt JP, de Munck JC, van Dijk BW, Berendse
HW, Scheltens P. Magnetoencephalographic evaluation of resting-state functional connectivity in Alzheimer's
disease. Neuroimage. 2006; 32(3):1335–1344.

Teipel SJ, Pruessner JC, Faltraco F, Born C, Rocha-Unold M, Evans A, Möller HJ, Hampel H. Comprehensive dis-
section of the medial temporal lobe in AD: measurement of hippocampus, amygdala, entorhinal, perirhinal
and parahippocampal cortices using MRI. Journal of Neurology. 2006; 253(6):794–800.

Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, Hill R, Hansen LA, Katzman R. Physical basis of cognitive
alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. Annals of
Neurology: Official Journal of the American Neurological Association and the Child Neurology Society. 1991;
30(4):572–580.

Tondelli M, Wilcock GK, Nichelli P, De Jager CA, Jenkinson M, Zamboni G. Structural MRI changes detectable up
to ten years before clinical Alzheimer's disease. Neurobiology of Aging. 2012; 33(4):825.e25–36.

Tsoi KK, Chan JY, Hirai HW, Wong SY, Kwok TC. Cognitive tests to detect dementia: a systematic review and
meta-analysis. JAMA Internal Medicine. 2015; 175(9):1450–1458.

Vogel JW, Young AL, Oxtoby NP, Smith R, Ossenkoppele R, Strandberg OT, La Joie R, Aksman LM, Grothe MJ,
Iturria-Medina Y, Initiative ADN, Pontecorvo MJ, Devous MD, Rabinovici GD, Alexander DC, Lyoo CH, Evans AC,
Hansson O. Four distinct trajectories of tau deposition identified in Alzheimer's disease. Nature Medicine.
2021; 27(5):871–881.

Wiesman AI, Castanheira JDS, Baillet S. Stability of spectral estimates in resting-state magnetoencephalogra-
phy: recommendations for minimal data duration with neuroanatomical specificity. NeuroImage. 2021; p.
118823.

Wiesman AI, Murman DL, May PE, Schantell M, Losh RA, Johnson HJ, Willet MP, Eastman JA, Christopher-Hayes
NJ, Knott NL, Houseman LL, Wolfson SL, Losh KL, Johnson CM, Wilson TW. Spatio-spectral relationships
between pathological neural dynamics and cognitive impairment along the Alzheimer's disease spectrum.
Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring. 2021; 13(1):e12200.

Xia M, Wang J, He Y. BrainNet Viewer: a network visualization tool for human brain connectomics. PLOS ONE.
2013; 8(7):e68910.

Young AL, Marinescu RV, Oxtoby NP, Bocchetta M, Yong K, Firth NC, Cash DM, Thomas DL, Dick KM, Cardoso
J, Swieten Jv, Borroni B, Galimberti D, Masellis M, Tartaglia MC, Rowe JB, Graff C, Tagliavini F, Frisoni GB,
Laforce Jr R, et al. Uncovering the heterogeneity and temporal complexity of neurodegenerative diseases
with Subtype and Stage Inference. Nature Communications. 2018; 9(1):1–16.

Young AL, Oxtoby NP, Daga P, Cash DM, Fox NC, Ourselin S, Schott JM, Alexander DC. A data-driven model of
biomarker changes in sporadic Alzheimer's disease. Brain. 2014; 137(9):2564–2577.

Zhou L, McInnes J, Wierda K, Holt M, Herrmann AG, Jackson RJ, Wang YC, Swerts J, Beyens J, Miskiewicz K, Vilain
S, Dewachter I, Moechars D, De Strooper B, Spires-Jones TL, De Wit J, Verstreken P. Tau association with
synaptic vesicles causes presynaptic dysfunction. Nature Communications. 2017; 8(1):15295.