Elucidating disease-associated mechanisms triggered by pollutants via the epigenetic landscape using large-scale ChIP-Seq data

Zhaonan Zou[1], Yuka Yoshimura[1], Yoshihiro Yamanishi[2] and Shinya Oki[1*]

[1] Department of Drug Discovery Medicine, Kyoto University Graduate School of Medicine, 53 Shogoin Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan

[2] Department of Complex Systems Science, Graduate School of Informatics, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8602, Japan

E-mail:

Zhaonan Zou   zou.zhaonan.56e@st.kyoto-u.ac.jp

Yuka Yoshimura   y-yoshimura1205@outlook.com

Yoshihiro Yamanishi   yamanishi@i.nagoya-u.ac.jp

Shinya Oki   oki.shinya.3w@kyoto-u.ac.jp

*Correspondence: oki.shinya.3w@kyoto-u.ac.jp; Tel: +81-75-366-7479

ABSTRACT

Background: Despite well-documented effects on human health, the action modes of

environmental pollutants are incompletely understood. Although transcriptome-based

approaches are widely used to predict associations between chemicals and disorders, the

molecular cues regulating pollutant-derived gene expression changes remain unclear.

Therefore, we developed a data-mining approach, termed "DAR-ChIPEA," to identify

transcription factors (TFs) playing pivotal roles in the action modes of pollutants.

Methods: Large-scale public ChIP-Seq data (human, n = 15,155; mouse, n = 13,156) were

used to predict TFs that are enriched in the pollutant-induced differentially accessible

genomic regions (DARs) obtained from epigenome analyses (ATAC-Seq). The resultant

pollutant–TF matrices were then cross-referenced to a repository of TF–disorder associations

to account for pollutant modes of action. We subsequently evaluated the performance of the

proposed method using a chemical perturbation dataset to compare the outputs of the

DAR-ChIPEA and our previously developed differentially expressed gene (DEG)-ChIPEA

methods using pollutant-induced DEGs as input. We then adopted the proposed method to

predict disease-associated mechanisms triggered by pollutants.

Results: The proposed approach outperformed other methods using the area under the

receiver operating characteristic curve score. The mean score of the proposed DAR-ChIPEA

was significantly higher than that of our previously described DEG-ChIPEA (0.7287 vs. 0.7060;

Q = 5.278 × 10^{-42}; two-tailed Wilcoxon rank-sum test). The proposed approach further

predicted TF-driven modes of action upon pollutant exposure, indicating that (1) TFs

regulating Th1/2 cell homeostasis are integral in the pathophysiology of tributyltin-induced

allergic disorders; (2) fine particulates (PM$_{2.5}$) inhibit the binding of C/EBPs, Rela, and Spi1 to

the genome, thereby perturbing normal blood cell differentiation and leading to immune

dysfunction; and (3) lead induces fatty liver by disrupting the normal regulation of lipid

metabolism by altering hepatic circadian rhythms.

Conclusion: Highlighting genome-wide chromatin change upon pollutant exposure to

elucidate the epigenetic landscape of pollutant responses outperformed our previously

described method that focuses on gene-adjacent domains only. Our approach has the

potential to reveal pivotal TFs that mediate deleterious effects of pollutants, thereby

facilitating the development of strategies to mitigate damage from environmental pollution.

Keywords:

ChIP-Seq, ATAC-seq, Transcription factor, Differentially accessible genomic region,

Epigenetic landscape, Action modes of environmental pollutants

BACKGROUND

Short- and/or long-term exposure to environmental pollutants has been associated with assorted adverse human health outcomes, including increased respiratory symptoms, heart or lung disease, and even early mortality [1–4]. Toxic pollutants in particular are implicated in carcinogenesis and other serious health effects, such as reproductive disorders and congenital anomalies [5–7]. To elucidate the modes of action (MoAs) by which chemicals, including environmental contaminants, elicit such toxic responses, computational strategies utilizing biomedical big data analysis have attracted the attention of researchers in recent decades [8–10]. For example, with the understanding that bioactive chemicals, such as pharmaceuticals and pollutants, affect human physiology by either maintaining or disrupting normal gene expression, respectively, comparative analyses of transcriptional profiles before and after chemical exposure, followed by gene ontology (GO) and pathway enrichment analyses, have been performed to elucidate chemical MoAs [11, 12]. However, although GO and pathway analyses can comprehensively extract functional characteristics from profiles of chemically induced differentially expressed genes (DEGs), considerable effort is required to identify specific primary literature that demonstrates the association between input DEGs and associated GOs or pathways. In addition, graph analytics (network analysis) is also frequently employed to estimate "molecular hubs" within chemically responsive gene regulatory networks, despite limitations in terms of the biological interpretability of the prediction outcomes [13].

Bioactive chemicals, including some pollutants, can target a wide variety of molecules, such as membrane transport proteins, ion channels, and metabolic enzymes. In particular, research on steroid receptors, which are activated by female (estrogens) and male (androgens) hormones, is actively pursued owing to the reported effects of endocrine-

disrupting chemicals, also known as environmental hormones, on reproduction and

development [14, 15]. In addition, heavy metals, including mercury and cadmium, which have

been associated with pollution-related disorders such as Minamata disease and Itai-itai

disease, may induce epigenetic changes as a component of their disease mechanisms [16,

17]. Furthermore, both intranuclear steroid receptors and enzymes that regulate epigenomic

states, such as DNA methylation and histone modifications, can function as transcription

factors (TFs) to directly control gene expression. Accordingly, certain pharmaceuticals such as

estrogen receptor inhibitors and histone deacetylase inhibitors have been shown to exert

anticancer effects by directly regulating TF function [18, 19]. Together, these observations

suggest that TFs that collectively regulate the on/off switches of genes whose expression is

significantly altered by exposure to environmental chemicals may function as direct targets

or mediators of chemical MoAs.

To identify TFs whose binding is enriched to a given set of genes or genomic regions,

binding motif-based methods (MotifEA) such as HOMER and MEME have widely been used

[20, 21]. This approach predicts TF binding by extracting shared short nucleotide sequences,

commonly referred to as "motifs", from multiple input genomic regions. These motifs are

then compared to known TF–DNA binding motifs by comparative analysis. As an alternative

method for identifying key TFs in chemical MoAs, we recently developed an approach that is

independent of TF–DNA binding motifs, which we termed the DEG-ChIPEA method [22]. This

method is based on the ChIP-Atlas database, which extracts actual TF binding sites on the

reference genome by curating and integrating tens of thousands of previously published

ChIP-Seq experimental data examining TF binding to genomic DNA [23]. The DEG-ChIPEA

method involves inputting chemically induced DEGs and performing enrichment analysis

based on all ChIP-Seq experiments obtained from ChIP-Atlas (i.e., ChIP-Seq-based

enrichment analysis; ChIPEA) to identify TFs whose binding is significantly enriched in the

vicinity of the input DEGs. These identified TFs can be considered promising candidate

pivotal factors that mediate the expression of therapeutic or adverse chemical effects.

However, despite the generally high accuracy of DEG-ChIPEA, a potential limitation is that

only TFs in the immediate vicinity of the transcription start site (transcription start site ± 5 kb)

of the input chemically induced genes are analyzed. Thus, potential contributions to gene

expression by TFs that bind to distal transcriptional regulatory regions, such as enhancers

and topologically associating domain boundaries located far from the gene locus, are not

taken into account.

In this study, we aimed to gain further insights into the MoAs of chemicals, particularly

those of environmental pollutants, with regard to health effects. To achieve this, we propose

a novel approach, termed "DAR-ChIPEA", that considers TF binding events throughout the

genome, including both proximal and distal transcriptional regulatory regions of genes, in

contrast to the limited purview of the DEG-ChIPEA method. The DAR-ChIPEA method

involves extracting differentially accessible regions (DARs), which are genomic regions with

significantly different chromatin accessibility prior to and following chemical exposure, using

data from environmental chemical perturbation epigenome experiments, such as ATAC-Seq

datasets. These DARs are then used as input for ChIPEA to generate a chemical–TF

association matrix, which allows the identification of TFs whose binding is significantly

enriched in the input DARs. In addition, we constructed a chemical–TF–disorder triadic

association by incorporating known TF–disease associations into the resulting chemical–TF

matrix. Our approach outperformed methods based on TF–DNA binding motifs, as well as

the previously developed DEG-ChIPEA method [22], which focuses only on TFs that bind in

the vicinity of chemically induced DEGs.

METHODS

Chemical- or pollutant-perturbed transcriptome and epigenome data

To identify genomic regions with significant differences in chromatin accessibility (DARs) in

response to chemical or pollutant exposure as determined through epigenome perturbation

experiments, we used ATAC-Seq datasets retrieved from the DBKERO (n = 2,078;

DRA006903–DRA006930) [24, 25] and TaRGET (n = 383; GSE146508) databases [24, 25]. In

addition, RNA-Seq datasets corresponding to the ATAC-Seq datasets were made available in

these databases and used to identify DEGs induced by chemical or pollutant exposure as

input to the DEG-ChIPEA or DEG-MotifEA analyses (Additional file 1: Supplementary Fig. S1;

DBKERO, n = 2,012, DRA006875–DRA006902; TaRGET, n = 468, GSE146508). DBKERO

harbors perturbation data of 99 chemicals in human cell lines, whereas TaRGET curates *in-*

vivo exposure data for seven pollutants in blood and liver samples from mice. All data used

in this analysis were archived in the National Center for Biotechnology Information Sequence

Read Archive (NCBI SRA) [26] and are associated with accession IDs as listed in

Supplementary Tables S1–S4 (Additional file 2).

Genome-wide TF binding and chromatin accessibility experimental data from ChIP-

Atlas

Genome-wide TF binding site information was obtained from the ChIP-Atlas database [23],

which comprehensively integrates nearly all publicly available ChIP-Seq data. Peak call data

in BED format (MACS2 [27]; Q-value <1 × 10^{-10}) were used in this study. Data from ChIP-Seq

experiments examining binding of 1,796 and 866 TFs on the human (GRCh38/hg38; hereafter

referred to as hg38) and mouse genomes (GRCm38/mm10; hereafter referred to as mm10),

respectively, were used in this study (hg38, n = 15,155; mm10, n = 13,156). In addition, the

information related to accessible chromatin used in this analysis to identify DARs induced by

chemical or pollutant exposure was also obtained from ChIP-Atlas, which also provides peak

call data for ATAC-Seq.

TF–disease associations

Association data between TFs and diseases were obtained from DisGeNET (v 7.0) [28], which

is a database of genetic and genomic information associated with human diseases. DisGeNET

integrates this information from other expert-curated repositories and PubMed literature.

Only manually curated data that were labeled with referenced PubMed IDs were retrieved,

resulting in a total of 12,857 associations between TFs and diseases, involving 1,004 TFs and

3,205 diseases.

Known chemical–disease associations as standard data

Manually curated chemical–disease association data were acquired from the Comparative

Toxicogenomics Database (CTD) [29]. In total, we incorporated 28,821 chemical–disease

associations involving 69 chemicals (DBKERO) and seven pollutants (TaRGET), and 2,150

diseases in this study.

Identification of chemical- or pollutant-induced DEGs

After obtaining the SRA accession IDs of the chemical or pollutant perturbation RNA-Seq

data from DBKERO and TaRGET, the raw sequencing data archived in NCBI SRA were

downloaded and decoded into FASTQ format using the "fastq-dump" command from the

SRA Toolkit (v3.0.0) [30]. Quality control was then performed using the "fastq_quality_filter"

command from the FASTX-Toolkit (v0.0.13) [31]. Alignment to the reference genomes

(DBKERO, hg38; TaRGET, mm10) was performed using HISAT2 (v2.1.0) [32], and expression

quantification was performed using featureCounts (v2.0.3) [33]. All commands were run with

default settings and parameters. DEGs were detected using the R Package "edgeR" (v3.40.2)

[34]. Information on the statistical significance used for identification of DEGs is detailed in

the Statistical Analyses section.

Identification of chemical- or pollutant-induced DARs

Referring to the SRA accession IDs for the chemical- or pollutant-perturbed ATAC-Seq data

obtained from DBKERO and TaRGET, the peak call data (in BED format) for the corresponding

ATAC-Seq experiments were acquired from ChIP-Atlas. Peak calling was performed using

MACS2 [27] in ChIP-Atlas; only peaks with a Q-value of $<1 \times 10^{-10}$ were considered as

statistically significant accessible chromatin regions. Next, common open chromatin regions

across all replicates for each chemical or pollutant perturbation condition were extracted

using BEDtools (v2.23.0) [35]. Differential analysis was then performed to identify DARs

induced by the chemicals or pollutants. Note that the term "differential analysis" in this

context refers to the identification of genomic regions with no overlap between groups with

and without chemical or pollutant exposure. In this study, DARs that exhibited relaxed or

condensed chromatin upon chemical or pollutant exposure are referred to as "opened" or

"closed" DARs, respectively.

TF enrichment analysis fully incorporating public ChIP-Seq experiments (ChIPEA)

Details of the ChIPEA procedure using the application programming interface are provided

online [22, 36]. To exploit the extensive ChIP-Seq data in ChIP-Atlas [23], we performed

ChIPEA on profile TFs whose binding sites were enriched around genomic regions or genes

of interest. In particular, using the genomic coordinates of opened and closed DARs, or the

gene symbols of up- and down-regulated genes by a query chemical or pollutant as input,

we first counted the number of intersects between the input data and peak-call data

(MACS2; Q-value $<1 \times 10^{-10}$) of all TF-related experiments archived in the ChIP-Atlas

database using the "intersectBed" command of BEDTools (v2.23.0) [35]. Note that 5 kb

upstream and downstream of the transcription start site were added for each gene when

chemically induced DEGs were queried. Enrichment scores ($-\log_{10}Q$) were then calculated

using the two-tailed Fisher's exact probability test to test whether the two datasets (opened

and closed DARs, or up- and down-regulated genes) overlapped with the ChIP-Seq peak-call

data with equivalent proportions before performing the Benjamini-Hochberg procedure for

multiple testing correction; fold enrichment values (opened DARs/closed DARs) were

concurrently obtained. If a chemical– or pollutant–TF association was indicated by multiple

ChIP-Seq experiments, only the highest enrichment score was adopted. Parameters used in

application programming interface-based ChIPEA were as follows: "genome=hg38 (DBKERO)

or mm10 (TaRGET); antigenClass=TFs and others; cellClass=All cell types; threshold=100."

Sample codes for DEG-ChIPEA and DAR-ChIPEA are provided as indicated in the Availability

of Data and Materials statement below.

TF enrichment analysis focusing on TF–DNA binding motifs (MotifEA)

In this method, TF binding to chemical- or pollutant-induced DARs and DEGs was estimated

using MotifEA with STREME and TOMTOM in MEME Suite (v5.4.1) [21], using default settings

termed DAR-MotifEA and DEG-MotifEA, respectively. STREME was first run to identify 1,000

sequence motifs from queried multiple genomic regions, i.e., chemical- or pollutant-induced

DARs or DEGs. Subsequently, TFs with enriched binding to the input genomic regions or

genes were estimated by matching the identified motifs with known TF binding motifs

obtained from HOCOMOCO v11 [37] using TOMTOM. Note that 5 kb upstream and

downstream of the transcription start site were added for each gene when chemically

induced DEGs were queried. Sample codes for DEG-MotifEA and DAR-MotifEA are provided

as indicated in the Availability of Data and Materials statement below.

Establishment of chemical– or pollutant–TF–disorder triadic associations

Using TFs enriched in chemical- or pollutant-induced DEGs and DARs as mediators, the TF–

disease associations retrieved from DisGeNET [28] were integrated with the chemical– or

pollutant–TF matrices obtained as outcomes of MotifEA or ChIPEA, thereby establishing

chemical– or pollutant–TF–disorder triadic associations. The enrichment scores obtained via

ChIPEA or MotifEA were continuously used as a quantitative measure to assess the degree of

newly constructed chemical– or pollutant–disorder associations.

Calculation of global area under the receiver operating characteristic curve (AUROC)

and area under the precision recall curve (AUPR) scores

All predicted chemical– or pollutant–disorder associations were first reorganized into a single

matrix containing enrichment scores with dimensions of $m \times n$, where m and n represent the

total number of chemical or pollutant perturbation conditions and the number of disorders,

respectively (Additional file 1: Supplementary Fig. S1). Among various TFs linking a specific

chemical–disease pair, we selectively adopted the most statistically significant TF, i.e., the one

with the highest enrichment score. We then utilized the maximum enrichment score as a

statistical measure to represent the association between this chemical and the disease. After

referencing known chemical–disease associations from CTD [29], chemical-wise ROC (which

plots the true positive rate against the false positive rate) and PR (which plots the precision

[positive predictive value] against the recall [sensitivity]) curves were generated (R Package

"ROCR" v1.0-11) [38]. The performance results were summarized into a global AUROC score,

in which 1 represents perfect classification and 0.5 represents random classification, and an

AUPR score, in which 1 represents perfect inference and the proportion of positive examples

in the standard data corresponds to random inference.

Mouse phenotype ontology enrichment analysis

GO enrichment analysis on opened DARs induced by pollutant exposure was performed

using the Genomic Regions Enrichment of Annotations Tool (GREAT) (v4.0.4) [39]. Default

parameters were used in this analysis; namely, "Species assembly: mm10; Association rule:

Basal+extension: 5000 bp upstream, 1000 bp downstream, 1000000 bp max extension,

curated regulatory domains included." Ontologies categorized as "Mouse Phenotype" were

then extracted. Parent category information for individual ontology terms was obtained with

reference to the Mouse Genome Informatics database [40].

Statistical Analyses

Statistical analysis was performed using R (v4.1.3). Unpaired two-tailed Wilcoxon rank-sum

tests were used to compare AUROC and AUPR scores between two groups, Bonferroni

correction was performed to presented adjusted P-values (Q-values). Fisher's exact test was

used for 2 × 2 contingency analysis, and the Benjamini-Hochberg procedure was used for

multiple testing correction. Sample sizes and Q-values are shown in the figures, and a Q-

value < 0.05 was considered statistically significant. NS denotes no significance. Chemical-

and pollutant-induced DEGs were identified using the R package edgeR (v3.40.2), and a P-

value <0.05 (DBKERO) or an adjusted P-value <0.1 (TaRGET) with an absolute value of $\log_2$

fold change >1 was recognized as statistically significant.

RESULTS

Study Design: Elucidation of Chemical MoAs Focusing on Chemically Reorganized

Chromatin Accessibility

To expand upon our prior approach (DEG-ChIPEA) [22], here, we present a novel analytical

approach to elucidate chemical MoAs focusing on chemically reorganized chromatin

assembly. Specifically, we utilize chemically induced DARs as input to cover the full spectrum

of TF binding to both proximal and distal transcription regulatory regions (e.g., promoters

and enhancers; Fig. 1a). Our approach identifies TFs that integratively regulate chemical- or

pollutant-induced DARs by comprehensively analyzing tens of thousands of publicly

available ChIP-Seq data obtained from the ChIP-Atlas database (Fig. 1b) [23]. In addition, we

integrated the predicted chemical–TF associations with disease-associated TF data obtained

from the DisGeNET database [28]. The resulting chemical–TF–disorder triadic associations

were then statistically validated using a known chemical–disease association dataset from

CTD as standard data (Additional file 1: Supplementary Fig. S1) [29].

Validation of the Proposed DAR-ChIPEA Approach to Predict TF-Driven Actions for _In_

Vitro Chemical Perturbation

Because TaRGET [25] contains information on only a few types of pollutants, it is not

sufficient to statistically evaluate the reliability of DAR-ChIPEA. To validate the proposed

DAR-ChIPEA approach to predict TF-driven actions for *in vitro* chemical perturbation, we first

evaluated the prediction accuracy of our method using the DBKERO database [24], which

contains data on hundreds of chemical perturbations to cell cultures, providing correlated

RNA-Seq and ATAC-Seq datasets. After retrieving pre-processed peak call datasets of ATAC-

Seq from cell lines with and without chemical treatment in the DBKERO database from ChIP-

Atlas, we performed a comparative analysis to detect DARs opened or closed by chemical

perturbation (Additional file 1: Supplementary Fig. S2). Next, we fully exploited almost all

publicly available human TF ChIP-Seq data in ChIP-Atlas to identify TFs enriched for binding

to the extracted DARs using ChIPEA (DAR-ChIPEA). This allowed us to generate chemical–TF

associations for each drug treatment condition. Furthermore, to evaluate the pivotality of TFs

in drug MoAs, chemical–TF–disorder associations were constructed by assigning TF-related

disease information obtained from DisGeNET to the resulting chemical–TF association matrix.

For comparison, in the chemical–TF association generation step, we also used chemically

induced DEGs as input to ChIPEA (DEG-ChIPEA), as previously reported (Additional file 1:

Supplementary Fig. S2) [22]. For the same purpose, a more conventional and commonly used

TF enrichment analysis method focusing on TF–DNA binding motifs (MotifEA) was also

performed with DEGs or DARs as input (hereafter referred to as DEG-MotifEA and DAR-

MotifEA; Fig. 2a).

We observed that the statistically significant maximum enrichment scores ($-\log_{10}Q$-value)

obtained from the proposed DAR-ChIPEA method were globally higher than those from our

previously developed DEG-ChIPEA method (Fig. 2b; mean maximum enrichment score =

10.25 and 8.139, respectively; $P = 3.013 \times 10^{-4}$; two-tailed Wilcoxon rank-sum test),

indicating that DAR-ChIPEA has a wider dynamic range than DEG-ChIPEA. Thus, DAR-ChIPEA

offers a significant advantage in discriminating associated and non-associated TFs with

regard to individual chemicals.

We then examined whether the chemical–TF–disorder triadic associations derived from

DAR-ChIPEA that exhibited higher statistical significance were congruent with existing

knowledge. The distribution of AUROC and AUPR scores for each dosing condition,

generated using manually curated chemical–disease association data obtained from CTD as

reference data (1,457, 1,711, 1,621, and 1,800 conditions for DEG-MotifEA, DAR-MotifEA,

DEG-ChIPEA, and DAR-ChIPEA, respectively), was depicted using bee swarm plots (Fig. 2c, d;

Additional file 2: Supplementary Table S5). The mean AUROC and AUPR scores of the DAR-

ChIPEA approach across all dosage conditions were 0.7287 and 0.4164, respectively. These

scores were significantly higher than those of other methods performed for comparison

(DEG-MotifEA: mean AUROC 0.6736; mean AUPR 0.3488, DAR-MotifEA: mean AUROC

0.6769; mean AUPR 0.3641, and DEG-ChIPEA: mean AUROC 0.7060; mean AUPR 0.3862;

$P<0.05$ for all; two-tailed Wilcoxon rank-sum test).

In addition, because the maximum enrichment scores varied drastically depending on the

given dosing conditions, we were concerned that the discrimination of items was limited

when comparing TFs within a dosing condition with low maximum enrichment scores.

Therefore, we also calculated "global" statistics using an inter-chemical merged enrichment

score vector to emphasize the significance of the actual values of the enrichment scores

(detailed in the Methods section). The global AUROC and AUPR scores of the proposed

approach were 0.7894 and 0.6137, respectively, significantly outperforming those of other

methods ($P<0.05$; two-tailed Wilcoxon rank-sum test; Fig. 2c, d). These results suggest that

the ChIPEA method using chemically induced DARs as input is a powerful approach for

identifying pivotal TFs, elucidating key factors for specific chemical MoAs and predicting

disorders or phenotypes associated with chemical treatments. The use of this approach may

ultimately lead to increased predictive power compared to that of the existing DEG-ChIPEA

and MotifEA-based computational methods.

Application of the DAR-ChIPEA Approach to Predict TF-Driven MoAs upon Pollutant

Administration *In Vivo*

We next employed the DAR-ChIPEA method to predict disorders caused by pollutants and

TF-driven MoAs upon pollutant administration *in vivo*. After retrieving ATAC-Seq peak call

datasets obtained with and without pollutant exposure from the TaRGET database available

in ChIP-Atlas, we identified opened and closed DARs induced by various pollutants, as

described in the Methods section (Additional file 1: Supplementary Fig. S2). We used publicly

available mouse TF ChIP-Seq data in ChIP-Atlas to generate pollutant–TF associations using

the proposed DAR-ChIPEA approach. We simultaneously performed analyses using the other

methods for comparison, namely, DEG-ChIPEA, DEG-MotifEA, and DAR-MotifEA (Fig. 3a). The

resulting pollutant–TF matrices were then used to generate pollutant–TF–disorder

associations, focusing on pivotal TFs, with reference to TF–disease associations from the

DisGeNET database. We evaluated the accuracy of the predicted pollutant–disorder

associations using ROC and PR curves using known chemical-disease association datasets

from CTD as a standard.

The maximum enrichment scores attained using the proposed DAR-ChIPEA approach

were globally higher than those achieved with our previously developed DEG-ChIPEA

method, similar to the results obtained using *in-vitro* input data, further supporting the wider

dynamic range for discrimination of associated TFs for individual pollutants (Fig. 3b; mean

maximum enrichment score = 76.77 and 11.57, respectively; $P = 8.421 \times 10^{-8}$; two-tailed

Wilcoxon rank-sum test).

In addition, we utilized a large set of publicly available Bisulfite-Seq experimental data

(mm10; $n = 34{,}793$) integrated into ChIP-Atlas to assess the DNA methylation status within

the DARs induced by representative pollutants. The results revealed that opened and closed

DARs were enriched by either hyper- or hypo-methylated regions, suggesting DNA

methylation change within pollutant-induced DARs (Additional file 1: Supplementary Fig. S3).

The distribution of AUROC and AUPR scores for each exposure condition (40, 28, 40, and

21 conditions for DEG-MotifEA, DAR-MotifEA, DEG-ChIPEA, and DAR-ChIPEA, respectively)

was visualized using bee swarm plots (Fig. 3c, d; Additional file 2: Supplementary Table S6).

The mean AUROC of the proposed DAR-ChIPEA approach across exposure conditions was

0.6748, significantly higher than those of the other methods (DEG-MotifEA, 0.6071; DAR-

MotifEA, 0.6147; DEG-ChIPEA, 0.6575; $P<0.05$; two-tailed Wilcoxon rank-sum test). However,

the DAR-ChIPEA AUPR score did not significantly ($P>0.05$) exceed those of the other

methods (DEG-MotifEA, 0.7656; DAR-MotifEA, 0.7908; DEG-ChIPEA, 0.7890; proposed: DAR-

ChIPEA, 0.7929). The global AUROC and global AUPR scores of DAR-ChIPEA were 0.7492 and

0.8491, respectively. These results suggest that the DAR-ChIPEA method performed

satisfactorily in terms of AUROC; the absence of significant differences regarding AUPR may

be attributed to the limited number of datasets analyzed.

Biological Interpretation of the Predicted Pollutant–TF–Disorder Associations

In the proposed DAR-ChIPEA method, high AUROC scores were obtained for tributyltin (TBT)

(blood samples), $PM_{2.5}$ (blood samples), and lead (liver samples; AUROC = 0.7028, 0.7030,

and 0.7349, respectively; Additional file 2: Supplementary Table S4, S6; highlighted with bold

red font), which motivated us to evaluate the results of putative pivotal TFs involved in the

MoAs of these pollutants in further detail, particularly with regard to the biological

interpretation of the predicted pollutant–TF–disorder associations (Fig. 4; Additional file 2:

Supplementary Tables S7–S9).

The organotin compound TBT is widely utilized in various industrial applications such as

pesticides, paints, preservatives, and heat stabilizers for plastics. Nevertheless, TBT has been

associated with a potentially increased risk of a number of lifestyle-related diseases,

including obesity, fatty liver, cardiovascular disease, and abnormal glucose metabolism [41–

44]. We initially conducted ontology enrichment analysis to discern any common nature of

genes located proximal to opened DARs of blood samples upon TBT exposure using

mammalian phenotype (MP) ontology. Notably, we found that the enriched gene sets were

prominently manifested in phenotypic features of the immune system and hematopoietic

system (upper left panel in Fig. 4). To further understand the transcriptional regulatory

mechanisms involved in the response to TBT exposure, we then performed DAR-ChIPEA

(upper right panel in Fig. 4). The genes encoding the top-20 enriched TFs exhibited robust

expression in both control and TBT-exposed samples, and the same applied for $PM_{2.5}$ and

lead, as detailed below (Additional file 1: Supplementary Fig. S4). The results of DAR-ChIPEA

showed that the opening of specific DARs susceptible to TBT exposure could enhance the

enriched binding of Pparg, a master regulator of adipogenesis [45]. This is consistent with

the *in-vitro* findings that TBT stimulates lipid accumulation by activating Pparg [46].

In addition, TBT can induce apoptosis in diverse tissues and cell categories, such as

human peripheral blood lymphocytes and human amnion cells, and high-dose exposure to

TBT has been reported to induce T lymphocyte apoptosis and thymic atrophy [47–49].

Furthermore, TBT suppresses the differentiation of naive CD4[+] cells into Th1 cells, while

stimulating differentiation into Th2 cells, thereby causing allergic disorders through the

overproduction of Th2 cells, although the MoAs remain incompletely understood [50]. Our

results showed that TBT exposure led to the enrichment of TFs such as Rela, Stat1, and Rbpj

in opened DARs, consistent with the pro-inflammatory nature of TBT. Conversely, the binding

of Bcl11b and Tbx21 was markedly enriched in closed DARs. BCL11B has apoptosis-inhibitory

properties and is involved in maintaining T cell differentiation, as supported by the reduced

numbers of T cells observed in *Bcl11b*-deficient mice [51, 52]. Notably, TBX21 also plays a

key role in Th1 cell differentiation, with *Tbx21*-deficient naive Th cells exhibiting distorted

Th2 cell polarization, a phenotype somewhat similar to that observed upon TBT exposure

[53, 54]. These findings suggest that Bcl11b and Tbx21 may contribute to TBT-induced

disruption of the Th1/Th2 balance.

$PM_{2.5}$ is fine particulate matter suspended in the atmosphere with a diameter of 2.5 μm

or less, which has the ability to transport pollutants in the environment. Exposure to $PM_{2.5}$ is

associated with respiratory diseases such as asthma, bronchitis, and pneumonia, as well as

heart diseases such as arrhythmias and atherosclerosis, and significantly increases mortality

from lung cancer [55–60]. Moreover, $PM_{2.5}$ can worsen bacterial infections such as

pneumonia caused by *Streptococcus pneumoniae* by impairing the function of macrophages

in the lungs [61]. Oxidative and inflammatory damage and increased cellular autophagy and

apoptosis have been implicated in the induction of various diseases by $PM_{2.5}$; however, the

underlying MoAs remain to be elucidated. Notably, the results of the MP ontology

enrichment analysis for $PM_{2.5}$ exhibited strong similarity to those of TBT, highlighting the

prominent enrichment of phenotypes related to the immune and hematopoietic systems

following $PM_{2.5}$ exposure, consistent with previous research (middle left panel in Fig. 4) [62].

Furthermore, the results of DAR-ChIPEA revealed that the binding of blood cell

differentiation-related factors such as Cebpe, Cebpb, Cebpa, Rela, and Spi1 was enriched in

the DARs of blood samples closed by $PM_{2.5}$ exposure, i.e., showed binding in the unexposed

samples (middle right panel in Fig. 4). Among the enriched TFs, the C/EBP family members

Cebpe and Cebpb are essential factors for differentiation and functional maintenance of

granulocytes and the monocyte–macrophage axis, respectively, whereas Cebpa contributes

to the regulation of growth arrest and differentiation of various cell types, including

progenitor cells in bone marrow [63–65]. In addition, Rela is important to erythropoiesis, and

Spi1 is important to early differentiation of B cells and bone marrow cells [66, 67].

Lead (Pb) has historically been used in a variety of applications, including paints and

cosmetic dyes, and remains a common component of water pipes, solder, and other

materials. Pb poisoning is typically a chronic condition that may not present with acute

symptoms. Regardless of the absence of acute symptoms, Pb poisoning ultimately exhibits

irreversible effects such as cognitive impairment, peripheral neuropathy, and progressive

renal dysfunction [68–70]. Chronic Pb exposure may also induce fatty liver disease, which

may be associated with changes in the gut microbiota [71]. Consistent with these

observations, MP ontology enrichment analysis on liver samples exposed to Pb corroborated

the significant enrichment of genes associated with the regulation of liver physiology, as well

as lipid homeostasis and metabolism, in the proximity of DARs induced by Pb administration

(lower left panel in Fig. 4). DAR-ChIPEA further revealed that Nr1d1 (also known as Rev-

erbα), an important member of the feedback loop regulating the circadian clock composed

of Clock and Arntl (also known as Bmal1) [72], was the most enriched TF in Pb-induced

opened DARs. In addition, Ep300 and CREB-binding protein (Crebbp), transcriptional

coactivators and members of the Clock/Bmal1 activator complex, were also found to be

enriched in opened DARs (lower right panel in Fig. 4). These findings suggest that circadian

rhythms in the liver may be disrupted by Pb exposure. Moreover, clock genes have been

shown to act on the promoter and introns of adipogenic Ppara [73, 74]. In the liver, Ppara

enhances lipid metabolism and is generally considered to have an inhibitory effect on

hepatic steatosis, although some reports suggest that Ppara may promote the pathogenesis

of non-alcoholic steatohepatitis [75, 76]. These findings, together with those from MP

ontology analysis, suggest that Pb induces fatty liver not only through the gut microbiota

pathway, but also by disrupting the normal regulation of lipid metabolism by circadian

rhythms in the liver.

DISCUSSION

In this study, we used the ATAC-Seq dataset, derived from samples perturbed by

environmental chemicals, from public databases to identify pivotal TFs involved in chemical

MoAs. We extracted DARs and applied ChIPEA, taking full advantage of large-scale ChIP-Seq

experiments, to identify TFs that were enriched for binding to these genomic regions. In

addition, we integrated the TF-associated disease dataset with the putative chemical–TF

associations to estimate key TFs mediating chemical-induced disorder induction and in turn

understand chemical MoAs. The proposed DAR-ChIPEA approach outperformed other

existing computational methods in terms of accuracy.

In general, when cells or tissues are exposed to chemical substances, a series of events

occur, initialized by pioneer factors, which are TFs that modulate chromatin accessibility.

Subsequently, various cooperative TFs, often acting in complexes (e.g., Arntl/Ep300/Cebp in

Fig. 4), bind to the newly opened or closed genome regions, causing alterations in the

expression of genes regulated by these TFs. Moreover, the DNA methylation status

undergoes modifications, contributing to the fixation of gene expression alteration. This

entire cascade ultimately leads to phenotypic changes induced by chemical substances. The

proposed method can provide insights into identifying key DNA-binding proteins involved in

the MoAs of chemicals, including both pioneer factors and cooperative TFs. In addition, the

public Bisulfite-Seq dataset in our previously developed ChIP-Atlas database can also be

utilized to infer DNA methylation status change upon chemical exposure (Supplementary

figure S2).

When dealing with health hazards caused by air pollution or industrial wastewater, it can

be difficult to isolate a single pollution-causing molecule from such complex contexts.

Application of approaches that focus on the detailed molecular structure of chemicals to

analyze their MoAs, such as docking simulation and supervised learning using molecular

structure as features, can thus be challenging [8–10, 77]. In this respect, the enrichment

analysis-based approaches presented in this study are advantageous in that they use omics

data as input and thus are better suited to analyze complex mixtures. In parallel with the

processes used for causative molecule isolation, in the proposed DAR-ChIPEA method,

chemically induced DAR information is obtained from experiments involving exposure to

polluted air and water and passed through the proposed analysis pipeline, allowing for rapid

and rough estimation of the key TFs involved in the MoAs of pollutants. This approach is

expected to serve as an efficient complementary approach for investigating the causes of

clinically-relevant pollution damage.

We found that the ChIPEA approach, an enrichment analysis method based on actual

ChIP-Seq experiments, was more accurate than MotifEA, a numerical analysis method based

on DNA binding motifs of TFs, in estimating chemical–disorder associations via key TFs. The

moderate performance of MotifEA likely occurred, in part, because the presence of a

particular binding motif does not necessarily indicate the actual binding of a TF, and

conversely, it is not uncommon for a TF to bind to genomic regions that are distinct from

known motifs. Furthermore, binding motifs can vary greatly depending on the tissue or cell

type. In comparison, ChIPEA is highly reliable for studying the interaction between TFs and

genomic DNA because it extracts TF binding sites in the genome based on a large number of

ChIP-Seq experiments reported in the literature in a motif-independent manner [22, 36].

However, it should be acknowledged that the number of ChIP-Seq data currently available

varies considerably across different tissues and cell types, and thus, ChIPEA may be less

effective in situations in which available data are insufficient to generate statistically

significant results.

Notably, the pivotal TFs found using our proposed approach may also serve as potential

targets for drug discovery aimed at mitigating health hazards posed by environmental

pollutants. Therefore, our approach is not only useful for elucidating the MoAs of

environmental chemicals, but also for screening candidate compounds for potential

pharmaceuticals. Owing to its ability to achieve an adequate depth of analysis with as few as

500 cells, ATAC-Seq provides the possibility to perform large-scale analyses of various

compounds using small-well plates [78]. RNA-Seq may also be employed simultaneously to

evaluate the pharmacological effects of the input compounds. In addition to the DBKERO

database used in this study, other databases are also available that focus on predicting the

medicinal effects and toxicity of various chemicals by analyzing changes in gene signatures

upon chemical perturbation. For example, the LINCS L1000 project and Open TG-GATEs have

obtained extensive transcriptome data on tens of thousands of drugs applied to various cell

lines and laboratory animals [79, 80]. To further clarify the molecular cues that mediate gene

expression changes in response to chemical administration, ATAC-Seq data should be

incorporated into the compound screening pipeline in these projects, thereby facilitating the

effective application of the proposed DAR-ChIPEA method. Furthermore, because ChIPEA is

already publicly available through the ChIP-Atlas website in both a user-friendly graphical

user interface and a command-based application programming interface, it could be easily

applied in various situations to elucidate drug MoAs. Realization of this potential application

could help identify TFs that are critical for the expression of drug effects, allowing the

formulation of strategies to minimize unexpected risk, especially for compounds with

potential adverse effects. In addition, identification of novel targets will allow the

repositioning of drugs that have successfully passed safety testing as therapeutic agents for

other diseases. Thus, the proposed DAR-ChIPEA approach could also contribute to lower

drug discovery costs and a more efficient drug repositioning in the future.

CONCLUSIONS

In this paper, we introduced a computational approach for highlighting genome-wide

chromatin change upon pollutant exposure to elucidate the epigenetic landscape of

pollutant responses, which outperformed other methods focusing on gene-adjacent

domains only. The proposed method can be adopted to identify pivotal TFs involved in the

MoAs of pollutants from an epigenetic perspective, thereby facilitating the development of

strategies to mitigate environmental pollution damage. The proposed DAR-ChIPEA approach

can also be used to further understand the MoAs of candidate drug compounds as well as to

discover unexpected therapeutic and side effects of approved drugs to reduce costs during

drug discovery.

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

We used freely available datasets; specific version numbers or dates for accessed or

downloaded datasets are indicated as appropriate in the reference list. Sample codes and all

raw results of the TF enrichment analyses (i.e., DEG-MotifEA, DAR-MotifEA, DEG-ChIPEA, and

DAR-ChIPEA) were deposited at https://github.com/zouzhaonan/dar_moa/.

Competing interests

The authors declare that they have no competing interests. The funders had no role in the

design of the study, the collection, analysis, and interpretation of data, or in writing the

manuscript.

Funding

This work was supported by Grants-in-Aid for Scientific Research (KAKENHI), Japan Society

for the Promotion of Science [22J15229 to ZZ, 23KF0048 to ZZ and SO, 22H02819, 23H04954

to SO]; Japan Science and Technology Agency [SPRING JPMJSP2110 to ZZ, PRESTO

JPMJPR1942, NBDC JPMJND2202, and ERATO JPMJER2101 to SO]; Japan Agency for Medical

Research and Development [PRIME JP22gm6710001, BINDS JP22ama121017 to SO], and

Kyoto University [MIP and KMS-FUND to ZZ]. Funding for open access charge: Japan Science

and Technology Agency [JPMJND2202].

Authors' contributions

Shinya Oki: Conceptualization, Supervision, Data curation, Funding acquisition, Writing–

original draft, Writing–review & editing. Zhaonan Zou: Formal analysis, Funding acquisition,

Writing–original draft, Writing–review & editing. Yuka Yoshimura: Formal analysis. Yoshihiro

Yamanishi: Validation, Writing–original draft, Writing–review & editing.

Acknowledgements

Computations were partially performed on the NIG supercomputer at ROIS National Institute

of Genetics, Japan. We thank Prof. Shuh Narumiya and the members of the Department of

Drug Discovery Medicine at Kyoto University for providing us with valuable advice.

REFERENCES

1. Lee SW, Yon DK, James CC, Lee S, Koh HY, Sheen YH et al. Short-term effects of multiple

outdoor environmental factors on risk of asthma exacerbations: Age-stratified time-series

analysis. J Allergy Clin Immunol 2019;144(6):1542–1550 e1541.

https://doi.org/10.1016/j.jaci.2019.08.037.

2. Kaufman JD, Adar SD, Barr RG, Budoff M, Burke GL, Curl CL et al. Association between air

pollution and coronary artery calcification within six metropolitan areas in the USA (the Multi-

Ethnic Study of Atherosclerosis and Air Pollution): a longitudinal cohort study. Lancet

2016;388(10045):696–704. https://doi.org/10.1016/S0140-6736(16)00378-0.

3. Yang BY, Qian Z, Howard SW, Vaughn MG, Fan SJ, Liu KK et al. Global association between

ambient air pollution and blood pressure: A systematic review and meta-analysis. Environ Pollut

2018;235:576–588. https://doi.org/10.1016/j.envpol.2018.01.001.

4. Zhu RX, Nie XH, Chen YH, Chen J, Wu SW, Zhao LH. Relationship Between Particulate Matter

(PM(2.5)) and Hospitalizations and Mortality of Chronic Obstructive Pulmonary Disease

Patients: A Meta-Analysis. Am J Med Sci 2020;359(6):354–364.

https://doi.org/10.1016/j.amjms.2020.03.016.

5.   Wong CM, Tsang H, Lai HK, Thomas GN, Lam KB, Chan KP et al. Cancer Mortality Risks from

Long-term Exposure to Ambient Fine Particle. Cancer Epidemiol Biomarkers Prev

2016;25(5):839–845. https://doi.org/10.1158/1055-9965.EPI-15-0626.

6.   Gianicolo EA, Bruni A, Rosati E, Sabina S, Guarino R, Padolecchia G et al. Congenital anomalies

among live births in a polluted area. A ten-year retrospective study. BMC Pregnancy Childbirth

2012;12:165. https://doi.org/10.1186/1471-2393-12-165.

7.   Mahalingaiah S, Hart JE, Laden F, Farland LV, Hewlett MM, Chavarro J et al. Adult air pollution

exposure and risk of infertility in the Nurses' Health Study II. Hum Reprod 2016;31(3):638–647.

https://doi.org/10.1093/humrep/dev330.

8.   Bosc N, Atkinson F, Felix E, Gaulton A, Hersey A, Leach AR. Large scale comparison of QSAR and

conformal prediction methods and their applications in drug discovery. J Cheminform

2019;11(1):4. https://doi.org/10.1186/s13321-018-0325-4.

9.   Jacob L, Vert JP. Protein-ligand interaction prediction: an improved chemogenomics approach.

Bioinformatics 2008;24(19):2149–2156. https://doi.org/10.1093/bioinformatics/btn409.

10.   Kolb P, Ferreira RS, Irwin JJ, Shoichet BK. Docking and chemoinformatic screens for new ligands

and targets. Curr Opin Biotechnol 2009;20(4):429–436.

https://doi.org/10.1016/j.copbio.2009.08.003.

11.   Pham TH, Qiu Y, Liu J, Zimmer S, O'Neill E, Xie L et al. Chemical-induced gene expression ranking

and its application to pancreatic cancer drug repurposing. Patterns (N Y) 2022;3(4):100441.

https://doi.org/10.1016/j.patter.2022.100441.

12.   Pilarczyk M, Fazel-Najafabadi M, Kouril M, Shamsaei B, Vasiliauskas J, Niu W et al. Connecting

omics signatures and revealing biological mechanisms with iLINCS. Nat Commun

2022;13(1):4678. https://doi.org/10.1038/s41467-022-32205-3.

13. Lee CW, Kim SM, Sa S, Hong M, Nam SM, Han HW. Relationship between drug targets and drug-

signature networks: a network-based genome-wide landscape. BMC Med Genomics

2023;16(1):17. https://doi.org/10.1186/s12920-023-01444-8.

14. DALYs GBD, Collaborators H. Global, regional, and national disability-adjusted life-years (DALYs)

for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and

territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017.

Lancet 2018;392(10159):1859-1922. https://doi.org/10.1016/S0140-6736(18)32335-3.

15. Quenby S, Gallos ID, Dhillon-Smith RK, Podesek M, Stephenson MD, Fisher J et al. Miscarriage

matters: the epidemiological, physical, psychological, and economic costs of early pregnancy

loss. Lancet 2021;397(10285):1658-1667. https://doi.org/10.1016/S0140-6736(21)00682-6.

16. Go S, Kurita H, Matsumoto K, Hatano M, Inden M, Hozumi I. Methylmercury causes epigenetic

suppression of the tyrosine hydroxylase gene in an in vitro neuronal differentiation model.

Biochem Biophys Res Commun 2018;502(4):435-441.

https://doi.org/10.1016/j.bbrc.2018.05.162.

17. Huang D, Zhang Y, Qi Y, Chen C, Ji W. Global DNA hypomethylation, rather than reactive oxygen

species (ROS), a potential facilitator of cadmium-stimulated K562 cell proliferation. Toxicol Lett

2008;179(1):43-47. https://doi.org/10.1016/j.toxlet.2008.03.018.

18. van Tilburg CM, Milde T, Witt R, Ecker J, Hielscher T, Seitz A et al. Phase I/II intra-patient dose

escalation study of vorinostat in children with relapsed solid tumor, lymphoma, or leukemia.

Clin Epigenetics 2019;11(1):188. https://doi.org/10.1186/s13148-019-0775-1.

19. Bardia A, Kaklamani V, Wilks S, Weise A, Richards D, Harb W et al. Phase I Study of Elacestrant

(RAD1901), a Novel Selective Estrogen Receptor Degrader, in ER-Positive, HER2-Negative

Advanced Breast Cancer. J Clin Oncol 2021;39(12):1360-1370.

https://doi.org/10.1200/JCO.20.02272.

20.   Duttke SH, Chang MW, Heinz S, Benner C. Identification and dynamic quantification of

regulatory elements using total RNA. Genome Res 2019;29(11):1836-1846.

https://doi.org/10.1101/gr.253492.119.

21.   Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. Nucleic Acids Res

2015;43(W1):W39-49. https://doi.org/10.1093/nar/gkv416.

22.   Zou Z, Iwata M, Yamanishi Y, Oki S. Epigenetic landscape of drug responses revealed through

large-scale ChIP-seq data analyses. BMC Bioinformatics 2022;23(1):51.

https://doi.org/10.1186/s12859-022-04571-8.

23.   Zou Z, Ohta T, Miura F, Oki S. ChIP-Atlas 2021 update: a data-mining suite for exploring

epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and Bisulfite-seq data. Nucleic

Acids Research 2022;50(W1):W175–W182. https://doi.org/10.1093/nar/gkac199.

24.   Suzuki A, Kawano S, Mitsuyama T, Suyama M, Kanai Y, Shirahige K et al. DBTSS/DBKERO for

integrated analysis of transcriptional regulation. Nucleic Acids Res 2018;46(D1):D229-D238.

https://doi.org/10.1093/nar/gkx1001.

25.   Wang T, Pehrsson EC, Purushotham D, Li D, Zhuo X, Zhang B et al. The NIEHS TaRGET II

Consortium and environmental epigenomics. Nat Biotechnol 2018;36(3):225-227.

https://doi.org/10.1038/nbt.4099.

26.   Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence Database C. The

sequence read archive. Nucleic Acids Res 2011;39(Database issue):D19-21.

https://doi.org/10.1093/nar/gkq1019.

27.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE et al. Model-based analysis of

ChIP-Seq (MACS). Genome biology 2008;9(9):R137. https://doi.org/10.1186/GB-2008-9-9-R137.

28.   Pinero J, Ramirez-Anguita JM, Sauch-Pitarch J, Ronzano F, Centeno E, Sanz F et al. The DisGeNET

knowledge platform for disease genomics: 2019 update. Nucleic Acids Res 2020;48(D1):D845–

855. https://doi.org/10.1093/nar/gkz1021.

29. Davis AP, Wiegers TC, Johnson RJ, Sciaky D, Wiegers J, Mattingly CJ. Comparative

Toxicogenomics Database (CTD): update 2023. Nucleic Acids Res 2023;51(D1):D1257-D1262.

https://doi.org/10.1093/nar/gkac833.

30. SRA ToolKit v3.0.0. https://github.com/ncbi/sra-tools. Accessed 31 March, 2022.

31. FASTX-Toolkit v0.0.13. http://hannonlab.cshl.edu/fastx_toolkit/. Accessed 31 March 2022.

32. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and

genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 2019;37(8):907-915.

https://doi.org/10.1038/s41587-019-0201-4.

33. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. Bioinformatics 2014;30(7):923-930.

https://doi.org/10.1093/bioinformatics/btt656.

34. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics 2010;26(1):139-140.

https://doi.org/10.1093/bioinformatics/btp616.

35. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics 2010;26(6):841–842. https://doi.org/10.1093/bioinformatics/btq033.

36. Oki S, Ohta T, Shioi G, Hatanaka H, Ogasawara O, Okuda Y et al. ChIP-Atlas: a data-mining suite

powered by full integration of public ChIP-seq data. EMBO Rep 2018;19(12):e46255.

https://doi.org/10.15252/embr.201846255.

37. Kulakovskiy IV, Vorontsov IE, Yevshin IS, Sharipov RN, Fedorova AD, Rumynskiy EI et al.

HOCOMOCO: towards a complete collection of transcription factor binding models for human

and mouse via large-scale ChIP-Seq analysis. Nucleic Acids Res 2018;46(D1):D252-D259.

https://doi.org/10.1093/nar/gkx1106.

38. ROCR v1.0-11. .

39. McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB et al. GREAT improves functional

interpretation of cis-regulatory regions. Nat Biotechnol 2010;28(5):495-501.

https://doi.org/10.1038/nbt.1630.

40. Blake JA, Baldarelli R, Kadin JA, Richardson JE, Smith CL, Bult CJ et al. Mouse Genome Database

(MGD): Knowledgebase for mouse-human comparative biology. Nucleic Acids Res

2021;49(D1):D981-D987. https://doi.org/10.1093/nar/gkaa1083.

41. Xu J, Ou K, Chen C, Li B, Guo J, Zuo Z et al. Tributyltin exposure disturbs hepatic glucose

metabolism in male mice. Toxicology 2019;425:152242.

https://doi.org/10.1016/j.tox.2019.152242.

42. Chamorro-Garcia R, Diaz-Castillo C, Shoucri BM, Kach H, Leavitt R, Shioda T et al. Ancestral

perinatal obesogen exposure results in a transgenerational thrifty phenotype in mice. Nat

Commun 2017;8(1):2012. https://doi.org/10.1038/s41467-017-01944-z.

43. Zuo Z, Chen S, Wu T, Zhang J, Su Y, Chen Y et al. Tributyltin causes obesity and hepatic steatosis

in male mice. Environ Toxicol 2011;26(1):79-85. https://doi.org/10.1002/tox.20531.

44. Mendes ABA, Motta NAV, Lima GF, Autran LJ, Brazao SC, Magliano DC et al. Evaluation of the

effects produced by subacute tributyltin administration on vascular reactivity of male wistar

rats. Toxicology 2022;465:153067. https://doi.org/10.1016/j.tox.2021.153067.

45. Lehrke M, Lazar MA. The many faces of PPARgamma. Cell 2005;123(6):993-999.

https://doi.org/10.1016/j.cell.2005.11.026.

46. Li X, Ycaza J, Blumberg B. The environmental obesogen tributyltin chloride acts via peroxisome

proliferator activated receptor gamma to induce adipogenesis in murine 3T3-L1 preadipocytes.

J Steroid Biochem Mol Biol 2011;127(1-2):9-15. https://doi.org/10.1016/j.jsbmb.2011.03.012.

47. Ueno S, Kashimoto T, Susa N, Asai T, Kawaguchi S, Takeda-Homma S et al. Reduction in

peripheral lymphocytes and thymus atrophy induced by organotin compounds in vivo. J Vet

Med Sci 2009;71(8):1041-1048. https://doi.org/10.1292/jvms.71.1041.

48. Zhang Y, Liang J, Sun L, Guo Z, Xu L. Inhibition of PP2A and the consequent activation of JNK/c-

Jun are involved in tributyltin-induced apoptosis in human amnionic cells. Environ Toxicol

2013;28(7):390-400. https://doi.org/10.1002/tox.20730.

49. Tada-Oikawa S, Murata M, Kato T. [Preferential induction of apoptosis in regulatory T cells by

tributyltin: possible involvement in the exacerbation of allergic diseases]. Nihon Eiseigaku

Zasshi 2010;65(4):530-535. https://doi.org/10.1265/jjh.65.530.

50. Kato T, Uchikawa R, Yamada M, Arizono N, Oikawa S, Kawanishi S et al. Environmental pollutant

tributyltin promotes Th2 polarization and exacerbates airway inflammation. Eur J Immunol

2004;34(5):1312-1321. https://doi.org/10.1002/eji.200324667.

51. Grabarczyk P, Przybylski GK, Depke M, Volker U, Bahr J, Assmus K et al. Inhibition of BCL11B

expression leads to apoptosis of malignant but not normal mature T cells. Oncogene

2007;26(26):3797-3810. https://doi.org/10.1038/sj.onc.1210152.

52. Ikawa T, Hirose S, Masuda K, Kakugawa K, Satoh R, Shibano-Satoh A et al. An essential

developmental checkpoint for production of the T cell lineage. Science 2010;329(5987):93-96.

https://doi.org/10.1126/science.1188995.

53. Szabo SJ, Kim ST, Costa GL, Zhang X, Fathman CG, Glimcher LH. A novel transcription factor, T-

bet, directs Th1 lineage commitment. Cell 2000;100(6):655-669. https://doi.org/10.1016/s0092-

8674(00)80702-3.

54. Zhu J, Jankovic D, Oler AJ, Wei G, Sharma S, Hu G et al. The transcription factor T-bet is induced

by multiple pathways and prevents an endogenous Th2 cell program during Th1 cell responses.

Immunity 2012;37(4):660-673. https://doi.org/10.1016/j.immuni.2012.09.007.

55. Kim Y, Park EH, Ng CFS, Chung Y, Hashimoto K, Tashiro K et al. Respiratory function declines in

children with asthma associated with chemical species of fine particulate matter (PM(2.5)) in

Nagasaki, Japan. Environ Health 2021;20(1):110. https://doi.org/10.1186/s12940-021-00796-x.

56. Chen PC, Mou CH, Chen CW, Hsieh DPH, Tsai SP, Wei CC et al. Roles of Ambient Temperature

and PM(2.5) on Childhood Acute Bronchitis and Bronchiolitis from Viral Infection. Viruses

2022;14(9). https://doi.org/10.3390/v14091932.

57. Shi W, Liu C, Annesi-Maesano I, Norback D, Deng Q, Huang C et al. Ambient PM(2.5) and its

chemical constituents on lifetime-ever pneumonia in Chinese children: A multi-center study.

Environ Int 2021;146:106176. https://doi.org/10.1016/j.envint.2020.106176.

58. He F, Yanosky JD, Fernandez-Mendoza J, Chinchilli VM, Al-Shaar L, Vgontzas AN et al. Acute

Impact of Fine Particulate Air Pollution on Cardiac Arrhythmias in a Population-Based Sample of

Adolescents: The Penn State Child Cohort. J Am Heart Assoc 2022;11(18):e026370.

https://doi.org/10.1161/JAHA.122.026370.

59. Chaulin AM, Sergeev AK. The Role of Fine Particles (PM 2.5) in the Genesis of Atherosclerosis

and Myocardial Damage: Emphasis on Clinical and Epidemiological Data, and Pathophysiological

Mechanisms. Cardiol Res 2022;13(5):268-282. https://doi.org/10.14740/cr1366.

60. Huang F, Pan B, Wu J, Chen E, Chen L. Relationship between exposure to PM2.5 and lung cancer

incidence and mortality: A meta-analysis. Oncotarget 2017;8(26):43322-43331.

https://doi.org/10.18632/oncotarget.17313.

61. Chen YW, Huang MZ, Chen CL, Kuo CY, Yang CY, Chiang-Ni C et al. PM(2.5) impairs macrophage

functions to exacerbate pneumococcus-induced pulmonary pathogenesis. Part Fibre Toxicol

2020;17(1):37. https://doi.org/10.1186/s12989-020-00362-2.

62. Ge J, Yang H, Lu X, Wang S, Zhao Y, Huang J et al. Combined exposure to formaldehyde and

PM(2.5): Hematopoietic toxicity and molecular mechanism in mice. Environ Int

2020;144:106050. https://doi.org/10.1016/j.envint.2020.106050.

63. Yamanaka R, Barlow C, Lekstrom-Himes J, Castilla LH, Liu PP, Eckhaus M et al. Impaired

granulopoiesis, myelodysplasia, and early lethality in CCAAT/enhancer binding protein epsilon-

deficient mice. Proc Natl Acad Sci U S A 1997;94(24):13187-13192.

https://doi.org/10.1073/pnas.94.24.13187.

64. Tanaka T, Akira S, Yoshida K, Umemoto M, Yoneda Y, Shirafuji N et al. Targeted disruption of the

NF-IL6 gene discloses its essential role in bacteria killing and tumor cytotoxicity by

macrophages. Cell 1995;80(2):353-361. https://doi.org/10.1016/0092-8674(95)90418-2.

65. Reckzeh K, Bereshchenko O, Mead A, Rehn M, Kharazi S, Jacobsen SE et al. Molecular and

cellular effects of oncogene cooperation in a genetically accurate AML mouse model. Leukemia

2012;26(7):1527-1536. https://doi.org/10.1038/leu.2012.37.

66. Grossmann M, Metcalf D, Merryfull J, Beg A, Baltimore D, Gerondakis S. The combined absence

of the transcription factors Rel and RelA leads to multiple hemopoietic cell defects. Proc Natl

Acad Sci U S A 1999;96(21):11848-11853. https://doi.org/10.1073/pnas.96.21.11848.

67. McKercher SR, Torbett BE, Anderson KL, Henkel GW, Vestal DJ, Baribault H et al. Targeted

disruption of the PU.1 gene results in multiple hematopoietic abnormalities. EMBO J

1996;15(20):5647-5658. https://doi.org/.

68. Lee H, Lee MW, Warren JR, Ferrie J. Childhood lead exposure is associated with lower cognitive

functioning at older ages. Sci Adv 2022;8(45):eabn5164.

https://doi.org/10.1126/sciadv.abn5164.

69. Rubens O, Logina I, Kravale I, Eglite M, Donaghy M. Peripheral neuropathy in chronic

occupational inorganic lead exposure: a clinical and electrophysiological study. J Neurol

Neurosurg Psychiatry 2001;71(2):200-204. https://doi.org/10.1136/jnnp.71.2.200.

70. Lin JL, Tan DT, Hsu KH, Yu CC. Environmental lead exposure and progressive renal insufficiency.

Arch Intern Med 2001;161(2):264-271. https://doi.org/10.1001/archinte.161.2.264.

71. Wan H, Wang Y, Zhang H, Zhang K, Chen Y, Chen C et al. Chronic lead exposure induces fatty

liver disease associated with the variations of gut microbiota. Ecotoxicol Environ Saf

2022;232:113257. https://doi.org/10.1016/j.ecoenv.2022.113257.

72. Allada R, Bass J. Circadian Mechanisms in Medicine. N Engl J Med 2021;384(6):550-561.

https://doi.org/10.1056/NEJMra1802337.

73. Oishi K, Shirai H, Ishida N. CLOCK is involved in the circadian transactivation of peroxisome-

proliferator-activated receptor alpha (PPARalpha) in mice. Biochem J 2005;386(Pt 3):575-581.

https://doi.org/10.1042/BJ20041150.

74. Inoue I, Shinoda Y, Ikeda M, Hayashi K, Kanazawa K, Nomura M et al. CLOCK/BMAL1 is involved

in lipid metabolism via transactivation of the peroxisome proliferator-activated receptor (PPAR)

response element. J Atheroscler Thromb 2005;12(3):169-174.

https://doi.org/10.5551/jat.12.169.

75. Kasano-Camones CI, Takizawa M, Ohshima N, Saito C, Iwasaki W, Nakagawa Y et al. PPARalpha

activation partially drives NAFLD development in liver-specific Hnf4a-null mice. J Biochem 2023.

https://doi.org/10.1093/jb/mvad005.

76. Klaunig JE, Babich MA, Baetcke KP, Cook JC, Corton JC, David RM et al. PPARalpha agonist-

induced rodent tumors: modes of action and human relevance. Crit Rev Toxicol

2003;33(6):655–780. https://doi.org/10.1080/713608372.

77. Stanton DT. QSAR and QSPR model interpretation using partial least squares (PLS) analysis. Curr

Comput Aided Drug Des 2012;8(2):107–127. https://doi.org/10.2174/157340912800492357.

78. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin

for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and

nucleosome position. Nature methods 2013;10(12):1213–1218.

https://doi.org/10.1038/NMETH.2688.

79. Igarashi Y, Nakatsu N, Yamashita T, Ono A, Ohno Y, Urushidani T et al. Open TG-GATEs: a large-

scale toxicogenomics database. Nucleic Acids Res 2015;43(Database issue):D921–D927.

https://doi.org/10.1093/nar/gku955.

80. Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X et al. A Next Generation

Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell 2017;171(6):1437–1452

e1417. https://doi.org/10.1016/j.cell.2017.10.049.

Supplementary Information

The online version contains supplementary material available at:

Additional file 1 (PDF): Supplementary Fig. S1: Workflow of the proposed approach. First,

we detected the pollutant-induced DARs using ATAC-Seq datasets obtained from the

TaRGET database (*in vivo* pollutant exposure). We then performed ChIPEA with pollutant-

induced DARs to predict TFs that play a pivotal role in the pharmacological effects of

pollutants on health. The prediction phase was completed upon construction of pollutant–

TF–disorder triadic associations integrating TF-disease associations retrieved from DisGeNET

with the pollutant–TF matrices obtained as outcomes of ChIPEA. Subsequently, in the

validation phase, we checked the authenticity of the predicted pollutant–disorder

associations (shown as blue-colored continuous matrix) by referencing known chemical–

disorder associations from CTD (shown as green-colored boolean matrix) and summarized

the results as AUROC and AUPR scores. The *in vitro* chemical perturbation dataset obtained

from the DBKERO database was also used to evaluate the prediction accuracy of the

proposed method. The performance of other methods that utilize chemically induced DEGs

as input or employ motif-based enrichment analysis (MotifEA) rather than ChIPEA were

simultaneously assessed using the same workflow. **Supplementary Fig. S2**: Summary of the

number of DEGs or DARs used as input to the enrichment analyses. Ranking plots are

illustrated for individual chemical dosing or pollutant exposure conditions, and are sorted by

the summation of the number of up- and down-regulated DEGs or opened and closed DARs.

Supplementary Fig. S3: Results of the Bisulfite-Seq enrichment analysis for three

representative pollutants. Dots indicate individual Bisulfite-Seq peak sets from ChIP-Atlas.

Peak sets obtained from Bisulfite-Seq on blood (in the case of TBT and PM2.5 exposure) and

liver (in the case of lead exposure) samples are colored (hyper-methylated regions, red;

hypo-methylated, blue). Positive and negative values of log2 fold enrichment (opened

DARs/closed DARs; X-axis) indicate peak sets enriched in opened and closed chromatin

induced by pollutant exposure, respectively. The enrichment score ($-\log_{10}Q$-value; Y-axis)

indicates the statistical significance of Bisulfite-Seq enrichment analysis. **Supplementary Fig.**

S4: Expression levels of TFs mentioned in Figure 4 in mouse tissues before and after

exposure to TBT, PM2.5, and lead. Genes encoding the top 20 enriched TFs in Figure 4 (right

panels) are colored (rank 1–10, orange; rank 11–20, blue). CPM, count per million.

Additional file 2 (XLSX): Supplementary Table S1: *In vitro* chemical perturbation RNA-Seq

dataset (DBKERO). **Supplementary Table S2:** *In vitro* chemical perturbation ATAC-Seq

dataset (DBKERO). **Supplementary Table S3:** *In vivo* pollutant exposure RNA-Seq dataset

(TaRGET). **Supplementary Table S4:** *In vivo* pollutant exposure ATAC-Seq dataset (TaRGET).

Data shown in bold font are those that yielded high AUROC scores, as reported in the main

text; see also Supplementary Table S6. **Supplementary Table S5:** AUROC and AUPR scores

for *in vitro* chemical perturbation datasets (DBKERO). **Supplementary Table S6:** AUROC and

AUPR scores for *in vivo* pollutant exposure datasets (TaRGET). Data shown in bold font are

those that yielded high AUROC scores, as reported in the main text; see also Supplementary

Table S4. **Supplementary Table S7:** Result of DAR-ChIPEA for exposing blood samples to

TBT. **Supplementary Table S8:** Result of DAR-ChIPEA for exposing blood samples to PM$_{2.5}$.

Supplementary Table S9: Result of DAR-ChIPEA for exposing liver samples to lead.

FIGURE LEGENDS

Fig. 1 Overview of the proposed DAR-ChIPEA approach. **a** Schematic diagram illustrating the

difference between our previously reported DEG-ChIPEA method [22] and the proposed

DAR-ChIPEA approach. The DEG-ChIPEA method is restricted to analyzing TF binding to the

vicinity of gene loci. In contrast, the DAR-ChIPEA approach comprehensively covers TF

binding to accessible chromatin regions (obtained from ATAC-seq) across the genome

including proximal (e.g., promoters) and distal (e.g., enhancers) regulatory elements. The

circles surrounded by dotted lines represent the TFs binding to distal regulatory elements

that are not detected in the DEG-ChIPEA method. **b** To demonstrate the feasibility of DAR-

ChIPEA, we identified the TFs enriched at pollutant-induced DARs by analyzing large-scale

ChIP-Seq data. Overlaps were evaluated between the DARs (green box) and peak-call data

(black lines) of TF-related ChIP-Seq experiments archived in ChIP-Atlas

Fig. 2 Validation of the proposed DAR-ChIPEA approach to predict TF-driven MoAs of *in*

vitro chemical perturbation. **a** Schematic diagram for TF enrichment analyses using *in vitro*

chemical perturbation datasets. DEGs or DARs upon chemical perturbation were detected

using data from RNA-Seq or ATAC-Seq, respectively, followed by enrichment analyses to

profile TF binding. Methods used for TF enrichment analysis are shown in blue; the proposed

method (DAR-ChIPEA) is highlighted in bold. **b** Ranking plot of maximum enrichment scores

($-\log_{10}Q$-value) for each dosing condition obtained using the DAR-ChIPEA and DEG-ChIPEA

methods. c and d Distribution of AUROC (**c**) and AUPR (**d**) scores for each chemical–TF–

disorder triadic association predicted using DEG-MotifEA, DAR-MotifEA, DEG-ChIPEA, and

ChIPEA. Orange dots represent mean scores; global scores are noted below the plots.

Differences between the methods are presented as Q-values (two-tailed Wilcoxon rank-sum

test) above the bee swarm plots

Fig. 3 Application of the DAR-ChIPEA approach to predict TF-driven modalities of pollutant

administration *in vivo*. **a** Schematic diagram for TF enrichment analyses using *in vivo*

pollutant exposure datasets. DEGs or DARs upon pollutant exposure were detected using

RNA-Seq or ATAC-Seq, respectively, followed by enrichment analyses to profile TF binding.

Methods used for TF enrichment analysis are shown in blue, with the proposed DAR-ChIPEA

method highlighted in bold. **b** Ranking plot of maximum enrichment scores ($-\log_{10}Q$-value)

for each exposure condition as obtained using the DAR-ChIPEA and DEG-ChIPEA methods. c

and d Distribution of AUROC (**c**) and AUPR (**d**) scores for each pollutant–TF–disorder triadic

association predicted using DEG-MotifEA, DAR-MotifEA, DEG-ChIPEA, and ChIPEA. Orange

dots represent mean scores; global scores are noted beneath the bee swarm plots.

Differences between the methods are presented as Q-values (two-tailed Wilcoxon rank-sum

test) above the bee swarm plots

Fig. 4 Biological interpretations of the predicted pollutant–TF–disorder associations for three

representative pollutants. Mouse phenotype ontologies enriched for the genes proximal to

pollutant-induced opened DARs are shown in the left panels. Bars representing individual

ontology terms are colored based on their parent categories, which were obtained from the

Mouse Genome Informatics database. Volcano plots illustrating the results of the proposed

DAR-ChIPEA approach are shown in the right panels. Dots indicate individual TFs and are

colored based on cell type classes obtained from ChIP-Atlas. Positive and negative values of

$\log_2$ fold enrichment (opened DARs/closed DARs; X-axis) indicate TFs enriched in relaxed and

condensed chromatin induced by pollutant exposure, respectively. The enrichment score ($-$

$\log_{10}Q$-value; Y-axis) indicates the statistical significance of ChIPEA.

a

DEG-ChIPEA

Enhancer · Promoter · Gene

DAR-ChIPEA

Enhancer · Promoter · Gene

b

Pollutants

TFs

Opened chromatin

Disorders

TF ChIP-Seq experiments

	TF Binding
TF1	+
TF2	−
TF3	+
TF4	+
TF5	−
TF6	+

Enhancer   Promoter   Gene

▬ Peaks called using MACS2

Figure 1

Figure 2

Figure 3

Figure 4