Disrupted memory T cell expansion in HIV-exposed uninfected infants is preceded by premature skewing of T cell receptor clonality.

Sonwabile Dzanibe[1], Aaron J. Wilk[2], Susan Canny[2,3], Thanmayi Ranganath[2], Berenice Alinde[1,4], Florian Rubelt[5,6], Huang Huang[5,6], Mark M. Davis[5,6,7], Susan Holmes[8], Heather B. Jaspan[1,9**], Catherine A. Blish[2,6,10**] and Clive M. Gray[1,4**]

Affiliations

1. Division of Immunology, Department of Pathology, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa.
2. Department of Medicine, School of Medicine, Stanford University, Stanford, CA.
3. Division of Rheumatology, Department of Pediatrics, Seattle Children's Hospital, Seattle, WA USA
4. Division of Molecular Biology and Human Genetics, Stellenbosch University, Cape Town, South Africa.
5. Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA, USA.
6. Institute for Immunity, Transplantation and Infection, Stanford University School of Medicine, Stanford, CA, USA.
7. Howard Hughes Medical Institute, School of Medicine, Stanford University, Stanford, CA.
8. Department of Statistics, Stanford University, Stanford, CA, USA
9. Seattle Children's Research Institute and Department of Paediatrics and Global Health, University of Washington, Seattle, WA.
10. Chan Zuckerberg Biohub, San Francisco, CA

**Corresponding authors: Clive M. Gray cgray@sun.ac.za; Catherine A. Blish cblish@stanford.edu; Heather B. Jaspan hbjaspan@gmail.com

Keywords: HIV exposure; infant immunity; T cell receptor, NK cels and antibody responses

Word count: Abstract: 151; Main text: 4993; Methods: 2039

Abstract

30
While preventing vertical HIV transmission has been very successful, the increasing

number of HIV-exposed uninfected infants (iHEU) experience an elevated risk to

infections compared to HIV-unexposed and uninfected infants (iHUU). Immune

developmental differences between iHEU and iHUU remains poorly understood and

here we present a longitudinal multimodal analysis of infant immune ontogeny that

highlights the impact of HIV/ARV exposure. Using mass cytometry, we show

alterations and differences in the emergence of NK cell populations and T cell memory

differentiation between iHEU and iHUU. Specific NK cells observed at birth were also

predictive of acellular pertussis and rotavirus vaccine-induced IgG and IgA responses,

respectively, at 3 and 9 months of life. T cell receptor Vβ clonotypic diversity was

significantly and persistently lower in iHEU preceding the expansion of T cell memory.

Our findings show that HIV/ARV exposure disrupts innate and adaptive immunity from

birth which may underlie relative vulnerability to infections.

Introduction.

Early life, especially in Africa, is often plagued by high infectious morbidity with infectious diseases accounting for 2.61 million deaths of which 45% occur duing the neonatal period[1]. This high mortality rate is most likely linked to the period during which the immune system adapts to extrauterine life. During this transition window, several factors such as maternal morbidity, microbial exposure, and vaccination significantly modulate infant immunity and consequently influences health and disease outcomes[2–5]. Investigating immune ontogeny and factors that impact immune trajectory during infancy can provide important insights into understanding how better to combat these early life immune stressors.

Current dogma is that the *in utero* environment is sterile and that newborn infants have limited antigen exposure prior to birth, with T cells being predominantly naïve with little T cell receptor (TCR) engagement[6,7]. This lack of pre-existing adaptive cellular memory early in life is likely to increase the vulnerability of infants to infectious agents and disease, especially if there is an absence of adequate breast-feeding to provide passive maternal antibody immunity[3,8,9]. Since the diversity and composition of the TCR is presumably dependent on prior antigen exposure, examining changes in the TCR repertoire during infancy could provide insight into immune development and how specific immune modulatory factors influence susceptibility to infectious diseases.

The advent of the Option B+ vertical transmission prevention program with effective antiretroviral (ARV) drugs has ensured that HIV transmission has been minimised, but also introduces the possible adverse effect of HIV and ARV exposure on neonatal development *in utero* and in the post-natal period. The intertwined nature of HIV and

ARV exposure makes it difficult to unravel, but it is known that HIV/ARV-exposed

uninfected infants (iHEU) have higher risk of infectious disease-related morbidity and

mortality compared to HIV-unexposed uninfected infants (iHUU) of a similar age,

suggesting disruptions to their immune maturation[10–15]. Many studies investigating

immunological disparities between iHEU and iHUU are limited to cross-sectional

analysis with few studies investigating longitudinal changes and thus there is a lack of

evidence on how T cells mature early in life in the context of HIV/ARV exposure. We

have previously shown that maternal HIV/ARV exposure alters the dynamics of the T

regulatory (Treg) to Th17 cell ratio resulting in a Th17/Treg imbalance associated with

gut damage[16]. In this paper, we extend this analysis to investigate HIV/ARV exposure

on T cell clonality, memory and NK cell maturation differences between iHEU and

iHUU.

Since early life is marked by inexperienced adaptive immunity, neonatal immune

defence is heavily reliant on innate immune cells such as NK cells to rapidly eliminate

infections. Compared to adults, however, neonatal NK cells display functional defects

such as reduced cytolytic activity including antibody mediated cell cytotoxicity[17,18],

decreased expression of adhesion molecules[19] and lower secretion of TNF-α and INF-

γ[20]. Earlier studies have further demonstrated that compared to iHUU, iHEU have a

lower proportion of NK cells measured at birth and 6 months, and having reduced IFN-

γ secretion and perforin expression[21,22]. In addition, the function of NK cells is

influenced by the combinatorial signalling through a diverse array of activating and

inhibitory receptors expressed on the cell surface[23]. The composition of these

receptors is driven by past viral exposure[20,24,25]. Given that NK cells also play a key

role in priming the adaptive immune system to respond to invading pathogens or to

tailor their immunogenicity towards specific vaccines[26], it is important to investigate

the maturation trajectory of NK cells and how this could shape adaptive immunity.

Whether early life exposure to HIV/ARV shapes the maturation and/or diversification

of NK cells remains unknown.

Here, we comprehensively investigated the relationship between adaptive and innate

immunophenotypes along with TCR diversity and the ability to mount an antibody

response to pertussis and rotavirus vaccination. Given the heightened infectious

morbidity risk in iHEU, we hypothesised that immune ontogeny is associated with a

narrowing TCR repertoire over time in parallel with expanded NK cell subsets in iHEU

relative to iHUU. We mapped the immune trajectory of NK and T cell phenotypic

clusters in the first 9 months of life and associated this with TCR diversity and antibody

titres. Using this approach, we show divergent adaptive immune maturation in iHEU

relative to iHUU occurring from 3 months of life with earlier phenotypic differences

evident in NK cells. Our data show a narrowing and persistent skewing of the TCR

repertoire from birth in iHEU that precedes altered CD4$^+$ and CD8$^+$ T cell memory

development.

Results

Immune cell transition from birth to 9 months of age.

We first analysed the immune ontogeny of innate and adaptive immune cell subsets and marker expression profiles longitudinally during the first 9 months in our iHUU and iHEU samples. Infant samples were analysed longitudinally using a mass cytometry antibody panel (Table S1). This included matched PBMC at birth (n=44) and at week 4 (n=53), 15 (n=52) and 36 (n=53), as well as in some cord blood mononuclear cells (n=5). Unsupervised cell clustering on all samples using the live singlet cell population (Figure S1A) revealed 11 cell clusters, shown in Figure 1A, consisting of B cells (cluster 1), T cells (CD4$^+$ T cluster 2 and CD8$^+$ T clusters 3, 4, and 9), monocytes (cluster 8), NK cells (cluster 10), and NKT-like cell (cluster 11). The three CD8$^+$ T cell clusters (3, 4 and 9) were delineated as naïve (13.9%, CD45RA$^+$CD27$^+$CCR7$^+$), a small effector memory population (0.07%, EM: CD45RA$^+$CD27$^+$CCR7$^-$) and effector cells (4.4%, Eff: CD45RA$^+$CD27$^-$CCR7$^-$). We also identified a lineage negative cluster (3.2%) and one cluster expressing HLA-DR only (1.2%) that appeared most closely related to the monocyte cluster (Figure 1A & S1B). Although we employed computational doublet removal from our clustering analysis, a minor miscellaneous population (0.65%) co-expressing typically mutually exclusive lineage markers was identified (Figure 1A & S1B). This small population is shown as a central cluster 7 in the dimensional reduction of the immune cell clusters (Figure 1A) and was excluded from downstream analysis.

When we examined these clusters over time, we observed an age dependent segregation of cells. Proportions of clusters changed over 36 weeks of life, where monocytes and CD4+ T cells diminished with age and B cells and effector memory

CD8+ T cells increased (Figure 1B). Monocyte (ρ=-0.4, p <0.001) and CD4$^+$ T cell (ρ=-

0.36, p <0.001) clusters were negatively correlated with infant age, whereas the EM

CD8$^+$ T cell and B cell clusters were positively correlated (ρ=0.66 and ρ=0.54

respectively, p value <0.001; Figure 1C). We further manually gated on CD4$^+$ and

CD8$^+$ T cells and targeted the NK cell population using lineage-exclusion manual

gating (Figure S1A). Using the median marker expression for each sample,

multidimensional scaling (MDS) revealed intra-individual variability associated with

infant age (Figure 1D). For both CD4$^+$ T cells and NK cells, the grouping of the samples

by marker expression displayed a converging trajectory along the MDS2-axis (Figure

1D). Both of these cell populations had a linear converging trajectory, although

stabilizing from week 15 for NK cells (Figure 1E). CD8$^+$ T cells, however, revealed a

divergent trajectory along the MDS1-axis (Figure 1D), with a significant expansion by

week 15 (Figure 1E).

Overall, these detailed unsupervised phenotypic changes in cell clustering illustrate

the archetypal progression and transition of infant immunity from neonatal to an adult-

like immune phenotype by 9 months of age, being characterised by a shift from innate

myeloid cells to an increase in lymphoid adaptive immune cells.

CD4 and CD8 T cell memory maturation is disrupted by HIV exposure after

three months of life.

To identify in more depth how HIV/ARV exposure disrupts T cell phenotypes, we

parsed manually gated CD4 and CD8 cells from all samples and time points using

unsupervised cell clustering which generated 13 CD4$^+$ T cell clusters and 7 CD8$^+$ T

cell clusters (Figures 2A and 2D). Tables 1 & 2 ascribe the phenotype for each of the

CD4$^+$ and CD8$^+$ T cell clusters, respectively. The marker expression patterns and

relationships between the cell clusters are shown using unsupervised hierarchical

clustering heatmaps in Figure S2A.

Across all time points, 13 clusters of CD4+ T cells existed (Figure 2A and Table 1)

where naïve cells (cluster 2, CD45RA$^+$CD27$^+$CCR7$^+$) made up the predominant

population with an overall frequency of 84.1% (Figure 2A & Table 1). The remaining

12 clusters for CD4+ T cells consisted of minor populations with the top three being

activated naïve-like cells (cluster 4, CD38$^+$HLA-DR$^+$CD45RA$^+$CD27$^+$CCR7$^+$), PD-1

expressing activated cells (cluster 12, PD-1+CD38+) and cytotoxic terminally

differentiated cells (cluster 3, CD45RA$^+$CD27$^-$CCR7$^-$CD57$^+$Per$^+$PD-1$^+$). When these

cell clusters were examined over infant age, there was a proportional increase in

cluster 1 (Th2-like, CCR4$^+$CCR6$^-$CXCR3$^-$), decrease in cluster 4 (activated cells,

CD38$^+$HLA-DR$^+$CD45RA$^+$CD27$^+$CCR7$^+$) and maintenance of cluster 2 (naïve,

CD45RA$^+$CD27$^+$CCR7$^+$) (Figure 2B). We also assessed these compositional

differences by performing Principal Component Analysis (PCA) on centred log-ratios

of the relative cluster abundance for each sample. Consistent with the observed

immunological trajectory shown in Figures 1D, phenotypic compositions were strongly

influenced by infant age, where early time points (cord, birth, and week 4) were

grouped together and distinguishable from week 15 and 36 time points (Figure 2C).

Looking at CD8$^+$ T cells in the same manner, we identified 7 cell clusters, being less

diverse than CD4$^+$ T cells, but a more distinct clustering between naïve and

effector/terminally differentiated memory (Figure 2D and Table 2). It was also clear

that naïve CD8$^+$ T cells were less populous and that clusters 1 (terminally

differentiated, Perforin$^+$CD45RA$^-$CD27$^-$CCR7$^-$CD57$^+$), 5 (cytotoxic cells,

Perforin$^+$CD57$^+$) and 6 (cytotoxic effector memory, Perforin$^+$CD45RA$^-$CD27$^-$CCR7$^-$

made up a higher proportion of cells than in CD4$^+$ T cell compartment. When

separating out these clusters over time, these perforin-expressing cell clusters (1, 5

and 6) increased over 9 months of life (Figure 2E), with a contraction of the naïve cell

population. Like CD4$^+$ T cells, there was an age dependent CD8$^+$ T cell compositional

change (Figure 2F).

We next determined whether there were any disruptions in T cell cluster compositions

due to HIV exposure. There were statistically significant differences in PCA-2 at weeks

15 and 36 for CD4$^+$ T cells between iHEU and iHUU (Figure 2G) and a converse

difference at week 36 for CD8$^+$ T cells (Figure 2H). To account for these divergent T

cell phenotypes, we used differential abundance testing to determine which of the cell

clusters were responsible for these differences. CD4 clusters 4, 5, 6, 7, 9 and 11 were

significantly lower in iHEU compared to iHUU at week 15, with clusters 5, 7 and 9

remaining lower in iHEU at week 36 (Figure 2I). These less frequent of cell clusters in

iHEU were characterised as either being activated (cluster 4: CD38$^+$HLA-DR$^+$),

expressing perforin (clusters 7, 9 and 11) or exhibiting terminal differentiation

(CD57$^+$/PD-1$^+$; clusters 5 and 6). In contrast, the CD4$^+$ Th2-like cluster 1 was

significantly higher in iHEU compared to iHUU at week 15 but was no longer

significantly elevated by week 36 (Figure 2I). Moreover, CD4$^+$ Th2 cells (cluster 1) and

cytotoxic terminally differentiated cells (cluster 3: CD45RA$^+$CD27$^-$CCR7$^-$

CD57$^+$Perforin$^+$PD-1$^+$) were also higher in iHEU compared to iHUU at week 4 (p=0.007

and p=0.013, respectively) although these differences were not statistically significant

following FDR correction (Figure S2B).

For CD8$^+$ T cells, there were significantly lower mean frequencies for clusters 2 (NKT-

like cells: CD56$^+$CD16$^+$NKp30$^+$NKp46$^+$NKG2A$^+$Per$^+$), 3 (Naïve-like activated cell:

CD45RA$^+$CD27$^+$CCR7$^+$CXCR3$^+$CCR4$^+$HLA-DR$^+$) and 7 (proliferating EM: Ki67$^+$

CD45RA$^+$CD27$^+$CCR7$^-$Per$^+$) in iHEU compared to iHUU at week 15 (Figure 2I). By

week 36, only the mean frequencies of clusters 2 (NKT-like cells) and 3 (Naïve-like

cells) remained significantly lower in iHEU compared to iHUU (Figure 2J).

A similar analytical approach was undertaken for NK cells, and of the 10 clusters

identified (Figure 3A), 9 were CD56$^{dim/+}$CD16$^{dim/+}$ (Table 3). This included

CD56^{Hi}CD16$^-$ (cluster 9), which had high NKG2A expression, six clusters that were

CD56$^{dim/+}$CD16$^{dim/+}$ (clusters 1, 4, 5, 6, 8 and 10) that were predominantly expressing

perforin, and 2 clusters that were CD56$^-$CD16$^+$ (clusters 3 and 7). Only cluster 2 was

characterised as CD56$^-$CD16$^-$ and expressed DNAM1 (Figure S2A & Table S6). The

NK cell phenotypic composition was also observed to be strongly associated with

infant age (Figure 3B and C), with cluster 1 (Per$^+$CD45RA$^+$Fc$_ε$RIIγ$^+$) expanding over

36 weeks and cluster 2 (CD56$^-$CD16$^-$DNAM1$^+$) contracting (Figure 3B). Of note, the

perforin-expressing cluster 5 was expanded at weeks 15 and 36. Compositional

differences in NK cell clusters between iHEU and iHUU were observed at week 4,

earlier than for T cells, and while this trend persisted at week 15, the differences were

not statistically significant (Figure 3D). Differential abundance testing revealed that at

week 4, iHEU had higher frequencies of cluster 1 (CD56^{dim}CD16^{dim}) co-expressing

Per$^+$CD45RA$^+$FcERIy+ (p=0.04, p.adj=0.2) and cluster 5 (CD56^{dim}CD16$^+$) co-

expressing Per$^+$CD57$^+$CD45RA$^+$CD38$^+$ (p=0.03, p.adj=0.2) compared to iHEU (Figure

S2C). This trend continued at week 15 for cluster 5 (p=0.041, p.adj=0.20). Also at

week 15, we observed lower frequencies of cluster 10 (CD56$^+$CD16$^+$: p=0.027,

p.adj=0.2) in iHEU compared to iHUU (Figure S2C). These findings show that that the

ontogeny of immune cells is ordered, and that HIV/ARV exposure sequentially disrupts

immune trajectory over time after birth beginning with that of innate (NK) cells and later

disrupting memory T cell maturation.

Early TCR repertoire skewing in iHEU.

To relate altered T cell memory maturation trajectories to T cell repertoire changes,

we characterised TCR clonotypes in fractionated naïve and memory T cells from

matching samples used to define memory lineage in the CyTOF panel. Each PBMC

sample was sorted into four T cell subsets: naïve (CD45RA$^+$CD27$^+$CCR7$^+$) CD4$^+$ and

CD8$^+$ cells and total memory (CD45RA$^{-/+}$CCR7$^{-/+}$) T cells (Figure S3A). We subjected

these sorted cell fractions (n=934 samples) to bulk sequencing of the TCRβ locus,

which enabled identification of TCR clones in 885 samples. The number of unique

clones identified per T cell subset positively correlated with the total number of cells

sequenced (Figure S3B). There were 24 samples that had either fewer than 10 clones

or the number of unique clones were greater than the number of cells sequenced, and

these were removed from downstream analysis. Therefore, we included 861 samples

having a total of 238,092 reads (range: 11-5451) and an overall median number of

unique clones of 81 (IQR: 42-156). The CDR3 lengths of the clonotypes were evenly

distributed across infant age and between iHEU and iHUU (Figure S3C).

The diversity and richness of the TCR clonotypes, determined by inverse Simpson

diversity index and Chao1 richness scores respectively, remained unchanged with

infant age (Figure S3D & S3E). However, when the TCR diversity scores were

stratified by infant HIV exposure status, iHEU TCR clonotypes had relatively lower

diversity compared to iHUU starting as early as birth (Figure 4A). These differences

were apparent for naïve CD4$^+$ T cells at weeks 15 and 36 and for naïve CD8$^+$ T cells

at birth and week 36. Differences in memory T cell diversity scores were observed to

occur only at birth, being significantly lower in iHEU for both CD4$^+$ and CD8$^+$ T cells

(Figure 4A). Similarly, richness (the number of unique TCR clones) was significantly

lower in iHEU relative to iHUU for both CD4$^+$ and CD8$^+$ T cells at birth (Figure 4B).

This remained statistically significant for naïve CD4$^+$ T cells at week 36 (Figure 4B).

When examining memory T cells, significant differences in TCR richness were

restricted to earlier timepoints, being lower in iHEU compared to iHUU at birth and

week 4 for CD4$^+$ T cells and only at birth for CD8$^+$ T cells. We also observed differences

in the overall structural overlap, as measured by Jaccard indices, of the memory CD4$^+$

TCR repertoire between iHEU and iHUU at birth, although this difference was less

prominent at later time points (Figure S3F). In contrast, the TCR structural overlap for

naïve CD8$^+$ T cells was lower in iHEU compared to iHUU only at week 36 (Figure

S3F). Differences in Vβ gene usage were observed between iHEU and iHUU with

most genes being used more frequently in iHEU (Figure S4). Together, these results

show expansion of specific TCR clones among iHEU, resulting in skewing of the TCR

repertoire relative to iHUU.

We next wished to determine if there was a relationship between TCR diversity and

the clusters of CD4$^+$ and CD8$^+$ T cells identified in Figures 2A-F. By using Spearman's

rank correlation between T cell clusters and the inverse Simpson TCR diversity scores

for naïve and memory CD4$^+$ and CD8$^+$ T cells, we found that the frequencies of CD4$^+$

T   cell   clusters   (4-6:   CD38+HLA-DR$^+$CD45RA$^+$CD27$^+$CCR7$^+$,

CCR4$^+$CCR6$^+$CXCR3$^+$CD57$^-$,   CCR4$^+$CCR6$^+$CXCR3$^+$CD57$^+$,   and   9-11:

Perforin$^+$Ki67$^+$CD25$^+$, Perforin$^+$Ki67$^-$HLA$^-$DR$^+$CD69$^+$CD39$^+$, Per$^+$Ki67$^+$) from iHEU

were positively correlated with the TCR diversity scores derived from sorted naïve

CD4+ T cells (Figure 4C). Similarly, frequencies of CD8+ T cells clusters (2, 3 and 7:

CD56+CD16+NKp46+NKG2A+, HLA-DR+, Ki67+CD45RA+CD27+CCR7-) were

positively correlated to the TCR diversity scores derived from the sorted naïve CD8+

T cells at either birth or week 4 (Figure 4D). No T cell clusters were significantly

correlated to TCR diversity derived from sorted memory T cells (data not shown).

Collectively, our findings show skewing of the TCR repertoire in iHEU relative to iHUU

from birth, in the naïve T cell compartment prior to phenotypically defined memory

maturation of CD4 and CD8 T cells.

Predicted TCR recognition of epitopes found only in iHUU.

To understand the predicted antigen specificities of the TCR clonotype differences

between iHEU and iHUU, we used GLIPH (grouping of lymphocyte interactions by

paratope hotspots)[27,28]. TCR specificities from sorted naïve CD4+ T cells in iHUU

appeared enriched with specificities for cytomegalovirus (CMV), Hepatitis C virus

(HCV), SARS-Cov2 and HIV-1 (Figure 4E). Enrichment of Influenzae A specific clones

was more apparent in memory CD4+ T cells at the week 36. For memory CD4+ T cells

at week 4, the shared specificity groups in iHUU were enriched for CMV clones.

Similarly, for CD8+ T cells, there were more specificities in the naïve compartment at

both earlier and later time points with less predictions in memory CD8+ T cells. TCR

specificities in sorted naïve CD4+ and CD8+ cells from iHUU appeared consistent

throughout 36 weeks, whereas specificities in the CD4 memory compartment

fluctuated and appeared predominantly at 4 and at 36 weeks (Figure 4E). Conversely

for iHEU sorted T cell populations, no enriched TCR specificities in either naïve or

memory CD4+ or CD8+ cells at any of the time points were predicted.

NK cell clusters are the strongest predictors of antibody responses.

To relate how early life T and NK cell clusters may associate with vaccine-induced antibody responses, we measured the response to acellular pertussis vaccination from birth to 36 weeks and to rotavirus at 36 weeks. We initially hypothesised that iHEU would have lower IgG and IgA responses compared with iHUU after vaccination with routine administration of these vaccines, which may explain a mechanism for higher infection rates of these pathogens in iHEU. To test this, IgG antibody levels against pertussis were measured at birth, weeks 4, 15 and 36, with rotavirus specific IgA and neutralization titres measured at week 36 (Figure 5B and C). The median anti-pertussis IgG OD values were lower in iHEU compared to iHUU at birth and week 4 (pre-vaccination at 6 weeks; Figure 5A). However, at week 15 (1 week post 3rd vaccination), iHEU showed a significantly higher IgG response relative to iHUU, persisting until week 36 (Figure 5A). No significant differences were observed in anti-rotaviral IgA levels or ability to neutralize rotavirus between iHEU and iHUU (Figure 5B, C).

We built a prediction model by grouping all infants (iHEU and iHUU) into those that had a protective pertussis-specific IgG titre (≥0.469 OD_{Abs}) denoted as responders and those that were non-responders (<0.469 OD_{Abs}). Most of the infants at birth and week 4 (89.7% and 94.1%, respectively) were determined to be non-responders as expected, since this was prior to vaccination. At week 15 and 36 of age, 46.4% and 34.0% infants respectively were determined to be responders (Figure 5A). Multivariate unbiased variable selection using MUVR[29] was performed to determine early life cell clusters that could distinguish later pertussis vaccine responders from non-responders, according to our classification. Certain cell cluster abundances at birth

and week 4 selected using MUVR were associated with low misclassification scores

of a pertussis vaccine response at weeks 15 and 36 (Figure S4). We then used these

abundances of the MUVR selected clusters to perform a partial least squares

discriminant analysis (PLS-DA) with the pertussis IgG responses at week 15 and 36

as response variables (Figures 5D and E, respectively). The latent variables (LV) of

the cell clusters as determined by PLS-DA were then used to compute area under the

curve (AUC) of the receiver operating curve (ROC) distinguishing pertussis

responders from non-responders. This approach revealed that NK cell clusters (1, 3,

4, 6, 8 and 10) at birth and at week 4 (clusters 1, 2, 3 and 9, Table 3) were ranked the

highest in the list of predictors of the pertussis IgG vaccine response at week 15

(Figure 5D, S4A & S4B, at birth: AUC = 0.93 and p=0.005; at week 4: AUC = 0.85 and

p=0.012). The cell clusters derived from CD4$^+$ or CD8$^+$ T cells were not significantly

predictive of an IgG pertussis vaccine response at weeks 15 and 36 (Figure 5D, S4B-

D, for CD4 clusters at birth: AUC=0.73, p=0.063; for CD8 clusters: AUC=0.69, p=0.12;

at week 4: AUC=0.70, p=0.06; for CD8 clusters: AUC=0.70, p=0.053).

Using the same approach for rotavirus responses, we used an arbitrary threshold of

overall median concentrations for IgA responses to dichotomize infants into those that

had concentrations below the median (low-responders) and those above the median

(high-responders; Figure 5B). In iHUU, 66.7% were high rotavirus vaccine responders

compared to 45.5% iHEU (p=0.11). We subsequently determined which immune cell

clusters at birth and week 4 were predictive of IgA responses measured at week 36

using MUVR (Figure 5F). PLS-DA analysis using the MUVR selected cell clusters

revealed that NK cells at birth were highly predictive of an IgA rotavirus response at

week 36 (Figure 5F, AUC = 0.92, p = 0.0007). Neither CD4$^+$ or CD8$^+$ T cell clusters

were predictive of a rotavirus response at week 36 (Figure 5F, for CD4 clusters at

birth: AUC=0.7, p=0.05; for CD8 clusters: AUC=0.55, p=0.59; at week 4: AUC=0.68,

p=0.042; for CD8 clusters: AUC=0.6, p=0.24).

We also tested whether contemporaneous cell clusters might be related to pertussis

responses. When we analysed the frequencies of the cell clusters at weeks 15 and 36

with pertussis antibody levels at weeks 15 and 36, CD4$^+$ T cell cluster 1 (Th2 cells)

was significantly correlated to pertussis antibody titres at week 15, while at week 36

CD4$^+$ T cell clusters 6, 7, 9-11, and 13 (Table 1) were inversely correlated to pertussis

antibody titres (Figure 5G). Only CD8$^+$ T cell cluster 7 (Proliferating EM:

Ki67$^+$CD45RA$^+$CD27$^+$CCR7$^-$) was observed to be inversely correlated to pertussis

antibody titres measured at week 36 (Figure 5G). Of note, there was no significant

contemporaneous correlation between the NK cell clusters and pertussis antibody

titres (Figure 5G).

Collectively, these data show that various NK cell phenotypes at birth and in the first

1 month of life can predict vaccine-induced antibody responses to acellular pertussis

and rotavirus at 9 months, regardless of infants being exposed to HIV.

Discussion

Herein, we describe immune maturation in the first 9 months of life and how HIV/ARV exposure alters this ontological trajectory. Our findings show ordered immune changes during infancy, typical of the transition from innate-like and naïve neonatal immunity towards memory differentiated adaptive infant immunity[30]. NK cell phenotype and cluster composition differed between iHEU and iHUU at birth and week 4 and was predictive of vaccine responses later in life, whereas T cell memory maturation was impacted after 15 weeks. There was a skewing of the TCR repertoire in naïve and memory cells in iHEU from birth, relative to iHUU, and the persistent lack of TCR diversity that could account for the increased vulnerability to infections reported among iHEU[10].

Although differentiated T cells are detected as early as 12-14 weeks in gestation[31,32], neonatal immunity is predominantly naïve and characterised by the abundance of CD45RA$^+$ T cells[7]. T cell memory expansion during infancy is therefore essential in protecting against pathogens[33]. The differences in the kinetics of CD4$^+$ and CD8$^+$ T cell memory maturation in our study could not be explained by changes in TCR clonotypes since for both T cell subsets, TCR diversity and richness did not vary over time for either iHEU or iHUU. We presume that the steady rate of CD4$^+$ T cell maturation is partially explained by the abundance of CD4$^+$ T cells early in life, responsible for regulating immunological responses and thus facilitating adaptation to an extrauterine life laden with microorganisms[6,34,35].

We show that the T cell compartment was significantly impacted by HIV/ARV exposure. The overall trajectory of T cell phenotypic composition in iHEU diverged from that of iHUU, particularly after week 15 of life. CD4$^+$ and CD8$^+$ T cell

compartments in iHEU had lower frequencies of activated cells and a minor cell

population expressing either perforin or PD-1 compared to iHUU. These findings

contradict earlier reports showing iHEU having heightened immune activation and/or

exhaustion[36,37], which may be related to mixedfeeding or non-breastmilk feeding that

could introduce water contaminants or alter infant microbiome[38]. Mothers in our study

were encouraged to exclusively breastfeed, likely accounting for lower levels of

activated T cells in iHEU compared to iHUU[39]. Nonetheless, reduced or impaired

memory T cells among iHEU compared to iHUU have been previously demonstrated

and associated with poor childhood vaccine responses and/or increased

hospitalization[16,39–41].

The 15 week period before significant expansion of T cell memory differentiation was

preceded by significantly reduced diversity and richness in memory T cell clonality

from birth in iHEU relative to iHUU. This was in agreement with others who reported

lower βTCR clonality in the cord blood of iHEU[42] and such reduced TCR diversity

found at birth infers that T cells have undergone clonal expansion *in utero*. Using

GLIPH[27,28], we could not detect any predicted antigen specificity enriched among

iHEU, in contrast to earlier studies showing high frequency of HIV-1 specific clones[42]

and reactivity of T cells towards HIV proteins in iHEU[43]. What was evident instead was

that naïve CD4$^+$ and CD8$^+$ TCR reactivities in iHUU were targeting CMV, EBV and

HCV antigens. There may be three possible reasons for naïve cells showing clonal

expansion and increased diversity in iHEU. Firstly, that our isolation of

CD45RA+CCR7+CD27+ naïve population was contaminated with some memory cells

during the sorting procedure. The contribution of sorting contamination would have

negligible outcomes as we obtained >95% purity after sorting and any inclusion of

memory cells would be of extremely low proportion. Additionally, the frequency of

naïve T cells was significantly higher compared to that of memory T cells throughout

infancy which could contribute to sampling bias resulting in enriched TCR clones

amongst naïve T cells compared to the memory fraction. Second, that these cell

populations are very early differentiated memory, and in the absence of CD95, we

could not account for persistence of CD4 or CD8 memory stem cell emergence at

birth[44–46]. The strong association between activated "naïve" CD4 and CD8 cells with

TCR diversity would give credibility to this possibility. We hypothesised that such an

outcome could be due either early differentiated cells that still expressed markers

characteristic of naïve cells or the inclusion of stem cell-like memory cells that co-

express CD27 and CD45RA[46]. Thirdly, these cells are not antigen-primed, but rather

akin to virtual T cells[47,48]. It has been shown that inflammatory conditions elicit virtual

CD8+ T cells in the absence of specific antigen[49]. Further investigations are required

to decipher the mechanism responsible for the diverse clonality found in naïve cells

and whether inflammatory conditions in iHEU maybe driving this.

Unlike the somatically rearranged TCR repertoire, NK cell diversity arises through

combinatorial expression of germline encoded receptors that sense expression of

MHC class I molecules through surface bound KIR[23]. Previous viral encounters have

been shown to augment NKR diversity and in some instances consequently reduce

the cytolytic capacity of NK cells[50,51]. We observed compositional differences of the

NK cell clusters between iHEU and iHUU at week 4 in which iHEU had marginally

higher memory-like/terminally differentiated NK cells (NKG2A-Per+CD57+) albeit not

significantly so after adjusting for multiple comparisons. Enhanced cytolytic activity

and elevated levels of the degranulation marker, CD107a, in iHEU compared to iHUU

at birth have been previously reported[21]. Certain NK cell clusters at birth were superior

in predicting both pertussis and rotavirus vaccine-induced IgG and IgA at 3-9 months

compared to CD4+ T cells. Depending on the immunological milieu, NK cells can either

negatively or positively regulate adaptive immunity, including promoting

immunoglobulin isotype switching and enhancing antibody production[52,53]. This finding

was found for both iHEU and iHUU and mechanistic investigations would need to

determine the role of NK cells in vaccine-induced antibodies.

The novelty of our study lies with the longitudinal follow-up of our infants with matched

samples for multiple analyses allowing us to detail the ontogeny of T and NK cells from

birth to 36 weeks of life and their relation with vaccination. The limitation of our study

is the small sample volumes collected from the infants and our consequent inability to

investigate functional changes in the immune subsets identified. A further limitation

was the inability to tease out the effect of HIV from ARV exposure since all pregnant

women living with HIV receive ARV treatment as a standard of care[54].

In conclusion, our data show a transitional immunity during infancy, which is impacted

by HIV/ARV exposure in a sequential manner starting with alteration of NK cells

followed by T cell differentiation. Of particular importance, is the finding that TCR

clonotypic diversity is significantly lower in iHEU from birth suggesting in utero skewing

well before T cell memory formation. In addition, our data show that the composition

of early life NK cells could predict vaccine responses in the first 9 months of life,

although how this is accomplished will require further studies. Lastly, we show here a

comprehensive phenotypic view of early life immune changes in response to HIV/ARV

exposure. These changes may well be linked to the observed disparities in co-

morbidities between iHEU and iHUU.

Methods

Study cohort

We analysed infants who were delivered by mothers living with and without HIV

infection who were enrolled prospectively from birth and followed up until 9 months of

age as previously described (Table S2)[55]. Briefly, pregnant women ≥18 years who

recently delivered <12 hr were enrolled in the study together with their respective

infants following signed informed consent. All mothers living with HIV received

combined antiretroviral treatment during pregnancy under the Option B programme.

The enrolment was restricted to infants having birth weight ≥ 2.5 kg, gestation ≥ 36

weeks and no complication experienced during delivery. Of those delivered by

mothers living with HIV, viral transmission was assessed by performing HIV DNA PCR

test after 6 weeks of life, and those who tested positive for perinatal HIV infection were

excluded from further analysis. The infants participating in this study received

childhood vaccines according to the South African Extended Program of Immunization

(EPI). This included administration of acellular pertussis vaccine at weeks 6, 10 and

14 and rotavirus vaccine at weeks 6 and 14. Blood samples were collected at birth,

weeks 4, 15 and 36 for isolation of plasma and peripheral blood mononuclear cells

(PBMC). Demographic characteristics of the mothers and their respective infants are

included in the study analysis and summarised in Table S2.

Plasma and peripheral blood mononuclear cell processing

Infant blood samples (0.5-3 mL) were collected into sodium heparin tubes and processed within 6hr. Plasma and PBMC were isolated using ficoll centrifugation. Plasma samples were stored at -80°C while PBMC were cryopreserved in 90% Fetal Calf serum (FCS) with 10% DMSO in liquid nitrogen.

Intracellular and surface staining of PBMC for mass cytometry

A total of 278 infant samples (including 27 duplicates) were used to assess the longitudinal changes of immune cells using mass cytometry and TCR sequencing. PBMC samples were retrieved from liquid nitrogen and thawed at 37°C before being transferred into RPMI 1640 media supplemented with 10% FCS and 10KU Benzonase. Cells were centrifuged, washed twice in PBS and counted using a TC20 cell counter (Biorad). We aimed to stain $2x10^6$ viable cells for mass cytometry, however cell recovery varied by infant and age, with fewer cells collected at earlier time points due to smaller blood volumes. PBMC samples were split into 2 aliquots with $2x10^6$ (or ¾ for cells with $<2x10^6$) used for mass cytometry staining and $1x10^6$ (or ¼ for cells with $<2x10^6$) used for TCR sequencing. Lyophilised surface and intracellular antibody mixtures[56] stored at 4°C were reconstituted in CyFACS buffer (PBS, 2% FCS) and permeabilization buffer (eBioscience permwash) respectively for cell labelling. Table S1 lists the antigen targets included in the mass cytometry antibody panel. Prior to antibody labelling, the cells were stained with cisplatin in PBS to determine cell viability followed by staining with surface antibodies for 20 min at room temperature. Cells were then fixed with 2% paraformaldehyde solution, permeabilised and stained with intracellular antibodies at 4°C for 45 min. Upon which cells were washed with CyFACS buffer and resuspended with 2% paraformaldehyde solution containing

Iridium DNA intercalator overnight at 4°C. Samples were then washed with PBS and

resuspended in MilliQ water containing EQ Four Element calibration beads (10% v,v,

Fluidigm) prior to acquisition using CyTOF 2 instrument (DVS Sciences).

Fluorescent activated cell sorting of naïve and memory T cells.

The remaining PBMC from each sample were used to sort for naïve and memory CD4⁺

and CD8⁺ T cells using fluorescent activated cell sorting. Cells were labelled by

surface staining using T cell markers including CD3, CD4 and CD8 and memory

markers CD27, CD45RA and CCR7 (Table S2 & Figure S3). Naïve cells were denoted

as those co-expressing CCR7, CD45RA and CD27 while the other remaining cells

were regarded as memory cells. A BD FACSAria Fusion (BD) was used for 4-way

sorting of CD4 and CD8 naïve and memory T cells from each sample and collected

directly into FCS. Sorted cells were centrifuged, resuspended in cell RNAProtect

(Qiagen) and stored at -40°C until processing for TCR sequencing.

<u>*Bulk TCR sequencing*</u>

RNA was purified from each sorted sample using the RNAeasy Plus Micro kit (Qiagen)

and libraries were prepared for TCR sequencing as described by Rubelt, et al.[57]. This

was performed at the University of Cape Town. Briefly, purified RNA was reverse

transcribed in a Rapid amplification of cDNA ends (RACE) reaction using

SMARTScribe Reverse Transcriptase using previously designed oligos (isoC-5′-

GTCAGATGTGTATAAGAGACAGnnnnnnnnnnnCGATAGrGrGrG -3′-C3_Spacer and

for the Poly A tail 5′- GTGTCACGTACAGAGTCATCtttttttttttttttttttttttttttttttttt -3′ VN) that

captures polyadenylated transcripts and introduces a 10bp unique molecular identifier

(UMI) at the 5' end of the product. cDNA was purified with AmpureXP beads (Beckman

Coulter) before whole transcriptome amplification using Advantage 2 Polymerase

(Clontech) using oligos that introduce Illumina Nextera Multiplex Identifier (MID) P5

Adapter sequences. The libraries were then sent to Stanford University (Department

of Microbiology and Immunology, Stanford University) in a blinded fashion, where

TCRβ-specific amplification was achieved with Q5 Hot Start Master Mix (NEB) using

constant region-specific oligos, simultaneously introducing P7 MID sequences. Final

sequencing libraries were purified using SPRISelect beads (Beckman Coulter),

quantified using the Agilent TapeStation and pooled for sequencing. Paired-end

sequencing was performed on an Illumina NovaSeq SP with 2x250 cycles, performed

by the Chan-Zuckerberg Biohub Initiative.

Quantification of vaccine antibody responses

Infant pertussis specific antibody responses were measured at birth, 4, 15 and 36

weeks of age using a commercial human IgG ELISA kit (Abcam). Plasma samples

were diluted 1:100 into sample diluent antibody titres measured in duplicates

according to manufacture instructions.

Rotavirus IgA titres were determined by enzyme immunoassay (EIA) and performed

in a blinded manner in the Division of Infectious Disease, Department of Pediatrics,

Cincinnati Childrens' Hospital Medical Centre, Cincinnati, Ohio. Briefly, purified rabbit

anti-rotavirus IgG were immobilised on microtiter plate as capture antibodies. Lysates

from rotavirus (strains RV3 and 8912) and mock infected cells (control) were added to

the immobilised capture antibodies to bind rotavirus antigens and any uncaptured

antigens were washed off with PBS, 0.05% Tween20. Reference standards, control

and test samples were diluted in PBS, 0.05% Tween20, 1% non-fat dry milk and 50

uL added to microtiter plate for antigen binding. Bound antibodies were detected by

biotyinylated goat anti-human IgA (Jackson Laboratories) and the addition of peroxidase conjugated avidin:biotin (Vector Laboratories, Inc., Burlingame, CA). Chromogenic signal was generated by addition of substrate O-phenylenediamine (Sigma) and reaction stopped after 30 min using 1M H_2SO_4. Absorbance was measured at 492_{nm} on a Molecular Devices SpectraMax Plus plate reader. The reference standard was assigned 1000 arbitrary units (AU) and a four-parameter logistic regression was used to extrapolate anti-rotavirus IgA tires using SoftMax software.

Rotavirus neutralisation titres were determined as previously described[58]. In this study we used Wa G1P8, 1076 G4P6 and DS-1 G2P4 virus strains obtained from the National Institute of Health (NIH). Plasma samples were serially diluted and incubated with the rotavirus strain for neutralization prior to adding the mixture to susceptible MA104 (monkey kidney) cell line for overnight incubation. Cell lysates were used to determine the level of un-neutralized rotavirus antigens using the EIA as described using guinea pig anti-rotavirus antiserum to measure captured rotavirus antigens. Rabbit anti-guinea pig IgG conjugated to horseradish peroxidase (Jackson ImmunoResearch) was used to detect bound antibodies, and chromogenic signal generated using OPD. The amount of rotavirus present in the resuspended lysate from each well was inversely related to the amount of neutralizing antibody present in the plasma. Each plasma dilution series as modelled using a logistic regression function. For each fitted curve the dilution which corresponds to a 40% response (ED_{40}), compared to the virus controls, was determined and reported as the neutralization titer. The ED_{40} represents the titre of the serum against a given virus, which represents a 60% reduction in amount of virus.

Data analysis

Mass cytometry

Processing

Post CyTOF acquisition, the files were normalized using Premessa R package and

target populations were manually gated using FlowJo software (version 10.5.3,

TreeStar). The target populations were exported as FCS files for downstream analysis

using R, an open-source statistical software[59]. All files for the target populations were

transformed by applying inverse hyperbolic sine with a cofactor of 5 and subjected to

doublet detection and removal using computeDoubletDensity from the scDblFinder

package[60].

Dimensional reduction

Median marker expressions for each sample were used to compute Euclidean

distance matrix to determine multidimensional scaling (MDS) coordinates using the

cmdscale function in R. MDS analysis were used to visualize immune cell trajectories.

Further, marker expression intensities were used to implement Uniform Manifold

Approximation and Projection (UMAP) while preserving local density distribution for

visualization using densMAP function from the denvis R package[61].

Cell Clustering

High resolution clustering of the target cell populations was performed using FlowSOM

algorithm and the resulting clusters grouped into metaclusters using

ConsensusClusterPlus available in the CATALYST R package[62]. The immune cell

clusters were visualised using hierarchical clustering and UMAP embedding. Centred

log-ratios of the relative abundance of the cell clusters per sample were determined

and used in Principal Component Analysis (PCA) for assessing immune cell

compositional differences.

TCR sequencing

Pre-processing of TCR sequencing data

Raw sequencing data was pre-processed by pRESTO[63]. Briefly, reads with mean

Phred quality scores less than 20 were discarded, UMIs were extracted from the first

10bp of read 1, and primer sequences masked. Next, a single consensus sequence

was constructed from reads sharing the same UMI barcode and paired-end consensus

sequences assembled into a full-length TCR sequence. In the case of non-overlapping

mate-pairs, the human TRB reference as used to properly space non-overlapping

reads. After removal of duplicate sequences, full-length reads were aligned and

clonotypes assembled using MiXCR[64,65].

T cell immune repertoire analysis

To perform quality control on our bulk TCR sequencing data, we removed samples

where fewer than 10 clones were detected ($n = 44$ samples) and where more clones

were detected than cells were sorted ($n = 18$ samples). This yielded a total of 861

samples for downstream analysis. The immunarch packag[66] operating in the open-

source statistical software R was used for all downstream immune repertoire analysis,

including calculation of CDR3β lengths and repertoire diversity by the Inverse Simpson

index and richness using Choa1.

Identification and annotation of T cell specificity groups

The GLIPH2 algorithm was used to establish T cell specificity groups, clusters of

CDR3 sequences that are predicted to bind to the same antigen, by discovering

groups of CDR3β sequences that share either global or local motifs[27,28]. To assign

specificity annotations to identified CDR3β sequence clusters, we adapted an

approach described by Chiou, et al[67]. Briefly, we collected human CDR3β sequences

with known specificity from VDJdb (https://vdjdb.cdr3.net/) and identified CDR3β

sequences within this database that could form TCR specificity groups. From 47,107

CDR3β sequences, this process identified 11,648 specificity groups comprised of

25,759 unique CDR3β sequences. We next combined these 25,759 annotated

sequences with 108,183 unique experimental CDR3β sequences from our bulk TCR

repertoire profiling cohort. This was performed separately for CD4 and CD8 T, as there

are differences between gene usage frequencies between CD4 and CD8 T cells that

can impact specificity predictions. We prioritized TCR specificity groups with at least 3

distinct CDR3β sequences. This process yielded 17,154 CDR3β sequences in 11,629

specificity groups for CD4 T cells, and 15,107 CDR3β sequences in 9,911 specificity

groups for CD8 T cells.

Statistical analysis

All statistical analysis and graphical visualization of the data was performed on the

open R software[59]. Spearman's rank correlation was used to test for the correlations

between the frequencies of the cell clusters with either infant age or antibody

responses and p-values adjusted for multiple comparisons using false discovery rates

(FDR). To compare differences in the abundance of cell clusters between iHEU and

iHUU the generalised linear model from the diffcyt package was used[68]. Pairwise

comparisons of medians between groups were performed using Wilcoxon-rank test or

Kruskal-Wallis for multiple group comparisons and the p-values adjusted for multiple comparisons using Benjamin Hochberg correction. Generalised linear mixed model (GLMM) with bootstrap resampling from the R package CytoGLMM[69] was used to identify NK cell markers that are predictive of HIV exposure at birth. To determine cell clusters that were predictive of antibody responses following vaccination, we used the multivariate modelling algorithm from MUVR that incorporates recursive variable selection using repeated double cross-validation within partial least squares modelling[29].

Acknowledgements

We would like to acknowledge the technical staff that assisted in the completion of this study. This includes Euan Johnston, Emily Tangie and Carine Kilola who assisted with the CyTOF experiments, Mikayla Stabile, Michelle Leong, Kassandra Pinedo, and Nicole Prins, for assistance with TCR RNA-seq processing, and Llyod Leach for assistance with measuring pertussis antibody responses.

Author contribution

Conceived by CMG, CAB and HBJ, Method set-up and validation by SD, AW, SC, TR, FR, HH, and MD. Study participant enrolment, sample collection and processing by HBJ and BA. Experimental investigations by SD. Data and statistical analysis by SD, AW and SH. Original draft by SD. Reviewed and edited by all authors.

Declaration of Interest

All authors declare no conflict of interest of any of the material included in this manuscript.

References

1.	Perin, J. *et al.* Global, regional, and national causes of under-5 mortality in 2000–19: an updated systematic analysis with implications for the Sustainable Development Goals. *The Lancet Child and Adolescent Health* **6**, 106–115 (2022).

2.	Zimmermann, P. & Jones, C. E. Factors That Influence Infant Immunity and Vaccine Responses. *The Pediatric infectious disease journal* **40**, S40–S46 (2021).

3.	Pasetti, M. F. *et al.* Maternal determinants of infant immunity: Implications for effective immunization and maternal-child health. *Vaccine* **38**, 4491–4494 (2020).

4.	Sanidad, K. Z. & Zeng, M. Y. Neonatal gut microbiome and immunity. *Current Opinion in Microbiology* **56**, 30–37 (2020).

5.	Goenka, A. & Kollmann, T. R. Development of immunity in early life. *Journal of Infection* **71**, S112–S120 (2015).

6.	Aagaard, K. *et al.* The Placenta Harbors a Unique Microbiome. *ScienceTranslationalMedicine* **6**, 229–262 (2014).

7.	Tosato, F. *et al.* Lymphocytes subsets reference values in childhood. *Cytometry Part A* **87**, 81–85 (2015).

8.	Howie, P. W., Forsyth, J. S., Ogston, S. A., Clark, A. & Du Florey, V. C. Protective effect of breat feeding against infection. *British Medical Journal* **300**, 11–16 (1990).

9.	Sankar, M. J. *et al.* Optimal breastfeeding practices and infant and child mortality: A systematic review and meta-analysis. *Acta Paediatrica, International Journal of Paediatrics* **104**, 3–13 (2015).

10.	Anderson, K. *et al.* Increased infectious-cause hospitalization among infants who are HIV-exposed uninfected compared with HIV-unexposed. *AIDS* **35**, 2327–9 (2021).

11.	Brennan, A. T. *et al.* A meta-analysis assessing all-cause mortality in HIVexposed uninfected compared with HIV-unexposed uninfected infants and children. *Aids* **30**, 2351–2360 (2016).

12.	Cohen, C. *et al.* Epidemiology of Acute Lower Respiratory Tract Infection in HIV- Exposed Uninfected Infants. *Pediatrics* **137**, e20153272 (2016).

13.	Slogrove, A. L., Goetghebuer, T., Cotton, M. F., Singer, J. & Bettinger, J. Pattern of Infectious Morbidity in HIV-Exposed Uninfected Infants and Children. *Frontiers in Immunology* **7**, 1–8 (2016).

14.	Brennan, A. T. *et al.* A meta-analysis assessing diarrhea and pneumonia in HIV-exposed uninfected compared to HIV-unexposed uninfected infants and children. *JAIDS Journal of Acquired Immune Deficiency Syndromes* **82**, 1–8 (2019).

15.	Slogrove, A. L., Powis, K. M., Johnson, L. F., Stover, J. & Mahy, M. Estimates of the global population of children who are HIV-exposed and uninfected, 2000–18: a modelling study. *The Lancet Global Health* **8**, e67–e75 (2020).

16.	Dzanibe, S. *et al.* Stereotypic Expansion of T Regulatory and Th17 Cells during Infancy Is Disrupted by HIV Exposure and Gut Epithelial Damage. *The Journal of Immunology* **208**, 27–37 (2022).

17.	Uksila, J., Lassila, O. & Hirvonen, T. Natural killer cell function of human neonatal lymphocytes. *Clinical and Experimental Immunology* **48**, 649–654 (1982).

18. Choi, S. S. *et al.* Interleukin-15 enhances cytotoxicity, receptor expression, and
expansion of neonatal natural killer cells in long-term culture. *Clinical and*
Diagnostic Laboratory Immunology **11**, 879–888 (2004).
19. Xing, D. *et al.* Cord blood natural killer cells exhibit impaired lytic
immunological synapse formation that is reversed with il-2 exvivo expansion.
Journal of Immunotherapy **33**, 684–696 (2010).
20. Strauss-Albee, D. M., Liang, E. C., Ranganath, T., Aziz, N. & Blish, C. A. The
newborn human NK cell repertoire is phenotypically formed but functionally
reduced. *Cytometry Part B - Clinical Cytometry* **92**, 33–41 (2017).
21. Smith, C. *et al.* Altered natural killer cell function in HIV-exposed uninfected
infants. *Frontiers in Immunology* **8**, 1–13 (2017).
22. Slyker, J. A. *et al.* The impact of HIV-1 infection and exposure on natural killer
(NK) cell phenotype in kenyan infants during the first year of life. *Frontiers in*
Immunology **3**, 1–16 (2012).
23. Vilches, C. & Parham, P. KIR: Diverse, rapidly evolving receptors of innate and
adaptive immunity. *Annual Review of Immunology* **20**, 217–251 (2002).
24. Sun, J. C., Beilke, J. N. & Lanier, L. L. Adaptive immune features of natural
killer cells. *Nature* **457**, 557–561 (2009).
25. Muntasell, A., Magri, G., Pende, D., Angulo, A. & López-Botet, M. Inhibition of
NKG2D expression in NK cells by cytokines secreted in response to human
cytomegalovirus infection. *Blood* **115**, 5170–5179 (2010).
26. Flommersfeld, S. *et al.* Fate mapping of single NK cells identifies a type 1
innate lymphoid-like lineage that bridges innate and adaptive recognition of
viral infection. *Immunity* **54**, 2288-2304.e7 (2021).
27. Glanville, J. *et al.* Identifying specificity groups in the T cell receptor repertoire.
Nature **547**, 94–98 (2017).
28. Huang, H., Wang, C., Rubelt, F., Scriba, T. J. & Davis, M. M. Analyzing the
Mycobacterium tuberculosis immune response by T-cell receptor clustering
with GLIPH2 and genome-wide antigen screening. *Nature Biotechnology* **38**,
1194–1202 (2020).
29. Shi, L., Westerhuis, J. A., Rosén, J., Landberg, R. & Brunius, C. Variable
selection and validation in multivariate modelling. *Bioinformatics* **35**, 972–980
(2019).
30. Olin, A. *et al.* Stereotypic Immune System Development in Newborn Children.
Cell **174**, 1277-1292.e14 (2018).
31. Haynes, B. Y. B. E., Martin, M. E., Kay, H. H. & Kurtzberg, J. Early events in
human T cell ontogeny. Phenotypic characterization and immunohistologic
localization of T cell precursors in early human fetal tissues. *The Journal of*
experimental medicine **168**, 1061–80 (1988).
32. Haynes, B. F. & Heinly, C. S. Early human T cell development: Analysis of the
human thymus at the time of initial entry of hematopoietic stem cells into the
fetal thymic microenvironment. *Journal of Experimental Medicine* **181**, 1445–
1458 (1995).
33. Scheible, K. M. *et al.* T cell developmental arrest in former premature infants
increases risk of respiratory morbidity later in infancy. *JCI insight* **3**, (2018).
34. Semmes, E. C. *et al.* Understanding Early-Life Adaptive Immunity to Guide
Interventions for Pediatric Health. *Frontiers in Immunology* **11**, 1–14 (2021).
35. Walker, R. W., Clemente, J. C., Peter, I. & Loos, R. J. F. The prenatal gut
microbiome: are we colonized with bacteria in utero? *Pediatric Obesity* **12**, 3–
17 (2017).

36. Rich, K. C., Siegel, J. N., Jennings, C., Rydman, R. J. & Landay, a L. Function
and phenotype of immature CD4+ lymphocytes in healthy infants and early
lymphocyte activation in uninfected infants of human immunodeficiency virus-
infected mothers. *Clinical and diagnostic laboratory immunology* **4**, 358–361
(1997).
37. Ono, E. *et al.* Imbalance of naive and memory T lymphocytes with sustained
high cellular activation during the first year of life from uninfected children born
to HIV-1-infected mothers on HAART. *Brazilian Journal of Medical and
Biological Research* **41**, 700–708 (2008).
38. Wood, L. F., Wood, M. P., Fisher, B. S., Jaspan, H. B. & Sodora, D. L. T cell
activation in South African HIV-exposed infants correlates with ochratoxin a
exposure. *Frontiers in Immunology* **8**, 1–8 (2017).
39. Garcia-Knight, M. A. *et al.* Altered memory T-cell responses to Bacillus
Calmette-Guerin and Tetanus Toxoid vaccination and altered cytokine
responses to polyclonal stimulation in HIV-exposed uninfected Kenyan infants.
PLoS ONE **10**, 1–19 (2015).
40. Smith, C. *et al.* Human Immunodeficiency Virus Exposure but Not Early
Cytomegalovirus Infection Is Associated With Increased Hospitalization and
Decreased Memory T-Cell Responses to Tetanus Vaccine. *Journal of
Infectious Diseases* **221**, 1167–1175 (2020).
41. Pérez, Y. B. *et al.* Impaired T helper cell responses in human
immunodeficiency virus - exposed uninfected newborns. *Immunity
Inflammation and Disease* **9**, 1541–1553 (2021).
42. Gabriel, B. *et al.* Analysis of the TCR Repertoire in HIV-Exposed but
Uninfected Infants. *Scientific Reports* **9**, (2019).
43. Holditch, S. J. *et al.* Decay kinetics of HIV-1 specific cell responses in
vertically HIV-1 exposed seronegative infants. *Frontiers in Immunology* **2**, 1–6
(2012).
44. Pušnik, J. *et al.* Expansion of Stem Cell-Like CD4 + Memory T Cells during
Acute HIV-1 Infection Is Linked to Rapid Disease Progression . *Journal of
Virology* **93**, (2019).
45. Ribeiro, S. P. *et al.* The CD8 + Memory Stem T Cell (T SCM) Subset Is
Associated with Improved Prognosis in Chronic HIV-1 Infection . *Journal of
Virology* **88**, 13836–13844 (2014).
46. Gattinoni, L. *et al.* A human memory T cell subset with stem cell-like
properties. *Nature Medicine* **17**, 1290–1297 (2011).
47. Savid-Frontera, C. *et al.* Exploring the immunomodulatory role of virtual
memory CD8+ T cells: Role of IFN gamma in tumor growth control. *Frontiers in
Immunology* **13**, 1–18 (2022).
48. Lee, J. Y., Hamilton, S. E., Akue, A. D., Hogquist, K. A. & Jameson, S. C.
Virtual memory CD8 T cells display unique functional properties. *Proceedings
of the National Academy of Sciences of the United States of America* **110**,
13498–13503 (2013).
49. Rolot, M. *et al.* Helminth-induced IL-4 expands bystander memory CD8+ T
cells for early control of viral infection. *Nature Communications* **9**, (2018).
50. Wilk, A. J. & Blish, C. A. Diversification of human NK cells: Lessons from deep
profiling. *Journal of Leukocyte Biology* **103**, 629–641 (2018).
51. Strauss-Albee, D. M. *et al.* Human NK cell repertoire diversity reflects immune
experience and correlates with viral susceptibility. *Science Translational
Medicine* **7**, (2015).

52. Rydyznski, C. E. & Waggoner, S. N. Boosting vaccine efficacy the natural
(killer) way. *Trends in Immunology* **36**, 536–546 (2015).
53. Rydyznski, C. E. *et al.* Affinity Maturation Is Impaired by Natural Killer Cell
Suppression of Germinal Centers. *Cell Reports* **24**, 3367-3373.e4 (2018).
54. Wessels, J. *et al.* The updated South African National Guideline for the
Prevention of Mother to Child Transmission of Communicable Infections
(2019). *Southern African Journal of HIV Medicine* **21**, 1–8 (2020).
55. Tchakoute, C. T. *et al.* Breastfeeding mitigates the effects of maternal HIV on
infant infectious morbidity in the Option B+ era. *AIDS (London, England)* **32**,
2383–2391 (2018).
56. Vendrame, E. *et al.* Profiling of the Human Natural Killer Cell Receptor-Ligand
Repertoire. *Journal of Vizualized Experiments* **165**, 139–148 (2020).
57. Rubelt, F. *et al.* Individual heritable differences result in unique cell lymphocyte
receptor repertoires of naïve and antigen-experienced cells. *Nature
Communications* **7**, 1–12 (2016).
58. Knowlton, D. R., Spector, D. M. & Ward, R. L. Development of an improved
method for measuring neutralizing antibody to rotavirus. *Journal of Virological
Methods* **33**, 127–134 (1991).
59. RCoreTeam. R: A language and environment for statistical computing. Preprint
at (2021).
60. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D.
Doublet identification in single-cell sequencing data using scDblFinder.
F1000Research **10**, 979 (2022).
61. Narayan, A., Berger, B. & Cho, H. Assessing single-cell transcriptomic
variability through density-preserving data visualization. *Nature Biotechnology*
39, 765–774 (2021).
62. Crowell, H. L., Zanotelli, V. R. T., Chevrier, S. & Robinson, M. D. CATALYST:
Cytometry dATa anALYSis Tools. Preprint at (2021).
63. Vander Heiden, J. A. *et al.* PRESTO: A toolkit for processing high-throughput
sequencing raw reads of lymphocyte receptor repertoires. *Bioinformatics* **30**,
1930–1932 (2014).
64. Bolotin, D. A. *et al.* MiXCR: Software for comprehensive adaptive immunity
profiling. *Nature Methods* **12**, 380–381 (2015).
65. Bolotin, D. A. *et al.* Antigen receptor repertoire profiling from RNA-seq data.
Nature biotechnology **35**, 908–911 (2017).
66. Nazarov, V. I. *et al.* immunarch: Bioinformatics Analysis of T-Cell and B-Cell
Immune Repertoires. Preprint at (2022).
67. Chiou, S. H. *et al.* Global analysis of shared T cell specificities in human non-
small cell lung cancer enables HLA inference and antigen discovery. *Immunity*
54, 586-602.e8 (2021).
68. Weber, L. M. diffcyt: Differential discovery in high-dimensional cytometry via
high-resolution clustering. Preprint at (2021).
69. Seiler, C. *et al.* CytoGLMM: conditional differential analysis for flow and mass
cytometry experiments. *BMC Bioinformatics* **22**, 1–14 (2021).
892

Figure 1: Immunophenotypic trajectory of infant peripheral blood mononuclear cells (PBMC) in the first 9 months of life. **A)** Dimensional reduction of infant PBMC Immune lineage clusters embedded on uniform manifold approximation and proximation (UMAP) with density preservation and table summarising immune cell cluster phenotype. **B)** Relative abundance of immune lineage clusters measured in cord blood (CB) and infant peripheral blood at birth (BTH), weeks (WK) 4, 15 and 36. **C)** Spearman's rank correlation between proportion of immune lineage cell clusters and infants age from birth until week 36. **D)** Age related maturation trajectory of CD4+ and CD8+ T cells and NK cells depicted using multidimensional scaling coordinates derived from median marker expressions for each infant samples. **E)** Boxplots summarizing MDS coordinate associated with infant age for CD4+ and CD8+ T cells and NK cells.

Figure 2: Divergent T cell memory differentiation in HIV-exposed uninfected infants (iHEU) compared to HIV-unexposed uninfected infants (iHUU). **A & D)** Uniform manifold approximation and proximation (UMAP) with density preservation showing

dimensional reduction of FlowSOM clusters of CD4+ and CD8 T cells **B &E)** Relative

abundances of CD4+ and CD8 + T cell clusters measured in cord blood (CB) and

infant peripheral blood at birth (BTH), weeks (WK) 4, 15 and 36. **C &F)** Age related

phenotypic composition of CD4+ and CD8+ T cells depicted as principal component

(PC) coordinates of centred log-odd ratios of FlowSOM clusters for each infant

samples and arrows indicating contribution of each cell cluster in scatter of PC

components. **G &H)** Boxplots comparing PC coordinates of CD4+ and CD8+ T cells

between iHEU and iHUU longitudinally. **I & J)** Generalized linear mixed model (GLMM)

comparing the abundances of CD4+ and CD8+ T cell clusters between iHEU and

iHUU at weeks 15 and 36.

Figure 3: Early life immunophenotypic alteration in NK cells in HIV-exposed uninfected

infants (iHEU) compared to HIV-unexposed uninfected infants (iHUU). **A)** Uniform

manifold approximation and proximation (UMAP) with density preservation showing

dimensional reduction of FlowSOM clusters on NK cells and table summarising

immune cell cluster phenotype. **B)** Relative abundances of NK cell clusters measured

in cord blood (CB) and infant peripheral blood at birth (BTH), weeks (WK) 4, 15 and

36. **C)** Age related phenotypic composition of NK cells depicted as principal

component (PC) coordinates of centred log-odd ratios of FlowSOM clusters for each

infant samples and arrows indicating contribution of each cell cluster in scatter of PC

components. **D)** Boxplots comparing PC coordinates of NK cells between iHEU and

iHUU

Figure 4: Premature CD4+ and CD8+ T cell receptor (TCR) repertoire skewing in HIV-

exposed uninfected infants (iHEU) relative to HIV-unexposed uninfected infants

(iHUU). **A)** Boxplots comparing Inverse Simpson TCR diversity scores between iHUU

and iHEU at birth and weeks 4, 15 and 36. **B)** Boxplots comparing Chao1 TCR

clonotype richness between iHUU and iHEU at birth and weeks 4, 15 and 36. **C &D**)

Spearman's rank correlation between naive CD4+ and CD8+ T cell Inverse Simpson

scores and frequencies of FlowSOM clusters for CD4+ and CD8+ T cell clusters

respectively. **E**) . GLIPH analysis showing antigen specificity groups of the top 100

TCR clones that are significantly different between iHEU and iHUU and between naïve

and memory CD4+ and CD8+ T cell subsets.

Figure 5 Association of vaccine antibody responses to immune cell phenotypes. **A)**

Comparing IgG levels against pertussis between HIV-exposed uninfected infants

(iHEU) and HIV-unexposed uninfected infants (iHUU), grey shaded area indicate

threshold IgG levels for protective pertussis vaccine response. **B & C**) Boxplots

comparing rotavirus specific IgA and neutralization titres between iHEU and iHUU at

week 36. **D, E & F)** Summary of ROC analysis using the latent variable axis-1 derived

from partial least square discriminate analysis (PLS-DA) of NK, CD4 and CD8 T cell

clusters determined to be best predictors at birth and week 4 of pertussis antibody

responses at weeks 15 and 36 and rotavirus antibody response at week 37. **G)**

Spearman's correlation between abundances FlowSOM clusters of CD4+ and CD8+

T cells and NK cells and anti-pertussis IgG titres measured at week 15 and 36.

Table 1: Phenotypic description of CD4$^+$ T cell clusters idenfied by unsupervised cell clustering.

Cluster	Phenotype	(%)
(1)	CCR4+CCR6-CXCR3- (Th2)	5.07
(2)	CD45RA+CD27+CCR7+ (Naïve)	84.1
(3)	CD45RA+CD27-CCR7-CD57+Per+PD-1+ (Cytotoxic TD)	1.3
(4)	CD38+HLA-DR+CD45RA+CD27+CCR7+ (Activated cells)	4.7
(5)	CCR4+CCR6+CXCR3+CD57-	0.7
(6)	CCR4+CCR6+CXCR3+CD57+	0.2
(7)	Per+Ki67+	0.3
(8)	CD127-CD25+CD39+TIGIT+ (Treg cells)	0.9
(9)	Per+Ki67+CD25+	0.3
(10)	Per+Ki67-HLA-DR+CD69+CD39+	0.1
(11)	Per+Ki67+CD27+Fc$_\varepsilon$RI$_\gamma$+	0.2
(12)	PD-1+CD38+	2.07
(13)	CD39+CD45RA+CCR7-CD27-	0.1

Table 2: Phenotypic description of CD8$^+$ T cell clusters identified by unsupervised cell clustering.

Cluster	Phenotype	(%)
(1)	Per+CD45RA-CD27-CCR7-CD57+ (TD)	7.5
(2)	CD56+CD16+NkpK6+NKG2A+ (NKT-like)	0.4
(3)	HLA-DR+	2.7
(4)	CD45RA+CD27+CCR7+ (Naïve)	55.3
(5)	Per+CD57+	15.6
(6)	Per+CD45RA-CD27-CCR7-	18.4
(7)	Ki67+CD45RA+CD27+	0.1

Table 3: Phenotypic description of NK cell clusters identified by unsupervised cell clustering.

Cluster	Phenotype	(%)
(1)	(CD56^{dim}CD16^{dim}) Per+CD45RA+FcERIy	30.5

(2)	(CD56⁻CD16⁻) DNAM1+	20.7
(3)	(CD56⁻CD16⁺) FcERIy	1.8
(4)	(CD56ᵈⁱᵐCD16⁺) Per+CD45RA+CD38+2B4+	11.7
(5)	(CD56ᵈⁱᵐCD16⁺) Per+CD45RA+CD38+CD57+	12.4
(6)	(CD56ᵈⁱᵐCD16ᵈⁱᵐ) FcERIy+DNAM1+HLA-DR+CXCR3+CCR4+	1.4
(7)	(CD56⁻CD16⁺) CD38+CD39+CD27+Ki67+	0.9
(8)	(CD56⁺CD16ᵇʳⁱ) Per+CD38+CD45RA+DNAM1+2B4+Nkp30+NKp46+NKG2A+Siglec-7+	15.0
(9)	(CD56ᵇʳⁱCD16⁻) DNAM1+NKG2D+Nkp46+NKG2A+	4.9
(10)	(CD56⁺CD16⁺) Per+CD38+CD45RA+LILRB1+NTBA+2B4+Nk30+NKG2D+HLA-DR+CCR4+CCR7+PD-1+TIGIT+NKg2C+	0.7

Figure 1: Immunophenotypic trajectory of infant peripheral blood mononuclear cells (PBMC) in the first 9 months of life. **A)** Dimensional reduction of infant PBMC Immune lineage clusters embedded on uniform manifold approximation and proximation (UMAP) with density preservation and table summarising immune cell cluster phenotype. **B)** Relative abundance of immune lineage clusters measured in cord blood (CB) and infant peripheral blood at birth (BTH), weeks (WK) 4, 15 and 36. **C)** Spearman's rank correlation between proportion of immune lineage cell clusters and infants age from birth until week 36. **D)** Age related maturation trajectory of CD4+ and CD8+ T cells and NK cells depicted using multidimensional scaling coordinates derived from median marker expressions for each infant samples. **E)** Boxplots summarizing MDS coordinate associated with infant age for CD4+ and CD8+ T cells and NK cells.

Figure 2: Divergent T cell memory differentiation in HIV-exposed uninfected infants (iHEU) compared to HIV-unexposed uninfected infants (iHUU). **A & D)** Uniform manifold approximation and proximation (UMAP) with density preservation showing dimensional reduction of FlowSOM clusters of CD4+ and CD8 T cells **B &E)** Relative abundances of CD4+ and CD8 + T cell clusters measured in cord blood (CB) and infant peripheral blood at birth (BTH), weeks (WK) 4, 15 and 36. **C &F)** Age related phenotypic composition of CD4+ and CD8+ T cells depicted as principal component (PC) coordinates of centred log-odd ratios of FlowSOM clusters for each infant samples and arrows indicating contribution of each cell cluster in scatter of PC components. **G &H)** Boxplots comparing PC coordinates of CD4+ and CD8+ T cells between iHEU and iHUU longitudinally. **I & J)** Generalized linear mixed model (GLMM) comparing the abundances of CD4+ and CD8+ T cell clusters between iHEU and iHUU at weeks 15 and 36, Positive log Fold change (FC) signified cell cluster frequencies were higher in iHEU compared to iHUU.

Figure 3: Early life immunophenotypic alteration in NK cells in HIV-exposed uninfected infants (iHEU) compared to HIV-unexposed uninfected infants (iHUU). **A)** Uniform manifold approximation and proximation (UMAP) with density preservation showing dimensional reduction of FlowSOM clusters on NK cells and table summarising immune cell cluster phenotype. **B)** Relative abundances of NK cell clusters measured in cord blood (CB) and infant peripheral blood at birth (BTH), weeks (WK) 4, 15 and 36. **C)** Age related phenotypic composition of NK cells depicted as principal component (PC) coordinates of centred log-odd ratios of FlowSOM clusters for each infant samples and arrows indicating contribution of each cell cluster in scatter of PC components. **D)** Boxplots comparing PC coordinates of NK cells between iHEU and iHUU.

Figure 4: Premature CD4+ and CD8+ T cell receptor (TCR) repertoire skewing in HIV-exposed uninfected infants (iHEU) relative to HIV-unexposed uninfected infants (iHUU). **A)** Boxplots comparing Inverse Simpson TCR diversity scores between iHUU and iHEU at birth and weeks 4, 15 and 36. **B)** Boxplots comparing Chao1 TCR clonotype richness between iHUU and iHEU at birth and weeks 4, 15 and 36. **C &D)** Spearman's rank correlation between naïve CD4+ and CD8+ T cell Inverse Simpson scores and frequencies of FlowSOM clusters for CD4+ and CD8+ T cell clusters respectively. **E)** . GLIPH analysis showing antigen specificity groups of the top 100 TCR clones that are significantly different between iHEU and iHUU and between naïve and memory CD4+ and CD8+ T cell subsets.

Figure 5 Association of vaccine antibody responses to immune cell phenotypes. **A)** Comparing IgG levels against pertussis between HIV-exposed uninfected infants (iHEU) and HIV-unexposed uninfected infants (iHUU), grey shaded area indicate threshold IgG levels for protective pertussis vaccine response. **B & C)** Boxplots comparing rotavirus specific IgA and neutralization titres between iHEU and iHUU at week 36. **D, E & F)** Summary of ROC analysis using the latent variable axis-1 derived from partial least square discriminate analysis (PLS-DA) of NK, CD4 and CD8 T cell clusters determined to be best predictors at birth and week 4 of pertussis antibody responses at weeks 15 and 36 and rotavirus antibody response at week 37. **G)** Spearman's correlation between abundances FlowSOM clusters of CD4+ and CD8+ T cells and NK cells and anti-pertussis IgG titres measured at week 15 and 36.