TARANIS interacts with VRILLE and PDP1 to modulate the

circadian transcriptional feedback mechanism in *Drosophila*

<u>Abbreviated Title</u>: Regulation of circadian rhythms by TARANIS

Oghenerukevwe Akpoghiran[1,2], Dinis J.S. Afonso[1,2], Yanan Zhang[1], and Kyunghee Koh[1*]

[1]Department of Neuroscience, the Farber Institute for Neurosciences, and Synaptic Biology

Center, Thomas Jefferson University, Philadelphia, USA. 19107

[2]These authors contributed equally.

*Correspondence should be addressed to Kyunghee Koh at <u>kyunghee.koh@jefferson.edu</u>

Number of pages: 47 pages

Number of figures and tables: 7 figures and 3 tables

Number of words: Abstract = 222/250; Introduction = 629/650; Discussion = 1237/1500.

<u>Conflict of Interest Statement</u>: The authors declare no competing financial interests.

<u>Acknowledgments:</u> This work was supported by grants from the National Institutes of Health

(R01NS086887 and R01NS109151 to K.K.). and funds from Thomas Jefferson University

Synaptic Biology Center (to K.K.). We thank the Bloomington *Drosophila* Stock Center and

National Institute of Genetics for fly stocks; *Drosophila* Genomics Research Center (supported

by NIH grant 2P40OD010949-10A1) for vectors; Drs. Amita Sehgal, Paul Hardin, Justin Blau,

and Jit Kong Cheong for antibodies and constructs; M. Boudinot and Dr. Francois Rouyer for the

FaasX software; Luke Kim and William Wei for technical assistance; and Drs. Yohei Kirino,

Joanna Chiu, Aaron Haeusler, Eun Young Kim, Sang Hyuk Lee, and James Jaynes, and

members of the Koh lab for helpful discussions.

Abstract

The molecular clock that generates daily rhythms of behavior and physiology consists of

interlocked transcription-translation feedback loops. In *Drosophila*, the primary feedback loop

involving the CLOCK-CYCLE transcriptional activators and the PERIOD-TIMELESS

transcriptional repressors is interlocked with a secondary loop involving VRILLE (VRI) and

PAR DOMAIN PROTEIN 1 (PDP1), a repressor and activator of *Clock* transcription,

respectively. Whereas extensive studies have found numerous transcriptional, translational, and

post-translational modulators of the primary loop, relatively little is known about the secondary

loop. In this study, we demonstrate that TARANIS (TARA), a *Drosophila* homolog of the TRIP-

Br/SERTAD family of transcriptional coregulators, functions with VRI and PDP1 to modulate

the circadian period and rhythm strength in *Drosophila*. Overexpression of TARA lengthens the

circadian period, whereas *tara* mutants exhibit reduced rhythmicity, a faster molecular clock, and

reduced expression of the PDF neuropeptide. We find that TARA can form a physical complex

with VRI and PDP1, enhancing their repressor and activator functions, respectively. The

conserved SERTA domain of TARA is required to regulate the transcriptional activity of VRI

and PDP1, and its deletion leads to reduced locomotor rhythmicity. Consistent with TARA's role

in enhancing VRI and PDP1 activity, overexpressing *tara* has a similar effect on the circadian

period and rhythm strength as simultaneously overexpressing *vri* and *Pdp1*. Together, our results

suggest that TARA modulates circadian behavior by enhancing the transcriptional activity of

VRI and PDP1.

Statement of Significance

Internal molecular clocks generating circadian rhythms of around 24 hours broadly impact

behavior and physiology, and circadian dysfunction is associated with various neurological and

metabolic diseases. The *Drosophila* circadian clock is a valuable model for understanding the

molecular mechanisms underlying daily rhythms. In this study, we identify a conserved gene,

taranis, as a novel regulator of the *Drosophila* molecular clock. We show that TARANIS

modulates circadian behavior by physically interacting with and enhancing the transcriptional

activity of clock proteins VRILLE and PDP1. Since mammalian homologs of VRILLE and

PDP1 also function in the molecular clock, our results have implications for understanding the

mammalian circadian clock.

Introduction

The molecular mechanisms of endogenous circadian oscillators are conserved across organisms, making *Drosophila melanogaster* a valuable model for understanding the genetics of the circadian clock (reviewed in Allada and Chung, 2010; Hardin, 2011; Tataroglu and Emery, 2015; Patke et al., 2020). The molecular clocks in flies and mammals consist of interlocked transcription-translation feedback loops that drive rhythmic gene expression. In the *Drosophila* clock, the CLOCK-CYCLE (CLK-CYC) heterodimer binds to the E-box elements in the *period* (*per*) and *timeless* (*tim*) promoters to activate their transcription (Allada et al., 1998), and the PER-TIM heterodimer inhibits CLK-CYC, suppressing their own transcription (Hardin et al., 1990). In an interlocked loop, CLK-CYC drives the transcription of *vri* and *Pdp1*. VRI and PDP1 competitively bind to the D-box elements in the *Clk* promoter to repress and activate *Clk* transcription, respectively (Cyran et al., 2003; Glossop et al., 2003). The mammalian clock involves analogous, although more complex, interlocked feedback loops, including nuclear receptors and mammalian homologs of VRI and PDP1 (reviewed in Partch et al., 2014; Takahashi, 2017; Patke et al., 2020). While many molecules underlying the transcriptional, translational, and post-translational modulation of the primary feedback loop have been identified (Allada and Chung, 2010; Zheng and Sehgal, 2012; Tataroglu and Emery, 2015; Patke et al., 2020), the mechanisms regulating the secondary interlocked loop are less well understood (Blau and Young, 1999; Cyran et al., 2003; Glossop et al., 2003; Benito et al., 2007; Zheng et al., 2009; Ling et al., 2012; Gunawardhana and Hardin, 2017; Gunawardhana et al., 2021).

VRI and PDP1 are bZIP transcriptional factors essential for proper development and circadian behavior (George and Terracol, 1997; Blau and Young, 1999; Reddy et al., 2000; Cyran et al., 2003; Glossop et al., 2003). Both regulate the circadian output pathway and the

molecular clock. Null mutants of *vri* and *Pdp1* are lethal, but knocking down either gene in clock

neurons leads to arrhythmic behavior despite a functioning molecular clock, suggesting their role

in the clock output (Blau and Young, 1999; Cyran et al., 2003; Gunawardhana and Hardin,

2017). Additionally, *vri* overexpression substantially lengthens the circadian period (Blau and

Young, 1999), while an isoform-specific *vri* deletion mutant exhibits a shorter period

(Gunawardhana et al., 2021), indicating VRI's role in regulating the clock speed. An isoform-

specific, arrhythmic *Pdp1ε* mutant exhibits reduced CLK expression (Zheng et al., 2009),

demonstrating that PDP1 also functions in the molecular clock. Although PDP1 overexpression

has little effect on the period length (Zheng et al., 2009), heterozygous *Pdp1* null mutants show

slightly longer periods (Cyran et al., 2003), suggesting that PDP1 also modulates the clock

speed.

We previously reported that TARA interacts with Cyclin A (CycA) and Cyclin-

dependent kinase 1 (Cdk1) to promote sleep in *Drosophila* (Afonso et al., 2015b)*.* In addition,

TARA and its mammalian homologs, Transcriptional Regulator Interacting with PHD-

Bromodomain (TRIP-Br, also known as SERTAD) family of proteins, function in diverse

biological processes, including cell cycle progression, metabolic regulation, and transcriptional

coregulation (Hsu et al., 2001; Calgaro et al., 2002; Liew et al., 2013; Dutta and Li, 2017).

TARA was initially categorized as a Trithorax group (TrxG) member of transcriptional

coactivators (Calgaro et al., 2002). However, a later study found that TARA enhances Polycomb

(Pc)-mediated transcriptional repression (Dutta and Li, 2017), suggesting that TARA may act as

a transcriptional coactivator or corepressor depending on the cellular context and interaction

partners. Here, we report a novel role of TARA in the transcriptional coregulation of VRI and

PDP1 in the molecular clock. Our results suggest that TARA modulates the pace of the

molecular clock and locomotor activity rhythms by physically interacting with and enhancing the

transcriptional activity of VRI and PDP1. Our study has implications for understanding the

mammalian circadian clock since mammalian homologs of VRI and PDP1 also function in the

molecular clock.

Materials and Methods

Fly stocks

Flies were raised on standard food containing molasses, cornmeal, and yeast at 25°C under 12 h-

12 h light-dark (LD) cycle. The following lines were obtained from the Bloomington *Drosophila*

Stock Center: UAS-*tara*-RNAi (JF01421, # 31634), UAS-*dicer2* (*dcr2,* #24650), *tara*[1] (#6403),

UAS-*vri* (#78077), *elav-Gal4* (#458)*, nos-Cas9* (#54591), *tim-Gal4* (#80941), *Pdf-Gal4*

(#80939) and the background control line *iso31* (*w*[1118], #3605). The UAS-*tara*-RNAi (6889R-1,

#NM_080255.2) line was obtained from the National Institute of Genetics, Japan. UAS-*Pdp1*

and *tara*[s132] were previously described (Zheng et al., 2009; Afonso et al., 2015b). UAS-*dcr2* was

combined with two RNAi lines (JF01421 and 6889R-1) to enhance the efficiency of *tara*

knockdown. TARA and its mammalian homologs have four conserved domains: CycA binding

homology (CAB), SERTA, PHD-Bromo binding (PBB), and C-terminal (Hirai et al.). UAS-*tara*

(containing the full-length *tara-B* coding sequence) and *tara* domain deletion (ΔCAB, ΔSERTA,

ΔPBB, and ΔCT) lines were generated in this study. All fly lines were outcrossed with *iso31* for

at least five generations, except for the UAS-*tara*-RNAi and *nos-Cas9* lines.

Circadian assays

To conduct the circadian assays, 3 to 5-day-old flies that had been entrained to an LD cycle at

25°C were placed individually in small glass tubes containing 5% sucrose and 2% agar. The

Drosophila Activity Monitoring System (Trikinetics) was used to record circadian locomotor

activity in constant darkness (DD). We employed the FaasX software (M. Boudinot and F.

Rouyer, Institute of Neurosciences Paris-Saclay, France) to calculate the circadian period and

rhythm strength of each fly. The software analyzes activity data collected in 30-min bins using

χ^2 analysis to determine the period length and rhythm strength. Based on the rhythm strength

value, each fly was classified as arrhythmic (<25), weakly rhythmic (25-50), or rhythmic (>50).

All flies were included in the statistical analysis of rhythm strength, while the analysis of period

length excluded arrhythmic flies.

Immunohistochemistry

Following dissection, adult brains were fixed in 4% paraformaldehyde for 30 min at room

temperature. The samples were blocked in 1% bovine serum albumin (BSA) for PER and PDF

antibody staining. The primary and secondary antibodies were diluted in 0.1% BSA. The primary

antibodies utilized were anti-PER (rabbit 13.1 from the Sehgal lab, University of Pennsylvania)

at a dilution of 1:8000 and mouse anti-PDF (mouse C7s from the Developmental Studies

Hybridoma Bank) at a dilution of 1:2000. The secondary antibodies, Alexa Fluor 647 goat anti-

rabbit and Alexa Fluor 488 goat anti-mouse (Invitrogen) were used at a dilution of 1:1000. Both

primary and secondary antibodies were incubated overnight at 4°C, and images were captured

using a Leica SP8 confocal microscope.

Plasmid constructs for cell-based assays

Actin-Clk, Actin-per, Actin-Gal4, per-enhanced luciferase (*per-luc*) (Darlington et al., 1998), and

pUAST-V5 vector were obtained from the Sehgal lab (University of Pennsylvania). *Clk-luc* (~3

kb *Clk* promoter region fused to a *luciferase* cDNA; Cyran et al., 2003)*, UAS-vri, and* CMV-

vri::VP16 were obtained from the Blau lab (New York University). UAS-*Pdp1* was previously

described (Zheng et al., 2009). CMV-FLAG::*hTRIP-Br1* and CMV-FLAG::*hTRIP-Br2* were a

gift from Dr. Jit Kong Cheong (National University of Singapore). *Actin-Renilla luciferease*

(*rluc*) and CMV-*rluc* (Promega) were used to control for transfection efficiency.

A full-length, untagged UAS-*tara* construct was generated by PCR-amplifying the entire

tara-B coding region from *tara* cDNA and inserting it into the pUAST vector (Brand and

Perrimon, 1993). UAS-HA*::tara* constructs were generated by inserting the *tara-B* coding

region, amplified from the untagged UAS-*tara* construct, into the pTHW Gateway vector

(*Drosophila* Genomics Resource Center) containing the UAST promoter and N-terminal 3xHA

tag. The TARA-B residues removed in the deletion constructs are as follows: ΔCAB:1-26,

ΔSERTA: 432-479, ΔPBB: 722-739, and ΔCT: 893-912. pUAST-attB-V5::*tara* was generated

by inserting the *tara-B* coding region into the pUAST-attB-V5 vector, which is derived from the

pUAST-attB vector (*Drosophila* Genomics Resource Center) for N-terminal V5 tagging. CMV-

tara was generated by digesting UAS-*tara* with BglII and XbaI, and inserting the *tara-B* coding

sequence into the BamHI and XbaI sites in the pCDNA3 vector (Invitrogen). UAS-*Pdp1*::V5

was generated by digesting UAS-*Pdp1* (Zheng et al., 2009) with NotI and XhoI, and inserting the

Pdp1 coding sequence into the corresponding sites in the pUAST-V5 vector, which is derived

from pUAST for C-terminal V5 tagging. UAS-*vri*::MYC was generated by inserting

the *vri* coding sequence, amplified from a UAS-*vri* construct (Cyran et al., 2003), into the

pTWM Gateway vector (*Drosophila* Genomics Resource Center). The constructs were validated

by sequencing (GENEWIZ).

Luciferase-based transcription assay

Drosophila S2 cells were cultured at 25°C in Schneider's Medium (Gibco), while HEK293 cells

were cultured at 37°C in DMEM Medium (Gibco). Both media were supplemented with 10%

heat-inactivated fetal bovine serum and 1% penicillin/streptomycin. Approximately 24 h prior to

transient transfection, 48-well plates were seeded with ~1.5 x 10^5 *Drosophila* S2 or HEK293

cells and maintained at their respective temperatures (25°C or 37°C). Transfections were

performed with Effectene Reagent (Qiagen) per the manufacturer's protocol. Transcriptional

activity was determined using the Dual Luciferase Assay Kit (Promega). Approximately 48 h

after transfection, cells were lysed in 1X Passive Lysis Buffer (PLB), and luciferase activity was

measured using the Luciferase Assay Reagent in Cytation 5 Luminometer (Biotek) following the

manufacturer's instructions.

For the PDP1- and VRI-dependent transcriptional assay, the following constructs were

transfected: *Clk-luc* (10 ng), *Actin-rluc* (10 ng), *Actin-Gal4* (70 ng), UAS-*Pdp1*::V5 (10 ng),

UAS-*vri*::MYC (35 ng), and UAS-HA*::tara* (1 ng, 5 ng, or 15 ng). An empty pUAST::V5 vector

was included to ensure an equal amount of total DNA (150 ng) in all conditions. For the CLK-

dependent transcriptional assays, the following constructs were used: *per-luc* (20 ng), *Actin-rluc*

(5 ng), *Actin-Gal4* (20 ng), *Actin-Clk* (1 ng), UAS-HA::*tara* (10 ng or 30 ng). Empty *pActin* and

pUAST::V5 vectors were used to control for the total DNA amounts (150 ng). For the PER-

dependent transcriptional assay, the following constructs were transfected: *per-luc* (20 ng),

Actin-rluc (15 ng), *Actin-Gal4* (20 ng), *Actin-Clk* (1 ng), *Actin-per* (2 ng, 5 ng, or 15 ng), and

UAS-HA*::tara* (30 ng). Empty *pActin* and pUAST::V5 vectors were used to control for the total

DNA amounts (150 ng). For the VRI::VP16-dependent transcriptional assay, the following

constructs were transfected: *Clk-luc* (10 ng), CMV-*rluc* (10 ng), *Actin-Gal4* (70 ng), CMV-

vri::VP16 (5 ng), CMV-*tara* (10ng or 30ng), CMV-HA::ΔSERTA (30 ng), CMV-

FLAG*::hTRIP-Br1* (30 ng) , CMV-FLAG*::hTRIP-Br2* (30 ng), and the pCDNA3 vector to

control for the total DNA amount (125 ng).

Western analysis and co-immunoprecipitation (co-IP)

To examine the expression levels of transfected proteins, *Drosophila* S2 cell lysates were

collected approximately 48 h after transfection. Cells were lysed in 1X PLB to release

cytoplasmic proteins, while the insoluble fractions were treated with a sample buffer containing

sodium dodecyl sulfate (SDS) to release nucleic proteins. For the co-IP assay, ~3.2×10^6 S2 cells

were seeded in each well of a 6-well plate and transfected ~24 h after seeding. Around 48 h after

transfection, cells were harvested and lysed in 1X RIPA buffer supplemented with Protease

Inhibitor Cocktail (Sigma). The cell lysates were pre-cleaned with Dynabeads Protein G

(Invitrogen) to avoid non-specific binding. The pre-cleaned lysates were incubated with anti-V5-

conjugated magnetic beads (Medical & Biological Laboratories) overnight at 4°C. To elute the

protein complexes bound to the beads, 2X SDS with 5% β-mercaptoethanol was used. Western

blot analysis was performed as described (Wu et al., 2010). Antibodies against TARA (TJR51,

1:1000; Afonso et al., 2015b), HA (Covance, 1:1000), PDP1 (1:1000; Cyran et al., 2003), β-

ACTIN (Abcam, 1:8000), and Histone H3 (Abcam, 1:3000) were used.

Generation of tara mutants with domain deletions and a UAS-tara transgenic line

To identify an optimal protospacer adjacent motif (PAM) sequence within each domain, we

utilized the flyCRISPR target finder (https://flycrispr.org/target-finder). The following PAM

(shown in bold) and their corresponding flanking sequences were identified, with the numbers

indicating the nucleotide positions in the *tara-b* transcript: CAB domain: 552-574 – **CCA** AGC

GGA AGC ATG AGC TGA CC; SERTA domain: 1781-1802 – G CCA ACA GGA GTT GCT

ACA **AGG**; PBB domain: 2672-2680 – TA CTA GCG ATT CGG GTT ATG **CGG**; CT

domain: 3186-3208 – **CCC** GCC TGC ACG ACA ACG AAC TG. To introduce the domain-

specific protospacer sequences into the pCFD4 vector (Addgene), we employed the ligation-

independent Gibson Assembly method (NEBuilder, New England BioLabs) according to the

manufacturer's protocol. Transgenic flies carrying domain-specific gRNAs were generated using

the standard embryo injection technique (Rainbow Transgenics). Domain-specific deletions were

generated by crossing fly lines carrying the domain specific gRNAs with a *nos-Cas9* fly line.

Mutant flies were identified by sequencing the genomic DNA (GENEWIZ). Additionally, a

transgenic fly line carrying the untagged UAS-*tara* construct was generated using the standard

embryo injection technique (Rainbow Transgenics).

RNA extraction and quantitative real-time PCR (qPCR)

Total RNA was extracted from 30 male fly heads using 1 mL of TRIzol Reagent (Ambion). We

employed 100 µL of 1-Bromo-3-Chloropropane (Molecular Research Center, Inc.) to separate

the phases, and RNA precipitation was achieved by adding 500 µL of 2-propanol. After

treatment with 2 µL of DNaseI, cDNA was amplified from 1 µg of RNA using a high-capacity

cDNA Reverse Transcription Kit (Applied Biosystems). For qPCR, we used SYBR Green PCR

Master Mix (Applied Biosystems) and the 7500 Real-Time PCR System (Applied Biosystems). 3

technical replicates of 6 independent biological samples were assayed for each genotype. The

following primers were used for *Pdf*: 5'-CCT TGT GCT TCT GGC CAT TT-3' and 5'-GTC

GAG GAG ATC CCG ATT GT-3'. *actin* mRNA was used for normalization.

Experimental design and statistical analyses

Throughout the study, we employed a between-subject design. The experimental design is

visually depicted in the figures and described in the corresponding figure legends. Behavioral

experiments were repeated at least twice using flies from independent crosses, while cell-based

experiments were conducted at least three times. Statistical analyses were performed using

GraphPad Prism 9. For datasets with two groups, Welch's *t*-tests were utilized for comparisons.

To compare multiple groups, one-way Brown-Forsythe and Welch ANOVA tests were

conducted, followed by Dunnett's T3 multiple comparisons tests. For experiments involving two

factors, two-way ANOVA tests were employed, followed by Tukey's post hoc tests or Sidak's

multiple comparisons tests.

Results

TARA functions in clock neurons to promote rhythmic behavior

We previously showed that *tara* mutants display arrhythmic locomotor behavior as well

as reduced sleep (Afonso et al., 2015b). We confirmed that the reduced rhythmicity maps to the

tara locus using RNAi-mediated *tara* knockdown in all neurons using *elav-Gal4* (Table 1, Fig. 1

A,B). The *Drosophila* brain contains ~150 circadian neurons in several clusters. A cluster of

clock neurons expressing the Pigment Dispensing Factor (PDF) neuropeptide, called small

ventral lateral neurons (s-LN$_v$s), is essential for free-running rhythms in DD (Renn et al., 1999).

To examine where TARA functions to promote rhythmic behavior, we employed *tim-Gal4*,

which targets all clock neurons, and *Pdf-Gal4*, which targets the PDF-expressing LN$_v$s. Male

flies are typically used for assaying locomotor rhythms because they display more robust and

consistent rhythms than females. However, because pan-neuronal *tara* knockdown led to

lethality in males, female flies were used in the pan-neuronal knockdown experiment. Both

males and females were assayed in experiments using *tim-Gal4* and *Pdf-Gal4*.

In both males and females, knockdown of *tara* in all clock neurons using *tim-Gal4*

resulted in a significant reduction of rhythm strength (Table 1, Fig. 1). In contrast, restricted *tara*

knockdown in PDF-expressing LN$_v$s did not significantly affect circadian behavior. This finding

suggests that TARA functions in multiple clock neuron clusters to regulate rhythmic behavior.

Comparing female data with *elav-Gal4* vs. *tim-Gal4*, we observed that pan-neuronal knockdown

of *tara* had a more substantial effect on rhythm strength than knockdown in clock neurons (Table

1, Fig 1A,B), suggesting TARA also functions in non-clock neurons to promote rhythmic

locomotor activity. The period length remained largely unaffected by *tara* knockdown (Table 1,

Fig. 1). Our results suggest that TARA promotes rhythmic locomotion in clock and non-clock

neurons.

TARA modulates the speed of the circadian clock

To investigate whether TARA influences molecular clock oscillation, we examined PER

protein expression in s-LN$_v$s in DD. Although over 80% of *tara*$^{s132/1}$ mutants are arrhythmic or

weakly rhythmic (Afonso et al., 2015b), they displayed robust daily cycling of PER levels in DD

(Fig. 2A,B), suggesting that TARA plays a role in the circadian output pathway. Interestingly,

the pattern of PER cycling suggests that TARA also contributes to setting the pace of the

molecular clock. Compared to control brains, *tara* mutant brains exhibited a significantly lower

PER signal at CT 3 and a substantially higher PER signal at CT 15 (Fig. 2A,B), consistent with a

faster molecular clock. We previously reported that *tara*$^{s132/1}$ mutants, which have the most

severe reduction in *tara* levels among viable *tara* mutants, did not exhibit shorter periods in

locomotor behavior (Afonso et al., 2015b). This may be because the more strongly affected

mutants were arrhythmic, leaving only the less severely affected mutants for period length

measurements. In addition to the PER cycling changes, we observed that PDF levels at the dorsal

projections of the s-LN$_v$s were considerably lower in *tara* mutants than in control flies (Fig.

2C,D). We also found a significant reduction in the total levels of *Pdf* mRNA in *tara* mutants

compared to the controls (Fig. 2E). However, the reduction in PDF protein was more pronounced

than that of the *Pdf* transcripts, suggesting that TARA impacts PDF levels at both the

transcriptional and post-transcriptional levels. *Pdf* null mutants display decreased rhythmicity

and shorter periods (Renn et al., 1999), suggesting that reduced PDF levels in *tara* mutants

contribute to their behavioral and molecular phenotypes.

To further examine the effect of TARA levels on the speed of the circadian clock, we

examined whether flies overexpressing TARA exhibit the opposite pattern of PER cycling as

tara mutants. Indeed, TARA overexpression in PDF-expressing (PDF+) neurons led to a

significantly lower PER signal at CT 21, consistent with a slower molecular clock (Fig. 3A,B).

Furthermore, flies overexpressing TARA in PDF+ neurons exhibited a significantly longer

circadian period than control flies (Table 1, Fig. 3 C-E). Collectively, our data suggest that *tara*

plays a role in setting the pace of the molecular clock and regulating the circadian output

pathway.

Evolutionarily conserved domains in TARA contribute to locomotor rhythmicity

TARA and its mammalian homologs have four conserved domains: CycA binding

homology (CAB), SERTA, PHD-Bromo binding (PBB), and C-terminal (Hirai et al.). To

identify the domains essential for circadian regulation, we utilized the CRISPR/Cas9 technique

to generate mutants in which the conserved domains were removed from the *tara* genomic locus

(Fig. 4A). Deletion of the SERTA domain led to lethality. In contrast, flies in which the CAB,

PBB, or CT domain was deleted were viable. We assayed SERTA deletion mutants in trans to

the hypomorphic $tara^{s132}$ allele and homozygous mutants for all other deletions for circadian

locomotor behavior. Compared with s132/+ heterozygous flies, the s132/ΔSERTA trans-

heterozygous mutants exhibited a substantial reduction in rhythm strength, but their circadian

period was not significantly altered (Table 2, Fig. 4 B-D). However, given that the s132 allele is

hypomorphic, and s132/ΔSERTA trans-heterozygotes retain some wild-type TARA protein, we

cannot rule out the possibility that the SERTA domain plays a role in determining the period

length. The lack of period change in s132/ΔSERTA mutants could be because the more strongly

affected mutants were arrhythmic and therefore excluded from the period analysis. Homozygous

CAB, PBB, or CT domain deletions had little effect on the period length (Table 2, Fig. 4B,C),

but they significantly reduced the strength of the locomotor rhythms (Table 2, Fig. 4B,D),

suggesting that they also contribute to rhythmic locomotor behavior. Our results indicate that all

four conserved domains play a role in maintaining rhythm strength.

TARA interacts with VRI and PDP1 to modulate *Clk* transcription

Considering the role mammalian homologs of TARA play in transcriptional coregulation

(Hsu et al., 2001), our discovery that TARA modulates the period length suggests that TARA

acts as a transcriptional coregulator of core clock components. We hypothesized that TARA

influences VRI activity, since *tara* and *vri* mutants exhibit similar circadian phenotypes.

Overexpression of *tara* or *vri* in PDF-expressing neurons leads to lengthened period (Table 1,

Fig. 3 C-E; Blau and Young, 1999), and both *tara* and *vri* mutants have decreased PDF

expression in s-LN$_v$s dorsal projections (Fig. 2C,D; Gunawardhana and Hardin, 2017).

Consequently, we investigated whether TARA regulates *Clk* transcription repression by VRI. For

this purpose, we employed *Drosophila* S2 cells transfected with a *Clk-luc* reporter that contains a

D-box element, a common binding site for PDP1 and VRI (Cyran et al., 2003). Previous work

has shown that VRI binds the same D-box element of the *Clk* promoter as PDP1, acting as a

competitive repressor (Cyran et al., 2003). Since repression by VRI would not be evident without

Clk transcription activation by PDP1, we co-transfected PDP1 and VRI at varying levels of

TARA.

A prior study used mammalian HEK293 cells to demonstrate PDP1-dependent *Clk*

transcription activation because it was not detectable in S2 cells (Cyran et al., 2003).

Intriguingly, we observed PDP1-dependent *Clk* transcription in *Drosophila* S2 cells when TARA

was co-transfected with PDP1 (Fig. 5A). Little PDP1 activity was observed without co-

transfected TARA, while PDP1 activity increased up to ~5-fold with rising TARA levels.

Transfection of TARA alone did not influence *Clk* transcription (Fig. 5A), in line with its lack of

a DNA binding domain and its function as a transcriptional coregulator rather than a

transcription factor. VRI co-transfection repressed PDP1-dependent *Clk* transcription, verifying

VRI's role as a competitive repressor of PDP1 (Fig. 5A). The degree of VRI-mediated

suppression of *Clk* transcription intensified with increasing levels of TARA. Intermediate TARA

levels, which significantly affected PDP1 activity, had little to moderate impact on VRI activity,

but higher TARA levels resulted in robust suppression of PDP1 activity by VRI (Fig. 5A).

However, since VRI suppressed nearly all PDP1-mediated *Clk* transcription at every level of

TARA, the smaller VRI-dependent suppression at lower levels of TARA may be attributed to

lower PDP1 activity rather than lower VRI activity.

To investigate the impact of TARA on VRI activity directly without co-transfected

PDP1, we utilized VRI fused to the viral VP16 transactivation domain (VRI::VP16), which

converts the repressor into a transcriptional activator (Cyran et al., 2003). We were unable to

express detectable amounts of VRI::VP16 in *Drosophila* S2 cells, possibly due to lethality

caused by its expression. Therefore, we turned to HEK293 cells, as a previous study

demonstrated *Clk* transcription by VRI::VP16 using the cell line (Cyran et al., 2003). Consistent

with the previous finding, VRI::VP16 promoted *Clk* transcription without co-transfected TARA

(Fig. 5B). Importantly, we discovered that co-transfected TARA upregulated VRI::VP16-

dependent *Clk* transcription up to ~4 fold in a dose-dependent manner, demonstrating TARA's

role in enhancing VRI activity. Moreover, TRIP-Br1 and TRIP-Br2, two mammalian homologs

of TARA, exhibited a moderate effect on VRI::VP16-dependent *Clk* transcription as well,

suggesting a conserved mechanism (Fig. 5B).

To evaluate whether TARA impacts *Clk* transcription by influencing PDP1 and VRI

protein levels, we conducted a Western analysis using cell lysates following transfection in S2

cells. Since transcriptional activity occurs in the nucleus, we examined nuclear fractions. Co-

transfected TARA did not influence PDP1 or VRI levels (Fig. 5C-G), suggesting that TARA

enhances the transcriptional activity of PDP1 and VRI, not their expression or protein stability.

Although co-transfected VRI increased TARA and PDP1 protein levels (Fig. 5C-G), these

effects would not account for the observed effects of TARA on *Clk* transcription. Our data

support TARA's function as a transcriptional coregulator of VRI and PDP1.

We then inquired whether the effects of TARA on VRI and PDP1 were mediated by

physical interaction. Co-immunoprecipitation (Co-IP) experiments in S2 cells demonstrated that

TARA can indeed form a physical complex with VRI or PDP1 (Fig. 5H). Collectively, our

results indicate that TARA can physically interact with VRI and PDP1 to enhance the

transcriptional suppression of *Clk* by VRI and its activation by PDP1.

TARA exhibits little impact on CLK and PER activity

Our findings indicate that TARA plays a role in the PDP1/VRI feedback loop of the

Drosophila molecular clock. To ascertain whether TARA also modulates the CLK-CYC/PER-

TIM feedback loop, we utilized a *per-luc* reporter in S2 cells and co-transfected CLK with

TARA. CLK-mediated transcription of *per* in S2 cells did not require co-transfected TARA, in

agreement with previous findings (Darlington et al., 1998), and was unaffected by increasing

TARA levels (Fig. 5I).

To investigate whether TARA regulates the suppression of CLK activity by PER, we co-

transfected CLK with varying levels of PER in S2 cells, with or without TARA. As previously

reported **(**Darlington et al., 1998**),** PER repressed CLK-dependent transcription of *per* in a dose-

dependent manner (Fig. 5J). Notably, the effect of TARA at each level of PER and the

interaction between the effects of TARA and PER were not significant, indicating that PER

suppression is independent of TARA (Fig. 5J). Together, our results reveal that TARA exerts

minimal impact on CLK's transcriptional activity and PER's repressive activity, demonstrating

the specificity in TARA's function as a transcriptional coregulator.

The SERTA domain is essential for transcriptional coregulation

To investigate the role of TARA's conserved domains in transcriptional regulation, we

generated deletion constructs for each domain: ΔCAB, ΔSERTA, ΔPBB, and ΔCT (Fig. 6A). We

transfected full-length or domain-deletion TARA along with *Clk-luc* and PDP1 in S2 cells and

conducted Luciferase assays. The results showed that TARA lacking the SERTA domain

(ΔSERTA) exerted a markedly diminished effect on PDP1 activity compared to full-length

TARA (Fig. 6B), indicating that the SERTA domain is critical for TARA to enhance PDP1's

transcriptional activity. Deletion of the other conserved domains had little effect on PDP1-

dependent *Clk* transcription. Next, we examined whether the SERTA domain is essential for

TARA's enhancement of VRI transcriptional activity using VRI::VP16 and HEK293 cells. We

discovered that the SERTA domain deletion eliminated TARA's influence on VRI::VP16

activity (Fig. 6C). To determine the expression levels of deletion constructs and their effects on

PDP1 expression, we performed a Western analysis. The results demonstrated that the deletion

constructs were expressed at levels similar to or slightly higher than the wild-type TARA (Figure

6D,E). Furthermore, PDP1 expression was not significantly affected by deletion constructs (Fig.

6D,F), indicating that the reduced *Clk* transcription upon ΔSERTA transfection is not due to

diminished PDP1 expression. Taken together, our findings demonstrate that the SERTA domain

of TARA is essential for enhancing PDP1 and VRI activity.

tara* genetically interacts with *vri* and *Pdp1* to modulate the molecular clock *in vivo

Since we discovered that TARA enhanced the transcriptional activity of VRI and PDP1

in cultured cells, we examined whether *tara* interacts with *vri* and *Pdp1* to modulate the

molecular clock *in vivo*. We overexpressed *vri* or *Pdp1* in PDF+ clock neurons, with or without

tara co-overexpression. We observed a pronounced increase in period length when *vri* was

overexpressed in LN$_v$s (Table 3, Fig. 7), consistent with previous findings (Blau and Young,

1999). Although overexpressing *tara* alone also significantly increased period length, co-

overexpressing both *tara* and *vri* shortened the period compared to overexpressing *vri* alone

(Table 3, Fig. 7), indicating a genetic interaction between *tara* and *vri*. TARA may lengthen the

circadian period by enhancing VRI activity when VRI is present at normal levels. However,

when VRI levels are elevated, additional TARA may not further amplify VRI's period-

lengthening effect, but instead, it may shorten the period through its effect on PDP1.

We observed a minor (~0.3 h) increase in period length when *Pdp1* was overexpressed

(Table 3, Fig. 7). This outcome may seem unexpected, given PDP1's role as a positive activator

of *Clk* transcription. However, the small effect is comparable to the previously reported ~0.2 h,

non-significant increase in the period length of *Pdp1*-overexpressing flies (Zheng et al., 2009).

Although overexpressing *tara* in PDF+ neurons led to a significantly longer period, co-

overexpression with *Pdp1* eliminated TARA's impact on the period length (Table 3, Fig. 7),

demonstrating that *tara* and *Pdp1* genetically interact to regulate the pace of the molecular clock.

Overexpression of *tara* alone had a small but significant effect on the robustness of

circadian rhythms, while overexpression of *vri* alone or in combination with *tara* had little effect

(Table 3, Fig. 7). Overexpression of *Pdp1* significantly weakened the strength of locomotor

rhythms, consistent with previous findings (Benito et al., 2007; Zheng et al., 2009). Co-

overexpression of *tara* and *Pdp1* further decreased the rhythm strength, but the decrease was not

statistically significant.

The effect of *tara* overexpression on the circadian period may appear more similar to that

of *vri* overexpression than *Pdp1* overexpression, as both *tara* and *vri* overexpression markedly

lengthen the period while *Pdp1* overexpression has only a minor effect. However, it is important

to note that this observation does not necessarily indicate that TARA has a greater impact on the

transcriptional activity of VRI compared to PDP1. Considering that TARA enhanced VRI and

PDP1 activity in S2 cells, the impact of *tara* overexpression on locomotor rhythms *in vivo* may

reflect the combined effects of enhancing both VRI and PDP1 activity. Indeed, when we co-

overexpressed *vri* and *Pdp1* in PDF+ clock neurons, we observed an increase in period length

similar to that of *tara* overexpression (Table 3, Fig. 7). Co-overexpressing *vri* and *Pdp1* also

resulted in a rhythm strength comparable to that of overexpressing *tara*. Collectively, our

findings indicate that *tara* interacts with *vri* and *Pdp1* to modulate the circadian period and

rhythm strength in *Drosophila*.

Discussion

The present study identifies TARA as a novel regulatory component of the interlocked

transcription-translation feedback loops in the *Drosophila* molecular clock. Whereas the primary

feedback loop involving CLK-CYC and PER-TIM in the *Drosophila* circadian clock has been

studied extensively (reviewed in Allada and Chung, 2010; Hardin, 2011; Tataroglu and Emery,

2015; Patke et al., 2020), the secondary loop, including VRI and PDP1, is relatively understudied

(Blau and Young, 1999; Cyran et al., 2003; Glossop et al., 2003; Benito et al., 2007; Zheng et al.,

2009; Ling et al., 2012; Gunawardhana and Hardin, 2017; Gunawardhana et al., 2021). Our

transcriptional assays reveal that TARA enhances the activity of both the negative and positive

arms of the secondary feedback loop. TARA has little effect on PER and CLK activity,

suggesting that TARA plays a more critical role in the secondary loop than the primary loop.

Consistent with the transcriptional assay results, our behavioral data demonstrate that *tara*

genetically interacts with *vri* and *Pdp1* to regulate the pace of the *Drosophila* circadian clock.

tara mutants display markedly impaired locomotor rhythmicity despite robust PER cycling,

suggesting that TARA plays an additional role in the circadian output pathway. These findings

are reminiscent of previous results showing that VRI and PDP1 play critical roles in the

molecular clock and the circadian output pathway (Blau and Young, 1999; Cyran et al., 2003;

Glossop et al., 2003; Benito et al., 2007; Zheng et al., 2009; Ling et al., 2012; Gunawardhana

and Hardin, 2017; Gunawardhana et al., 2021), supporting the view that TARA functions with

VRI and PDP1 to regulate circadian rhythms. We previously reported that TARA promotes sleep

through its interaction with CycA and Cdk1 in a subset of neurons in the *pars lateralis*, a region

considered analogous to the mammalian hypothalamus (Afonso et al., 2015b). Collectively, these

findings suggest that TARA regulates circadian rhythms and sleep by interacting with different

molecular partners in distinct neuronal populations.

Given that TARA enhances both the activator and suppressor of *Clk* transcription, *tara*

overexpression might be expected to have little effect on the circadian period. However, *vri*

overexpression leads to a markedly long period, while *Pdp1* overexpression has a slight period-

lengthening effect. Therefore, the combined effect of enhancing both VRI and PDP1 is expected

to be a longer period compared to control flies. Indeed, we find that *tara* overexpression

lengthens the circadian period, mirroring the consequence of simultaneously overexpressing *vri*

and *Pdp1*. It is unclear why *vri* overexpression affects the clock speed more than *Pdp1*

overexpression. A previous study demonstrated that VRI expression peaks a few hours before

PDP1 through an as-yet unidentified mechanism (Cyran et al., 2003). The earlier accumulation

of VRI may be linked to a more potent effect of *vri* overexpression on the circadian period.

Considering that PDP1 is an activator of *Clk* transcription and *Clk* overexpression shortens the

circadian period (Zhao et al., 2003), it is unclear why *Pdp1* overexpression leads to a slightly

lengthened period. PDP1 likely has multiple target genes, and the combined effects of PDP1

overexpression on *Clk* and other target genes may be a small increase in the circadian period.

The components of the primary feedback loop in the molecular clock are highly

conserved between flies and mammals (reviewed in Allada and Chung, 2010; Hardin, 2011;

Tataroglu and Emery, 2015; Patke et al., 2020). Mammalian CLK, BMAL1 (a CYC homolog),

and PER1/2/3 perform similar functions as their *Drosophila* orthologs, while

CRYPTOCHROME (CRY)1/2 are functionally equivalent to *Drosophila* TIM. However, the

functions of the secondary feedback loop involving VRI and PDP1 in *Drosophila* are performed

by more complex, multi-layered feedback loops in mammals. The layered loops include nuclear

receptors, REV-ERBα/β (also known as NR1D1/2), and Retinoid-related Orphan Receptor

(ROR)α/β/δ, which competitively bind the RRE element in the promotor of RRE-containing

genes, including *BMAL1*, to activate and suppress its transcription (Preitner et al., 2002; Ueda et

al., 2002; Sato et al., 2004). REV-ERBs and RORs are not homologs of VRI and PDP1, but

mammalian homologs of VRI and PDP1, named Nuclear Factor Interleukin-3-Regulated Protein

(NFIL3, also known as E4BP4) and D-BOX BINDING PROTEIN (DBP), respectively, also

participate in the interlocked feedback loops. Like their *Drosophila* homologs, NFIL3 and DBP

competitively bind the D-box element to inhibit and activate the transcription of D-box-

containing genes, including RORs (Mitsui et al., 2001). RORs activate *NFIL3* transcription,

while NFIL3 inhibits the transcription of *ROR*s, forming a feedback loop (Mitsui et al., 2001;

Gachon et al., 2004; Ueda et al., 2005). Taken together, REV-ERBs and NFIL3 suppress *BMAL1*

transcription, while RORs and DBP activate *BMAL1* transcription. These findings suggest that

VRI performs the combined functions of REV-ERBs and NFIL3, while PDP1 performs the

combined functions of RORs and DBP. Initial studies demonstrating that VRI and PDP1 regulate

Clk transcription were performed in mammalian HEK293 cells because PDP1 did not activate

Clk transcription in S2 cells (Cyran et al., 2003). Our data show that TARA is the missing factor

necessary for *Clk* transcription by PDP1 in *Drosophila* S2 cells. HEK293 cells may express

sufficient levels of TRIP-Br proteins, the mammalian homologs of TARA, to allow PDP1

activity at the *Clk* promoter. These considerations and our finding that TRIP-Br1/2 enhance VRI

activity in HEK293 cells raise the intriguing possibility that TRIP-Br proteins regulate the

mammalian circadian period through their interactions with NFIL3 and DBP for the transcription

of D-box-containing genes.

TARA and TRIP-Br proteins contain four conserved domains. The function of the

SERTA domain is not well understood (Calgaro et al., 2002; Manansala et al., 2013). The

present study uncovers a novel role of the SERTA domain in enhancing PDP1 and VRI activity

and regulating the pace of the *Drosophila* circadian clock. We also find that TARA is a critical

regulator of rhythm amplitude and that all four domains contribute to robust locomotor rhythms.

It would be interesting to investigate further the molecular functions of the conserved domains

and their role in circadian rhythms.

TARA is ubiquitously expressed in neurons (Afonso et al., 2015b), but what regulates

TARA expression or activity is largely unknown. Notably, chromatin immunoprecipitation tiling

array assays (Abruzzi et al., 2011) detected cyclic CLK binding to a promoter region of *tara*,

suggesting that it is a direct transcriptional target of CLK. Although we did not observe daily

cycling of TARA protein levels in s-LN$_v$s (Afonso et al., 2015a), TARA activity may exhibit

circadian cycling, since its partners VRI and PDP1 undergo circadian cycling. TARA levels or

activity may also respond to changing environmental and internal conditions, such as the

metabolic state. Earlier findings suggest that mammalian homologs of TARA are involved in

metabolic regulation. TRIP-Br2 is involved in lipid and oxidative metabolism in adult mice

(Liew et al., 2013), and prolonged nutrient starvation dramatically decreases TRIP-Br3 to induce

apoptosis (Li et al., 2015). Circadian regulation is closely linked to metabolism (Bailey et al.,

2014; Asher and Sassone-Corsi, 2015; Guan and Lazar, 2021), and TARA may play a role in

integrating circadian and metabolic regulation. Further studies will need to be performed to

examine the potential role of TARA in linking circadian rhythms and metabolism.

Our findings reveal a novel role for *tara* in regulating the pace of the circadian clock

through its interaction with *vri* and *Pdp1* in the secondary feedback loop of the *Drosophila*

circadian clock. They provide a rationale for investigating the role of the TRIP-Br proteins in

regulating circadian rhythms and their potential interaction with NFIL3 and DBP in the

mammalian molecular clock.

References

Abruzzi KC, Rodriguez J, Menet JS, Desrochers J, Zadina A, Luo W, Tkachev S, Rosbash M

(2011) Drosophila CLOCK target gene characterization: implications for circadian tissue-

specific gene expression. Genes Dev 25:2374-2386.

Afonso DJ, Machado DR, Koh K (2015a) Control of sleep by a network of cell cycle genes. Fly

(Austin) 9:165-172.

Afonso DJ, Liu D, Machado DR, Pan H, Jepson JE, Rogulja D, Koh K (2015b) TARANIS

Functions with Cyclin A and Cdk1 in a Novel Arousal Center to Control Sleep in

Drosophila. Curr Biol 25:1717-1726.

Allada R, Chung BY (2010) Circadian organization of behavior and physiology in Drosophila.

Annu Rev Physiol 72:605-624.

Allada R, White NE, So WV, Hall JC, Rosbash M (1998) A mutant Drosophila homolog of

mammalian Clock disrupts circadian rhythms and transcription of period and timeless.

Cell 93:791-804.

Asher G, Sassone-Corsi P (2015) Time for food: the intimate interplay between nutrition,

metabolism, and the circadian clock. Cell 161:84-92.

Bailey SM, Udoh US, Young ME (2014) Circadian regulation of metabolism. J Endocrinol

222:R75-96.

Benito J, Zheng H, Hardin PE (2007) PDP1epsilon functions downstream of the circadian

oscillator to mediate behavioral rhythms. J Neurosci 27:2539-2547.

Blau J, Young MW (1999) Cycling vrille expression is required for a functional Drosophila

clock. Cell 99:661-671.

Brand AH, Perrimon N (1993) Targeted gene expression as a means of altering cell fates and

generating dominant phenotypes. Development 118:401-415.

Calgaro S, Boube M, Cribbs DL, Bourbon HM (2002) The Drosophila gene taranis encodes a

novel trithorax group member potentially linked to the cell cycle regulatory apparatus.

Genetics 160:547-560.

Cyran SA, Buchsbaum AM, Reddy KL, Lin MC, Glossop NR, Hardin PE, Young MW, Storti

RV, Blau J (2003) vrille, Pdp1, and dClock form a second feedback loop in the

Drosophila circadian clock. Cell 112:329-341.

Darlington TK, Wager-Smith K, Ceriani MF, Staknis D, Gekakis N, Steeves TD, Weitz CJ,

Takahashi JS, Kay SA (1998) Closing the circadian loop: CLOCK-induced transcription

of its own inhibitors per and tim. Science 280:1599-1603.

Dutta P, Li WX (2017) The SERTAD protein Taranis plays a role in Polycomb-mediated gene

repression. PLoS One 12:e0180026.

Gachon F, Fonjallaz P, Damiola F, Gos P, Kodama T, Zakany J, Duboule D, Petit B, Tafti M,

Schibler U (2004) The loss of circadian PAR bZip transcription factors results in

epilepsy. Genes Dev 18:1397-1412.

George H, Terracol R (1997) The vrille gene of Drosophila is a maternal enhancer of

decapentaplegic and encodes a new member of the bZIP family of transcription factors.

Genetics 146:1345-1363.

Glossop NR, Houl JH, Zheng H, Ng FS, Dudek SM, Hardin PE (2003) VRILLE feeds back to

control circadian transcription of Clock in the Drosophila circadian oscillator. Neuron

37:249-261.

Guan D, Lazar MA (2021) Interconnections between circadian clocks and metabolism. J Clin

Invest 131.

Gunawardhana KL, Hardin PE (2017) VRILLE Controls PDF Neuropeptide Accumulation and

Arborization Rhythms in Small Ventrolateral Neurons to Drive Rhythmic Behavior in

Drosophila. Curr Biol 27:3442-3453 e3444.

Gunawardhana KL, Rivas GBS, Caster C, Hardin PE (2021) Crosstalk between vrille transcripts,

proteins, and regulatory elements controlling circadian rhythms and development in

Drosophila. iScience 24:101893.

Hardin PE (2011) Molecular genetic analysis of circadian timekeeping in Drosophila. Adv Genet

74:141-173.

Hardin PE, Hall JC, Rosbash M (1990) Feedback of the Drosophila period gene product on

circadian cycling of its messenger RNA levels. Nature 343:536-540.

Hirai T, Ramos TL, Lin P-Y, Simonetta F, Su LL, Picton LK, Baker J, Lin J-X, Li P, Seo K,

Lohmeyer JK, Bolivar-Wagers S, Mavers M, Leonard WJ, Blazar BR, Garcia KC, Negrin

RS (2021) Selective expansion of regulatory T cells using an orthogonal IL-2/IL-2

receptor system facilitates transplantation tolerance. The Journal of clinical investigation

131.

Hsu SI, Yang CM, Sim KG, Hentschel DM, O'Leary E, Bonventre JV (2001) TRIP-Br: a novel

family of PHD zinc finger- and bromodomain-interacting proteins that regulate the

transcriptional activity of E2F-1/DP-1. EMBO J 20:2273-2285.

Li C, Jung S, Lee S, Jeong D, Yang Y, Kim KI, Lim JS, Cheon CI, Kim C, Kang YS, Lee MS

(2015) Nutrient/serum starvation derived TRIP-Br3 down-regulation accelerates

apoptosis by destabilizing XIAP. Oncotarget 6:7522-7535.

Liew CW, Boucher J, Cheong JK, Vernochet C, Koh HJ, Mallol C, Townsend K, Langin D,

612   Kawamori D, Hu J, Tseng YH, Hellerstein MK, Farmer SR, Goodyear L, Doria A,

613   Bluher M, Hsu SI, Kulkarni RN (2013) Ablation of TRIP-Br2, a regulator of fat lipolysis,

614   thermogenesis and oxidative metabolism, prevents diet-induced obesity and insulin

615   resistance. Nat Med 19:217-226.

616   Ling J, Dubruille R, Emery P (2012) KAYAK-alpha modulates circadian transcriptional

617   feedback loops in Drosophila pacemaker neurons. J Neurosci 32:16959-16970.

618   Manansala MC, Min S, Cleary MD (2013) The Drosophila SERTAD protein Taranis determines

619   lineage-specific neural progenitor proliferation patterns. Dev Biol 376:150-162.

620   Mitsui S, Yamaguchi S, Matsuo T, Ishida Y, Okamura H (2001) Antagonistic role of E4BP4 and

621   PAR proteins in the circadian oscillatory mechanism. Genes Dev 15:995-1006.

622   Partch CL, Green CB, Takahashi JS (2014) Molecular architecture of the mammalian circadian

623   clock. Trends Cell Biol 24:90-99.

624   Patke A, Young MW, Axelrod S (2020) Molecular mechanisms and physiological importance of

625   circadian rhythms. Nat Rev Mol Cell Biol 21:67-84.

626   Preitner N, Damiola F, Lopez-Molina L, Zakany J, Duboule D, Albrecht U, Schibler U (2002)

627   The orphan nuclear receptor REV-ERBalpha controls circadian transcription within the

628   positive limb of the mammalian circadian oscillator. Cell 110:251-260.

629   Reddy KL, Wohlwill A, Dzitoeva S, Lin MH, Holbrook S, Storti RV (2000) The Drosophila

630   PAR domain protein 1 (Pdp1) gene encodes multiple differentially expressed mRNAs

631   and proteins through the use of multiple enhancers and promoters. Dev Biol 224:401-414.

Renn SC, Park JH, Rosbash M, Hall JC, Taghert PH (1999) A pdf neuropeptide gene mutation

and ablation of PDF neurons each cause severe abnormalities of behavioral circadian

rhythms in Drosophila. Cell 99:791-802.

Sato TK, Panda S, Miraglia LJ, Reyes TM, Rudic RD, McNamara P, Naik KA, FitzGerald GA,

Kay SA, Hogenesch JB (2004) A functional genomics strategy reveals Rora as a

component of the mammalian circadian clock. Neuron 43:527-537.

Takahashi JS (2017) Transcriptional architecture of the mammalian circadian clock. Nat Rev

Genet 18:164-179.

Tataroglu O, Emery P (2015) The molecular ticks of the Drosophila circadian clock. Curr Opin

Insect Sci 7:51-57.

Ueda HR, Hayashi S, Chen W, Sano M, Machida M, Shigeyoshi Y, Iino M, Hashimoto S (2005)

System-level identification of transcriptional circuits underlying mammalian circadian

clocks. Nat Genet 37:187-192.

Ueda HR, Chen W, Adachi A, Wakamatsu H, Hayashi S, Takasugi T, Nagano M, Nakahama K,

Suzuki Y, Sugano S, Iino M, Shigeyoshi Y, Hashimoto S (2002) A transcription factor

response element for gene expression during circadian night. Nature 418:534-539.

Wu MN, Joiner WJ, Dean T, Yue Z, Smith CJ, Chen D, Hoshi T, Sehgal A, Koh K (2010)

SLEEPLESS, a Ly-6/neurotoxin family member, regulates the levels, localization and

activity of Shaker. Nat Neurosci 13:69-75.

Zhao J, Kilman VL, Keegan KP, Peng Y, Emery P, Rosbash M, Allada R (2003) Drosophila

clock can generate ectopic circadian clocks. Cell 113:755-766.

Zheng X, Sehgal A (2012) Speed control: cogs and gears that drive the circadian clock. Trends

Neurosci 35:574-585.

Zheng X, Koh K, Sowcik M, Smith CJ, Chen D, Wu MN, Sehgal A (2009) An isoform-specific

mutant reveals a role of PDP1 epsilon in the circadian oscillator. J Neurosci 29:10920-

10927.

Figure Legends

Figure 1. Pan-neuronal and clock-neuron-specific knockdown of *tara* degrades rhythm

strength. (A) Representative circadian actograms of females of the following genotypes in DD:

pan-neuronal knockdown (*elav > tara* RNAi), clock-neuron-specific knockdown (*tim > tara*

RNAi), LN$_v$-specific knockdown (*Pdf > tara* RNAi), and their respective *Gal4* driver controls

(*elav > +*, *tim > +*, and *Pdf > +*). In this figure and subsequent figures, the gray and black bars

above the actogram indicate subjective day and night, respectively. **(B)** Period length (top) and

rhythm strength (bottom) of females of indicated genotypes. N = 29-61 for period length; n = 29-

71 for rhythm strength. Red labels indicate female flies, while blue labels indicate male flies. The

horizontal dotted line represents the mean period length (top) and rhythm strength (bottom) of

the common control (*+ > tara* RNAi). **(C)** Representative circadian actograms of males of

indicated genotypes in DD. **(D)** Period length (top) and rhythm strength (bottom) of males of

indicated genotypes. N = 31-47 for period length and rhythm strength. Mean ± SEM is shown.

****$p < 0.0001$, ns: not significant; Brown-Forsythe and Welch ANOVA tests followed by

Dunnett's T3 multiple comparisons test relative to both parental controls (B and D).

Figure 2. PER cycling and PDF accumulation are altered in the s-LN$_v$ of *tara* mutants. (A)

Representative expression patterns of PER and PDF in the s-LN$_v$s of control and *tara*[s132/1] males

at the indicated circadian times on the 3rd day in DD. The scale bar is 10 μm. **(B)** Quantification

of PER levels in s-LN$_v$s for control and *tara*[s132/1] males (n = 11-19). **(C)** Representative PDF

expression patterns in the dorsal projections of s-LN$_v$s at the indicated circadian times on the 3rd

day in DD in control and *tara*[s132/1] males. Signal intensity is indicated using the "royal" lookup

table, where black represents the minimum intensity, and white represents the maximum

intensity. The scale bar is 20 μm. **(D)** Quantification of PDF levels in the dorsal projection of s-

LN$_v$s for control and *tara*$^{s132/1}$ males (n = 15-18). **(E)** Quantification of *Pdf* mRNA levels from

whole head extracts for control and *tara*$^{s132/1}$ males. Six independent biological samples, each

from 30 fly heads, were assayed. Mean ± SEM is shown. ****$p < 0.0001$, ***$p < 0.001$, **$p <$

0.01, *$p < 0.05$, ns: not significant; two-way ANOVA followed by Sidak's multiple comparisons

tests (B and D); Welch's *t*-test (E).

Figure 3. Overexpression of TARA in LN$_v$s results in slower PER cycling and lengthened

circadian period. (A) Representative expression patterns of PER and PDF in s-LN$_v$s of flies

overexpressing TARA (*Pdf > tara*) and parental controls at the indicated circadian times on the

3rd day in DD. The scale bar is 10 μm. **(B)** PER signal quantification in s-LN$_v$s (n = 8-16). **(C)**

Representative circadian actograms of male flies overexpressing *tara* in PDF-positive LN$_v$s (*Pdf*

> tara) and their parental controls. **(D-E)** Period length (D) and rhythm strength (E) of males of

indicated genotypes. Mean ± SEM is shown. ****$p < 0.0001$, ns: not significant; two-way

ANOVA followed by Tukey's post hoc test (B); Brown-Forsythe and Welch ANOVA tests

followed by Dunnett's T3 multiple comparisons tests (D and E).

Figure 4. Deletion of conserved TARA domains significantly reduces rhythm strength. (A)

Amino acid sequences of conserved TARA domains (boxed) and their flanking sequences. The

numbers indicate the positions in the TARA-B sequence. Conserved amino acids are shown in

bold. The dashes represent deleted amino acids, and gray indicates substituted amino acids. **(B)**

Representative circadian actograms of indicated genotypes. **(C-D)** Period length (C) and rhythm

strength (D) of males of indicated genotypes. Mean ± SEM is shown. ****p<0.0001, **p<0.001,

ns: not significant; one-way Brown-Forsythe and Welch ANOVA tests followed by Dunnett's T3

multiple comparisons test (C and D).

Figure 5. TARA interacts with PDP1 and VRI to regulate *Clk* transcription

(A) Luciferase activity of the *Clk-luc* reporter in *Drosophila* S2 cells transfected with UAS-*Pdp1*

(10 ng), UAS-*vri* (35 ng), and UAS-HA::*tara* (5 ng, 10 ng, or 15 ng) as indicated. Luciferase

activity was normalized to the condition that includes UAS-*Pdp1* and 15 ng of UAS-HA::*tara*

(+++). **(B)** Luciferase activity of the *Clk-luc* reporter in HEK293 cells transfected with CMV-

vri::VP16 (5 ng), CMV-*tara* (10 ng or 30 ng), and CMV-*TRIP-Br1* or CMV-*TRIP-Br2* (30 ng)

as indicated. Relative luciferase activity compared to the condition in which CMV-*vri::VP16*

alone is transfected is presented. **(C-D).** Representative Western blot of *Drosophila* S2 cells

transfected with the *Clk-luc* reporter, UAS-*Pdp1*, UAS-*vri*, and UAS-HA::*tara*, as indicated.

Antibodies against HA, PDP1, VRI, and Histone H3 were used. **(E-G)** Quantification of TARA

(E), PDP1 (F), and VRI (G) expression levels. **(H)** TARA forms a physical complex with VRI

and PDP1 in S2 cells. V5::TARA was immunoprecipitated with an anti-V5 antibody, and

antibodies against TARA, PDP1, and VRI were used for Western blotting. **(I)** Luciferase activity

of the *per-luc* reporter in *Drosophila* S2 cells transfected with UAS-HA::*tara* (10 ng or 30 ng),

and UAS-*Clk* (1 ng). Relative luciferase activity compared to the condition in which UAS-*Clk*

alone is transfected is presented. **(J)** The assay was conducted as in (I), except with three doses

of co-transfected UAS-*per* (2 ng, 5 ng, or 10 ng). Mean ± SEM is shown. ***p<0.001, **p<0.01,

*p<0.05, ns: not significant; two-way ANOVA followed by Tukey's post hoc tests (A); Welch's

t-test (E); one-way Brown-Forsythe and Welch ANOVA followed by Dunnett's T3 multiple

comparisons test (F-G, and I); two-way ANOVA followed by Sidak's multiple comparisons tests

(J). The effect of PER levels on *per-luc* activity was significant irrespective of the presence (+)

or absence (-) of TARA. None of the pairwise comparisons of the +TARA vs. -TARA conditions

at each level of PER revealed a significant difference (J).

Figure 6. TARA enhances the transcriptional activity of PDP1 and VRI and forms a

physical complex with them. (A) Schematic of TARA domain deletion mutants. **(B)** Luciferase

activity of the *Clk-luc* reporter in *Drosophila* S2 cells transfected with 30 ng of full-length UAS-

HA::*tara* (WT) or HA-tagged TARA domain deletion constructs (ΔCAB, ΔSERTA, ΔPBB, and

ΔCT). Luciferase activity compared to the condition in which UAS-*Pdp1* is transfected with full-

length TARA is presented. **(C)** Luciferase activity of the *Clk-luc* reporter in HEK293 cells

transfected with 5 ng of CMV-*vri::VP16* and 30 ng of full-length CMV-*tara* (WT) or a SERTA

domain deletion construct (ΔSERTA). Relative luciferase activity compared to the condition in

which CMV-*vri::VP16* is transfected with full-length CMV-*tara* is presented. **(D)** Western blot

of *Drosophila* S2 cells transfected with a *Clk-luc* reporter with 30 ng of full-length UAS-

HA::*tara* (WT) or HA-tagged domain deletion constructs (ΔCAB, ΔSERTA, ΔPBB, and ΔCT)

and 15 ng UAS-*Pdp1* as indicated. **(E-F)** Quantification of TARA (E) and PDP1 (F) expression

levels. Mean ± SEM is shown. ****$p<0.0001$, ***$p<0.001$, **$p<0.01$, *$p<0.05$, ns: not

significant; one-way Brown-Forsythe and Welch ANOVA followed by Dunnett's T3 multiple

comparisons test relative to a control condition (B,C,E, and F).

Figure 7. *tara* regulates the clock speed through genetic interactions with *vri* and *Pdp1*. (A)

Representative circadian actograms of males of indicated genotypes. **(B)** Period length (top) and

rhythm strength (bottom) of indicated genotypes. N = 36-64 for period length, n = 40-67 for

rhythm strength. The horizontal dotted line represents the mean period length (top) and rhythm

strength (bottom) for the common parental control (*Pdf* > +). Mean ± SEM is shown. ****p <

0.0001, ***p < 0.001, **p<0.01, *p < 0.05, ns: not significant; one-way Brown-Forsythe and

Welch ANOVA followed by Dunnett's T3 multiple comparisons test.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

763 **Table 1. Circadian locomotor rhythm phenotypes in DD**

Genotype	N	% R	% WR	% AR	tau (h) ± SEM	Power ± SEM
Knockdown						
<u>Female</u>						
+ > *tara* RNAi	63	96.8	1.6	1.6	24.55 ± 0.05	114.1 ± 3.57
elav > +	53	88.7	3.8	7.5	24.31 ± 0.17	98.3 ± 5.14
elav > *tara* RNAi	71	26.8	21.1	52.1	24.03 ± 0.09	30.6 ± 4.86
tim > +	47	83.1	14.8	2.1	24.64 ± 0.05	91.7 ± 5.05
tim > *tara* RNAi	45	60	28.9	11.1	24.36 ± 0.07	62.8 ± 4.90
Pdf > +	29	100	0	0	24.86 ± 0.05	106.0 ± 4.57
Pdf > *tara* RNAi	42	100	0	0	24.79 ± 0.06	114.6 ± 3.32
<u>Male</u>						
+ > *tara* RNAi	38	100	0	0	23.66 ± 0.05	153.6 ± 3.40
tim > +	47	100	0	0	24.39 ± 0.05	135.6 ± 3.23
tim > *tara* RNAi	43	72	14	14	24.26 ± 0.08	69.38 ± 5.90
Pdf > +	31	100	0	0	24.31 ± 0.06	137.6 ± 2.98
Pdf > *tara* RNAi	47	100	0	0	24.09 ± 0.05	131.6 ± 3.04
Overexpression						
<u>Male</u>						
Pdf > +	74	98.6	0	1.4	24.09 ± 0.05	132.6 ± 3.03
+ > *tara*	62	100	0	0	23.97 ± 0.06	130.8 ± 3.09
Pdf > *tara*	54	100	0	0	24.73 ± 0.04	116.3 ± 2.44

765 N: number of flies; R: rhythmic; WR: weakly rhythmic; AR: arrhythmic.

766 tau: free-running period; Power: a measure of rhythm strength

Table 2. Circadian locomotor rhythm phenotypes of TARA domain deletion mutants.

Genotype	N	% R	% WR	% AR	tau (h) ± SEM	Power ± SEM
control	28	100	0	0	23.67 ± 0.06	137.0 ± 5.87
ΔCAB	52	71.2	5.8	23.1	23.75 ± 0.07	79.4 ± 8.03
s132/+	66	95.5	1.5	3.0	23.56 ± 0.03	114.7 ± 5.46
s132/ΔSERTA	31	58.1	22.6	19.4	23.63 ± 0.05	56.5 ± 7.76
control	70	97.1	1.4	1.4	23.58 ± 0.03	142.5 ± 4.20
ΔPBB	67	94.0	1.5	4.5	23.61 ± 0.03	121.5 ± 4.39
control	30	96.8	3.2	0	23.54 ± 0.06	134.4 ± 6.59
ΔCT	31	71.6	15.6	12.5	23.56 ± 0.08	85.0 ± 7.57

N: number of flies; R: rhythmic; WR: weakly rhythmic; AR: arrhythmic.

tau: free-running period; Power: a measure of rhythm strength

771   **Table 3. Circadian locomotor rhythm phenotypes of TARA, VRI, and PDP1**

772   **overexpression in PDF+ neurons**

Genotype	N	% R	% WR	% AR	tau (h) ± SEM	Power ± SEM
Pdf > +	64	95.2	3.2	1.6	23.88 ± 0.06	113.8 ± 3.94
+ > tara	67	92.5	6	1.5	23.61 ± 0.03	115.3 ± 4.59
Pdf > tara	61	93.5	4.9	1.6	24.57 ± 0.05	95.4 ± 4.15
+ > vri	42	100	0	0	23.64 ± 0.05	145.6 ± 3.85
Pdf > vri	40	97.5	2.5	0	25.23 ± 0.07	134.5 ± 4.21
+ > vri, tara	46	100	0	100	23.90 ± 0.06	143.6 ± 2.39
Pdf > vri, tara	40	95	5	0	24.84 ± 0.05	127.6 ± 4.50
+ > Pdp1	55	96.4	1.8	1.8	23.68 ± 0.07	129.8 ± 5.22
Pdf > Pdp1	59	67.8	18.6	13.6	24.21 ± 0.08	79.4 ± 5.81
+ > Pdp1, tara	47	97.9	2.1	0	23.62 ± 0.04	130.4 ± 3.79
Pdf > Pdp1, tara	44	59.1	22.7	18.2	24.07 ± 0.11	58.5 ± 4.80
+ > Pdp1, vri	56	93	3.5	3.5	23.76 ± 0.04	104.3 ± 5.27
Pdf > Pdp1, vri	55	94.6	3.6	1.8	24.72 ± 0.09	102.7 ± 4.16

775   N: number of flies; R: rhythmic; WR: weakly rhythmic; AR: arrhythmic.

776   tau: free-running period; Power: a measure of rhythm strength.