DNA methylation-environment interactions in the human genome

Rachel A. Johnston[1,2,3], Katherine A. Aracena[4], Luis B. Barreiro[4,5,6], Amanda J. Lea[7,8], Jenny Tung[1,8, 9,10,11]

[1]Department of Evolutionary Anthropology, Duke University, Durham, NC 27708, USA
[2]Zoo New England, Boston, MA 02121, USA
[3]Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
[4]Department of Human Genetics, University of Chicago, Chicago, IL 60637, USA
[5]Section of Genetic Medicine, Department of Medicine, University of Chicago, Chicago, IL 60637, USA
[6]Committee on Immunology, University of Chicago, Chicago, IL 60637, USA
[7]Department of Biological Sciences, Vanderbilt University, Nashville, TN 37240, USA
[8]Canadian Institute for Advanced Research, Toronto, Canada M5G 1Z8
[9]Duke Population Research Institute, Duke University, Durham, NC 27708, USA
[10]Department of Biology, Duke University, Durham, NC 27708, USA
[11]Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Saxony, Germany

*Corresponding author: E-mail: jtung@eva.mpg.de; racheljohnston7@gmail.com

Abstract

Previously we showed that a massively parallel reporter assay, mSTARR-seq, could be used to simultaneously test for both enhancer-like activity and DNA methylation-dependent enhancer activity for millions of loci in a single experiment (Lea *et al*., 2018). Here we apply mSTARR-seq to query nearly the entire human genome, including almost all CpG sites profiled either on the commonly used Illumina Infinium MethylationEPIC array or via reduced representation bisulfite sequencing. We show that fragments containing these sites are enriched for regulatory capacity, and that methylation-dependent regulatory activity is in turn sensitive to the cellular environment. In particular, regulatory responses to interferon alpha (IFNA) stimulation are strongly attenuated by methyl marks, indicating widespread DNA methylation-environment interactions. In agreement, methylation-dependent responses to IFNA identified via mSTARR-seq predict methylation-dependent transcriptional responses to challenge with influenza virus in human macrophages. Our observations support the idea that pre-existing DNA methylation patterns can influence the response to subsequent environmental exposures—one of the tenets of biological embedding. However, we also find that, on average, sites previously associated with early life adversity are not more likely to functionally influence gene regulation than expected by chance.

Introduction

DNA methylation is sensitive to the environment, can remain stable across cell divisions, and in some contexts, can alter gene regulation. Consequently, it has received a high level of attention as a potential pathway linking environmental exposures to downstream trait variation, especially when these exposures are separated temporally, as proposed by the "biological embedding" hypothesis (Hertzman, 1999; Hertzman, 2012). In a canonical example, gestational manipulation of methyl donors in mice is causally implicated in changes in DNA methylation and expression levels of the *agouti* gene, which in turn affects susceptibility to obesity and insulin resistance later in life (Dolinoy *et al.*, 2006). However, the mechanism underlying the *agouti* case—"metastable" methylation of an inserted intracisternal A particle [IAP] class retrovirus—is now thought to be unusual (Kazachenka *et al.*, 2018). Thus, it remains unclear whether the many reports of environment-DNA methylation associations are likely to represent links along a causal pathway to phenotype, or are better viewed as passive biomarkers of exposure.

Experimental tests of the causal impact of DNA methylation on gene regulation indicate that both scenarios are possible. *In vitro* studies reveal that transcription factor binding is frequently (but not ubiquitously) sensitive to DNA methylation (O'Malley *et al.*, 2016; Yin *et al.*, 2017). More recently, epigenomic editing of DNA methylation marks has been shown to alter the activity of transcription factors important in disease and development *in vivo* (Greenberg and Bourc'his, 2019; Yim *et al.*, 2020), as well as the formation of CTCF-mediated gene loops (Lei *et al.*, 2017). However, other epigenetic editing studies show that, even within the same regulatory element, changes in DNA methylation have functional consequences for gene expression at only a subset of sites (e.g. Maeder *et al.*, 2013). Indeed, in the native genome, enhancer activity appears to be insensitive to DNA methylation levels in the majority of cases (Kreibich *et al.*, 2023). Time series analyses in human dendritic cells have also shown that, in response to a pathogen challenge, nearly all changes in gene expression precede changes in DNA methylation (Pacis *et al.*, 2019). Such methylation changes may nevertheless be functionally relevant if they influence the speed and magnitude of subsequent challenges (e.g., by marking the accessibility of latent enhancers across cell divisions) (Kamada *et al.*, 2018; Sun and Barreiro, 2020). This hypothesis may explain, for example, enhanced transcriptional responses to glucocorticoid stimulation in the descendants of hippocampal progenitor cells previously challenged with glucocorticoids (Provençal *et al.*, 2020).

Together, these findings suggest that environment-associated DNA methylation marks are mixed sets that include functionally silent sites, sites that are constitutively important for gene regulation, and sites in which DNA methylation levels affect gene regulation only under certain conditions. Disentangling which sites belong to which set is important for interpreting and prioritizing the results of a growing body of studies in the biological, social, and health sciences, especially those that test hypotheses that assume a functional role for DNA methylation, such as biological embedding (Hertzman, 2012; Demetriou *et al.*, 2015; Aristizabal *et al.*, 2020). To facilitate this process, we recently introduced a massively parallel reporter assay, mSTARR-seq, that is capable of testing the functional effects of DNA methylation on regulatory activity in high-throughput (Lea *et al.*, 2018). This method allowed us to test for methylation-dependent regulatory activity at millions of CpG sites in the human genome. However, our previous data set did not investigate how DNA methylation marks affect the response to environmental challenge. It also did not explicitly target the CpG sites that are the

most commonly assayed in the human genome (i.e., those featured on the Illumina
MethylationEPIC array).

Here, we address these gaps by using mSTARR-seq to construct a genome-wide map of
DNA methylation-dependent enhancer activity for 27.3 million CpG sites in the human genome,
accomplished by assessing 600 bp windows encompassing 93.5% of CpGs in the human genome
as a whole. This set includes 99.3% of CpG sites present on the Illumina MethylationEPIC array,
the most commonly used platform for profiling DNA methylation in humans, and 99.4% of CpG
sites accessible via reduced-representation bisulfite sequencing (RRBS; Meissner *et al.*, 2005).
To evaluate the importance of DNA methylation in responses to environmental challenge, we
performed mSTARR-seq under baseline conditions (i.e., no challenge) and following exposure to
two physiologically relevant environmental challenges: dosage with the synthetic glucocorticoid
dexamethasone, which plays a central role in metabolic regulation and stress-related
homeostasis, and interferon alpha (IFNA), a cytokine that elicits genomic and immunological
responses associated with viral infection.

We used these data to pursue three goals. First, we describe overall patterns of regulatory
activity and methylation-dependent regulatory activity across profiled sites, including how sites
targeted by the EPIC array or by RRBS compare to the genome as a whole. The results of this
analysis produce a new resource: a map of DNA methylation-dependent enhancer activity across
the human genome. Second, we test the degree to which methylation-dependent regulatory
activity is affected by exposure to steroid hormone or immune defense-related signaling
molecules. Our results yield insight into the frequency of DNA methylation-environment
interactions genome-wide, and provide support for the hypothesis that DNA methylation
potentiates the cellular response to external stressors. Finally, we illustrate the applicability of
this resource by testing two predictions of the biological embedding hypothesis: that pre-existing
DNA methylation levels can affect the transcriptional response to subsequent environmental
challenge (Provençal *et al.*, 2020; Sun and Barreiro, 2020; Fanucchi *et al.*, 2021), and that sites
linked to early life adversity (ELA) are likely to be functionally important for shaping gene
expression. Our results suggest that mSTARR-seq can identify DNA methylation-environment
interactions that also occur *in vivo* in human populations and can therefore help prioritize
environment- and ELA-associated loci for future follow-up.

Results

mSTARR-seq captures enhancer activity and methylation-dependent enhancer activity
genome-wide
We assessed regulatory activity and methylation-dependent regulatory activity for
4,558,475 600-base pair windows of the genome by pairing hybridization capture of targeted loci
in the human genome with the massively parallel reporter assay, mSTARR-seq (Lea *et al.*,
2018). In brief, mSTARR-seq performs enzymatic manipulation of DNA methylation across
hundreds of thousands to millions of reporter DNA fragments simultaneously. By measuring the
ability of fragments to self-transcribe (as in Arnold *et al.*, 2013; Shlyueva *et al.*, 2014; reviewed
in Gallego Romero and Lea, 2022), it generates estimates of the enhancer-like regulatory
potential for each fragment when that fragment is in an unmethylated versus a methylated state
(Figure 1A, B). Notably, results from the unmethylated condition are akin to those from a
conventional STARR-seq experiment, in that they assess regulatory activity irrespective of CpG
content or methylation status. To focus on the CpG sites most likely to be included in DNA

methylation studies in humans, we performed custom capture with SeqCap EZ Prime Choice
Probes (Roche), targeting all CpG sites on the Illumina Infinium MethylationEPIC array and
those likely to be profiled using reduced representation bisulfite sequencing, which enriches for
CpG sites near targets of *MspI* restriction enzyme digestion (Figure 1-figure supplement 1).
Because of substantial attention to potential environmental and early life effects on DNA
methylation at the glucocorticoid receptor gene (*NR3C1*, reviewed in Liu and Nusslock, 2018),
we also targeted the 6.5 Mb in and flanking *NR3C1*. Finally, to assess background expectations
for regulatory and methylation-dependent regulatory activity in the human genome, we targeted
a control set of 100,000 loci for capture, chosen at random across the human genome after
excluding centromeres, gaps, and uncalled bases in the hg38 genome.
We generated an mSTARR-seq library from captured DNA from the GM12878 cell line,
followed by transfection into the K562 erythroleukemic cell line and co-purification of plasmid-
derived RNA and DNA (6 replicates per condition: methylated versus unmethylated; see
Materials and Methods; Supplementary File 1). Based on our minimal criteria for assessment
(600 basepair windows where at least half of the plasmid-derived DNA samples had non-zero
reads covering the window; see Materials and Methods), 90.2% of the human genome was
included in the plasmid DNA library purified at the end of the mSTARR-seq experiment (mean
sequencing depth per replicate = 30.4 million reads; mean read coverage per window = 12.723 $\pm$
41.696 s.d.). Within this set, which also included off-target regions, windows were ~15-fold
enriched for targeted regions in the genome (Fisher's Exact Test $\log_2$(OR) [95% confidence
interval (CI)] = 3.971 [3.934, 4.009], p < 1.0 x 10^{-300}]). Thus, in windows that passed our
minimal assessment criteria, we successfully targeted 99.3% of CpG sites on the
MethylationEPIC array and 99.4% of sites likely to be included in RRBS libraries. Across target
sets, read depth was not predicted by CpG density, indicating no systematic power differences in
assessing methylation-dependent activity due to differences in CpG number per window (R^2 =
0.099, p = 0.6852; Supplementary file 2). Our results are comparable to published fragment
diversity levels achieved with this method (Lea *et al.*, 2018; Figure 1-figure supplement 2).
Because demethylation or remethylation of DNA fragments could occur within cells during the
experiment, we performed bisulfite sequencing at the end of the experiment. We confirmed that
DNA methylation levels were substantially higher in the methylated condition samples than in
the unmethylated samples, where methylation levels were near zero (mean methylated = 0.885,
mean unmethylated = 0.066, Figure 1-figure supplement 3; unpaired t-test: t = -14.66, df =
15,124, p = 2.39 x 10^{-10}). We note that the observed small deviations from 0% and 100%
methylation would reduce our power to detect methylation-dependent activity, but should not
incur false positive results.
To test for regulatory potential, we focused on those windows where at least 3 replicate
samples produced non-zero RNA-seq reads in either the methylated condition or in the
unmethylated condition (n = 216,091 non-overlapping 600 bp windows, 4.3% of the human
genome; mean DNA-seq read coverage per window = 93.907 $\pm$ 10.091 s.d.; mean RNA-seq read
coverage per window = 165.093 $\pm$ 1008.816). These inclusion criteria focused our analysis on
the subset of windows where we were able to detect minimal evidence of transcription from the
plasmid. The proportion of the human genome showing evidence of regulatory activity in
STARR-seq family assays is expected to be small based on previous work. For example, Johnson
and colleagues estimated that 0.165% of the human genome had regulatory potential in
unstimulated A549 cells in a genome-wide STARR-seq assay, based on the proportion of hg38
basepairs that fell in regions with regulatory activity in their assay (Johnson *et al.*, 2018). Among

the regions analyzed here, 3,721 windows (1.7%) showed evidence for regulatory activity, where
the amount of RNA generated from a window significantly exceeded the amount of DNA input
sequenced for the same window (FDR < 1%; Supplementary File 3). Sequencing depth does not
appear to constrain the ability to detect regulatory regions, as rarefaction analyses suggest that
increasing sequencing depth of RNA or DNA libraries would add only a small number of
additional regions with any evidence of transcriptional activity (Figure 1-figure supplement 4).
Further, among analyzed windows, DNA coverage per fragment explains minimal variance in
mSTARR-seq enhancer activity ($R^2 = 0.029$, $p < 1.0 \times 10^{-300}$).
Regions with regulatory activity were enriched for enhancers and active promoters
annotated by the ENCODE Consortium for the K562 cell line ($\log_2(OR) = 0.584 – 3.337$; all $p <$
1×10^{-3}; Figure 1C; Supplementary file 4 (ENCODE, 2012; Ernst and Kellis, 2012); note that
previous research has found that promoters can show enhancer-like activity in STARR-seq-type
assays (e.g. Klein *et al.*, 2020). In support of assay reproducibility, regulatory regions also were
highly consistent with previous mSTARR-seq data generated in K562 cells using a previously
published, independently generated DNA library that we reanalyzed using the current pipeline
(overlap among windows with FDR < 1% in each data set analyzed independently: $\log_2(OR)$
[95% CI] = 6.212 [6.086, 6.440], $p < 1.0 \times 10^{-300}$; Figure 1-figure supplements 5-6;
Supplementary Files 5-6) (Lea *et al.*, 2018).
Regions with mSTARR-seq-annotated regulatory activity also exhibited a high degree of
consistency between cell types. In a comparison against newly generated mSTARR-seq data
from the HepG2 liver cell line, using the same mSTARR-seq library as in the previously
published K562 experiments (Lea *et al.*, 2018), regulatory regions detected in this study were
enriched among regulatory regions detected in HepG2 cells ($\log_2(OR)$ [95% CI] = 3.534 [3.381,
3.684], $p = 5.21 \times 10^{-307}$; Figure 1-figure supplements 5 and 7; Supplementary File 7). They also
overlap with regulatory regions identified in a conventional STARR-seq experiment in A549
lung epithelial cells ($\log_2(OR)$ [95% CI] = 2.451 [2.442, 2.461], $p < 1.0 \times 10^{-300}$) (Johnson *et al.*,
2018). Although we prioritized a blood-derived cell line because of the frequency of
environmental epigenetic studies done in blood, these results suggest that our findings can be
generalized, to some extent, to other cell and tissue types.
Among the 3,721 windows with regulatory activity in either the methylated or
unmethylated condition, 1,768 windows (47.5% of regulatory windows; FDR < 1%) were
differentially active depending on condition, pointing to DNA methylation-dependent regulatory
activity. This result is also concordant when we apply the current pipeline to previously
published data from K562s and our new HepG2 data, which exhibit correlated effects of DNA
methylation on regulatory activity in the windows examined in both cases (K562: $R^2 = 0.286$ for
1250 regulatory windows with FDR < 1% in both data sets, $p = 3.19 \times 10^{-93}$; HepG2: $R^2 = 0.277$
for 511 regulatory windows with FDR < 1% in both data sets, $p = 8.87 \times 10^{-38}$ Figure 1-figure
supplement 8). Among methylation-dependent regulatory regions, the majority of regions (1744
windows, 98.6%) were more active in the unmethylated condition than the methylated condition
(Figure 1-figure supplements 9, 10). Consistent with previous findings (Lea *et al.*, 2018),
regulatory regions with more CpGs are more likely to be repressed by DNA methylation (Figure
1-figure supplement 11; Spearman's rho = 0.370, $p = 9.865 \times 10^{-121}$; n = 3,721 regions with
mSTARR-seq regulatory activity). Regulatory windows showing higher activity in the
methylated condition (24 windows) were enriched for binding motifs of the transcription factors
p53, which has been previously reported to have increased binding affinity to methylated DNA
relative to unmethylated DNA (Kribelbauer *et al.*, 2017) ($\log_2(OR)$ [95% CI] = 2.352 [0.714,

3.767], p = 2.91 x 10^{-3}; Supplementary File 8), and Tfcp2I1, which has been found to recruit
Tet2 to mediate enhancer demethylation (Sardina *et al.*, 2018) ($\log_2$(OR) [95% CI] = 2.615
[0.580, 4.229], p = 6.84 x 10^{-3}).

Figure 1. Study design and identification of enhancer activity. (A) Sheared DNA from the GM12878
cell line was subjected to enrichment via capture with probes targeting loci selected in reduced
representation bisulfite sequencing (RRBS) workflows (*MspI* targets), CpG sites on the Infinium EPIC
array, the gene *NR3C1* and flanking regions, and 100,000 randomly distributed control regions. Note that
a single 600 bp window can contain multiple target types (Figure 1-figure supplement 1). Library
diversity summaries are shown in Figure 1-figure supplement 2. **(B)** Captured loci were cloned into the
mSTARR-seq vector, *pmSTARRseq1*, treated with either the CpG methylating enzyme *M. SssI* or a sham
treatment, and transfected into K562 cells. Methylation levels post-transfection and rarefaction analyses
of sequencing depth of replicate samples are shown in Figure 1-figure supplements 3, 4. Right panel
shows an example of DNA methylation-dependent regulatory activity near the first exon of the *TTC32*
gene, where the methylation-dependent regulatory element overlaps an active promoter chromatin state
(red horizontal bar denotes active promoter as defined by ENCODE: ENCODE, 2012). **(C)** mSTARR-seq
regulatory activity in the baseline condition is strongly enriched in ENCODE-defined enhancers and some
classes of promoters (indicated in blue), and depleted in repressed, repetitive, and heterochromatin states.
See Supplementary file 4 for full results of this analysis. Regions with mSTARR-seq regulatory activity
detected in this experiment also significantly overlap with regions with regulatory activity in other
mSTARR-seq and conventional STARR-seq datasets (Figure 1-figure supplement 5; see also Figure 1-
figure supplement 6 for estimates of concordance across technical replicates). **(D)** Left column shows,
under the baseline condition (i.e. unstimulated cells), the proportion of 600 bp windows that exhibited
minimal regulatory activity (at least 3 replicate samples produced non-zero RNA-seq reads in either the
methylated condition or the unmethylated condition) in the mSTARR-seq assay (pink) versus those with
detectable input DNA but no evidence of regulatory activity (blue), for windows containing sites from
each target set. Right column shows the proportion of windows with regulatory capacity (i.e., the subset
of the windows represented in pink on the left that produce excess RNA relative to the DNA input at FDR

< 1%) that are also methylation-dependent (dark brown). Within each column, pie charts are scaled by the
total numbers of windows represented. See Figure 1-figure supplements 7 and 8 for comparisons to
regulatory regions in other datasets. Figure 1-figure supplements 9 and 10 show window-level RNA to
DNA ratios. Figure 1-figure supplement 11 shows the relationship between CpG density and methylation-
dependent regulatory activity.
254
255   **Commonly studied CpG sites are enriched for methylation-dependent regulatory activity**
256   We compared methylation-dependent regulatory activity between CpGs targeted on the
EPIC array, CpGs typically profiled in RRBS libraries (i.e., near *MspI* cut sites), the 6.5 Mb in or
flanking the glucocorticoid receptor gene *NR3C1*, and the control set of genome-representative
loci. As expected, windows that contained EPIC CpG sites, sites associated with RRBS, or that
were located in or near *NR3C1* were significantly more likely to show at least some degree of
transcriptional activity (i.e., show plasmid-derived RNA reads in at least half the sham or
methylated replicates) than the genomic background (i.e., 98,967 randomly distributed loci: see
Materials and Methods), and thus were more likely to be included in our analysis set (n =
216,091 windows) for regulatory activity (EPIC: $\log_2$(OR) [95% CI] = 2.189 [2.152, 2.227], p <
1.0×10^{-300}; RRBS: $\log_2$(OR) [95% CI] = 4.126 [4.086, 4.163], p < 1.0×10^{-300}; *NR3C1*:
$\log_2$(OR) [95% CI] = 3.259 [3.193, 3.325], p < 1.0×10^{-300}; Figure 1D). However, conditional on
minimal transcriptional activity, windows containing EPIC array sites were not more likely to
exhibit *significant* enhancer-like regulatory activity (FDR < 1%; n = 3,721 windows) than
background windows of the genome, and RRBS- and *NR3C1*-associated sites were in fact
slightly less likely to do so (EPIC: $\log_2$(OR) [95% CI] = 0.112 [-0.168, 0.405], p = 0.47; RRBS:
$\log_2$(OR) [95% CI] = -0.493 [-0.779, -0.195], p = 1.11×10^{-3}; *NR3C1*: $\log_2$(OR) [95% CI] = -
0.730 [-1.263, -0.217], p = 3.89×10^{-3}). This result is likely explained by our inclusion criteria,
as only regions with evidence for minimal RNA transcription were retained prior to formal
analysis.
275   In contrast, among regions with detectable regulatory activity, 63.3% of those containing
RRBS-associated CpG sites exhibited methylation-dependent regulatory activity, compared to
45.9% of control background loci (RRBS against control: $\log_2$(OR) [95% CI] = 1.025 [0.430,
1.624], p = 4.39×10^{-4}; Figure 1D). Note that of 109 regulatory windows in the control set, 108
contain at least 1 CpG, so this difference is not because the control set is CpG-free (mean
number CpGs per fragment in control set = 13.291 ± 11.179 s.d.; EPIC = 19.641 ± 14.471;
RRBS = 20.870 ± 13.883; *NR3C1* = 7.405 ± 9.409). Neither windows with EPIC CpGs nor
windows in or near *NR3C1* were enriched for methylation-dependent regulatory activity relative
to the genomic background (EPIC against control: $\log_2$(OR) [95% CI] = 0.284 [-0.297, 0.871], p
= 0.33; *NR3C1* versus control: $\log_2$(OR) [95% CI] = -0.948 [-2.129, 0.178], p = 0.11).
Consequently, differential methylation identified through RRBS is more likely to be capable of
driving differences in gene regulation, as detectable by mSTARR-seq, than differential
methylation elsewhere in the genome. We caution, however, that even among RRBS sites, 98.8%
do not occur in regions of the genome with detectable regulatory activity in either the methylated
or unmethylated conditions, at least in the K562 cell type, and 35.0% of those that fall in putative
regulatory elements exhibit no evidence for methylation-dependent activity.
291
292   **Environmental perturbation reveals cryptic regulatory elements and cryptic effects of DNA**
293   **methylation**
294   Enhancer activity can be cell type- or environment-dependent (e.g., Ostuni *et al.*, 2013;
Johnson *et al.*, 2018; Chaudhri *et al.*, 2020). DNA methylation-dependent enhancer activity may

show similar context-dependence, thus potentially accounting for the large number of sites that
fall in functionally silent regions described above. However, this possibility has not been
systematically tested. To do so, we next compared regulatory activity between the baseline
unchallenged condition and cells challenged with interferon alpha (IFNA) or dexamethasone
(dex) (Figure 2A; Figure 2-figure supplements 1-2; Supplementary file 9). Regions that were
identified to have regulatory potential in the baseline condition were highly likely to retain
regulatory potential after cells were challenged with IFNA or dex (Figure 2-figure supplement 3;
IFNA $\log_2$(OR) [95% CI] = 8.639 [8.499, 8.812], p < 1.0 x 10^{-300}; dex $\log_2$(OR) [95% CI] =
9.640 [9.483, 9.776], p < 1.0 x 10^{-323}; Supplementary Files 10-11). However, environmental
challenges also revealed thousands of putative regulatory regions that were undetectable at
baseline but active post-stimulation (Figure 2B; 1614 IFNA-specific; 1131 dex-specific). Of
4,632 IFNA regulatory regions (<1% FDR), 44.1% are not detectable at baseline at a 1% FDR
threshold in the baseline condition, and even with a relaxed baseline FDR of 10%, 25.1% remain
undetectable. Of 4,217 dex regulatory regions (<1% FDR), 40.2% are not detectable at baseline
(1% FDR), and 31.5% remain undetectable at a baseline FDR of 10%.
311   Regulatory windows specific to the IFNA treatment were enriched for 33 transcription
factor binding motifs (TFBMs) (Supplementary File 12), with strong enrichments detected for
TFBMs involved in innate immune defense in general, and interferon signaling specifically
(Figure 2C). For example, the most enriched motif was the canonical DNA target of interferon
signaling, known as IFN-stimulated response elements (ISRE; $\log_2$(OR) [95% CI] = 3.158
[2.953, 3.358], Bonferroni corrected p = 7.31 x 10^{-133}), followed by binding motifs for several
IFN-regulatory factors (IRF1, IRF2, IRF3, IRF4, IRF8; all OR > 1.5 and Bonferroni corrected p
< 1 x 10^{-15}) (Chen *et al.*, 2017). Regulatory windows specific to the dex-stimulated condition
were significantly enriched for 28 transcription factor binding motifs, including the
glucocorticoid response element IR3 and binding motifs of several transcription factors known to
interact with or be modulated by the glucocorticoid receptor (AP-1, CEBP:CEBP, CEBP:AP1,
JunB, Jun-AP1, GATA1, STAT3, and STAT5; all OR > 1.3, Bonferroni corrected p < 0.01;
Supplementary File 13) (Cain and Cidlowski, 2017). Results were qualitatively similar if we
used a more stringent definition of IFNA-specific and dex-specific regulatory activity (e.g.
"IFNA-specific" defined as FDR < 1% in IFNA condition and FDR > 10% in the other two
conditions; Supplementary Files 14-15).
327   To evaluate the relevance of these regions to *in vivo* gene regulation, we also generated
matched mRNA-seq data for the endogenous K562 genome from the same experiments. These
data showed that expressed genes located closest to mSTARR-seq-annotated, IFNA-specific
enhancers were more strongly upregulated after IFNA stimulation than the set of expressed
genes located closest to shared mSTARR-seq-annotated enhancers (i.e., those identified in both
the IFNA and at least one other condition, considering genes within 100 kb maximum distance
from each enhancer element; unpaired t-test; t = 3.268, df = 601.88, p = 1.15 x 10^{-3}). We found
similar results when using inferred enhancer-gene linkages from the EnhancerAtlas 2.0 (Gao *et*
al., 2016) and restricting the set of IFNA-specific enhancers to those with external experimental
support for ISRE binding (n = 1005 windows; ChIP-Seq data from ENCODE [2012], relative to
all other genes in the gene expression dataset; unpaired t-test; t = 3.579, df = 118.36, p = 5.01 x
10^{-4}; Figure 2D; Supplementary File 16). Despite a weaker overall gene expression response to
dex, genes closest to dex-specific mSTARR-seq enhancers were also more strongly upregulated
after dex stimulation than genes near shared enhancers (unpaired t-test; t = 3.477, df = 479.6, p =
5.53 x 10^{-4}).

342   For the IFNA challenge, the DNA methylation state of each window appears to play an
343 important role in shaping condition-specific responses to stimulation in the mSTARR-seq data
344 set. Nearly twice as many regulatory windows (81.4%; 1314 of 1614) exhibited methylation-
345 dependent regulatory activity in the IFNA-specific condition than in the baseline or dex-specific
346 condition (47.5% and 48.4% respectively; two-sided binomial test for IFNA compared to
347 baseline: $p = 3.58 \times 10^{-312}$). Further, regulatory regions that harbor TFBMs for TFs central to the
348 interferon response (ISRE, IRF1, IRF2, IRF3, IRF4, IRF8; N = 663 windows) strongly
349 responded to IFNA challenge if in an unmethylated state, but mounted systematically attenuated
350 responses if in a methylated state. As a result, 562 of these 663 windows (84.7%) exhibited
351 significant methylation-dependent regulatory activity and 561 of them (99.8%) were more active
352 in the IFN-challenged state when unmethylated. This pattern is recapitulated when focusing on
353 analyzed windows with external experimental support for ISRE binding (n = 1005 windows;
354 ChIP-Seq data from ENCODE [2012]). These windows show no evidence for methylation-
355 dependent regulatory activity in the baseline condition (paired t-test; t = -0.792, df = 1004, p =
356 0.43), primarily because they show no regulatory activity at all without IFNA stimulation. After
357 IFNA stimulation, though, they exhibit strong methylation-dependence. Specifically, only
358 unmethylated windows are capable of a response (paired t-test; t = 31.748, df = 1004, $p = 1.02 \times$
359 10^{-153}; Figure 2E).

enrichment results. **(D)** Genes targeted by ISRE enhancers (ISRE enhancers identified from ENCODE
ChIP-seq data; gene targets identified from enhancer-gene linkages from EnhancerAtlas 2.0: Gao *et al.*,
2016) that are also identified as IFNA condition-specific mSTARR-seq enhancers show stronger K562
endogenous gene expression responses to IFNA stimulation than non-ISRE targets (unpaired t-test: t =
3.58, df = 118.36, p = 5.01 x 10^{-4}; Supplementary File 16). Each box represents the interquartile range,
with the median value depicted as a horizontal bar. Whiskers extend to the most extreme values within
1.5x of the interquartile range. **E)** mSTARR-seq regulatory activity for windows containing ISRE targets
(n = 1005 windows) interacts strongly with exposure to IFNA. These windows are capable of mounting a
strong response to IFNA stimulation when unmethylated (dashed line; paired t-test: t = 23.02, df = 1004,
p = 1.78 x 10^{-94}) but not when methylated (solid line; paired t-test: t = -1.74, df = 1004, p = 0.082). Dots
show the mean beta corresponding to enrichment of RNA reads versus DNA reads across windows;
whiskers show the standard error. Because y-axis values correspond to model estimates, they can be
positive (i.e., more mSTARR-seq RNA reads than input DNA reads) or negative values (i.e., fewer
mSTARR-seq RNA reads than mSTARR-Seq input DNA reads, indicating no regulatory activity).
390
Methylation levels at mSTARR-seq IFNA-specific enhancers predict the transcriptional
response to influenza virus in human macrophages
Our findings show that, in the context of mSTARR-seq, pre-existing DNA methylation
state can capacitate or constrain the regulatory response to IFNA stimulation. This result
suggests that DNA methylation-environment interactions may be an important determinant of
gene expression levels *in vivo*. To test this possibility, we drew on matched whole genome
bisulfite sequencing (WGBS) and RNA-seq data collected from human monocyte-derived
macrophages (n = 35 donors), with and without infection with the influenza A virus (IAV),
commonly known as flu (Figure 3A; Aracena *et al.*, in press). We asked whether variation in the
gene expression response to flu is predicted by DNA methylation levels in the baseline (non-
infected) condition, especially at loci where the response to IFNA is affected by DNA
methylation in mSTARR-seq. Importantly, flu and IFNA challenges induce similar innate
immune responses (Killip *et al.*, 2015).
Within each individual in the macrophage data set, mean DNA methylation levels at
baseline significantly predict the mean gene expression response to flu across the full set of
mSTARR-seq enhancer windows detected in the IFNA condition: higher methylation at baseline
predicts an attenuated gene expression response, on average (n = 35 individuals at 2769 enhancer
windows: mean Pearson's r = -0.105 ± 0.006 s.d., all Bonferroni-corrected p < 3 x 10^{-5};
Supplementary File 17). This effect is largely driven by the subset of mSTARR-seq IFNA-
specific enhancers. Specifically, the correlation between baseline DNA methylation levels and
the gene expression response to flu is 3.44-fold stronger in IFNA-specific enhancers than for
enhancers identified in both the IFNA condition and at least one other condition (Figure 3B;
IFNA-specific enhancers: n = 1033, mean Pearson's r = -0.170 ± 0.009 s.d., all Bonferroni-
corrected p < 2 x 10^{-5}; shared enhancers: n = 1736, mean Pearson's r = -0.049 ± 0.01 s.d., all
Bonferroni-corrected p > 0.1). These results appear to be driven by strong methylation-
dependence in flu-infected cells, as the average within-individual correlation between DNA
methylation and gene expression is 2.44 times as large after infection (r = -0.261 ± 0.006) than at
baseline (r = -0.106 ± 0.008) in IFNA-specific mSTARR-seq enhancers (Figure 3C;
Supplementary File 18).
The limited sample size of the macrophage data set makes it better suited for analyses of
the overall relationship between DNA methylation and gene expression within an individual,
rather than locus-specific analyses of interindividual variation (especially as locus-specific

variance in DNA methylation across individuals is low: mean = 0.004, standard deviation =
0.008). Nevertheless, across 1382 testable loci (600 bp windows containing at least 1 CpG with
interindividual variance > 0.01), we identified one IFNA-specific, methylation-dependent
mSTARR-seq enhancer where endogenous variation in DNA methylation levels across
individuals clearly predicts the response of the nearest gene's transcriptional response to flu
(Figure 3D-E; p = 6.05 x 10^{-5}, q-value = 0.0837; Supplementary File 19). This mSTARR-seq
enhancer (chr1:1013400-1014000) overlaps the promoter of interferon-stimulated gene 15
(*ISG15*; Figure 3E), and its average methylation explains 38% of the variance in the *ISG15*
transcriptional response to flu across individuals. The magnitude of *ISG15* transcriptional
response to flu appears primarily driven by the enhancer's effect on gene expression at baseline.
Lower enhancer methylation at baseline is associated with higher *ISG15* expression at baseline,
ultimately resulting in a shallower fold change response to flu (Figure 3-figure supplement 1).
This example illustrates the value of integrating observational data on DNA methylation and
gene expression with mSTARR-seq. The population data support the *in vivo* relevance of the
mSTARR-seq data, while the mSTARR-seq data suggest that baseline variation in *ISG15*
methylation *in vivo* is causally meaningful to the response to influenza. Notably, *ISG15* plays
critical roles in regulating the type I interferon response and modulating host immunity to both
viral and bacterial infections (reviewed in Perng and Lenschow, 2018).

Figure 3. DNA methylation in mSTARR-seq enhancers predicts *in vivo* gene expression in
macrophages. (A) Study design of the in *vivo* experiment, in which matched macrophage samples from
35 individuals were either left non-infected or infected with influenza A virus (IAV) for 24 hours and

processed for RNA-seq and whole genome bisulfite sequencing (WGBS; Aracena *et al.*, in press). **(B)**
Within individuals, DNA methylation (DNAm) levels at mSTARR-seq enhancers in non-infected cells
are negatively correlated with the nearest genes' transcriptional responses to IAV, but only in mSTARR-
seq enhancers that were specific to the IFNA condition (IFNA-specific enhancers: n = 1033, mean
Pearson's r = -0.170 ± 0.009 s.d., all Bonferroni-corrected $p < 2 \times 10^{-5}$; shared enhancers: n = 1736, mean
Pearson's r = -0.049 ± 0.01 s.d., all Bonferroni-corrected p > 0.1). Each colored line represents an
individual, and vertical gray lines represent 95% confidence intervals (see Supplementary File 17 for full
results). **(C)** The average within-individual correlation (r) between DNA methylation and gene expression
(GE) is 2.44 times as large after infection (r = -0.261 ± 0.006) than at baseline (r = -0.106 ± 0.008) in
IFNA-specific mSTARR-seq enhancers (right panel), but much less affected by infection at mSTARR-
seq enhancers that are shared across conditions (left panel). Within each panel, each colored line
represents an individual for the same set of enhancers (see Supplementary File 18 for full results), without
and with IAV infection. **(D)** Across individuals, the *ISG15* transcriptional response to IAV is significantly
correlated with average DNAm at the mSTARR-seq enhancer chr1:1013400-1014000 in non-infected
cells ($R^2 = 0.381$, $p = 6.05 \times 10^{-5}$, q = 0.084). Each dot represents an individual (see Supplementary File
19 for full results and Figure 3-figure supplement 1 for condition-specific results). **(E)** The mSTARR-seq
enhancer predictive of *ISG15* response to IAV (dark green bar) is located in the active promoter of *ISG15*
(as defined by ENCODE [ENCODE, 2012]; red denotes active promoter, pink denotes weak promoter,
orange denotes strong enhancer, yellow denotes weak enhancer). Three adjacent, methylation-dependent,
IFNA-specific mSTARR-seq enhancers were identified (light green), but do not significantly predict
ISG15 response to IAV (q > 10%). The bottom 6 tracks depict non-normalized, raw read pileups for
mSTARR-seq RNA replicates in either the unmethylated (open circle) or methylated (filled circle)
condition, with all y-axis maximums set to 20,000.
467
Most CpG sites associated with early life adversity do not show regulatory activity in K562s
Finally, because changes in DNA methylation are of particular interest to research on the
biological embedding of early life experience (Hertzman and Boyce, 2010), we tested whether
DNA methylation differences associated with early life adversity (ELA) translate to functional
effects on gene regulation in the mSTARR-seq data. We first performed a literature search to
compile CpG sites that have previously been associated with ELA in humans using the Illumina
EPIC array or one of its precursors (Infinium Human Methylation 450K and 27K BeadChips).
Our search resulted in a total of 27 studies (Supplementary File 20), which together identified
8,525 unique ELA-associated sites.
For 26 of 27 studies, ELA-associated CpG sites were not more likely to occur within
putative regulatory windows (detected in either the methylated condition, unmethylated
condition, or both) than background chance (Figure 4). This pattern was qualitatively consistent
regardless of whether we considered baseline, IFNA-, or dex-challenged samples
(Supplementary File 20). The only exception was for a set of differentially methylated regions in
the children of mothers exposed to objective hardship (e.g., living in a shelter, loss of electricity)
who were pregnant during or within 3 months of the 1998 Quebec Ice Storm (Cao-Lei *et al.*,
2014). In this study, ELA-associated sites were more likely to fall in windows with regulatory
potential in our sample ($\log_2$(OR) [95% CI] = 1.343 [0.715, 1.909], $p = 3.39 \times 10^{-5}$). Among
these sites, 53.85% were also detected to have methylation-dependent activity, which is slightly,
but not significantly higher than the proportion of methylation-dependent sites on the Illumina
Methylation450K chip as a whole ($\log_2$(OR) [95% CI] = 0.884 [-0.338, 2.130], p = 0.16).
Consequently, ELA-associated sites in this study were not more likely to exhibit methylation-
dependent activity than chance. We speculate that regulatory enrichment in this data set is due to
its focus on intermediately methylated CpG sites with substantial interindividual variance in

DNA methylation levels, which tends to enrich for enhancer elements. Indeed, ELA-associated
sites in this study were more strongly enriched in enhancer regions than the union set of sites in
other studies we investigated (log$_2$(OR) [95% CI] = 1.295 [0.543, 2.015], p = 6.64 x 10^{-4}).

Enrichment of regulatory activity in ELA candidata CpGs

Figure 4. Early life adversity-associated CpG sites are not enriched for mSTARR-seq regulatory
activity. Log$_2$-tranformed odds ratios from Fisher's Exact Tests for enrichment relative to the background
set of sites on each array platform, for 27 studies of early life adversity-DNA methylation level
correlations (see Supplementary File 20 for full FET results). Whiskers show standard error. Only Cao-
Lei *et al*., 2014 shows significant enrichment for regulatory activity (log$_2$(OR) [95% CI] = 1.343 [0.715,
1.909], p = 3.39 x 10^{-5}), but these sites are not more likely to exhibit methylation-dependent activity than
expected by chance (log$_2$(OR) [95% CI] = 0.884 [-0.338, 2.130], p = 0.16). For details on the source
tissue and measures of ELA, see Figure 4-figure supplement 1.

Discussion

Using mSTARR-seq, we assessed the functional role of DNA methylation across more
than 99% of CpG sites assessed by two commonly used methods to measure methylation (the
EPIC array and RRBS). Compared to our previous work, we identify a higher rate of
methylation-dependence in the present data set (47.5% of analyzed windows in the current data
set versus 10% of analyzed windows from Lea *et al.* (2018) when we apply the current pipeline
to analyze both datasets; Supplementary Files 3, 6). This difference likely stems from lower
variance between replicates in the present study, which increases power (Figure 1-figure

512   supplement 6). Thus, our findings reveal that DNA methylation often significantly attenuates
513   regulatory activity in K562 enhancer and promoter elements.
514   The results also support the idea that CpG sites identified by environment-DNA
methylation association are a mixed bag. For example, despite the pervasive assumption in the
literature and in popular science that early adversity causally impacts downstream outcomes
through persistent epigenetic modification (e.g., Dubois *et al.*, 2019), 98% of CpG sites
associated with early life adversity in the literature fell in windows with no discernable
regulatory potential in K562 cells, irrespective of their methylation status or cellular state. Our
findings suggest that many ELA-associated sites may be better treated as passive biomarkers of
exposure rather than links on the causal pathway between early life disadvantage and later life
health outcomes. However, other cell types and cellular contexts should be tested to further
evaluate this hypothesis (we used the erythroleukemic K562 cell line, as ELA-associated sites
are most commonly assessed in blood). For instance, while we included the dexamethasoné
condition here because glucocorticoid dysregulation is commonly invoked in association with
the response to early life adversity, the relationship between glucocorticoid signaling and early
life adversity is complex (Eisenberger and Cole, 2012; Koss and Gunnar, 2018) and may not be
well-modeled by acute glucocorticoid exposure. Given that the repertoire of poised and active
enhancers often differs across cell types and cellular states, expanding experimental approaches
like mSTARR-seq to other contexts can therefore serve a valuable role in prioritizing CpG sites
identified in observational studies of differential methylation (e.g., CpG sites associated with
disease).
533   Our results dovetail with work supporting the importance of CpG sites within enhancer
elements in responding to environmental perturbations (Pacis *et al.*, 2015; Husquin *et al.*, 2018).
Indeed, one of the most interesting findings from our analysis is the observation that DNA
methylation-environment interactions are widespread in the human genome. Across thousands of
genomic regions, both regulatory activity and methylation-dependent effects on regulatory
activity were only detectable upon stimulation, consistent with a model in which DNA
methylation contributes to epigenomic priming by modulating responsivity to environmental
stimuli (Kamada *et al.*, 2018; Sun and Barreiro, 2020). In support of this idea, IFNA-specific
enhancers detected in mSTARR-seq are able to nonrandomly identify collections of loci in the
genome where baseline DNA methylation also predicts the endogenous gene expression
response to flu infection (Figure 3B-C). This relationship is easier to discern in within-individual
comparisons across loci than in locus-by-locus analyses of interindividual variation. However,
we were also able to pinpoint one specific gene, *ISG15*, in which baseline DNA methylation
explains a large fraction of population variation in the gene expression response to flu (Figure
3D-E). Notably, the effect size for *ISG15* is very large, accounting for over a third of total
variance in the fold-change gene expression response to flu. Unfortunately, a comparable *in vivo*
dataset for dex challenge was unavailable to conduct parallel validation studies. However, our
results overall suggest that mSTARR-seq could help identify additional *in vivo* DNA
methylation-environment interactions, given sufficiently large sample sizes and data on the
environmental exposures of interest.
553   Applying mSTARR-seq in additional cell types may therefore help resolve whether DNA
methylation is responsible for exaggerated transcriptional responses to repeated challenges, as
previously suggested for glucocorticoid exposure in hippocampal progenitor cells (Provençal *et*
al., 2020), or whether differential methylation after pathogen infection contributes to the
emerging paradigm of "trained immunity" in innate immune cells (reviewed in Fanucchi *et al.*,

2021). More generally, while our findings agree with the idea that many differences in DNA
methylation—even extreme ones (~0% versus ~100%) like those tested here—are silent with
respect to transcription factor binding and gene expression (e.g., Maeder *et al.*, 2013; Kreibich *et*
al., 2023), they also suggest that the functional importance of DNA methylation is likely to be
underestimated without considering its interaction with the cellular environment.
563
Materials and Methods
DNA capture and mSTARR-seq
We used the DNEasy Blood and Tissue Kit (Qiagen) to extract 5 ug DNA from the
GM12878 lymphoblastoid cell line. We sheared the extracted DNA on a Covaris S2 with the
following parameters: intensity=3; duty cycle=5%, cycles/burst=200, treatment time=40 seconds,
temperature=4° C; intensifier=yes. We then performed agarose gel size selection of ~600-700 bp
DNA fragments followed by purification with the Qiaquick Gel Extraction Kit (Qiagen). We
note that we intentionally targeted longer fragments (~600-700 bp) than those targeted in (Lea *et*
al., 2018) (~300-700 bp) because our previous work showed that longer fragments are more
likely to drive regulatory activity (Lea *et al.*, 2018).
We used Roche's NimbleDesign Software to design probes to capture the following
genomic regions: (i) EPIC CpGs targeted by the Illumina Infinium MethylationEPIC microarray
(862,831 CpGs); (ii) predicted CpG cut sites from an *in silico Msp1* digest of the hg38 genome,
followed by *in silico* size selection to 100-500 bp, which together simulates the first step in
reduced representation bisulfite sequencing protocols (318,929 CpGs); (iii) 6.5 Mb centered
around the *NR3C1* glucocorticoid receptor gene; and (iv) 100,099 loci included as a control for
genomic background that were randomly distributed across the hg38 human genome, excluding
centromeres, gaps, and Ns. Although the Y chromosome was included during the design of
control loci, we excluded the Y chromosome in analyses, resulting in 99,999 control loci which
fell into 98,967 unique 600 bp windows. Of these 98,967 windows, 94,679 contained CpG sites
and 4,288 did not, reflecting the sequence composition of the human genome as a whole. We
note that a single 600 bp genomic window could simultaneously include EPIC CpGs, *Msp1* CpG
cut sites, the *NR3C1* region, or control sites (i.e., assignment of windows to these compartments
are not mutually exclusive; Figure 1-figure supplement 1).
We then captured target regions from the sheared GM12878 DNA using SeqCap EZ
Prime Choice Probes (Roche), following the SeqCap EZ HyperCap protocol version 2.3. We
performed two capture reactions. We used 6 cycles for the Pre-Capture LM-PCR program and 16
cycles for the Post-Capture LM-PCR program. For the Post-Capture LM-PCR reaction, we used
the mSTARR_primerF2 and mstarr_primerR2 primers from Lea *et al.* (2018) to prepare the
DNA for ligation into the mSTARR plasmid. The resulting captured DNA was used as input for
the mSTARR-seq protocol (Lea *et al.*, 2018), beginning at the mSTARR-seq published protocol
step "Generate linearized mSTARR backbone for large-scale cloning".
We performed 4 and 6 replicate Gibson assemblies and transformations for the first and
second capture reactions, respectively. We then sequenced each transformation replicate on a
MiSeq to measure library diversity per replicate, with the following modifications to the
published mSTARR-seq protocol: we used 25 ul Kapa HiFi HotStart ReadyMix instead of 25 ul
NEB Q5 Hot start master mix, with the following cycle program: 98 C for 45 sec, 12 cycles of
98 C for 15 seconds, 60 C for 30 seconds, and 72 C for 60 seconds, followed by a final extension
at 72 C for 5 minutes. We measured library diversity of each transformation using the ENCODE
Project's PCR bottleneck coefficient (PBC), which is the percentage of non-duplicate mapped

reads from the total number of mapped reads (Landt *et al.*, 2012) (Supplementary File 21). We
pooled the ten transformations by weighting their molarities according to the PBCs to create the
final library. We then performed 8 re-transformations to expand the pooled library.
Replicate methyltransferase reactions and parallel mock methyltransferase reactions, in
which the *M. SssI* enzyme was replaced with water, were performed in 500 ul reactions, cleaned
with Ampure beads, and then pooled. The methylated DNA library and unmethylated DNA
library were each transfected into 18 replicate T75 flasks, each containing 12 million K562s,
with Lipofectamine 3000 (ThermoFisher Scientific) following the manufacturer's instructions.
42 hours post-transfection, replicates were treated with 2,000 U/mL IFNA2b (Thermo Fisher
Scientific), 1 uM dex (Sigma-Aldrich), or vehicle control (media). 48 hours post-transfection (6
hours post-treatment), cells were harvested for mSTARR-seq sequencing library generation. At
harvest, 5×10^5 cells were aliquoted separately into Buffer RLT to measure endogenous RNA
response to treatment, and 2 million cells were aliquoted for plasmid DNA extraction to measure
input DNA in each replicate. mSTARR RNA-seq and DNA-seq libraries were generated
following Lea *et al.* (2018). To measure endogenous gene expression from K562s in each
condition, we extracted RNA from the separately aliquoted cells using the Qiagen RNEasy kit
and prepared RNA-seq libraries using the NEBNext Ultra II RNA Library Prep Kit for Illumina.
621
Comparison of mSTARR MD regulatory activity across cell types and experiments
To assess cell type-specificity of methylation-dependent regulatory activity, we
transfected the mSTARR-seq library from Lea *et al.* (2018), comprised of 1:3 sheared genomic
DNA:*MspI*-digested DNA, into the HepG2 cell line. For the HepG2 experiment, we performed
one methyltransferase reaction and one sham methyltransferase reaction. Transfections were
performed with Lipofectamine 3000 (ThermoFisher Scientific) following the manufacturer's
instructions, with reagent quantities scaled to the following per replicate: 6.9 million HepG2
cells, 46 ug of DNA, 138 ul of Lipofectamine 3000, and 180 ul of P3000 (transfection enhancer
reagent). We performed all regulatory and MD regulatory analyses of HepG2 cells, as well as
analyses of K562s from a previously published smaller mSTARR-seq experiment (Lea *et al.*,
2018), following the same bioinformatics pipeline used for our main K562 experiment, described
below. To compare against previously published STARR-seq data in A549 cells, we identified
the union set of all regulatory peaks identified across six replicate STARR-seq experiments
collected at time point 0 in Johnson *et al.* (2018). To test for significant overlap between the
current K562 results and the other three data sets, we used Fisher's exact tests against a
background set of windows tested in both the current K562 experiment and each of the other data
sets.
639
mSTARR-seq regulatory and MD regulatory analyses
mSTARR RNA-seq and DNA-seq libraries were sequenced on the Illumina NovaSeq
platform as 100 basepair, paired-end reads (Supplementary File 1). Reads were trimmed with
Trim Galore (version 0.6.4) (Krueger, 2015) to remove basepairs at the ends of reads with Phred
scores less than 20 and stretches of 2 or more basepairs that matched adapter sequences.
Trimmed reads with a minimum length of 25 basepairs were retained. We mapped reads using
bwa (version 0.7.12) (Li and Durbin, 2009) using default settings. We retained read pairs that
mapped to a single best location using the *samtools* (version 1.3.1) (Li *et al.*, 2009) package
'view' command with options -q 10 -f 0x2. We segmented the genome into 600 bp, non-
overlapping windows, and used the bedtools (version 2.25.0) (Quinlan and Hall, 2010) *coverage*

function to count the number of RNA and DNA fragments overlapping each 600 bp window. We
chose 600 bp for the genomic window size to accommodate our library's DNA fragment size
(600-700 bp; Supplementary File 1), and because using smaller windows resulted in the
identification of many adjacent windows as separate regulatory elements when they likely
represent a single true enhancer (the median length of ENCODE-annotated enhancers is 600 bp).
For each of the 3 treatment conditions (baseline, IFNA, and dex), we reduced our dataset
to windows for which at least three DNA samples in the methylated condition and three DNA
samples in the unmethylated condition (i.e., 6 DNA samples total), and three RNA samples in
either the methylated or unmethylated condition, had nonzero counts. One mSTARR RNA-seq
sample in the baseline condition (sample ID L31250) was removed from further analysis because
it had an unusually high proportion of zero counts in the testable windows; the corresponding
paired DNA sample was therefore also removed prior to analysis (Supplementary File 1). Next,
we retained only windows that showed high repeatability across DNA samples, following Lea *et*
al. (2018). In brief, for each window we calculated the pairwise difference of read abundance for
every pair of samples and created a distribution of all pairwise differences; windows were
removed if at least 25% of pairs fell outside the central 90^{th} percentile of the distribution (37,410
windows, 14.8% of the overall data set). On this reduced data set of testable windows, we
performed voom normalization using the limma (version 3.44.3) *voomWithQualityWeights*
function (Smyth, 2005; Law *et al.*, 2014), with methylation status included as a covariate in the
design. We then used the *limma* function *lmFit* to apply the following model for each window:
670
671    $$y_i = \mu + m_i\beta_1 + s_i\beta_2 * I(m = 0) + s_i\beta_3 * I(m = 1) + \varepsilon_i$$
672
where y_i is the vector of normalized counts for n = 24 samples (12 RNA and 12 DNA samples;
22 samples for the baseline condition); μ is the intercept; m_i is methylation condition (0 =
unmethylated; 1 = methylated) and β_1 its effect size; and I is an indicator variable in which β_2 and
β_3 are the effects of sample type (s_i; 0 = DNA; 1 = RNA) in the unmethylated (m = 0) and
methylated conditions (m =1), respectively. ε_i is the residual error.
To estimate the false discovery rate for identifying windows with regulatory activity (in
either the methylated or unmethylated condition, or both), we compared the observed results to
empirical null distributions generated using 100 permutations of RNA/DNA labels within each
pair of samples following Storey and Tibshirani (2003). Unlike in the real data, in which most
windows show a strong bias towards more reads in the DNA condition (because most windows
exhibit no regulatory activity, and therefore show many more input DNA reads than RNA reads),
permuted data results in relatively balanced effect sizes. Using the overall distribution of p-
values to construct the null therefore substantially inflates the number of significant windows,
but specifically because most observed windows have systematically *less* activity in the RNA
condition than the DNA condition, not more (the direction of interest). Thus, retaining all p-
values to construct the null leads to highly miscalibrated false discovery rates because the
distribution of observed values is skewed towards smaller values—because of windows with
"significantly" no regulatory activity—compared to the permuted data. To address this problem,
for each permutation, we subsampled N windows with a positive sample type beta
(corresponding to larger read counts in RNA samples versus DNA samples) in either the
methylated condition, unmethylated condition, or both. Here, N is the number of windows with a
positive sample type beta in the observed data (e.g., N = 17,461 in the baseline dataset). The p-

values from these windows were used to generate the empirical null. To define putative enhancer
elements, we used an FDR cutoff of 1%.
697   To identify methylation-dependent enhancer activity, we focused on the windows that
exhibited a positive sample type beta (i.e., exhibited more RNA reads than DNA reads) in either
the methylated or unmethylated condition (or both) and tested for significant differences between
β_2 and β_3 in the equation above (i.e., an interaction effect between RNA/DNA condition and the
methylated/unmethylated condition). To estimate the false discovery rate for identifying
methylation-dependent windows, we re-ran our main nested model on the same 100 sets of N
windows each, in this case permuting methylation condition (unmethylated versus methylated)
across sample pairs to generate the empirical null for the interaction effect of sample type (RNA
versus DNA) * methylation condition (methylated or unmethylated). To define methylation-
dependent regulatory elements, we again used an FDR cutoff of 1%.
707   To assess whether mSTARR RNA-seq and DNA-seq libraries were sequenced deeply
enough to saturate detection of unique 600 bp windows included in the formal analyses, we used
seqtk (https://github.com/lh3/seqtk) to randomly subset sequence reads from the raw fastq files
for all RNA-seq or all DNA-seq replicates in the baseline dataset. We then applied to these data
subsets the same filtering pipeline described above to generate a reduced data set of testable
windows. The results of this rarefaction analysis (shown in Figure 1-figure supplement 4) shows
that our sequencing depth is sufficient to capture all analyzable windows based on both the DNA
read filter and the RNA read filter.
715
716   *Enrichment of transcription factor binding motifs*
717   To identify enrichment of potential transcription factor binding sites in the methylated
718   condition, we used HOMER and motifs defined in the HOMER database (Heinz *et al.*, 2010) and
719   the baseline treatment dataset to compare motifs within regulatory regions in the methylated
720   condition, relative to motifs across all regulatory regions.
721   To identify potential transcription factor binding sites in windows with condition-specific
regulatory or methylation-dependent regulatory activity, we first identified regions showing
regulatory activity uniquely in the dex or IFNA conditions relative to both other conditions
(based on an FDR of 1% to define regulatory activity). We used HOMER and motifs defined in
the HOMER database to test for enrichment of transcription factor binding motifs within
windows showing condition-specific regulatory activity relative to all windows tested for
regulatory activity in that condition. We set a threshold of Bonferroni-corrected p-value < 0.01 to
identify significantly enriched binding motifs.
729
730   *Endogenous gene expression response of K562s to dex and IFNA*
731   Endogenous RNA-seq libraries from K562 cells challenged with dex or IFNA were
sequenced on an Illumina NovaSeq 6000 S1 flow cell as 100 base pair, paired-end reads. Reads
were trimmed with Trim Galore (Trim Galore version 0.6.4_dev; Cutadapt version 2.3) to
remove basepairs with a Phred score less than 20, and end sequences that matched at least two
basepairs of the adapter sequence. Only trimmed reads longer than 25 basepairs were retained.
We used the STAR package (version 2.5.0) (Dobin *et al.*, 2013) two-pass mapping to map the
filtered reads to the hg38 genome. We retained uniquely mapped reads by filtering the output
SAM file to keep reads with MAPQ = 255. We then used *htseq* (version 0.6.0) (Anders *et al.*,
2014) to quantify read counts per gene. We only retained genes that had TPM > 3 in at least 3 of

6 samples in at least one of the three conditions (baseline, IFNA, or dex). Only protein-coding
genes were retained for final analysis, resulting in a total of 10,676 testable genes.
We performed differential expression analysis separately for IFNA and dex, by subsetting
the data to the IFNA and baseline samples, or subsetting the data to the dex and baseline
samples. For each subset, we *voom* normalized the data (Law *et al.*, 2014), and used the *limma*
(Smyth, 2005) function *lmFit* to model the normalized gene expression as a function of
treatment, controlling for the percent of reads mapping to annotated features (to account for
technical variation in the efficiency of RNA purification during library prep) and methylation
condition for the associated sample. We calculated the FDR using the q-value approach of
(Storey and Tibshirani, 2003), based on an empirical null distribution that was derived from 100
permutations of treatment condition in the model.
To test whether genes were more responsive to IFNA stimulation if they were predicted
targets of IFNA-specific ISRE enhancers, we first used the bedtools *intersect* function to identify
the union set of ChIP-seq peaks for STAT1 and STAT2 (two of the three components of the
ISGF3 transcription factor that binds ISRE motifs; ENCODE accession numbers
ENCFF478XGE and ENCFF394KTR; ChIP-Seq data were not available for the third ISGF3
component, IRF9). We then reduced these regions to those showing significant, IFNA condition-
specific regulatory activity in the mSTARR-seq dataset (FDR < 1%). We used K562 enhancer-
gene links from EnhancerAtlas 2.0 (Gao *et al.*, 2016) to link the resulting IFNA-specific ISRE
enhancers to their target genes. Finally, we performed a two-sided, unpaired t-test to compare the
endogenous expression responses to IFNA for genes associated with putative IFNA-specific
ISRE enhancers relative to genes that are not associated with IFNA-specific ISRE enhancers.
762
Endogenous gene expression and methylation in human macrophages
To assess effects of DNA methylation-environment interactions on gene expression *in*
vivo, we evaluated endogenous methylation and gene expression from matched whole genome
bisulfite sequencing (WGBS) and RNA-seq data collected from human monocyte-derived
macrophages (n = 35 donors), with and without infection with the influenza A virus (IAV;
Aracena *et al.*, in press). Unsmoothed methylation counts were obtained for 19,492,906 loci in
both non-infected and IAV-infected samples (total n = 70) as described (Aracena *et al.*, in press).
We filtered loci to require coverage of ≥ 4 sequence reads in at least half of the non-infected or
IAV-infected samples. In the RNA-seq dataset for the same 35 individuals, we excluded any
genes that did not have an average RPKM > 2 in non-infected or IAV-infected samples. This
resulted in a total of 19,041,420 CpG sites and 14,122 genes used in downstream analyses.
We calculated normalization factors using calcNormFactors in edgeR (v 3.28.1)
(Robinson *et al.*, 2010) to scale the raw library sizes. We then used the voom function in limma
(v 3.42.2) (Smyth, 2005; Ritchie *et al.*, 2015) to apply the normalization factors, estimate the
mean-variance relationship, and convert raw read counts to log(CPM) values. Sequencing
batches were regressed out using ComBat from the sva Bioconductor package (v 3.34.0) (Leek *et*
al., 2012), which fit a model that also included age (mean-centered) and admixture estimates. We
subsequently regressed out age effects using limma. Individual-wise fold-change (FC) matrices
were built by subtracting non-infected counts from IAV-infected counts for each individual using
weights calculated as in (Harrison *et al.*, 2019; Aracena *et al.*, in press).
For comparison of the mSTARR-seq dataset to the dataset from Aracena *et al.* (in press),
GrCh38/hg38 coordinates were lifted over to GRCh37/hg19 using the UCSC liftOver tool. We
required a 0.95 minimum ratio of bases that remap, excluded loci that output multiple regions,

set the minimum hit size in query to 0, and set the minimum chain size in the target to 0. The
GRCh37/hg19 coordinates were used in the following analyses.
We first sought to assess, within each of the 35 individuals, the correlation between
mSTARR-seq enhancer methylation levels at baseline (i.e., in non-infected cells) and
transcriptional responses of their nearest genes to IAV. To find overlaps between enhancer
regions and CpG loci, we used the *bedtools* intersect function (v2.29.1) with the "left outer join"
option. For each 600 bp mSTARR-seq enhancer region in each individual, we calculated the
mean methylation level across all overlapping CpGs. Each enhancer was linked to its nearest
gene, with a maximum distance of 100 kb, as described above. If an enhancer had multiple
linked genes (e.g., an enhancer that overlapped more than one gene), we took the mean
transcriptional response of all linked genes. For each individual, we calculated the Pearson's
correlation coefficient (r) between the DNA methylation levels within the mSTARR-seq-defined
enhancer windows, in non-infected cells, and transcriptional responses of their linked genes to
IAV infection. We also investigated the correlation between mean mSTARR-seq enhancer
window methylation and gene expression *within* non-infected and flu-infected samples
separately.
Finally, for each mSTARR-seq enhancer-gene pair, we sought to test the extent to which
average methylation level in the enhancer in non-infected samples explained the gene's
transcriptional response to IAV, across individuals. Here, we calculated the average CpG
methylation level for each 600 bp enhancer, after excluding CpGs with methylation variance less
than 0.01 across individuals. Thus, enhancers that did not contain any CpGs with appreciable
interindividual variation in DNA methylation levels were excluded from the analysis. This
filtering step resulted in 1,382 enhancer-gene pairs for this analysis. For each enhancer-gene pair,
we calculated the Pearson's correlation coefficient (and R^2) between the average methylation
level of the enhancer and the transcriptional response of the linked gene. P-values were corrected
for multiple hypothesis testing using the q-value method in R (Storey and Tibshirani, 2003).
812
CpG methylation associated with early life adversity
We performed a literature search to identify studies for which CpG methylation
differences have been linked to adverse conditions during early life. We considered all journal
articles that contained the words "early life adversity" and "Infinium" on
https://scholar.google.com on May 10, 2021, which produced 269 results. We required that the
study evaluated CpG methylation using an Illumina Infinium array (27K, 450K, or EPIC), and
that the article provided Infinium CpG probe IDs or CpG genomic coordinates of candidate
CpGs. Articles that reported no significant CpG sites, but still reported "top" CpG sites, were
retained. Articles that performed analyses to identify differentially methylated regions (DMRs;
as opposed to CpG site-by-site analysis), and then reported individual CpG sites within the
candidate DMRs, were retained. We considered early life adversity to encompass both social and
nonsocial sources of environmental adversity (e.g., exposure to severe weather), any time from
prenatal development to 18 years of age. We did not impose criteria for cell type or subject age
at time of sampling. See Figure 4-figure supplement 1 and Supplementary File 20 for a summary
of the resulting studies.
For each study dataset, we performed a Fisher's exact test to test whether ELA candidate
sites, relative to the total set of CpG sites on the Illumina array used in that study (450K, EPIC,
or 27K array), were enriched in the mSTARR-seq regulatory windows we identified in the
baseline, dex, or IFNA condition.

Acknowledgments

We thank B.J. Nielsen, Tawni Voyles, and Tina Del Carpio for their contributions to data generation, Terrie Moffitt and the Reddy lab at Duke University for project feedback, Graham Johnson for sharing the A549 data, Courtney Karner for use of equipment, Xiang Zhou for constructive advice on statistical modeling, and two anonymous reviewers for constructive feedback on an earlier version of this manuscript. This study was supported by the National Institutes of Health (R01HD088558 to J.T. and F32HD095616 to R.A.J.), the Canadian Institutes for Advanced Research Child Brain and Development Program, a Sloan Foundation Early Career Research Fellowship to J.T., and a Foerster-Bernstein Postdoctoral Fellowship to R.A.J. High-performance computing resources were supported by the North Carolina Biotechnology Center (Grant Number 2016-IDG-1013 and 2020-IIG-2109).

Data Availability

mSTARR-seq RNA and DNA sequencing data generated in this study for K562 and HepG2 cells are available in the NCBI Sequence Read Archive (SRA; BioProject accession number PRJNA922490). K562 endogenous RNA-seq data are available in the NCBI Gene Expression Omnibus (GEO series accession GSE222643). The previously published data from (Lea *et al.*, 2018) used in this study are available in NCBI's SRA accession number SRP120556. The previously published A549 data from (Johnson *et al.*, 2018) were kindly provided by Graham Johnson. The sequencing data for the macrophage RNA-seq and WGBS datasets are available at the European Genome-Phenome Archive (EGA) under accession numbers EGAD00001008422 and EGAD00001008359, respectively. The zenodo repository at https://zenodo.org/record/7949036#.ZGZ5UnbMJq9 provides the code used for analyses, as well as track files (compatible with the UCSC genome browser) for mSTARR-seq hg38 windows with regulatory activity, and windows with methylation-dependent regulatory activity, in all three conditions (baseline, IFNA, and dex). The mSTARR-seq plasmid, *pmSTARRseq1* is available through AddGene.

Figure Supplements

Figure 1-figure supplement 1. Overlap of target genomic regions with each other.

Figure 1-figure supplement 2. mSTARR-seq library diversity.

Figure 1-figure supplement 3. Methylation levels of mSTARR-seq DNA, pre- and post-transfection.

Figure 1-figure supplement 4. Rarefaction curve showing total number of windows formally tested for regulatory activity, as a function of number of reads sequenced per DNA or RNA replicate.

Figure 1-figure supplement 5. Overlap of regulatory activity across datasets.

Figure 1-figure supplement 6. Correlations between RNA and DNA replicates.

Figure 1-figure supplement 7. Library diversity and regions of regulatory activity in HepG2

878   cells.
879
880   Figure 1-figure supplement 8. Methylation-dependent regulatory activity across datasets.
881
882   Figure 1-figure supplement 9. RNA to DNA ratios in the methylated and unmethylated replicates
883   in the baseline dataset.
884
885   Figure 1-figure supplement 10. Histograms of RNA to DNA ratios in the baseline dataset.
886
887   Figure 1-figure supplement 11. Relationship between CpG density and methylation-dependent
888   regulatory activity.
889
890   Figure 2-figure supplement 1. Filtering results across datasets.
891
892   Figure 2-figure supplement 2. Overlap of tested genomic windows across datasets.
893
894   Figure 2-figure supplement 3. Overlap of regulatory activity and effects of methylation across
895   environmental conditions.
896
897   Figure 2-figure supplement 4. *IFIT5* endogenous gene expression is responsive to IFNA
898   stimulation.
899
900   Figure 3-figure supplement 1. Across individuals, methylation in the mSTARR-seq annotated
901   enhancer chr1:1013400-1014000 predicts the *ISG15* gene expression response to flu.
902
903   Figure 4-figure supplement 1. Summary of early-life adversity (ELA) studies.
904
905   **Supplementary Files**
906   Supplementary File 1. Table summarizing the K562 mSTARR-seq DNA and RNA libraries
907   generated in this study.
908
909   Supplementary File 2. Table providing sequencing depth and CpG composition statistics for the
910   target regions (EPIC array, RRBS, *NR3C1*, and control) in the baseline dataset.
911
912   Supplementary File 3. Table of results from the model testing for methylation-dependent
913   regulatory activity in K562 cells at baseline.
914
915   Supplementary File 4. Table of enrichment results of mSTARR-seq regulatory regions for
916   ENCODE-defined chromosome states.
917
918   Supplementary File 5. Table providing sequencing depth and CpG composition statistics for the
919   baseline dataset and Lea *et al.*, 2018 dataset.
920
921   Supplementary File 6. Table of results from the model testing for methylation-dependent
922   regulatory activity in K562 cells at baseline condition using the dataset from (Lea *et al.*, 2018).
923

924 Supplementary File 7. Table of results from the model testing for methylation-dependent
925 regulatory activity in HepG2 cells at baseline.
926
927 Supplementary File 8. Table of enrichment results of transcription factor binding motifs for
928 regulatory windows with higher activity in the methylated relative to the sham condition.
929
930 Supplementary File 9. Table providing the number of 600 bp windows passing each filter for the
931 baseline, IFNA, dex, and HepG2 datasets.
932
933 Supplementary File 10. Table of results from the model testing for methylation-dependent
934 regulatory activity in K562 cells stimulated with IFNA.
935
936 Supplementary File 11. Table of results from the model testing for methylation-dependent
937 regulatory activity in K562 cells stimulated with dex.
938
939 Supplementary File 12. Table of enrichment results of transcription factor binding motifs for
940 regulatory windows specific to the IFNA treatment (<1% FDR in IFNA and >1% FDR in
941 baseline and Dex).
942
943 Supplementary File 13. Table of enrichment results of transcription factor binding motifs for
944 regulatory windows specific to the Dex treatment (<1% FDR in Dex and >1% FDR in baseline
945 and IFNA).
946
947 Supplementary File 14. Table of enrichment results of transcription factor binding motifs for
948 regulatory windows specific to the IFNA treatment (<1% FDR in IFNA and >10% FDR in
949 baseline and Dex).
950
951 Supplementary File 15. Table of enrichment results of transcription factor binding motifs for
952 regulatory windows specific to the Dex treatment (<1% FDR in Dex and >10% FDR in baseline
953 and IFNA).
954
955 Supplementary File 16. Table of results from the model testing for the K562 endogenous gene
956 expression response to IFNA.
957
958 Supplementary File 17. Table of Pearson's correlation results, within individuals, between
959 mSTARR-seq enhancer DNA methylation in non-infected macrophages and transcriptional
960 response of nearest genes upon IAV infection.
961
962 Supplementary File 18. Table of Pearson's correlation results, within individuals, between
963 mSTARR-seq enhancer DNA methylation in non-infected macrophages and gene expression
964 level in non-infected or IAV-infected macrophages.
965
966 Supplementary File 19. Table of Pearson's correlation results, across individuals, between
967 mSTARR-seq enhancer DNA methylation in non-infected macrophages and transcriptional
968 response of nearest genes upon IAV infection.
969

970   Supplementary File 20. Table of enrichment results of regulatory activity in early-life adversity
971   associated CpGs relative to regulatory activity in all CpGs on the associated Illumina array.
972
973   Supplementary File 21. Library diversity of replicate transformations using the ENCODE
974   Project's PCR bottleneck coefficient (PBC).

References

Anders S, Pyl PT, Huber W (2014) HTSeq–A Python framework to work with high-throughput sequencing data. Bioinformatics, btu638.

Aracena KA, Lin YL, Luo K, Pacis A, Gona S, Mu Z, Yotova V, Sindeaux R, Pramatarova A, Simon MM, Chen X, Groza C, Lougheed D, Gregoire R, Brownlee D, Boyce C, Pique-Regi R, Li Y, He X, Bujold D, Pastinen T, Bourque G, Barreiro LB. Epigenetic variation impacts individual differences in the transcriptional response to influenza infection. Nature Genetics. **In press**.

Aristizabal MJ, Anreiter I, Halldorsdottir T, Odgers CL, McDade TW, Goldenberg A, Mostafavi S, Kobor MS, Binder EB, Sokolowski MB (2020) Biological embedding of experience: A primer on epigenetics. Proceedings of the National Academy of Sciences, **117**, 23261-23269.

Arnold CD, Gerlach D, Stelzer C, Boryń ŁM, Rath M, Stark A (2013) Genome-wide quantitative enhancer activity maps identified by STARR-seq. Science, **339**, 1074-1077.

Cain DW, Cidlowski JA (2017) Immune regulation by glucocorticoids. Nature Reviews Immunology, **17**, 233-247.

Cao-Lei L, Massart R, Suderman MJ, Machnes Z, Elgbeili G, Laplante DP, Szyf M, King S (2014) DNA methylation signatures triggered by prenatal maternal stress exposure to a natural disaster: Project Ice Storm. PloS one, **9**, e107653.

Chaudhri VK, Dienger-Stambaugh K, Wu Z, Shrestha M, Singh H (2020) Charting the cis-regulome of activated B cells by coupling structural and functional genomics. Nature Immunology, **21**, 210-220.

Chen K, Liu J, Cao X (2017) Regulation of type I interferon signaling in immunity and inflammation: A comprehensive review. Journal of autoimmunity, **83**, 1-11.

Demetriou CA, van Veldhoven K, Relton C, Stringhini S, Kyriacou K, Vineis P (2015) Biological embedding of early‐life exposures and disease risk in humans: a role for DNA methylation. European journal of clinical investigation, **45**, 303-332.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics, **29**, 15-21.

Dolinoy DC, Weidman JR, Waterland RA, Jirtle RL (2006) Maternal genistein alters coat color and protects Avy mouse offspring from obesity by modifying the fetal epigenome. Environmental health perspectives, **114**, 567-572.

Dubois M, Louvel S, Le Goff A, Guaspare C, Allard P (2019) Epigenetics in the public sphere: interdisciplinary perspectives. Environmental Epigenetics, **5**, dvz019.

Eisenberger NI, Cole SW (2012) Social neuroscience and health: neurophysiological mechanisms linking social ties with physical health. Nature neuroscience, **15**, 669-674.

ENCODE (2012) An integrated encyclopedia of DNA elements in the human genome. Nature, **489**, 57.

Ernst J, Kellis M (2012) ChromHMM: automating chromatin-state discovery and characterization. Nature methods, **9**, 215-216.

Fanucchi S, Domínguez-Andrés J, Joosten LA, Netea MG, Mhlanga MM (2021) The intersection of epigenetics and metabolism in trained immunity. Immunity, **54**, 32-43.

Gallego Romero I, Lea AJ (2022) Leveraging massively parallel reporter assays for
evolutionary questions. arXiv e-prints, arXiv: 2204.05857.
Gao T, He B, Liu S, Zhu H, Tan K, Qian J (2016) EnhancerAtlas: a resource for enhancer
annotation and analysis in 105 human cell/tissue types. Bioinformatics, **32**, 3543-
3551.
Greenberg MV, Bourc'his D (2019) The diverse roles of DNA methylation in mammalian
development and disease. Nature reviews Molecular cell biology, **20**, 590-607.
Harrison GF, Sanz J, Boulais J, Mina MJ, Grenier J-C, Leng Y, Dumaine A, Yotova V, Bergey
CM, Nsobya SL (2019) Natural selection contributed to immunological differences
between hunter-gatherers and agriculturalists. Nature ecology & evolution, **3**, 1253-
1264.
Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass
CK (2010) Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. Molecular
cell, **38**, 576-589.
Hertzman C (1999) The biological embedding of early experience and its effects on health
in adulthood. Annals of the New York Academy of Sciences, **896**, 85-95.
Hertzman C (2012) Putting the concept of biological embedding in historical perspective.
Proceedings of the National Academy of Sciences, **109**, 17160-17167.
Hertzman C, Boyce T (2010) How experience gets under the skin to create gradients in
developmental health. Annual review of public health, **31**, 329-347.
Husquin LT, Rotival M, Fagny M, Quach H, Zidane N, McEwen LM, MacIsaac JL, Kobor MS,
Aschard H, Patin E (2018) Exploring the genetic basis of human population
differences in DNA methylation and their causal impact on immune gene regulation.
Genome biology, **19**, 1-17.
Johnson GD, Barrera A, McDowell IC, D'Ippolito AM, Majoros WH, Vockley CM, Wang X,
Allen AS, Reddy TE (2018) Human genome-wide measurement of drug-responsive
regulatory activity. Nature communications, **9**, 1-9.
Kamada R, Yang W, Zhang Y, Patel MC, Yang Y, Ouda R, Dey A, Wakabayashi Y, Sakaguchi K,
Fujita T (2018) Interferon stimulation creates chromatin marks and establishes
transcriptional memory. Proceedings of the National Academy of Sciences, **115**,
E9162-E9171.
Kazachenka A, Bertozzi TM, Sjoberg-Herrera MK, Walker N, Gardner J, Gunning R, Pahita E,
Adams S, Adams D, Ferguson-Smith AC (2018) Identification, characterization, and
heritability of murine metastable epialleles: implications for non-genetic
inheritance. Cell, **175**, 1259-1271. e1213.
Killip MJ, Fodor E, Randall RE (2015) Influenza virus activation of the interferon system.
Virus research, **209**, 11-22.
Klein JC, Agarwal V, Inoue F, Keith A, Martin B, Kircher M, Ahituv N, Shendure J (2020) A
systematic evaluation of the design and context dependencies of massively parallel
reporter assays. Nature Methods, **17**, 1083-1091.
Koss KJ, Gunnar MR (2018) Annual research review: Early adversity, the hypothalamic–
pituitary–adrenocortical axis, and child psychopathology. Journal of Child
Psychology and Psychiatry, **59**, 327-346.

Kreibich E, Kleinendorst R, Barzaghi G, Kaspar S, Krebs AR (2023) Single-molecule
footprinting identifies context-dependent regulation of enhancers by DNA
methylation. Molecular Cell.
Kribelbauer JF, Laptenko O, Chen S, Martini GD, Freed-Pastor WA, Prives C, Mann RS,
Bussemaker HJ (2017) Quantitative analysis of the DNA methylation sensitivity of
transcription factor complexes. Cell reports, **19**, 2383-2395.
Krueger F (2015) Trim Galore.
http://wwwbioinformaticsbabrahamacuk/projects/trim_galore/.
Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, Batzoglou S, Bernstein BE, Bickel
P, Brown JB, Cayting P (2012) ChIP-seq guidelines and practices of the ENCODE and
modENCODE consortia. Genome research, **22**, 1813-1831.
Law CW, Chen Y, Shi W, Smyth GK (2014) Voom: precision weights unlock linear model
analysis tools for RNA-seq read counts. Genome biology, **15**, R29.
Lea AJ, Vockley CM, Johnston RA, Del Carpio CA, Barreiro LB, Reddy TE, Tung J (2018)
Genome-wide quantification of the effects of DNA methylation on human gene
regulation. Elife, **7**, e37513.
Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD (2012) The sva package for removing
batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics, **28**, 882-883.
Lei Y, Zhang X, Su J, Jeong M, Gundry MC, Huang Y-H, Zhou Y, Li W, Goodell MA (2017)
Targeted DNA methylation in vivo using an engineered dCas9-MQ1 fusion protein.
Nature communications, **8**, 1-10.
Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler
transform. Bioinformatics, **25**, 1754-1760.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R
(2009) The sequence alignment/map format and SAMtools. Bioinformatics, **25**,
2078-2079.
Liu PZ, Nusslock R (2018) How stress gets under the skin: early life adversity and
glucocorticoid receptor epigenetic regulation. Current genomics, **19**, 653-664.
Maeder ML, Angstman JF, Richardson ME, Linder SJ, Cascio VM, Tsai SQ, Ho QH, Sander JD,
Reyon D, Bernstein BE (2013) Targeted DNA demethylation and activation of
endogenous genes using programmable TALE-TET1 fusion proteins. Nature
biotechnology, **31**, 1137-1142.
Meissner A, Gnirke A, Bell GW, Ramsahoye B, Lander ES, Jaenisch R (2005) Reduced
representation bisulfite sequencing for comparative high-resolution DNA
methylation analysis. Nucleic acids research, **33**, 5868-5877.
O'Malley RC, Huang S-sC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker
JR (2016) Cistrome and epicistrome features shape the regulatory DNA landscape.
Cell, **165**, 1280-1292.
Ostuni R, Piccolo V, Barozzi I, Polletti S, Termanini A, Bonifacio S, Curina A, Prosperini E,
Ghisletti S, Natoli G (2013) Latent enhancers activated by stimulation in
differentiated cells. Cell, **152**, 157-171.
Pacis A, Mailhot-Léonard F, Tailleux L, Randolph HE, Yotova V, Dumaine A, Grenier J-C,
Barreiro LB (2019) Gene activation precedes DNA demethylation in response to
infection in human dendritic cells. Proceedings of the National Academy of Sciences,
116, 6938-6943.

Pacis A, Tailleux L, Morin AM, Lambourne J, MacIsaac JL, Yotova V, Dumaine A, Danckaert A,
Luca F, Grenier J-C (2015) Bacterial infection remodels the DNA methylation
landscape of human dendritic cells. Genome research, **25**, 1801-1811.
Perng Y-C, Lenschow DJ (2018) ISG15 in antiviral immunity and beyond. Nature Reviews
Microbiology, **16**, 423-439.
Provençal N, Arloth J, Cattaneo A, Anacker C, Cattane N, Wiechmann T, Röh S, Ködel M,
Klengel T, Czamara D (2020) Glucocorticoid exposure during hippocampal
neurogenesis primes future stress response by inducing changes in DNA
methylation. Proceedings of the National Academy of Sciences, **117**, 23280-23285.
Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics, **26**, 841-842.
Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers
differential expression analyses for RNA-sequencing and microarray studies.
Nucleic acids research, **43**, e47-e47.
Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. bioinformatics, **26**,
139-140.
Sardina JL, Collombet S, Tian TV, Gómez A, Di Stefano B, Berenguer C, Brumbaugh J,
Stadhouders R, Segura-Morales C, Gut M (2018) Transcription factors drive Tet2-
mediated enhancer demethylation to reprogram cell fate. Cell stem cell, **23**, 727-
741. e729.
Shlyueva D, Stampfel G, Stark A (2014) Transcriptional enhancers: from properties to
genome-wide predictions. Nature Reviews Genetics, **15**, 272-286.
Smyth GK (2005) Limma: linear models for microarray data. In: Bioinformatics and
computational biology solutions using R and Bioconductor, pp. 397-420. Springer.
Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. Proc Natl
Acad Sci U S A, **100**, 9440-9445.
Sun S, Barreiro LB (2020) The epigenetically-encoded memory of the innate immune
system. Current opinion in immunology, **65**, 7-13.
Yim YY, Teague CD, Nestler EJ (2020) In vivo locus-specific editing of the neuroepigenome.
Nature Reviews Neuroscience, **21**, 471-484.
Yin Y, Morgunova E, Jolma A, Kaasinen E, Sahu B, Khund-Sayeed S, Das PK, Kivioja T, Dave
K, Zhong F (2017) Impact of cytosine methylation on DNA binding specificities of
human transcription factors. Science, **356**, eaaj2239.
1144
1145

Figure 1-figure supplement 1. Overlap of target genomic regions with each other. Upset plot
showing the degree to which 600 bp non-overlapping genomic windows are shared between the
four target genomic regions (EPIC CpGs, *Msp1* CpG cut sites, the *NR3C1* region, or control
sites). Overlap occurs because a single 600 bp genomic window can simultaneously include
EPIC CpGs, *Msp1* CpG cut sites, the *NR3C1* region, and/or control sites. This plot includes
722,472 unique windows, reflecting the set of windows containing at least 1 basepair of
sequence in the target loci.

Figure 1-figure supplement 2. mSTARR-seq library diversity. Comparison of diversity of
unique mSTARR-seq DNA and RNA fragments from the library generated in this study
(transfected into K562 cells) relative to the library published in Lea *et al*., 2018 (independently
transfected into K562 cells). Each dot represents an experimental replicate. Each box represents
the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to
the most extreme values within 1.5x of the interquartile range.

Figure 1-figure supplement 3. Methylation levels of mSTARR-seq DNA, pre- and post-
transfection. (A) Bisulfite sequencing shows that DNA methylation on the mSTARR-seq
plasmid is maintained until the end of the experiment (i.e., 48 hours after transfection), with
significantly higher methylation levels in the replicates from the methyltransferase reaction
relative to the replicates from the sham methyltransferase reaction (mean methylated = 0.885,
mean unmethylated = 0.066; unpaired t-test: t = -14.66, df = 15,124, p = 2.39 x 10^{-10}). Each dot
represents an experimental replicate. Red dots indicate post-transfection DNA samples; the
single blue dot per condition indicates pre-transfection DNA methylation levels. Methylation
estimates are based on the CpG at the position 2294, which is located in the plasmid region used
for Gibson assembly. We assessed methylation of this CpG, rather than across CpGs genome-
wide, because the genomic coverage of our bisulfite sequencing data across replicates was too
variable to perform reliable site-by-site analysis of DNA methylation levels before and after the
48 hour experiment. One sample from the dex sham reaction, L31395, shows an unexpectedly
high level of methylation, which appears to be due to an error during generation of the bisulfite
sequencing library (e.g., mislabeled tube or poor bisulfite conversion), and not the experimental
replicate of cells itself, as the mSTARR-seq RNA library (L31244) from the same replicate
clusters with the unmethylated sham replicates as expected (panel B). **(B)** The first two principal
components summarizing overall counts of mSTARR-seq reads for the dex-treated RNA
samples (i.e., the raw readout of overall regulatory activity). Each dot represents an experimental
replicate, with red and black indicating sham and methylated replicates, respectively. Overall
regulatory activity of sample L31244 (indicated by arrow) clusters with the sham replicates as
expected, suggesting that this replicate was indeed transfected with sham-treated mSTARR-seq
DNA.

Figure 1-figure supplement 4. Rarefaction curve showing total number of windows
formally tested for regulatory activity, as a function of number of reads sequenced per
DNA or RNA replicate. Sequencing reads from the (A) DNA replicates or (B) RNA replicates
of the baseline dataset were rarefied to the values shown on the x-axis before running the data
processing steps and applying the filtering criteria described in the Materials and Methods for the
full data set. Dashed vertical lines represent the mean number of sequenced reads per DNA
replicate (mean [SD] = 30.375 million [3.335 million]) or RNA replicate (mean [SD] = 31.712
million [8.194 million]) in the full baseline dataset. These analyses show that our sequencing
effort saturated the number of formally analyzable windows based on either our criteria for
inclusion based on DNA library sequencing depth or RNA library sequencing depth.

Overlap of regulatory activity with other datasets

Figure 1-figure supplement 5. Overlap of regulatory activity across datasets. Regulatory
regions (in either the unmethylated sham condition, the methylated condition, or both) identified
via mSTARR-seq in this study significantly overlap with: K562 regulatory regions (in either the
unmethylated sham or methylated condition, or both) from a previously generated mSTARR-seq
dataset reanalyzed with our pipeline (Lea et al., 2018) ($\log_2$(OR) [95% CI] = 6.212 [6.086,
6.440], p < 1.0 x 10^{-300}); regulatory regions (in either the unmethylated sham or methylated
condition, or both) from an mSTARR-seq experiment in HepG2 liver cells ($\log_2$(OR) [95% CI] =
3.534 [3.381, 3.684], p = 5.21 x 10^{-307}); and regulatory regions from a conventional STARR-seq
experiment (i.e., an unmethylated condition) in A549 lung epithelial cells (Johnson et al., 2018)
($\log_2$(OR) [95% CI] = 2.451 [2.442, 2.461], p < 1.0 x 10^{-300}). Bars represent 95% confidence
intervals.

Figure 1-figure supplement 6. Correlations between RNA and DNA replicates. Pearson
correlations (r) of raw counts between RNA replicates (A-D) and between DNA replicates (E-H)
within the windows we formally analyzed for enhancer activity in the baseline dataset reported
here and in Lea *et al*., 2018, following a uniform data processing pipeline. All replicate pairs
(both RNA and DNA, in both sham and methylated conditions) in the baseline dataset show
correlations ≥ 0.89, demonstrating replicate reproducibility comparable to other STARR-seq
studies (e.g., Klein et al., 2020). For RNA libraries, replicates in the baseline dataset are more
correlated than in the Lea *et al*., 2018 dataset (RNA replicates: baseline mean r = 0.926; Lea *et*
al., 2018 mean r = 0.347), although DNA replicates show similar inter-replicate consistency
(baseline mean r = 0.992; *Lea et al*., 2018 mean r = 0.991).

Figure 1-figure supplement 7. Library diversity and regions of regulatory activity in
HepG2 cells. (A) Comparison of diversity of unique mSTARR-seq DNA and RNA fragments
from the library published in Lea *et al*., 2018 (transfected into K562 cells) versus the same
library transfected into HepG2 cells in this study. Each dot represents an experimental replicate.
Each box represents the interquartile range, with the median value depicted as a horizontal bar.
Whiskers extend to the most extreme values within 1.5x of the interquartile range. **(B)**
mSTARR-seq regulatory activity in HepG2 cells is strongly enriched in ENCODE-defined
enhancers (indicated in blue) and some classes of promoters, and depleted in repressed and
repetitive states.

Figure 1-figure supplement 8. Methylation-dependent regulatory activity across datasets.
Effects of methylation on regulatory activity estimated in this study in the baseline dataset are
consistent with methylation effects in K562s estimated from a previously generated mSTARR-
seq dataset (Lea *et al*., 2018) and with methylation effects estimated in HepG2 liver cells (Lea *et*
al., 2018: Pearson's r = 0.534 for 1250 windows with FDR < 1% in both data sets, R^2 = 0.286, p
= 3.19 x 10^{-93}; HepG2: Pearson's r = 0.526 for 511 windows with FDR < 1% in both data sets,
R^2 = 0.277, p = 8.87 x 10^{-38}). Each dot represents a 600 bp regulatory window identified, in
either the sham or methylated states, in both datasets (FDR < 1%; note that not all regulatory
windows show significant methylation dependence [MD]). Dashed lines are the best fit lines. In
all cases, negative effect sizes correspond to reduced activity in the methylated condition and
positive effect sizes correspond to increased activity in the methylated condition.

Figure 1-figure supplement 9. RNA to DNA ratios in the methylated and unmethylated
replicates in the baseline dataset. Mean RNA (in counts per million) to DNA ratios
formethylated replicates (x-axis) versus unmethylated replicates (y-axis). A constant of 0.5 was
added to the initial raw counts to ensure no denominator values were 0. Each dot represents a
600 bp window that was formally tested for enhancer activity (**A**), exhibited significant
regulatory activity (**B**), or exhibited significant methylation-dependent regulatory activity (**C**) in
the baseline dataset. Solid diagonal lines represent y = x, and dashed lines represent the best fit
lines. As expected, 600 bp windows tend to show higher RNA to DNA ratios in the
unmethylated condition relative to the methylated condition.

Figure 1-figure supplement 10. Histograms of RNA to DNA ratios in the baseline dataset.
The x-axis represents the $\log_2$(mean RNA [in counts per million] to DNA ratios) for the baseline
dataset. A constant of 0.5 was added to the initial raw counts to prevent 0 from being in the
denominator. Lower values on the x-axis indicate windows showing lower regulatory activity.

Figure 1-figure supplement 11. Relationship between CpG density and methylation-
dependent regulatory activity. CpG-dense mSTARR-seq regulatory regions are more likely to
be repressed by DNA methylation (positive y-axis value; Spearman's rho = 0.370, p = 9.865 x
10^{-121}; n = 3,721 regions with mSTARR-seq regulatory activity). Each dot represents a 600 bp
window that showed significant regulatory activity (FDR < 1%). Red and blue dots represent
regulatory windows where methylation-dependent activity was or was not detected, respectively.
The dashed line represents the best fit line.

Figure 2-figure supplement 1. Filtering results across datasets. Each of the 5 datasets began
with 5,051,776 600 bp genomic windows. For each of the 5 datasets, we reduced the dataset to
windows that had nonzero counts in at least three DNA samples in the methylated condition *and*
three DNA samples in the unmethylated condition (i.e., 6 DNA samples total; "DNA filter"). We
then reduced the dataset to windows that had nonzero counts in at least three RNA samples in
either the methylated *or* unmethylated condition ("RNA filter"). Finally, we retained only
windows that showed high repeatability across DNA samples, following (Lea *et al*., 2018)
("DNA repeatability"). Numbers correspond to million windows that passed or failed each filter
for which the arrow points to. Note that one mSTARR RNA-seq sample in the baseline condition
[sample ID L31250] was removed from further analysis because it had an unusually high
proportion of zero counts in the testable windows; we therefore also removed the corresponding
paired DNA sample prior to analysis). See Supplementary File 9 for the precise window numbers
corresponding to the plots.

Figure 2-figure supplement 2. Overlap of tested genomic windows across datasets. Upset
plots showing the degree to which 600 bp non-overlapping genomic windows are shared
between five datasets (baseline null, IFNA, dex, HepG2, and Lea *et al*., 2018; all datasets were
analyzed following the same pipeline). Note that the same input library was used in Lea *et al*.,
2018 and the HepG2 experiment, which differs from the library used here for baseline, dex, and
IFNA experiments.

Figure 2-figure supplement 3. Overlap of regulatory activity and effects of methylation across environmental conditions. (A) Regulatory regions in the baseline condition are highly likely to retain regulatory activity upon challenge with IFNA or dex (IFNA $\log_2$(OR) [95% CI] = 8.639 [8.499, 8.812], $p < 1.0 \times 10^{-300}$; dex $\log_2$(OR) [95% CI] = 9.640 [9.483, 9.776], $p < 1.0 \times 10^{-300}$). Regulatory regions also significantly overlap between IFNA- and dex-challenged cells ($\log_2$(OR) [95% CI] = 8.554 [8.420, 8.698], $p < 1.0 \times 10^{-300}$). (B) Regulatory windows identified in two environmental conditions tend to share significant effects of DNA methylation on regulatory activity (i.e., interaction effects between methylation and regulatory activity) across the two environmental conditions (baseline and IFNA $\log_2$(OR) [95% CI] = 6.345 [5.673, 7.102], $p < 7.33 \times 10^{-239}$; baseline and dex $\log_2$(OR) [95% CI] = 5.982 [5.599, 6.384], $p < 1.0 \times 10^{-300}$; IFNA and dex $\log_2$(OR) [95% CI] = 5.576 [5.123, 6.055], $p < 4.70 \times 10^{-266}$).

**Figure 2-figure supplement 4. *IFIT5* endogenous gene expression is responsive to IFNA
stimulation.** Tracks show non-normalized, raw read pile-ups of endogenous *IFIT5*
(ENSG00000152778) gene expression in either the unmethylated (open circle) or methylated
(filled circle) condition, with all y-axis maximums set to 100. The replicates for endogenous
gene expression shown here are the same replicates used for measuring regulatory activity shown
in Figure 2A. *IFIT5* is only detectably expressed after IFNA stimulation (note that the difference
in peak heights between the IFNA-stimulated unmethylated and methylated conditions is
because the plot shows raw reads; there is no effect of methylation treatment on endogenous
IFIT5 gene expression after normalization for library size: $p = 0.489$).

Figure 4-figure supplement 1. Summary of early-life adversity (ELA) studies. Colored
rectangles indicate ranges of age at adversity. Circles represent mean ages at sample collection
and are colored according to the tissue type used to measure methylation levels. Ranges and
standard errors of ages at sample collection, when reported, are indicated by gray rectangles and
horizontal black solid lines, respectively. See Supplementary File 20 for results of Fisher's exact
tests assessing enrichment of ELA-associated CpGs for regulatory activity.