1

5   **Distinct brain network features predict internalizing and externalizing traits in children**

6   **and adults**

8   Yueyue Lydia Qu [1], Jianzhong Chen [2,3,4,5], Angela Tam [2,3,4,5], Leon Qi Rong Ooi [2,3,4,5,6], Elvisha

9   Dhamala [1,7,8], Carrisa Cocuzza [1], Connor Lawhead [1], B. T. Thomas Yeo [2,3,4,5,6,9], Avram J.

10   Holmes [1,10,11]

12   [1] Department of Psychology, Yale University, New Haven, CT, USA

13   [2] Center for Sleep and Cognition, National University of Singapore, Singapore, Singapore

14   [3] Center for Translational MR Research, National University of Singapore, Singapore, Singapore

15   [4] Department of Electrical and Computer Engineering, National University of Singapore,

16   Singapore, Singapore

17   [5] N.1 Institute for Health & Institute for Digital Medicine (WisDM), National University of

18   Singapore, Singapore, Singapore

19   [6] Integrative Sciences and Engineering Programme (ISEP), National University of Singapore,

20   Singapore, Singapore

21   [7] Institute of Behavioral Sciences, Feinstein Institutes for Medical Research, Manhasset, USA

22   [8] Kavli Institute for Neuroscience, Yale University, New Haven, USA

23   [9] Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, MA,

24   USA

25   [10] Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

26   [11] Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway, NJ, USA

Abstract
Internalizing and externalizing traits are two distinct classes of behaviors in psychiatry.
However, whether shared or unique brain network features predict internalizing and
externalizing behaviors in children and adults remain poorly understood. Using a sample of
2262 children from the Adolescent Brain Cognitive Development (ABCD) study and 752 adults
from the Human Connectome Project (HCP), we show that network features predicting
internalizing and externalizing behavior are, at least in part, dissociable in children, but not in
adults. In ABCD children, traits within internalizing and externalizing behavioral categories are
predicted by more similar network features concatenated across task and resting states than
those between different categories. We did not observe this pattern in HCP adults. Distinct
network features predict internalizing and externalizing behaviors in ABCD children and HCP
adults. These data reveal shared and unique brain network features accounting for individual
variation within broad internalizing and externalizing categories across developmental stages.

Introduction

A classic distinction in child and adolescent psychiatry has been the study of "internalizing" and "externalizing" behaviors[1]. These two broad classes of psychopathology were first proposed by T.M. Achenbach from a factor analysis of symptoms in children and adolescents with psychiatric illness[2]. Internalizing behaviors are internally directed towards the individual and manifest in their extreme form as sadness, withdrawal, somatic complaints, and anxiety, while externalizing behaviors are directed towards the external environment and involve disruptive, aggressive, impulsive, and defiant behaviors[3]. The expressions of internalizing and externalizing behaviors exhibit cross-generational associations between parents and children[4–6]. These behaviors have also been linked with reduced school engagement and an increased risk for suicide attempts in childhood and adolescence[7–9], as well as worse work performance and lower cognitive abilities in adulthood[10,11]. However, the neural underpinnings associated with internalizing and externalizing behaviors across distinct developmental stages remain poorly understood.

Throughout development, functional connectivity patterns within and between large-scale brain networks can predict individual differences in cognition[12], impulsivity[13] and psychiatric symptoms[14,15]. While individual-level variability in the functioning of large-scale brain networks can predict individual differences within broad categories of cognition, personality and mental health in both children and adults[16,17], macroscale patterns of brain functioning are dynamic across the lifespan[18–20]. The transition from childhood through adolescence to adulthood reflects critical neurodevelopmental stages characterized by a protracted period of synaptic pruning, intracortical myelination, cortical thinning, and functional network segregation[18,21]. Therefore, it is unclear if the specific brain-behavior relationships observed in childhood mirror those identified in adulthood. Furthermore, although shared network features account for individual variation within broad classes of behavior[16], individual-specific patterns of functional network connections may predict even finer-grained categories, such as internalizing and externalizing behaviors. Here, we aimed to examine the extent that functional network-based predictors of internalizing versus externalizing behaviors were similar across a large sample of children and their parents. We further tested whether such patterns can be observed in an independent sample of young adults.

In the present study, we predicted internalizing and externalizing measures of psychopathology in a sample of children (and their parents) from the Adolescent Brain Cognitive Development (ABCD) study[22] using children's functional connectivity patterns across four brain states: resting-state, monetary incentive delay (MID) task[23], stop signal task (SST)[24] and

emotional N-Back task[25]. We further explored functional connectivity predictors of internalizing

and externalizing behavior in an independent cohort of young adults from the Human

Connectome Project (HCP)[26], using resting-state fMRI connectivity matrices. Both single- (KRR)

and multi-kernel ridge regression (multiKRR) models revealed network-based features that were

predictive of behaviors within the same category were more correlated with each other than with

those across different categories in ABCD children and parents, while KRR models showed a

lack of categorical distinction in HCP adults. Moreover, predictive network features were distinct

across the two samples. These results support internalizing and externalizing behaviors as

distinct factors of psychopathology and suggest that brain-based predictive features may

change across the lifespan.

<div align="center">

Results

</div>

ABCD Results

We used fMRI data acquired across three task states, including monetary-incentive

delay (MID), stop-signal task (SST) and N-back, as well as resting state fMRI from N=11,875

typically-developing children (ABCD 2.0.1 release[22]). Our analyses considered 33 dimensional

measures from the available mental health assessments collected from child participants and

their parents[27], comprised of 15 measures of internalizing problems, 10 measures of

externalizing problems, 2 measures of thought problems and 6 measures of attention problems

(Supplementary Table 1). The final analytical sample consisted of n=2,262 unrelated children

who passed fMRI quality control and had complete data (see Methods).

Multi-kernel ridge regression predicts most behavioral measures

We defined 400 cortical and 19 subcortical regions-of-interest (ROIs) based on Schaefer

Parcellation[28,29] and computed a 419 by 419 functional connectivity (FC) matrix for each brain

state. Following prior work[16], we used multi-kernel ridge regression (multiKRR) models to

predict each behavioral measure from child-specific FC matrices concatenated across brain

states. To evaluate predictive accuracy, we performed nested cross-validation procedures with

120 folds (see Methods). Pearson's correlation between predicted and actual behavioral scores

and coefficient of determination (COD; see Supplementary method S3) were used as accuracy

metrics. Statistical significance of prediction accuracy was assessed by permutation testing.

Prediction accuracies -- given by Pearson's correlation -- of the models trained on

children's functional connectivity data are shown in Fig. 1A (for behavioral predictions in ABCD

children) and Fig. 1B (for behavioral predictions in ABCD parents). Most behavioral measures

were predicted better than chance after FDR correction (q<0.05), except for child somatic

complaints and somatic problems, parent intrusive behavior, parent ADHD problems and parent

inattention (see Methods). Parent ADHD problems became significantly predicted after FDR

correction when COD was used as the accuracy metric. Prediction accuracies were broadly

stable across both metrics (see Supplementary Fig. 1 for COD results). Notably, these findings

demonstrate that patterns of FC specific to each child can significantly predict their parent's self-

reported internalizing and externalizing behaviors (Fig. 1B).

Figure 1. Cross-validated prediction performance using the multi-kernel ridge regression (multiKRR) model, using functional connectivity matrices concatenated across four brain states (resting state, MID, SST and N-back) from children's neuroimaging data to predict (A) parent-reported child behavior and (B) self-reported parent behavior. Prediction performance was calculated as the mean Pearson's correlation between observed and predicted values across 120 cross-validation folds for each behavioral measure from the ABCD dataset. For each boxplot, the top and bottom edges represent upper and lower quartiles of correlation coefficient (*r*) distributions, and the horizontal lines mark the corresponding median. Outliers are plotted as circles and were defined as data points outside of the interquartile range. The whiskers extend to the most extreme data points not considered as outliers. Asterisks (*) denote above-chance significance after correcting for multiple comparisons (FDR $q<0.05$).

Predictive brain network features are more similar within behavioral categories

There is broad consistency in the brain network features predictive of mental health-

relevant traits[16]. Here, we sought to determine if internalizing and externalizing behaviors

exhibited unique predictive network markers in childhood. At each cross-validation fold, we

quantified "feature importance" (i.e., how important a given network-based predictor was to the

model) of each interregional FC edge predicting each behavior using Haufe-transformed (see

Methods) predictive feature weights[30], yielding a 419 by 419 predictive feature matrix for each

behavior and for each brain state.

Next, we analyzed whether predictive feature weights computed from multiKRR model

outputs were more similar among behaviors within than between categories (Fig. 2). The

predictive feature weight vector for each behavioral measure was averaged across all four brain

states and correlated with all other measures. Focusing on each of the four internalizing and

externalizing categories (Child Internalizing, Child Externalizing, Parent Internalizing and Parent

Externalizing), the difference between mean correlation within each category ("within-category

mean correlation") and mean correlation with all other three categories ("between-category

mean correlation") was computed 10000 times and used to generate a null distribution of mean

differences (Fig. 3; see Methods). Mean within-category correlations of predictive feature

weights were significantly higher than mean between-category correlations (FDR qs≤0.002; Fig.

3). The above analyses were rerun using KRR models using only resting-state fMRI and yielded

similar results (Supplementary Fig. 2-3). Notably, the similarity pattern of predictive feature

weights across behavioral measures was highly correlated with the similarity pattern of these

behavioral measures on the behavioral level (Supplementary Fig. 4; r=0.97).

Figure 2. Predictive-network features are similar within behavioral categories. Pearson's

correlation (*r*) of predictive feature weights between all pairs of behavioral measures

significantly predicted by multiKRR models in the ABCD study. Behavioral measures from the

same behavioral categories are grouped together. Warmed colors indicate stronger positive

correlations of the mean predictive feature weights between a pair of behavioral measures,

indicating that these behavioral measures were predicted by similar functional connectivity

patterns.

Figure 3. Correlations of predictive feature weights computed from multiKRR outputs were significantly stronger across behavioral measures within the same category than between different categories. Differences between within- and between-category mean correlations for child and parent internalizing and externalizing categories were significantly greater than the null distributions (FDR qs ≤0.002). Correlation values were converted to z-scores using Fisher's r-to-z transformation prior to averaging. Histograms display null distributions of mean differences generated through 10000 permutations with shuffled behavioral labels. Dashed lines represent observed mean differences for each of the four categories.

Distinct brain network features in children predict internalizing and externalizing

behaviors in both children and their parents

Prior work suggests the presence of shared network features across broad categories of

mental health[16]. Our analyses revealed unique parcel-level FC profiles predicting distinct

aspects of psychopathology. Next, we examined the extent that some predictive-network

features may be shared across behavioral categories. Predictive feature matrices were

averaged across all behavioral measures within each category, resulting in 32 predictive feature

matrices (one for each behavioral category and each brain state). To limit the number of

multiple comparisons, predictive feature weights were averaged within and between 18

networks (following the 17-network partition in Yeo et al., 2011[31] plus one subcortical network[28])

at each permutation. Permutation testing was performed on mean predictive feature weights

from each of the resulting 171 unique network blocks. We also conducted a conjunction analysis

to extract the predictive feature weights that were not only statistically significant but also

exhibited consistent directionality (positive or negative) across all brain states, and then

averaged these predictive feature weights across all brain states (Fig. 4A). These analyses

yielded predictive feature weights that are both shared across behavioral measures within a

category and across brain states (Fig. 4B). Predictive feature weights were summed across

each row in Fig. 4A and plotted on brain surface in Fig. 4C for the positive weights and Fig. 4D

for the negative weights. These figures reveal that both shared and unique FC patterns predict

distinct behavioral categories in both children and their parents.

To examine the extent to which these predictive features are similar across behavioral

categories in children and their parents, we next calculated the proportion of overlapping

network blocks which significantly predicted each pair of behaviors (Fig. 4E). Two network

blocks were counted as overlapping if sums of predictive feature weights within these network

blocks exhibited consistent directionality. Of note, the observed predictive features were not fully

distinct across children and parents. The largest proportion of overlap was 0.66 between parent

internalizing and externalizing categories, while the lowest proportion of overlap was 0.42

between child internalizing and parent externalizing categories. Proportions of overlap between

other four category pairs ranged from 0.50 to 0.60, demonstrating the presence of both common

and distinct patterns of predictive-network features across categories. As one example, the

proportion of network blocks that exhibit the same directionality across child and parent

internalizing categories was 0.54.

Figure 4. Shared and unique functional network features predict internalizing and externalizing

behaviors in children and their parents. (A) Matrices of predictive feature weights, averaged

across all behavioral measures within each child and parental internalizing and externalizing

categories, and averaged across all brain states. Only weights that were statistically significant

and that exhibited the same directionality across all brain states are shown. Rows and columns:

predictive weights based on FC estimates of all pairwise cortical regions. For visualization

purposes, all predictive feature weights were divided by their standard deviation. (B) Predictive

feature weights averaged based on network assignment in panel (A). (C) Positive predictive

feature weights summed across rows of panel (A) for each cortical region. A more positive value

indicates that stronger functional connectivity associated with a given cortical parcel predicts

higher behavioral scores in a behavioral category. (D) Negative predictive feature weights

summed across rows of panel (A) for each cortical region. A more negative value indicates that

weaker functional connectivity associated with a given cortical region predicts higher behavioral

scores in a behavioral category. In both panels (C) and (D), the color of each cortical region

indicates the percentile of predictive feature weights among 400 regions. (E) The 2D grid

displays the proportion of network blocks that exhibit the same directionality across each pair of

child behavioral categories relative to the behavioral category represented by each column.

Here, each within- and between-network block was coded as 1, 0 or -1 depending on whether

sum of predictive feature weights within that block is greater than, equal to or lesser than 0,

resulting in an 18 by 18 matrix for each behavioral category. The number of network blocks

having the same non-zero entries across both matrices associated with each pair of behavioral

categories was counted and divided by the total number of non-zero significantly predictive

network blocks.

Single-kernel ridge regression predicted most behavioral measures in adults

To examine brain-based predictive network features in adults, we used resting-state

fMRI data from the HCP WU-Minn S1200 sample (n=752) and analyzed 18 dimensional

measures from the Achenbach Self-Report[32]. There were 8 measures of internalizing problems,

5 measures of externalizing problems, 1 measure of thought problems and 4 measures of

attention problems (Supplementary Table 2; see Methods). All analysis steps were performed

as above, except that single-kernel ridge regression (KRR) models were used to predict each

behavioral measure from subject-specific resting-state FC due to the lack of task fMRI data in

the HCP. Given that the HCP was not collected across different sites, we implemented 60

random initiations of 10-fold nested cross-validation.

Prediction accuracies – given by Pearson's correlation – of the KRR models across all

behaviors are shown in Supplementary Fig. 6. Although most behavioral measures were

predicted better than chance after FDR correction ($q<0.05$), only two out of eight behavioral

measures under the internalizing category survived FDR correction (Supplementary Fig. 6).

When COD was used as the accuracy measure, only withdrawn, aggressive behavior, and

attention problems reached better-than-chance accuracy after FDR correction (Supplementary

Fig. 7).

Predictive brain network features are similar across behavioral categories in adulthood

As above, we examined similarity patterns of predictive feature weights calculated from

KRR model outputs across behaviors within and between different categories (Fig. 5). In

contrast to the ABCD analyses, predictive feature weights were highly correlated across

categories (Fig. 5; Supplementary Fig. 9). We then conducted a permutation test similar to the

ABCD analyses, focusing on adult internalizing and externalizing categories. Mean within-

category correlations of predictive feature weights were not significantly different from mean

between-category correlations (FDR qs>0.12; Fig. 5B). These results suggest that predictive

network features associated with internalizing and externalizing behavior in adults are broadly

consistent between behavioral categories. Although network features predicting intrusive

behavior were weakly correlated with those predicting other measures, it is not surprising given

the weak correlations between intrusive behavior and other measures on the behavioral level.

The observed similarity pattern of predictive feature weights across behaviors was moderately

correlated with the similarity pattern of these measures on the behavioral level (Supplementary

Fig. 8; r=0.59).

275
276

Figure 5. Predictive brain network features are similar across conceptually-linked behavioral categories in adulthood. (A) The correlation matrix displays Pearson's correlation *r* of predictive feature weights between all pairs of behavioral measures significantly predicted in the HCP study. Behavioral measures are grouped within associated behavioral categories. Higher intensity colors indicate higher positive (red) and negative (blue) correlations of the mean predictive feature weights between a pair of behavioral measures. (B) Differences between mean correlations of predictive feature weights across behavioral measures within the same category and between different categories were not significantly greater than the null

distributions in the adult internalizing and externalizing categories (FDR qs>0.12). Correlation

values were converted to z-scores using Fisher's r-to-z transformation prior to averaging.

Graphing conventions are similar to that of Figure 3.

Predictive feature weights are largely distinct across ABCD and HCP datasets

To investigate if functional network features predicting internalizing and externalizing

behaviors are distinct across development, we assessed the similarity of predictive network

features computed from KRR model outputs between ABCD and HCP datasets. We found that

FC features predicting internalizing and externalizing problems in ABCD children and HCP

adults were only weakly correlated (Fig. 6). We then ran permutation tests to compare the

difference in the mean correlation within each category and the mean correlation between each

category and the corresponding category in the other age group. As the adult internalizing

category contained only two significantly predicted measures, we only focused on child

internalizing and externalizing and adult externalizing behaviors. The difference was significantly

greater than its null distribution (FDR qs≤0.0146) for the two child categories but failed to reach

statistical significance for the adult externalizing category (FDR q=0.0601; Fig. 6B). From the

predictive feature matrices associated with child and adult internalizing and externalizing

categories, we computed the proportion of overlapping network blocks which significantly

predicted each pair of categories (Fig. 6C). Proportions of overlap were distinctly higher for pairs

of behavioral categories within the same dataset than between the two datasets. These results

suggest that although shared brain network features account for individual variation within broad

categories of internalizing and externalizing problems in childhood, functional network predictors

may change throughout the lifespan, exhibiting distinct fingerprints across developmental

stages.

Despite the broad distinction between FC patterns predicting internalizing and

externalizing behaviors across these two datasets, shared predictive patterns may still be

present within select edges. Here, the shared predictive network features associated with both

child and adult internalizing categories primarily involved the default, control and visual

networks, while the shared predictive network features associated with child and adult

externalizing primarily involved somato/motor, ventral and dorsal attention networks

(Supplementary Fig. 10).

Figure 6. Distinct functional network features predict internalizing and externalizing behaviors in

children from the ABCD study and adults from the HCP study. (A) The correlation matrix

displays the Pearson's correlation r of predictive feature weights between all pairs of behavioral

measures significantly predicted by KRR models in the ABCD and HCP studies. Measures from

the same behavioral category are grouped together. Colors indicate positive (red) and negative

(blue) correlations of the mean predictive feature weights between behavioral measures and

populations. (B) Predictive feature weights associated with child internalizing and externalizing

categories were significantly more correlated within categories than with the corresponding adult

categories (FDR qs≤0.0186). The permutation test was not applicable for the adult internalizing

category because only one correlation can be computed between two behavioral measures in

the category. Correlation values were converted to z-scores using Fisher's r-to-z transformation

prior to averaging. Graphing conventions are similar to that of Figure 3. (C) The 2D grid displays

the proportion of network blocks that exhibit the same directionality across each pair of child and

adult behavioral categories relative to the behavioral category represented by each column.

Discussion

In this study, we first used estimates of individual-specific, functional connectivity from a

large, diverse sample of children to predict internalizing- and externalizing-related behavioral

measures in children and their parents. Predictive feature weights were more correlated across

behaviors within the same categories than with those from different categories. Of note, our

analyses revealed that brain data specific to each child can be used to predict self-reported

internalizing and externalizing behaviors in their parents. We repeated these analyses in an

independent sample of healthy young adults and the opposite pattern was observed, where

predictive feature weights were similarly correlated across distinct mental health-linked

behavioral categories. Moreover, predictive feature weights associated with internalizing- and

externalizing-related behavioral measures were distinct across children and adults, suggesting

that brain-based predictors of internalizing and externalizing behaviors may change across the

lifespan.

Internalizing and externalizing symptoms reflect distinct factors across various mental

disorders, irrespective of demographic and collection method[33–37]. Predictive network features

are similar across behaviors within the broad categories of mental health[16]. Although large-scale

networks can be mechanistically informative for studying neurocognitive processes[38,39] and

psychiatric phenotypes[15,40–42], the similarity of whole-brain FC patterns predicting measures of

internalizing and externalizing behavior has not been directly assessed. Through the use of both

KRR and multiKRR models[16,43], we were able to predict most mental health measures in

children and their parents from children's resting-state functional connectome. Here, we

demonstrated that the whole-brain patterns of functional connectivity in children can be used to

predict internalizing and externalizing measures in their parents. Our results highlight that the

predictive utility of functional connectomes may extend beyond the individual, and provide a

robust entry point for future work on shared environmental and contextual factors, broader

behavioral patterns within family systems, and/or the heritability of internalizing and

externalizing traits.

Consistent with prior work by Chen et al. 2022 (which used the same dataset), we

observed that predictive features are generally similar across measures of internalizing and

externalizing behaviors. However, above and beyond this broad pattern of similarity, predictive

feature weights were more correlated within than between behavioral categories in ABCD

children. These findings are consistent with theoretical models that consider internalizing and

externalizing behaviors as distinct constructs of psychopathology under a general

psychopathology *p* factor[44,45]. Behavioral measures associated with different categories are

characterized by both common and distinct network predictors in children. On average, higher

behavioral scores in both child internalizing and externalizing categories were predicted by more

positive FC between default, control and limbic networks, between somato/motor and salience

networks and more negative FC between default and somato/motor networks. Beyond these

shared features, there was substantial heterogeneity in the FC patterns predicting internalizing-

and externalizing-related behaviors. These results align with previous neuroimaging studies

implicating frontoparietal[46,47], default[47–49], salience[49,50], limbic[49] and somato/motor[49,51] network

disruptions across psychiatric disorders.

Contrary to the similarity pattern observed in ABCD children, mean correlations of

predictive feature weights across all pairs of behavioral measures within internalizing and

externalizing categories were not significantly different from mean correlations between different

categories in HCP adults. Our findings suggest that diffuse functional network patterns may

predict a more general psychopathology factor in adults, while more specific FC patterns may

differentially predict behaviors associated with specific categories of psychopathology in

children. One consideration is that KRR models used in HCP analyses reached better-than-

chance predictive accuracy for only two out of eight measures assigned to adult internalizing

category. This may have biased the results for the permutation test assessing statistical

significance of mean correlation differences within and between adult internalizing and

externalizing categories. In addition to different correlation patterns across behavioral categories

between the two datasets, we also observed distinct FC features predicting same categories of

internalizing and externalizing behaviors in children compared to adults. Weak correlations of

predictive feature weights associated with internalizing and externalizing behavior across the

two samples may be attributable to development of functional network organization from

childhood through adolescence and then adulthood[19,20,52–54]. However, such differences may

also be attributable to site differences between the two collection efforts. Of note, our

interpretations are limited by the cross-sectional nature of the available data. Future work

should further characterize the longitudinal trajectories of brain development and associated

brain-based predictions across the lifespan. Another limitation of our study is that we did not test

our models separately in each sex. Previous studies have suggested brain-based predictive

models often fail to generalize across sexes[55], and future work should test sex-/gender- specific

models of behavior[56].

Taken together, our study found that predictive network features cluster within the same

categories of internalizing and externalizing behavior in ABCD children. Intriguingly, the utility of

brain-based predictive models in children extended to capture behaviorally relevant signals in

their parents. Finally, although most behaviors were predicted better than chance in children

and adults, analyses revealed distinct network predictors across datasets. Future work will

benefit from the longitudinal study of common and distinct brain-based predictive features

across childhood, adolescence, and adulthood.

Methods

Participants

11,875 typically-developing children and their parents across 21 sites in the United

States participated in the ABCD study at baseline (ABCD release 2.0.1). The final analytical

sample consisted of 2,262 unrelated children who passed strict preprocessing quality control,

had complete fMRI data across all brain states and complete scores across all behavioral

measures. Similar to Chen et al., 2022, we combined the 22 ABCD sites into 10 "site-

categories" to reduce sample size variability across sites (Supplementary Table 5). Subjects

within the same site were also in the same site-category. Detailed demographic information can

be found in Supplementary Table 6.

1,206 healthy adults participated in the HCP study (HCP S1200 Data Release). After

pre-processing quality control of imaging data, participants were filtered from Li's set of 953

participants[57] based on the availability of a complete set of structural and resting-state fMRI

scans, as well as all behavioral scores of interest. Our main analysis comprised 752 adult

participants, who fulfilled all selection criteria[17]. Detailed demographic information can be found

in Supplementary Table 7.

Neuroimaging

Data acquisition

For the ABCD study, all T1w images and fMRI data was acquired using protocols

harmonized across three 3 tesla(T) scanner platforms (i.e., Phillips, Siemens Prisma and

General Electric 750) at 21 sites. Twenty minutes of resting-state fMRI data, consisting of four

5-minute runs, was collected from each ABCD child participant. For each of the three tasks

(MID, SST and N-Back)[23–25], fMRI data was acquired over two runs with 2.4mm isotropic

resolution with a TR of 800ms. The structural T1 scans were acquired with 1mm isotropic

resolution with a TR of 2500ms. For full details of imaging acquisition can be found elsewhere[58].

The fMRI data in the HCP data was acquired using an optimized protocol with 2mm

isotropic resolution and a TR of 700ms. Each HCP subject goes through one structural MRI

session and two fMRI sessions. Each fMRI session consists of two 15-minute resting-state

scans with opposite phase encoding directions (L/R and R/L). The structural T1 scans were

acquired using 0.7mm isotropic resolution and a TR of 2400ms. Full details of the acquisition

protocol can be found elsewhere[26].

Data processing

444   Minimally preprocessed T1w images[59] in the ABCD study were further processed using

445 FreeSurfer v5.3.0[60–65]. The cortical surface meshes were then registered a common spherical

446 coordinate system[62,63]. Subjects who failed recon-all QC were subsequently excluded[59]. The

447 minimally preprocessed fMRI data[59] were subsequently processed in the following manner. The

448 first four frames were removed[59]. Slice time correction was performed with the FSL library[66,67].

449 Motion correction was performed using rigid body translation and rotation with the FSL package.

450 The resulting fMRI images were then aligned with the processed T1w images[68] with FsFast

451 (https://surfer.nmr.mgh.harvard.edu/fswiki/FsFast), and only runs with registration costs less

452 than 0.6 were retained. Framewise displacement (FD)[67] and voxel-wise differentiated signal

453 variance (DVARS)[69] were computed by fsl_motion_outliers. Volumes with FD > 0.3 mm or

454 DVARS > 50, along with one volume before and two volumes after, were flagged as outliers. A

455 bandstop filter was applied to remove respiratory pseudomotion[70]. Uncensored segments of

456 data having fewer than 5 contiguous volumes were also flagged as outliers and censored[71].

457 Runs with more than half of the volumes flagged as outliers were discarded. Participants with

458 less than 4 minutes of data for each fMRI state (rest, MID, N-Back, SST) were excluded from

459 further analysis. Nuisance regressors, including global signal, six motion correction parameters,

460 averaged ventricular signal, averaged white matter signal, and their temporal derivatives (18

461 regressors in total), were regressed out of the fMRI time series from the unflagged volumes.

462 Data were interpolated across censored frames[72], band-pass filtered at 0.009 Hz$\leq$f$\leq$0.08 Hz,

463 projected onto FreeSurfer fsaverage6 surface space, and smoothed using a 6mm full-width half

464 maximum kernel.

465   For the HCP study, minimally preprocessed T1w images[73] went through bias- and

466 distortion- correction using the *PreFreeSurfer* pipeline and registered to MNI space. Cortical

467 surface reconstruction was conducted using FreeSurfer v5.2 using recon-all adapted for high-

468 resolution images. The reconstructed surface meshes were then registered to the Conte69

469 surface template[74]. After preprocessing, the fMRI data were corrected for gradient-nonlinearity-

470 induced distortions. The fMRI time series in each frame were then realigned to the single-band

471 reference image to correct for subject motion using rigid body transformation[67,75] with FSL. The

472 resulting single-band image underwent spline interpolation to correct for distortions and was

473 then registered to the T1w image[68]. Native fMRI volumes went through nonlinear registration to

474 the MNI space and mapped to the standard CIFTI grayordinate coordinate space. Further

475 details about the preprocessing and processing pipelines of structural and functional images

476 can be found elsewhere[73].

<u>Functional connectivity</u>

We used 400 cortical regions of interest[29] (ROIs) and 19 subcortical ROIs[28]. Functional

connectivity (FC) was measured by Pearson's *r* correlations between the mean time series of

each pair of ROIs. Censored frames were ignored when computing functional connectivity. In

the ABCD study, the average FC matrix across all runs in each subject from each state (rest,

MID, N-back, SST) was used for subsequent analyses. To match processing across resting and

task states, task activations were not regressed from the task-state data. For the HCP study, the

average FC matrix across all runs in each subject was only computed from the resting state and

used for subsequent analyses.

Measures of internalizing and externalizing behaviors

We included 25 dimensional measures of internalizing and externalizing in our analyses,

selected from all available mental health relevant assessments taken from child participants and

their parents[27] in the ABCD study. This consisted of 15 internalizing measures and 10

externalizing measures. Thought and attention problems are related to both internalizing and

externalizing psychopathology[76,77]. Accordingly, we included 2 measures of thought problems

and 6 measures of attention. Participants without available data across all behavioral measures

were excluded from analysis. The complete list of the included variables can be found in Tables

S1 and S2. Behavioral measures were grouped into four categories: Internalizing, Externalizing,

Thought Problems and ADHD Problems for both children and their parents, resulting in eight

behavioral categories in total (Supplementary Table 1).

In data from the HCP, we analyzed 18 dimensional measures of internalizing,

externalizing, thought and attention problems from the Achenbach Self-Report (ASR)

questionnaire, resulting in four behavioral categories (Supplementary Table 2).

Statistical analysis

Consistent with prior work[16], we used multi-kernel ridge regression (multiKRR) with l_2

regularization to predict each behavioral measure from participant-specific FC matrices across

all brain states (rest, MID, N-back, SST) jointly in the ABCD study. Behavioral measures in the

HCP study were predicted from resting-state FC using kernel ridge regression (KRR) with l_2

regularization. Details about KRR and multiKRR models can be found in the Supplement (see

Supplementary methods S1 and S2). Age and mean FD were entered as covariates. Both

models assume that participants with more similar FC patterns have more similar behavioral

measures. Models were implemented with nested cross-validation procedures similar to Ooi et

al., 2022. Head motion (mean FD and DVARS) were regressed from each behavioral measure

prior to cross-validation.

In the ABCD analyses, we performed leave-3-site-clusters-out nested cross-validation

for each behavioral measure. At each fold, a different set of 3 site-categories served as the test

set, and the remaining 7 site-categories were used as the training set, resulting in 120 folds in

total. In the HCP analyses, we implemented 60 random initiations of 10-fold nested cross-

validation. Participants from the same family were assigned to either training or testing sets and

were never split across training and test sets in any cross-validation fold.

Across both datasets, model and regularization parameters were estimated from the

training set at each fold. The estimated parameters were then applied to the unseen participants

from the test set and evaluated for accuracy by both correlating predicted and actual

measures[78], and by coefficient of determination (COD). To assess whether model prediction

performed better than chance, statistical significance of prediction accuracy was assessed by a

permutation test whereby the entire cross-validation procedure was rerun on behavior measures

randomly reshuffled across participants in each dataset. Care was taken to avoid shuffling

between families or sites.

Model Interpretation

To interpret the predictive importance of each FC feature, we used an approach from

Haufe and colleagues (2014) to transform predictive feature weights associating each FC edge

to the behavioral measure. Predictive feature weight was computed by the covariance between

the predicted behavioral measure and the FC edge. This resulted in a 419 x 419 predictive

feature matrix for each brain state and each behavioral measure. A positive (or negative)

predictive feature weight indicates that higher FC predicts greater (or lower) behavioral values.

Statistical significance of these predictive feature weights was tested with permutation tests and

corrected for multiple comparison using FDR (q<0.05). To reduce the number of multiple

comparisons, predictive feature weights were averaged within and between 18 large-scale

functional networks[28,29] before conducting the permutation test.

Acknowledgements

Data used in the preparation of this article were obtained, in part, from the Adolescent Brain

Cognitive Development SM (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive

(NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children aged

9-10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the

National Institutes of Health and additional federal partners under award numbers

U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037,

U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134,

U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038,

U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of

supporters is available at abcdstudy.org/federal-partners.html. A list of participating sites and a

complete list of the study investigators can be found at abcdstudy.org/consortium_members/.

ABCD consortium investigators designed and implemented the study and/or provided data but

did not necessarily participate in the analysis or writing of this report. Additional data were

provided, in part, by the Human Connectome Project, WU-Minn Consortium (Principal

Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH

Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the

McDonnell Center for Systems Neuroscience at Washington University. This manuscript reflects

the views of the authors and may not reflect the opinions or views of the NIH or the ABCD and

HCP consortia investigators.

Funding Sources

This work was supported by the National Institute of Mental Health (R01MH120080 and

R01MH123245 to AJH). This work was also supported by the following awards to BTTY: the

Singapore National Research Foundation (NRF) Fellowship (Class of 2017), the NUS Yong Loo

Lin School of Medicine (NUHSRO/2020/124/TMR/LOA), the Singapore National Medical

Research Council (NMRC) LCG (OFLCG19May-0035), the NMRC STaR (STaR20nov-0003)

and the Singapore Ministry of Health (MOH) Centre Grant (CG21APR1009). This work was also

supported by the following awards to ED: the Kavli Institute for Neuroscience at Yale University

(Postdoctoral Fellowship for Academic Diversity), the Feinstein Institutes for Medical Research

Advancing Women in Science and Medicine (Career Development Award and Barbara Zucker

Emerging Scientist Award). Any opinions, findings and conclusions or recommendations

expressed in this material are those of the authors and do not reflect the views of the funders.

Financial Disclosures

All authors reported no biomedical financial interests or potential conflicts of interest.

References

1. Achenbach, T. M. The Child Behavior Profile: I. Boys aged 6-11. *J Consult Clin Psychol*
46, 478–488 (1978).
2. Achenbach, T. M. The classification of children's psychiatric symptoms: a factor-analytic
study. *Psychol Monogr* **80**, 1–37 (1966).
3. Achenbach, T. M. *Manual for the child behavior checklist/ 4-18 and 1991 profile.*
(University of Vermont, Department of Psychiatry, 1991).
4. Hammen, C., Shih, J. H. & Brennan, P. A. Intergenerational transmission of depression:
test of an interpersonal stress model in a community sample. *J Consult Clin Psychol* **72**,
511–522 (2004).
5. Conger, R. D., Neppl, T., Kim, K. J. & Scaramella, L. Angry and aggressive behavior
across three generations: a prospective, longitudinal study of parents and children. *J*
Abnorm Child Psychol **31**, 143–160 (2003).
6. Kim, H. K., Capaldi, D. M., Pears, K. C., Kerr, D. C. R. & Owen, L. D. Intergenerational
Transmission of Internalising and Externalising Behaviours Across Three Generations:
Gender-Specific Pathways. *Crim Behav Ment Health* **19**, 125 (2009).
7. Duprey, E. B., Oshri, A. & Liu, S. Developmental pathways from child maltreatment to
adolescent suicide-related behaviors: The internalizing and externalizing comorbidity
hypothesis. *Dev Psychopathol* **32**, 945–959 (2020).
8. Commisso, M. *et al.* Childhood externalizing, internalizing and comorbid problems:
distinguishing young adults who think about suicide from those who attempt suicide.
Psychol Med (2021) doi:10.1017/S0033291721002464.
9. Olivier, E., Morin, A. J. S., Langlois, J., Tardif-Grenier, K. & Archambault, I. Internalizing
and Externalizing Behavior Problems and Student Engagement in Elementary and
Secondary School Students. *J Youth Adolesc* **49**, 2327–2346 (2020).
10. Papachristou, E. & Flouri, E. The codevelopment of internalizing symptoms, externalizing
symptoms, and cognitive ability across childhood and adolescence. *Dev Psychopathol*
32, 1375–1389 (2020).
11. Narusyte, J., Ropponen, A., Alexanderson, K. & Svedberg, P. Internalizing and
externalizing problems in childhood and adolescence as predictors of work incapacity in
young adulthood. *Soc Psychiatry Psychiatr Epidemiol* **52**, 1159 (2017).
12. Sripada, C. *et al.* Prediction of neurocognition in youth from resting state fMRI. *Molecular*
Psychiatry 2019 25:12 **25**, 3413–3421 (2019).
13. Shannon, B. J. *et al.* Premotor functional connectivity predicts impulsivity in juvenile
offenders. *Proc Natl Acad Sci U S A* **108**, 11241–11245 (2011).
14. Uddin, L. Q. *et al.* Salience Network–Based Classification and Prediction of Symptom
Severity in Children With Autism. *JAMA Psychiatry* **70**, 869 (2013).
15. Lake, E. M. R. *et al.* The Functional Brain Organization of an Individual Allows Prediction
of Measures of Social Abilities Transdiagnostically in Autism and Attention-
Deficit/Hyperactivity Disorder. *Biol Psychiatry* **86**, 315–326 (2019).
16. Chen, J. *et al.* Shared and unique brain network features predict cognitive, personality,
and mental health scores in the ABCD study. *Nature Communications 2022 13:1* **13**, 1–
17 (2022).
17. Ooi, L. Q. R. *et al.* Comparison of individualized behavioral predictions across
anatomical, diffusion and functional connectivity MRI. *Neuroimage* **263**, 119636 (2022).
18. Sydnor, V. J. *et al.* Neurodevelopment of the association cortices: Patterns, mechanisms,
and implications for psychopathology. *Neuron* **109**, 2820–2846 (2021).
19. Hwang, K., Hallquist, M. N. & Luna, B. The Development of Hub Architecture in the
Human Functional Brain Network. *Cerebral Cortex* **23**, 2380–2393 (2013).

20. Dong, H. M., Margulies, D. S., Zuo, X. N. & Holmes, A. J. Shifting gradients of
macroscale cortical organization mark the transition from childhood to adolescence. *Proc*
Natl Acad Sci U S A **118**, (2021).
21. Bethlehem, R. A. I. *et al.* Brain charts for the human lifespan. *Nature 2022* 1–11 (2022)
doi:10.1038/s41586-022-04554-y.
22. Volkow, N. D. *et al.* The conception of the ABCD study: From substance use to a broad
NIH collaboration. *Dev Cogn Neurosci* **32**, 4–7 (2018).
23. Knutson, B., Westdorp, A., Kaiser, E. & Hommer, D. FMRI visualization of brain activity
during a monetary incentive delay task. *Neuroimage* **12**, 20–27 (2000).
24. Logan, G. D. On the ability to inhibit thought and action: A users' guide to the stop signal
paradigm. (1994).
25. Cohen, A. O., Conley, M. I., Dellarco, D. V & Casey, B. J. The impact of emotional cues
on short-term and long-term memory during adolescence. *Proceedings of the Society for*
Neuroscience. San Diego, CA. November (2016).
26. Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An Overview.
Neuroimage **80**, 62 (2013).
27. Barch, D. M. *et al.* Demographic, physical and mental health assessments in the
adolescent brain and cognitive development study: Rationale and description. *Dev Cogn*
Neurosci **32**, 55–66 (2018).
28. Fischl, B. *et al.* Whole brain segmentation: automated labeling of neuroanatomical
structures in the human brain. *Neuron* **33**, 341–355 (2002).
29. Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
Functional Connectivity MRI. *Cereb Cortex* **28**, 3095–3114 (2018).
30. Haufe, S. *et al.* On the interpretation of weight vectors of linear models in multivariate
neuroimaging. *Neuroimage* **87**, 96–110 (2014).
31. Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by
intrinsic functional connectivity. *J Neurophysiol* **106**, 1125–1165 (2011).
32. Achenbach, T. M. & Rescorla, L. A. *Manual for the ASEBA adult forms & profiles.*
(University of Vermont, Research Center for Children, Youth, & Families, 2003).
33. Krueger, R. F., Markon, K. E., Patrick, C. J., Benning, S. D. & Kramer, M. D. Linking
antisocial behavior, substance use, and personality: an integrative quantitative model of
the adult externalizing spectrum. *J Abnorm Psychol* **116**, 645–666 (2007).
34. Eaton, N. R. *et al.* An invariant dimensional liability model of gender differences in mental
disorder prevalence: evidence from a national sample. *J Abnorm Psychol* **121**, 282–288
(2012).
35. Achenbach, T. M. & Edelbrock, C. S. Psychopathology of Childhood.
https://doi.org/10.1146/annurev.ps.35.020184.001303 **35**, 227–256 (2003).
36. Kessler, R. C. *et al.* Development of Lifetime Comorbidity in the World Health
Organization World Mental Health Surveys. *Arch Gen Psychiatry* **68**, 90–100 (2011).
37. Ringwald, W. R., Forbes, M. K. & Wright, A. G. C. Meta-analytic tests of measurement
invariance of internalizing and externalizing psychopathology across common
methodological characteristics. *Journal of psychopathology and clinical science* **131**,
847–856 (2022).
38. Rosenberg, M. D., Finn, E. S., Scheinost, D., Constable, R. T. & Chun, M. M.
Characterizing Attention with Predictive Network Models. *Trends Cogn Sci* **21**, 290–302
(2017).
39. Rosenberg, M. D. *et al.* A neuromarker of sustained attention from whole-brain functional
connectivity. *Nature Neuroscience 2016 19:1* **19**, 165–171 (2015).
40. Satterthwaite, T. D. *et al.* Connectome-wide network analysis of youth with Psychosis-
Spectrum symptoms. *Molecular Psychiatry 2015 20:12* **20**, 1508–1515 (2015).

41. Pornpattananangkul, N., Leibenluft, E., Pine, D. S. & Stringaris, A. Association Between
Childhood Anhedonia and Alterations in Large-scale Resting-State Networks and Task-
Evoked Activation. *JAMA Psychiatry* **76**, 624–633 (2019).

42. Karcher, N. R., O'Brien, K. J., Kandala, S. & Barch, D. M. Resting-State Functional
Connectivity and Psychotic-like Experiences in Childhood: Results From the Adolescent
Brain Cognitive Development Study. *Biol Psychiatry* **86**, 7–15 (2019).

43. He, T. *et al.* Deep neural networks and kernel regression achieve comparable accuracies
for functional connectivity prediction of behavior and demographics. *Neuroimage* **206**,
116276 (2020).

44. Brislin, S. J. *et al.* Differentiated nomological networks of internalizing, externalizing, and
the general factor of psychopathology ('p factor') in emerging adolescence in the ABCD
study. *Psychol Med* **52**, 3051–3061 (2022).

45. Caspi, A. *et al.* The p Factor: One General Psychopathology Factor in the Structure of
Psychiatric Disorders? *Clin Psychol Sci* **2**, 119 (2014).

46. Marek, S. & Dosenbach, N. U. F. The frontoparietal network: function, electrophysiology,
and importance of individual precision mapping. *Dialogues Clin Neurosci* **20**, 133 (2018).

47. Baker, J. T. *et al.* Functional connectomics of affective and psychotic pathology. *Proc Natl
Acad Sci U S A* **116**, 9050–9059 (2019).

48. Doucet, G. E. *et al.* Transdiagnostic and disease-specific abnormalities in the default-
mode network hubs in psychiatric disorders: A meta-analysis of resting-state functional
imaging studies. *European Psychiatry* **63**, (2020).

49. Huang, C. C. *et al.* Transdiagnostic and Illness-Specific Functional Dysconnectivity
Across Schizophrenia, Bipolar Disorder, and Major Depressive Disorder. *Biol Psychiatry
Cogn Neurosci Neuroimaging* **5**, 542–553 (2020).

50. Sheffield, J. M. *et al.* Transdiagnostic associations between functional brain network
integrity and cognition. *JAMA Psychiatry* **74**, 605–613 (2017).

51. Kebets, V. *et al.* Somatosensory-Motor Dysconnectivity Spans Multiple Transdiagnostic
Dimensions of Psychopathology. *Biol Psychiatry* **86**, 779–791 (2019).

52. Grayson, D. S. & Fair, D. A. Development of large-scale functional networks from birth to
adulthood: a guide to neuroimaging literature. *Neuroimage* **160**, 15 (2017).

53. Galván, A. & Tottenham, N. Adolescent Brain Development. *Developmental
Psychopathology* 1–36 (2016) doi:10.1002/9781119125556.DEVPSY218.

54. Casey, B. J., Galván, A. & Somerville, L. H. Beyond simple models of adolescence to an
integrated circuit-based account: A commentary. *Dev Cogn Neurosci* **17**, 128 (2016).

55. Dhamala, E., Yeo, B. T. T. & Holmes, A. J. One Size Does Not Fit All: Methodological
Considerations for Brain-Based Predictive Modeling in Psychiatry. *Biol Psychiatry* **93**,
717–728 (2023).

56. Dhamala, E. *et al.* Brain-based predictions of psychiatric illness-linked behaviors across
the sexes. *Biol Psychiatry* (2023) doi:10.1016/J.BIOPSYCH.2023.03.025.

57. Li, J. *et al.* Global signal regression strengthens association between resting-state
functional connectivity and behavior. *Neuroimage* **196**, 126–141 (2019).

58. Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD) study: Imaging
acquisition across 21 sites. *Dev Cogn Neurosci* **32**, 43 (2018).

59. Hagler, D. J. *et al.* Image processing and analysis methods for the Adolescent Brain
Cognitive Development Study. *Neuroimage* **202**, 116091 (2019).

60. Fischl, B., Liu, A. & Dale, A. M. Automated manifold surgery: constructing geometrically
accurate and topologically correct models of the human cerebral cortex. *IEEE Trans Med
Imaging* **20**, 70–80 (2001).

61. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical surface-based analysis. I. Segmentation
and surface reconstruction. *Neuroimage* **9**, 179–194 (1999).

62. Fischl, B., Sereno, M. I., Tootell, R. B. H. & Dale, A. M. High-resolution intersubject
averaging and a coordinate system for the cortical surface. *Hum Brain Mapp* **8**, 272
(1999).
63. Fischl, B., Sereno, M. I. & Dale, A. M. Cortical surface-based analysis. II: Inflation,
flattening, and a surface-based coordinate system. *Neuroimage* **9**, 195–207 (1999).
64. Ségonne, F. *et al.* A hybrid approach to the skull stripping problem in MRI. *Neuroimage*
22, 1060–1075 (2004).
65. Ségonne, F., Pacheco, J. & Fischl, B. Geometrically accurate topology-correction of
cortical surfaces using nonseparating loops. *IEEE Trans Med Imaging* **26**, 518–529
(2007).
66. Smith, S. M. *et al.* Advances in functional and structural MR image analysis and
implementation as FSL. *Neuroimage* **23 Suppl 1**, (2004).
67. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved Optimization for the Robust
and Accurate Linear Registration and Motion Correction of Brain Images. *Neuroimage* **17**,
825–841 (2002).
68. Greve, D. N. & Fischl, B. Accurate and Robust Brain Image Alignment using Boundary-
based Registration. *Neuroimage* **48**, 63 (2009).
69. Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious
but systematic correlations in functional connectivity MRI networks arise from subject
motion. *Neuroimage* **59**, 2142–2154 (2012).
70. Gratton, C. *et al.* Removal of high frequency contamination from motion estimates in
single-band fMRI saves data without biasing functional connectivity. *Neuroimage* **217**,
116866 (2020).
71. Gordon, E. M. *et al.* Generation and Evaluation of a Cortical Area Parcellation from
Resting-State Correlations. *Cerebral Cortex* **26**, 288–303 (2016).
72. Power, J. D. *et al.* Methods to detect, characterize, and remove motion artifact in resting
state fMRI. *Neuroimage* **84**, 320–341 (2014).
73. Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome
Project. *Neuroimage* **80**, 105–124 (2013).
74. Van Essen, D. C., Glasser, M. F., Dierker, D. L., Harwell, J. & Coalson, T. Parcellations
and hemispheric asymmetries of human cerebral cortex analyzed on surface-based
atlases. *Cereb Cortex* **22**, 2241–2262 (2012).
75. Jenkinson, M. & Smith, S. A global optimisation method for robust affine registration of
brain images. *Med Image Anal* **5**, 143–156 (2001).
76. Jacob, C. *et al.* Internalizing and externalizing behavior in adult ADHD. *Atten Defic*
Hyperact Disord **6**, 101–110 (2014).
77. Arslan, İ. B., Lucassen, N., van Lier, P. A. C., de Haan, A. D. & Prinzie, P. Early
childhood internalizing problems, externalizing problems and their co-occurrence and
(mal)adaptive functioning in emerging adulthood: a 16-year follow-up study. *Soc*
Psychiatry Psychiatr Epidemiol **56**, 193 (2021).
78. Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using
patterns of brain connectivity. *Nature Neuroscience 2015 18:11* **18**, 1664–1671 (2015).
799