Identification of novel interactors of PP2A-B'ζ

Ahmed Elshobaky[1,2], Cathrine Lillo[1], Kristian Persson Hodén[3], Amr R.A. Kataya[1,4]

[1]University of Stavanger, Centre for Organelle Research, Faculty of Science and Technology, N-4036 Stavanger, Norway
[2]Botany Department, Faculty of Science, Mansoura University, 35516 Mansoura, Egypt
[3]Department of Plant Biology, Uppsala BioCenter, Linnéan Center for Plant Biology, Swedish University of Agricultural Sciences, P.O. Box 7080, 75007, Uppsala, Sweden.
[4]Department of Biological Sciences, University of Calgary, Calgary, T2N 1N4, Canada.

Author for correspondence: Amr R.A. Kataya
Email: amr.kataya@ucalgary.ca

Figures: 1-5, Supplementary Figure S1 and S2.
Tables: Supplementary Tables 1-5.

Abstract

Protein phosphatase 2A (PP2A) is a heterotrimeric conserved serine/threonine phosphatase complex including a catalytic, scaffolding, and regulatory subunit. Three A Subunits, 17 B subunits, and five C subunits are encoded by the Arabidopsis genome, allowing 255 possible PP2A holoenzyme combinations. The regulatory subunits are crucial for substrate specificity and PP2A complex localization and are classified into B, B', and B'' non-related families in land plants. In *Arabidopsis*, the close homologs B'η, B'θ, B'γ, and B'ζ are further classified into a subfamily of B' called B'η. Previous studies suggest a role of mitochondrial targeted PP2A subunit (B'ζ) in energy metabolism. Potentially, the PP2A-B'ζ holoenzyme is involved in the regulation of the mitochondrial succinate/fumarate translocator or affects enzymes involved in energy metabolism. To investigate this hypothesis, the interaction between PP2A-B'ζ and enzymes involved in the mitochondrial energy flow was investigated using bimolecular fluorescence complementation in tobacco and onion cells. Interaction of B'ζ subunit was proven with the Krebs cycle proteins Succinate-fumarate translocator (mSFC1), Malate dehydrogenase (mMDH2), and Aconitase (ACO3). Additional interacting candidates were identified by comparing the total phosphoproteomes of wild type and B'ζ mutants: the mitochondrial regulator Arabidopsis pentatricopeptide repeat 6 (PPR6) and the two metabolic enzymes Phosphoenolpyruvate carboxylase (PPC3) and Phosphoenolpyruvate carboxykinase (PCK1). Overall, this study identified potential PP2A substrates and highlighted the role of PP2A in regulating energy metabolism in mitochondria.

Introduction

Protein post-translational modifications (PTMs) are covalent additions or modification of chemical groups to a protein after its synthesis. PTMs add an additional level of regulation, on top of gene expression and mRNA translation. Phosphorylation is an abundant PTM and a main regulator of cell signaling that is involved in a vast number of cellular processes (von Stechow 2016). Reversible phosphorylation is achieved through the balance between protein kinases, which act by adding a phosphate group to amino acids in the protein chain, and phosphatases, which remove phosphorylated groups from proteins (Bheri et a., 2021). Attesting to the importance of protein phosphorylation events, the *Arabidopsis thaliana* genome contains around 940 kinases and 150 phosphatases (Li and Liu, 2021). Protein phosphorylation and dephosphorylation processes regulate multiple downstream cellular processes such as, e.g., transcription, energy metabolism and development, by affecting protein attributes such as, e.g., localization, interactions and activity (von Stechow 2016; Li & Liu, 2021).

Meta-analysis of phosphoproteomics data revealed that serine (Ser) is the most abundantly phosphorylated residue in plants (~85%), followed by threonine (Thr; ~15%) (van Wijk, 2014). Consequently, one of the major phosphatase families in plants is composed of the serine/threonine-specific phosphoprotein phosphatases (PPPs). Among the PPPs, PP1 and PP2A are the largest groups in *Arabidopsis* (Uhrig et al., 2013, Bheri et a., 2021). Plant PP2A enzymes have been shown to be involved in various functions such as regulation of light signaling, hormone signaling, metabolism, other PTMs and defense signaling (Uhrig et al., 2013; Farkas et al., 2007). PP2A enzymes are heterotrimeric proteins composed of a catalytic C subunit with regulatory (B) and scaffolding (A) subunits. The regulatory subunits are essential for substrate specificity and localization of the complex. Complexity of PP2A enzymes is achieved through combination of a low number of C subunits with A and B subunits (Matre et al., 2009). Thus, despite having only five C subunits, *Arabidopsis* can theoretically attain over 250 PP2A variants (three A subunits and 17 B subunits). The plant B family is further divided into the B, B' and B'' subfamilies, of which the B' gene family contains nine members: α, β, γ, δ, ε, ζ, η, θ and κ (Bheri, 2021). We here focus on the B' subunit ζ (Z).

The plant immune system needs to be under tight negative regulation to avoid over-activation and autoimmunity. Protein kinases and phosphatases have been suggested as key players in plant immunity regulation, and some emergent studies have implicated PP2A in this role (Segonzac et al., 2014; Kataya et al., 2015b; Rasool et al., 2014). More specifically, all four members of the B'η subgroup (η, γ, θ and ζ) were shown to be negative regulators of plant PAMP-triggered immunity. Increased resistance to the virulent bacterial strain *Pseudomonas syringae* DC3000 was observed in knockout mutants of *b'θ*, *b'ζ* and *b'η*. Furthermore, the PP2A-B'ζ and PP2A-B'η complexes control the activation of cell-surface pattern recognition receptors by regulating the phosphostatus of the co-receptor BAK1 (Segonzac et al., 2014). The *b'γ* knockdown, on the other hand, displayed constitutive defense responses, and, in line with this, was implicated in resistance to a variety of infectious agents; aphids, hemibiotrophic bacteria and necrotrophic fungi (Rasool et al., 2014; Trotta et al., 2011).

Notably, the constitutive defense activation observed in the *b'γ* mutant seems to be explained by an involvement of the PP2A-B'γ complex in methionine recycling. By performing BiFC experiments, Rahikainen et al. (2016) first showed that B'γ interacts with indole glucosinolate methyl transferases. Further, B'γ was found to negatively regulate the metabolism of glucosinolates, which are important secondary metabolites, acting as repellents for aphids and microbial plant pathogens in *Arabidopsis*. Increased build-up of specific compounds such as 4-methoxy-indol-3-yl-methyl glucosinolate is therefore a likely reason for the elevated defense responses of the *b'γ* knockout.

Mitochondria are double membrane-bound eukaryotic organelles. Reversible phosphorylation involving mitochondrial phosphatases has been largely neglected, although mitochondria are increasingly recognized as a hub for cell signaling, and several mitochondrial phosphatases have been reported to play vital functions. Nonetheless, evidence for the localization and activities of these reported phosphatases, as well as their functions and their mitochondrial translocation mechanism, are still poorly understood (Lim et al.,2016).

Quantitative phosphoproteomics is a method that can be used to identify and characterise PTM sites in proteins (Urban, 2022). When performed on samples collected from suitable mutants, it can reveal novel substrates of kinases and phosphatases. Performing tandem mass tag (TMT) labeling prior to fragmentation increases throughput by enabling multiplex quantification of relative differences between samples. In a typical phosphoproteomics

workflow, proteins are first fragmented into peptides by trypsin digestion, then separated by liquid chromatography (LC) and finally quantified by mass spectrometry (MS). Phosphorylated sites are identified using bioinformatic methods applied to the MS data.

PP2A regulatory subunit (B'ζ) was reported to target mitochondria (Matre et al. 2009), and when B'ζ is mutated, plants not only show negative regulation of plant innate immunity, but also have a sugar dependence phenotype (Kataya et al., 2015b; Segonzac et al., 2014). This indicates a defect in energy metabolism in the *pp2a-b'ζ* mutant. As seedlings of PP2A-B'θ, the nearest homolog of PP2a-B'ζ, have shown a defect in peroxisomal β-oxidation, and is assumed to block the supply of succinate to the mitochondrial Krebs cycle during early seedling establishment (Kataya et al., 2015a), we investigated a putative role of PP2A-B'ζ in the regulation of energy flow in mitochondria. We studied *in vivo* interactions with proteins involved in the energy flow to mitochondria and identified potential substrates of PP2A-B'ζ employing comparative quantitative phosphoproteomics.

Methodology

Plant material and growth conditions

Seeds of *Arabidopsis thaliana* wild type (Col-0) and two *pp2a-b'ζ* knockout lines (SALK_150586C; "Z1" and SALK_107944C; "Z2") were surface-sterilized and stratified. Plants were grown under short day or long day conditions as described in Kataya et al. (2020). Mutant seeds were obtained from the European Arabidopsis Stock Centre (Nottingham, UK) and genotyped by PCR using T-DNA insertion primers obtained from SALK institute website SIGnAL (http://www.signal.salk.edu/tdnaprimers.2.html) as shown in Supplementary Table S1.

Gene cloning for *in planta* expression *in vivo*

Transient expression of constructs for bimolecular fluorescence complementation (BiFC) was done in two systems: *Agrobacterium* infiltration into *Nicothiana benthamiana* leaves and gene gun delivery into onion epidermal cells. BiFC constructs for onion transformation were created by cloning the genes for candidate interactors into pUC-VYNE$_{N173}$ and the gene for B'ζ (At2g21650) into pUC-VYCE$_{C155}$. For *Agrobacterium* constructs, the plasmids hygII-VYNE$_{N173}$ and kanII-VYCE$_{C155}$ were used for interactors and B'ζ, respectively (Supplementary Fig. S1). For *mMDH2* and *CSY5*, RNA was extracted from *Arabidopsis* flower tissue (Qiagen RNeasy Plant Mini, 74004) and reverse transcribed using SuperScript IV (ThermoFisher, 18090010) and PCR-amplified using Phusion polymerase (ThermoFisher, F531L). Genes from rest of the interactor candidates were obtained in plasmids from ABRC (https://abrc.osu.edu). Correctness of the inserts was verified by Sanger sequencing. All cloning primers are listed in Supplementary Table S1.

N. benthamiana plants were grown in soil under long day conditions and transformed as described in Schweiger and Schwenkert (2014). Gene gun transformation into onion epidermal cells was done as in Kataya et al. (2020). Interactions were analysed two days post infiltration for tobacco leaves and one to two days after bombardment for onion cells as described by Kataya et al. (2015a). Each interaction was analysed in at least three independent experiments.

Sugar dependence assay

Seedlings were grown on Linsmeier and Skoog (LS) for 7 days with or without sucrose (1%) under short-day conditions (light:dark 8h:16h) or in continuous darkness. The seedlings were

scanned using a CANON scanner, and hypocotyl length was measured using ImageJ (https://rsb.info.nih.gov/ij/).

Quantitative phosphoproteomics

Total protein was extracted from 170 mg tissue collected from 6-day-old seedlings grown on Murashige and Skoog (MS) without sugar (triplicate samples each from WT, Z1 and Z2) as described by Roitinger et al. (2015). The proteins were precipitated using the TCA/acetone method, collected by centrifugation, and processed for trypsin digestion using filter-aided sample preparation. Digestion was done overnight at 37°C using 0.22 µg/µL MS-grade trypsin (Promega, PRV5111) and the samples were purified on peptide desalting spin columns (Thermo Fisher Scientific, 89851). The phosphopeptides were labeled with nine isobaric tandem mass tags, using the TMT10plex kit (Thermo Fisher Scientific, 90110), and enriched by Sequential enrichment of Metal Oxide Affinity Chromatography (High-Select SMOAC; Thermo Fisher Scientific, A32993).

Liquid chromatography-tandem mass spectrometry (LC-MS/MS)

Tryptic peptides were analysed by LC-MS/MS at the Southern Alberta Mass Spectrometry Facility using an Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Fisher Scientific) controlled by Xcalibur (v4.0.21.10) and connected to a Thermo Scientific Easy-nLC 1200 system. Briefly, the Orbitrap first performed a full MS scan at a resolution of 120000 FWHM to detect the precursor ion (m/z from 375 to 1575, charge of +2 to +4). The AGC (auto gain control) and the max injection time were respectively set at 4e5 and 50 ms. The most intense precursor ions that showed a peptidic isotopic profile and had an intensity threshold over 2e4 were isolated using the quadrupole (window 0.7) and fragmented using HCD (38% collision energy) in the ion routing multipole. Fragment ions (MS2) were analyzed at a resolution of 15000, setting the first mass to 100 in order to acquire the TMT reporter ions (AGC 1e5 ms, max injection time 105 ms). To avoid acquisition of the same precursor ion having a similar m/z (plus or minus 10 ppm), dynamic exclusion was enabled for 45 sec.

Bioinformatics

Phosphopeptides that are more abundant in the two B'ζ mutants than in WT were identified using the DEqMS method (Zhu et al., 2020). Data was normalized applying equal median normalization before applying DEqMS. Furthermore, unmodified peptides and peptides only

oxidized were removed from the dataset. In the comparison between WT, Z1 and Z2, peptides were considered differentially phosphorylated if $p_{adj} < 0.05$ and absolute $\log_2$FC > 1.

Proteins involved in energy metabolism were subjected to gene expression analysis by Genevestigator (Zimmermann et al., 2004).

Mass spectrometry-verified phosphorylation sites were searched for in candidate PP2A-B'ζ interactors using the PhosPhAt4 database (https://phosphat.uni-hohenheim.de/; Heazlewood et al., 2008).

Subcellular localization predictions were done with the tool TargetP 2.0 (https://services.healthtech.dtu.dk/service.php?TargetP-2.0; Almagro et al., 2019).

Enriched Gene Ontology (GO) terms (Ashburner et al., 2020; Gene Ontology Consortium, 2021) were identified in the combined phosphoproteome datasets from WT and Z1. The GO analysis was performed using topGO v. 2.50.0 (Alexa & Rahnenfuhrer 2023) with the adjusted p-values from the DEqMS test of the WT vs Z1 data. Parameters for the GO analysis test were the default weight01 algorithm and Kolmogorov-Smirnov statistics. Annotations used were from the org.At.tair.db R package (Carlson, 2019).

Results and Discussion

Our previous work showed that B'ζ mutant seedlings displayed a growth retardation phenotype on sucrose-free medium (Kataya et al., 2015b), indicating a defect in energy metabolism. In addition, B'ζ was reported to be dually targeted to the cytosol and mitochondrion (Matre et al., 2009). We hypothesise that the PP2A-B'ζ holoenzyme is involved in regulation of the mitochondrial succinate/fumarate translocator or affect enzymes involved in the Krebs cycle. To test this, we set out to investigate the potential substrates of PP2A-B'ζ by bimolecular fluorescence complementation (BiFC) and phosphoproteomics analysis.

Sugar dependence assay points to a role in energy metabolism

To further investigate the function of PP2A-B'ζ, we employed two homozygotic knockout lines (SALK_150586C and SALK_107944C), hereafter referred to as Z1 and Z2. The knockout genotypes of both lines were confirmed by PCR. Line Z2 was previously observed to have impaired growth on sucrose-free medium (Kataya et al., 2015b), whereas line Z1 was added to the present study to strengthen our phenotypic and functional observations. Both mutants displayed impaired hypocotyl elongation on sucrose-free medium, especially when grown under continuous darkness (Fig. 1), clearly indicating a role for the B'ζ subunit in energy metabolism. No phenotype was apparent in Z1 or Z2 when grown on sucrose-containing MS or in soil, which is aligned with observations from Matre et al. (2009).

This observation made us look further into the enzymes of the Krebs cycle, with the aim of identifying potential substrates for PP2A-B'ζ-dependent dephosphorylation. Seven candidates were chosen for further investigations, based on their documented key roles in the Krebs cycle: Citrate synthase (CSY5), Aconitase (ACO3), Succinate dehydrogenase (SDH2-1), Mitochondrial succinate-fumarate carrier (mSFC1), Succinyl-CoA-ligase (SCS), Fumarase (FUM1) and Mitochondrial malate dehydrogenase (mMDH2). Since all candidates except mSFC are part of multigene families in *Arabidopsis*, we carefully chose for each protein the paralog that was most relevant for our study. As example, ACO3 carries the main aconitase activity out of the three *Arabidopsis* ACOs (de Bellis et al., 2020) and FUM1 was preferred over FUM2 since it is mitochondrion-targeted and essential (i.e., FUM2 cannot compensate for the lack of FUM1) (Pracharoenwattana et al., 2010).

Candidate B'ζ interactors carry Ser and Thr phosphorylation

First, we analysed the gene expression pattern of *B'ζ* and the genes coding for the seven metabolic proteins using GeneVestigator (Zimmermann et al., 2004). All genes were found to be expressed throughout development, but the *mSFC1*, *ACO3* and *B'ζ* genes are up-regulated under leaf senescence (Supplementary Fig. S2).

Next, we looked for evidence for Ser and Thr phosphorylation sites in the candidate substrates using the PhosPhAt4 database (Heazlewood et al., 2008). Experimentally verified phospho-Ser and phospho-Thr residues were identified in ACO3, FUM1 and SCS, making them plausible targets for dephosphorylation by PP2A-B'ζ (Supplementary Table S2). On the other hand, mSFC1, SDH2-1, mMDH2 and CSY5 each contain several predicted phosphorylation hotspots (Supplementary Table S2).

Verification of B'ζ interactors involved in energy flow to mitochondria

The genes encoding each enzyme and the carrier protein were cloned and the proteins were tested *in vivo* for their interaction with B'ζ by BiFC, a method that allows detection of protein-protein interactions that under natural conditions are transient and thus hard to observe. Two heterologous systems for plant transient expression were used: tobacco leaves (agroinfiltration) and onion epidermal cells (biolistic bombardment). Since mitochondrial localization of B'ζ depends on a free N-terminus (Matre et al., 2009), we tagged B' ζ with part of the fluorescent protein Venus in its C terminus. The other part of the fluorescent protein was linked to the C terminus of each candidate interactor (Supplementary Fig. S1).

In the first system, transient expression in tobacco leaves, the B'ζ fusion protein was seen to interact with mMDH2, in a location that appears to be the mitochondria (Fig. 2a). This association of B'ζ with a key enzyme of the Krebs cycle provides a first link between PP2A and the mitochondrial energy flow. mMDH2 is one out of two mitochondrial-localized MDH enzymes in *Arabidopsis* and catalyzes the conversion of malate into oxaloacetate in Krebs cycle. The mMDH enzymes play key roles in the early life stages in *Arabidopsis*, as evidenced from the severe phenotype of the *mmdh1mmdh2* double knockout (Sew et al., 2016). We also found a positive interaction of B'ζ with mSFC1 (Fig. 2b). This association takes place in the protruding parts of the tobacco cells, similar to the localization of YFP-tagged B'ζ in *Arabidopsis* (Matre et al., 2009).

A positive signal was seen between B'ζ and ACO3 in onion epidermal cells (Fig. 3a). In addition, physical interaction between B'ζ and mSFC1 was confirmed in this second system (Fig. 3b). To determine the localization of the positive interactions, the mitochondrial marker pWEN95 was used. The mitochondrial marker harbors the mitochondrial import peptide from the β-subunit of the mitochondrial ATP synthase gene (Boutry et al., 1987), fused with red fluorescent protein (RFP). The import peptide targets RFP to mitochondrial matrix. The succinate-fumarate carrier, mSFC1, was shown to be colocalized with the mitochondrial marker (Fig. 3c), suggesting that the interaction of PP2A-B'ζ and mSFC1 takes place in mitochondria or on the mitochondrial membrane in onion cells. Due to the mechanism of mitochondrial protein import, which requires protein unfolding for passage through the mitochondrial import system (Ford et al., 2022), the PP2A holoenzyme would need to disassemble before entry and then reform inside the mitochondrion. A mitochondrial targeting signal in one of the subunits would therefore not be able to direct the PP2A complex into the matrix. More likely, therefore, the interaction between mSFC1 and PP2A takes place on the mitochondrial surface. This interpretation agrees with the previous observation of mSFC1-GFP fusion protein at the mitochondrial membrane in tobacco epidermal cells (Brito et al., 2019).

The ACO3 interaction, on the other hand, was seen to be cytoplasmic (Fig. 3d). Overall, our BiFC data agree with previously reported localization of B'ζ to both the mitochondrion and the cytoplasm (Matre et al., 2009). ACO is an iron-sulfur-containing hydratase that is dually targeted to the mitochondrion and the cytoplasm in plants and yeast (Lushchak et al., 2014; de Bellis et al., 2020, Ben-Menachem et al., 2011). Mitochondrial ACO is active in the Krebs cycle, where it catalyzes the stereo-specific isomerization of citrate to isocitrate. The cytosolic form of *Arabidopsis* ACO3 was implicated in the regulation of organellar reactive oxygen species (ROS) homeostasis and was reported to bind to the PP2A-B'γ subunit (Konert et al., 2015). Our observation of cytosolic interaction between ACO3 and B'ζ indicates that PP2A-B'ζ could potentially be involved in ROS signalling as well, and that ACO3 is dephosphorylated by two different PP2A isozymes.

Co-localization of B'ζ with mSFC1 and ACO3 is in line with concordant up-regulation of their three respective genes in senescent leaves (Supplementary Fig. S2). No interaction was seen between PP2A-B'ζ and CSY5, SDH2-1, SCS or FUM1 in any of the two systems. Possibly, dephosphorylation of these metabolic enzymes takes place in other developmental

stages, in different tissue types or under specific conditions not tested here. In summary, BiFC revealed cytoplasmic interaction of PP2A-B'ζ with the metabolic enzyme ACO3 and the transporter protein mSFC1. The latter most likely takes place at the mitochondrial surface. All observations were strengthened from multiple independent transformation events and in two heterologous plant expression systems.

The mSFC1 carrier protein plays an important role in the exchange of metabolites between organelles (Catoni et al., 2003). By inference from the homologous protein in yeast, *Arabidopsis* mSFC1 was first believed to import peroxisomal-produced succinate into the mitochondrion in exchange for fumarate, which in turn via multiple downstream reactions would be used in gluconeogenesis or in the glyoxylate cycle (Catoni et al., 2003; Pracharoenwattana et al., 2010). However, in the absence of biochemical evidence for succinate/fumarase exchange of mSFC1, its function remained obscure, until more recently, it was revealed that mSFC1 transports citrate, isocitrate and aconitate more efficiently than fumarate and succinate. Thus, the main role of mSFC1 was proposed to be the exchange of cytosolic citrate for mitochondrial isocitrate (Brito et al., 2019). mSFC1 was found to play an important function in seed germination and seedling development, which is consistent with its expression in cotyledons, hypocotyls and root tips (Brito et al., 2019; Catoni et al., 2003). Physical interaction between B'ζ and mSFC1 suggests that dephosphorylation mediated by the PP2A-B'ζ complex plays a role in regulation of metabolite exchange and cellular energy production in *Arabidopsis*.

Additional metabolic enzymes revealed by phosphoproteomics

Encouraged by the results from BiFC, ACO3 and mSFC1 were cloned into an expression vector for *in vitro* interaction studies using microscale thermophoresis (MST). Recombinant proteins were produced in *Escherichia coli* and purified, but technical problems hampered the MST analysis. We therefore sought to employ an alternative method for studying B'ζ interactors: quantitative phosphoproteomics. In order to identify substrates affected by the absence of B'ζ, total proteins were extracted from seedlings of WT and the two knockout mutants, Z1 and Z2, and the three phosphoproteomes were compared. In this analysis, substrates were expected to be more abundantly phosphorylated in Z1 and Z2 than in the WT.

By applying differential expression analysis (DEqMS) on the phosphoproteomics data, we identified two novel B'ζ interactors directly involved in plant energy metabolism (Fig. 4,):

Phosphoenolpyruvate carboxylase 3 (PPC3) and Phosphoenolpyruvate carboxykinase 1 (PCK1). These two enzymes, plus a third protein, Pentotricopeptide repeat protein 6 (PPR6), had increased phosphorylation in the Z1 mutant compared to WT (Supplementary Table S3, Supplementary Table S4), as expected from abolished PP2A-B'ζ function in the *pp2a-b'ζ* knockout.

PPC3 catalyses the carboxylation of phosphoenolpyruvate (PEP) into oxaloacetate (OAA), whereas PCK1 acts in the opposite direction (i.e., forms PEP by OAA decarboxylation). PPC3 and PCK1 thereby play an important role in the regulation of the Krebs cycle. In fact, PCK1 serves as a cataplerotic enzyme, i.e., it removes intermediate metabolites from the Krebs cycle (Yang et al., 2009). Interestingly, the two enzymes are known to be regulated by phosphorylation, and PP2A was found to be the phosphatase responsible for dephosphorylating both PPC3 and PCK1 (Walker and Leegood, 1995; Carter et al., 1990; Shane et al., 2013). The identity of the regulatory subunit was, however, not known in these previous studies.

Meanwhile, the Ser and Thr sites that are specifically more abundantly phosphorylated in the Z1 mutant are supported by MS data deposited in the PhosPhAt4 repository (Supplementary Table S2). The detected phosphorylation in PPC3 (Ser 11) is also in agreement with a previous reporting PP2A-medited dephosphorylation at an N-terminal Ser (Moraes and Plaxton, 2001; Gregory et al., 2009).

PPR proteins play important roles in post-transcriptional regulation of mitochondrial genes (Manavski et al., 2012; Wang et al., 2021). They regulate many different aspects of the RNA life cycle, such as RNA stability, translation, processing, editing and splicing. As such, PPR proteins are important for cellular respiration in plants. PPR6 is predicted to contain a mitochondrial target peptide (TargetP 2.0, likelihood 0.9365), and therefore likely exerts its function inside mitochondria. The maize homolog of PPR6 is responsible for translation initiation and 5' end maturation of the mitochondrial ribosomal protein S3 (*rps3*) mRNA, with reduced translation and extended *rps3* mRNA 5' end in the *mppr6* mutant (Manavski et al., 2012). Since *Arabidopsis PPR6* can complement the maize phenotype, the function of PPR6 seems to be conserved. We therefore hypothesize that PP2A-B'ζ, via dephosphorylation of the PPR6 protein, affects mitochondrial mRNA stability, with consequences for mitochondrial energy metabolism.

ACO3 and mSFC1 were not found to the differentially phosphorylated in our comparison between the Z1, Z2 and WT phosphoproteomes. Potentially, PP2A-B'ζ shares these two substrates with PP2A holoenzymes containing other regulatory B subunits, which would dephosphorylate ACO3 and mSFC1 in the absence of B' ζ. The difference between our BiFC observations and phosphoproteomics data could also be due to the different tissue types being used – leaves from adult tobacco plants and onion epidermal cells vs. 6-day old whole *Arabidopsis* seedlings.

To get additional insights into which pathways and processes that might be regulated by protein phosphorylation-dephosphorylation processes, a GO term enrichment analysis was done. Since the number of differentially phosphorylated proteins in the Z1 mutant was rather low, the combined phosphoproteomic datasets from WT and Z1 were included in the pathway analysis. Notably, the terms phosphoenolpyruvate carboxylase activity (GO:0008964) and phosphoenolpyruvate carboxykinase activity (GO:0004612) were among the most significantly enriched terms in the Molecular Function domain (Supplementary Table S5), supporting the important role of reversible phosphorylation in PEP ← → OAA conversion and the regulation of energy metabolism.

In summary, our phosphoproteomics analysis revealed for the first time that the PP2A-B'ζ subunit is crucial for dephosphorylation of the metabolic enzymes PPC3, PCK1 and the mitochondrial gene regulator PPR6.

Conclusion

Knockout mutants of *PP2A-B'ζ* display impaired growth upon sucrose starvation, suggesting a deficiency in energy metabolism when the B'ζ regulatory subunit is removed. Along this line, the Krebs cycle enzymes PCK1 and PPC3, as well as the proposed mitochondrial gene expression regulator PPR6, were differentially phosphorylated in the *pp2a-b'ζ1* mutant. Interaction between PP2A-B'ζ and the metabolic enzymes mMDH2 and ACO3 was proven by BiFC. In addition, the mitochondrial metabolite carrier mSFC1 was shown to interact with B'ζ on the mitochondrial membrane. Taken together, results from this work (Fig. 5) suggest that the B'ζ-containing PP2A complex dephosphorylates several proteins involved in mitochondrial energy flow and thus plays an important role in plant metabolism.

Data availability

Unprocessed LC-MS/MS phosphosite data generated in this study is attached (supplementary dataset.xlsb).

References

Aggarwal KK, Saluja D, Sachar RC (1993) *Phytochemistry* **34**. Phosphorylation of rubisco in *Cicer arietinum*: non-phosphoprotein nature of rubisco in *Nicotiana tabacum*. https://doi.org/10.1016/0031-9422(93)80004-C

Alexa A, Rahnenfuhrer J (2023). *topGO: Enrichment Analysis for Gene Ontology*. R package version 2.52.0.

Almagro Armenteros JJ, Salvatore M, Emanuelsson O, Winther O, von Heijne G, Elofsson A, Nielsen H. (2019) *Life Sci Alliance* **2**. Detecting sequence signals in targeting peptides using deep learning. https://doi.org/10.26508/lsa.201900429

Andrés C, Lurin C, Small ID (2006) *Physiologica Plantarum* **129**. The multifarious roles of PPR proteins in plant mitochondrial gene expression. https://doi.org/10.1111/j.1399-3054.2006.00766.x

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. (2000) *Nat Genet.* **25**(1). Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. https://doi.org/10.1038/75556

Ben-Menachem R, Regev-Rudzki N, Pines O. (2011) *J Mol Biol.* **409**. The aconitase C-terminal domain is an independent dual targeting element. https://doi.org/10.1016/j.jmb.2011.03.045

Bheri M, Mahiwal S, Sanyal SK, Pandey GK (2021) *FEBS J.* **288**(3) Plant protein phosphatases: What do we know about their mechanism of action? https://doi.org/10.1111/febs.15454

Boutry M, Nagy F, Poulsen C, Aoyagi K, Chua NH. (1987) *Nature* **328**. Targeting of bacterial chloramphenicol acetyltransferase to mitochondria in transgenic plants. https://doi.org/10.1038/328340a0

Brito DS, Agrimi G, Charton L, Brilhaus D, Bitetto MG, Lana-Costa J, Messina E, Nascimento CP, Feitosa-Araújo E, Pires MV, Pérez-Díaz JL, Obata T, Porcelli V, Palmieri L, Araújo WL, Weber APM, Linka N, Fernie AR, Palmieri F, Nunes-Nesi A. (2020) *Biochem J.* **477**. Biochemical and functional characterization of a mitochondrial citrate carrier *in Arabidopsis thaliana*. https://doiorg/10.1042/BCJ20190785

Carlson M (2019). *org.At.tair.db: Genome wide annotation for Arabidopsis*. R package version 3.8.2.

Catoni E, Schwab R, Hilpert M, Desimone M, Schwacke R, Flügge UI, Schumacher K, Frommer WB. (2003) *FEBS Lett.* **534**. Identification of an Arabidopsis mitochondrial succinate-fumarate translocator. https://doi.org/10.1016/s0014-5793(02)03782-1

Carter PJ, Nimmo HG, Fewson CA, Wilkins MB (1990) *FEBS Lett* **263**. Bryophyllum fedtschenkoi protein phosphatase type 2A can dephosphorylate phosphoenolpyruvate carboxylase. https://doi.org/10.1016/0014-5793(90)81381-W

Chen X., Zhang, W., Zhang, B. et al. (2011) *Proteome Sci* **9**:37. Phosphoproteins regulated by heat stress in rice leaves. https://doi.org/10.1186/1477-5956-9-37

de Bellis L, Luvisi A, Alpi A. (2020) *Biology* **9**. Aconitase: to be or not to be inside plant glyoxysomes, that is the question. https://doi.org/10.3390/biology9070162

Engholm-Keller K, Larsen MR. (2011) *J Proteomics* **75**(2). Titanium dioxide as chemo-affinity chromatographic sorbent of biomolecular compounds--applications in acidic modification-specific proteomics. https://doi.org/10.1016/j.jprot.2011.07.024

Farkas I, Dombrádi V, Miskei M, Szabados L, Koncz C. (2007) *Trends Plant Sci.* **12**(4). *Arabidopsis* PPP family of serine/threonine phosphatases. https://doi.org/10.1016/j.tplants.2007.03.003

Ford HC, Allen WJ, Pereira GC, Liu X, Dillingham MS, Collinson I. (2022) *ELife* **11**:e75426. Towards a molecular mechanism underlying mitochondrial protein import through the TOM and TIM23 complexes. https://doi.org/10.7554/eLife.75426

Gene Ontology Consortium. The Gene Ontology resource: enriching a GOld mine. (2021) *Nucleic Acids Res.* **49**(D1):D325-D334. https://doi.org/10.1093/nar/gkaa1113

Grabsztunowicz M, Koskela MM, & Mulo P. (2017) Frontiers in Plant Science 8. Post-translational Modifications in Regulation of Chloroplast Function: Recent Advances. https://doi.org/10.3389/fpls.2017.00240

Gregory AL, Hurley BA, Tran HT, Valentine AJ, She YM, Knowles VL, Plaxton WC. (2009) *Biochem J.* **420**(1). *In vivo* regulatory phosphorylation of the phosphoenolpyruvate carboxylase AtPPC1 in phosphate-starved *Arabidopsis thaliana*. https://doi.org/10.1042/BJ20082397

Heazlewood JL, Durek P, Hummel J, Selbig J, Weckwerth W, Walther D, Schulze WX (2008) *Nucleic Acids Research* **36**. PhosPhAt: A Database of phosphorylation sites in *Arabidopsis thaliana* and a plant specific phosphorylation site predictor. https://doi.org/10.1093/nar/gkm812

Hodges M, Jossier M, Boex-Fontvieille E, & Tcherkez G. (2013) Plant Biol. 15. Protein phosphorylation and photorespiration. https://doi.org/10.1111/j.1438-8677.2012.00719.x

Kataya AR, Heidari B, Hagen L, Kommedal R, Slupphaug G, Lillo C. (2015a) *Plant Physiol*. **167**(2). Protein phosphatase 2A holoenzyme is targeted to peroxisomes by piggybacking and positively affects peroxisomal β-oxidation. https://doi.org/10.1104/pp.114.254409

Kataya AR, Heidari B, Lillo C. (2015b) *Plant Signal Behav*. **10**(5). Protein phosphatase 2A regulatory subunits affecting plant innate immunity, energy metabolism, and flowering time--joint functions among B'η subfamily members. https://doi.org/10.1080/15592324.2015.1026024

Kataya ARA, Elshobaky A, Heidari B, Dugassa NF, Thelen JJ, Lillo C. (2020) *Planta* **251**. Multi-targeted trehalose-6-phosphate phosphatase I harbors a novel peroxisomal targeting signal 1 and is essential for flowering and development. https://doi.org/10.1007/s00425-020-03389-z

Konert G, Trotta A, Kouvonen P, Rahikainen M, Durian G, Blokhina O, Fagerstedt K, Muth D, Corthals GL, Kangasjärvi S. (2015) *New Phytol*. **205**(3). Protein phosphatase 2A (PP2A) regulatory subunit B'γ interacts with cytoplasmic ACONITASE 3 and modulates the abundance of AOX1A and AOX1D in *Arabidopsis thaliana*. https://doi.org/10.1111/nph.13097

Li B, Zhang X-Q, Chollet R. (1996) *Plant Physiology* 111. Phosphoenolpyruvate carboxylase kinase in tobacco leaves is activated by light in a similar but not identical way as in maize. https://doi.org/10.1104/pp.111.2.497

Li P, Liu J (2021). *Protein Phosphorylation in Plant Cell Signaling*. In: Wu, X.N. (eds) *Plant Phosphoproteomics. Methods in Molecular Biology* **2358**. Humana, New York, NY. https://doi.org/10.1007/978-1-0716-1625-3_3

Lim S, Smith KR, Lim ST, Tian R, Lu J, Tan M. (2016) Cell Biosci 6. Regulation of mitochondrial functions by protein phosphorylation and dephosphorylation. https://doi.org/10.1186/s13578-016-0089-3

Lushchak OV, Piroddi M, Galli F, Lushchak VI. (2014) *Redox Rep*. **19**. Aconitase post-translational modification as a key in linkage between Krebs cycle, iron homeostasis, redox signaling, and metabolism of reactive oxygen species. https://doi.org/10.1179/1351000213Y.0000000073

Manavski N, Guyon V, Meurer J, Wienand U, Brettschneider R. (2012) Plant Cell **24**. An essential pentatricopeptide repeat protein facilitates 5' maturation and translation initiation of rps3 mRNA in maize mitochondria. https://doi.org/10.1105/tpc.112.099051

Matre P, Meyer C, Lillo C. (2009) *Planta* 230. Diversity in subcellular targeting of the PP2A B'η subfamily members. https://doi.org/10.1007/s00425-009-0998-z

Menz J, Li Z, Schulze WX, Ludewig U. (2016) *Plant J*. **88**(5):Early nitrogen-deprivation responses in Arabidopsis roots reveal distinct differences on transcriptome and (phospho-) proteome levels between nitrate and ammonium nutrition. https://doi.org/10.1111/tpj.13272

Moraes TF & Plaxton WC. (2000) *European Journal of Biochemistry* **267**. Purification and characterization of phospho*enol*pyruvate carboxylase from *Brassica napus* (rapeseed) suspension cell cultures. https://doi.org/10.1046/j.1432-1327.2000.01495.x

Pracharoenwattana I, Zhou W, Keech O, Francisco PB, Udomchalothorn T, Tschoep H, Stitt M, Gibon Y, Smith SM. (2010) *Plant J*. **62**. *Arabidopsis* has a cytosolic fumarase required for the massive allocation of photosynthate into fumaric acid and for rapid plant growth on high nitrogen. https://doi.org/10.1111/j.1365-313X.2010.04189.x

Qin T, Zhao P, Sun J et al. (2021) *Front Genet*. **12**:765580. Research Progress of PPR Proteins in RNA Editing, Stress Response, Plant Growth and Development. http://doi.org/10.3389/fgene.2021.765580

Rahikainen M, Trotta A, Alegre S, Pascual J, Vuorinen K, Overmyer K, Moffatt B, Ravanel S, Glawischnig E, Kangasjärvi S. (2017) *Plant J*. **89**(1). PP2A-B'γ modulates foliar trans-methylation capacity and the formation of 4-methoxy-indol-3-yl-methyl glucosinolate in *Arabidopsis* leaves. https://doi.org/10.1111/tpj.13326.

Rappsilber J, Ishihama Y, Mann M. (2003) *Anal Chem*. **75**(3). Stop and go extraction tips for matrix-assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. https://doi.org/10.1021/ac026117i

Rasool B, Karpinska B, Konert G, Durian G, Denessiouk K, Kangasjärvi S, Foyer CH. (2014) *Front Plant Sci*. **5**:405. Effects of light and the regulatory B-subunit composition of protein phosphatase 2A on the susceptibility of *Arabidopsis thaliana* to aphid (*Myzus persicae*) infestation. https://doi.org/10.3389/fpls.2014.00405.

Roitinger E, Hofer M, Köcher T, Pichler P, Novatchkova M, Yang J, Schlögelhofer P, Mechtler K. (2015) *Mol Cell Proteomics* **14**(3). Quantitative phosphoproteomics of the ataxia telangiectasia-mutated (ATM) and ataxia telangiectasia-mutated and rad3-related (ATR) dependent DNA damage response in *Arabidopsis thaliana*. https://doi.org/10.1074/mcp.M114.040352.

Schweiger R, Schwenkert S (2014) *J Vis Exp*. **85**. Protein-protein interactions visualized by bimolecular fluorescence complementation in tobacco protoplasts and leaves. https://doi.org/10.3791/51327

Segonzac C, Macho AP, Sanmartín M, Ntoukakis V, Sánchez-Serrano JJ, Zipfel C. (2014) *EMBO J*. **33**(18). Negative control of BAK1 by protein phosphatase 2A during plant innate immunity. https://doi.org/10.15252/embj.201488698.

Sew YS, Ströher E, Fenske R, Millar AH. (2016) *Plant Physiol*. **171**(2). Loss of mitochondrial malate dehydrogenase activity alters seed metabolism impairing seed maturation and post-germination growth in *Arabidopsis*. https://doi.org/10.1104/pp.16.01654

Shane MW, Fedosejevs ET, Plaxton WC (2013) *Plant Physiology* **161**. Reciprocal control of anaplerotic phospho*enol*pyruvate carboxylase by *in vivo* monoubiquitination and phosphorylation in developing proteoid roots of phosphate-deficient Harsh Hakea. https://doi.org/10.1104/pp.112.213496

Tenenbaum D, Maintainer B (2023). *KEGGREST: Client-side REST access to the Kyoto Encyclopedia of Genes and Genomes (KEGG)*. R package version 1.40.0.

Trotta A, Wrzaczek M, Scharte J, Tikkanen M, Konert G, Rahikainen M, Holmström M, Hiltunen HM, Rips S, Sipari N, Mulo P, Weis E, von Schaewen A, Aro EM, Kangasjärvi S. (2011) *Plant Physiol*. **156**(3). Regulatory subunit B'gamma of protein phosphatase 2A prevents unnecessary defense reactions under low light in *Arabidopsis*. https://doi.org/10.1104/pp.111.178442

Uhrig RG, Labandera AM, Moorhead GB (2013) *Trends Plant Sci*. **18**(9) Arabidopsis PPP family of serine/threonine protein phosphatases: many targets but few engines. http://doi.org/10.1016/j.tplants.2013.05.004

Urban J (2022) *Anal Chim Acta* **1199** A review on recent trends in the phosphoproteomics workflow. From sample preparation to data analysis. https://doi.org/10.1016/j.aca.2021.338857

van Wijk KJ, Friso G, Walther D, Schulze WX. (2014) *Plant Cell* **26**. Meta-analysis of *Arabidopsis thaliana* phospho-proteomics data reveals compartmentalization of phosphorylation motifs. https://doiorg/10.1105/tpc.114.125815

von Stechow L (2016) *Phospho-Proteomics Methods and Protocols* **2,** Springer, New York, NY. https://doi.org/10.1007/978-1-4939-3049-4

Walker, RP & Leegood, RC (1995) FEBS Lett. 362 Purification, and phosphorylation in vivo and in vitro, of phosphoenolpyruvate carboxykinase from cucumber cotyledons. https://doi.org/10.1016/0014-5793(95)00212-R

Wang X, An Y, Xu P, Xiao J. (2021) *Front Plant Sci*. **12**. Functioning of PPR proteins in organelle RNA metabolism and chloroplast biogenesis. https://doiorg/10.3389/fpls.2021.627501

Wu XN, Sanchez Rodriguez C, Pertl-Obermeyer H, Obermeyer G, Schulze WX (2013) *Mol Cell Proteomics*. 12(10) Sucrose-induced receptor kinase SIRK1 regulates a plasma membrane aquaporin in Arabidopsis. https://doi.org/10.1074/mcp.M113.029579

Wu X, Sklodowski K, Encke B, Schulze WX. (2014) *J Proteome Res*. **13**(7) A kinase-phosphatase signaling module with BSK8 and BSL2 involved in regulation of sucrose-phosphate synthase. https://doi.org/10.1021/pr5003164

Yang J, Kalhan SC, & Hanson RW (2009). *J. Biol. Chem.* **284** What is the metabolic role of phosphoenolpyruvate carboxykinase? https://doi.org/10.1074/jbc.R109.040543

Zhang H, Zhou H, Berke L, Heck AJ, Mohammed S, Scheres B, Menke FL (2013) *Mol Cell Proteomics* **12** Quantitative phosphoproteomics after auxin-stimulated lateral root induction identifies an SNX1 protein phosphorylation site required for growth. https://doi.org/10.1074/mcp.M112.021220

Zhu Y, Orre LM, Zhou Tran Y, et al. (2020) Molecular & Cellular Proteomics **19** DEqMS: a method for accurate variance estimation in differential protein expression analysis. https://doi.org/10.1074/mcp.TIR119.001646.

Zhu Q, Dugardeyn J, Zhang C, Tet al. (2012) *Plant J.* **71**(5) SLO2, a mitochondrial pentatricopeptide repeat protein affecting several RNA editing sites, is required for energy metabolism. https://doi.org/10.1111/j.1365-313X.2012.05036.x

Zhu X, Shen G, Wijewardene I, Cai Y, Esmaeili N, Sun L, Zhang H (2021) *Plant Physiol Biochem.* **169** The B'ζ subunit of protein phosphatase 2A negatively regulates ethylene signaling in Arabidopsis. https://doi.org/10.1016/j.plaphy.2021.10.037

Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W. (2004) *Plant Physiol.* **136** GENEVESTIGATOR. Arabidopsis microarray database and analysis toolbox. https://doi.org/10.1104/pp.104.046367

Figure 1. The *pp2a-b'ξ* knockout mutants show reduced growth on medium without sucrose. Hypocotyl length was measured on seedlings grown on MS with or without sucrose in **a** 8 h light/16 h darkness or **b** continuous darkness. Ten roots were measured per genotype and the experiment was repeated four times. * *P* < 0.05 and ** *P* < 0.01; two- tailed t-test with equal variance (no sucrose vs 1% sucrose). Error bars indicate standard deviation.

Figure 2. Interaction of PP2A-B'ζ with mMDH2 and mSFC1. BiFC constructs were transiently expressed in tobacco leaf epidermal cells. The C-terminal part of Venus was fused to B'ζ and the N-terminal Venus fragment was fused to **a** mMDH2 and **b** mSFC1. **i** Venus; **ii** brightfield. Both interactions are suggested to be mitochondrial.

Figure 3. Visualization of PP2A-B'ζ interaction with mSFC1 and ACO3 in onion epidermal cells via BiFC. B'ζ was tagged-terminally with the C-terminal part of Venus while the N-terminal portion of Venus was fused to the C terminus of ACO3 (**a** and **b**) and mSFC1(**c** and **d**). **i** Venus; **ii** brightfield; **iii** RFP (mitochondrial marker pWEN95); **iv** overlay of Venus and RFP; **v** overlay of Venus, RFP and brightfield. In b, the signals from Venus and the mitochondrial marker overlap, indicating mitochondrial localization of the mSFC1-B'ζ interaction. The absence of overlap with the mitochondrial marker in d suggests cytoplasmic localisation of the ACO3-B'ζ interaction.

Figure 4. Volcano plots of quantitative phosphoproteomics. Phosphosites with fold-changes >2 and adjusted *p* value <0.05 were regarded as significanlty differentially phosphorylated. The AGI identifiers corresponding to the corresponding three proteins are shown. **a**: WT vs. Z1. **b**: WT vs. Z2. **c**: Z1 vs. Z2.

Figure 5. Models of PP2A subunit composition and proposed function of PP2A-B'ζ. **a** The PP2A holoenzyme is composed of a scaffolding subunit ("A") a regulatory subunit ("B") and a catalytic subunit (C). The number of genes encoding each subunit in *Arabidopsis* is shown in parenthesis. The focus of this study, B'ζ, belongs to the B' protein family. **b** PP2A-B'ζ dephosphorylates multiple metabolic proteins. BiFC-validated substrate interactions of B'ζ are indicated with brown dashed lines and B'ζ -dependent phosphorylation sites revealed by phosophoproteomics are shown with blue dashed lines. Residues known to be phosphorylated are shown in the six substrates. S; serine, T; threonine, *, predicted phosphorylation. PCK1 catalyses the conversion of oxaloacetate (OAA) into phosphoenolpyruvate (PEP), whereas PPC3 catalyses the reverse reaction. Dephosphorylation of the mitochondrion-localised substrates could take place before entry into the mitochondrion or at the mitochondrial membrane (as is most likely for the carrier protein mSFC1).