Focusing attention in human working memory and long-term memory: benefits through dissociable processes

Dongyu Gong[1,2], Dejan Draschkow[1,2], Anna C. Nobre[1,2]

[1] Department of Experimental Psychology, University of Oxford, Oxford, UK

[2] Oxford Centre for Human Brain Activity, Wellcome Centre for Integrative Neuroimaging, Department of Psychiatry, University of Oxford, Oxford, UK

Author notes

Correspondence concerning this article should be addressed to Dongyu Gong (dongyu.gong@psy.ox.ac.uk), Dejan Draschkow (dejan.draschkow@psy.ox.ac.uk), or Anna C. Nobre (kia.nobre@psy.ox.ac.uk). Address: Oxford Centre for Human Brain Activity, Warneford Hospital, OX3 7JX, Oxford, UK.

ABSTRACT

We developed a new experimental approach to compare directly how attentional orienting

facilitates retrieval in human working memory and long-term memory, and how selective attention

within these different memory timescales impacts incoming sensory information processing. In

two experiments with healthy young adults (*N*s = 30 and 44), retrospective attention cues

prioritised an item encoded in working memory or long-term memory. Participants then retrieved

a memory item or performed a perceptual discrimination task. The cue was informative for the

retrieval task but not for the perceptual task. Attentional orienting improved memory retrieval for

both memory types and also enhanced discrimination for visual stimuli at the location matching

the prioritised working memory or long-term memory item. Eye-tracking data revealed a striking

dissociation in gaze biases related to orienting in working memory vs. long-term memory. The

findings suggest potent and at least partly dissociable attention-orienting processes for different

memory timescales.

Keywords: visual attention, internal attention, selective retrieval, retrocue, perceptual

discrimination, microsaccades, gaze bias

Selective attention is fundamental to human cognition, delivering information that is relevant to ongoing behaviour among boundless competing signals[1-4]. While standard models emphasise external sensory inputs as a primary target of attentional control[1,5], the arrow of attention can also point inward[6,7] to flexibly prioritise and select items within working memory (WM). Retrospective cues presented during WM delay period that predict goal-relevant items have proved highly useful for investigating the modulatory mechanisms of internal selective attention[8-10].

Interestingly, like during attentional selection in the external world[11-13], orienting attention within visual WM is accompanied by subtle gaze shifts[14-16]. Studies have also shown that selecting an item in WM leads to the capture of attention by sensory stimuli with matching locations or features[17-19]. These capture effects appear obligatory, operating even when they are counter-productive for the perceptual task[20], thus underscoring the strong role of WM in guiding external attention.

While there is a solid appreciation for the strong interrelation between attention and WM, a similar symbiotic relationship has been proposed for attention and long-term memory (LTM). LTM has long been considered to guide sensory processing, as exemplified by Helmholtz's view that perception results from unconscious inference based on prior memories[21]. Experimental tasks manipulating prior experience with stimulus contexts have since confirmed the role LTM plays in guiding attention[22-28]. In turn, researchers have posited that selective attention can also operate within LTM, guiding retrieval. Suggestive evidence comes from the observation that brain areas controlling external attentional orienting are activated in tasks of episodic memory retrieval[29-32]. Studies showing similar modulation of alpha-band neural activity during visual memory retrieval as during external visual attention have further hinted at a role for attentional selection in guiding LTM retrieval[33,34].

To build on this promising trail and test whether and how selective attention can modulate LTM retrieval, we need experimental tasks that manipulate the goal relevance of items in LTM. Such tasks should test directly whether orienting attention to a specific item in LTM yields performance benefits relative to retrieving other items associated with the same context. If positive evidence suggests the ability to orient attention in LTM flexibly and selectively, it would be interesting to probe how similar the processes and consequences are compared to orienting in WM. Would

orienting in LTM also be accompanied by gaze biases? Would the selective attention in LTM spill

over to modulate sensory processing? Answering these questions would inform enduring debates

regarding the relationship between WM and LTM[35-39].

Here, we introduce a new experimental framework to test the ability to orient attention selectively

among competing LTM contents and to compare directly selective attention in WM and LTM. We

borrow from retrospective cue (referred to as "retrocue" hereafter) designs to direct attention to a

specific item of a pre-learned array of competing items in LTM or to a specific item of an encoded

WM array presented earlier in the trial. In two studies, we measured the consequences of

attentional orienting in WM vs. LTM for memory retrieval and additionally tested for spill-over

effects on sensory processing. At the end of each trial, participants either retrieved a memory item

or performed a perceptual discrimination task. The retrocue was informative for the retrieval task

but not for the perceptual task. This allowed us to investigate whether attentional selection in WM

and LTM enhanced the retrieval of relevant memoranda and biased perceptual processing in an

unrelated task in similar ways.

Our first study showed that orienting attention in LTM conferred significant benefits to retrieval

and spill-over advantages for discriminating visual stimuli at matching locations, like orienting in

WM. The second study replicated and extended the pattern of results by probing for different

stimulus dimensions and incorporating eye tracking. Measuring subtle directional biases in gaze

allowed us to investigate if overt manifestations of orienting behaviour accompanied attentional

selection towards LTM items, as they do for WM contents. The results revealed surprising

differences between oculomotor engagement when orienting attention in LTM compared to WM.

EXPERIMENT 1

Methods

Participants. Thirty individuals (23 females, 7 males, $M = 25.13$ years, $SD = 4.15$) with reported

normal or corrected visual acuity volunteered and received monetary compensation for

participation. The experiment was approved by the Central University Research Ethics Committee

of the University of Oxford, and all participants provided informed consent before any

experimental procedure began.

Apparatus and stimuli. Participants sat in front of a 27″ monitor (1920 × 1080 pixels, 100 Hz)
and rested their chin on a chinrest placed 95 cm away from the monitor. The experiment was
programmed in MATLAB (MathWorks, Natick, MA, U.S.A.) using the Psychophysics Toolbox[40].
Stimuli appeared overlaid on a grey background. Throughout the experiment, four squares (2.5°
in diameter) were always presented as placeholders at the four quadrants, at 5° horizontally and
vertically from the central fixation to the centre of each square. The stimuli consisted of four
equiluminant colours (brown [183.5, 113, 19], green [65, 143, 110.5], lilac [138, 117.5, 190],
magenta [245.5, 37, 112.5]) drawn from a circle in CIELAB colour space.

Procedure and design. The experiment included a learning session and a testing session,
separated by a 5-minute break.

During the learning session, participants were trained to encode two colours and their
corresponding locations into LTM (**Fig. 1**). These two colours were randomly selected from the
four colours defined above, and they were always located along one of the two pairs of diagonal
locations (i.e., top left and bottom right, or top right and bottom left, counterbalanced across
participants). We refer to this pair of locations as LTM locations. Each learning trial began with a
fixation display lasting randomly between 800 and 1000 ms, after which the LTM display was
presented for 150 ms. Following a delay of 850 ms, participants were probed to reproduce either
the colour at one location or the location of one colour. On colour reproduction trials, a colour
wheel (containing 360 colours) was presented at the centre, and participants responded by rotating
the dial and selecting a colour along the wheel. The colour wheel was presented in a random
orientation on every colour reproduction trial. On location reproduction trials, one of the two
colours was presented at the centre, and participants responded by pressing one of four keys
mapped to the four locations. Performance feedback was presented after both colour and location
reproduction trials. Each to-be-learned attribute (two colours and two locations) was probed on 20
trials, resulting in a total of 80 learning trials presented in random order.

During the testing session, participants performed either a memory recall task or a perceptual
discrimination task on each trial (**Fig. 2A**). Each testing trial began with a fixation display (800-
1000 ms). A WM display followed, where the two colours unused in the learning session were

presented for 150 ms at the unused pair of diagonal locations. We refer to this pair of locations as

WM locations. To make sure the contents in WM were not fixed across trials, each WM colour

was randomly assigned to one of the WM locations on every trial. Following a delay of 850 ms, a

retrocue that was either neutral or informative was presented for 200 ms. The retrocue was neutral

on one-third of the trials; the fixation display changed to white, providing no information about

the item to be probed. The retrocue was informative on two-thirds of the trials; the fixation display

changed to one of the four colours, matching either a WM or LTM item with equal probability.

Informative retrocues indicated the item to be probed in the memory recall task with 100% validity.

Following a second delay of 800 ms after the retrocue, the memory recall task and the perceptual

discrimination tasks were equally likely to be presented.

In the memory recall task, participants were required to recall the location of the probed item. On

trials containing an informative retrocue, participants reported the location of the WM or LTM

item indicated by the retrocue. On trials containing a neutral cue, the probed item was indicated

by a centrally presented colour, chosen randomly from the WM or LTM colours. Responses were

delivered by pressing one of four keys mapped to the four locations (the same keys as used in the

location reproduction task during the learning session). Feedback was then presented for 500 ms

indicating whether the response was correct or wrong.

In the perceptual discrimination task, four arrows (length: 1.25°, width of the tail: 0.625°, RGB

value: [128, 128, 128]) were presented in the placeholders for 100 ms, after which randomly

generated Gaussian noise masks were applied to the four locations for 100 ms. Following the mask,

one location was probed, and participants pressed one of the arrow keys to report the arrow

direction at that location. Each of the four locations had an equal possibility of being probed. The

arrow direction at each location was independently drawn from four possible directions (↑ , ←,

↓ , →), resulting in $4^4 = 256$ possible combinations.

Participants were informed that the memorised items and the retrocue bore no predictive relation

concerning the location of the perceptual item to be discriminated. When retrocues were

informative, however, participants always needed to maintain the retrocued item in mind for

potential future use because of the randomisation of the memory recall and perceptual

discrimination trial order, which allowed us to examine any spill-over benefits elicited by the

retrocue on the perceptual discrimination task. The relationship between the perceptual
discrimination task and the retrocue was totally incidental because all locations were equally likely
to be probed no matter which location the retrocue would preferentially bias attention to. As a
result, only 25% of informative-retrocue trials in the perceptual discrimination task were
"matching" trials on which the probed sensory location coincided with the location of the retrocued
item.

The testing session consisted of 480 trials divided into 10 blocks (each including 48 trials). To
become familiarised with the task, participants performed an additional 48 practice trials before
testing.

Behavioural Analysis. Behavioural data were analysed in MATLAB. For the learning session, we
examined the average colour reproduction errors and location reproduction accuracy respectively,
by sorting the learning trials of each type into 4 bins (each containing 10 trials). Colour
reproduction errors (in units of degrees) were calculated by taking the absolute difference between
the angle of the target colour and the reproduced colour on the colour wheel. One-way repeated-
measures ANOVAs with linear contrast weights ([-3, -1, 1, 3]) across four bins tested for the
efficacy of training.

For the testing session, data from the memory recall and perceptual discrimination tasks were
analysed separately. During pre-processing, we excluded trials on which RTs were 3 SD above the
individual mean across all conditions in either task. After this exclusion step, an average of 98.35%
(SD = 0.44%) trials were retained in the analyses.

To test for benefits of internal selective attention on WM and LTM retrieval, we analysed the
average RTs and accuracy for the memory recall task as a function of retrocueing (neutral vs.
informative) and the memory timescale of probed items (WM vs. LTM). To examine whether
orienting attention to a memory item benefited subsequent perceptual processing at matching
locations, we compared perceptual discrimination accuracy on retrocue matching trials vs. neutral
trials (matching vs. neutral) when locations associated with WM or LTM items were probed (WM
vs. LTM). To gauge the quality of perception, accuracy was the dependent variable of interest, but
RTs were also evaluated for completeness. Additional analyses regarding the putative costs of
misdirected selective attention in WM and LTM for perceptual discrimination are presented in the

Supplementary Information.

When comparing behavioural performance between conditions, we applied a repeated-measures

ANOVA and reported partial η^2 as a measure of effect size. For post hoc *t*-tests, we reported

Bonferroni-corrected *p* values that we denoted as "$p_{\text{Bonferroni}}$". We reported Cohen's *d* as a measure

of effect size for all the *t*-tests. Where relevant, the within-subject standard error of the mean (*SEM*)

was calculated from normalised data[41]. When evaluating potential perceptual benefits elicited by

WM and LTM retrocues, we applied one-sample *t*-tests against 0.

Fig. 1. LTM training in Experiment 1. On every learning trial, participants memorised the same
colours and locations of two squares at diagonal locations. When prompted, they reproduced the
colour at the probed location or the location of the probed colour. Colour reproduction trials and
location reproduction trials were presented in random order. Learning performance was aggregated
into 4 bins (of 10 trials each) and shown in the form of errors for colour reproduction and accuracy
for location reproduction, respectively. Error bars represent ±1 *SEM* (*n* = 30).

Results

We first tested whether our LTM training in the learning session was effective, by evaluating colour

reproduction errors and location reproduction accuracy (**Fig. 1**). Trial bins exerted a significant

main effect on colour reproduction errors: a linear contrast analysis showed that participants

reproduced the LTM colours with smaller errors on late as compared to early trials (F (1, 26) =

16.156, $p < .001$, partial $\eta^2 = 0.157$). For the location reproduction task, participants remained near

ceiling-level accuracy across the four bins: first bin: 0.990 ± 0.005 ($M \pm SEM$), last bin: 0.997 ±

0.002, F (1, 26) = 0.480, $p = .490$. Although the location reproduction task was too easy to show

the location learning process, the improvement of colour reproduction performance suggests that

participants indeed learned the two items and stored them into LTM.

Attentional orienting improves WM and LTM recall

We then assessed the effects of WM and LTM retrocues on memory recall performance (**Fig. 2B**).

Both retrocueing (neutral vs. informative) and the memory timescale of probed items (WM vs.

LTM) significantly impacted retrieval speed: RTs in the memory recall task were faster when

retrocues were informative (F (1, 29) = 83.205, $p < .001$, partial $\eta^2 = 0.742$) and when WM items

were probed (F (1, 29) = 8.239, $p = .008$, partial $\eta^2 = 0.221$). Crucially, retrocueing and memory

timescale also interacted (F (1, 29) = 18.627, $p < .001$, partial $\eta^2 = 0.391$). Both WM and LTM

retrocues conferred a significant benefit (WM: t (29) = 9.217, $p_{Bonferroni} < .001$, $d = 1.521$; LTM: t

(29) = 6.575, $p_{Bonferroni} < .001$, $d = 0.833$), but retrocue benefits were stronger for the speed of

retrieving WM items (t (29) = 4.316, $p < .001$, $d = 0.748$).

Similar analyses on retrieval quality showed significant main effects of retrocueing (F (1, 29) =

4.340 $p = .046$, partial $\eta^2 = 0.130$) and memory timescale (F (1, 29) = 4.667, $p = .039$, partial $\eta^2 =$

0.139), as well as a significant interaction (F (1, 29) = 12.115, $p = .002$, partial $\eta^2 = 0.295$). Post-

hoc comparisons revealed a significant improvement in recall accuracy by retrocues for WM items

(t (29) = 2.981, $p_{Bonferroni} = .012$, $d = 0.465$) but no significant effect of retrocues for LTM items (t

(29) = 1.530, $p_{Bonferroni} = .274$). This could possibly be explained by the overall very high recall

accuracy, particularly so for LTM (0.977 ± 0.005), leaving little room for improvement.

Fig. 2. Prioritising WM and LTM representations guides recall and perception in Experiment 1. A, task schematic of the testing trials in Experiment 1. Participants first encoded two items into WM, with different colours and locations from those in LTM. After the first delay, neutral retrocues were uninformative, whereas informative retrocues (WM or LTM) indicated which location would be relevant in the memory recall task. After a second delay, the memory recall task and the perceptual discrimination task were equally likely to be presented. In the recall task, participants recalled the location of a randomly chosen colour on neutral-retrocue trials and the location of the retrocued colour on informative-retrocue trials. In the perceptual task, participants discriminated the direction of one of the four arrows that briefly appeared at the four quadrants before being masked. The location of the probed arrow was indicated by a post-cue. **B**, memory recall mean RTs and accuracy for WM and LTM items, grouped by neutral-retrocue and informative-retrocue trials. **C**, perceptual discrimination accuracy at WM and LTM locations, grouped by retrocue neutral and matching trials. **D**, means and distributions of perceptual accuracy benefits for WM and LTM locations matching retrocues. Error bars in **B-D** represent ±1 *SEM* (*n* = 30).

Attending to WM and LTM items benefits perceptual processing at matching locations

We then examined whether the selective prioritisation of WM and LTM memoranda by retrocues

also incidentally impacted perceptual discrimination of items occurring at the matching location

(see **Fig. 2C**).

Perceptual discrimination accuracy showed a main effect of retrocue matching (F (1, 29) = 31.541,

$p < .001$, partial $\eta^2 = 0.521$). Accuracy was superior for discriminating the arrow direction at the

retrocue matching location (0.699 ± 0.029) than for items on neutral-retrocue trials (0.592 ± 0.027).

Surprisingly, we also found a main effect of the memory timescale associated with probed

locations (F (1, 29) = 4.887, $p = .035$, partial $\eta^2 = 0.144$). Accuracy was higher when LTM

locations were probed (0.655 ± 0.026) compared to WM locations (0.601 ± 0.032). There was no

interaction between retrocue matching and memory timescale of the probed location, F (1, 29) =

0.926, $p = .344$. **Fig. 2D** shows that incidental retrocue benefits on perceptual discrimination were

significantly larger than zero for both memory timescales (WM: 0.094 ± 0.020, t (29) = 4.804, p

$< .001$, $d = 0.877$; LTM: 0.121 ± 0.028, t (29) = 4.362, $p < .001$, $d = 0.797$) but did not differ in

size (t (29) = 0.962, $p = .344$).

For completeness, analysis of RT revealed a main effect of retrocue matching on perceptual

discrimination RTs: participants were faster for items at matching vs. neutral locations (F (1, 29)

= 5.469, $p = .027$, partial $\eta^2 = 0.159$). There was no main effect of the memory timescale (F (1, 29)

= 0.194, $p = .663$) or interaction between the factors (F (1, 29) = 0.541, $p = .468$).

EXPERIMENT 2

Experiment 2 was designed to replicate and extend the general pattern of findings from Experiment

1 using more complex stimuli and a recall task that did not emphasize associations with the spatial

location of items.

In Experiment 1, items were defined by colour and location, with the memory recall task requiring

participants to reproduce the location of an item based on its colour. The recall task thus naturally

encouraged the prioritisation of the retrocued item's location in anticipation of the recall probe.

This follow-up experiment tested whether the location of the memoranda would be prioritised

automatically and incidentally benefit perceptual discrimination, even when the recall task did not

require retrieval of the item's location. Improvements in discriminating spatially matching visual

stimuli under such conditions would add to the growing evidence that visual memories preserve a

spatial layout based on the sensory layout at encoding[42-44] and that spatial locations can play a

special role in scaffolding memory contents[45-47]. To test for incidental spatial prioritisation in non-

spatial WM and LTM recall tasks, Experiment 2 used more complex stimuli defined by unique

shapes in addition to colours and required reporting of non-spatial features on memory recall trials.

Previous research demonstrated that systematic small shifts in gaze position (gaze biases)

accompany shifts in the internal focus of attention within working memory[14-16]. In Experiment 2,

we employed eye tracking to compare directly whether and how the involvement of the oculomotor

system might differ when WM and LTM representations are prioritised for potential use in the

memory recall task.

Methods

Participants. A total of 44 volunteers (27 females, 17 males, $M = 25.93$ years, $SD = 4.51$) with

reported normal or corrected-to-normal visual acuity were recruited. The sample size was

calculated using G*Power[48] to achieve 90% power for the one-sample t-tests performed to test the

significance of perceptual benefits following WM and LTM retrocues. The effect sizes for these

comparisons in Experiment 1 were 0.877 and 0.797. We assumed a conservative approach and

aimed to power for the detection of a medium effect size (0.5) because we expected that the

manipulation in Experiment 2 would lead to a smaller effect due to the incidental nature of the

spatial attributes in the task.

Most of the experimental setup was identical to Experiment 1, with the following modifications.

Apparatus and stimuli. Eye movements were recorded with the EyeLink 1000 Desktop Mount

(SR Research, Ottawa, ON, Canada) at 1000 Hz. Eye-tracker calibration used the built-in

calibration and validation protocols from the EyeLink software. When possible, horizontal and

vertical gaze positions were continuously recorded for both eyes. For some participants ($N = 13$),

only one eye was tracked due to a lack of good-quality binocular tracking (mostly because of

wearing glasses).

Stimuli appeared on a white background. Four shapes were equidistantly sampled from the

Validated Circular Shape (VCS) space[49], and then randomly assigned to each WM and LTM item,

adding a new feature dimension to the existing configurations in Experiment 1. These same four

shapes were used across all participants, but the shapes assigned to WM and LTM items were

randomised across participants.

Procedure and design. During the learning session, participants were trained to memorise the

colours and shapes of the two LTM items (**Fig. 3**). On every trial, they were probed to reproduce

either the colour or the shape of one item. At the response stage, either a colour wheel or a shape

wheel was presented at the centre, indicating the feature dimension to be reproduced in this trial.

Both the colour and shape wheels were presented in a random orientation each time. The shape

wheel consisted of 360 shapes from the VCS space. To avoid clustering, eight shapes sampled

from equidistant positions on the wheel were displayed along the cardinal axes (i.e., every 45

degrees). These eight shapes served as visual anchors, which were also randomly chosen every

time in accordance with the orientation of the shape wheel. Participants responded using a

computer mouse that controlled the dial on the wheel. Participants had unlimited time to retrieve

the item from memory and to decide what to reproduce. However, once they started moving the

dial, they had only 2500 ms to complete their reproduction. This was intended to encourage

participants to recall the exact colour or shape before moving the dial. The position of the dial

when participants clicked the left mouse button or when the time limit was reached was taken as

the response. Immediately after their response, participants received feedback for 500 ms. Each

colour and shape were probed on 20 trials, resulting in a total of 80 learning trials presented in

random order.

During the testing session, each WM shape was randomly combined with one of the WM colours

on every trial (**Fig. 4A**). As in Experiment 1, participants performed either a memory recall task

or a perceptual discrimination task on each trial equiprobably. In the memory recall task,

participants reproduced the shape of the item matching the retrocued colour or a randomly probed

colour when the retrocue was neutral. The shape wheel was identical to that used during the

learning session and was randomly rotated across trials. Uninformative, grey neutral retrocues

appeared on one-fifth of the trials. The remaining trials contained informative coloured retrocues

matching each of the LTM or WM colours with equal probability. The perceptual discrimination

task was the same as that in Experiment 1.

The testing session consisted of 600 trials divided into 10 blocks (each including 60 trials). To

become familiarised with the task, participants performed an additional 30 practice trials before

testing.

Behavioural Analysis. The analyses of interest were basically the same as in Experiment 1, with

location reproduction in the learning and testing sessions replaced by shape reproduction. Shape

reproduction errors (in units of degrees) were calculated by taking the absolute difference between

the angle of the target shape and the reproduced shape on the shape wheel. The RTs in the memory

recall task were calculated as the time from probe onset to when the response was recorded, either

when participants clicked the left mouse button or when the time limit was reached. After

excluding memory recall and perceptual discrimination trials on which RTs were 3 SD above the

individual mean across all conditions, an average of 98.66% (SD = 0.47%) trials were retained in

the analyses.

Eye-tracking Analysis. Data were first converted from edf to asc format and subsequently read

into RStudio. For binocularly tracked participants, data from the left and right eyes were averaged

to obtain a single horizontal and a single vertical gaze position channel. Blinks were marked by

detecting NaN clusters in the eye-tracking data and then interpolated using a linear interpolation

procedure. Data were epoched from 250 ms before to 1000 ms after cue onset. To make our

analyses more robust to drift of the eyes during fixation, we obtained the average gaze position

within the 250 ms window before cue onset for every trial and participant, and subtracted it from

every corresponding time course. We performed analyses on horizontal and vertical channels

separately. For both channels, we only included trials on which gaze position remained within $\pm$

50% from fixation (with 100% denoting the centres of the original item locations at a $\pm$ 5° visual

angle) throughout the course of the trial, as previous work showed that the gaze bias phenomenon

is constituted by a bias in gaze around fixation[14-16]. For the horizontal channel, data for all 44

participants were used, with an average of 9.6% ± 1.7% ($M ± SEM$) trials excluded per participant.

For the vertical channel, six participants had to be removed due to a high number of excluded trials

(> 50%). For the 38 participants retained, an average of 14.6% ± 2.1% trials were excluded. Gaze

time courses were smoothed using a 25-ms average moving window.

For the horizontal channel, we compared trial-averaged gaze-position time courses between

conditions in which the retrocued item occupied the left side (top/bottom left location) or the right

side (top/bottom right location) during encoding, separately for trials with WM retrocues and trials

with LTM retrocues. For the vertical channel, we did the same between conditions in which the

retrocued item occupied the bottom side (bottom left/right location) or the top side (top left/right

location) during encoding, also separately for trials with WM and LTM retrocues. For both

channels, trial-averaged gaze-position time courses were also obtained for trials with neutral

retrocues. To increase sensitivity, we also constructed a measure of towardness separately for trials

with WM and LTM retrocues, which expressed the gaze bias toward the side of the retrocued item

in a single value[14-16]. We did this for both horizontal and vertical channels to obtain towardness in

horizontal and vertical directions, respectively.

Statistical evaluation of the towardness time courses used a cluster-based permutation approach[50]

implemented in the *permuco* package[51], which is ideally suited to evaluate physiological effects

across multiple time points while retaining high sensitivity.

Results

As in Experiment 1, we first confirmed that participants learned the colours and shapes of LTM

items over training (**Fig. 3**). For the colour reproduction task, although the result of the linear

contrast analysis on colour reproduction errors was not significant ($F(1, 40) = 2.304$, $p = .132$),

the errors remained relatively low with a numerically decreasing trend across bins (first bin: 23.107°

± 2.295, last bin:19.182° ± 1.612). Learning performance improved significantly for the shape

reproduction task, with errors decreasing across bins ($F(1, 40) = 17.939$, $p < .001$, partial $\eta^2 =$

0.122).

394

Fig. 3. LTM training in Experiment 2. On every learning trial, participants memorised two consistent items at diagonal locations defined by unique colours and shapes. When prompted, they reproduced the colour or shape at one location. Colour reproduction trials and shape reproduction trials appeared in random order. Learning performance was sorted into 4 bins (with 10 trials each) and shown in the form of errors for both colour and shape reproduction. Error bars represent ±1 *SEM* ($n = 44$).

401

Attentional orienting improves WM and LTM recall

We examined the effects of WM and LTM retrocues on memory recall performance (**Fig. 4B**). Retrocueing significantly shortened RTs ($F (1, 43) = 179.198$, $p < .001$, partial $\eta^2 = 0.807$). Although RTs were equivalent for both memory timescales ($F (1, 43) = 0.436$, $p = .513$), there was a significant interaction between the two factors ($F (1, 43) = 24.568$, $p < .001$, partial $\eta^2 = 0.364$). RT benefits were present for both WM ($t (43) = 11.657$, $p_{Bonferroni} < .001$, $d = 0.594$) and LTM ($t (43) = 9.016$, $p_{Bonferroni} < .001$, $d = 0.273$) but were stronger for WM ($t (43) = 4.957$, $p < .001$, $d = 0.929$).

For recall accuracy, both retrocueing ($F (1, 43) = 8.971$, $p = .005$, partial $\eta^2 = 0.173$) and memory timescale ($F (1, 43) = 32.995$, $p < .001$, partial $\eta^2 = 0.434$) exerted significant main effects. The

412   two factors also interacted (F (1, 43) = 7.441, p = .006, partial η^2 = 0.163). Post-hoc comparisons

413   revealed that WM retrocues significantly reduced shape reproduction errors (t (43) = 4.322,

414    $p_{Bonferroni}$ < .001, d = 0.429) but LTM retrocues did not (t (43) = 1.250, $p_{Bonferroni}$ = .436). Shape

415   reproduction was very accurate and errors were smaller when LTM shapes were reproduced

416   (16.610° ± 1.519) as compared to WM shapes (27.603° ± 2.417), possibly reflecting ceiling effects.

Attending to WM and LTM items benefits perceptual processing at matching locations

We then examined whether retrocueing memory items incidentally benefitted items at matching

locations in the perceptual discrimination task, even though retrieving spatial locations was never

required for either task (**Fig. 4C**).

Perceptual discrimination accuracy was sensitive to both experimental factors. Discrimination was

more accurate for arrows at retrocue matching locations than items on neutral-retrocue trials (F (1,

43) = 9.504, p = .004, partial η^2 = 0.181) and for arrows appearing at LTM locations (F (1, 43) =

55.213, p < .001, partial η^2 = 0.562). Retrocue matching and memory timescale did not interact (F

(1, 43) = 0.067, p = .797). **Fig. 4D** shows that perceptual benefits for items matching WM and

LTM retrocue locations were both significantly larger than zero (WM: 0.037 ± 0.003, t (43) =

2.191, p = .034, d = 0.330; LTM: 0.032 ± 0.002, t (43) = 2.176, p = .035, d = 0.328) and did not

differ in size (t (43) = 0.259, p = .797).

RTs during perceptual discrimination were faster for arrows appearing at LTM locations (main

effect of memory timescale: F (1, 43) = 14.297, p < .001, partial η^2 = 0.250). Retrocue matching

had no significant effect (F (1, 43) = 0.638, p = .429) and the factors did not interact (F (1, 43) =

3.353, p = .074).

When comparing the size of perceptual benefits in Experiments 1 and 2, we found a main effect

of Experiment (1 vs. 2), F (1, 72) = 13.357, p < .001, partial η^2 = 0.157, indicating a significantly

larger benefit size in Experiment 1 compared to Experiment 2. No main effect of memory timescale

(WM vs. LTM, F (1, 72) = 0.560, p = .457) or interaction between Experiment and memory

timescale (F (1, 72) = 1.151, p = .287) was found.

Fig. 4. Prioritising WM and LTM representations guides recall and perception in Experiment 2. A, task schematic of the testing trials in Experiment 2. Participants first encoded two items in WM. Their colours, shapes, and locations differed from those in LTM. After the first delay, neutral retrocues were uninformative, whereas informative retrocues (WM or LTM) indicated which item shape would be relevant in the memory recall task. After a second delay, the memory recall task and the perceptual discrimination task were equally likely to be presented. In the recall task, participants recalled the shape of the item matching a randomly chosen colour on neutral-retrocue trials and the shape of the item matching the retrocued colour on informative-retrocue trials. In the perceptual task, participants discriminated the direction of one of the four arrows that briefly appeared at the four quadrants before being masked. The location of the probed arrow was indicated by a post-cue. **B**, memory recall mean RTs and accuracy for WM and LTM items, grouped by neutral-retrocue and informative-retrocue trials. **C**, perceptual discrimination accuracy at WM and LTM locations, grouped by retrocue neutral and matching trials. **D**, means and distributions of perceptual accuracy benefits for WM and LTM locations matching retrocues. Error bars in **B-D** represent ± 1 *SEM* ($n = 44$).

Orienting in WM but not LTM elicits directional gaze biases

Eye gaze measurements revealed a striking dissociation between orienting attention in WM and LTM. Replicating previous findings[14,15,45], WM retrocues elicited significant gaze biases in the

Fig. 5. Attention-related gaze biases following WM but not LTM retrocues in Experiment 2.
A, time courses of horizontal gaze biases following neutral, WM, and LTM retrocues. For WM and LTM retrocues, time courses are shown separately for conditions in which the retrocued item occupied the left side (top/bottom left location) or the right side (top/bottom right location) during encoding. **B**, time courses of horizontal gaze bias towardness on trials with WM and LTM retrocues, corresponding to the data in **A**. **C**, time courses of vertical gaze biases following neutral, WM, and LTM retrocues. For WM and LTM retrocues, time courses were shown separately for conditions in which the retrocued item occupied the top side (top left/right location) or the bottom side (bottom left/right location) during encoding. **D**, time courses of vertical gaze bias towardness on trials with WM and LTM retrocues, corresponding to the data in **C**. Horizontal lines below the time courses in **B** and **D** indicate significant temporal clusters (green: WM compared to zero; black: WM-LTM difference). No significant clusters were detected on trials with LTM retrocues in either horizontal or vertical directions.

direction of attentional orienting. In contrast, no evidence of similar gaze biases occurred when

orienting attention in LTM. **Fig. 5A** and **5C** show the horizontal and vertical gaze biases,

respectively, following neutral, WM, and LTM retrocues. Eye traces after neutral cues showed a

characteristic pattern of leftward and downward shifts in eye position. Compared to this baseline

pattern of gaze shifts, systematic changes were observed for trials with WM retrocues depending

on the direction of the retrocued item, both horizontally and vertically. No significant changes

occurred for trials with LTM retrocues, on which the baseline pattern of gaze shifts was observed

regardless of the retrocued item direction.

These differences between gaze biases following WM and LTM retrocues were evident in the

associated towardness time courses (**Fig. 5B** and **5D**). There were significant clusters in both

horizontal and vertical towardness time courses following WM retrocues (cluster-based

permutation tests, horizontal: $ps < .045$; vertical: $ps < .047$). Critically, there were significant

differences between the towardness time courses following WM and LTM retrocues along both

horizontal and vertical directions (cluster-based permutation tests, horizontal: approximately 450-

1000 ms after cue onset, $p = .002$; vertical: approximately 420-870 ms after cue onset, $p = .003$).

There were no significant clusters in either time course following LTM retrocues.

DISCUSSION

We demonstrated that internal attention-directing cues benefit both WM and LTM recall in humans

and that selective prioritisation of WM and LTM items also brought significant, spill-over

advantages for discriminating visual stimuli at the location matching the selected memory item.

Eye tracking revealed a striking difference in the mechanisms for internal shifts of attention in

WM vs. LTM. Our findings reinforce the increasing recognition of the strong and multidirectional

relationship between attention and memories of different timescales[52,53] as well as the plurality of

mechanisms supporting their interactions[7].

Our data demonstrate the capacity to orient attention flexibly to a specific item in LTM,

highlighting similarities with orienting in WM. In our tasks, orienting attention to contents in LTM

significantly improved retrieval speed in a similar, though less pronounced way as for WM. No

benefits were found for the accuracy of LTM, in contrast to WM, which could be related to the

very high retrieval accuracy in our tasks, especially for LTM. It will be important, therefore, to

explore possible consequences to retrieval accuracy in more challenging memory tasks. The

current pattern of results suggests that attention can, at least, enhance the accessibility or

"readiness" to act on LTM memory representations. The modulations observed for WM compared

to LTM retrieval were stronger, shortening response times to a greater degree and affecting

retrieval accuracy. The findings could point to differences in the types of mechanisms of internal

attention within these different memory domains or could, instead, reflect variations in the strength

of the memory representations themselves and the resulting retrieval demands. Future studies can

examine these interesting emerging possibilities systematically.

Across the two experiments, we also revealed spill-over effects that orienting in WM and LTM has

for sensory processing. In the perceptual discrimination task, prioritising items within the spatial

layout of either WM or LTM boosted the discrimination accuracy of location-matching stimuli.

While previous reports of this phenomenon focused on the influence of WM[54,55], our results

showed a comparable effect from prioritised LTM representations. There was no strategic benefit

for using the retrocue information in the perceptual task since the location of the retrocued item

only matched the probed location on 25% of the trials. Consequences of memory selection for

sensory processing were therefore likely incidental.

Moreover, when locations were never probed for reproduction in Experiment 2 (colour-shape

bindings were sufficient to perform well in the memory recall task), we still observed

discrimination accuracy benefits following WM and LTM retrocues. This suggests memory

retrieval involves prioritization of location information even when this is not required, thereby

supporting prior research indicating the privileged role of spatial locations in binding features held

in memory[45-47]. Furthermore, shape reporting on a randomly oriented wheel in Experiment 2 also

controlled for the possible effects of motor preparation in Experiment 1 related to the consistent

response mappings of location reports. Thus, retrocues necessarily enhanced recall performance

by improving the accessibility to and/or quality of sensory information. Future research should

address in detail the various possible targets and modulatory mechanisms underlying internal

attention in WM and LTM[7,52].

Our eye-tracking data yielded the most suggestive evidence for a functional dissociation between
the mechanisms for orienting attention in WM and LTM. We found a striking difference in
oculomotor involvement on trials with WM and LTM retrocues by capitalising on subtle
directional biases in human gaze behaviour[11-13]. Consistent with previous work[14-16], we observed
that gaze was biased toward retrocued WM items. Critically, this bias was more pronounced in
WM compared to LTM. No significant gaze biases were detected following LTM retrocues. The
findings thus reveal differential involvement of oculomotor mechanisms when orienting in WM
vs LTM.

An unexpected finding in our experiments was the better perceptual discrimination performance
when LTM locations were probed compared to WM locations in both experiments (higher accuracy
for LTM locations in Experiment 1, and both higher accuracy and faster RTs for LTM locations in
Experiment 2). It will be fascinating to explore whether this finding points to a primacy of LTM
to guide perception[20] or rather reflects a peculiarity of our experimental conditions. For example,
the greater consistency of feature mappings to LTM locations than WM locations in our tasks may
have decreased competition among memories and lead to stronger sensory biases. Or, instead, the
results may be tapping into something more fundamental, such as relevant LTM representations
exerting stronger and more automatic biases, akin to sensory salience effects[56,57].

In summary, we compared the effects of focusing attention within human WM vs. LTM and
discovered that both bring significant benefits to retrieval and subsequent sensory processing, most
likely through dissociable mechanisms. The strikingly distinct oculomotor signatures of covert
spatial attention in WM and LTM corroborate a plurality of functional properties when memories
of different timescales guide adaptive behaviour and open novel opportunities for furthering our
understanding of the relationship between WM and LTM.

Code availability

Code is available from the authors on reasonable request.

Data availability

All data are publicly available through the Open Science Framework at https://osf.io/n629s/.

REFERENCES

1. Desimone, R., & Duncan, J. S. Neural Mechanisms of Selective Visual Attention. *Annual*
Review of Neuroscience **18**, 193–222 (1995).

2. *The handbook of attention* (MIT Press, 2015).

3. Nobre, A. C. Attention. in *Stevens' Handbook of Experimental Psychology and Cognitive*
Neuroscience (ed. Wixted, J. T.) 1–75 (Wiley, 2018).

4. *The Oxford handbook of attention* (Oxford University Press, 2014).

5. Corbetta, M., & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the
brain. *Nature Reviews Neuroscience* **3**, 201–215 (2002).

6. Chun, M. M., Golomb, J. D., & Turk-Browne, N. B. A Taxonomy of External and Internal
Attention. *Annual Review of Psychology* **62**, 73–101 (2011).

7. van Ede, F., & Nobre, A. C. Turning Attention Inside Out: How Working Memory Serves
Behavior. *Annual review of psychology* **74**, 137-165 (2023).

8. Griffin, I. C., & Nobre, A. C. Orienting Attention to Locations in Internal Representations.
Journal of Cognitive Neuroscience **15**, 1176–1194 (2003).

9. Landman, R., Spekreijse, H., & Lamme, V. A. F. Large capacity storage of integrated objects
before change blindness. *Vision Research* **43**, 149–164 (2003).

10. Souza, A. S., & Oberauer, K. In search of the focus of attention in working memory: 13 years
of the retrocue effect. *Attention, Perception, & Psychophysics* **78**, 1839–1860 (2016).

11. Engbert, R., & Kliegl, R. Microsaccades uncover the orientation of covert attention. *Vision*

Research **43**, 1035–1045 (2003).

12. Hafed, Z. M., & Clark, J. J. Microsaccades as an overt measure of covert attention shifts. *Vision Research* **42**, 2533–2545 (2002).

13. Corneil, B. D., & Munoz, D. P. Overt Responses during Covert Orienting. *Neuron*, *82*, 1230–1243 (2014).

14. Draschkow, D., Nobre, A. C., & van Ede, F. Multiple spatial frames for immersive working memory. *Nature Human Behaviour* **6**, 536-544 (2022).

15. van Ede, F., Chekroud, S. R., & Nobre, A. C. Human gaze tracks attentional focusing in memorized visual space. *Nature Human Behaviour* **3**, 462–470 (2019).

16. van Ede, F., Board, A., & Nobre, A. Goal-directed and stimulus-driven selection of internal representations. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 24590–24598 (2020).

17. Awh, E., & Jonides, J. Overlapping mechanisms of attention and spatial working memory. *Trends in Cognitive Sciences* **5**, 119–126 (2001).

18. Downing, P. E. Interactions between visual working memory and selective attention. *Psychological Science* **11**, 467–473 (2000).

19. Soto, D., Humphreys, G. W., & Rotshtein, P. Dissociating the neural mechanisms of memory-based guidance of visual selection. *Proceedings of the National Academy of Sciences* **104**, 17186–17191 (2007).

20. Soto, D., & Humphreys, G. W. Working Memory Biases in Human Vision. in *The Oxford Handbook of Attention* (eds. Nobre, A. C. & Kastner, S.) 753–776 (Oxford University Press, 2014).

21. Helmholtz, H. *Handbuch der physiologischen Optik*. In *Allgemeine Encyklopädie der Physik* Vol. 9 (ed. Karsten, G.) 37–51 (Voss, 1867).

22. Brockmole, J. R., Castelhano, M. S., & Henderson, J. M. Contextual cueing in naturalistic scenes: Global and local contexts. *Journal of Experimental Psychology: Learning, Memory, and*

Cognition **32**, 699–706 (2006).

23. Chun, M. M., & Jiang, Y. Contextual cueing: Implicit learning and memory of visual context
guides spatial attention. *Cognitive Psychology* **36**, 28–71 (1998).

24. Fan, J. E., & Turk-Browne, N. B. Incidental biasing of attention from visual long-term
memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition* **42**, 970–977
(2016).

25. Stokes, M. G., Atherton, K. E., Patai, E. Z., & Nobre, A. C. Long-term memory prepares
neural activity for perception. *Proceedings of the National Academy of Sciences of the United*
States of America **109**, 1838–1839 (2012).

26. Summerfield, J. J., Lepsien, J., Gitelman, D. R., Mesulam, M.-M., & Nobre, A. C. Orienting
attention based on long-term memory experience. *Neuron* **49**, 905–916 (2006).

27. Theeuwes, J., Bogaerts, L., & van Moorselaar, D. What to expect where and when: How
statistical learning drives visual selection. *Trends in Cognitive Sciences* **26**, 860–872 (2022).

28. Võ, M. L.-H., Boettcher, S. E., & Draschkow, D. Reading scenes: How scene grammar guides
attention and aids perception in real-world environments. *Current Opinion in Psychology* **29**, 205–
210 (2019).

29. Cabeza, R., Ciaramelli, E., Olson, I. R., & Moscovitch, M. The parietal cortex and episodic
memory: An attentional account. *Nature Reviews Neuroscience* **9**, 613–625 (2008).

30. Chun, M. M., & Johnson, M. K. Memory: Enduring Traces of Perceptual and Reflective
Attention. *Neuron* **72**, 520–535 (2011).

31. Ciaramelli, E., Grady, C. L., & Moscovitch, M. Top-down and bottom-up attention to
memory: A hypothesis (AtoM) on the role of the posterior parietal cortex in memory retrieval.
Neuropsychologia **46**, 1828–1851 (2008).

32. Sestieri, C., Shulman, G. L., & Corbetta, M. The contribution of the human posterior parietal
cortex to episodic memory. *Nature Reviews Neuroscience* **18**, 183-192 (2017).

33. Sabo, M., & Schneider, D. Pattern reinstatement and attentional control overlap during

episodic long-term memory retrieval. *Scientific Reports* **12**, 10739 (2022).

34. Sutterer, D. W., Foster, J. J., Serences, J. T., Vogel, E. K., & Awh, E. Alpha-band

oscillations track the retrieval of precise spatial representations from long-term memory. *Journal*

of Neurophysiology **122**, 539–551 (2019).

35. Atkinson, R. C., & Shiffrin, R. M. Human memory: a proposed system and its control

processes. *Psychology of Learning and Motivation* **2**, 89–195 (1968).

36. Cowan, N. What are the differences between long-term, short-term, and working

memory?. *Progress in brain research* **169**, 323-338 (2008).

37. Fukuda, K., & Woodman, G. F. Visual working memory buffers information retrieved from

visual long-term memory. *Proceedings of the National Academy of Sciences of the United States*

of America **114**, 5306–5311 (2017).

38. Hirschstein, Z., & Aly, M. Long-term memory and working memory compete and cooperate

to guide attention. *Attention, Perception, & Psychophysics*, 1-33 (2022).

39. Logie, R. H. *Visuo-spatial Working Memory* (Psychology Press, 1995).

40. Brainard, D. H. The Psychophysics Toolbox. *Spatial Vision* **10**, 433–436 (1997).

41. Cousineau, D. Confidence intervals in within-subject designs: A simpler solution to Loftus

and Masson's method. *Tutorials in quantitative methods for psychology* **1**, 42-45 (2005).

42. Dell'Acqua, R., Sessa, P., Toffanin, P., Luria, R., & Jolicœur, P. Orienting attention to objects

in visual short-term memory. *Neuropsychologia* **48**, 419–428 (2010).

43. Groen, I. I. A., Dekker, T. M., Knapen, T., & Silson, E. H. Visuospatial coding as ubiquitous

scaffolding for human cognition. *Trends in Cognitive Sciences* **26**, 81–96 (2022).

44. Kuo, B.-C., Rao, A., Lepsien, J., & Nobre, A. C. Searching for Targets within the Spatial

Layout of Visual Short-Term Memory. *Journal of Neuroscience* **29**, 8032–8038 (2009).

45. Pertzov, Y., & Husain, M. The privileged role of location in visual working memory. *Attention,*

Perception, & Psychophysics **76**, 1914–1924 (2014).

46. Schneegans, S., & Bays, P. M. Neural Architecture for Feature Binding in Visual Working
Memory. *Journal of Neuroscience* **37**, 3913–3925 (2017).

47. Treisman, A., & Zhang, W. Location and Binding in Visual Working Memory. *Memory &*
Cognition **34**, 1704–1719 (2006).

48. Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*
Methods **39**, 175–191 (2007).

49. Li, A. Y., Liang, J. C., Lee, A. C. H., & Barense, M. D. The validated circular shape space:
Quantifying the visual similarity of shape. *Journal of Experimental Psychology: General* **149**,
949–966 (2020).

50. Maris, E., & Oostenveld, R. Nonparametric statistical testing of EEG- and MEG-data.
Journal of Neuroscience Methods **164**, 177–190 (2007).

51. Frossard, J., & Renaud, O. Permutation Tests for Regression, ANOVA, and Comparison of
Signals: The permuco Package. *Journal of Statistical Software* **99**, 1–32 (2021).

52. Nobre, A. C., & Stokes, M. G. Premembering Experience: A Hierarchy of Time-Scales for
Proactive Attention. *Neuron* **104**, 132–146 (2019).

53. Nobre, A. C., & Stokes, M. G. Memory and attention: the back and forth. In *The Cognitive*
Neurosciences (ed. Gazzaniga, M.S.) 291–300 (MIT Press, 2020).

54. Olivers, C. N. L., Peters, J., Houtkamp, R., & Roelfsema, P. R. Different states in visual
working memory: When it guides attention and when it does not. *Trends in Cognitive Sciences*
15, 327–334 (2011).

55. Soto, D., Hodsoll, J., Rotshtein, P., & Humphreys, G. W. Automatic guidance of attention
from working memory. *Trends in Cognitive Sciences* **12**, 342–348 (2008).

56. Theeuwes, J. Goal-driven, stimulus-driven, and history-driven selection. *Current opinion in*

psychology **29**, 97-101 (2019).

57. Wolfe J. M. Guided Search 6.0: An updated model of visual search. *Psychonomic bulletin &*
review **28**, 1060–1092 (2021).

Acknowledgements

This research was funded by a Clarendon Scholarship, a Medical Research Council Studentship,
and a New College-Yeotown Scholarship to D.G.; a Wellcome Trust Senior Investigator Award
(104571/Z/14/Z) and a James S. McDonnell Foundation Understanding Human Cognition
Collaborative Award (220020448) to A.C.N.; and by the NIHR Oxford Health Biomedical
Research Centre. The Wellcome Centre for Integrative Neuroimaging is supported by core funding
from the Wellcome Trust (203139/Z/16/Z). For the purpose of open access, the author has applied
a CC BY public copyright licence to any Author Accepted Manuscript version arising from this
submission.

Author contributions

696   D.G., D.D. and A.C.N. conceived and designed the experiments. D.G. acquired and analysed the
697   data. D.G., D.D. and A.C.N. interpreted the data. D.G., D.D. and A.C.N. drafted and revised the
698   manuscript.